

COMPLETELY INTEGRABLE SYSTEMS WITH A SYMMETRY IN COORDINATES*

TOSHIO OSHIMA[†]

Abstract. We explicitly construct the integrals of completely integrable quantum or classical systems whose potential functions are invariant under the action of a classical Weyl group. Our potential functions and integrals are expressed by the Weierstrass elliptic function.

1. Introduction. Many completely integrable quantum or classical dynamical systems have been constructed in connection with root systems (cf. [OP1], [OP2], [In]). Consequently most of them are invariant under the action of the corresponding Weyl groups. Our study is to determine all the completely integrable systems with this invariant property.

Let W be the Weyl group of type A_{n-1} with $n \geq 3$ or of type B_n with $n \geq 2$ or of type D_n with $n \geq 4$. We identify W with the group of the coordinate transformations

$$(x_1, \dots, x_n) \mapsto (\varepsilon_1 x_{\sigma(1)}, \dots, \varepsilon_n x_{\sigma(n)})$$

of \mathbb{R}^n , where σ are the elements of the n -th permutation group \mathfrak{S}_n and

$$\begin{cases} \varepsilon_1 = \dots = \varepsilon_n = 1 & \text{if } W \text{ is of type } A_{n-1}, \\ \varepsilon_1 = \pm 1, \dots, \varepsilon_n = \pm 1 & \text{if } W \text{ is of type } B_n, \\ \varepsilon_1 = \pm 1, \dots, \varepsilon_n = \pm 1 \text{ and } \#\{i; \varepsilon_i = -1\} \text{ is even} & \text{if } W \text{ is of type } D_n. \end{cases}$$

We study the Schrödinger operator

$$(1.1) \quad P = -\frac{1}{2} \sum_{1 \leq j \leq n} \frac{\partial^2}{\partial x_j^2} + R(x)$$

on \mathbb{R}^n with a W -invariant potential function $R(x)$ which has enough W -invariant commuting differential operators assuring the complete integrability of P . To be precise we assume that there exist W -invariant differential operators P_1, \dots, P_n with

$$(1.2) \quad [P_i, P_j] = 0 \quad \text{for } 1 \leq i < j \leq n$$

and

$$(1.3) \quad P \in \mathbb{C}[P_1, \dots, P_n]$$

such that

$$(1.4) \quad P_j = \sum_{1 \leq i_1 < \dots < i_j \leq n} \partial_{i_1} \dots \partial_{i_j} + R_j \quad \text{with } \text{ord } R_j < j \quad \text{for } 1 \leq j \leq n$$

or

$$(1.5) \quad P_j = \sum_{1 \leq i_1 < \dots < i_j \leq n} \partial_{i_1}^2 \dots \partial_{i_j}^2 + R_j \quad \text{with } \text{ord } R_j < 2j \quad \text{for } 1 \leq j \leq n$$

* Received November 1, 1998; accepted for publication February 9, 1999.

[†] Department of Mathematical Sciences, University of Tokyo, Komaba, Tokyo 153-8914, Japan (oshima@ms.u-tokyo.ac.jp).

or

$$(1.6) \quad \begin{cases} P_n = \partial_1 \cdots \partial_n + R_n & \text{with } \text{ord } R_n < n, \\ P_j = \sum_{1 \leq i_1 < \cdots < i_j \leq n} \partial_{i_1}^2 \cdots \partial_{i_j}^2 + R_j & \text{with } \text{ord } R_j < 2j \text{ for } 1 \leq j < n \end{cases}$$

if the type of W is A_{n-1} or B_n or D_n , respectively. Here $\mathbb{C}[P_1, \dots, P_n]$ is the commutative algebra generated by P_1, \dots, P_n , $\text{ord } R_j$ denote the orders of differential operators R_j and for simplicity we put $\partial_i = \frac{\partial}{\partial x_i}$.

We assume that the coefficients of the differential operators are extended to holomorphic functions on a Zariski open subset of an open connected neighborhood of the origin of the complexification \mathbb{C}^n of \mathbb{R}^n .

The main result of our previous paper [OS] is the following:

If W is of type A_{n-1} with $n \geq 3$, then

$$(1.7) \quad R(x) = \sum_{1 \leq i < j \leq n} u(x_i - x_j)$$

with

$$(1.8) \quad u(t) = C_1 \wp(t) + C_2.$$

If W is of type B_n with $n \geq 2$, then

$$(1.9) \quad R(x) = \sum_{1 \leq i < j \leq n} \left(u(x_i - x_j) + u(x_i + x_j) \right) + \sum_{1 \leq j \leq n} v(x_j).$$

Here if $n \geq 3$, we have

$$(1.10) \quad \begin{cases} u(t) = C_1 \wp(t) + C_2, \\ v(t) = \frac{C_3 \wp(t)^4 + C_4 \wp(t)^3 + C_5 \wp(t)^2 + C_6 \wp(t) + C_7}{\wp'(t)^2} \end{cases}$$

or

$$(1.11) \quad u(t) = C_1 t^{-2} + C_2 t^2 + C_3 \quad \text{and} \quad v(t) = C_4 t^{-2} + C_5 t^2 + C_6$$

or

$$(1.12) \quad u(t) = C_1 \quad \text{and} \quad v(t) \text{ is any even function.}$$

If W is of type D_n with $n \geq 4$, then (1.9) holds with $v = 0$ and u is given by (1.10) or (1.11).

Here C_1, C_2, \dots are complex numbers and $\wp(t)$ is the Weierstrass elliptic function $\wp(t|2\omega_1, 2\omega_2)$ with primitive half-periods ω_1 and ω_2 , which are allowed to be infinity.

The purpose of this paper is to construct the operators P_1, \dots, P_n mentioned above when u or (u, v) is given by (1.8) or (1.10) for any complex numbers C_1, C_2, \dots and for any periods of the elliptic function (cf. Theorem 7.2, 7.3 and 7.5), which was announced in [OOS]. Hence we shall have the complete integrability of the corresponding Schrödinger operator (1.1). We remark that if W is of type A_{n-1} , the complete integrability and the operators P_1, \dots, P_n are already known (cf. [Ca], [Su], [OP2], [OS], [Et], Theorem 3.2 in this paper).

Taking the ‘‘classical limit’’, we shall also obtain the integrals of the Hamiltonian corresponding to the Schrödinger operator (1.1) because of our simple expression of the operators P_1, \dots, P_n .

When W is of type B_2 , our argument in this paper is valid but there exist other potentials which assure the complete integrability. This is caused by a symmetry between u and v . We shall treat this case in another paper (cf. [OOS], [OO], [Oc]).

If u or (u, v) is given by (1.11), the operators P_1, \dots, P_n do not exist in general and then we need W -invariant operators of higher orders (cf. [OP2]), which will be discussed in future.

If (u, v) is given by (1.12), the algebra $\mathbb{C}[P_1, \dots, P_n]$ equals the totality of \mathfrak{S}_n -invariants of $\mathbb{C}[-\frac{1}{2}\partial_1^2 + v(x_1), \dots, -\frac{1}{2}\partial_n^2 + v(x_n)]$.

We note that if $2\omega_1 = \sqrt{-1}\pi$ and $\omega_2 = \infty$, then (1.10) is reduced to

$$(1.13) \quad \begin{cases} u(t) = C'_1 \sinh^{-2} t + C'_2, \\ v(t) = C'_3 \sinh^{-2} t + C'_4 \sinh^{-2} 2t + C'_5 \sinh^2 t + C'_6 \sinh^2 2t + C'_7 \end{cases}$$

with complex numbers C'_1, \dots, C'_7 . The system studied by Heckman-Opdam ([He1], [He2], [HO], [Op1] and [Op2]) corresponds to this trigonometric case with $C'_5 = C'_6 = 0$ and they proved its complete integrability. When $C'_5 = C'_6 = 0$, an explicit form of P_1, \dots, P_n is given by [De].

Moreover if $\omega_1 = \omega_2 = \infty$, then (1.10) is reduced to

$$(1.14) \quad \begin{cases} u(t) = C'_1 t^{-2} + C'_2, \\ v(t) = C'_3 t^{-2} + C'_4 t^2 + C'_5 t^4 + C'_6 t^6 + C'_7. \end{cases}$$

Here we quote a result in [OS] for the operator which commutes with the Schrödinger operator P :

If there exists a nonzero constant ω such that the W -invariant differential operators P_1, \dots, P_n are invariant by the parallel translation $x_1 \mapsto x_1 + \omega$, then any W -invariant differential operator Q that is also invariant by the same parallel translation is contained in $\mathbb{C}[P_1, \dots, P_n]$ if $[P, Q] = 0$.

After this paper [Os] was written, [Ch] proved the completely integrability of the Schrödinger operator (1.1) with the elliptic potential function attached to the root system. If the root system is of type B_n in our situation, the potential considered in [Ch] corresponds to the case where $v(t) = C_3\wp(t) + C_4$ or $v(t) = C_3\wp(2t) + C_4$ in (1.10). The method is quite interesting but different from this note constructing explicitly all the integrals.

Lastly we give a brief overview of the following sections.

In §2 preliminary remarks are made and two results employed throughout are established.

In §3 the two fundamental operators Δ and Δ_n for A_{n-1} and D_n are introduced and their commutativity is proved by the results in §2. An expansion of Δ_n gives the commuting differential operators for A_{n-1} .

In §4 the Schrödinger operator is allowed to have a term in the potential only depending on the particle position through a given function v . A functional differential equation (4.4) is established that will ensure the commutativity of the fundamental operators P and P_n for B_n given by (4.2).

In §5, using the lemmas in §2, we establish solutions of the functional differential equation with the assumption $u = w$ which corresponds to the form (1.9).

In §6 we look at various rational and trigonometric degenerations of the solutions of the functional differential equation.

In §7 we bring the results of the previous sections together and establish the commuting differential operators P_1, \dots, P_n for B_n and D_n .

2. Preliminaries.

First we introduce some notation used in this paper. For an element w of the permutation group \mathfrak{S}_n of the set of indices $\{1, \dots, n\}$, we define $w(i) = i$ for any $i \in \mathbb{Z}$ satisfying $i < 1$ or $i > n$ and we identify \mathfrak{S}_n with a subgroup of the group of bijective transformations of \mathbb{Z} . Then we have naturally $\mathfrak{S}_k \subset \mathfrak{S}_n$ if $k < n$.

When we distinguish the Weyl group that we are looking at, we denote it by $W(A_{n-1})$, $W(B_n)$ or $W(D_n)$ according to its type. Then $W(A_{n-1}) \simeq \mathfrak{S}_n$ and $W(A_{n-1}) \subset W(D_n) \subset W(B_n)$. We define a homomorphism ε of $W(B_n)$ to $\{\pm 1\}$ by

$$(2.1) \quad \varepsilon(w) = \begin{cases} 1 & \text{if } w \in W(D_n), \\ -1 & \text{if } w \notin W(D_n). \end{cases}$$

For the coordinate system (x_1, \dots, x_n) of \mathbb{R}^n we put

$$\partial_i = \frac{\partial}{\partial x_i}, \quad \partial^\alpha = \partial_1^{\alpha_1} \dots \partial_n^{\alpha_n} \quad \text{and} \quad |\alpha| = \alpha_1 + \dots + \alpha_n.$$

Here $\alpha = (\alpha_1, \dots, \alpha_n)$ with non-negative integers α_i .

Let $P = \sum p_\alpha(x) \partial^\alpha$ be a differential operator. Then we put

$$(2.2) \quad {}^tP = \sum (-1)^{|\alpha|} \partial^\alpha p_\alpha(x)$$

and we say that P is *self-adjoint* if ${}^tP = P$ and *skew self-adjoint* if ${}^tP = -P$. For $w \in W$ and a differential operator P , we denote by $w(P)$ the differential operator corresponding to P under the coordinate transformation w of \mathbb{R}^n . In particular we define $P^- = w^-(P)$ by $w^- \in W(B_n)$ with $w^-(x_1, \dots, x_n) = (-x_1, \dots, -x_n)$ and we call P has an *even parity* if $P^- = P$ and an *odd parity* if $P^- = -P$. Then we note that

$${}^t({}^tP) = (P^-)^- = P, \quad {}^t(P^-) = ({}^tP)^-, \quad {}^t(PQ) = {}^tQ{}^tP, \quad (PQ)^- = P^-Q^-.$$

In general the suffix $\{1, \dots, k\}$ of a function or an operator (eg. $Q_{\{1, \dots, k\}}$) means that it is a function or an operator of the variables x_1, \dots, x_k invariant under $w \in \mathfrak{S}_k$. And for a function or an operator $Q_{\{1, \dots, k\}}$ and a subset I of $\{1, \dots, n\}$, we define $Q_I = w(Q_{\{1, \dots, k\}})$ if there exist $w \in W(A_{n-1}) \simeq \mathfrak{S}_n$ with $w(\{1, \dots, k\}) = I$.

Now we review the Weierstrass elliptic function (cf. [WW]), which is

$$(2.3) \quad \wp(z|2\omega_1, 2\omega_2) = \frac{1}{z^2} + \sum_{\omega \neq 0} \left(\frac{1}{(z - \omega)^2} - \frac{1}{\omega^2} \right),$$

where the sum ranges over all periods $\omega = 2m_1\omega_1 + 2m_2\omega_2$ ($m_1, m_2 \in \mathbb{Z}$) except 0. We define

$$(2.4) \quad \omega_3 = -(\omega_1 + \omega_2) \quad \text{and} \quad \omega_4 = 0.$$

The Weierstrass elliptic function $\wp(t)$ satisfies the differential equation

$$(2.5) \quad \begin{aligned} (\wp')^2 &= 4\wp^3 - g_2\wp - g_3 \\ &= 4(\wp - e_1)(\wp - e_2)(\wp - e_3). \end{aligned}$$

Here

$$(2.6) \quad \begin{aligned} g_2 &= 60 \sum_{\omega \neq 0} \omega^{-4}, \quad g_3 = 140 \sum_{\omega \neq 0} \omega^{-6}, \\ \wp(\omega_\nu) &= e_\nu \quad \text{for } \nu = 1, 2, 3, \\ e_1 + e_2 + e_3 &= 0, \quad e_1 e_2 + e_2 e_3 + e_3 e_1 = -\frac{g_2}{4}, \quad e_1 e_2 e_3 = \frac{g_3}{4}. \end{aligned}$$

Moreover we have important formulas

$$(2.7) \quad \begin{vmatrix} \wp(x) & \wp'(x) & 1 \\ \wp(y) & \wp'(y) & 1 \\ \wp(z) & \wp'(z) & 1 \end{vmatrix} = 0 \quad \text{if } x + y + z = 0,$$

$$(2.8) \quad \wp(z + \omega_i) = e_i + \frac{(e_i - e_j)(e_i - e_k)}{\wp(z) - e_i} \quad \text{if } \{i, j, k\} = \{1, 2, 3\}$$

and

$$(2.9) \quad \wp(2z) = \frac{(12\wp(z)^2 - g_2)^2}{16\wp'(z)^2} - 2\wp(z).$$

In this paper the periods are allowed to be infinity and hence g_1 and g_2 or e_1 and e_2 take any complex numbers. Then the condition

$$(2.10) \quad (e_1 - e_2)(e_2 - e_3)(e_3 - e_1) \neq 0$$

holds if and only if the both periods are finite. On the other hand, if $e_1 = e_2 = \frac{1}{3}\lambda^2$ and $e_3 = -\frac{2}{3}\lambda^2$ with $\lambda \in \mathbb{C}$, then

$$(2.11) \quad \wp(z|\sqrt{-1}\lambda^{-1}\pi, \infty) = \lambda^2 \sinh^{-2} \lambda z + \frac{1}{3}\lambda^2.$$

In particular, if $e_1 = e_2 = e_3 = 0$, we have

$$(2.12) \quad \wp(z|\infty, \infty) = z^{-2}.$$

We note that if (2.10) holds, the function $v(t)$ given by (1.10) is rewritten into

$$(2.13) \quad v(t) = C_5' + \sum_{j=1}^4 C_j' \wp(t + \omega_j)$$

with suitable complex numbers C_1', \dots, C_5' (cf. (2.5) and (2.8)). Moreover for any complex numbers C_1'', C_2'' and C_3'' , it follows from (2.9) that

$$(2.14) \quad v(t) = C_1'' \wp(t) + C_2'' \wp(2t) + C_3''$$

is a special case of (1.10) and the complete integrability of the corresponding Schrödinger operator was a question in [OP2].

Now we prepare

LEMMA 2.1. *Let $\tilde{v}_k(t)$, $\tilde{u}_{ij}(t)$ and $\tilde{w}_{ij}(t)$ be functions with a single variable for $1 \leq i < j \leq 3$ and $1 \leq k \leq 3$. Put $v_k = \tilde{v}_k(x_k)$, $v_k' = \tilde{v}_k'(x_k)$, $u_{ij} = \tilde{u}_{ij}(x_i - x_j)$, $u_{ij}' = \tilde{u}_{ij}'(x_i - x_j)$, $w_{ij} = \tilde{w}_{ij}(x_i + x_j)$ and $w_{ij}' = \tilde{w}_{ij}'(x_i + x_j)$ for $1 \leq i < j \leq 3$ and*

$1 \leq k \leq 3$. Then we have

$$(2.15) \quad \begin{aligned} & \partial_1((u_{12} - w_{12})(u_{13} - w_{13})) + \partial_2((u_{12} - w_{12})(u_{23} - w_{23})) + \partial_3((u_{13} - w_{13})(u_{23} - w_{23})) \\ &= \begin{vmatrix} u_{12} & u'_{12} & 1 \\ u_{23} & u'_{23} & 1 \\ u_{13} & -u'_{13} & 1 \end{vmatrix} + \begin{vmatrix} u_{12} & u'_{12} & 1 \\ w_{13} & -w'_{13} & 1 \\ w_{23} & w'_{23} & 1 \end{vmatrix} + \begin{vmatrix} u_{23} & u'_{23} & 1 \\ w_{12} & -w'_{12} & 1 \\ w_{13} & w'_{13} & 1 \end{vmatrix} + \begin{vmatrix} u_{13} & -u'_{13} & 1 \\ w_{23} & -w'_{23} & 1 \\ w_{12} & w'_{12} & 1 \end{vmatrix}, \end{aligned}$$

$$(2.16) \quad \begin{aligned} & \partial_1((u_{12} + w_{12})(u_{13} - w_{13})) + \partial_2((u_{12} + w_{12})(u_{23} - w_{23})) + \partial_3((u_{13} + w_{13})(u_{23} + w_{23})) \\ &= \begin{vmatrix} u_{12} & u'_{12} & 1 \\ u_{23} & u'_{23} & 1 \\ u_{13} & -u'_{13} & 1 \end{vmatrix} + \begin{vmatrix} u_{12} & u'_{12} & 1 \\ w_{13} & -w'_{13} & 1 \\ w_{23} & w'_{23} & 1 \end{vmatrix} - \begin{vmatrix} u_{23} & u'_{23} & 1 \\ w_{12} & -w'_{12} & 1 \\ w_{13} & w'_{13} & 1 \end{vmatrix} - \begin{vmatrix} u_{13} & -u'_{13} & 1 \\ w_{23} & -w'_{23} & 1 \\ w_{12} & w'_{12} & 1 \end{vmatrix} \end{aligned}$$

and

$$(2.17) \quad \begin{aligned} & v'_1(u_{12} - w_{12}) + 2v_1(u'_{12} - w'_{12}) + \partial_2((v_1 + v_2)(u_{12} + w_{12}) - 2v_1v_2) \\ &= \begin{vmatrix} v_1 & v'_1 & 1 \\ v_2 & -v'_2 & 1 \\ u_{12} & -u'_{12} & 1 \end{vmatrix} + \begin{vmatrix} v_1 & -v'_1 & 1 \\ v_2 & -v'_2 & 1 \\ w_{12} & w'_{12} & 1 \end{vmatrix}. \end{aligned}$$

Proof. If we note that $u'_{ij} = \partial_i u_{ij} = -\partial_j u_{ij}$ and $w'_{ij} = \partial_i w_{ij} = \partial_j w_{ij}$, then equalities (2.15), (2.16) and (2.17) are clear by direct calculations. \square

In the case when $u(t) = C_1\wp(t) + C_2$, the function $u(t)$ is even and satisfies

$$(2.18) \quad \begin{vmatrix} u(x) & u'(x) & 1 \\ u(y) & u'(y) & 1 \\ u(z) & u'(z) & 1 \end{vmatrix} = 0 \quad \text{for } x + y + z = 0,$$

which is clear from (2.7). Hence we have

COROLLARY 2.2. For given even functions $u(t)$, $v(t)$ and $w(t)$, put

$$(2.19) \quad \begin{aligned} & \phi_{ij} = u(x_i - x_j) + w(x_i + x_j), \\ & \psi_{ij} = u(x_i - x_j) - w(x_i + x_j), \\ & v_k = v(x_k). \end{aligned}$$

Then clearly

$$(2.20) \quad \phi_{ji} = \phi_{ij}, \quad \psi_{ji} = \psi_{ij} \quad \text{and} \quad \partial_i \phi_{ij} + \partial_j \psi_{ij} = 0.$$

i) If

$$(2.21) \quad u(t) = w(t) = C_1\wp(t) + C_2$$

or

$$(2.22) \quad u(t) = C_1\wp(t) + C_2 \quad \text{and} \quad w(t) = C_3,$$

then

$$(2.23) \quad \partial_i(\psi_{ij}\psi_{ik}) + \partial_j(\psi_{ij}\psi_{jk}) + \partial_k(\psi_{ik}\psi_{jk}) = 0$$

and

$$(2.24) \quad \partial_i(\phi_{ij}\psi_{ik}) + \partial_j(\phi_{ij}\psi_{jk}) + \partial_k(\phi_{ik}\phi_{jk}) = 0.$$

ii) If

$$(2.25) \quad u(t) = v(t) = w(t) = C_1 \wp(t) + C_2$$

or

$$(2.26) \quad u(t) = w(t) = C_1 \wp(t) + C_2 \quad \text{and} \quad v(t) = C_3,$$

then

$$(2.27) \quad (\partial_i v_i) \psi_{ij} + 2v_i (\partial_i \psi_{ij}) + \partial_j ((v_i + v_j) \phi_{ij} - 2v_i v_j) = 0.$$

Here the indices i, j and k are mutually different and C_1, C_2 and C_3 are any complex numbers.

Remark 2.3. The equation (2.18) for u and its generalization are studied by [BP], [BBy], [OS] in connection with integrable systems and equations similar to those in Corollary 2.2 are discussed in [BBu]

3. A fundamental integral of type A_{n-1} and D_n .

In this section we use the notation in Corollary 2.2. Put

$$(3.1) \quad \psi_{\{1, \dots, 2k\}} = \frac{1}{2^k k!} \sum_{w \in \mathfrak{S}_{2k}} w(\psi_{12} \psi_{34} \psi_{56} \cdots \psi_{2k-1, 2k}).$$

We sometimes denote by $\psi_{i,j}$ in place of ψ_{ij} to distinguish the suffices.

Define

$$(3.2) \quad \Delta = -\frac{1}{2} \sum_{j=1}^n \partial_j^2 + \sum_{1 \leq i < j \leq n} \phi_{ij}$$

and

$$(3.3) \quad \Delta_n = \sum_{0 \leq \nu \leq \lfloor \frac{n}{2} \rfloor} \frac{1}{(2\nu)!(n-2\nu)!} \sum_{w \in \mathfrak{S}_n} w(\psi_{\{1, \dots, 2\nu\}} \partial_{2\nu+1} \cdots \partial_n).$$

Let $\bar{\Delta}$ and $\bar{\Delta}_n$ be the functions of (x, ξ) obtained by replacing ∂_i by ξ_i for $i = 1, \dots, n$ in (3.2) and (3.3), respectively.

The Poisson bracket of functions $f(x, \xi)$ and $g(x, \xi)$ is defined by

$$(3.4) \quad \{f, g\} = \sum_{j=1}^n \left(\frac{\partial f}{\partial \xi_j} \frac{\partial g}{\partial x_j} - \frac{\partial g}{\partial \xi_j} \frac{\partial f}{\partial x_j} \right).$$

Then we have

PROPOSITION 3.1. *Suppose u and w are given by (2.21) or (2.22). Then*

$$[\Delta_n, \Delta] = \{\bar{\Delta}_n, \bar{\Delta}\} = 0.$$

Proof. Put $Q = [\Delta_n, \Delta]$ and suppose $Q \neq 0$. Since ${}^t \Delta_n = (-1)^n \Delta_n$ and ${}^t \Delta = \Delta$, ${}^t Q = {}^t [\Delta_n, \Delta] = -[{}^t \Delta_n, {}^t \Delta] = (-1)^{n+1} Q$. Hence the order of Q equals $n - 2m - 1$ with a suitable positive integer m . Then by using (2.20) and (2.23), the coefficient of $\partial_{2m+2} \cdots \partial_n$ in the \mathfrak{S}_n -invariant operator Q equals

$$\sum_{i=1}^{2m+1} \psi_{\{1, \dots, 2m+1\} \setminus \{i\}} \partial_i \sum_{\mu < \nu} \phi_{\mu\nu} + \sum_{j=2m+2}^n \partial_j \psi_{\{1, \dots, 2m+1, j\}}$$

$$\begin{aligned}
 &= \sum_{i=1}^{2m+1} \sum_{\substack{j \neq i \\ 1 \leq j \leq n}} \psi_{\{1, \dots, 2m+1\} \setminus \{i\}} \partial_i \phi_{ij} + \sum_{j=2m+2}^n \sum_{i=1}^{2m+1} \psi_{\{1, \dots, 2m+1\} \setminus \{i\}} \partial_j \psi_{ij} \\
 &= - \sum_{i=1}^{2m+1} \sum_{\substack{j \neq i \\ 1 \leq j \leq 2m+1}} \psi_{\{1, \dots, 2m+1\} \setminus \{i\}} \partial_j \psi_{ij} \\
 &= - \sum_{i=1}^{2m+1} \sum_{\substack{j \neq i \\ 1 \leq j \leq 2m+1}} \sum_{\substack{k \neq i, k \neq j \\ 1 \leq k \leq 2m+1}} \psi_{\{1, \dots, 2m+1\} \setminus \{i, j, k\}} \psi_{jk} \partial_j \psi_{ij} \\
 &= - \sum_{1 \leq i < j < k \leq 2m+1} \psi_{\{1, \dots, 2m+1\} \setminus \{i, j, k\}} (\partial_i(\psi_{ij} \psi_{ik}) + \partial_j(\psi_{ij} \psi_{jk}) + \partial_k(\psi_{ik} \psi_{jk})) \\
 &= 0,
 \end{aligned}$$

which contradicts to the fact that the order of Q equals $n - 2m - 1$. Thus we have $[\Delta_n, \Delta] = 0$ and by the same calculation we have also $\{\bar{\Delta}_n, \bar{\Delta}\} = 0$. \square

The following theorem is known but we repeat it here for the completeness.

THEOREM 3.2 (Type A_{n-1} . [OP2], [OS, Theorem 5.2 and Remark 5.3]). *Put*

$$u(t) = C_1 \wp(t) + C \quad \text{and} \quad w(t) = 0.$$

Regard Δ_n as a polynomial function of C , denote it by $P_n(C)$ and put $P_{n-1}(C) = [P_n(C), x_1 + \dots + x_n]$. Then

$$(3.5) \quad [P_n(C), P_n(C')] = [P_n(C), P_{n-1}(C')] = [P_{n-1}(C), P_{n-1}(C')] = 0$$

for any $C, C' \in \mathbb{C}$.

Defining P_k by $P_n(C) = \sum_{0 \leq \nu \leq [\frac{n}{2}]} P_{n-2\nu} C^\nu$ and $P_{n-1}(C) = \sum_{0 \leq \nu \leq [\frac{n-1}{2}]} (2\nu + 1) P_{n-2\nu-1} C^\nu$, we have

$$\begin{aligned}
 (3.6) \quad P_k &= \sum_{0 \leq j \leq [\frac{k}{2}]} \frac{C_1^j}{2^j j! (k - 2j)!} \sum_{w \in \mathfrak{S}_n} w(\wp(x_1 - x_2) \wp(x_3 - x_4) \cdots \\
 &\quad \cdot \wp(x_{2j-1} - x_{2j}) \partial_{2j+1} \cdots \partial_k)
 \end{aligned}$$

and P_1, \dots, P_n are the required operators for the Schrödinger operator (1.1) with (1.7) and (1.8) when W is of type A_{n-1} .

By replacing ∂_i and $[\ , \]$ by ξ_i and $\{ \ , \ \}$, respectively, we have the same claim for the corresponding Hamiltonian system.

Proof. Put $Q = [P_n(C), P_n(C')]$ and suppose $Q \neq 0$. Since ${}^t P_n(C) = P_n(C)^-$, we have $-{}^t Q = Q^-$. By Jacobi's identity for $[\ , \]$, we have $[Q, \Delta] = 0$, which implies that the coefficients of the terms of highest order in Q are polynomial functions of x (cf. [Be, Lemma 2.5] or [OS, Lemma 3.1 and Lemma 3.5]). Hence if ω_1 is finite, the coefficients are constant because of their periodicity. Moreover by the analytic continuation we can conclude that the coefficients are constant even if $\omega_1 = \infty$. This contradicts to $-{}^t Q = Q^- \neq 0$. Hence we have $[P_n(C), P_n(C')] = 0$.

Note that $[P_{n-1}(C), \Delta] = -[\partial_1 + \dots + \partial_n, P_n(C)] + [[P_n(C), \Delta], x_1 + \dots + x_n] = 0$. Hence the same argument as above shows (3.5) by replacing $(P_n(C), P_n(C'))$ by $(P_n(C), P_{n-1}(C'))$ or $(P_{n-1}(C), P_{n-1}(C'))$.

The remaining part of the theorem is clear from the definition of $P_n(C)$. \square

4. A functional differential equation. Retain the notation in Corollary 2.2 and the previous section and put

$$(4.1) \quad \Delta_{\{1, \dots, k\}} = \sum_{0 \leq \nu \leq \lfloor \frac{k}{2} \rfloor} \frac{1}{(2\nu)!(k-2\nu)!} \sum_{w \in \mathfrak{S}_k} w(\psi_{\{1, \dots, 2\nu\}} \partial_{2\nu+1} \cdots \partial_k)$$

for $k = 1, \dots, n$ (cf. (3.3)). Then we have easily

LEMMA 4.1.

$$\begin{aligned} [\Delta_{\{1, \dots, k\}}, x_k] &= \Delta_{\{1, \dots, k-1\}}, \\ \Delta_{\{1, \dots, k\}} &= \Delta_{\{1, \dots, k-1\}} \partial_k + \sum_{1 \leq \nu \leq k-1} \psi_{\nu k} \Delta_{\{1, \dots, k-1\} \setminus \{\nu\}}. \end{aligned}$$

Let $q_{\{1, \dots, k\}}$ be suitable symmetric functions of (x_1, \dots, x_k) for $k = 1, \dots, n$ and put $q_\emptyset = 1$. For even functions $v_j = v(x_j)$, we examine the condition such that the operators

$$(4.2) \quad \begin{aligned} P &= -\frac{1}{2} \sum_{j=1}^n \partial_j^2 + \sum_{1 \leq i < j \leq n} \phi_{ij} + \sum_{j=1}^n v_j, \\ P_n &= \sum_{k=0}^n \frac{1}{k!(n-k)!} \sum_{w \in \mathfrak{S}_n} w(q_{\{1, \dots, k\}} \Delta_{\{k+1, \dots, n\}}^2) \end{aligned}$$

satisfy $[P_n, P] = 0$. We denote by $\bar{\Delta}_{\{1, \dots, k\}}$ and \bar{P} and \bar{P}_n the functions of (x, ξ) obtained by replacing ∂_i by ξ_i in the above definition of the corresponding operators. We introduce symmetric functions $T_{\{1, \dots, k\}}$ of (x_1, \dots, x_k) such that

$$(4.3) \quad q_{\{1, \dots, k\}} = \sum_{I_1 \amalg \dots \amalg I_\nu = \{1, \dots, k\}} T_{I_1} \cdots T_{I_\nu},$$

where the sum runs over all different partitions of $\{1, \dots, k\}$. For example

$$\begin{aligned} q_\emptyset &= T_\emptyset = 1, \quad q_{\{1\}} = T_{\{1\}}, \quad q_{\{1,2\}} = T_{\{1\}}T_{\{2\}} + T_{\{1,2\}}, \\ q_{\{1,2,3\}} &= T_{\{1\}}T_{\{2\}}T_{\{3\}} + T_{\{1\}}T_{\{2,3\}} + T_{\{2\}}T_{\{3,1\}} + T_{\{3\}}T_{\{1,2\}} + T_{\{1,2,3\}}. \end{aligned}$$

THEOREM 4.2. *Retain the above notation. Suppose*

$$(4.4) \quad \begin{cases} T_{\{1\}} &= -2v_1, \\ \partial_k T_{\{1, \dots, k\}} &= \sum_{j=1}^{k-1} \left(2T_{\{1, \dots, k-1\}} (\partial_j \psi_{jk}) + (\partial_j T_{\{1, \dots, k-1\}}) \psi_{jk} \right) \\ &\text{for } k = 2, \dots, n. \end{cases}$$

Then $[P_n, P] = \{\bar{P}_n, \bar{P}\} = 0$.

Proof. It follows from Proposition 3.1 that

$$\begin{aligned} [P_n, P] &= \left[\sum_{k=0}^n \frac{1}{k!(n-k)!} \sum_{w \in \mathfrak{S}_n} w(q_{\{1, \dots, k\}} \Delta_{\{k+1, \dots, n\}}^2), P \right] \\ &= \sum_{k=0}^n \frac{1}{k!(n-k)!} \sum_{w \in \mathfrak{S}_n} w \left([q_{\{1, \dots, k\}} \Delta_{\{k+1, \dots, n\}}^2, P] \right) \end{aligned}$$

$$\begin{aligned}
 & -\frac{1}{2}\partial_1^2 - \dots - \frac{1}{2}\partial_k^2 + \sum_{\nu=k+1}^n \left(v_\nu + \sum_{\mu=1}^k \phi_{\mu\nu} \right) \Big] \\
 = & \sum_{k=0}^n \frac{1}{k!(n-k)!} \sum_{w \in \mathfrak{S}_n} w \left(q_{\{1, \dots, k\}} \left[\Delta_{\{k+1, \dots, n\}}^2, \sum_{\nu=k+1}^n \left(v_\nu + \sum_{\mu=1}^k \phi_{\mu\nu} \right) \right] \right. \\
 & \left. + \frac{1}{2} \left[\partial_1^2 + \dots + \partial_k^2, q_{\{1, \dots, k\}} \right] \Delta_{\{k+1, \dots, n\}}^2 \right).
 \end{aligned}$$

Hence by Lemma 4.1 and (2.20) we have

$$\begin{aligned}
 \{\bar{P}_n, \bar{P}\} &= \sum_{k=0}^{n-1} \frac{1}{k!(n-k-1)!} \sum_{w \in \mathfrak{S}_n} w \left(2q_{\{1, \dots, k\}} (v'_{k+1} + \sum_{j=1}^k \partial_{k+1} \phi_{j, k+1}) \right. \\
 & \quad \left. \cdot \bar{\Delta}_{\{k+2, \dots, n\}} \bar{\Delta}_{\{k+1, \dots, n\}} \right) \\
 & \quad + \sum_{k=1}^n \frac{1}{(k-1)!(n-k)!} \sum_{w \in \mathfrak{S}_n} w \left((\partial_k q_{\{1, \dots, k\}}) \bar{\Delta}_{\{k+1, \dots, n\}}^2 \xi^k \right) \\
 = & \sum_{k=1}^n \frac{1}{(k-1)!(n-k)!} \sum_{w \in \mathfrak{S}_n} w \left(2q_{\{1, \dots, k-1\}} \left(v'_k + \sum_{j=1}^{k-1} \partial_k \phi_{jk} \right) \right. \\
 & \quad \left. \cdot \bar{\Delta}_{\{k+1, \dots, n\}} \bar{\Delta}_{\{k, \dots, n\}} \right) \\
 & \quad + \sum_{k=1}^n \frac{1}{(k-1)!(n-k)!} \sum_{w \in \mathfrak{S}_n} w \left((\partial_k q_{\{1, \dots, k\}}) (\bar{\Delta}_{\{k, \dots, n\}} \right. \\
 & \quad \left. - \sum_{j=k+1}^n \psi_{kj} \bar{\Delta}_{\{k+1, \dots, n\} \setminus \{j\}}) \bar{\Delta}_{\{k+1, \dots, n\}} \right) \\
 = & \sum_{k=1}^n \frac{1}{(k-1)!(n-k)!} \sum_{w \in \mathfrak{S}_n} w \left(\left(2q_{\{1, \dots, k-1\}} v'_k \right. \right. \\
 & \quad \left. \left. - \sum_{j=1}^{k-1} (2q_{\{1, \dots, k-1\}} \partial_j \psi_{jk} + (\partial_j q_{\{1, \dots, k-1\}}) \psi_{jk}) \right. \right. \\
 & \quad \left. \left. + \partial_k q_{\{1, \dots, k\}} \right) \bar{\Delta}_{\{k+1, \dots, n\}} \bar{\Delta}_{\{k, \dots, n\}} \right).
 \end{aligned}$$

Here the last equality follows from

$$\begin{aligned}
 & \sum_{k=1}^n \frac{1}{(k-1)!(n-k)!} \sum_{w \in \mathfrak{S}_n} w \left((\partial_k q_{\{1, \dots, k\}}) \sum_{j=k+1}^n \psi_{kj} \bar{\Delta}_{\{k+1, \dots, n\} \setminus \{j\}} \bar{\Delta}_{\{k+1, \dots, n\}} \right) \\
 = & \sum_{\ell=1}^n \frac{1}{(\ell-1)!(n-\ell)!} \sum_{w \in \mathfrak{S}_n} w \left(\sum_{i=1}^{\ell-1} (\partial_i q_{\{1, \dots, \ell-1\}}) \psi_{i\ell} \bar{\Delta}_{\{\ell+1, \dots, n\}} \bar{\Delta}_{\{\ell, \dots, n\}} \right).
 \end{aligned}$$

Hence if $q_{\{1, \dots, k\}}$ satisfy

$$\begin{aligned}
 \partial_k q_{\{1, \dots, k\}} &= -2q_{\{1, \dots, k-1\}} v'_k \\
 (4.5) \quad & + \sum_{j=1}^{k-1} \left(2q_{\{1, \dots, k-1\}} (\partial_j \psi_{jk}) + (\partial_j q_{\{1, \dots, k-1\}}) \psi_{jk} \right)
 \end{aligned}$$

for $k = 1, \dots, n$, then $\{\bar{P}_n, \bar{P}\} = 0$. Under the assumption of the theorem, the right hand side of (4.5) equals

$$\begin{aligned} & \sum_{I_1 \amalg \dots \amalg I_\nu = \{1, \dots, k-1\}} \left(T_{I_1} \cdots T_{I_\nu} \partial_k T_{\{k\}} \right. \\ & \quad \left. + \sum_{j=1}^{k-1} \left(2T_{I_1} \cdots T_{I_\nu} (\partial_j \psi_{jk}) + (\partial_j (T_{I_1} \cdots T_{I_\nu})) \psi_{jk} \right) \right) \\ & = \sum_{I_1 \amalg \dots \amalg I_\nu = \{1, \dots, k-1\}} \partial_k \left(T_{I_1} \cdots T_{I_\nu} T_{\{k\}} + \sum_{\mu=1}^\nu T_{I_1} \cdots T_{I_\mu \cup \{k\}} \cdots T_{I_\nu} \right), \end{aligned}$$

which equals the left hand side of (4.5) and hence we have $\{\bar{P}_n, \bar{P}\} = 0$.

Thus we have

$$\begin{aligned} [P_n, P] &= \sum_{k=0}^n \frac{1}{k!(n-k)!} \sum_{w \in \mathfrak{S}_n} w \left(q_{\{1, \dots, k\}} \sum_{\nu=k+1}^n (\partial_\nu^2 v_\nu + \sum_{1 \leq \mu \leq k} \partial_\nu^2 \phi_{\mu\nu}) \right. \\ & \quad \left. + \Delta_{\{k+1, \dots, n\} \setminus \{\nu\}}^2 + \frac{1}{2} \sum_{\nu=1}^k (\partial_\nu^2 q_{\{1, \dots, k\}}) \Delta_{\{k+1, \dots, n\}}^2 \right) \end{aligned}$$

and therefore $[P_n, P]$ is clearly self-adjoint. Since P and P_n are self-adjoint, $[P_n, P]$ is skew self-adjoint. Hence we can conclude $[P_n, P] = 0$. \square

REMARK 4.3. Let $T_{\{1, \dots, k\}}^1$ and $T_{\{1, \dots, k\}}^2$ be solutions of (4.4) for $(u, v, w) = (f, g_1, h)$ and (f, g_2, h) , respectively. Then $C^{k-1} C' T_{\{1, \dots, k\}}^1 + C^{k-1} C'' T_{\{1, \dots, k\}}^2$ are solutions for $(u, v, w) = (Cf, C'g_1 + C''g_2, Ch)$. Here C, C' and C'' are any complex numbers and $k = 1, \dots, n$.

5. Solutions of the functional differential equation.

In this section we shall construct elliptic solutions of (4.4) in the case when

$$(5.1) \quad u(t) = w(t) = C\varphi(t) + C'$$

with $C, C' \in \mathbb{C}$ (cf. (2.19)).

Retain the notation in the previous section and assume (5.1).

LEMMA 5.1. Under the notation in Corollary 2.2, the functions

$$\begin{cases} \Phi_0 = 1, \\ \Phi_n = (-1)^n \sum_{w \in \mathfrak{S}_n} w(\phi_{01} \phi_{12} \cdots \phi_{n-1, n}) \quad \text{for } n \geq 1 \end{cases}$$

satisfy

$$(5.2) \quad \partial_n \Phi_n = \partial_0 (\Phi_{n-1} \psi_{0n}) + \sum_{j=1}^{n-1} \left(2\Phi_{n-1} (\partial_j \psi_{jn}) + (\partial_j \Phi_{n-1}) \psi_{jn} \right).$$

Proof. For $j = 1, \dots, n - 1$

$$\begin{aligned}
 & (-1)^{n-1} \left(2\Phi_{n-1}(\partial_j \psi_{jn}) + (\partial_j \Phi_{n-1})\psi_{jn} \right) \\
 &= \sum_{\substack{w \in \mathfrak{S}_{n-1} \\ w(n-1)=j}} (\partial_j \psi_{jn}) w(\phi_{01} \phi_{12} \cdots \phi_{n-2, n-1}) \\
 &+ \sum_{\substack{w \in \mathfrak{S}_{n-1} \\ w(n-1)=j}} (\partial_j(\phi_{w(n-2)j} \psi_{jn})) w(\phi_{01} \phi_{12} \cdots \phi_{n-3, n-2}) \\
 &+ \sum_{\substack{w \in \mathfrak{S}_{n-1} \\ w(n-1) \neq j}} (\partial_j(\phi_{w(w^{-1}(j)-1)j} \psi_{jn})) \prod_{\substack{i \neq w(j) \\ 1 \leq i \leq n-1}} w(\phi_{i-1, i}) \\
 &+ \sum_{\substack{w \in \mathfrak{S}_{n-1} \\ w(n-1) \neq j}} (\partial_j(\phi_{jw(w^{-1}(j)+1)} \psi_{jn})) \prod_{\substack{i \neq w(j) \\ 1 \leq i \leq n-1}} w(\phi_{i, i+1}).
 \end{aligned}$$

Hence it follows from (2.20) and (2.24) that the right hand side of (5.2) equals

$$\begin{aligned}
 & (-1)^n \left(\sum_{j=1}^{n-1} \sum_{\substack{w \in \mathfrak{S}_{n-1} \\ w(n-1)=j}} \partial_n \left(\phi_{jn} \prod_{i=1}^{n-1} w(\phi_{i-1, i}) \right) \right. \\
 & \left. - \sum_{w \in \mathfrak{S}_{n-1}} \sum_{k=1}^{n-1} w \left(\partial_k(\phi_{k-1, k} \psi_{k, n}) + \partial_{k-1}(\phi_{k-1, k} \psi_{k-1, n}) \right) \prod_{\substack{i \neq k \\ 1 \leq i \leq n-1}} w(\phi_{i-1, i}) \right) \\
 &= (-1)^n \left(\sum_{\substack{w \in \mathfrak{S}_n \\ w(n)=n}} \partial_n \left(\prod_{i=1}^n w(\phi_{i-1, i}) \right) \right. \\
 & \left. + \sum_{w \in \mathfrak{S}_{n-1}} \sum_{k=1}^{n-1} w \left(\partial_n(\phi_{k-1, n} \phi_{n, k}) \prod_{\substack{i \neq k \\ 1 \leq i \leq n-1}} \phi_{i-1, i} \right) \right) \\
 &= \partial_n \Phi_n.
 \end{aligned}$$

Thus we have the lemma. \square

LEMMA 5.2. *Suppose there exist a symmetric function $g(s, t)$ of (s, t) such that*

$$(5.3) \quad 2v_1(\partial_1 \psi_{12}) + (\partial_1 v_1)\psi_{12} = \partial_2(2g_{12} - (v_1 + v_2)\phi_{12})$$

by denoting $g_{ij} = g(x_i, x_j)$. Put

$$(5.4) \quad \begin{aligned}
 & S_{\{1\}}^o = -2v_1, \\
 & S_{\{1, \dots, k\}}^o = 2(-1)^k \sum_{w \in \mathfrak{S}_k} w(v_1 \phi_{12} \phi_{23} \cdots \phi_{k-1, k}) \quad \text{for } k \geq 1.
 \end{aligned}$$

Then

$$(5.5) \quad \begin{aligned} \partial_k (S_{\{1, \dots, k\}}^o - (-1)^k \sum_{w \in \mathfrak{S}_{k-1}} 4w(g_{1k}\phi_{12}\phi_{23} \cdots \phi_{k-2, k-1})) \\ = \sum_{j=1}^{k-1} \left(2S_{\{1, \dots, k-1\}}^o (\partial_j \psi_{jk}) + (\partial_j S_{\{1, \dots, k-1\}}^o) \psi_{jk} \right). \end{aligned}$$

i) If

$$(5.6) \quad 2v_1(\partial_1 \psi_{12}) + (\partial_1 v_1) \psi_{12} = \partial_2(2\lambda v_1 v_2 - (v_1 + v_2)\phi_{12})$$

with a complex number λ , then

$$(5.7) \quad \begin{cases} S_{\{1\}} & = -2v_1, \\ \partial_k S_{\{1, \dots, k\}} & = \sum_{j=1}^{k-1} \left(2S_{\{1, \dots, k-1\}}^o (\partial_j \psi_{jk}) + (\partial_j S_{\{1, \dots, k-1\}}^o) \psi_{jk} \right) \end{cases}$$

by putting

$$(5.8) \quad S_{\{1, \dots, k\}} = \sum_{I_1 \amalg \dots \amalg I_\nu = \{1, \dots, k\}} (-\lambda)^{\nu-1} (\nu-1)! S_{I_1}^o \cdots S_{I_\nu}^o \quad \text{for } k \geq 1.$$

ii) If there exist even functions $f(t)$ and $h(t)$ and complex numbers λ, λ' and λ'' such that

$$(5.9) \quad \begin{cases} 2v_1(\partial_1 \psi_{12}) + (\partial_1 v_1) \psi_{12} = \partial_2(2\lambda' f_1 f_2 + 2\lambda''(h_1 + h_2) - (v_1 + v_2)\phi_{12}), \\ 2f_1(\partial_1 \psi_{12}) + (\partial_1 f_1) \psi_{12} = \partial_2(-(f_1 + f_2)\phi_{12}), \\ 2h_1(\partial_1 \psi_{12}) + (\partial_1 h_1) \psi_{12} = \partial_2(2\lambda(f_1 + f_2) - (h_1 + h_2)\phi_{12}), \end{cases}$$

the following functions $S_{\{1, \dots, k\}}$ satisfy (5.7).

$$\begin{aligned} S_{\{1, \dots, k\}} &= S_{\{1, \dots, k\}}^o - \sum_{I_1 \amalg I_2 = \{1, \dots, k\}} (\lambda' S_{I_1}' S_{I_2}' + \lambda'' S_{I_1}'' D_{I_2} + \lambda'' D_{I_1} S_{I_2}'') \\ &\quad + \sum_{I_1 \amalg I_2 \amalg I_3 = \{1, \dots, k\}} 2\lambda \lambda'' (S_{I_1}' D_{I_2} D_{I_3} + D_{I_1} S_{I_2}' D_{I_3} + D_{I_1} D_{I_2} S_{I_3}'), \\ S_{\{1, \dots, k\}}' &= 2(-1)^k \sum_{w \in \mathfrak{S}_k} w(f_1 \phi_{12} \phi_{23} \cdots \phi_{k-1, k}), \\ S_{\{1, \dots, k\}}'' &= 2(-1)^k \sum_{w \in \mathfrak{S}_k} w(h_1 \phi_{12} \phi_{23} \cdots \phi_{k-1, k}), \\ D_{\{1, \dots, k\}} &= 2(-1)^k \sum_{w \in \mathfrak{S}_k} w(\phi_{12} \phi_{23} \cdots \phi_{k-1, k}). \end{aligned}$$

Here we put $f_j = f(x_j)$ and $h_j = h(x_j)$ for $j \geq 1$ and $S_{\{1\}}' = -2f_1, S_{\{1\}}'' = -2h_1$ and $D_{\{1\}} = -2$.

Proof. Owing to (5.3) and (2.20) and Lemma 5.1, the right hand side of (5.5) equals

$$2(-1)^{k-1} \sum_{w \in \mathfrak{S}_{k-1}} w \left(2 \sum_{j=1}^{k-1} v_1 \phi_{12} \cdots \phi_{k-2, k-1} (\partial_j \psi_{jk}) \right)$$

$$\begin{aligned}
 & + (\partial_1 v_1) \phi_{12} \cdots \phi_{k-2,k-1} \psi_{1k} + \sum_{j=1}^{k-1} v_1 (\partial_j (\phi_{12} \cdots \phi_{k-2,k-1})) \psi_{jk} \\
 = & 2(-1)^k \sum_{w \in \mathfrak{S}_{k-1}} w \left(\partial_k (v_k \phi_{k1} \phi_{12} \cdots \phi_{k-2,k-1} - 2g_{1k} \phi_{12} \cdots \phi_{k-2,k-1}) \right. \\
 & - v_1 \phi_{12} \cdots \phi_{k-2,k-1} (\partial_1 \psi_{1k}) - 2 \sum_{j=2}^{k-1} v_1 \phi_{12} \cdots \phi_{k-2,k-1} (\partial_j \psi_{jk}) \\
 & \left. - \sum_{j=1}^{k-1} v_1 (\partial_j (\phi_{12} \cdots \phi_{k-2,k-1})) \psi_{jk} \right) \\
 = & 2(-1)^k \sum_{w \in \mathfrak{S}_{k-1}} w \left(\partial_k (v_k \phi_{k1} \phi_{12} \cdots \phi_{k-2,k-1} - 2g_{1k} \phi_{12} \cdots \phi_{k-2,k-1} \right. \\
 & \left. + \frac{1}{(k-1)!} \sum_{\substack{w' \in \mathfrak{S}_k \\ w'(1)=1}} w' (v_1 \phi_{12} \phi_{23} \cdots \phi_{k-1,k}) \right) \\
 = & \partial_k (S_{\{1, \dots, k\}}^{\circ} - (-1)^k \sum_{w \in \mathfrak{S}_{k-1}} 4w (g_{1k} \phi_{12} \phi_{23} \cdots \phi_{k-2,k-1})).
 \end{aligned}$$

Hence we have (5.5) and if (5.6) holds, we have

$$(-1)^k \sum_{w \in \mathfrak{S}_{k-1}} 4w (g_{1k} \phi_{12} \phi_{23} \cdots \phi_{k-2,k-1}) = \lambda S_{\{1, \dots, k-1\}}^{\circ} S_{\{k\}}^{\circ}$$

and therefore the right hand side of the second equation of (5.7) equals

$$\begin{aligned}
 & \sum_{j=1}^{k-1} \sum_{I_1 \amalg \cdots \amalg I_\nu = \{1, \dots, k-1\}} (-\lambda)^{\nu-1} (\nu-1)! \left(2S_{I_1}^{\circ} \cdots S_{I_\nu}^{\circ} (\partial_j \psi_{jk}) \right. \\
 & \left. + (\partial_j S_{I_1}^{\circ} \cdots S_{I_\nu}^{\circ}) \psi_{jk} \right) \\
 = & \sum_{I_1 \amalg \cdots \amalg I_\nu = \{1, \dots, k-1\}} (-\lambda)^{\nu-1} (\nu-1)! \sum_{\mu=1}^{\nu} \partial_k \left(S_{I_1}^{\circ} \cdots S_{I_\mu \cup \{k\}}^{\circ} \cdots S_{I_\nu}^{\circ} \right. \\
 & \left. - \lambda S_{I_1}^{\circ} \cdots S_{I_\mu}^{\circ} \cdots S_{I_\nu}^{\circ} S_{\{k\}}^{\circ} \right) \\
 = & \partial_k \left(\sum_{I_1 \amalg \cdots \amalg I_\nu = \{1, \dots, k\}} (-\lambda)^{\nu-1} (\nu-1)! S_{I_1}^{\circ} \cdots S_{I_\nu}^{\circ} \right) \\
 = & \partial_k S_{\{1, \dots, k\}}.
 \end{aligned}$$

Now suppose (5.9). Then by denoting

$$\begin{aligned}
 S_{\{1, \dots, k\}}^1 & = S_{\{1, \dots, k\}}^{\circ}, \\
 S_{\{1, \dots, k\}}^2 & = \sum_{I_1 \amalg I_2 = \{1, \dots, k\}} S'_{I_1} S'_{I_2}, \\
 S_{\{1, \dots, k\}}^3 & = \sum_{I_1 \amalg I_2 = \{1, \dots, k\}} (S''_{I_1} D_{I_2} + D_{I_1} S''_{I_2}),
 \end{aligned}$$

$$S^4_{\{1,\dots,k\}} = \sum_{I_1 \amalg I_2 \amalg I_3 = \{1,\dots,k\}} (S'_{I_1} D_{I_2} D_{I_3} + D_{I_1} S'_{I_2} D_{I_3} + D_{I_1} D_{I_2} S'_{I_3})$$

and

$$F_k^\nu = (-1)^k \left(\partial_k S^\nu_{\{1,\dots,k\}} - \sum_{j=1}^{k-1} (2S^\nu_{\{1,\dots,k-1\}} \partial_j \psi_{jk} + (\partial_j S^\nu_{\{1,\dots,k-1\}}) \psi_{jk}) \right)$$

for $\nu = 1, \dots, 4$, it follows from (5.5) that

$$\begin{aligned} F_k^1 &= \partial_k (\lambda' S'_{\{1,\dots,k-1\}} S'_{\{k\}} + \lambda'' S''_{\{1,\dots,k-1\}} D_{\{k\}} + \lambda'' D_{\{1,\dots,k-1\}} S''_{\{k\}}), \\ F_k^2 &= \partial_k (S'_{\{1,\dots,k-1\}} S'_{\{k\}}), \\ F_k^3 &= \partial_k \left(S''_{\{1,\dots,k-1\}} D_{\{k\}} + D_{\{1,\dots,k-1\}} S''_{\{k\}} \right. \\ &\quad \left. + \lambda \sum_{I_1 \amalg I_2 = \{1,\dots,k-1\}} ((S'_{I_1} D_{\{k\}} + D_{I_1} S'_{\{k\}}) D_{I_2} + D_{I_1} (S'_{I_2} D_{\{k\}} + D_{I_2} S'_{\{k\}})) \right), \\ F_k^4 &= \partial_k \left(\sum_{I_1 \amalg I_2 = \{1,\dots,k-1\}} (S'_{I_1} D_{I_2} D_{\{k\}} + D_{I_1} S'_{I_2} D_{\{k\}} + D_{I_1} D_{I_2} S'_{\{k\}}) \right). \end{aligned}$$

Since $\partial_k ((S'_{I_1} D_{I_2} + D_{I_1} S'_{I_2}) D_{\{k\}}) = 0$ in the above, we have

$$F_k^1 - \lambda' F_k^2 - \lambda'' F_k^3 + 2\lambda \lambda'' F_k^4 = 0,$$

which implies (5.7). Thus we have completed the proof of the lemma. \square

DEFINITION 5.3. For given even functions f and g of t , we define

$$\begin{aligned} \Phi_{\{1,\dots,k\}}(f, g) &= \sum_{w \in W(B_k)} w(f(x_1)g(x_1 - x_2)g(x_2 - x_3) \cdots g(x_{k-1} - x_k)), \\ \Theta_{\{1,\dots,k\}}(f, g) &= \sum_{I_1 \amalg \cdots \amalg I_\nu = \{1,\dots,k\}} (-1)^{\nu-1} (\nu - 1)! \Phi_{I_1}(f, g) \cdots \Phi_{I_\nu}(f, g) \end{aligned}$$

for $k \geq 1$. Here we note that $\Theta_{\{1\}}(f, g) = \Phi_{\{1\}}(f, g) = 2f(x_1)$ and $\Theta_\emptyset(f, g) = \Phi_\emptyset(f, g) = 0$.

PROPOSITION 5.4. *Suppose*

$$u(t) = w(t) = C_5 \wp(t), \quad v(t) = \sum_{j=1}^4 C_j \wp(t + \omega_j) - \frac{C_0}{2}$$

with $C_0, \dots, C_5 \in \mathbb{C}$. Then (4.4) holds by putting

$$T_{\{1,\dots,k\}} = (-C_5)^{k-1} \left(\frac{C_0}{2} \Theta_{\{1,\dots,k\}}(1, \wp(t)) - \sum_{j=1}^4 C_j \Theta_{\{1,\dots,k\}}(\wp(t + \omega_j), \wp(t)) \right).$$

Proof. Suppose $C_5 = 1$. If $v(t) = \wp(t + \omega_\nu)$ with $\nu = 1, \dots, 4$, the assumption (5.3) in Lemma 5.2 holds with $g_{12} = v_1 v_2$. In fact (2.27) means (5.3) if $v(t) = \wp(t)$ and then the coordinate transformation $x_j \mapsto x_j + \omega_\nu$ for $j = 1, \dots, n$ implies the case when $v(t) = \wp(t + \omega_\nu)$. If v is constant, then (5.3) is also valid with $g_{12} = v_1 v_2$. Hence the proposition follows from Lemma 5.2 and Remark 4.3. \square

Remark 5.5. Since we may put $g_{12} = 0$ in the above proof when v is constant, we may replace $\Theta_{\{1,\dots,k\}}(1, \wp(t))$ by $\Phi_{\{1,\dots,k\}}(1, \wp(t))$ in Proposition 5.4.

6. Degenerate solutions of the functional differential equation.

We give trigonometric and rational solutions of (4.4):

PROPOSITION 6.1. *For complex numbers λ, C_0, \dots, C_5 with $\lambda \neq 0$, put*

$$\begin{aligned} u(t) &= w(t) = C_5 \sinh^{-2} \lambda t, \\ v(t) &= C_1 \sinh^{-2} \lambda t + C_2 \cosh^{-2} \lambda t + C_3 \sinh^2 \lambda t + \frac{C_4}{4} \sinh^2 2\lambda t - \frac{C_0}{2}, \\ g(s, t) &= C_5 \left(C_1 \sinh^{-2} \lambda s \cdot \sinh^{-2} \lambda t - C_2 \cosh^{-2} \lambda s \cdot \cosh^{-2} \lambda t \right. \\ &\quad \left. + C_4 (\sinh^2 \lambda s + \sinh^2 \lambda t + 2 \sinh^2 \lambda s \cdot \sinh^2 \lambda t) \right). \end{aligned}$$

Then (5.3) holds. Moreover we have (4.4) with

$$\begin{aligned} T_I &= (-C_5)^{\#I-1} \left(\frac{C_0}{2} T_I^\circ(1) - C_1 T_I^\circ(\sinh^{-2} \lambda t) - C_2 T_I^\circ(\cosh^{-2} \lambda t) \right. \\ &\quad \left. - C_3 T_I^\circ(\sinh^2 \lambda t) - C_4 T_I^\circ\left(\frac{1}{4} \sinh^2 2\lambda t\right) \right) \end{aligned}$$

by putting

$$\begin{aligned} T_I^\circ(1) &= \Phi_I(1, \rho), \\ T_I^\circ(\sinh^{-2} \lambda t) &= \Theta_I(\sinh^{-2} \lambda t, \rho), \\ T_I^\circ(\cosh^{-2} \lambda t) &= -\Theta_I(-\cosh^{-2} \lambda t, \rho), \\ T_I^\circ(\sinh^2 \lambda t) &= \Phi_I(\sinh^2 \lambda t, \rho), \\ T_I^\circ\left(\frac{1}{4} \sinh^2 2\lambda t\right) &= \Phi_I\left(\frac{1}{4} \sinh^2 2\lambda t, \rho\right) - \sum_{I_1 \amalg I_2 = I} \left(2\Phi_{I_1}(\sinh^2 \lambda t, \rho) \cdot \Phi_{I_2}(\sinh^2 \lambda t, \rho) \right. \\ &\quad \left. + \Phi_{I_1}(\sinh^2 \lambda t, \rho) \cdot \Phi_{I_2}(1, \rho) + \Phi_{I_1}(1, \rho) \cdot \Phi_{I_2}(\sinh^2 \lambda t, \rho) \right), \end{aligned}$$

where $I \subset \{1, \dots, n\}$, $\rho = \sinh^{-2} \lambda t$ and the last sum runs over different partitions.

Proof. We can prove (5.3) by direct calculations but here we do it in a different way. First note that we may assume that C_5 and one of the numbers C_1, \dots, C_4 equal 1 and that the other 4 numbers are 0. Also by a simple change of coordinates we may assume $\lambda = 1$.

Now put $u(t) = \wp(t) - e_1$. Then if $v(t) = \wp(t + \omega_j) - e_1$ we have (5.3) with $g_{12} = v_1 v_2$ for $j = 1, \dots, 4$. If $e_1 = e_2 = \frac{1}{3}$ and $e_3 = -\frac{2}{3}$, then $\wp(t) - e_1 = \sinh^{-2} t$ and

$$\begin{aligned} \wp(t + \omega_3) - e_1 &= e_3 - e_1 + \frac{(e_3 - e_1)(e_3 - e_2)}{\wp(t) - e_3} \\ &= -1 + \frac{1}{\sinh^{-2} t + 1} = -\cosh^{-2} t. \end{aligned}$$

Hence if $u(t) = \sinh^{-2} t$ and $v(t) = \sinh^{-2} t$ or $-\cosh^{-2} t$, we have (5.3) with $g_{12} = v_1 v_2$.

Put $e_1 = \frac{1}{3}$, $e_2 = \frac{1}{3} - \varepsilon$, $e_3 = -\frac{2}{3} + \varepsilon$ with $0 < |\varepsilon| \ll 1$. Then it follows from (2.8) that

$$\wp(t + \omega_1) - e_1 = \frac{(e_1 - e_2)(e_1 - e_3)}{\wp(t) - e_1} = \varepsilon \sinh^2 t + o(\varepsilon).$$

Hence putting $v(t) = \wp(t + \omega_1) - e_1$, the coefficients of ε in (5.3) proves (5.3) for $(u, v) = (\sinh^{-2} t, \sinh^2 t)$ with $g(s, t) = 0$.

Next suppose $v(t) = (\wp(t + \omega_1) - e_1) + (\wp(t + \omega_2) - e_1) + (e_1 - e_2)$. Then

$$\begin{aligned} v(t) &= \varepsilon \left(\frac{e_1 - e_3}{\wp(t) - e_1} - \frac{e_2 - e_3}{\wp(t) - e_2} \right) = \varepsilon^2 \frac{\wp(t) - e_3}{(\wp(t) - e_1)(\wp(t) - e_2)} \\ &= \varepsilon^2 \frac{\sinh^{-2} t + 1}{\sinh^{-4} t} + o(\varepsilon^2) = \frac{\varepsilon^2}{4} \sinh^2 2t + o(\varepsilon^2) \end{aligned}$$

and (5.3) holds with

$$\begin{aligned} g(s, t) &= (\wp(s + \omega_1) - e_1)(\wp(t + \omega_1) - e_1) + (\wp(s + \omega_2) - e_1)(\wp(t + \omega_2) - e_1) \\ &= (\varepsilon \sinh^2 s)(\varepsilon \sinh^2 t) + (-\varepsilon \sinh^2 s - \varepsilon)(-\varepsilon \sinh^2 t - \varepsilon) + o(\varepsilon^2) \\ &= \varepsilon^2(1 + \sinh^2 s + \sinh^2 t + 2 \sinh^2 s \cdot \sinh^2 t) + o(\varepsilon^2). \end{aligned}$$

Hence we have (5.3) if

$$\begin{cases} g(s, t) = \sinh^2 s + \sinh^2 t + 2 \sinh^2 s \cdot \sinh^2 t, \\ (u, v) = (\sinh^{-2} t, \frac{1}{4} \sinh^2 2t) \end{cases}$$

The remaining part of the proposition is clear from Lemma 5.2 and Remark 4.3. We can also get it from Proposition 5.4 by considering the limit as above. \square

Remark 6.2. Since

$$\begin{aligned} \sinh^2 \lambda s + \sinh^2 \lambda t + 2 \sinh^2 \lambda s \cdot \sinh^2 \lambda t \\ = \sinh^2 \lambda s \cdot \sinh^2 \lambda t + \cosh^2 \lambda s \cdot \cosh^2 \lambda t - 1, \end{aligned}$$

we may put

$$\begin{aligned} T_I^\circ(\frac{1}{4} \sinh^2 2\lambda t) &= \Phi_I(\frac{1}{4} \sinh^2 2\lambda t, \rho) \\ &- \sum_{I_1 \amalg I_2 = I} \left(\Phi_{I_1}(\sinh^2 \lambda t, \rho) \cdot \Phi_{I_2}(\sinh^2 \lambda t, \rho) + \Phi_{I_1}(\cosh^2 \lambda t, \rho) \cdot \Phi_{I_2}(\cosh^2 \lambda t, \rho) \right) \end{aligned}$$

in Proposition 6.1.

PROPOSITION 6.3. *For complex numbers C_0, \dots, C_5 , put*

$$\begin{aligned} u(t) &= w(t) = C_5 t^{-2}, \\ v(t) &= C_1 t^{-2} + C_2 t^2 + C_3 t^4 + C_4 t^6 - \frac{C_0}{2}, \\ g(s, t) &= C_5 (C_1 s^{-2} t^{-2} + C_3 (s^2 + t^2) + C_4 (s^4 + t^4 + 3s^2 t^2)). \end{aligned}$$

Then (5.3) holds. Moreover we have (4.4) with

$$T_I = (-C_5)^{\#I-1} \left(\frac{C_0}{2} T_I^\circ(1) - C_1 T_I^\circ(t^{-2}) - C_2 T_I^\circ(t^2) - C_3 T_I^\circ(t^4) - C_4 T_I^\circ(t^6) \right)$$

by putting

$$\begin{aligned} T_I^\circ(1) &= \Phi_I(1, \rho), \\ T_I^\circ(t^{-2}) &= \Theta_I(t^{-2}, \rho), \\ T_I^\circ(t^2) &= \Phi_I(t^2, \rho), \\ T_I^\circ(t^4) &= \Phi_I(t^4, \rho) - \sum_{I_1 \amalg I_2 = \{1, \dots, k\}} (\Phi_{I_1}(t^2, \rho) \cdot \Phi_{I_2}(1, \rho) + \Phi_{I_1}(1, \rho) \cdot \Phi_{I_2}(t^2, \rho)), \end{aligned}$$

$$\begin{aligned}
 T_I^o(t^6) &= \Phi_I(t^6, \rho) - \sum_{I_1 \amalg I_2 = \{1, \dots, k\}} (3\Phi_{I_1}(t^2, \rho) \cdot \Phi_{I_2}(t^2, \rho) \\
 &\quad + \Phi_{I_1}(t^4, \rho) \cdot \Phi_{I_2}(1, \rho) + \Phi_{I_1}(1, \rho) \cdot \Phi_{I_2}(t^4, \rho)) \\
 &+ \sum_{I_1 \amalg I_2 \amalg I_3 = \{1, \dots, k\}} 6(\Phi_{I_1}(t^2, \rho) \cdot \Phi_{I_2}(1, \rho) \cdot \Phi_{I_3}(1, \rho) \\
 &\quad + \Phi_{I_1}(1, \rho) \cdot \Phi_{I_2}(t^2, \rho) \cdot \Phi_{I_3}(1, \rho) + \Phi_{I_1}(1, \rho) \cdot \Phi_{I_2}(1, \rho) \cdot \Phi_{I_3}(t^2, \rho)),
 \end{aligned}$$

where $I \subset \{1, \dots, n\}$, $\rho = t^{-2}$ and the sums run over different partitions.

Proof. Note that the proof proceeds in the same way as in the proof of Proposition 6.1. Put $u(t) = \lambda^2 \sinh^{-2} \lambda t$. Then for

$$\begin{cases} v(t) &= \lambda^2 \sinh^{-2} \lambda t, \\ g(s, t) &= \lambda^4 \sinh^{-2} \lambda s \cdot \sinh^{-2} \lambda t \end{cases}$$

or

$$\begin{cases} v(t) &= \lambda^{-2} \sinh^2 \lambda t, \\ g(s, t) &= 0 \end{cases}$$

or

$$\begin{cases} v(t) &= \lambda^{-4} (\frac{1}{4} \sinh^2 2\lambda t - \sinh^2 \lambda t), \\ g(s, t) &= \lambda^{-2} (\sinh^2 \lambda s + \sinh^2 \lambda t + 2 \sinh^2 \lambda s \cdot \sinh^2 \lambda t) \end{cases}$$

or

$$\begin{cases} v(t) &= \lambda^{-6} (1 - 2 \sinh^2 \lambda t + \frac{1}{4} \sinh^2 2\lambda t - \cosh^{-2} \lambda t), \\ g(s, t) &= \lambda^{-4} (\cosh^2 \lambda s \cdot \cosh^2 \lambda t + \sinh^2 \lambda s \cdot \sinh^2 \lambda t \\ &\quad + \cosh^{-2} \lambda s \cdot \cosh^{-2} \lambda t - 2), \end{cases}$$

we have (5.3). By the analytic continuation of these $u(t)$, $v(t)$ and $g(s, t)$ to $\lambda = 0$, we have (5.3) for $u(t) = t^{-2}$ and $v(t) = t^{-2}$ or t^2 or t^4 or t^6 with $g(s, t) = s^{-2}t^{-2}$ or 0 or $s^2 + t^2$ or $s^4 + t^4 + 3s^2t^2$, respectively. In fact, for example, we have

$$\begin{aligned}
 &\lambda^{-6} (1 - 2 \sinh^2 \lambda t + \frac{1}{4} \sinh^2 2\lambda t - \cosh^{-2} \lambda t) \\
 &= \lambda^{-6} (1 - \sinh^2 \lambda t + \sinh^4 \lambda t - (1 + \sinh^2 \lambda t)^{-1}) \\
 &= \lambda^{-6} \sinh^6 \lambda t + o(\lambda) = t^6 + o(\lambda), \\
 &\lambda^{-4} (\cosh^2 \lambda s \cdot \cosh^2 \lambda t + \sinh^2 \lambda s \cdot \sinh^2 \lambda t + \cosh^{-2} \lambda s \cdot \cosh^{-2} \lambda t - 2) \\
 &= \lambda^{-4} \left((1 + \sinh^2 \lambda s)(1 + \sinh^2 \lambda t) + (1 + \sinh^2 \lambda s)^{-1} (1 + \sinh^2 \lambda t)^{-1} \right. \\
 &\quad \left. + \sinh^2 \lambda s \cdot \sinh^2 \lambda t - 2 \right) \\
 &= \lambda^{-4} \left((\sinh^2 \lambda s + \sinh^2 \lambda t)^2 + \sinh^2 \lambda s \cdot \sinh^2 \lambda t \right) + o(\lambda) \\
 &= s^4 + t^4 + 3s^2t^2 + o(\lambda).
 \end{aligned}$$

The remaining part of the proposition is clear from Lemma 5.2 and Remark 4.3. We can also get it from Proposition 6.1 by taking the limit at $\lambda = 0$. □

7. Integrals of type B_n and D_n .

The argument in the preceding sections gives the integrals when W is of type B_n or D_n .

DEFINITION 7.1. For given even function $u(t)$ and symmetric functions $T_{\{1, \dots, k\}}$ of (x_1, \dots, x_k) for $k = 1, \dots, n$, define $W(B_n)$ -invariant differential operator

$$P(u, T) = \sum_{k=0}^n \frac{1}{k!(n-k)!} \sum_{w \in \mathfrak{S}_n} w \left(q_{\{1, \dots, k\}} \Delta_{\{k+1, \dots, n\}}^2 \right)$$

by

$$\begin{aligned} \Delta_{\{1, \dots, k\}} &= \sum_{0 \leq j \leq \lfloor \frac{k}{2} \rfloor} \frac{1}{2^k j! (k-2j)!} \sum_{w \in W(B_k)} \varepsilon(w) w \left(u(x_1 - x_2) u(x_3 - x_4) \cdots \right. \\ &\quad \left. \cdot u(x_{2j-1} - x_{2j}) \partial_{2j+1} \partial_{2j+2} \cdots \partial_k \right), \\ q_{\{1, \dots, k\}} &= \sum_{I_1 \amalg \dots \amalg I_\nu = \{1, \dots, k\}} T_{I_1} \cdots T_{I_\nu}, \end{aligned}$$

where

$$\begin{aligned} q_\emptyset &= 1, \quad q_{\{1\}} = T_{\{1\}}, \quad q_{\{12\}} = T_{\{1\}} T_{\{2\}} + T_{\{1,2\}}, \dots \\ T_{w(\{1, \dots, k\})} &= w(T_{\{1, \dots, k\}}), \quad \Delta_{w(\{1, \dots, k\})} = w(\Delta_{\{1, \dots, k\}}) \quad \text{for } w \in \mathfrak{S}_n. \end{aligned}$$

Replacing ∂_i by ξ_i for $i = 1, \dots, n$ in the definition of $\Delta_{\{1, \dots, k\}}$ and $P(u, T)$, we define functions $\bar{\Delta}_{\{1, \dots, k\}}$ and $\bar{P}(u, T)$ of (x, ξ) , respectively.

THEOREM 7.2 (Elliptic Potentials: Generic cases of Type B_n). Put

$$(7.1) \quad \begin{cases} u(t) = C_5 \wp(t), \\ v(t) = \sum_{j=1}^4 C_j \wp(t + \omega_j) - \frac{C_0}{2} \end{cases}$$

and define $P_n(C_0) = P(u, T)$ and $\bar{P}_n(C_0) = \bar{P}(u, T)$ by

$$\begin{aligned} T_{\{1, \dots, k\}} &= (-C_5)^{k-1} \left(\frac{C_0}{2} T_{\{1, \dots, k\}}^\circ(1) - \sum_{j=1}^4 C_j T_{\{1, \dots, k\}}^\circ(\wp(t + \omega_j)) \right), \\ T_{\{1, \dots, k\}}^\circ(\psi) &= \sum_{I_1 \amalg \dots \amalg I_\nu = \{1, \dots, k\}} (-1)^{\nu-1} (\nu-1)! S_{I_1}(\psi) \cdots S_{I_\nu}(\psi), \\ S_{\{1, \dots, k\}}(\psi) &= \sum_{w \in W(B_k)} w \left(\psi(x_1) \wp(x_1 - x_2) \wp(x_2 - x_3) \cdots \wp(x_{k-1} - x_k) \right). \end{aligned}$$

Then

$$(7.2) \quad [P_n(C), P_n(C')] = \{\bar{P}_n(C), \bar{P}_n(C')\} = 0$$

for $C, C' \in \mathbb{C}$.

Let P_j be the coefficient of C_0^{n-j} in $P_n(C_0)$. Then P_1, \dots, P_n are the required commuting differential operators (1.5) for the Schrödinger operator

$$(7.3) \quad P = -\frac{1}{2} \sum_{j=1}^n \frac{\partial^2}{\partial x_j^2} + \sum_{1 \leq i < j \leq n} (u(x_i - x_j) + u(x_i + x_j)) + \sum_{k=1}^n v(x_k)$$

in the case when W is of type B_n .

By using $\bar{P}_n(C_0)$ in place of $P_n(C_0)$, we have integrals \bar{P}_j of the Hamiltonian

$$(7.4) \quad \bar{P} = -\frac{1}{2} \sum_{j=1}^n \xi_j^2 + \sum_{1 \leq i < j \leq n} (u(x_i - x_j) + u(x_i + x_j)) + \sum_{k=1}^n v(x_k),$$

where $\bar{P}_1, \dots, \bar{P}_n$ are functionally independent and satisfy $\{\bar{P}_i, \bar{P}_j\} = 0$.

Proof. Theorem 4.2 and Proposition 5.4 imply

$$(7.5) \quad [P_n(C_0), P] = 0.$$

Fix $C, C' \in \mathbb{C}$ and put $Q = [P_n(C), P_n(C')]$. Then we have $[Q, P] = 0$ and $Q^- = -{}^tQ = Q$ and therefore we have $Q = 0$ as in the proof of Theorem 3.2.

Since $q_{\{1, \dots, k\}}$ is a monic polynomial of C_0 with degree k , it is clear that P_j for $j = 1, \dots, n$ satisfy (1.5). The remaining part of the theorem is also clear. \square

THEOREM 7.3 (Type D_n). *Suppose W is of type D_n . Then by putting $C_1 = C_2 = C_3 = C_4 = 0$, the operators P_1, \dots, P_{n-1} in Theorem 7.2 and $P_n = \Delta_{\{1, \dots, n\}}$ are the required commuting differential operators (1.6) for the Schrödinger operator*

$$(7.6) \quad P = -\frac{1}{2} \sum_{j=1}^n \frac{\partial^2}{\partial x_j^2} + \sum_{1 \leq i < j \leq n} (u(x_i - x_j) + u(x_i + x_j))$$

with the function $u(t)$ given by (7.1). Here the periods of $\wp(t)$ are allowed to be infinity.

Proof. Theorem 7.2 and Proposition 3.1 prove $[P_j, P] = 0$ for $j = 1, \dots, n$. Then the commutators $Q_j = [P_j, P_n]$ satisfy $Q_j^- = -{}^tQ_j = (-1)^n Q_j$ and $[Q_j, P] = 0$ and hence $Q_j = 0$ as in the proof of Theorem 3.2. \square

Remark 7.4. i) In Theorem 7.2 we have $P_n = P_n(0)$ and

$$(7.7) \quad P_{n-k} = \sum_{i=k}^n \sum_{j=i}^n \frac{1}{i!(j-i)!(n-j)!} \sum_{w \in \mathfrak{S}_n} \sum_{I_1 \amalg \dots \amalg I_k = \{1, \dots, i\}} w((-C_5)^{i-k} 2^{-k} T_{I_1}^o(1) \cdots T_{I_k}^o(1) q_{\{i+1, \dots, j\}} \Delta_{\{j+1, \dots, n\}}^2)$$

for $k = 1, \dots, n-1$, where $q_{\{i+1, \dots, j\}}$ are defined by putting $C_0 = 0$.

ii) Because of the uniqueness of $\mathbb{C}[P_1, \dots, P_n]$ in terms of (u, v) (cf. [OS, Theorem 6.5]), the existence of the commuting differential operators P_1, \dots, P_n for (1.10) which satisfy (1.5) is guaranteed by the analytic continuation of the parameters g_2 and g_3 of $\wp(t)$ even if ω_1 or ω_2 is infinite. We have explicitly given the analytic continuation. In fact Theorem 4.2, Proposition 6.1, Proposition 6.3 and the proof of Theorem 7.2 imply the following theorem.

THEOREM 7.5 (Degenerate cases of Type B_n). *Suppose*

i) *Trigonometric Potentials:*

$$(7.8) \quad \begin{cases} u(t) = C_5 \sinh^{-2} \lambda t, \\ v(t) = C_1 \sinh^{-2} \lambda t + C_2 \cosh^{-2} \lambda t + C_3 \sinh^2 \lambda t + \frac{C_4}{4} \sinh^2 2\lambda t - \frac{C_0}{2} \end{cases}$$

with a non-zero complex number λ or

ii) *Rational Potentials:*

$$(7.9) \quad \begin{cases} u(t) = C_5 t^{-2}, \\ v(t) = C_1 t^{-2} + C_2 t^2 + C_3 t^4 + C_4 t^6 - \frac{C_0}{2}. \end{cases}$$

Then for the function $T_{\{1, \dots, k\}}$ defined in Proposition 6.1 or Proposition 6.3, we have the same statements as in Theorem 7.2.

REFERENCES

- [Be] F. A. BEREZIN, *Laplace operators on semisimple Lie groups*, Proc. Mosc. Math. Soc., 6 (1971), pp. 371–463. (Russian)
- [BBy] H. W. BRADEN AND J. G. B. BYATT-SMITH, *On a functional differential equation of determinant type*, math.CA/9804082, preprint.
- [BBu] H. W. BRADEN AND V. M. BUCHSTABER, *Integrable systems with pairwise interactions and functional equations*, Reviews in Mathematics and Mathematical Physics, 10(2) (1997), pp. 121–166.
- [BP] V. M. BUCHSTABER AND A. M. PERELOMOV, *On the functional equation related to the quantum three-body problem*, Contemporary Mathematical Physics, Amer. Math. Soc. Transl. Ser. 2, 175 (1996), pp. 15–34.
- [Ca] F. CALOGERO, *Solution of the one dimensional n -body problem with quadratic and/or inverse quadratic pair potentials*, J. Math. Phys., 12 (1971), pp. 419–436.
- [Ch] I. CHEREDNIK, *Elliptic quantum many-body problem and double affine Knizhnik-Zamolodchikov equation*, Commun. Math. Phys., 169 (1995), pp. 441–461.
- [De] A. DEBIARD, *Système différentiel hypergéométrique et parties radiales des espaces symétriques de type BC_p* , Springer Lecture Notes in Math., 1296 (1988), pp. 42–124.
- [Et] P. I. ETINGOF, *Quantum integrable systems and representations of Lie algebras*, J. Math. Phys., 36 (1995), pp. 2636–2651.
- [He1] G. J. HECKMAN, *Root system and hypergeometric functions II*, Comp. Math., 64 (1987), pp. 353–373.
- [He2] ———, *An elementary approach to the hypergeometric shift operators of Opdam*, Invent. Math., 103 (1991), pp. 341–350.
- [HO] G. J. HECKMAN AND E. M. OPDAM, *Root system and hypergeometric functions I*, Comp. Math., 64 (1987), pp. 329–352.
- [In] V. I. INOZEMTSEV, *Lax representation with spectral parameter on a torus for Integrable particle systems*, Lett. Math. Phys., 17 (1989), pp. 11–17.
- [Oc] H. OCHIAI, *Commuting differential operators of rank two*, Indag. Math. (N.S.), 7 (1996), pp. 243–255.
- [OO] H. OCHIAI AND T. OSHIMA, *Commuting differential operators with B_2 symmetry*, UTMS 94–65, Dept. of Mathematical Sciences, Univ. of Tokyo, 1994, pp. 1–31, preprint.
- [OOS] H. OCHIAI, T. OSHIMA, AND H. SEKIGUCHI, *Commuting families of symmetric differential operators*, Proc. Japan Acad., 70A (1994), pp. 62–66.
- [OP1] M. A. OLSHANETSKY AND M. A. PERELOMOV, *Classical integrable finite dimensional systems related to Lie algebras*, Phys. Rep., 71 (1981), pp. 313–400.
- [OP2] ———, *Quantum integrable systems related to Lie algebras*, Phys. Rep., 94 (1983), pp. 313–404.
- [Op1] E. M. OPDAM, *Root system and hypergeometric functions III*, Comp. Math., 67 (1988), pp. 21–49.
- [Op2] ———, *Root system and hypergeometric functions IV*, Comp. Math., 67 (1988), pp. 191–209.
- [Os] T. OSHIMA, *Completely integrable systems with a symmetry in coordinates*, UTMS 94–6, Dept. of Mathematical Sciences, Univ. of Tokyo, 1994, pp. 1–22, preprint.
- [OS] T. OSHIMA AND H. SEKIGUCHI, *Commuting families of differential operators invariant under the action of a Weyl group*, J. Math. Sci. Univ. Tokyo, 2 (1995), pp. 1–75.
- [Sj] J. SEKIGUCHI, *Zonal spherical functions on some symmetric spaces*, Rubl. RIMS Kyoto Univ., 12 Suppl. (1977), pp. 455–459.
- [Su] B. SUTHERLAND, *Exact results for a quantum many-body problem in one dimension*, Phys. Rev., A5 (1972), pp. 1372–1376.
- [WW] E. T. WHITTAKER AND G. N. WATSON, *A Course of Modern Analysis, Fourth Edition*, Cambridge University Press, 1927.

