

EQUIVARIANT DE RHAM THEORY AND GRAPHS*

V. GUILLEMIN[†] AND C. ZARA[‡]

Abstract. Goresky, Kottwitz and MacPherson have recently shown that the computation of the equivariant cohomology ring of a G -manifold can be reduced to a computation in graph theory. This opens up the possibility that many of the fundamental theorems in equivariant de Rham theory may, on closer inspection, turn out simply to be theorems about graphs. In this paper we show that for some familiar theorems, this is indeed the case.

Introduction. This article will consist of two essentially disjoint parts. Part 1 is an exposition of (mostly) well-known results about G -manifolds. In section 1.1–1.3 we review the definition of the equivariant de Rham cohomology ring of a G -manifold and recall the statements of the two fundamental “localization theorems” in equivariant de Rham theory: the Atiyah-Bott-Berline-Vergne theorem and the Jeffrey-Kirwan theorem. In section 1.4 we discuss the “Smith” problem for G -manifolds (which is concerned with the question: Given a G -manifold with isolated fixed points, what kinds of representations can occur as isotropy representations at the fixed points?) Then in sections 1.5–1.7 we report on some very exciting recent results of Goresky-Kottwitz-MacPherson which have to do with the tie-in between “equivariant de Rham theory” and “graphs” alluded to in our title. These results show that for a large class of G -manifolds, M , with M^G finite, the equivariant cohomology ring of M is isomorphic to the equivariant cohomology ring of a pair (Γ, α) , where Γ is the intersection graph of a necklace of embedded S^2 's, each of which is equipped with a circle action (i.e., an axis of symmetry), and α is an “axial” function which describes the directions in which the axes of these S^2 's are tilted. Finally, in section 1.8 we discuss a Morse theoretic recipe for computing the Betti numbers of M in terms of the pair (Γ, α) .

The second part of this article is concerned with the combinatorial invariants of a pair (Γ, α) , Γ being *any* finite simple d -valent graph and α an abstract analogue of the axial function alluded to above. In particular, for such a pair we will prove combinatorial versions of the theorems described in sections 1.2-1.3 and 1.8. These combinatorial “localization” theorems help to shed some light on the role of the localization theorems in Smith theory: From the localization theorems one can generate a lot of complicated identities among the weights of the isotropy representations. However, the question of whether one can extract from these identities any new information about the isotropy representations themselves has been an open question for a long time. Our graph theoretical results seem to indicate that one can't.

This article is the first of a series of two articles on graphs and equivariant cohomology. In the second article in this series we will discuss K-theoretical analogues of the results above and give a purely combinatorial proof of the so-called “quantization commutes with reduction” conjecture.

1.1. Equivariant de Rham theory. Let G be an n -dimensional Lie group which is compact, connected and abelian, i.e., an n -dimensional torus. Let \mathfrak{g} be its

*Received November 11, 1998; accepted for publication November 30, 1998.

[†]Department of Mathematics, MIT, Cambridge, MA 02139 (vvg@math.mit.edu). Research Supported in part by NSF grant DMS 890771.

[‡]Department of Mathematics, MIT, Cambridge, MA 02139 (czara@math.mit.edu).

Lie algebra and \mathfrak{g}^* the vector space dual of \mathfrak{g} . We will fix a basis ξ^1, \dots, ξ^n of \mathfrak{g} and let x_1, \dots, x_n be the dual basis. Using this basis, the symmetric algebra $S(\mathfrak{g}^*)$ can be identified with the polynomial ring $\mathbb{C}[x_1, \dots, x_n]$.

Let M be a $2d$ -dimensional manifold and τ an action of G on M . From τ one gets an infinitesimal action, $\partial\tau$, of \mathfrak{g} on M which associates to every element ξ of \mathfrak{g} a vector field ξ_M . Let $\Omega(M)$ be the usual complex of de Rham forms on M and $\Omega(M)^G$ the subcomplex of G -invariant de Rham forms. One defines the *equivariant* de Rham complex of M to be the tensor product

$$(1.1) \quad \Omega_G(M) = \Omega(M)^G \otimes S(\mathfrak{g}^*)$$

with the coboundary operator

$$(1.2) \quad d_G(\omega \otimes f) = d\omega \otimes f + \sum \iota(\xi_M^i)\omega \otimes x_i f.$$

The *equivariant cohomology ring* of M , $H_G(M)$, is the cohomology ring of this complex. A few properties of this ring which we will need below are:

1. $H_G(M)$ is an $S(\mathfrak{g}^*)$ -module. (This follows from the fact that $\Omega_G(M)$ is an $S(\mathfrak{g}^*)$ module by (1.1) and d_G is an $S(\mathfrak{g}^*)$ module morphism by (1.2).)
2. $H_G(pt) = S(\mathfrak{g}^*)$.
3. Suppose M is compact and oriented. Then there is an integration operation

$$(1.3) \quad \int : \Omega_G(M) \longrightarrow S(\mathfrak{g}^*)$$

defined by

$$\int (\omega \otimes f) = f \int \omega.$$

It is easily checked that $\int d_G = 0$ and hence that this integration operation induces an integration operation on cohomology

$$(1.4) \quad \int : H_G(M) \longrightarrow S(\mathfrak{g}^*).$$

4. One can write d_G as a sum, $d_1 + d_2$, d_1 and d_2 being the first and second terms on the right hand side of (1.2). Thus $\Omega_G(M)$ is a *bi-complex*, and the additive structure of $H_G(M)$ can be computed by the spectral sequence of this bi-complex. The E_1 term in this spectral sequence is the d_1 -cohomology of $\Omega_G(M)$, namely

$$(1.5) \quad H(M) \otimes S(\mathfrak{g}^*).$$

One says that M is *equivariantly formal* if the spectral sequence is trivial, i.e. if, as vector spaces,

$$(1.6) \quad H_G(M) = H(M) \otimes S(\mathfrak{g}^*).$$

One can show, by the way, that if (1.6) holds as an identity of vector spaces, it also holds as an identity of $S(\mathfrak{g}^*)$ -modules. However, (1.6) doesn't, in general, tell one very much about the *ring* structure of $H_G(M)$ (about which we will have more to say in §1.6).

The property of being equivariantly formal is a bit technical; however there are a number of interesting assumptions on M which will imply this property. (See [GKM].) Of these assumptions the one that will be of most interest to us is the following:

THEOREM(Kirwan). *If M is a symplectic manifold and the action τ is Hamiltonian, M is equivariantly formal.*

1.2. The Atiyah-Bott-Berline-Vergne localization theorem. Let M be compact and oriented and, also, to simplify the statement of the localization theorem, let M^G be finite. For $p \in M^G$ one has an isotropy representation τ_p of G on T_p and we will denote the weights of this representation by $\alpha_{i,p}, i = 1, \dots, d$. Since τ_p is a real representation, these weights are, strictly speaking, only defined up to sign; however, since M is oriented, the product

$$\alpha_{1,p} \dots \alpha_{d,p}$$

is well-defined as an element of $S^d(\mathfrak{g}^*)$. Let

$$j_p : pt \rightarrow M$$

be the mapping “ pt ” onto p and note that if c is in $H_G(M)$, j_p^*c is in $H_G(pt)$ and thus in $S(\mathfrak{g}^*)$. The localization theorem asserts that, for every equivariant cohomology class $c \in H_G(M)$,

$$(1.7) \quad \int c = \sum_{p \in M^G} \frac{j_p^*c}{\prod \alpha_{i,p}}.$$

There are many deep and beautiful applications of (1.7) but the focus of our interest in this article is that (1.7) implies a lot of complicated identities among the weights $\alpha_{i,p}$. For instance, for $c = 1$, it implies

$$(1.8) \quad \sum (\prod \alpha_{i,p})^{-1} = 0.$$

What are these identities? In particular, are there simpler identities of which they are formal consequences? We will show in part 2 of this article that one can shed some light on these questions by looking at a graph-theoretical analogue of (1.7).

1.3. The Jeffrey-Kirwan theorem. Another interesting source for identities of type (1.7) is the Jeffrey-Kirwan theorem ([JK]): Let K be a one-dimensional connected closed subgroup of G with Lie algebra \mathfrak{k} and let $\xi \in \mathfrak{k}$ be a basis vector of the group lattice of K . Suppose M possesses a G -invariant symplectic form ω and that the action of K on M is Hamiltonian, i.e.

$$(1.9) \quad \iota(\xi_M)\omega = -df,$$

f being a G -invariant function. In addition suppose that

$$M^K = M^G$$

and hence that the critical points of f coincide with the fixed points of G . Let a be a regular value of f and let $M_a = f^{-1}(a)$. By the remark above, M_a contains no K -fixed points, so the action of K on M_a is locally free and the quotient space

$$M_a/K =: M_{red}$$

is an orbifold. Moreover, from the action of G on M_a one gets an inherited action of the quotient group

$$G/K =: G_1$$

on M_{red} . Let j be the inclusion of M_a into M and π the projection of M_a onto M_{red} . By the Marsden-Weinstein theorem there exists a symplectic form ω_{red} on M_{red} satisfying $\pi^*\omega_{red} = j^*\omega$. In particular, M_{red} is *oriented*. If \mathfrak{g}_1 is the Lie algebra of G_1 , its vector space dual, \mathfrak{g}_1^* , can be identified with the annihilator \mathfrak{g}_ξ^* of ξ in \mathfrak{g} ; so there is an integration operator

$$(1.10) \quad \int : H_{G_1}(M_{red}) \longrightarrow S(\mathfrak{g}_\xi^*).$$

Also, since the action of K on M_a is locally free, the map π induces an isomorphism

$$\pi^* : H_{G_1}(M_{red}) \longrightarrow H_G(M_a)$$

so one gets a map

$$(\pi^*)^{-1}j^* =: \mathcal{K}$$

of $H_G(M)$ into $H_{G_1}(M_{red})$. The Jeffrey-Kirwan theorem asserts that for every equivariant cohomology class $c \in H_G(M)$,

$$(1.11) \quad \int \mathcal{K}(c) = \text{Res}_\xi \left(\sum_{f(p) > a} \frac{j_p^* c}{\prod \alpha_{j,p}} \right), \quad p \in M^G,$$

Res_ξ being the residue of the rational function in brackets with respect to the “ ξ -coordinate” on \mathfrak{g}^* , the other coordinates being held fixed. (This residue can be defined intrinsically to be an element of $S(\mathfrak{g}_\xi^*)$. See §2.6)

1.4. The Smith problem. The Smith conjecture asserts that if M^G consists of two points, p and q , the isotropy representation of G at p is isomorphic (as a representation over \mathbb{R}) to the isotropy representation of G at q . The first complete proof of this theorem (for G an arbitrary compact Lie group) is due to Atiyah, Bott and Milnor (See [AB2], Theorem 3.83). If the cardinality of M^G is greater than two, the question of how the isotropy representations of G at distinct fixed points are related to each other is still open and is known as the “Smith problem”. In this section we will describe a few of the more obvious relations:

1. Relations of type J .

Suppose that M admits a G -invariant almost-complex structure. Then for every $p \in M^G$, the isotropy representations of G on T_p is a complex representation, so the weights of this representation,

$$(1.12) \quad \alpha_{i,p}, \quad i = 1, \dots, d$$

are unambiguously defined (not just defined up to sign).

For every closed subgroup H of G let \mathfrak{h} be the Lie algebra of H and

$$\rho_H : \mathfrak{g}^* \longrightarrow \mathfrak{h}^*$$

the transpose of the inclusion map $\mathfrak{h} \rightarrow \mathfrak{g}$. Let E be a connected component of M^H and p and q elements of E^G . We claim that the weights (1.12) can be ordered so that

$$(1.13) \quad \rho_H \alpha_{i,p} = \rho_H \alpha_{i,q}.$$

Proof. Let x be an arbitrary point of E and consider the isotropy representation of H on its normal space to E at x . This representation is a complex representation, so the weights of this representation are unambiguously defined and can't vary as x varies in E . Thus, in particular, they have to be the same at p and at q , implying (1.13). \square

2. Relations of type ω .

Assuming that M admits a G -invariant almost-complex structure, J , is equivalent to assuming that M admits a G -invariant "almost-symplectic" structure, i.e., a two-form, ω , which is everywhere of maximal rank. Suppose that ω is actually a symplectic form and the action τ is *Hamiltonian*, or, in other words, that there exists a moment map

$$\Phi : M \rightarrow \mathfrak{g}^*.$$

From the convexity theorem ([A], [GS]) one gets

THEOREM. *Let Δ be the set of regular values of Φ in $\Phi(M)$. Then Δ is a disjoint union of open convex polytopes. Moreover, the vertices of these polytopes are the images of the fixed points $p \in M^G$ and the edges going out of these vertices are pointing in the directions of the vectors $\alpha_{i,p}$.*

The one-skeleton, Γ , of this configuration is called the *moment graph* of M . (See [Gu].) It exhibits a lot of relations among the $\alpha_{i,p}$'s which are probably not much simpler than the relations (1.7) but have the virtue of being of a more geometric character.

1.5. The Goresky-Kottwitz-MacPherson graph. We will assume from now on that M admits a G -invariant almost-complex structure. Thus, for very $p \in M^G$, the weights, $\alpha_{i,p} \in \mathfrak{g}^*$, are unambiguously defined. In addition we will assume: if $i \neq j$, $\alpha_{i,p}$ and $\alpha_{j,p}$ are linearly independent. This "GKM hypothesis" has the following consequence: Let $\mathfrak{h} = \mathfrak{h}_i$ be the annihilator of $\alpha_{i,p}$ in \mathfrak{g} and let H be the $(n-1)$ -dimensional subtorus of G with Lie algebra, \mathfrak{h} . Clearly, $p \in M^G \subseteq M^H$.

PROPOSITION. *Let E be the connected component of M^H containing p . Then*

$$E \cong \mathbb{C}P^1 \cong S^2$$

and the action of G on E is the standard action of the circle, S^1 , on S^2 by "rotation about the z -axis". In particular E contains just two G -fixed points (one of which is p).

Proof. The tangent space to E at p is the 2-dimensional subspace of T_p on which G acts with weight $\alpha_{i,p}$ so E itself is 2-dimensional. Since E is compact and the action of G/H is non-trivial, E is diffeomorphic to S^2 , and this action is the standard action of S^1 on S^2 by the Korn-Lichtenstein theorem. \square

Let q be the other fixed point of G in E . Let $\alpha_{p,e} = \alpha_{i,p}$ be the weight of the isotropy representation of G on $T_p E$ and let $\alpha_{q,e} = \alpha_{j,q}$ be the weight of the isotropy representation of G on $T_q E$. From the fact that the action of G on E is diffeomorphic to the standard action of S^1 on S^2 it follows that

$$(1.14) \quad \alpha_{p,e} = -\alpha_{q,e}.$$

For each of the weights $\alpha_{i,p}$ one gets an embedded $\mathbb{C}P^1$ of the type above, and we can represent these $\mathbb{C}P^1$'s graphically by d lines issuing from p . Each of these lines joins p to another fixed point, q , and the $\mathbb{C}P^1$'s associated with the weights, $\alpha_{j,q}$, can also be represented graphically by d lines issuing from q . One of these will be the line from p to q , but the remaining $d - 1$ lines will join q to other fixed points. By repeating this construction over and over until one runs out of fixed points, one obtains a finite d -valent graph, Γ , the vertices of which are the fixed points of G and the edges of which correspond to embedded $\mathbb{C}P^1$'s, each of these $\mathbb{C}P^1$'s being a connected component of the fixed point set of an $(n - 1)$ -dimensional subtorus of G . We will call Γ the Goresky-Kottwitz-MacPherson (GKM) graph of M .

EXAMPLE. Suppose M is a Hamiltonian G -manifold whose moment map, Φ , maps M^G injectively into \mathfrak{g}^* . Then Φ embeds the GKM graph into \mathfrak{g}^* , and its image is the moment graph.

1.6. The Goresky-Kottwitz-MacPherson theorem. The graph Γ is equipped with an additional piece of structure. Namely let I_Γ be the incidence relation of this graph: the set of all pairs (p, e) , p being a vertex and e an edge containing p . Then one has a map

$$(1.15) \quad \alpha : I_\Gamma \longrightarrow \mathfrak{g}^* - \{0\}$$

mapping (p, e) to the weight $\alpha_{p,e}$. We will call this the *axial function* of Γ . It has the following properties (the first two of which we have already commented on):

1. If e is an edge and p and q are the vertices joined by e

$$(1.16) \quad \alpha_{p,e} = -\alpha_{q,e}.$$

2. If p is a vertex and e_1, \dots, e_d are the edges containing p , the vectors

$$(1.17) \quad \alpha_{p,e_i}, \quad i = 1, \dots, d$$

are pair-wise linearly independent.

3. Let e be an edge and p and q the vertices joined by e . Let

$$\mathfrak{g}_e = \{\xi \in \mathfrak{g}, \alpha_{p,e}(\xi) = 0\}$$

and let

$$\rho_e : \mathfrak{g}^* \longrightarrow \mathfrak{g}_e^*$$

be the transpose of the inclusion map, $\mathfrak{g}_e \longrightarrow \mathfrak{g}$. Let $e_i, i = 1, \dots, d$ and $e'_i, i = 1, \dots, d$ be the edges containing p and q respectively, with $e_d = e'_d = e$. Then the e_i 's can be ordered so that

$$(1.18) \quad \rho_e \alpha_{p,e_i} = \rho_e \alpha_{q,e'_i}.$$

Proof. (1.18) is just a special case of (1.13), H being the subtorus of G with Lie algebra \mathfrak{g}_e . \square

From the data (Γ, α) one can construct a graded ring

$$(1.19) \quad H(\Gamma, \alpha) = \bigoplus H^{2k}(\Gamma, \alpha)$$

as follows. For each edge e , the map $\rho_e : \mathfrak{g}^* \rightarrow \mathfrak{g}_e^*$ extends to a ring morphism

$$(1.20) \quad \rho_e : S(\mathfrak{g}^*) \rightarrow S(\mathfrak{g}_e^*).$$

Let V_Γ be the set of vertices of Γ and let $H^{2k}(\Gamma, \alpha)$ be the set of all maps

$$f : V_\Gamma \rightarrow S^k(\mathfrak{g}^*)$$

satisfying the compatibility condition:

$$(1.21) \quad \rho_e f(p) = \rho_e f(q)$$

for all vertices, p and q , and edges, e , joining p to q . $H(\Gamma, \alpha)$ can be given a ring structure by pointwise multiplication

$$(f_1 f_2)(p) = f_1(p) f_2(p).$$

(Notice that if f_1 and f_2 satisfy (1.21) so does $f_1 f_2$ since ρ_e is a ring morphism.) In addition, $H(\Gamma, \alpha)$ contains $S(\mathfrak{g}^*)$ as a subring: the ring of constant maps of V_Γ into $S(\mathfrak{g}^*)$. In particular, $H(\Gamma, \alpha)$ is a module over $S(\mathfrak{g}^*)$.

THEOREM([GKM]). *If M is equivariantly formal, $H_G(M)$ is isomorphic, as a graded ring, to $H(\Gamma, \alpha)$.*

We will sketch a proof of this at the end of §1.8.

1.7. Holonomy. Let \mathbb{P} be the complex projective line, let z be the standard coordinate function on $\mathbb{C} = \mathbb{P}_0 = \mathbb{P} - \{\infty\}$ and $w = z^{-1}$ the corresponding coordinate function on $\mathbb{P}_\infty = \mathbb{P} - \{0\}$. The multiplicative group, $\mathbb{C}^\# = \mathbb{C} - \{0\}$, acts on \mathbb{C} by homotheties:

$$(a, z) \rightarrow az$$

and this extends to a holomorphic action, ρ , of $\mathbb{C}^\#$ on \mathbb{P} . Let E be a holomorphic, rank r vector bundle over \mathbb{P} and suppose ρ lifts to an action, ρ , of $\mathbb{C}^\#$ on E by vector bundle automorphisms. Let E_0 and E_∞ be fibers of E over 0 and ∞ . The isotropy representations of $\mathbb{C}^\#$ at 0 and ∞ decompose these spaces into invariant one-dimensional subspaces

$$(1.22) \quad E_0 = V_1 \oplus \cdots \oplus V_r$$

and

$$(1.23) \quad E_\infty = V'_1 \oplus \cdots \oplus V'_r.$$

Let m_1, \dots, m_r be the weights of the representations of $\mathbb{C}^\#$ on V_1, \dots, V_r . We will assume that these weights are all distinct and that $m_1 < m_2 < \cdots < m_r$. Similarly

we will assume that the weights m'_i of the representations of $\mathbb{C}^\#$ on V'_1, \dots, V'_r are all distinct (but we won't require that $m'_1 < \dots < m'_r$). By a theorem of Birkhoff-Grothendieck ([Kl]) there is an equivariant decomposition of E into line-bundles

$$(1.24) \quad E = \mathbb{L}_1 \oplus \dots \oplus \mathbb{L}_r$$

such that the fiber of \mathbb{L}_i over 0 is V_i . Moreover this decomposition is *unique up to isomorphism*. (To see this, let

$$E = \mathbb{L}'_1 \oplus \dots \oplus \mathbb{L}'_r$$

be another decomposition with these properties: Over \mathbb{P}_0 , one can find a trivializing section, s_i , of \mathbb{L}_i which transforms under $\mathbb{C}^\#$ according to the law

$$(1.25) \quad \rho_a^* s_i = a^{m_i} s_i$$

and a trivializing section, s'_i , of \mathbb{L}'_i with the same transformation property. Moreover, one can assume that $s_i(0) = s'_i(0)$. We claim that $s_1 \equiv s'_1$ (and hence $\mathbb{L}_1 = \mathbb{L}'_1$). To see this we note that because of the transformation law (1.25)

$$s'_1 = s_1 + \sum_{i>1} c_i z^{m_1 - m_i} s_i$$

on \mathbb{P}_0 . However, since s'_1 is holomorphic near 0 and $m_1 - m_i$ is strictly negative, the constants, c_i , are all zero.

Applying this argument to the quotient bundle, E/\mathbb{L}_1 , one concludes by induction that $\mathbb{L}_i \cong \mathbb{L}'_i$ for all i . Q.E.D.)

In particular, coming back to the isotropy decompositions (1.22) and (1.23), one has a canonical map of the r -element set

$$(1.26) \quad \{V_1, \dots, V_r\}$$

onto the r -element set

$$(1.27) \quad \{V'_1, \dots, V'_r\}$$

which maps $V_i = (\mathbb{L}_i)_0$ onto $(\mathbb{L}_i)_\infty$; and we can relabel the V'_i 's so that this map maps the element, V_i , of the set (1.26) onto the element, V'_i , of the set (1.27). We will call the map the *holonomy* map.

Let us apply these remarks to one of the $\mathbb{C}P^1$'s in §1.5; i.e., let H be a subtorus of G of codimension one and $X = \mathbb{C}P^1$ a 2-dimensional connected component of M^H . Let NX be the normal bundle of X and let β_1, \dots, β_N be the weights of the isotropy representation of H on a typical fiber of X . The fiber of N over $x \in X$ splits into a direct sum of vector subspaces

$$(1.28) \quad (E_1)_x \oplus \dots \oplus (E_N)_x$$

such that the weight of the representation of H on $(E_i)_x$ is β_i . Since the β_i 's can occur with multiplicities, the dimension of $(E_i)_x$ doesn't have to be one; however the assignment, $x \rightarrow (E_i)_x$, defines a complex vector bundle, E_i , over X . Moreover, this bundle can be given a holomorphic structure, and the action of $S^1 = G/H$ on E_i can

be extended to a holomorphic action of $\mathbb{C}^\#$. In particular one can define for E_i a *holonomy* map of the type we described at the beginning of this section.

Now let $X^G = \{p, q\}$, and $\alpha_{i,p}$ and $\alpha_{i,q}$ be the weights of the isotropy representations of G on T_pM and T_qM . By assumption these weights occur with multiplicity one, so one gets decompositions of T_pM and T_qM into direct sums of one-dimensional weight spaces

$$T_p^{\alpha_{1,p}} \oplus \dots \oplus T_p^{\alpha_{d,p}}$$

and

$$T_q^{\alpha_{1,q}} \oplus \dots \oplus T_q^{\alpha_{d,q}}.$$

Let ρ be the projection of \mathfrak{g}^* onto \mathfrak{h}^* . By (1.18) we can reorder the $\alpha_{i,q}$'s so that $\rho(\alpha_{i,p}) = \beta_i = \rho(\alpha_{i,q})$; so, if the β_i 's are all distinct, one has a *canonical* map of the set

$$(1.29) \quad \{\alpha_{1,p}, \dots, \alpha_{d,p}\}$$

onto the set

$$(1.30) \quad \{\alpha_{1,q}, \dots, \alpha_{d,q}\}$$

mapping $\alpha_{i,p}$ onto $\alpha_{i,q}$. However, this canonical map can even be defined when the β_i 's are *not* distinct (that is, when the vector bundles, E_i , are *not* of rank one) by using the holonomy structure on these bundles. In other words there exists a canonical holonomy mapping from the set (1.29) to the set (1.30) which, when the β_i 's are distinct, is defined trivially by the recipe

$$\alpha_{i,p} \longrightarrow \rho(\alpha_{i,p}) = \beta_i = \rho(\alpha_{i,q}) \longrightarrow \alpha_{i,q},$$

but, when the β_i 's are not distinct, involves some topological properties of the bundles, E_i . We will denote this map by $\theta_{p,e}$.

One can give a slightly more "graphical" description of this map: As in §1.6, let Γ be the GKM graph, let V_Γ be its vertices and let I_Γ be the incidence relation. The projection, $I_\Gamma \rightarrow V_\Gamma$, can be regarded as a fiber bundle over V_Γ , the fiber, E_p , over $p \in V_\Gamma$ being the set of all pairs, (p, e) in I_Γ (i.e., the set of all oriented edges of Γ pointing out from p). In this "fiber bundle" picture, $\theta_{p,e}$ is just a map

$$(1.31) \quad \theta_{p,e} : E_p \longrightarrow E_q$$

with the properties

$$(1.32) \quad \theta_{p,e}(p, e) = (q, e)$$

and

$$(1.33) \quad \theta_{p,e} = \theta_{q,e}^{-1},$$

e being the edge joining p to q . A family of maps, θ , with the properties (1.32)–(1.33) is called a *connection* on Γ (cf. [CS]). In terms of this connection, one can reformulate (1.18) more precisely:

THEOREM 1.1. *Let e be an edge of Γ joining the vertex, p , to the vertex, q . Then for every $(p, e_i) \in E_p$*

$$(1.34) \quad \rho_e(\alpha_{p,e_i}) = \rho_e(\alpha_{q,e'_i}),$$

(q, e'_i) being the image with respect to $\theta_{p,e}$ of (p, e_i) .

1.8. Betti numbers. The theorem of Goresky-Kottwitz-MacPherson described in § 1.6 implies that the odd Betti numbers of M are zero. The even Betti numbers can be computed as follows. As in §1.3 let K be a one-dimensional closed connected subgroup of G with $M^G = M^K$ and let ξ be a basis vector of \mathfrak{k} . For every $p \in V_\Gamma$ let σ_p be the number of edges, e , with one vertex p and with $\alpha_{p,e}(\xi) < 0$.

THEOREM. *The $2k$ -th Betti number, β_k , of M is equal to the number of points, $p \in V_\Gamma$, with $\sigma_p = k$.*

If M possesses a G -invariant symplectic form having the properties described in §1.3, this theorem can be proved by Morse theory : Let f be the function defined by (1.9). The critical points of this function coincide with the fixed points of G and it is not difficult to show that the index of the Hessian of f at $p \in M^G$ is just $2\sigma_p$.

We will now sketch a proof of the Goresky-Kottwitz-MacPherson theorem. It is clear that

$$H_G(M^G) = H^0(M^G) \otimes S(\mathfrak{g}^*) = \text{Maps}(V_\Gamma, S(\mathfrak{g}^*));$$

so if $i : M^G \rightarrow M$ is the inclusion, there is an induced map

$$i^* : H_G(M) \rightarrow \text{Maps}(V_\Gamma, S(\mathfrak{g}^*)).$$

It is easy to see that the image of i^* is contained in $H(\Gamma, \alpha)$. Moreover, by a classical theorem of Borel, i^* is an isomorphism modulo torsion; in particular, the kernel of i^* is a torsion submodule of $H_G(M)$. However, by (1.6), $H_G(M)$ is a free $S(\mathfrak{g}^*)$ -module; so $i^* : H_G^{2k}(M) \rightarrow H^{2k}(\Gamma, \alpha)$ is injective and

$$\dim H_G^{2k}(M) = \sum \beta_r \dim S^{k-r}(\mathfrak{g}^*).$$

Therefore, in order to prove that

$$(1.35) \quad \dim H^{2k}(\Gamma, \alpha) = \sum \beta_r \dim S^{k-r}(\mathfrak{g}^*),$$

it suffices to show that the dimension of $H^{2k}(\Gamma, \alpha)$ is less than or equal to the right hand side of (1.35), and we will prove this (by a very simple algebraic argument) in section 2.9.

2.1. Graphs and axial functions. In part 2 of this article, \mathfrak{g} will simply be a vector space over \mathbb{R} of dimension n and \mathfrak{g}^* its vector space dual. (In particular \mathfrak{g} will *not* necessarily be the Lie algebra of a group, G .) Let Γ be a finite simple d -valent graph and I_Γ its incidence relation.

DEFINITION. 1. An *axial function* on Γ is a map $\alpha : I_\Gamma \rightarrow \mathfrak{g}^* - \{0\}$ having the properties (1.16)–(1.18).

Note that by axiom (1.16) an axial function can be thought as a function

$$\alpha : E_\Gamma^\pm \rightarrow \mathfrak{g}^*$$

on the set of *oriented* edges of Γ , having the property that if e^+ and e^- are the two oriented edges of Γ associated with an unoriented edge e , then

$$\alpha(e^+) = -\alpha(e^-).$$

DEFINITION. A *connection*, θ , on Γ is a collection of maps, $\theta_{p,e}$, $(p, e) \in I_\Gamma$, satisfying the axioms (1.31)–(1.33); an axial function, α , is *compatible* with θ if it satisfies the stronger version, (1.34) of axiom (1.18).

For instance, suppose that for every vertex, p , and every edge, e , containing p , no two of the vectors (1.18) are equal. Then, as we showed in section 1.7, there is a *unique* connection on Γ which is compatible with α .

We will list below a few examples of axial functions and connections on graphs and describe some of their functorial properties:

EXAMPLE 1. *The complete graph on N vertices.* The vertices of this graph are the elements of the N -element set $\{1, \dots, N\}$ and each pair of elements, (i, j) , $i \neq j$, is joined by an edge. Thus the set of oriented edges is just the set

$$\{(i, j); 1 \leq i, j \leq N, i \neq j\}.$$

Let α_i , $i = 1, \dots, N$ be non-zero elements of \mathfrak{g}^* . Then the function $(i, j) \rightarrow \alpha_i - \alpha_j$ satisfies (1.16) and (1.18) and hence it is an axial function iff, for every i , the $N - 1$ vectors $\alpha_i - \alpha_j$, $i \neq j$ are pairwise linearly independent. Conversely, we claim that every axial function is of this form. (Proof: Let $(i, j) \rightarrow \alpha_{i,j}$ be an axial function. Then if i, j and k are distinct

$$\alpha_{i,k} = \alpha_{j,k} + c_{i,j}\alpha_{i,j} \quad c_{i,j} = c_{j,i}.$$

Hence

$$\alpha_{k,i} = \alpha_{j,i} + c_{k,j}\alpha_{k,j} = \alpha_{k,j} + c_{i,j}\alpha_{j,i}$$

so $(1 - c_{i,j})\alpha_{j,i} = (1 - c_{k,j})\alpha_{k,j}$ and $c_{i,j} = c_{k,j} = 1$. Now let $\alpha_1 = 0$ and let $\alpha_i = \alpha_{i,1}$ for $i > 1$.)

An α -compatible connection, θ , can be defined as follows: For every oriented edge, (i, j) , of Γ let $\theta_{i,j}$ be the map of the set of edges of Γ containing i onto the set of edges of Γ containing j which maps (i, j) onto (j, i) and, for $k \neq i, j$, maps (i, k) onto (j, k) .

EXAMPLE 2. *Sub-objects.* Let Γ_1 be an r -valent sub-graph of Γ and j the embedding of V_{Γ_1} into V_Γ . From j one gets an embedding $i: I_{\Gamma_1} \rightarrow I_\Gamma$ and one can pull-back the axial function α to I_{Γ_1} . In general $i^*\alpha$ won't be an axial function; however if it is, we will say that Γ_1 is *compatible* with α and call $(\Gamma_1, i^*\alpha)$ a *sub-object* of (Γ, α) .

EXAMPLE 3. *Sub-objects which are "totally geodesic" with respect to a connection.* Let θ be a connection on Γ , and let Γ' be an r -valent subgraph of Γ . For every vertex, p , of Γ let E_p and E'_p be the oriented edges of Γ and of Γ' issuing from p . We will say that Γ' is totally geodesic with respect to θ if, for every oriented edge, (p, q) , of Γ' the restriction of the holonomy map, $\theta_{p,q}: E_p \rightarrow E_q$, to E'_p maps E'_p onto E'_q .

If this happens, this restriction defines an induced connection, θ' , on Γ' . Moreover, if $\alpha: I_\Gamma \rightarrow \mathfrak{g}^*$ is an axial function which is compatible with θ , the restriction of α to $I_{\Gamma'}$ is an axial function and is compatible with θ' .

EXAMPLE 4. *The totally geodesic sub-objects of Γ_N .* Let Γ_N be the complete graph on N vertices and for every subset, S , of $\{1, \dots, N\}$ let Γ_S be the graph

whose vertices are the elements of S and whose oriented edges are the pairs (s_1, s_2) , $(s_i \in S)$, $s_1 \neq s_2$. It is obvious that Γ_S is a totally geodesic with respect to the connection we defined in example 1; and, in fact, every totally geodesic sub-object of Γ_N is a Γ_S for some subset S . (To see this, let Γ' be a connected totally geodesic subgraph and let (p_1, p_2) be an oriented edge of Γ' . If q is a vertex of Γ' distinct from p_1 and p_2 and (p_1, q) is an oriented edge of Γ' , (p_2, q) has to be an oriented edge of Γ' , so it follows from the connectivity of Γ' that, for every pair of vertices, p and q , of Γ' (p, q) is an oriented edge of Γ' .)

EXAMPLE 5. *The graph $\Gamma_{\mathfrak{h}}$.* Let \mathfrak{h} a vector subspace of \mathfrak{g} and let $\rho_{\mathfrak{h}} : \mathfrak{g}^* \rightarrow \mathfrak{h}^*$ be the transpose of the inclusion map $\mathfrak{h} \rightarrow \mathfrak{g}$. Let $\Gamma_{\mathfrak{h}}$ be the subgraph of Γ whose edges are the edges, e , of Γ for which

$$(2.1) \quad \rho_{\mathfrak{h}} \alpha_{p,e} = -\rho_{\mathfrak{h}} \alpha_{q,e} = 0,$$

p and q being the vertices of e . Each connected component of this graph is k -valent for some k and is a sub-object in the sense of item 2. Moreover, if p and q are in the same connected component of $\Gamma_{\mathfrak{h}}$ and e_i and e'_i , $i = 1, \dots, d$, are the edges of Γ containing p and q , one can order the e_i 's so that

$$(2.2) \quad \rho_{\mathfrak{h}} \alpha_{p,e_i} = \rho_{\mathfrak{h}} \alpha_{q,e'_i}$$

(compare with (1.13)).

EXAMPLE 6. *Product objects.* Let Γ_1 be a graph of valence d_1 and Γ_2 a graph of valence d_2 . The vertices of the *product* graph, $\Gamma_1 \times \Gamma_2$, are the pairs (p, q) , $p \in V_{\Gamma_1}$ and $q \in V_{\Gamma_2}$; two vertices (p, q) and (p', q') are joined by an edge if either $p = p'$ and q and q' are joined by an edge in Γ_2 or $q = q'$ and p and p' are joined by an edge in Γ_1 . Thus this product graph is a $d_1 + d_2$ -valent graph and its set of oriented edges is the disjoint union

$$(2.3) \quad E_{\Gamma_1}^{\pm} \amalg E_{\Gamma_2}^{\pm}$$

of the set of oriented edges of Γ_1 and the set of oriented edges of Γ_2 . If

$$\alpha_i : E_{\Gamma_i}^{\pm} \rightarrow \mathfrak{g}^*, \quad i = 1, 2$$

is an axial function on Γ_i , one defines the product function α on $\Gamma_1 \times \Gamma_2$ to be the function which is equal to α_1 on the first summand of (2.3) and equal to α_2 on the second summand. Then α satisfies (1.16) - (1.18) and it is called *the product axial function*. If in addition, θ_1 and θ_2 are connections on Γ_1 and Γ_2 one can define a product connection, θ , on $\Gamma_1 \times \Gamma_2$ by letting

$$\theta_{(p,q;p',q)} = \theta_{p,p'} \times (\text{Identity})_{q,q}$$

and letting

$$\theta_{(p,q;p,q')} = (\text{Identity})_{p,p} \times \theta_{q,q'}.$$

If α_i and θ_i are compatible, for $i = 1$ and 2 , the product axial function which we defined above is compatible with θ .

EXAMPLE 7. *Blowing-up*. This operation can be defined for any sub-object of a graph-axial function pair; however, for simplicity, we will only consider here the special case when the sub-object is a point. Let Γ be a finite simple d -valent graph and let p_0 be an arbitrary vertex of Γ . Let e_i , $i = 1, \dots, d$ be the edges of Γ containing p_0 and let q_i be the vertex joined by e_i to p_0 . From this data one can construct a new graph, $\Gamma^\#$, as follows. Replace the vertex p_0 by d new vertices, p_i , $i = 1, \dots, d$ (which one should think of as being the “baricenters” of the edges e_i) and to each of these new vertices adjoin d edges; one edge going from p_i to q_i (which one should think of as being a replacement for the old edge e_i) and one edge going from p_i to each of the p'_j 's, $j \neq i$. Let

$$(2.4) \quad \beta : V_{\Gamma^\#} \longrightarrow V_\Gamma$$

be the map which sends $\{p_1, \dots, p_d\}$ to p_0 and is the identity on the complement of $\{p_1, \dots, p_d\}$. We will call this map the *blowing-down* map. The set $\{p_1, \dots, p_d\}$, which is the pre-image of p_0 with respect to β , is the set of vertices of a sub-graph, Γ_0 , of $\Gamma^\#$ (the complete graph on d vertices) which we will call *the singular locus* of the blowing-down map β .

Now let $\alpha : \Gamma \longrightarrow \mathfrak{g}^*$ be an axial function and let α_i , $i = 1, \dots, d$ be the values of α on the vertex-edge pairs (p_0, e_i) . Let us assume that for each i the $d - 1$ vectors $\alpha_j - \alpha_i$, $j \neq i$ are pairwise linearly independent. We can then define an axial function, $\alpha^\#$, on $\Gamma^\#$, as follows:

1. On the oriented edges, e , of Γ , not containing p_0 , $\alpha^\#(e) = \alpha(e)$.
2. On the oriented edges, $e = (p_i, q_i)$, $\alpha^\#(e) = \alpha_i$.
3. On the oriented edges, $e = (p_i, p_j)$, $\alpha^\#(e) = \alpha_j - \alpha_i$.

This defines $\alpha^\#$ on *all* edges of $\Gamma^\#$ and it is easy to check that $\alpha^\#$ satisfies the axioms (1.16) - (1.18).

Finally, if θ is a connection on Γ there is a unique connection, $\theta^\#$, on $\Gamma^\#$ with the following properties.

1. The restriction of θ to Γ_0 is the connection described in example one.
2. $\theta^\#_{(p_i, p_j)}$ maps the oriented edge, (p_i, q_i) , to the oriented edge (p_j, q_j) .
3. $\theta^\#_{(p_i, q_i)}$ maps the oriented edge (p_i, q_i) to the oriented edge $\theta_{(p_i, q_i)}(e_j)$.
4. If q is not equal to one of the p_i 's and q and q_i are joined by an edge,

$$\theta^\#_{(q_i, q)}(e_i^\#) = \theta_{(q_i, q)}(e_i^-),$$

$e_i^\#$ being the edge joining q_i to p_i and e_i^- the edge joining q_i to p . On the other edges of $\Gamma^\#$ issuing from q_i , $\theta^\#_{(q_i, q)} = \theta_{(q_i, q)}$.

5. If q and q' are not equal to one of the q_i 's or one of the p_i 's, $\theta^\#_{(q, q')} = \theta_{(q, q')}$.

EXAMPLE 8. *The case $d = \dim \mathfrak{g}^* = 2$* . Let Γ be a finite connected 2-valent graph with N vertices, \mathfrak{g}^* a 2-dimensional vector space and α an axial function. We will orient the edges of Γ so that for each vertex, p , one of the edges containing p is pointing in the direction of p and the other is pointing away from p (since Γ is connected there are clearly just two ways of orienting the edges so that their orientations have this property.). Let $p_1, \dots, p_N, p_{N+1} = p_1$ be an enumeration of the vertices of Γ such that the outward pointing edge at p_i joins p_i to p_{i+1} and let α_i be the value of α on

(p_i, p_{i+1}) . Then (1.18) is equivalent to

$$(2.5) \quad \alpha_{i+1} \wedge \alpha_i = \alpha_i \wedge \alpha_{i-1}$$

for all i . (For example for $N = 4k$, let $\{\alpha_1, \alpha_2\}$ be a basis of \mathfrak{g}^* . Then a solution of (2.5) is obtained by letting $\alpha_1 = -\alpha_3 = \alpha_5 = \dots$ and $\alpha_2 = -\alpha_4 = \alpha_6 = \dots$).

2.2. Orientations. Let (Γ, α) be a graph-axial function pair and let

$$\mathcal{P} = \{\xi \in \mathfrak{g}, \alpha_{p,e}(\xi) \neq 0 \text{ for all } (p, e) \in I_\Gamma\}.$$

Then for every $\xi \in \mathcal{P}$, the axial function α defines an *orientation* of Γ ; in other words, for each edge e , it fixes an ordering of the vertices of e . Namely if p and q are the vertices of e , one orders them so that

$$(2.6) \quad p < q \Leftrightarrow \alpha_{p,e}(\xi) > 0.$$

It is clear that this orientation doesn't depend on ξ but only on the connected component of \mathcal{P} in which ξ is contained. On the other hand it is clear that different components will give rise to different orientations (for instance, replacing ξ by $-\xi$ reverses all the orientations). We will say that Γ satisfies the *no-cycle condition* if, for at least one of these orientations, Γ has no cycles.

DEFINITION. Given $\xi \in \mathcal{P}$, a function $f : V_\Gamma \rightarrow \mathbb{R}$ is *positively oriented with respect to ξ* if, for every pair of vertices, p and q , and edge, e , joining p to q , the ratio of $f(p) - f(q)$ to $\alpha_{q,e}(\xi)$ is positive.

If f is positively oriented with respect to ξ , the orientation of Γ associated with ξ can't have closed cycles since f has to be strictly increasing along any oriented path. We will prove that the converse is true:

THEOREM. *If the orientation of Γ associated with ξ has no cycles, there exists a function $f : V_\Gamma \rightarrow \mathbb{R}$ which is positively oriented with respect to ξ .*

Proof. Given $p \in V_\Gamma$, consider the *longest* oriented path with initial point p , i.e., the longest sequence

$$(2.7) \quad p = p_1, p_2, \dots, \quad p_i \in V_\Gamma$$

with the property that p_i and p_{i+1} are the vertices of a common edge and, relative to the orientation on this edge, $p_i < p_{i+1}$. If Γ has no cycles this longest path has to be of finite length, i.e., has to terminate at some point, p_N . Now set $f(p) = -N$. It is easy to check that this function is positively oriented with respect to ξ . \square

REMARKS.

1. The vertices, p , where $f(p) = 0$ have the property that all edges containing p are pointing "into" p , i.e. p is a "maximum" of the oriented graph Γ . In particular, if $f(p) = -N$, this is true of the vertex p_N in the sequence (2.7); so the argument above shows that every vertex can be joined by an oriented path to a maximal vertex.

2. One can perturb f so that it remains positively oriented with respect to ξ and, in addition, takes on distinct values at distinct vertices. Namely suppose that

$$f^{-1}(k) = \{p_1, \dots, p_r\}.$$

Redefine f on the set $\{p_1, \dots, p_r\}$ by setting f equal to $k + \epsilon_i$ on p_i where $\epsilon_i \neq \epsilon_j$ for $i \neq j$ and the ϵ_i 's are small. This redefined function is still positively oriented with respect to ξ but now takes distinct values at p_1, \dots, p_r .

2.3. The cohomology ring of (Γ, α) . We define the *cohomology ring* of (Γ, α) to be the ring $H(\Gamma, \alpha)$ which we defined in §1.6. As (Γ, α) is no longer the GKM data associated with a G -manifold, it is, perhaps, a misnomer to refer to this ring as a “cohomology ring”; however, there are other reasons for using this terminology. For instance, if Γ is the one-skeleton of a simplicial polytope, $H(\Gamma, \alpha)$ is just the Stanley-Reissner cohomology ring of the dual polytope. (We are indebted to Mark Goresky for this observation.) We will describe below a few properties of this ring.

1. As we pointed out in §1.6, $H(\Gamma, \alpha)$ contains $S(\mathfrak{g}^*)$ as a subring.
2. *Chern classes:* For each $p \in V_\Gamma$, let e_1, \dots, e_d be the edges containing p and let $c_k(p)$ be the k -th elementary symmetric function in the monomials $\alpha_{p,e_1}, \dots, \alpha_{p,e_d}$. The function $p \rightarrow c_k(p)$ defines an element c_k of $H^{2k}(\Gamma, \alpha)$ which can be thought of as the k -th Chern class of the “tangent bundle” of Γ .
3. *Symplectic structures:* An element of $H^2(\Gamma, \alpha)$ is just a map $c : V_\Gamma \rightarrow \mathfrak{g}^*$ satisfying

$$(2.8) \quad c(p) - c(q) = \lambda_e \alpha_{q,e}$$

for every pair of vertices, p, q , and edge, e , joining p to q . We will call c *symplectic* if, for every edge e , λ_e is positive. The existence of a symplectic structure implies that for every $\xi \in \mathcal{P}$, the orientation of Γ associated with ξ has the no-cycle property. (Proof: It follows from (2.8) that the ξ -component of c is an \mathbb{R} -valued function on V_Γ which is positively oriented with respect to ξ .)

4. *Thom classes:* Fix a vertex p and let e_1, \dots, e_d be the edges containing p . Let $\tau : V_\Gamma \rightarrow S^d(\mathfrak{g}^*)$ be the map which is zero at $q \neq p$ and at p is equal to $\alpha_{p,e_1} \dots \alpha_{p,e_d}$. Then τ is in $H^{2d}(\Gamma, \alpha)$.
5. *Sub-objects:* Let Γ_1 be a sub-graph of Γ which is compatible with α . Then the inclusion map $j : V_{\Gamma_1} \rightarrow V_\Gamma$ induces a map

$$j^* : H(\Gamma, \alpha) \rightarrow H(\Gamma_1, \alpha_1), \quad \alpha_1 = i^* \alpha.$$

6. *Gysin maps:* Suppose that Γ_1 is compatible with α . The *Thom class* of Γ_1 is the map $\tau : V_\Gamma \rightarrow S^{d-r}(\mathfrak{g}^*)$ which is zero on the vertices of Γ which are not vertices of Γ_1 and on vertices, p , of Γ_1 , is equal to

$$\tau(p) = \alpha_{p,e_1} \dots \alpha_{p,e_s}$$

where $s = d - r$ and e_1, \dots, e_s are the edges of Γ at p which don't belong to Γ_1 .

From τ one gets a Gysin map

$$H^{2k}(\Gamma_1, \alpha_1) \rightarrow H^{2(k+s)}(\Gamma, \alpha)$$

mapping f to τf . (Since τ is supported on V_Γ , this map is well-defined.)

7. *The cohomology of blow-ups:* Let Γ be a d -valent graph and $\alpha : I_\Gamma \rightarrow \mathfrak{g}^*$ an axial function. Let p_0 be a vertex of Γ and $(\Gamma^\#, \alpha^\#)$ the blow-up of (Γ, α) at p_0 (See §2.1). From the blowing down map (2.4) one gets a pull-back map on cohomology

$$\beta^* : H(\Gamma, \alpha) \rightarrow H(\Gamma^\#, \alpha^\#)$$

which embeds $H(\Gamma, \alpha)$ as a subring of $H(\Gamma^\#, \alpha^\#)$. Moreover the singular locus, Γ_0 , of β , is a sub-object of $\Gamma^\#$ (in the sense of example 2 above) and its Thom class,

$$\tau \in H^2(\Gamma^\#, \alpha^\#)$$

generates $H(\Gamma^\#, \alpha^\#)$ over the sub-ring $H(\Gamma, \alpha)$, subject to the relation

$$\tau^d - c_1\tau^{d-1} + c_2\tau^{d-2} - \dots \pm c_d,$$

the c_i 's being the Chern classes of Γ for $i < d$ and c_d being the Thom class of p_0 .

2.4. The Atiyah-Bott-Berline-Vergne localization theorem. We have just discussed functoriality for sub-objects of Γ . What about quotient objects? To take the most extreme case let “ pt ” be the trivial zero-valent graph consisting of one vertex, pt , and no edges and let

$$\pi : V_\Gamma \rightarrow pt$$

be the constant map. We have already seen (see §2.3 item 1) that there is a functorial map

$$\pi^* : H^{2k}(pt) \rightarrow H^{2k}(\Gamma, \alpha).$$

However, does there exist a Gysin map

$$\pi_* : H^{2k}(\Gamma, \alpha) \rightarrow H^{2(k-d)}(pt)?$$

Such a map, if it existed, would have to have the following property. Let p be a vertex of Γ and let $j_p : pt \rightarrow V_\Gamma$ be the map $pt \rightarrow p$. Then, by functoriality, π_* would have to satisfy

$$\pi_*(j_p)_* = \text{identity}$$

and, by items 4 and 6 of §2.3, π_* would have to have the form

$$(2.9) \quad \pi_* f = \sum \frac{f(p)}{\prod \alpha_{p,e}}.$$

However, it is by no means obvious that this map is well defined, i.e. that the right hand side of (2.9) is in $S(\mathfrak{g}^*)$. We will prove that it is :

THEOREM 2.2. π_* maps $H^{2k}(\Gamma, \alpha)$ into $S^{k-d}(\mathfrak{g}^*)$.

Proof. Let $f \in H^{2k}(\Gamma, \alpha)$; then $\pi_* f$ can be written as

$$(2.10) \quad \pi_* f = \frac{g}{\prod_{j=1}^N \alpha_j},$$

where $g \in S^{k-d+N}(\mathfrak{g}^*)$ and $\alpha_1, \dots, \alpha_N$ are pair-wise linearly independent. We will show that α_1 divides g .

The vertices of Γ can be divided into two categories:

1. The first subset, V_1 , contains the vertices $p \in V_\Gamma$ for which none of the $\alpha_{p,e}$'s is a multiple of α_1
2. The second subset, V_2 , contains the vertices $p \in V_\Gamma$ for which there exists an edge e such that $\alpha_{p,e}$ is a multiple of α_1 . (Notice that 1.17 implies that there will be exactly one such edge.)

The part of (2.9) corresponding to vertices in the first category will then be of the form

$$(2.11) \quad \sum_{p \in V_1} \frac{f(p)}{\prod \alpha_{p,e}} = \frac{g_1}{\prod_{j=2}^N \alpha_j}$$

with $g_1 \in S(\mathfrak{g}^*)$.

If $p \in V_2$ then there exists an edge e from p such that $\alpha_{p,e} = \lambda \alpha_1$ with $\lambda \in \mathbb{C} - \{0\}$; let q be the other vertex of e . Since $\alpha_{q,e} = -\alpha_{p,e}$ it results that $q \in V_2$ as well and thus the vertices in V_2 can be paired as above.

Let $e_i, i = 1, \dots, d$ and $e'_i, i = 1, \dots, d$ be the edges containing p and q respectively, with $e_d = e'_d = e$. Then the e_i 's can be ordered (cf. 1.18) so that

$$(2.12) \quad \alpha_{p,e_i} \equiv \alpha_{q,e'_i} \pmod{\alpha_1}.$$

Also (1.21) implies that

$$(2.13) \quad f(q) \equiv f(p) \pmod{\alpha_1}.$$

The part of (2.9) corresponding to p and q is given by

$$(2.14) \quad \frac{f(p)}{\prod_{j=1}^d \alpha_{p,e_j}} + \frac{f(q)}{\prod_{j=1}^d \alpha_{q,e'_j}} = \frac{f(p)\alpha_{q,e'_1} \dots \alpha_{q,e'_{d-1}} - f(q)\alpha_{p,e_1} \dots \alpha_{p,e_{d-1}}}{\lambda \alpha_1 \alpha_{p,e_1} \dots \alpha_{q,e'_{d-1}}}.$$

But multiplying together the congruences (2.12) and (2.13) we obtain that α_1 divides the numerator of (2.10) so that

$$(2.15) \quad \frac{f(p)}{\prod_{j=1}^d \alpha_{p,e_j}} + \frac{f(q)}{\prod_{j=1}^d \alpha_{q,e'_j}} = \frac{g_{p,q}}{\prod_{j=2}^N \alpha_j}$$

with $g_{p,q} \in S(\mathfrak{g}^*)$. Therefore

$$(2.16) \quad \sum_{p \in V_2} \frac{f(p)}{\prod \alpha_{p,e}} = \frac{g_2}{\prod_{j=2}^N \alpha_j}$$

with $g_2 \in S(\mathfrak{g}^*)$. Adding (2.11) and (2.16) we get that

$$\frac{g}{\prod_{j=1}^N \alpha_j} = \frac{g_1 + g_2}{\prod_{j=2}^N \alpha_j}$$

with $g_1 + g_2 \in S(\mathfrak{g}^*)$, i.e. that α_1 divides g . The same argument can be used to show that each α_j divides g and therefore $\pi_* f \in S^{k-d}(\mathfrak{g}^*)$, as desired. \square

2.5. The Kirwan map. From now on we will assume that (Γ, α) satisfies the no-cycle condition. Let ξ be an element of \mathcal{P} which gives an orientation of Γ without cycles and let $\phi : V_\Gamma \rightarrow \mathbb{R}$ be positively oriented with respect to ξ ; without loss of generality we can assume that ϕ is injective. For $c \in \mathbb{R} - \phi(V_\Gamma)$, we define the *c-cross section*, Γ_c , of Γ to be the set of edges, e , of Γ with the property that, for one of the vertices, p , of e , $\phi(p) > c$ and for the other vertex, q , $\phi(q) < c$. Let \mathfrak{g}_ξ^* be the annihilator of ξ in \mathfrak{g}^* and let $H^{2k}(\Gamma_c, \alpha)$ be the set of all maps

$$f : \Gamma_c \rightarrow S^k(\mathfrak{g}_\xi^*).$$

The sum

$$H(\Gamma_c, \alpha) = \bigoplus H^{2k}(\Gamma_c, \alpha)$$

is a graded ring under point-wise multiplication and we will define a morphism of graded rings

$$\mathcal{K}_c : H(\Gamma, \alpha) \rightarrow H(\Gamma_c, \alpha)$$

as follows: For $e \in \Gamma_c$ let p and q be the vertices of e . The projection $\mathfrak{g}^* \rightarrow \mathfrak{g}_e^*$ maps \mathfrak{g}_ξ^* bijectively onto \mathfrak{g}_e^* since $\alpha_{p,e}(\xi) \neq 0$, so one has a composite map

$$\mathfrak{g}^* \rightarrow \mathfrak{g}_e^* \leftrightarrow \mathfrak{g}_\xi^*$$

and hence an induced morphism of graded rings:

$$S(\mathfrak{g}^*) \rightarrow S(\mathfrak{g}_e^*) \leftrightarrow S(\mathfrak{g}_\xi^*).$$

If f is in $H(\Gamma, \alpha)$, the images of f_p and f_q in $S(\mathfrak{g}_e^*)$ are the same by (1.21) and hence so are their images in $S(\mathfrak{g}_\xi^*)$. We define $\mathcal{K}_c(f)(e)$ to be this common image and call the map \mathcal{K}_c *the Kirwan map*.

Next we will define a morphism, γ_c , of $S(\mathfrak{g}_\xi^*)$ -modules, mapping $H(\Gamma_c, \alpha)$ into the quotient field of $S(\mathfrak{g}_\xi^*)$. To define γ_c , let e , as above, be an element of Γ_c and let p and q be the vertices of e . We will assume that $\phi(p) > c$ and $\phi(q) < c$ and, hence, that $\alpha_{q,e}(\xi) > 0$. Let

$$e_i^+, \quad i = 1, \dots, d-1$$

be the other edges of Γ (other than e) intersecting at p and

$$e_i^-, \quad i = 1, \dots, d-1$$

be the other edges of Γ intersecting at q . By the compatibility axiom we can assume that α_{p,e_i^+} and α_{q,e_i^-} have the same image in \mathfrak{g}_e^* and, hence, under the identification, $\mathfrak{g}_e^* \leftrightarrow \mathfrak{g}_\xi^*$, have the same image in \mathfrak{g}_ξ^* . This implies that

$$\alpha_{i,e}^\# =: \alpha_{p,e_i^+} - m_{i,e}^+ \alpha_{p,e} = \alpha_{q,e_i^-} - m_{i,e}^- \alpha_{q,e},$$

where

$$m_{i,e}^+ = \frac{\alpha_{p,e_i^+}(\xi)}{\alpha_{p,e}(\xi)} \quad \text{and} \quad m_{i,e}^- = \frac{\alpha_{q,e_i^-}(\xi)}{\alpha_{q,e}(\xi)}.$$

Let

$$m_e = \alpha_{q,e}(\xi) = -\alpha_{p,e}(\xi)$$

and note that, since $\phi(p) > \phi(q)$, $m_e = |m_e| > 0$.

We now define, for $f \in H^{2k}(\Gamma_c, \alpha)$,

$$(2.17) \quad \gamma_c f = \sum_{e \in \Gamma_c} \frac{1}{m_e} \frac{f(e)}{\prod_i \alpha_{i,e}^\#}$$

and we define \mathfrak{p}_c to be the composition

$$(2.18) \quad \mathfrak{p}_c = \gamma_c \mathcal{K}_c.$$

THEOREM 2.3. \mathfrak{p}_c is a map of $H^{2k}(\Gamma, \alpha)$ into $S^{k-d+1}(\mathfrak{g}_\xi^*)$.

We will prove this by obtaining an explicit “residue formula” for $\mathfrak{p}_c(f)$, $f \in H^{2k}(\Gamma, \alpha)$. This residue formula can be viewed as a kind of combinatorial version of the Jeffrey-Kirwan theorem described in §1.3. It is also closely related to the localization theorem proved by Jaap Kalkman and the first author in [GK], and the residue results which we will describe in the next section are mostly taken from [GK].

2.6. Residues. Let $\alpha_1, \dots, \alpha_d$ be elements of \mathfrak{g}^* and ξ be an element of \mathfrak{g} with the property that $\alpha_i(\xi) \neq 0$ for all i . Given $f \in S^k(\mathfrak{g}^*)$ we define an element

$$(2.19) \quad \text{Res}_\xi \frac{f}{\prod \alpha_i}$$

of $S^{k-d+1}(\mathfrak{g}_\xi^*)$ as follows: Choose a basis x, y_1, \dots, y_{n-1} of \mathfrak{g}^* such that y_1, \dots, y_{n-1} is a basis of \mathfrak{g}_ξ^* and $x(\xi) = 1$. Let

$$(2.20) \quad \alpha_i = m_i x - \sum_{j=1}^{n-1} a_{ij} y_j$$

and let

$$f(x) = \sum_{r=0}^k x^r f_r(y)$$

and

$$(2.21) \quad \beta_i = m_i^{-1} \sum a_{ij} y_j;$$

then

$$\frac{f}{\alpha_1 \dots \alpha_d} = (\prod m_i)^{-1} x^{-d} \sum x^r f_r \left(1 - \frac{\beta_1}{x}\right)^{-1} \dots \left(1 - \frac{\beta_d}{x}\right)^{-1}.$$

Now replace $(1 - \frac{\beta_i}{x})^{-1}$ by the power series

$$\sum_{k=0}^{\infty} x^{-k} \beta_i^k$$

and define (2.19) to be the coefficient of x^{-1} in the product on the right. It is easy to see that this definition doesn't depend on the choice of x, y_1, \dots, y_n . If the α_i 's are pairwise linearly independent (i.e., if, for $i \neq j$, α_i and α_j are not multiples of each other) there is a relatively simple formula for (2.19).

LEMMA 1. *Let A be a graded commutative algebra over \mathbb{C} and let $f = f(x)$ be a polynomial in x with coefficients in A . Then for indeterminants z_1, \dots, z_d*

$$(2.22) \quad \text{Res}_x \frac{f(x)}{(x - z_1) \cdots (x - z_d)} = \sum_{i=1}^d \frac{f(z_i)}{\prod_{j \neq i} (z_i - z_j)}.$$

Proof. The conclusion follows immediately from the decomposition in simple fractions

$$\frac{f(x)}{(x - z_1) \cdots (x - z_d)} = F(x) + \sum_{i=1}^d \frac{f(z_i)}{\prod_{j \neq i} (z_i - z_j)} \frac{1}{x - z_i}$$

where $F(x)$ is a polynomial term in x . \square

Let

$$h = \frac{f}{\prod_{j=1}^d (x - z_j)} \quad \text{and} \quad h_j = \frac{f(z_j)}{\prod_{r \neq j} (z_j - z_r)}, \quad \forall j.$$

LEMMA 2. *$h \in A[x]$ if and only if $\text{Res}_x(x^k h) = 0, \forall k \geq 0$.*

Proof. From (2.22) we get that

$$\text{Res}_x(x^k h) = \sum_{j=1}^d (z_j)^k h_j.$$

Then the fact that $\text{Res}_x(x^k h) = 0, \forall k = 1, \dots, d$ can be written as

$$\begin{pmatrix} z_1^1 & \cdots & z_j^1 & \cdots & z_d^1 \\ \vdots & & \vdots & & \vdots \\ z_1^k & \cdots & z_j^k & \cdots & z_d^k \\ \vdots & & \vdots & & \vdots \\ z_1^d & \cdots & z_j^d & \cdots & z_d^d \end{pmatrix} \begin{pmatrix} h_1 \\ \vdots \\ h_j \\ \vdots \\ h_d \end{pmatrix} = 0.$$

Since the corresponding Vandermonde determinant is non-zero we deduce that $h_1 = \cdots = h_d = 0$, i.e. $f(z_j) = 0, \forall j = 1, \dots, d$, from which we obtain that $h \in A[x]$; the other implication is clear. \square

We will now apply lemma 1 to the evaluation of (2.19). Let

$$m_i = \alpha_i(\xi)$$

and for $i \neq j$ let

$$\alpha_{j,i}^\# = \alpha_j - m_{j,i}\alpha_i$$

with

$$m_{j,i} = \frac{\alpha_j(\xi)}{m_i}.$$

Note that $\alpha_{j,i}^\# \in \mathfrak{g}_\xi^*$ since $\alpha_{j,i}^\#(\xi) = \alpha_j(\xi) - \alpha_j(\xi) = 0$. Let

$$\mathfrak{g}_i^* = \mathfrak{g}^* / \{c\alpha_i; c \in \mathbb{R}\}.$$

The projection map: $\mathfrak{g}^* \rightarrow \mathfrak{g}_i^*$ is bijective on \mathfrak{g}_ξ^* , so one gets a composite map

$$\mathfrak{g}^* \rightarrow \mathfrak{g}_i^* \xrightarrow{\cong} \mathfrak{g}_\xi^*$$

as in §2.5 and hence a ring morphism

$$\mathcal{K}_i : S(\mathfrak{g}^*) \rightarrow S(\mathfrak{g}_\xi^*).$$

THEOREM 2.4. For $f \in S^k(\mathfrak{g}^*)$

$$(2.23) \quad \text{Res}_\xi \frac{f}{\alpha_1 \dots \alpha_d} = \sum_i \frac{1}{m_i} \frac{\mathcal{K}_i f}{\prod_{j \neq i} \alpha_{j,i}^\#}.$$

Proof. With the notations (2.20)-(2.21)

$$\frac{f}{\prod \alpha_i} = \left(\prod m_k \right)^{-1} \frac{f(x, y)}{\prod (x - \beta_k(y))}.$$

Thus by Lemma 1

$$\begin{aligned} \text{Res}_\xi \frac{f}{\prod \alpha_i} &= \left(\prod m_k \right)^{-1} \sum_i \frac{f(\beta_i, y)}{\prod_{k \neq i} (\beta_i - \beta_k)} \\ &= \sum_i \frac{1}{m_i} \frac{f(\beta_i, y)}{\prod_{k \neq i} m_k (\beta_i - \beta_k)}. \end{aligned}$$

But $m_k(\beta_i - \beta_k) = \alpha_{k,i}^\#$ and the map

$$\mathcal{K}_i : S(\mathfrak{g}^*) \rightarrow S(\mathfrak{g}_\xi^*)$$

maps x to β_i and y_k to itself, so $\mathcal{K}_i f(x, y) = f(\beta_i, y)$. Thus the sum on the right is identical with the right hand side of (2.22). \square

2.7. The Jeffrey-Kirwan theorem. We will prove Theorem 2.3 by deducing it from the following result:

THEOREM 2.5. For $f \in H^{2k}(\Gamma, \alpha)$

$$(2.24) \quad \mathfrak{p}_c(f) = \sum_{\phi(p) < c} \text{Res}_\xi \frac{f_p}{\prod \alpha_{p,e}}.$$

In particular, $\mathfrak{p}_c(f)$ is in $S^{k-d+1}(\mathfrak{g}_\xi^*)$.

Proof. Choose $c = c_0 > c_1 > c_2 > \dots > c_N$ in $\mathbb{R} - \phi(V)$ so that $\phi(V) \subset (c_N, \infty)$ and $\forall r \geq 0$ there is exactly one vertex p_r with $\phi(p_r) \in (c_{r+1}, c_r)$. Inspection of (2.18) shows that

$$(2.25) \quad \mathfrak{p}_{c_r}(f) - \mathfrak{p}_{c_{r+1}}(f) = \sum \frac{1}{m_i} \frac{\mathcal{K}_i f}{\prod_{j \neq i} \alpha_{j,i}^\#},$$

where $e_i, i = 1, \dots, d$, are the edges of Γ containing p_r and $\alpha_i = \alpha_{p_r, e_i}$. On the other hand, by Theorem 2.4, the right hand side of (2.25) is just

$$\text{Res}_\xi \frac{f(p_r)}{\prod \alpha_{p_r, e}}.$$

The conclusion follows since $\mathfrak{p}_{c_N}(f) = 0$. \square

COROLLARY 2.1. If π_* is the map given by (2.9) then, for $f \in H^{2k}(\Gamma, \alpha)$,

$$(2.26) \quad \text{Res}_\xi(\pi_* f) = 0.$$

We conclude this section by observing that combining corollary 2.1 and lemma 2 we obtain a new proof of theorem 2.2 for graphs that satisfy the no-cycle condition:

Let $f \in H^{2k}(\Gamma, \alpha)$. Then, as in (2.10),

$$\pi_* f = \frac{g}{\prod_{j=1}^N \alpha_j},$$

where $g \in S^{k-d+N}(\mathfrak{g}^*)$ and $\alpha_1, \dots, \alpha_N$ are pair-wise linearly independent.

Let ξ generate an orientation of Γ with no cycles and choose $\theta \in \mathfrak{g}^*$ such that $\theta(\xi) = 1$ and θ is not equal to any of $\alpha_1, \dots, \alpha_N$. Then $\theta^r f \in H^{2(k+r)}(\Gamma, \alpha)$ and

$$\pi_*(\theta^r f) = \frac{\theta^r g}{\prod_{i=1}^N \alpha_i}.$$

But (2.26) implies that $\text{Res}_\xi(\pi_*(\theta^r f)) = 0 \forall r \geq 0$ and it follows now from lemma 2 that $\pi_* f \in S^{k-d}(\mathfrak{g}^*)$.

2.8. The Betti numbers of the pair (Γ, α) . For $\xi \in \mathcal{P}$ and $p \in V_\Gamma$, let $\sigma_p = \sigma_p(\xi)$ be the number of edges e , with one vertex p , for which $\alpha_{p,e}(\xi) < 0$. Let β_k be the number of vertices p with $\sigma_p = k$. Since σ_p depends on ξ , it is surprising to find that these ‘‘Betti numbers’’ don’t.

THEOREM 2.6. *The β_k 's don't depend on ξ ; i.e. they are combinatorial invariants of (Γ, α) .*

Proof. Let \mathcal{P}_i , $i = 1, \dots, N$, be the connected components of \mathcal{P} and consider an $(n - 1)$ -dimensional wall separating two adjacent \mathcal{P}_i 's. This wall is defined by an equation of the form

$$(2.27) \quad \alpha_{p,e}(\xi) = 0$$

for some $(p, e) \in I_\Gamma$. Let q be the other vertex of e and lets compute the changes in σ_p and σ_q as ξ passes through this wall: Let e_i , $i = 1, \dots, d$ be the edges meeting at p and e'_i , $i = 1, \dots, d$ be the edges meeting at q (with $e_d = e'_d = e$). By (1.18) we can order the e_i 's so that, for $i \leq d - 1$,

$$\alpha_{p,e_i} = \alpha_{q,e'_i} + c_i \alpha_{p,e}.$$

From (1.17) follows that for every $i = 1, \dots, d - 1$,

$$\dim(\ker \alpha_{p,e} \cap \ker \alpha_{p,e_i}) = n - 2.$$

Therefore there exists ξ_0 such that $\alpha_{p,e}(\xi_0) = 0$ but $\alpha_{p,e_i}(\xi_0) = \alpha_{q,e'_i}(\xi_0) \neq 0$, for all $i = 1, \dots, d - 1$.

Then there exists a neighborhood U of ξ_0 in \mathfrak{g} such that for all $i = 1, \dots, d - 1$ and $\xi \in U$, $\alpha_{p,e_i}(\xi)$ and $\alpha_{q,e'_i}(\xi)$ have the same sign and this common sign doesn't depend on $\xi \in U$. Such a neighborhood will intersect both regions created by the wall (2.27). Now suppose that $\xi \in U$ and that r of the numbers $\alpha_{p,e_i}(\xi)$, $i = 1, \dots, d - 1$, are negative. Since $\alpha_{p,e}(\xi) = -\alpha_{q,e}(\xi)$, it follows that for $\alpha_{p,e}(\xi)$ positive

$$\sigma_p = r \quad \text{and} \quad \sigma_q = r + 1$$

and for $\alpha_{p,e}(\xi)$ negative

$$\sigma_p = r + 1 \quad \text{and} \quad \sigma_q = r.$$

In either case, as ξ passes through the wall (2.27), the Betti numbers don't change. \square

(For this simple and beautiful proof of the well-definedness of the Betti numbers we are indebted to Ethan Bolker.)

2.9. Betti numbers and cohomology. Simple examples show that the formula (1.35) is not true for an arbitrary graph-axial function pair (Γ, α) . However, we will prove that if (Γ, α) has the no-cycle property for some $\xi \in \mathcal{P}$ then the equality (1.35) can be replaced by the inequality

$$(2.28) \quad \dim H^{2k}(\Gamma, \alpha) \leq \sum \beta_r \dim S^{k-r}(\mathfrak{g}^*).$$

In addition we will show that, for k large,

$$(2.29) \quad \dim H^{2k}(\Gamma, \alpha) = \sum \beta_r \dim S^{k-r}(\mathfrak{g}^*) + O(k^{n-3}).$$

(Note that since

$$\dim S^k(\mathfrak{g}^*) = \binom{k+n-1}{n-1} = \frac{1}{(n-1)!} \left(k^{n-1} + \binom{n}{2} k^{n-2} + O(k^{n-3}) \right),$$

the first term on the right hand side is strictly greater than the error term.) In particular, if $n = 2$, the formula (2.29) asserts that

$$(2.30) \quad \dim H^{2k}(\Gamma, \alpha) = \sum \beta_r \dim S^{k-r}(\mathfrak{g}^*)$$

for all k greater than some fixed k_0 .

Proof. Let $\alpha_i \in \mathfrak{g}^*$, $i = 1, \dots, N$, be a pairwise linearly independent set of vectors with the property that every one of the vectors $\alpha_{p,e}$, $(p, e) \in I_\Gamma$, is a multiple of a vector in this set. Let I be the graded ideal in $S(\mathfrak{g}^*)$ generated by the monomials

$$g_i = \alpha_1 \cdots \widehat{\alpha}_i \cdots \alpha_N.$$

LEMMA 3. *The algebraic dimension of the quotient ring $S(\mathfrak{g}^*)/I$ is $n - 2$.*

Proof. This follows trivially from the fact that the algebraic variety defined by I is the union of the sets $\alpha_i = \alpha_j = 0$, $i \neq j$. \square

As a corollary of this lemma we get the bound

$$(2.31) \quad \dim S^k(\mathfrak{g}^*)/I^k = O(k^{n-3}).$$

Now let $\phi : V_\Gamma \rightarrow \mathbb{R}$ be a function which is positively oriented with respect to ξ and let $H_c(\Gamma, \alpha)$ be the subring of $H(\Gamma, \alpha)$ consisting of all maps $f : V_\Gamma \rightarrow S(\mathfrak{g}^*)$ with support on the set $\phi \geq c$. Let $p \in V_\Gamma$ with $\phi(p) = c$ and suppose that there are no points $q \in V_\Gamma$ with $\phi(q)$ on the interval (c, c') . Letting $\sigma_p = r$ we will prove the Morse inequality

$$(2.32) \quad \dim H_c^{2k}(\Gamma, \alpha) - \dim H_{c'}^{2k}(\Gamma, \alpha) \leq \dim S^{k-r}(\mathfrak{g}^*)$$

and an inequality in the opposite direction:

$$(2.33) \quad \dim H_c^{2k}(\Gamma, \alpha) - \dim H_{c'}^{2k}(\Gamma, \alpha) \geq \dim I^{k-r}.$$

To prove (2.32) let e_i , $i = 1, \dots, d$, be the edges of Γ containing p and let $\alpha_i = \alpha_{p, e_i}$. We will order the α_i 's so that $\alpha_i(\xi) < 0$ for $1 \leq i \leq r$ and $\alpha_i(\xi) > 0$ for $r+1 \leq i \leq d$. Then if $f \in H_c^{2k}(\Gamma, \alpha)$, $f(p)$ must be a multiple of $\alpha_1, \dots, \alpha_r$ so the image of the restriction map

$$(2.34) \quad H_c^{2k}(\Gamma, \alpha) \rightarrow S^k(\mathfrak{g}^*), \quad f \rightarrow f(p),$$

is contained in $S^{k-r}(\mathfrak{g}^*)\alpha_1 \cdots \alpha_r$. Since the kernel of this map is $H_{c'}^{2k}(\Gamma, \alpha)$, this proves (2.32). We will prove the inequality (2.33) by showing that if $h \in I^{k-r}$ then $h\alpha_1 \cdots \alpha_r$ is in the image of (2.34). Indeed, if $h \in I^{k-r}$ then $h\alpha_1 \cdots \alpha_r$ can be written as a sum

$$\sum_{i=r+1}^N h_i \alpha_1 \cdots \widehat{\alpha}_i \cdots \alpha_N.$$

Let p_j be the vertex joined to p by e_j for $j = r+1, \dots, d$, and, for fixed $j_0 \in \{r+1, \dots, d\}$, define $f : V_\Gamma \rightarrow S^k(\mathfrak{g}^*)$ to be the map which takes the value $h\alpha_1 \cdots \alpha_r$

at p , the value $h_{j_0}\alpha_1 \cdots \widehat{\alpha_{j_0}} \cdots \alpha_N$ at p_{j_0} and zero elsewhere. It is easily checked that $f \in H_c^{2k}(\Gamma, \alpha)$ and $f(p) = h\alpha_1 \cdots \alpha_r$. This proves (2.33).

Next let c and c' be any pair of real numbers with $c < c'$. From (2.32)-(2.33) one gets, by a simple induction, the Morse inequalities

$$\dim H_c^{2k}(\Gamma, \alpha) - \dim H_{c'}^{2k}(\Gamma, \alpha) \leq \sum \beta_r(c, c') \dim S^{k-r}(\mathfrak{g}^*)$$

and

$$\dim H_c^{2k}(\Gamma, \alpha) - \dim H_{c'}^{2k}(\Gamma, \alpha) \geq \sum \beta_r(c, c') \dim I^{k-r},$$

where $\beta_r(c, c')$ is the number of points $p \in V_\Gamma$ with $\sigma_p = r$ and $c \leq \phi(p) < c'$. In particular, for $c' \gg 0$ and $c \ll 0$, one gets from these estimates and from (2.31) the inequalities (2.28) and (2.29). \square

2.10. The role of the zeroth Betti number. An example of a graph-axial function pair that fails to satisfy (1.35) is the $d = n = 2$ example described at the end of §2.1. The graph in this example is a connected graph; so its topological zeroth Betti number is 1. However, its graph theoretical zeroth Betti number, β_0 , is N . A simple computation shows that, for this example, the identity (1.35) holds for all $k > 0$. But for $k = 0$ the left hand side of (1.35) is 1 (since the graph is connected) whereas the right hand side, β_0 , is N . From this example one can generate examples of graph-axial function pairs, (Γ, α) , for which the estimate (2.29) is “best possible” by taking Cartesian products of this graph with graphs which do satisfy (1.35).

However, by making some additional assumptions on the pair (Γ, α) one can considerably improve (2.29). The assumptions we will make are of two kinds:

1. To avoid the problem posed by the example we have just described, we will assume that the zeroth Betti numbers of certain connected subgraphs of Γ are equal to 1.
2. For every $p \in V_\Gamma$ we will make certain “general position” hypothesis about the vectors $\alpha_{p,e}$, $(p, e) \in I_\Gamma$. To formulate these hypothesis we introduce the following refinement of the notion of “pairwise linearly independent”:

DEFINITION. A collection of vectors $\alpha_i \in \mathfrak{g}^*$, $i = 1, \dots, N$, is *l-independent* if, for every sequence $1 \leq i_1 < i_2 < \dots < i_l \leq N$, the vectors $\alpha_{i_1}, \dots, \alpha_{i_l}$ are linearly independent.

Now let (Γ, α) be a graph-axial function pair which satisfies the no-cycle condition for some $\xi \in \mathcal{P}$. The main result of this section is the following sharpening of (2.29):

THEOREM 2.7. *Suppose the following hypotheses hold:*

1. *For every subspace \mathfrak{h} of \mathfrak{g} of codimension strictly less than l , the zeroth Betti number of the connected components of $\Gamma_{\mathfrak{h}}$ are equal to 1.*
2. *For every vertex p of Γ , the vectors $\alpha_{p,e}$, $(p, e) \in I_\Gamma$, are l -independent.*

Then:

$$(2.35) \quad \dim H^{2k}(\Gamma, \alpha) = \sum \beta_r \dim S^{k-r}(\mathfrak{g}^*) + O(k^{n-1-l}).$$

REMARK. For $l = 2$ the above conditions are always satisfied; (2.29) is the particular case of (2.35) corresponding to $l = 2$.

For $l = n$ this theorem says that the left hand side of (2.35) is equal to the first term on the right for k greater than some fixed k_0 . This result can be slightly improved.

THEOREM 2.8. *If the hypotheses of theorem 2.7 hold for $l = n$ then*

$$(2.36) \quad \dim H^{2k}(\Gamma, \alpha) = \sum \beta_r \dim S^{k-r}(\mathfrak{g}^*)$$

for $k > d - n$.

We will prove these two results by refining the Morse inequalities (2.33). For this we will need the following generalization of Lemma 3 of §2.9:

LEMMA 4. *Let $\gamma_1, \dots, \gamma_N$ be a collection of vectors in \mathfrak{g}^* which are l -independent and let I_l be the ideal in $S(\mathfrak{g}^*)$ generated by the monomials*

$$(2.37) \quad \frac{\gamma_1 \cdots \gamma_N}{\gamma_{i_1} \cdots \gamma_{i_{l-1}}}, \quad 1 \leq i_1 < \cdots < i_{l-1} \leq N.$$

Then the algebraic dimension of $S(\mathfrak{g}^)/I_l$ is $n - l$. Moreover, if $N \geq n$ and $n = l$ then $S^m(\mathfrak{g}^*) = I_l^m$ for $m > N - n$.*

(For the proof of this lemma see the appendix at the end of this section.)

Proof of theorem 2.7. Let $\phi : V_\Gamma \rightarrow \mathbb{R}$ be a strictly monotone function which is positively oriented with respect to ξ . Let p be an arbitrary vertex of Γ , let $c = \phi(p)$ and assume that there are no points $q \in V_\Gamma$ with $c < \phi(q) < c'$. Let e_1, \dots, e_d be the edges of Γ containing p and let $\alpha_i = \alpha_{p, e_i}$. We can order these vectors so that $\alpha_i(\xi) < 0$ for $1 \leq i \leq r$ and $\alpha_i(\xi) > 0$ for $r + 1 \leq i \leq d$, where $r = \sigma_p$. Let $N = d - r$.

Suppose $l \leq N$. We will prove that

$$(2.38) \quad \dim H_c^{2k}(\Gamma, \alpha) - \dim H_{c'}^{2k}(\Gamma, \alpha) \geq \dim I_l^{k-r},$$

where I_l is the ideal of $S(\mathfrak{g}^*)$ constructed as in lemma 4 using the $N = d - r$ vectors $\alpha_{r+1}, \dots, \alpha_d$ (which are l -independent, by hypothesis 2), i.e. I_l is the ideal in $S(\mathfrak{g}^*)$ generated by the monomials

$$\frac{\alpha_{r+1} \cdots \alpha_d}{\alpha_{i_1} \cdots \alpha_{i_{l-1}}}, \quad r + 1 \leq i_1 < \dots < i_{l-1} \leq N.$$

To show (2.38), consider (as in §2.9) the restriction map

$$(2.39) \quad H_c^{2k}(\Gamma, \alpha) \longrightarrow S^{k-r}(\mathfrak{g}^*) \alpha_1 \cdots \alpha_r$$

given by $f \rightarrow f(p)$. The kernel of this map is $H_{c'}^{2k}(\Gamma, \alpha)$; so, to prove (2.38) it suffices to show that the image of this map contains $I_l^{k-r} \alpha_1 \cdots \alpha_r$. Consider the set of vectors $\alpha_{i_1}, \dots, \alpha_{i_{l-1}}$, $r + 1 \leq i_1 < \dots < i_{l-1} \leq d$. These vectors are linearly independent; so

their annihilator, \mathfrak{h} , is of codimension $l - 1$ in \mathfrak{g} . Let Γ_1 be the connected component of $\Gamma_{\mathfrak{h}}$ containing p . Since the numbers $\alpha_{i_1}(\xi), \dots, \alpha_{i_{l-1}}(\xi)$ are greater than zero, p is a local minimum point for the restriction of ϕ to $\Gamma_{\mathfrak{h}}$; so, by hypothesis 1, p is also a global minimum. Therefore the vertices of Γ_1 are contained in the set $\phi \geq c$, and, hence the Thom class, τ_1 , of Γ_1 , is supported in this set, i.e. is an element of $H_c(\Gamma, \alpha)$. However, at p , τ_1 is equal to

$$\left(\frac{\alpha_{r+1} \cdots \alpha_d}{\alpha_{i_1} \cdots \alpha_{i_{l-1}}} \right) \alpha_1 \cdots \alpha_r.$$

This argument shows that, for all generators, f , of I_l , $f\alpha_1 \cdots \alpha_r$ is in the image of the restriction map (2.39). Hence the image of this restriction map contains $I_l^{k-r}\alpha_1 \cdots \alpha_r$, which proves (2.38).

Now suppose that $l > N$. In this case we can simply take \mathfrak{h} to be the annihilator of $\alpha_{r+1}, \dots, \alpha_d$, and, by the same argument as above, conclude that the image of the restriction map is equal to $S^{k-r}(\mathfrak{g}^*)\alpha_1 \cdots \alpha_r$. Hence for $l > N$,

$$(2.40) \quad \dim H_c^{2k}(\Gamma, \alpha) - \dim H_c^{2k}(\Gamma, \alpha) = \dim S^{k-r}(\mathfrak{g}^*).$$

The proof of the estimate (2.35) via (2.38), (2.40) and lemma 4 is the same as the proof of the estimate (2.29) via (2.33) and lemma 3. We will omit details. \square

Proof of theorem 2.8. Let $l = n$. If $N < n$ the equality (2.40) holds for all k (as we have just seen). If $N \geq n$ then, by lemma 4 and by (2.38), the equality (2.40) holds if $k - r > N - n$, i.e. if $k > d - n$. Thus for $l = n$, (2.35) can be sharpened to (2.36). \square

Appendix: The proof of Lemma 4. We will prove by induction that the algebraic variety defined by I_l is the union of the sets

$$\gamma_{i_1} = \cdots = \gamma_{i_l} = 0, \quad 1 \leq i_1 < \cdots < i_l \leq N.$$

Let x be a point on this variety. Since $I_{l-1} \subset I_l$, the variety defined by I_{l-1} contains the variety defined by I_l ; so it follows by induction that there exists a sequence $1 \leq i_1 < \cdots < i_{l-1} \leq N$ with $\gamma_{i_1} = \cdots = \gamma_{i_{l-1}} = 0$ at x . However, since I_l contains the quotient of $\gamma_1 \cdots \gamma_N$ by $\gamma_{i_1} \cdots \gamma_{i_{l-1}}$, there exists some $j \neq i_1, \dots, i_{l-1}$ such that $\gamma_j = 0$ at x . This proves the first assertion of lemma 4. Now let $l = n$. We will prove that

$$S^m(\mathfrak{g}^*) = I_n^m$$

for $m > N - n$ by a double induction on n and N .

The equality above is true if $N = n$ or if $n = 1$, as can be easily checked. Consider now a pair (N, n) with $N > n$. We now assume that the assertion is true for $(N - 1, n - 1)$ and for $(N - 1, n)$.

Let \mathfrak{h} be the annihilator of γ_N in \mathfrak{g} . The restriction map

$$S(\mathfrak{g}^*) \longrightarrow S(\mathfrak{h}^*)$$

maps $\gamma_1, \dots, \gamma_{N-1}$ onto vectors $\beta_1, \dots, \beta_{N-1}$, which are $(n - 1)$ -independent in \mathfrak{h}^* ; so, by induction, every element of $S^m(\mathfrak{h}^*)$, $m > N - n$, is in the ideal generated by the

monomials

$$\frac{\beta_1 \cdots \beta_{N-1}}{\beta_{i_1} \cdots \beta_{i_{n-2}}}, \quad 1 \leq i_1 < \dots < i_{n-2} \leq N-1.$$

Since the kernel of the map $S(\mathfrak{g}^*) \rightarrow S(\mathfrak{h}^*)$ is the ideal generated by γ_N , it follows that for $m > N - n$, every element of $S^m(\mathfrak{g}^*)$ can be written as a linear combination of

$$\frac{\gamma_1 \cdots \gamma_{N-1} \gamma_N}{\gamma_{i_1} \cdots \gamma_{i_{n-2}} \gamma_N}, \quad 1 \leq i_1 < \dots < i_{n-2} \leq N-1,$$

with polynomial coefficients, plus a term of the form $f\gamma_N$, $f \in S^{m-1}(\mathfrak{g}^*)$. By induction the theorem is true in dimension n for the vectors $\gamma_1, \dots, \gamma_{N-1}$. Then f is in the ideal generated by the monomials

$$\frac{\gamma_1 \cdots \gamma_{N-1}}{\gamma_{i_1} \cdots \gamma_{i_{n-1}}}, \quad 1 \leq i_1 < \dots < i_{n-1} \leq N-1$$

and hence $f\gamma_N$ is in the ideal generated by

$$\frac{\gamma_1 \cdots \gamma_N}{\gamma_{i_1} \cdots \gamma_{i_{n-1}}}, \quad 1 \leq i_1 < \dots < i_{n-1} \leq N-1.$$

REFERENCES

- [A] M. F. ATIYAH, *Convexity and commuting Hamiltonians*, Bull. London Math. Soc., 14 (1982), pp. 1–15.
- [AB1] M. F. ATIYAH AND R. BOTT, *The moment map and equivariant cohomology*, Topology, 23 (1984), pp. 1–28.
- [AB2] M. F. ATIYAH AND R. BOTT, *A Lefschetz fixed point formula for elliptic complexes, I*, Ann. of Math., 86 (1967), pp. 374–407.
- [BV] N. BERLINE AND M. VERGNE, *Classes caractéristiques équivariantes*, C.R. Acad. Sci., Paris, 295 (1982), pp. 539–541.
- [Br] M. BRION, *Equivariant cohomology and equivariant intersection theory*, in Representation Theory and Algebraic Geometry, Kluwer Acad. Publ., Dordrecht, 1998, pp. 1–37.
- [CS] F. R. K. CHUNG AND S. STERNBERG, *Laplacian and Vibrational Spectra for Homogeneous Graphs*, J. of Graph Theory, 16 (1992), pp. 605–627.
- [GK] V. GUILLEMIN AND J. KALKMAN, *The Jeffrey-Kirwan localization theorem and residue operations in equivariant cohomology*, Journal reine angew. Math., 470 (1996), pp. 123–140.
- [GKM] M. GORESKY, R. KOTTWITZ, AND R. MACPHERSON, *Equivariant cohomology, Koszul duality and the localization theorem*, Invent. Math., 131 (1998), pp. 25–83.
- [GS] V. GUILLEMIN AND S. STERNBERG, *Convexity properties of the moment mapping*, Invent. Math., 67 (1982), pp. 491–513.
- [Gu] V. GUILLEMIN, *Deformations of a Hamiltonian action of a compact Lie group*, in Integrable systems (Luminy 1991), Progress in Math., 115, Birkhäuser, Boston, 1993, pp. 227–233.
- [JK] L. JEFFREY AND F. KIRWAN, *Localization for non-abelian group actions*, Topology, 34 (1995), pp. 291–327.
- [Kl] A. A. KLYACHKO, *Equivariant bundles on toric varieties*, Math USSR Izvestiya, 35:2 (1990), pp. 337–375.