

HAAR-TYPE MULTI-WAVELET BASES AND SELF-AFFINE MULTI-TILES*

TIM FLAHERTY[†] AND YANG WANG[‡]

Abstract. Gröchenig and Madych showed that a Haar-type wavelet basis of $L^2(\mathbf{R}^n)$ can be constructed from the characteristic function χ_Ω of a compact set Ω if and only if Ω is an integral self-affine tile of Lebesgue measure one. In this paper we generalize their result to the multiwavelet settings. We give a complete characterization of *Haar-type* scaling function vectors $\chi_\Omega(x) := [\chi_{\Omega_1}(x), \dots, \chi_{\Omega_r}(x)]^T$, where $\Omega = (\Omega_1, \dots, \Omega_r)$ is an r -tuple of compact sets in \mathbf{R}^n . We call Ω a *self-affine multi-tile* because Ω_i 's tile \mathbf{R}^n by translation and have the property that each affine image $A(\Omega_i)$ is the union of translates of some Ω_j 's. We also construct associated *Haar-type multiwavelets*, and present examples using various dilation matrices A .

1. Introduction. Let A be an expanding matrix in $M_n(\mathbf{Z})$, that is, one with integer entries and all eigenvalues $|\lambda_i(A)| > 1$. A compactly supported nonzero function $f(x) \in L^2(\mathbf{R}^n)$ is called a *scaling function* of a multiresolution analysis with dilation factor A if it has the following properties:

(i) $f(x)$ satisfies a refinement equation

$$(1.1) \quad f(x) = \sum_{\alpha \in \mathbf{Z}^n} c_\alpha f(Ax - \alpha),$$

where the coefficients c_α are real and $\sum_{\alpha \in \mathbf{Z}^n} c_\alpha = |\det(A)|$.

(ii) The integer translates $f(x - \alpha)$, $\alpha \in \mathbf{Z}^n$, are orthogonal in $L^2(\mathbf{R}^n)$.

It is well known that given any scaling function of a multiresolution analysis, an orthonormal wavelet basis can be constructed from that scaling function, see [5].

In [13] Gröchenig and Madych studied Haar-type scaling functions and wavelet bases. A *Haar-type* scaling function is a scaling function of the form χ_Ω , where Ω is a compact set in \mathbf{R}^n , and a Haar-type wavelet basis is the wavelet basis constructed from a Haar-type scaling function. Gröchenig and Madych showed that for a given expanding matrix $A \in M_n(\mathbf{R})$ the function χ_Ω is a scaling function of a multiresolution analysis with dilation A if and only if Ω satisfies the following conditions:

(i) Ω is a \mathbf{Z}^n -tile of \mathbf{R}^n , i.e.,

$$\bigcup_{\alpha \in \mathbf{Z}^n} (\Omega + \alpha) = \mathbf{R}^n$$

and the union is measure disjoint.

(ii) Ω satisfies a set-valued equation (up to a measure zero set)

$$(1.2) \quad A(\Omega) = \bigcup_{d \in \mathcal{D}} (\Omega + d)$$

for some finite digit set $\mathcal{D} \in \mathbf{Z}^n$ of cardinality $|\det(A)|$.

*Received September 23, 1998; accepted for publication December 8, 1998.

[†]Mathematical Sciences Department, Carnegie Mellon University, Pittsburgh, PA 15213, USA (tim@andrew.cmu.edu).

[‡]School of Mathematics, Georgia Institute of Technology, Atlanta, GA 30332, USA (wang@math.gatech.edu). Research supported in part by the National Science Foundation, grant DMS-9706793.

We call a compact set Ω with positive Lebesgue measure that satisfies the condition (ii) a *self-affine tile*. It is known that such an Ω must have nonempty interior, and must tile \mathbf{R}^n by translation by some subset of \mathbf{Z}^n . If in addition Ω is a \mathbf{Z}^n -tile of \mathbf{R}^n , then the digit set \mathcal{D} must form a complete residue system (mod A), i.e., a complete set of coset representatives of the group $\mathbf{Z}^n/A(\mathbf{Z}^n)$. Conversely, if the digit set \mathcal{D} is a complete residue system (mod A) then there is a unique compact set Ω satisfying (1.2), and Ω must be a self-affine tile, namely it must have positive Lebesgue measure. Note that there is an obviously equivalent formulation of (1.2) in the form of a refinement equation, which is

$$(1.3) \quad \chi_\Omega(x) = \sum_{d \in \mathcal{D}} \chi_\Omega(Ax - d).$$

Self-affine tiles have been extensively studied in recent years. More results on them can be found in Bandt [1], Gröchenig and Haas [11], Kenyon [18], and Lagarias and Wang [23], [19], [20].

In this paper we study Haar-type scaling function vectors and multiwavelet bases. Scaling function vectors are generalizations of scaling functions to vector valued functions. More precisely, a vector valued function $f(x) = [f_1(x), \dots, f_r(x)]^T$, with each compactly supported $f_i(x) \in L^2(\mathbf{R}^n)$, is a *scaling function vector* if the following two conditions are met:

- (a) $f(x)$ satisfies a vector refinement equation

$$(1.4) \quad f(x) = \sum_{\alpha \in \mathbf{Z}^n} c_\alpha f(Ax - \alpha),$$

where the coefficients c_α are matrices in $M_r(\mathbf{R})$. r is called the *vector multiplicity* of $f(x)$.

- (b) $v_f := \int_{\mathbf{R}^n} f(x) dx \neq 0$ and for all $\alpha, \beta \in \mathbf{Z}^n$,

$$(1.5) \quad \int_{\mathbf{R}^n} f(x - \alpha) f^T(x - \beta) dx = \delta_{\alpha - \beta} \Lambda$$

where Λ is an $r \times r$ diagonal matrix with positive diagonal elements and δ_γ is the standard Kronecker symbol with $\delta_0 = 1$ and $\delta_\gamma = 0$ otherwise.

Note that $v_f \neq 0$ implies that v_f is a $|\det(A)|$ -eigenvector of the matrix $c := \sum_{\alpha \in \mathbf{Z}^n} c_\alpha$.

As with the case of a scaling function, an orthonormal basis of $L^2(\mathbf{R}^n)$ called a *multiwavelet basis* can be constructed from a scaling function vector. Multiwavelets have received considerable attention recently, after the construction of multiwavelets by Donovan, Geronimo, Hardin, Kessler and Massopust using fractal interpolation functions [15], [9], [6], and by Goodman and Lee [10]. One advantage of multiwavelets is that they can be made to combine smoothness with small supports. Although these properties are achieved at the cost of using more wavelets, their potential importance in numerical applications may outweigh the cost.

DEFINITION 1.1. *We say a function vector $f : \mathbf{R}^n \rightarrow \mathbf{R}^r$ is of Haar-type if each component of $f(x)$ is the characteristic function of some compact set.*

Now, let $\Omega = (\Omega_1, \dots, \Omega_r)$ be an r -tuple of compact sets in \mathbf{R}^n , and let $\chi_\Omega(x) := [\chi_{\Omega_1}(x), \dots, \chi_{\Omega_r}(x)]^T$. It is then natural to ask whether $\chi_\Omega(x)$ is a scaling function vector for some given dilation A . Our main theorems, stated below, classify all Haar-type scaling function vectors.

THEOREM 1.1. *Let $A \in M_n(\mathbf{Z})$ be an expanding matrix and $\Omega = (\Omega_1, \dots, \Omega_r)$ be an r -tuple of compact sets in \mathbf{R}^n . Suppose that $\chi_\Omega(x)$ is a scaling function vector satisfying the vector refinement equation*

$$(1.6) \quad \chi_\Omega(x) = \sum_{\alpha \in \mathbf{Z}^n} c_\alpha \chi_\Omega(Ax - \alpha).$$

Then

- (i) *Each coefficient matrix $c_\alpha \in M_r(\mathbf{R})$ is a zero-one matrix.*
- (ii) *For each $\alpha \in \mathbf{Z}^n$ the matrix*

$$b_\alpha = \sum_{\beta \in \mathbf{Z}^n} c_{\alpha+A\beta}$$

is a zero-one matrix and contains exactly one entry of 1 in each column.

- (iii) *The nonnegative matrix $c = \sum_{\alpha \in \mathbf{Z}^n} c_\alpha$ is irreducible.*
- (iii') *The nonnegative matrix $c = \sum_{\alpha \in \mathbf{Z}^n} c_\alpha$ is primitive.¹*
- (iv) *$\{\Omega_i : 1 \leq i \leq r\}$ are measure disjoint, and $\cup_{i=1}^r \Omega_i$ is a \mathbf{Z}^n -tile of \mathbf{R}^n .*

We use the term *self-affine multi-tile* for Ω because \mathbf{R}^n can be tiled by the translates of $\Omega_1, \dots, \Omega_r$, and for all $1 \leq i \leq r$ we have

$$(1.7) \quad A(\Omega_i) = \bigcup_{j=1}^r (\Omega_j + \mathcal{D}_{ij}),$$

where each \mathcal{D}_{ij} is a finite, possibly empty, set in \mathbf{Z}^n , with all unions in (1.7) measure disjoint.

We will now briefly describe the construction of *Haar-type multiwavelet bases* which correspond to the scaling function vectors of Theorem 1.1. Let $f = \chi_\Omega$ be a scaling function vector satisfying (1.6). Let $\mathcal{E}_i = \{(\alpha, j) : (c_\alpha)_{ij} = 1\}$ for all $1 \leq i \leq r$, and set $\gamma_i = |\mathcal{E}_i|$. For each i define an arbitrary bijection I_i mapping \mathcal{E}_i onto $\{1, 2, \dots, \gamma_i\}$. We now normalize the scaling functions, setting $\phi_i = \|f_i\|_2^{-1} f_i$. Next let M_i be an orthogonal $\gamma_i \times \gamma_i$ matrix with the first row $\gamma_i^{-1/2} [1, 1, \dots, 1]$. Define, for each $1 \leq i \leq r$, and $2 \leq j \leq \gamma_i$ the following functions

$$(1.8) \quad g_{ij}(x) = |\det(A)| \sum_{(\alpha, k) \in \mathcal{E}_i} M_i(j, I_i(\alpha, k)) \phi_k(Ax - \alpha).$$

We have that g_{ij} is supported on the set Ω_i , using (1.6). Hence, we only need to establish orthonormality of g_{ij} to g_{ik} , and of g_{ij} to ϕ_k . This readily follows by computing these inner products, applying the refinement equations, and using the orthogonality of the M_i . Finally, observe that we have a total of $\gamma_1 + \dots + \gamma_r - r = |\det(A)|r - r$ wavelet functions, precisely the number of wavelets needed.

THEOREM 1.2. *Let $A \in M_n(\mathbf{Z})$ be an expanding matrix. Suppose that the coefficient matrices c_α of the vector refinement equation*

$$(1.9) \quad f(x) = \sum_{\alpha \in \mathbf{Z}^n} c_\alpha f(Ax - \alpha)$$

satisfy properties (i)-(iii) of Theorem 1.1. Then up to scalar multiplications the vector refinement equation has a unique compactly supported solution, which is of Haar-type $f(x) = \chi_\Omega(x)$. Furthermore, $\Omega = (\Omega_1, \dots, \Omega_r)$ satisfies

¹Note that (iii') is stronger than (iii). We include (iii) to simplify the statement of Theorem 1.2.

- (A) Each Ω_i has nonempty interior, and $\Omega_i = \overline{\Omega_i}^{\circ}$.²
 (B) There exist subsets $\mathcal{J}_1, \dots, \mathcal{J}_r$ of \mathbf{Z}^n such that

$$(1.10) \quad \bigcup_{i=1}^r \bigcup_{\gamma_i \in \mathcal{J}_i} (\Omega_i + \gamma_i)$$

is a tiling of \mathbf{R}^n .

We remark that the solution χ_{Ω} in Theorem 1.2 is not necessarily a scaling function vector, even when $n = 1$ and $r = 1$. The simplest counter-example is the refinement equation

$$f(x) = f(2x) + f(2x - 3),$$

which admits the Haar-type solution $f(x) = \chi_{[0,3)}(x)$ that is not a scaling function. Checking whether a solution χ_{Ω} is a scaling function vector can be done in finitely many steps using an appropriate generalization of an algorithm in Lawton [24]. For a solution χ_{Ω} that is not a scaling function vector, it is not known whether $\bigcup_{i=1}^r \Omega_i$ still tiles \mathbf{R}^n by translation. We state here as a conjecture that it does.

The rest of this paper are organized as follows: In §2 we state and prove some general results concerning scaling function vectors and certain generalization of iterated function systems. We then use these results to prove our main theorems in §3. In §4 we present some examples of self-affine multi-tiles.

The second author would like to thank Professor Ka-Sing Lau for the invitation to visit the Chinese University of Hong Kong, where this research was conducted primarily. Both authors thank the mathematics department of CUHK for the generous support, and Professor Lau and Dr. Sze-Man Ngai in particular for their hospitality and stimulating discussions. After we completed this paper we received a preprint by Gröchenig, Haas and Raugi [12] that contains some overlapping results.

2. General Results. Consider the vector refinement equation

$$(2.1) \quad f(x) = \sum_{\alpha \in \mathbf{Z}^n} c_{\alpha} f(Ax - \alpha)$$

where $f(x) = [f_1(x), \dots, f_r(x)]^T$, $c_{\alpha} \in M_r(\mathbf{R})$ and $c_{\alpha} \neq 0$ for only finitely many α . We define the *symbol* of the vector refinement equation to be

$$(2.2) \quad m(\xi) = \frac{1}{|\det(A)|} \sum_{\alpha \in \mathbf{Z}^n} c_{\alpha} e^{-i\langle \alpha, \xi \rangle}.$$

Let $B := A^T$. It verifies easily that the Fourier transform of $f(x)$ satisfies

$$(2.3) \quad \widehat{f}(\xi) = m(B^{-1}\xi) \widehat{f}(B^{-1}\xi).$$

Some standard properties for non-vector refinement equations are known to generalize to the vector refinement equation (2.1). For example, if $f(x)$ is integrable and $v_f := \int_{\mathbf{R}^n} f(x) dx \neq 0$ then v_f is a 1-eigenvector of the matrix $m(0)$. But unlike the non-vector case, a given vector refinement equation may have more than one independent

²If we interpret the equality in (1.9) as almost everywhere, then $\Omega_i = \overline{\Omega_i}^{\circ}$ up to a measure zero set.

compactly supported L^1 solution. In fact, under the assumption that the spectral radius $\rho(m(0)) \leq 1$ each 1-eigenvector v of $m(0)$ leads to a compactly supported solution $f(x)$ (in the sense of tempered distribution) to the vector refinement equation (2.1) via

$$(2.4) \quad \widehat{f}(\xi) = \left(\prod_{j=1}^{\infty} m(B^{-j}\xi) \right) v,$$

see Cohen, Daubechies and Plonka [4].

For a vector valued function $f(x)$ we define

$$v_f := \int_{\mathbf{R}^n} f(x) dx, \quad \Lambda_f := \int_{\mathbf{R}^n} f(x)f^T(x) dx.$$

For a scaling function vector $f(x)$ the matrix Λ_f is a diagonal matrix with positive diagonal entries.

LEMMA 2.1. *Suppose that $f(x)$ satisfies (2.1) and is a scaling function vector. Then*

$$(2.5) \quad \sum_{\alpha \in \mathbf{Z}^n} c_\alpha \Lambda_f c_{\alpha+A\beta}^T = \delta_\beta |\det(A)| \Lambda_f.$$

Proof. Let $q = |\det(A)|$. By definition of a scaling function vector,

$$\begin{aligned} \delta_\beta |\det(A)| \Lambda_f &= q \int_{\mathbf{R}^n} f(x)f^T(x + \beta) dx \\ &= q \sum_{\alpha \in \mathbf{Z}^n} \sum_{\gamma \in \mathbf{Z}^n} c_\alpha \left(\int_{\mathbf{R}^n} f(Ax - \alpha)f^T(Ax + A\beta - \gamma) dx \right) c_\gamma^T \\ &= \sum_{\alpha \in \mathbf{Z}^n} \sum_{\gamma \in \mathbf{Z}^n} c_\alpha \left(\int_{\mathbf{R}^n} f(y - \alpha)f^T(y + A\beta - \gamma) dy \right) c_\gamma^T \\ &= \sum_{\alpha \in \mathbf{Z}^n} c_\alpha \left(\int_{\mathbf{R}^n} f(y - \alpha)f^T(y - \alpha) dy \right) c_{\alpha+A\beta}^T \\ &= \sum_{\alpha \in \mathbf{Z}^n} c_\alpha \Lambda_f c_{\alpha+A\beta}^T. \quad \square \end{aligned}$$

DEFINITION 2.1. *Let Λ be a diagonal matrix with positive diagonal entries. The vector refinement equation (2.1) is said to satisfy the orthogonal coefficients condition (with respect to Λ) if*

$$\sum_{\alpha \in \mathbf{Z}^n} c_\alpha \Lambda c_{\alpha+A\beta}^T = \delta_\beta |\det(A)| \Lambda.$$

THEOREM 2.2. *Suppose that the vector refinement equation (2.1) satisfies the orthogonal coefficients condition. Suppose further that $v_0 \in \mathbf{R}^r$ is a 1-eigenvector of $m(0) = |\det(A)|^{-1} \sum_{\alpha \in \mathbf{Z}^n} c_\alpha$, and that $\rho(m(0)) \leq 1$. Then there exists a nontrivial compactly supported L^2 solution $f(x) = [f_1(x), \dots, f_r(x)]^T$ of the vector refinement equation (2.1) such that $v_f = v_0$.*

Proof. The orthogonal coefficients condition implies the existence of a diagonal matrix $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_r)$ with $\lambda_i > 0$ such that

$$\sum_{\alpha \in \mathbf{Z}^n} c_\alpha \Lambda c_{\alpha+A}^T = \delta_\beta |\det(A)| \Lambda.$$

We partition the cube $[0, 1]^n$ into disjoint sets $\Omega_1, \dots, \Omega_r$ of positive Lebesgue measures. Let $g(x) = [g_1(x), \dots, g_r(x)]^T$ such that: (1) each $g_i(x)$ is supported on Ω_i ; (2) $v_g = v_0$; (3) $\Lambda_g = t\Lambda$ for some $t > 0$. Without loss of generality we assume that $t = 1$, $\Lambda_g = \Lambda$.

Let \mathcal{T} be the operator defined by

$$(2.6) \quad \mathcal{T}h(x) = \sum_{\alpha \in \mathbf{Z}^n} c_\alpha h(Ax - \alpha).$$

Then $\widehat{\mathcal{T}h}(\xi) = m(B^{-1}\xi)\widehat{h}(B^{-1}\xi)$ where $B = A^T$. Denote $g_k = \mathcal{T}^k g$ for $k \geq 0$. By induction on k it is easily verified that

$$(2.7) \quad v_{g_k} = v_0 \quad \text{and} \quad \int_{\mathbf{R}^n} g_k(x)g_k(x - \alpha)^T dx = \delta_\alpha \Lambda.$$

Therefore there exists a constant $K > 0$ such that

$$(2.8) \quad \int_{\mathbf{R}^n} |\widehat{g}_k(\xi)|^2 d\xi \leq K,$$

where $|\cdot|$ denotes the Euclidean norm on \mathbf{C}^r . Now

$$\widehat{g}_k(\xi) = \left(\prod_{j=1}^k m(B^{-j}\xi) \right) \widehat{g}(B^{-k}\xi).$$

So by Theorem 3.2 of Cohen, Daubechies and Plonka [4], $\widehat{g}_k(\xi)$ converges uniformly on compact sets to the entire function $\widehat{f}(\xi) := \left(\prod_{j=1}^\infty m(B^{-j}\xi) \right) v_0^3$. The function $f(x)$, which is the inverse Fourier transform of $\widehat{f}(\xi)$, is compactly supported and satisfies the vector refinement equation (2.1) because $\widehat{f}(\xi) = m(B^{-1}\xi)\widehat{f}(B^{-1}\xi)$. Moreover, $f(x)$ is L^2 because $\widehat{f}(\xi)$ is. Finally, $v_f = \widehat{f}(0) = v_0$. \square

Our next two theorems concern certain generalization of iterated function systems (IFS). Let \mathcal{C}_n denote the space of all nonempty compact subsets of \mathbf{R}^n . Let $\|\cdot\|$ be a norm on \mathbf{R}^n . We define the *Hausdorff metric* on \mathcal{C}_n with respect to the norm by

$$(2.9) \quad d_H(D, D') := \max \left\{ \sup_{x \in D} \inf_{x' \in D'} \|x - x'\|, \sup_{y' \in D'} \inf_{y \in D} \|y - y'\| \right\}.$$

It is well known that (\mathcal{C}_n, d_H) is a complete metric space. Now let d_H^r be the metric defined on \mathcal{C}_n^r , the space of all r -tuples of nonempty compact subsets of \mathbf{R}^n , given by

$$(2.10) \quad d_H^r(\Omega, \Omega') := \max_{1 \leq i \leq r} \{d_H(\Omega_i, \Omega'_i), \dots, d_H(\Omega_r, \Omega'_r)\}.$$

³The result was established in Theorem 3.2 for $A = 2I$. But the proof obviously generalizes to an arbitrary expanding matrix A .

Then (C_n^r, d_H^r) is also a complete metric space. The following result is essentially due to Mauldin and Williams [27]. We include a proof for self-containment.

PROPOSITION 2.3. *Let A be an expanding matrix in $M_n(\mathbf{R})$ and let \mathcal{D}_{ij} , $1 \leq i, j \leq r$, be finite subsets of \mathbf{R}^n , with $\cup_{j=1}^r \mathcal{D}_{ij}$ nonempty. Then there exists a unique $\Omega = (\Omega_1, \dots, \Omega_r) \in C_n^r$ such that*

$$(2.11) \quad A(\Omega_i) = \bigcup_{j=1}^r (\Omega_j + \mathcal{D}_{ij}), \quad 1 \leq i \leq r.$$

Proof. Since A is expanding, there exists a norm $\|\cdot\|$ on \mathbf{R}^n such that $\|A^{-1}x\| \leq \lambda\|x\|$ where $0 < \lambda < 1$ for all $x \in \mathbf{R}^n$. Let the Hausdorff metric d_H be defined using this norm. We now consider the map $\Phi : C_n^r \rightarrow C_n^r$ defined by

$$\Phi_i(\Omega) = \bigcup_{j=1}^r (A^{-1}(\Omega_j) + A^{-1}(\mathcal{D}_{ij})), \quad 1 \leq i \leq r.$$

We show that Φ is contractive in (C_n^r, d_H^r) . Note that

$$d_H\left(\bigcup_{j=1}^m E_j, \bigcup_{j=1}^m E'_j\right) \leq \max_{1 \leq j \leq m} d_H(E_j, E'_j),$$

whenever E_j, E'_j are all in C_n , see Barnsley [2]. Set $\Gamma_i = \{j : \mathcal{D}_{ij} \neq \emptyset\}$. Now for any $\Omega, \Omega' \in C_n^r$ we have

$$\begin{aligned} d_H(\Phi_i(\Omega), \Phi_i(\Omega')) &\leq \max_{j \in \Gamma_i} d_H(A^{-1}(\Omega_j + \mathcal{D}_{ij}), A^{-1}(\Omega'_j + \mathcal{D}_{ij})) \\ &\leq \lambda \max_{j \in \Gamma_i} d_H(\Omega_j + \mathcal{D}_{ij}, \Omega'_j + \mathcal{D}_{ij}) \\ &\leq \lambda \max_{j \in \Gamma_i} \max_{\gamma \in \mathcal{D}_{ij}} d_H(\Omega_j + \gamma, \Omega'_j + \gamma) \\ &= \lambda \max_{j \in \Gamma_i} d_H(\Omega_j, \Omega'_j) \\ &\leq \lambda \max_{1 \leq j \leq r} d_H(\Omega_j, \Omega'_j) \\ &= \lambda d_H^r(\Omega, \Omega'). \end{aligned}$$

Hence

$$d_H^r(\Phi(\Omega), \Phi(\Omega')) = \max_{1 \leq i \leq r} d_H(\Phi_i(\Omega), \Phi_i(\Omega')) \leq \lambda d_H^r(\Omega, \Omega').$$

So Φ is contractive, and it follows that there exists a unique $\Omega \in C_n^r$ such that $\Phi(\Omega) = \Omega$, proving the theorem. \square

We remark that the irreducibility condition on $c = \sum_{\alpha \in \mathbf{Z}^n} c_\alpha$ implies that $\cup_{j=1}^r \mathcal{D}_{ij}$ is nonempty, where $\mathcal{D}_{ij} = \{\alpha : (c_\alpha)_{ij} = 1\}$.

THEOREM 2.4. *Let A be an expanding matrix in $M_n(\mathbf{Z})$ and let \mathcal{D}_{ij} , $1 \leq i, j \leq r$, be finite (possibly empty) subsets of \mathbf{Z}^n . Let $\Omega = (\Omega_1, \dots, \Omega_r) \in C_n^r$ be the solution to*

$$A(\Omega_i) = \bigcup_{j=1}^r (\Omega_j + \mathcal{D}_{ij}), \quad 1 \leq i \leq r.$$

Suppose that $\Omega = \cup_{i=1}^r \Omega_i$ has positive Lebesgue measure. Then $\Omega + \mathbf{Z}^n = \mathbf{R}^n$ and Ω has nonempty interior.

Proof. Let $\pi : \mathbf{R}^n \rightarrow \mathbf{T}^n := \mathbf{R}^n/\mathbf{Z}^n$ be the canonical covering map. Denote $A_* := \pi \circ A \circ \pi^{-1}$ and $\bar{\Omega} = \pi(\Omega)$. Then

$$A_*(\bar{\Omega}) = \bigcup_{i=1}^r \bigcup_{j=1}^r \pi(\Omega_j + \mathcal{D}_{ij}) \subseteq \bigcup_{i=1}^r \pi(\Omega_i) = \bar{\Omega}.$$

Note that we only have $A_*(\bar{\Omega}) \subseteq \bar{\Omega}$ because some of the \mathcal{D}_{ij} 's may be empty. So $\bar{\Omega}$ is invariant under the ergodic map A on \mathbf{T}^n . This implies that either $\mu(\bar{\Omega}) = 0$ or $\bar{\Omega} = \mathbf{T}^n$ up to a measure zero subset. But $\mu(\Omega) > 0$, so $\bar{\Omega} = \mathbf{T}^n$. Hence Ω contains a fundamental domain of the lattice \mathbf{Z}^n , and $\Omega + \mathbf{Z}^n = \mathbf{R}^n$ up to a measure zero set. But each $\Omega + \alpha$, $\alpha \in \mathbf{Z}^n$, is a compact set. So any $x_0 \in \mathbf{R}^n$ must lie in the closure of some $\Omega + \alpha$, and hence in $\Omega + \alpha$. Thus $\Omega + \mathbf{Z}^n = \mathbf{R}^n$.

It remains to be shown that Ω has nonempty interior. Let U be the closure of the unit ball $B_1(0)$ of \mathbf{R}^n . So

$$U = \bigcup_{\alpha \in \mathbf{Z}^n} (\Omega + \alpha) \cap U.$$

Clearly $(\Omega + \alpha) \cap U \neq \emptyset$ for only finitely many $\alpha \in \mathbf{Z}^n$, and each $(\Omega + \alpha) \cap U$ is closed. So $U = \bigcup_{i=1}^m U_i$ where $U_i = (\Omega + \alpha_i) \cap U$. Hence at least one of the U_i 's has nonempty interior, and so Ω has nonempty interior. \square

3. Proofs of Main Theorems. We first state some facts concerning nonnegative matrices, that is, matrices with nonnegative entries. We call a nonnegative matrix $B = [b_{ij}] \in M_r(\mathbf{R})$ *irreducible* if for all $1 \leq i, j \leq r$ there exist $1 \leq k_1, \dots, k_m \leq r$ such that

$$b_{ik_1} b_{k_1 k_2} \cdots b_{k_{m-1} k_m} b_{k_m j} > 0.$$

Otherwise B is called *reducible*. B is *primitive* if B^k are irreducible for all $k \geq 1$. The Perron-Frobenius Theorem states that if a nonnegative B is irreducible then the spectral radius $\rho = \rho(B)$ is a simple eigenvalue of B , and B has a ρ -eigenvector v that is positive. On the other hand, if B is reducible then B can be block triangularized by a permutation matrix P , that is

$$PBP^{-1} = \begin{bmatrix} B_{11} & 0 \\ B_{21} & B_{22} \end{bmatrix}, \quad B_{11}, B_{22} \text{ are nonempty.}$$

We now go back to the vector refinement equation

$$(3.1) \quad f(x) = \sum_{\alpha \in \mathbf{Z}^n} c_\alpha f(Ax - \alpha).$$

As before, here $A \in M_n(\mathbf{Z})$ is expanding, $c_\alpha \in M_r(\mathbf{R})$ and $c_\alpha \neq 0$ for only finitely many α .

LEMMA 3.1. *Suppose that all c_α in (3.1) are zero-one matrices and $\rho(c) = |\det(A)|$ where $c := \sum_{\alpha \in \mathbf{Z}^n} c_\alpha$. If c is reducible then the vector refinement equation (3.1) cannot admit a Haar-type scaling function vector solution.*

Proof. Since c is reducible we may assume without loss of generality that

$$c = \begin{bmatrix} \hat{c} & 0 \\ * & \tilde{c} \end{bmatrix}, \quad \hat{c} \in M_m(\mathbf{R}), \quad \tilde{c} \in M_{r-m}(\mathbf{R}),$$

for some $0 < m < r$. The nonnegativity forces every c_α to be of the form

$$c_\alpha = \begin{bmatrix} \hat{c}_\alpha & 0 \\ * & \tilde{c}_\alpha \end{bmatrix}, \quad \hat{c}_\alpha \in M_m(\mathbf{R}), \quad \tilde{c}_\alpha \in M_{r-m}(\mathbf{R}).$$

Now assume that χ_Ω is a Haar-type scaling function vector satisfying (3.1) for some $\Omega = (\Omega_1, \dots, \Omega_r) \in \mathcal{C}_n^r$. Let $\hat{\Omega} = (\Omega_1, \dots, \Omega_m) \in \mathcal{C}_n^m$. Then

$$(3.2) \quad \chi_{\hat{\Omega}}(x) = \sum_{\alpha \in \mathbf{Z}^n} \hat{c}_\alpha \chi_{\hat{\Omega}}(Ax - \alpha).$$

Define $\hat{\mathcal{D}}_{ij} = \{\alpha \in \mathbf{Z}^n : (\hat{c}_\alpha)_{ij} = 1\}$ for $1 \leq i, j \leq m$. Then $\hat{\Omega}$ satisfies

$$(3.3) \quad A(\Omega_i) = \bigcup_{j=1}^m (\Omega_j + \hat{\mathcal{D}}_{ij}), \quad 1 \leq i \leq m.$$

It follows from Theorem 2.4 that either $\mu(\cup_{i=1}^m \Omega_i) = 0$ or $\mu(\cup_{i=1}^m \Omega_i) \geq 1$. But all $\mu(\Omega_i) > 0$ by definition, we must thus have $\mu(\cup_{i=1}^m \Omega_i) \geq 1$. However, this forces $\mu(\cup_{i=1}^r \Omega_i) > 1$ and hence the orthogonality of $\{\chi_{\Omega_i}(x - \alpha) : 1 \leq i \leq r, \alpha \in \mathbf{Z}^n\}$ cannot hold, a contradiction. \square

Proof of Theorem 1.1. (i) Let $\langle \cdot, \cdot \rangle$ denote the standard inner product in $L^2(\mathbf{R}^n)$. The scaling function vector χ_Ω gives

$$\langle \chi_{\Omega_i}(x), \chi_{\Omega_j}(x - \alpha) \rangle = 0, \quad i \neq j \text{ or } \alpha \neq 0.$$

This means that Ω_i and $\Omega_j + \alpha$ are measure disjoint for $i \neq j$ or $\alpha \neq 0$. The vector refinement equation (1.6) now yields $c_\alpha \in M_r(\{0, 1\})$ immediately.

(ii) By Lemma 2.1 there exists a diagonal matrix $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_r)$ with all $\lambda_i > 0$ such that

$$(3.4) \quad \sum_{\alpha \in \mathbf{Z}^n} c_{\alpha+A\beta_1} \Lambda c_{\alpha+A\beta_2}^T = \delta_{\beta_1-\beta_2} |\det(A)| \Lambda.$$

It follows that for $\beta_1 \neq \beta_2$ the matrices $c_{\alpha+A\beta_1}$ and $c_{\alpha+A\beta_2}$ cannot have entries of 1 in a common column. In addition, no c_α can have two or more entries of 1 in a column, or it would create a positive off-diagonal entry in $c_\alpha \Lambda c_\alpha^T$ to contradict (3.4). Therefore $b_\alpha = \sum_{\beta \in \mathbf{Z}^n} c_{\alpha+A\beta}$ is a zero-one matrix with at most one entry of 1 in each column.

We show that b_α cannot have a zero column. Let $\alpha_1, \dots, \alpha_q$ be a complete residue system (mod A), where $q = |\det(A)|$. Note that

$$\begin{aligned} \sum_{j=1}^q b_{\alpha_j} \Lambda b_{\alpha_j}^T &= \sum_{j=1}^q \sum_{\beta_1 \in \mathbf{Z}^n} \sum_{\beta_2 \in \mathbf{Z}^n} c_{\alpha_j+A\beta_1} \Lambda c_{\alpha_j+A\beta_2}^T \\ &= \sum_{\gamma \in \mathbf{Z}^n} \sum_{\alpha \in \mathbf{Z}^n} c_\alpha \Lambda c_{\alpha+A\gamma}^T \\ &= q\Lambda. \end{aligned}$$

Now assume that some b_{α_i} , say b_{α_1} , has a zero column. Then for $v = [1, \dots, 1]^T$ we have

$$v^T b_{\alpha_1} \Lambda b_{\alpha_1}^T v < \sum_{i=1}^r \lambda_i.$$

Therefore

$$v^T q \Lambda v = \sum_{i=1}^r v^T b_{\alpha_i} \Lambda b_{\alpha_i}^T v < q \sum_{i=1}^r \lambda_i,$$

contradicting $v^T q \Lambda v = q \sum_{i=1}^r \lambda_i$. So b_{α_i} has no zero column. Now for any $\alpha \in \mathbf{Z}^n$ we have $b_\alpha = b_{\alpha_i}$ for some i . This proves (ii).

(iii) This is Lemma 3.1.

(iii') Iterating (1.6) yields

$$\chi_\Omega(x) = \sum_{\alpha \in \mathbf{Z}^n} \sum_{\beta \in \mathbf{Z}^n} c_\alpha c_\beta \chi_\Omega(A^2 x - A\alpha - \beta).$$

For the new equation, $\sum_{\alpha \in \mathbf{Z}^n} \sum_{\beta \in \mathbf{Z}^n} c_\alpha c_\beta = c^2$. By (iii) c^2 is irreducible. By the same argument, all c^k are irreducible. Hence c is primitive.

(vi) The measure disjointness of $\{\Omega_1, \dots, \Omega_r\}$ is clear. Now by Theorem 2.4 the set $\Omega := \cup_{i=1}^r \Omega_i$ has the property that $\Omega + \mathbf{Z}^n = \mathbf{R}^n$. But the orthogonality condition forces $\mu(\Omega) \leq 1$. Hence $\mu(\Omega) = 1$ and therefore $\mathbf{R}^n = \Omega + \mathbf{Z}^n$ is a tiling. \square

REMARK. In general $\Omega := \cup_{i=1}^r \Omega_i$ is not itself a self-affine tile. It is not hard to check that Ω is a self-affine tile if and only if $\cup_{i=1}^r \mathcal{D}_{ij} = \mathcal{D}$ for all $1 \leq j \leq r$, where $\mathcal{D} \in \mathbf{Z}^n$ with $|\mathcal{D}| = |\det(A)|$ and $\mathcal{D}_{ij} = \{\alpha : (c_\alpha)_{ij} = 1\}$. We omit the proof here.

LEMMA 3.2. *Let D_1, \dots, D_m be measure disjoint compact sets in \mathbf{R}^n such that $\cup_{i=1}^m D_i = \overline{B_K(x_0)}$. Then for any D_i having positive Lebesgue measure we have $D_i^\circ \neq \emptyset$ and $\overline{D_i^\circ} = D_i$ up to a measure zero set.*

Proof. Suppose that some D_i , say D_1 , has positive Lebesgue measure but no interior point. Then for any $y \in D_1$ there exists a sequence $\{y_k\}$ such that $\lim_{k \rightarrow \infty} y_k = y$ and $y_k \notin D_1$. Hence $y \in \cup_{i=2}^m D_i$, and so $D_1 \subseteq \cup_{i=2}^m D_i$. This contradicts the measure disjointness of D_i 's. Thus $D_i^\circ \neq \emptyset$ for all D_i with positive Lebesgue measure. Now using the same argument we have $\partial D_i \subseteq \cup_{j \neq i} D_j$. Hence by measure disjointness again $\mu(\partial D_i) = 0$ for all i . Thus $D_i = \overline{D_i^\circ}$ up to a measure zero set. \square

Proof of Theorem 1.2. The assumed condition (ii) implies that the sum of the entries in each column of c is $q = |\det(A)|$. So the nonnegative matrix $m(0) = q^{-1}c$ is column stochastic. Therefore $\rho(m(0)) = 1$. By Theorem 2.2 there exists a compactly supported $f(x) = [f_1(x), \dots, f_r(x)]^T$, $f(x) \neq 0$ and each $f_i(x) \in L^2(\mathbf{R}^n)$, such that

$$f(x) = \sum_{\alpha \in \mathbf{Z}^n} c_\alpha f(Ax - \alpha).$$

Let $\mathbf{U} = (U_1, \dots, U_r)$ where each U_i is the essential support of $f_i(x)$. The vector refinement equation (1.9) implies that

$$(3.5) \quad U_i \subseteq \bigcup_{j=1}^r A^{-1}(U_j + \mathcal{D}_{ij}), \quad 1 \leq i \leq r$$

where $\mathcal{D}_{ij} = \{\alpha : (c_\alpha)_{ij} = 1\}$. It follows by iterating (3.5) while replacing U_i in the equation by its closure that $U_i \subseteq \Omega_i$, where $\Omega = (\Omega_1, \dots, \Omega_r) \in \mathcal{C}_n^r$ is the unique solution to

$$(3.6) \quad A(\Omega_i) = \bigcup_{j=1}^r (\Omega_j + \mathcal{D}_{ij}), \quad 1 \leq i \leq r.$$

Since $f \neq 0$ at least one of the U_i 's has positive Lebesgue measure. It follows from Theorem 2.4 that $\Omega = \cup_{i=1}^r \Omega_i$ has nonempty interior.

We show that χ_Ω satisfies the vector refinement equation (1.9). Note that condition (ii) implies that $\sum_{i=1}^r |\mathcal{D}_{ij}| = |\det(A)|$. Therefore

$$(3.7) \quad \sum_{i=1}^r \mu\left(\bigcup_{j=1}^r (\Omega_j + \mathcal{D}_{ij})\right) \leq \sum_{i=1}^r \sum_{j=1}^r |\mathcal{D}_{ij}| \mu(\Omega_i) = |\det(A)| \sum_{i=1}^r \mu(\Omega_i).$$

But

$$(3.8) \quad |\det(A)| \sum_{i=1}^r \mu(\Omega_i) = \sum_{i=1}^r \mu(A(\Omega_i)) = \sum_{i=1}^r \mu\left(\bigcup_{j=1}^r (\Omega_j + \mathcal{D}_{ij})\right).$$

Thus all inequalities in (3.7) are equalities, and so all unions in $\bigcup_{j=1}^r (\Omega_j + \mathcal{D}_{ij})$ are measure disjoint. This forces

$$\chi_\Omega(x) = \sum_{\alpha \in \mathbf{Z}^n} c_\alpha \chi_\Omega(Ax - \alpha).$$

Note that the vector $v = \int_{\mathbf{R}^n} \chi_\Omega(x) dx$ is a $|\det(A)|$ -eigenvector of c . It follows from the irreducibility of c that v is a positive vector. So all $\mu(\Omega_i) > 0$.

We now prove property (B). $\Omega^o \neq \emptyset$ so $\Omega_i^o \neq \emptyset$ for some i , say $\Omega_1^o \neq \emptyset$. Since A is expanding, for some sufficiently large $N > 0$ the set $A^N(\Omega_1)$ contains an interior point $x_0 \in \mathbf{Z}^n$. So 0 is an interior point of $\tilde{\Omega}_1 := A^N(\Omega_1) - x_0$. This means that for each $k > 0$ there exists an $m_k > 0$ such that the ball $B_k(0)$ is covered by $A^{m_k}(\tilde{\Omega}_1)$. It follows from the inflation property (3.6) that there exist finite subsets $\mathcal{J}_1^k, \dots, \mathcal{J}_r^k$ of \mathbf{Z}^n such that

$$(3.9) \quad B_k(0) \subseteq \bigcup_{i=1}^r (\Omega_i + \mathcal{J}_i^k),$$

where all unions in (3.9) are measure disjoint. Now for any given $K > 0$ and each i there are only finitely many distinct $\mathcal{J}_i^k \cap B_K(0)$. So we can find a subsequence $\mathcal{J}_1^{m_k}, \dots, \mathcal{J}_r^{m_k}$ such that $\mathcal{J}_i^{m_k} \rightarrow \mathcal{J}_i$ in the sense that for each $K > 0$, $\mathcal{J}_i^{m_k} \cap B_K(0) = \mathcal{J}_i \cap B_K(0)$ for sufficiently large m_k . By (3.9)

$$(3.10) \quad \mathbf{R}^n = \bigcup_{i=1}^r (\Omega_i + \mathcal{J}_i)$$

with all unions measure disjoint, proving (B).

To prove (A), take a sufficiently large $K > 0$ so that $\overline{B_K(0)}$ contains at least one translate of each Ω_i in the tiling (3.10). So $\overline{B_K(0)}$ is the measure disjoint union of compact sets that include translates of each Ω_i . Since all Ω_i have positive Lebesgue measure, by Lemma 3.2 Ω_i all have nonempty interior and $\Omega_i = \overline{\Omega_i^o}$ up to a measure zero set. This proves (A). \square

4. Examples. We now look at some examples of self-affine multi-tiles and Haar-type scaling function vectors. We first select an expanding integer matrix A , and then choose coefficient matrices c_α satisfying properties (i)-(iii) of Theorem 1.1. In Examples 1-3 our expanding matrix A is the quincunx matrix (also referred to as the “dragon” matrix),

$$A = \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}.$$

EXAMPLE 1. We let A be the quincunx matrix and choose the coefficient matrices to be

$$c_{(0,0)} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}, \quad c_{(1,0)} = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, \quad c_{(0,1)} = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}.$$

The self-affine multi-tile $\Omega = (\Omega_1, \Omega_2)$ is shown in Figure 1.

EXAMPLE 2. We again choose A to be the quincunx matrix and $r = 3$. The coefficient matrices are

$$c_{(0,0)} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 1 \end{bmatrix}, \quad c_{(1,0)} = \begin{bmatrix} 0 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad c_{(0,1)} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

The self-affine multi-tile $\Omega = (\Omega_1, \Omega_2, \Omega_3)$ is shown in Figure 2.

EXAMPLE 3. We let A be the quincunx matrix and $r = 4$. The coefficient matrices are chosen as

$$c_{(0,0)} = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad c_{(1,0)} = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \quad c_{(0,1)} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}.$$

$\Omega = (\Omega_1, \Omega_2, \Omega_3)$ is shown in Figure 3.

EXAMPLE 4. In this last example we let

$$A = \begin{bmatrix} 2 & 1 \\ -1 & 1 \end{bmatrix},$$

which has $\det(A) = 3$. We choose the coefficient matrices to be

$$c_{(0,0)} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad c_{(0,-1)} = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \quad c_{(1,-1)} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}, \quad c_{(1,0)} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}.$$

$\Omega = (\Omega_1, \Omega_2)$ is shown in Figure 4.

We have obtained these examples somewhat by trial and error, by generating the plots of the tiles, and choosing ones that looked nice. In some cases the plots of the different tiles were very dispersed, and hard to visualize. There are many other possibilities for Haar-type scaling function vectors, using other dilation matrices A .




Figure 1: Ω_1 (light), Ω_2 (dark)




Figure 2: $\Omega_1, \Omega_2, \Omega_3$ (light ... dark)




Figure 3: $\Omega_1, \Omega_2, \Omega_3, \Omega_4$, (light ... dark)




Figure 4: Ω_1 (light), Ω_2 (dark)

REFERENCES

- [1] C. BANDT, *Self-similar sets 5. Integer matrices and fractal tilings of \mathbf{R}^n* , Proc. Amer. Math. Soc., 112 (1991), pp. 549–562.
- [2] M. BARNSLEY, *Fractals everywhere*, Academic Press, 1988.
- [3] M. A. BERGER AND Y. WANG, *Multidimensional two-scale dilation equations*, in Wavelets—A Tutorial in Theory and Applications, C. K. Chui, ed., 1992, pp. 295–323.
- [4] A. COHEN, I. DAUBECHIES, AND G. PLONKA, *Regularity of refinable function vectors*, J. Fourier Anal. Appl., 3 (1997), pp. 295–324.
- [5] I. DAUBECHIES, *Ten Lectures on Wavelets*, SIAM, Philadelphia, PA, 1992.
- [6] G. DONOVAN, J. GERONIMO, D. HARDIN, AND P. MASSOPUST, *Construction of orthogonal wavelets using fractal interpolation functions*, SIAM J. Math. Anal., 27 (1996), pp. 1158–1192.
- [7] K. J. FALCONER, *The Geometry of Fractal Sets*, Cambridge Univ. Press, Cambridge, 1985.
- [8] J. GERONIMO AND D. HARDIN, *Fractal interpolation surfaces and a related 2-D multiresolution analysis*, J. Math. Anal. Appl., 176 (1993), pp. 561–586.
- [9] J. GERONIMO, D. HARDIN, AND P. MASSOPUST, *Fractal functions and wavelet expansions based on several scaling functions*, J. Approx. Th., 78 (1994), pp. 373–401.
- [10] T. GOODMAN AND S. L. LEE, *Wavelets of multiplicity r* , Trans. Amer. Math. Soc., 342 (1994), pp. 307–324.
- [11] K. GRÖCHENIG AND A. HAAS, *Self-similar lattice tilings*, J. Fourier Analysis, 1 (1994), pp. 131–170.
- [12] K. GRÖCHENIG, A. HAAS, AND A. RAUGI, *Self-affine tilings with several tiles I*, Preprint.
- [13] K. GRÖCHENIG AND W. MADYCH, *Multiresolution analysis, Haar bases, and self-similar tilings*, IEEE Trans. Inform. Th., 38:2, Part 2 (1992), pp. 558–568.
- [14] B. GRÜNBAUM AND G. C. SHEPHARD, *Tilings and Patterns*, Freeman, New York, 1987.
- [15] D. HARDIN, B. KESSLER, AND P. MASSOPUST, *Multiresolution analysis based on fractal functions*, J. Approx. Th., 71 (1992), pp. 373–401.
- [16] C. HEIL AND D. COLLELA, *Matrix refinement equations: existence and uniqueness*, J. Fourier Anal. Appl., 2 (1996), pp. 363–377.
- [17] J. E. HUTCHINSON, *Fractals and self-similarity*, Indiana Univ. Math. J., 30 (1981), pp. 713–747.
- [18] R. KENYON, *Self-replicating tilings*, in Symbolic Dynamics and Applications, P. Walters, ed., Contemporary Math., 135, AMS, Providence, RI, 1992, pp. 239–264.
- [19] J. C. LAGARIAS AND Y. WANG, *Self-affine tiles in \mathbf{R}^n* , Adv. in Math., 121 (1996), pp. 21–49.
- [20] J. C. LAGARIAS AND Y. WANG, *Integral self-affine tiles in \mathbf{R}^n I. Standard and nonstandard digit sets*, J. London Math. Soc., 53 (1996), pp. 161–179.
- [21] J. C. LAGARIAS AND Y. WANG, *Integral self-affine tiles in \mathbf{R}^n , part II: lattice tilings*, J. Fourier Anal. and Appl., 3 (1997), pp. 83–101.
- [22] J. C. LAGARIAS AND Y. WANG, *Haar bases in \mathbf{R}^n and algebraic number theory*, J. Number Theory, 57 (1996), pp. 181–197.
- [23] J. C. LAGARIAS AND Y. WANG, *Haar type orthonormal wavelet basis in \mathbf{R}^2* , J. Fourier Analysis and Appl., 2 (1995), pp. 1–14.
- [24] W. LAWTON, *Necessary and sufficient conditions for constructing orthonormal wavelet bases*, J. Math. Phys., 32 (1991), pp. 57–61.
- [25] W. LAWTON, S. L. LEE, AND Z. SHEN, *An algorithm for matrix extension and wavelet construction*, Math. Comp., 65, pp. 723–737.
- [26] S. MALLAT, *Multiresolution analysis and wavelets*, Tans. Amer. Math. Soc., 315 (1989), pp. 69–88.
- [27] D. MAULDIN AND S. WILLIAMS, *Hausdorff dimension in graph direct constructions*, Trans. Amer. Math. Soc., 309 (1988), pp. 811–829.
- [28] R. S. STRICHARTZ, *Wavelets and Self-Affine Tilings*, Constructive Approximation, 9 (1993), pp. 327–346.
- [29] R. S. STRICHARTZ, *Self-similarity in harmonic analysis*, J. Fourier Anal. Appl., 1 (1994), pp. 1–37.
- [30] W. P. THURSTON, *Groups, Tilings and Finite State Automata*, AMS Colloquium Lect. Notes, 1989.
- [31] A. VINCE, *Replicating tessellations*, SIAM J. Discrete Math., 3 (1993), pp. 501–521.