

## PAIRS OF INTERSECTING REAL MANIFOLDS IN COMPLEX SPACE \*

S. M. WEBSTER†

**Introduction.** In this work we consider two real  $n$ -dimensional submanifolds of complex  $n$ -space which intersect at the origin,

$$(0.1) \quad 0 \in M_1 \cap M_2, \quad M_1, M_2 \subset \mathbf{C}^n.$$

We shall usually assume that the  $M_i$  are totally real, and that the intersection is isolated and transverse, though other cases are eventually interesting. We are primarily interested in the holomorphic equivalence problem, i. e. to find a biholomorphic map fixing the origin and taking  $M_1$  and  $M_2$  simultaneously into some canonical form, but we also consider some specific geometric questions. For example, do there exist complex analytic curves through 0 and cutting each  $M_i$  in a real curve? Do there exist analytic annuli in  $\mathbf{C}^n$  with one bounding circle on  $M_1$  and the other on  $M_2$ ? What is the precise local hull of holomorphy of  $M_1 \cup M_2$  near 0?

The study of a pair of intersecting real curves in the complex plane was perhaps begun by E. Kasner in 1912, see Pfeiffer [6]. The case of two totally real linear  $n$ -planes in  $\mathbf{C}^n$  was studied by Weinstock [12].

We shall consider three main cases: the *generic* case, the *real Lagrangian* case, and the *holomorphically reversible* case. By the generic case we shall mean that there is no additional structure imposed. In the real Lagrangian case we consider such real submanifolds  $M_i^{2n}$  in  $\mathbf{C}^{2n}$  with its (standard) *holomorphic* symplectic form  $\omega$ , where  $\text{Re}(\omega)$  vanishes when restricted to each  $M_i^{2n}$ . The pair is holomorphically reversible if there exists a *holomorphic* involution  $\tau$  near 0, with  $\tau M_1 = M_2$ .

In this paper we shall assume that  $M_1$  and  $M_2$  are real analytic, and use power series methods. Each  $M_i$  is locally the fixed point set of an anti-holomorphic involution  $\rho_i$ ,

$$(0.2) \quad M_i = FP(\rho_i), \quad \sigma = \rho_1 \rho_2.$$

We shall focus on the holomorphic normalization of the pair  $\rho_1, \rho_2$ , and then derive results about  $M_1, M_2$ . The holomorphic map  $\sigma$ , which will be central to our study, is anti-holomorphically reversible (briefly, *anti-reversible*), in that it is conjugate to its inverse by an anti-holomorphic involution:

$$(0.3) \quad \sigma^{-1} = \rho_2 \rho_1 = \rho_1^{-1} \sigma \rho_1.$$

The square  $\sigma^2$  measures the extent to which  $\rho_1$  and  $\rho_2$  fail to commute. The above mentioned “cutting curves”  $C$  satisfy  $\rho_1 C = \rho_2 C = C$ . We now have the additional concept of a pair of “switched curves”,  $\rho_i C_1 = C_2, i = 1, 2$ . Both are then invariant by  $\sigma$ .

In the holomorphically reversible case,

$$(0.4) \quad \rho_2 = \tau \rho_1 \tau, \quad \sigma = \tau_1 \tau, \quad \tau_1 = \rho_1 \tau \rho_1,$$

---

\*Received November 2, 2002; accepted for publication June 3, 2003.

†Department of Mathematics, University of Chicago, 5734 S. University Avenue, Chicago, Illinois 60637, USA (webster@math.uchicago.edu).

which says that  $\sigma$  is also the product of two *holomorphic* involutions. A special case of this was the key to the theory of analytic real  $n$ -manifolds in  $\mathbf{C}^n$  with nondegenerate complex tangents developed in [5]. In the real Lagrangian case we have

$$(0.5) \quad \rho_i^* \omega = -\bar{\omega}, \quad \sigma^* \omega = \omega, \quad \omega = \sum_{j=1}^n dz_j \wedge dz_{n+j}.$$

We emphasize that we are using a holomorphic symplectic form, and not the more usual real part of the Kähler form.

Section 1 is concerned with the linear case. The eigenvalues of a linear anti-reversible map  $\sigma$  occur in pairs. A linear transformation which diagonalizes  $\sigma$  also takes the  $\rho_i$  into the appropriate canonical form. Section 2 treats the non-linear case as an analytic or formal power series perturbation. A procedure is given (prop 1) to derive certain properties of  $\rho_1, \rho_2$  from those of  $\sigma$ , under certain “non-resonance” conditions on the linear part of  $\sigma$ .

In section 3 we collect some known results on convergence relative to a general map  $\sigma$  and adapt them to our needs here. In particular, theorem (2) gives an existence result for invariant submanifolds with linearization for  $\rho_1, \rho_2$  in the generic case. Proposition (2) gives a convergence result for an invariant submanifold without linearization for a symplectic map  $\sigma$ , in the “integrable” hyperbolic case. The argument parallels that for flows given in [9].

In section 4 we derive symplectic (or unimodular) normal forms for anti-reversible symplectic maps  $\sigma$  in  $\mathbf{C}^2$ . The theory splits into two cases (see (1.12)) according to whether the eigenvalues of  $\sigma$  are (i) unimodular (4.5), (4.6), (4.9); or (ii) real (4.12), (4.13). In general these normal forms yield *implicit* normal forms for the real Lagrangian surfaces  $M_1, M_2$ . We also give an application to the local hull of holomorphy of  $M_1 \cup M_2$  in the higher dimensional real Lagrangian case.

Section 5 shows that our pair of real Lagrangian surfaces in  $\mathbf{C}^2 \ni (z, p)$ ,  $\omega = dp \wedge dz$ , may be given in terms of two analytic real functions  $r_i$  as

$$(0.6) \quad \begin{aligned} M_1 : \quad & p = \partial_z r_1, \quad r_1 = az^2 + bz\bar{z} + \bar{a}\bar{z}^2 + \dots, \\ M_2 : \quad & p = \partial_z r_2, \quad r_2 = z\bar{z} + \dots, \end{aligned}$$

where  $b \neq 0$ , and  $(a, b) \neq (0, 1)$ , and the dots represent terms of order three or higher. In case (ii) of (1.12), we apply the normal form for the involutions  $\rho_i$  to write (0.6) with functions  $r_i$  of the form (see (4.16), (4.17) below)

$$(0.7) \quad r_i = \lambda_i^{-1} |z|^2 + \hat{r}_i(|z|^2), \quad \hat{r}_i(s) = O(s^2), \quad i = 1, 2,$$

where

$$(0.8) \quad \lambda_i = \bar{\lambda}_i \neq 0, \pm 1, \quad \lambda_1 \lambda_2 = \epsilon = \pm 1.$$

One consequence of the results of sections 4 and 5 is the following.

**THEOREM 1.** *Suppose that*

$$(0.9) \quad \frac{b^2 - 4|a|^2 + 1}{2b} > 1 \text{ in (0.6), or } \epsilon = +1 \text{ in (0.8).}$$

*Then there exists a real analytic 1-parameter family of analytic annuli  $A_c \subset \mathbf{C}^2$ ,  $0 < c < c_0$ , bounding on  $M_1 \cup M_2$ , and shrinking to the origin as  $c \rightarrow 0$ . These annuli*

sweep out a three dimensional Levi-flat manifold lying on a real analytic set, which is biholomorphic to  $\text{Im}(zp) = 0$ , with  $\text{Re}(zp) > 0$ .

The moduli of these annuli determine to large extent the invariants of the pair of surfaces (see (4.19) below). The union of these annuli  $A_c$  contribute to the local holomorphic hull of  $M_1 \cup M_2$ . It is perhaps interesting to note the instability inherent in this result. If either  $M_1$  or  $M_2$  is slightly perturbed so as to destroy the real Lagrangian condition, then all these analytic annuli may be lost, and  $M_1 \cup M_2$  may actually become holomorphically convex. This is reminiscent of fixed point results for area preserving mappings [9], which may be lost under generic perturbation.

A holomorphically reversible analogue of the foregoing is contained in [5]. In fact, under certain conditions on the tangent planes to the  $M_i$  at the origin, the holomorphic involution  $\tau$  is the covering involution of a 2-fold branched cover which maps  $M_1 \cup M_2$  to a real surface  $N$  with elliptic complex tangent in  $\mathbf{C}^2$ . The holomorphic normal form for  $M_1, M_2$  yields the normal form for  $N$ . That the corresponding annuli  $A_c$  are mapped to the Bishop analytic discs [1] bounding on  $N$  was first pointed out to the author by N. Sibony [8]. Furthermore, a global case of holomorphic reversibility occurs in [11] in the study of the Kobayashi extremal discs of an ellipsoidal domain. Therefore, we have concentrated here on the real Lagrangian case. A preliminary study indicates that theorem one should also hold in the smooth case, by a direct construction of analytic annuli. This will be taken up in a future work.

X. Gong has recently shown that there exist anti-reversible maps  $\sigma$  which are formally holomorphically reversible, but that the involution  $\tau$  cannot have a positive radius of convergence [3]. Thanks are due to him for discussions leading up to this work.

**1. The linear theory.** In this section we consider the relevant aspects of a pair of linear  $n$ -planes  $M_1, M_2$  passing through the origin of  $\mathbf{C}^n$ . With  $M_2$  totally real and  $M_1$  transverse to it, we may write

$$(1.1) \quad \begin{aligned} M_2 &= \{y = 0\} = \{z = \bar{z}\}, \\ M_1 &= \{x = Ay\} = \{z = B\bar{z}\}, \end{aligned}$$

$$z = x + iy, \quad \bar{z} = x - iy.$$

The two matrices  $A, B$  are related by the Cayley transform [13],  $C : \mathcal{A} \rightarrow \mathcal{B}$ ,

$$(1.2) \quad \begin{aligned} B &= C(A) = (A + iI)(A - iI)^{-1} \\ A &= C^{-1}(B) = i(B + I)(B - I)^{-1} \end{aligned}$$

$$\mathcal{B} = \{B | B\bar{B} = I\}, \quad \mathcal{A} = T_I \mathcal{B} = \{A | A = \bar{A}\}.$$

The ‘‘spectral mapping theorem’’ relates the eigenvalues,

$$(1.3) \quad Av = \lambda v \Leftrightarrow Bv = \mu v, \quad \mu = \frac{\lambda + i}{\lambda - i}, \quad \lambda = i \frac{\mu + 1}{\mu - 1}.$$

In particular,  $\lambda \mapsto \bar{\lambda}$  corresponds to  $\mu \mapsto \bar{\mu}^{-1}$ , the real line to the unit circle, and the imaginary axis to the real axis. If  $M_1$  contains some non-zero  $z$  and  $iz$ , then this gives  $A$  an eigenvector to  $\lambda = \pm i$ , i. e.  $\mu = 0, \infty$ . We exclude this case, so that  $B$

is non-singular. Then  $M_1, M_2$  are both totally real and are the fixed point sets of anti-linear involutions  $\rho_1, \rho_2$ ,

$$(1.4) \quad \rho_1(z) = B\bar{z}, \quad \rho_2(z) = \bar{z}, \quad \sigma(z) = \rho_1\rho_2(z) = Bz.$$

We briefly consider the case of two totally real  $n$ -planes with non-transverse intersection  $M_0 = M_1 \cap M_2$ . If again  $\rho_i$  are the involutions, then

$$M_0 = \{z | \rho_1(z) = \rho_2(z) = z\} \subset \{z | \sigma(z) = z\}.$$

Thus,  $M_0$  is the fixed point set of  $\rho_1(z) \equiv \rho_2(z)$  as acting on the  $+1$  eigenspace of  $\sigma$ . Conversely, if this eigenspace is of positive dimension, we get such an  $M_0$ . The case  $\dim M_0 = n - 2$  occurs in connection with a quadratic real  $n$ -manifold in  $\mathbf{C}^n$  with a non-degenerate complex tangent [5].

**a)** Now we consider the generic case of the linear theory. Let  $e$  be an eigenvector of  $\sigma$  with eigenvalue  $\mu$ ,

$$(1.5) \quad \sigma e = \mu e, \Rightarrow \rho_2 e = \bar{\mu} \rho_1 e = \bar{\mu} \sigma \rho_2 e.$$

Thus,  $\rho_1 e$  and  $\rho_2 e$  are dependent eigenvectors of  $\sigma$  with the eigenvalue  $1/\bar{\mu}$ . We consider two cases,

$$(1.6) \quad (i) \mu = \bar{\mu}^{-1}, \quad (ii) \mu \neq \bar{\mu}^{-1}.$$

We make the assumption here and in general that all eigenspaces are one-dimensional.

In case (i) we have

$$(1.7) \quad \rho_i e = \lambda_i e, \quad \lambda_i \bar{\lambda}_i = 1, \quad i = 1, 2, \quad \mu = \lambda_1 \bar{\lambda}_2.$$

The complex line spanned by  $e$  cuts each  $M_i$  in a real line. If we make the change  $e \mapsto ce$ , then  $\lambda_i \mapsto (\bar{c}/c)\lambda_i$ ,  $i = 1, 2$ . Thus, we may arrange either  $\lambda_2 = 1$ , or more symmetrically

$$(1.8) \quad \lambda_1 \lambda_2 = 1.$$

In case (ii) we set

$$(1.9) \quad \begin{aligned} \sigma e_1 &= \mu e_1, & \sigma e_2 &= \bar{\mu}^{-1} e_2, \\ \rho_i e_1 &= \lambda_i e_2, & \rho_i e_2 &= \bar{\lambda}_i^{-1} e_1, & \mu &= \bar{\lambda}_2 / \bar{\lambda}_1. \end{aligned}$$

The complex lines spanned by  $e_1$  and  $e_2$  are switched by the involutions  $\rho_i$ . Under the change  $e_i \mapsto c_i e_i$ , we have  $\lambda_i \mapsto (\bar{c}_1/c_2)\lambda_i$ . So again, we could arrange either  $\lambda_2 = 1$  or (1.8).

The involutions  $\rho_i$  act on the set of complex lines through the origin in  $\mathbf{C}^n$ . For  $n = 2$ , this is the Riemann sphere, on which each  $\rho_i$  fixes the points of a circle  $K_i$ . In case (i)  $K_1 \cap K_2$  is two points representing the two cutting lines. In case (ii) the  $K_i$  are disjoint. If they are taken concentric, centered at 0, then 0 and  $\infty$  represent the two switched lines.

**b)** Next we consider the real Lagrangian case. Since  $\sigma$  is symplectic, it has eigenvalue  $\mu^{-1}$  along with  $\mu$ ; we assume  $\mu \neq \pm 1$ . The above reasoning gives four eigenvalues and vectors,

$$(1.10) \quad \sigma e_1 = \mu e_1, \quad \sigma e_2 = \mu^{-1} e_2, \quad \sigma e_3 = \bar{\mu}^{-1} e_3, \quad \sigma e_4 = \bar{\mu} e_4.$$

We assume the vectors normalized by  $\omega(e_1, e_2) = 1$ ,  $\omega(e_3, e_4) = 1$ , and  $\omega(e_i, e_j) = 0$  otherwise. Then from (1.9) we get

$$(1.11) \quad \begin{aligned} \rho_i e_1 &= \lambda_i e_3, & \rho_i e_3 &= \bar{\lambda}_i^{-1} e_1, \\ \rho_i e_2 &= -\lambda_i^{-1} e_4, & \rho_i e_4 &= -\bar{\lambda}_i e_2, & \mu &= \overline{\lambda_2/\lambda_1}. \end{aligned}$$

By scaling the eigenvectors while preserving the  $\omega(e_i, e_j)$ , we can again arrange  $\lambda_2 = 1$  or (1.8). This is the basic 4-fold case.

There are two special (2-fold) cases,

$$(1.12) \quad (i) \mu = \bar{\mu}^{-1}, \quad (ii) \mu = \bar{\mu},$$

in which we again assume one-dimensional eigenspaces. In case (i)

$$(1.13) \quad \begin{aligned} \sigma e_1 &= \mu e_1, & \sigma e_2 &= \mu^{-1} e_2, & \mu &= \lambda_1 \bar{\lambda}_2, \\ \rho_i e_1 &= \lambda_i e_1, & \rho_i e_2 &= -\lambda_i^{-1} e_2, & \lambda_i \bar{\lambda}_i &= 1, \quad i = 1, 2. \end{aligned}$$

Again we can achieve (1.8) by scaling.

In case (ii)

$$(1.14) \quad \begin{aligned} \sigma e_1 &= \mu e_1, & \sigma e_2 &= \mu^{-1} e_2, & \mu &= \lambda_2/\lambda_1, \\ \rho_i e_1 &= \lambda_i e_2, & \rho_i e_2 &= \lambda_i^{-1} e_1, & \lambda_i &= \bar{\lambda}_i, \quad i = 1, 2, \end{aligned}$$

the latter following from (1.7). This time the change  $e_i \mapsto c_i e_i$ ,  $c_1 c_2 = 1$  results in  $\lambda_i \mapsto |c_1|^2 \lambda_i$ . The change  $(e_1, e_2) \mapsto (e_2, -e_1)$  results in  $\mu \mapsto \mu^{-1}$  and  $\lambda_i \mapsto -\lambda_i^{-1}$ . Thus, it is possible to change the sign of  $\lambda_2$ , but the sign of  $\lambda_1 \lambda_2$ , which by (1.14) is the sign of  $\mu$ , is invariant. It follows that we can achieve either  $\lambda_2 = 1$ , or

$$(1.15) \quad \lambda_1 \lambda_2 = \epsilon \equiv \pm 1, \quad \epsilon = \text{sgn}(\mu).$$

In this last case assume that  $n = 2$  and take coordinates relative to the above determined basis:  $z = z_1 e_1 + z_2 e_2$ . Then

$$(1.16) \quad \sigma(z) = (\mu z_1, \mu^{-1} z_2), \quad \rho_i(z) = (\lambda_i^{-1} \bar{z}_2, \lambda_i \bar{z}_1), \quad i = 1, 2.$$

It follows that the complex curve  $z_1 z_2 = c$  is invariant under both  $\rho_i$  if  $c = \bar{c}$ . Then

$$(1.17) \quad \{z_1 z_2 = c\} \cap FP(\rho_i) = \{z_1 z_2 = c, \quad z_1 \bar{z}_1 = c/\lambda_i\}, \quad i = 1, 2.$$

Thus  $z_1 z_2 = c$  contains an analytic annulus bounding on  $M_1 \cup M_2$ , if both  $c/\lambda_i > 0$ . This is possible if  $\epsilon = +1$ . Then taking  $c$ 's with  $c/\lambda_1 > 0$  gives a one-parameter family of such annuli  $A_c$  lying on the 3-dimensional algebraic set  $\text{Im}(z_1 z_2) = 0$ . As  $c$  varies the bounding circles of  $A_c$  sweep out  $M_1$  and  $M_2$ , minus the origin, while the  $A_c$  themselves sweep out a 3-dimensional manifold contained in the polynomial hull of  $M_1 \cup M_2$  ( see also [12]).

c) The complex-linearly reversible case is similar. Now we have (1.10) with

$$(1.18) \quad \tau e_1 = e_2, \quad \tau e_3 = e_4.$$

From (0.4) we get

$$(1.19) \quad \begin{aligned} \rho_1(e_1, e_2, e_3, e_4) &= (\lambda_1 e_3, \lambda_2 e_4, \bar{\lambda}_1^{-1} e_1, \bar{\lambda}_2^{-1} e_2), \\ \rho_2(e_1, e_2, e_3, e_4) &= (\lambda_2 e_3, \lambda_1 e_4, \bar{\lambda}_2^{-1} e_1, \bar{\lambda}_1^{-1} e_2). \end{aligned}$$

A change  $e_i \mapsto c_i e_i$  with  $c_1 = c_2$  and  $c_3 = c_4$  preserves (1.10) and allows us to achieve (1.8).

We may again have the two 2-fold cases (1.12). In case (i) we have

$$\rho_1(e_1, e_2) = (\lambda_1 e_1, \lambda_2 e_2), \quad \rho_2(e_1, e_2) = (\lambda_2 e_1, \lambda_1 e_2),$$

$$|\lambda_i| = 1, \quad \mu = \lambda_1 \bar{\lambda}_2,$$

and a change of basis will give (1.8). In case (ii)

$$\rho_i(e_1, e_2) = (\lambda_i e_2, \bar{\lambda}_i^{-1} e_1), \quad \bar{\lambda}_1 \lambda_2 = 1, \quad \mu = \overline{\lambda_2 / \lambda_1}.$$

A change of basis as above allows us to get  $\lambda_1 > 0$ , and hence (1.8). We refer to [5] for more details on the holomorphically reversible case.

**2. Non-linear involutions.** We consider a pair of non-linear anti-holomorphic involutions  $\rho_1, \rho_2$  and the anti-reversible map  $\sigma$ , which may be given by either convergent or formal power series at the origin of  $\mathbf{C}^n$ . We shall assume that the results of the previous section have been applied to the linear parts of these maps as needed, and write

$$(2.1) \quad \begin{aligned} \sigma(z) &= Mz + S(z), & S(z) &= \sum_{|J| \geq 2} b_J z^J, \\ \rho_i(z) &= L_i \bar{z} + R_i(z), & R_i(z) &= \sum_{|J| \geq 2} c_J \bar{z}^J, \quad i = 1, 2. \end{aligned}$$

Writing out the relations  $\sigma = \rho_1 \rho_2$  and  $\rho_i^2 = I$  gives

$$(2.2) \quad \begin{aligned} S &= L_1 \bar{R}_2 + R_1 \circ \rho_2, & M &= L_1 \bar{L}_2, & I &= L_i \bar{L}_i, \\ 0 &= L_1 \bar{R}_1 + R_1 \circ \rho_1, & 0 &= L_2 \bar{R}_2 + R_2 \circ \rho_2. \end{aligned}$$

To eliminate  $R_2$  and  $\bar{R}_2$ , multiply the first equation on the left by  $\bar{L}_1$ , and substitute this and its conjugate into the third equation. Then conjugation, left multiplication by  $L_1$ , and using the first equation of the second line gives the first of the following relations.

$$(2.3) \quad \begin{aligned} R_1 - MR_1 \circ \sigma &= S \circ \rho_2 + ML_1 \bar{S}, \\ R_2 \circ \sigma^{-1} - MR_2 &= S \circ \rho_2 + L_1 \bar{S} \circ \sigma^{-1}. \end{aligned}$$

The second equation is similarly derived.

As a first application in the generic case, suppose that  $\sigma$  has already been linearized,

$$(2.4) \quad \sigma(z) = Mz, \quad M = \text{diag}(\mu_1, \dots, \mu_n).$$

By (1.6) the eigenvalues (i) have modulus one, or (ii) occur in pairs. We have zero on the right hand sides of (2.3), and we may compose the second equation with  $\sigma$ . Substitution of the series in (2.1) into (2.3) gives

$$(2.5) \quad (1 - \bar{\mu}^J \mu_\alpha) c_{i\alpha J} = 0, \quad 1 \leq \alpha \leq n, \quad |J| \geq 2, \quad i = 1, 2,$$

$$\bar{\mu}^J = \bar{\mu}_1^{j_1} \dots \bar{\mu}_n^{j_n}.$$

The matrix  $M$  is *non-resonant*, for the full linearization problem, if none of the coefficients of the  $c_{i\alpha J}$ , i. e. the *divisors*, in (2.5) vanishes. It then follows that all the coefficients  $c_{i\alpha J}$  vanish, and that the  $\rho_i$  are anti-linear.

More generally, we assume that  $\sigma$  admits an invariant complex submanifold  $N$ , passing through the origin of  $\mathbf{C}^n$ , on which it may or may not be linearizable. As the switched curves show,  $N$  may not be left invariant by the  $\rho_i$ ; some further conditions are needed.

We choose coordinates

$$(2.6) \quad z = (z', z'') \in \mathbf{C}^l \times \mathbf{C}^{n-l},$$

so that  $N$  is the linear space  $z'' = 0''$ . Our further assumption is that this splitting is also preserved by the linear parts of the  $\rho_i$ . Then we may write

$$(2.7) \quad \sigma(z) = \begin{pmatrix} M'z' + S'(z', z'') \\ M''z'' + S''(z', z'') \end{pmatrix}, \quad \rho_i(z) = \begin{pmatrix} L'_i z' + R'_i(z', z'') \\ L''_i z'' + R''_i(z', z'') \end{pmatrix}.$$

We are assuming that  $S''(z', 0'') = 0''$ , and we want to show that  $R''_i(z', 0'') = 0''$ ,  $i = 1, 2$ . Taking the ''-components in (2.7), restricting to  $z'' = 0''$ , and using the condition on  $S''$  gives

$$(2.8) \quad \begin{aligned} (R''_1 - M''R''_1 \circ \sigma)_{z''=0''} &= (S'' \circ \rho_2)_{z''=0''}, \\ (R''_2 \circ \sigma^{-1} - M''R''_2)_{z''=0''} &= (S'' \circ \rho_2)_{z''=0''}. \end{aligned}$$

Suppose that  $R''_2(z', 0'') = O(|z'|^k)$ . Since  $S''(L_2 \bar{z}) = 0''$  and  $\bar{S}''(\sigma^{-1}(z)) = 0''$  for  $z'' = 0''$ , the second equation in (2.8) shows that the terms of order  $k$  satisfy

$$(2.9) \quad R''_{2,k}(M'^{-1}z', 0'') - M''R''_{2,k}(z', 0'') = 0''.$$

It follows that if we have the non-resonance conditions

$$(2.10) \quad \bar{\mu}^{-J} - \mu_\alpha = \bar{\mu}_1^{-j_1} \cdots \bar{\mu}_l^{-j_l} - \mu_\alpha \neq 0, \quad j_1 + \cdots + j_l \geq 2, \quad l < \alpha \leq n,$$

then we must have  $R''_{2,k}(z', 0'') = 0''$ ; and by induction  $R''_2(z', 0'') = 0''$ . A similar argument with the first equation in (2.8) now shows that  $R''_1(z', 0'') = 0''$ .

Suppose that, in addition,  $\sigma$  is linear on the invariant submanifold; that is  $S'(z', 0'') = 0'$  and  $S''(z', 0'') = 0''$ . Then the previous arguments show that  $R''_i(z', 0'') = 0''$ , and  $R'_i(z', 0'') = 0'$ , provided that we have the stronger non-resonance conditions

$$(2.11) \quad \bar{\mu}^{-J} - \mu_a = \bar{\mu}_1^{-j_1} \cdots \bar{\mu}_l^{-j_l} - \mu_a \neq 0, \quad j_1 + \cdots + j_l \geq 2, \quad 1 \leq a \leq n.$$

This proves the following.

PROPOSITION 1. *Suppose that  $\sigma$  and the linear parts of  $\rho_1$  and  $\rho_2$  are as in (2.7), leaving invariant the complex submanifold  $z'' = 0''$ .*

a) *If the conditions (2.10) hold, then  $\rho_1$  and  $\rho_2$  also leave  $z'' = 0''$  invariant.*

b) *If  $\sigma$  is also linear on  $z'' = 0''$ , and conditions (2.11) hold, then  $\rho_1$  and  $\rho_2$  are anti-linear on  $z'' = 0''$ .*

**3. Results on the map  $\sigma$ .** The formal arguments of the last section reduce some problems about the pair of involutions  $\rho_i$  to properties of the map  $\sigma$ , which we now take up.

We assume that  $\sigma$  has the form in (2.1), so that the linear part of  $\sigma$  preserves  $z'' = 0''$ . We want to find an  $l$ -dimensional  $\sigma$ -invariant submanifold tangent to  $z'' = 0''$  at 0. It will be given as the image of a map  $f : \mathbf{C}^l \rightarrow \mathbf{C}^n$ ,

$$(3.1) \quad f(w) = \begin{pmatrix} w + F'(w) \\ 0 + F''(w) \end{pmatrix}, \quad F(w) = \sum_{|J| \geq 2} c_J w^J,$$

with an induced map  $\hat{\sigma}$  on  $\mathbf{C}^l$ ,

$$(3.2) \quad \hat{\sigma}(w) = M'w + \hat{S}(w).$$

Writing out the functional equation  $\sigma \circ f = f \circ \hat{\sigma}$  gives

$$(3.3) \quad \begin{aligned} F'(\hat{\sigma}(w)) - M'F'(w) &= S'(f(w)) - \hat{S}(w), \\ F''(\hat{\sigma}(w)) - M''F''(w) &= S''(f(w)). \end{aligned}$$

For the problem of an invariant submanifold with linearization,  $\hat{S} = 0$ , we have, on the lefthand sides of (3.3), the divisors

$$(3.4) \quad \mu'^J - \mu_a = \mu_1^{j_1} \cdots \mu_l^{j_l} - \mu_a, \quad 1 \leq a \leq n, \quad |J| \geq 2.$$

If these are all non-zero, then a unique formal solution  $F$  exists. To state sufficient conditions for convergence of the type given by A. D. Brjuno [2], [7], we define

$$(3.5) \quad \omega(m) = \min\{|\mu'^J - \mu_a| : 1 \leq a \leq n, 2 \leq |J| \leq m\},$$

and choose an increasing sequence of integers  $1 = q_0 < q_2 < \cdots < q_j < \cdots$ , say  $q_j = 2^j$ . Brjuno's condition is then

$$(3.6) \quad \sum_{j=0}^{\infty} -q_j^{-1} \log \omega(q_{j+1}) < \infty.$$

By a theorem of Pöschel [7] the condition (3.6) implies that the power series solution  $f$  has some positive radius of convergence.

By combining these considerations on  $\sigma$  with proposition (1b), we get the following result.

**THEOREM 2.** *Suppose that  $\sigma = \rho_1\rho_2$ , where the (convergent) maps are as in (2.7), and the linear part of  $\sigma$  satisfies (2.4), (2.11), (3.5), and (3.6). Then there exists an  $l$ -dimensional analytic submanifold  $N$  tangent to the space  $z'' = 0''$  at 0 in  $\mathbf{C}^n$ , which is invariant by  $\sigma$ ,  $\rho_1$ , and  $\rho_2$ . These maps are simultaneously holomorphically (anti-)linearizable on  $N$ .*

For the existence of a cutting curve, we take  $l = 1$ , and  $|\mu_1| = 1$  in the theorem. This is related to Klingenberg's construction of asymptotic curves [4]. Even in the case  $n = 1$ , we do not know if the Brjuno condition is necessary for the result, i. e. whether an analogue of Yoccoz's theorem [14] holds in the anti-reversible category.

For the existence of switched curves, we take  $l = 2$  and  $\mu_2 = \bar{\mu}_1^{-1} \neq \mu_1$ , thus

$$(3.7) \quad \mu_1 = re^{i\theta}, \quad \mu_2 = r^{-1}e^{i\theta}, \quad \mu_\alpha = r_\alpha e^{i\theta_\alpha}, \quad 3 \leq \alpha \leq n,$$



with  $r > 1$ . We may simplify and sharpen the conditions for convergence. Since all positive or negative powers of  $r$  will be bounded away from the  $r_\alpha$ , we need only consider  $a = 1, 2$ , and  $|j_1 - j_2| = 1$  in (2.11) and (3.4). A simple consideration of all cases shows that we may replace (3.5) by

$$(3.8) \quad \omega(m) = \min\{|e^{2j\theta i} - 1| : 1 \leq j \leq m\}.$$

Of course, the theorem gives linearization on a 2-dimensional submanifold, which is a-priori stronger than just the existence of a pair of switched curves.

For the invariant submanifold problem without linearization, the question of convergence is much more delicate. Following [9] and [5] we consider a special situation, which we shall need later. We assume  $l = 2$  in (2.6) and

$$(3.9) \quad M' = \text{diag}(\mu_1, \mu_2), \quad \mu_1 \mu_2 = 1,$$

and set

$$(3.10) \quad F = (F', F'') = \sum c_{j_1 j_2} w_1^{j_1} w_2^{j_2}, \quad F' = (F_1, F_2), \quad F'' = (F_\alpha).$$

We assume that the terms of degree  $< k$  in  $F$  and in  $\hat{S} = (\hat{S}_1, \hat{S}_2)$  have been determined, and analyze the as yet undetermined terms of order  $k$  in (3.3). We find the divisors

$$(3.11) \quad \mu_1^{k-2j} - \mu_1, \quad \mu_1^{k-2j} - \mu_1^{-1}, \quad \mu_1^{k-2j} - \mu_\alpha,$$

for the coefficients of  $F_1, F_2$ , and  $F_\alpha$ , respectively. The first two vanish for  $k = 2j + 1$  and  $k = 2j - 1$ , respectively, and we assume that all the other divisors are non-zero. Then  $\hat{S}$  is uniquely determined at  $k$ -th order to make up for the lost terms. Also, we normalize  $F$  to make it unique.

More precisely, we introduce the “type” of a power series:  $\text{type}(w_1^{j_1} w_2^{j_2}) = j_1 - j_2$ , and write for  $k \in \mathbf{Z}$

$$(3.12) \quad [F]_k = \sum_{j_1 - j_2 = k} c_{j_1 j_2} w_1^{j_1} w_2^{j_2}.$$

Then we require

$$(3.13) \quad [F_1]_{+1} = 0, \quad [F_2]_{-1} = 0, \quad \hat{S}_1 = [\hat{S}_1]_{+1}, \quad \hat{S}_2 = [\hat{S}_2]_{-1}.$$

It follows that there is a unique so normalized formal map  $f$  satisfying  $\sigma \circ f = f \circ \hat{\sigma}$  and

$$(3.14) \quad \hat{\sigma}(w) = (\mu(t)w_1, \nu(t)w_2), \quad t = w_1 w_2,$$

where  $\mu(0) = \mu_1, \nu(0) = \mu_2$ , are the given eigenvalues.

In general, there is no convergence argument for  $f$ , even if  $\sigma$  is convergent [9]. However, if  $\sigma$  is convergent, and if

$$(3.15) \quad \mu(t)\nu(t) = 1, \quad |\mu_1| \neq 1,$$

where the first is a formal power series relation, then  $f$  does converge. In fact, the functional equation (3.3) is now of the form of (4.1) in [5], with  $\xi = w_1, \eta = w_2$ , and  $\zeta = 0$ . The powers of  $\mu_1$  and  $\mu_1^{-1}$  tend to 0 or  $\infty$ , and we assume that none equals

any  $\mu_\alpha$ . Then there are no small divisors, and the argument proving theorem (4.1) in [5] applies directly to give convergence here. The case  $l = n = 2$  is in [9].

The condition (3.15) may be interpreted as saying that the map  $\sigma$  is “integrable”, since it implies that the function  $w_1 w_2$  is  $\sigma$ -invariant.

The foregoing can be applied, if the map  $\sigma$  is symplectic, by the following.

**PROPOSITION 2.** *Let the holomorphic symplectic map  $\sigma$  have the form (2.6), (2.7), (2.4) with  $l = 2$ , and eigenvalues satisfying  $\mu_1 \mu_2 = 1$ ,  $|\mu_1| \neq 1$ , and no integral power of  $\mu_1$  equal to any  $\mu_\alpha$ . Suppose that the  $(z_1, z_2)$ -plane is symplectic at the origin. Then the formal transformation  $f$  just described converges, giving an invariant 2-manifold on which  $\sigma$  corresponds to the Birkhoff normal form map*

$$(3.16) \quad \hat{\sigma}(w) = (\mu(t)w_1, \mu(t)^{-1}w_2), \quad t = w_1 w_2, \quad \mu(0) = \mu_1.$$

For the proof it suffices to verify the first condition in (3.15). From  $\sigma^* \omega = \omega$  and  $\sigma \circ f = f \circ \hat{\sigma}$ , we see that the 2-form  $f^* \omega$  is  $\hat{\sigma}^*$ -invariant. To simplify notation we denote it by  $\omega$  and work on  $\mathbf{C}^2$ . By our assumptions,

$$(3.17) \quad \omega = a(w)dw_1 \wedge dw_2, \quad a(0) \neq 0.$$

From (3.14) the substitution  $\hat{\sigma}^* \omega = \omega$  gives

$$(3.18) \quad a(\mu(t)w_1, \nu(t)w_2)\varphi'(t) = a(w_1, w_2), \quad \varphi(t) = \mu(t)\nu(t)t.$$

We have  $\varphi'(0) = 1$ . By taking the part of type 0 in (3.18), we get an equation  $A(\varphi(t))\varphi'(t) = A(t)$ ,  $A(0) \neq 0$ . Integrating,  $B'(t) = A(t)$ ,  $B(0) = 0$ , gives

$$(3.19) \quad B(\varphi(t)) = B(t) = \sum_{j=1}^{\infty} b_j t^j, \quad b_1 \neq 0.$$

If  $\varphi(t) = t + c_k t^k + \dots$ , then substitution into (3.19) gives  $b_k = b_1 c_k + b_k$ . Hence,  $c_k = 0$ , and so  $\varphi(t) \equiv t$ , which proves (3.15).

The analogue of this result for flows is already in [9].

**4. Anti-reversible symplectic maps in  $\mathbf{C}^2$ .** We take  $w = (w_1, w_2) \in \mathbf{C}^2$  with  $\omega = dw_1 \wedge dw_2$ , and  $\rho_1, \rho_2, \sigma$  satisfying (0.5). We assume that  $\sigma$  is in normal form (3.16), and derive the form of  $\rho_1$  and  $\rho_2$ , in the two cases (1.12), using the formulae (2.3). In (2.1) we denote the two components by  $S = (S', S'')$  and  $R_i = (R'_i, R''_i)$ .

In case (i) the second equation in (2.3) is

$$(4.1) \quad \begin{aligned} R'_2 \circ \sigma^{-1} - \mu_1 R'_2 &= S' \circ \rho_2 + \lambda_1 \bar{S}' \circ \sigma^{-1}, \\ R''_2 \circ \sigma^{-1} - \mu_1^{-1} R''_2 &= S'' \circ \rho_2 - \lambda_1^{-1} \bar{S}'' \circ \sigma^{-1}. \end{aligned}$$

We use the type decomposition (3.12), but with respect to the conjugate variables  $(\bar{w}_1, \bar{w}_2)$ . Since  $[Q \circ \sigma^{-1}]_m = \overline{\mu(t)}^{-m} [Q]_m$ , multiplying through by  $\overline{\mu(t)}^m$  gives

$$(4.2) \quad \begin{aligned} (1 - \overline{\mu_1 \mu(t)^m}) [R'_2]_m &= \overline{\mu(t)^m} [S' \circ \rho_2]_m, \quad m \neq +1, \\ (1 - \overline{\mu_1^{-1} \mu(t)^m}) [R''_2]_m &= \overline{\mu(t)^m} [S'' \circ \rho_2]_m, \quad m \neq -1, \end{aligned}$$

because of the form (3.16) of  $\sigma$ .

We assume that  $[R'_2]_m, m \neq +1$ , and  $[R''_2]_m, m \neq -1$  have no terms of order  $< k$ . Since  $S \circ \rho_2(z) = S(L_2 \bar{z}) + O(R_2)$ , there are no terms of order  $k$  on the right hand

side. Since  $1 - \mu_1 \bar{\mu}_1^m \neq 0$ , if  $m \neq +1$ , and  $1 - \mu_1^{-1} \bar{\mu}_1^m \neq 0$ , if  $m \neq -1$ , it follows that the terms of order  $k$  vanish. Thus,  $\rho_2$  has the form

$$(4.3) \quad \rho_2(w) = (\lambda_2(\bar{t})\bar{w}_1, \kappa_2(\bar{t})\bar{w}_2).$$

A similar but simpler argument with the first equation in (4.1) shows that

$$(4.4) \quad \rho_1(w) = (\lambda_1(\bar{t})\bar{w}_1, \kappa_1(\bar{t})\bar{w}_2).$$

Using (4.3), (4.4) in the conditions  $\rho_i^* \omega = -\bar{w}$  gives  $\partial_{\bar{t}}(\lambda_i(\bar{t})\kappa_i(\bar{t})\bar{t}) = -1$ . Hence  $\lambda_i(\bar{t})\kappa_i(\bar{t}) = -1$ , and

$$(4.5) \quad \begin{aligned} \rho_i(w) &= (\lambda_i(\bar{t})\bar{w}_1, -\lambda_i(\bar{t})^{-1}\bar{w}_2), \quad i = 1, 2, \\ \mu(t) &= \lambda_1(-t)\bar{\lambda}_2(t). \end{aligned}$$

Similarly, the condition  $\rho_i^2 = I$  gives

$$(4.6) \quad \lambda_i(t) = \lambda_i^*(t), \quad \lambda_i^*(t) \equiv 1/\bar{\lambda}_i(-t),$$

where the bar means to conjugate the coefficients, and  $*$  is an involutive multiplicative homomorphism.

For further simplification, we conjugate by a transformation

$$(4.7) \quad f(w) = (\alpha(t)w_1, \alpha(t)^{-1}w_2),$$

which preserves  $\omega$  and commutes with  $\sigma$ . This preserves the form (4.5) and results in

$$(4.8) \quad \lambda_i(t) \mapsto \lambda_i(t)\beta(t)\beta^*(t), \quad i = 1, 2,$$

where  $\beta(t) = \alpha(-t)$ . If we choose  $\beta(t) = \lambda_2(t)^{-1/2}$ , or  $\beta(t) = (\lambda_1(t)\lambda_2(t))^{-1/2}$ , the properties of  $*$  show that we get, respectively  $\lambda_2(t) = 1$ , or

$$(4.9) \quad \lambda_1(t)\lambda_2(t) = 1.$$

Now we turn to case (ii). The equations (4.1) are now replaced by

$$(4.10) \quad \begin{aligned} R_2' \circ \sigma^{-1} - \mu_1 R_2' &= S' \circ \rho_2 + \lambda_1^{-1} \bar{S}'' \circ \sigma^{-1}, \\ R_2'' \circ \sigma^{-1} - \mu_1^{-1} R_2'' &= S'' \circ \rho_2 + \lambda_1 \bar{S}' \circ \sigma^{-1}. \end{aligned}$$

We now have (4.2) with  $m \neq -1$  in the first equation and  $m \neq +1$  in the second. The same arguments give

$$(4.11) \quad \rho_i(w) = (\lambda_i(\bar{t})\bar{w}_2, \kappa_i(\bar{t})\bar{w}_1).$$

Again the conditions  $\rho_i^* \omega = -\bar{w}$  and  $\rho_i^2 = I$  give

$$(4.12) \quad \begin{aligned} \rho_i(w) &= (\lambda_i(\bar{t})\bar{w}_2, \lambda_i(\bar{t})^{-1}\bar{w}_1), \quad \bar{\lambda}_i(t) = \lambda_i(t), \quad i = 1, 2, \\ \mu(t) &= \lambda_1(t)/\lambda_2(t). \end{aligned}$$

Conjugation by  $f$  in (4.7) again preserves the form (4.12) and results in

$$(4.13) \quad \lambda_i(t) \mapsto \lambda_i(t)(\alpha(t)\bar{\alpha}(t))^{-1}, \quad \lambda_1(t)\lambda_2(t) \mapsto \lambda_1(t)\lambda_2(t)(\alpha(t)\bar{\alpha}(t))^{-2}.$$

To preserve the reality condition in (4.12), we restrict to  $\alpha(t) = \bar{\alpha}(t)$ . The symplectic change of coordinates  $(w_1, w_2) \mapsto (w_2, -w_1)$  results in

$$(4.14) \quad \mu(t) \mapsto \mu(-t)^{-1}, \quad \lambda_i(t) \mapsto -\lambda_i(-t)^{-1}.$$

By extracting a real square root or fourth root, we can achieve either  $\lambda_2(t) = 1$ , or

$$(4.15) \quad \lambda_1(t)\lambda_2(t) = \epsilon = \pm 1, \quad \epsilon = \text{sgn}(\mu_1).$$

If we restrict to the surface  $M_i = FP(\rho_i)$  in case (ii), then  $w = \rho_i(w)$ , and (4.12) gives  $w_2 = \lambda_i(\bar{t})^{-1}\bar{w}_1$ , and  $\lambda_i(\bar{t}) = \lambda_i(t)$ . If we multiply the second component of (4.12) by  $w_1$  and set  $s = |w_1|^2$ , then we get  $s = t\lambda_i(t)$ , which is an invertible power series with real coefficients. It follows that  $t = w_1w_2$  is real when restricted to  $M_i$ . Let  $t = s\phi_i(s)$  be the inverse function, and define a real function of  $s$  by

$$(4.16) \quad r_i(s) = \int_0^s \lambda_i(s\phi_i(s))^{-1} ds.$$

This results in the following equation for  $M_i$

$$(4.17) \quad M_i : w_2 = \partial_{w_1} r_i, \quad r_i = r_i(|w_1|^2),$$

which is precisely (0.6), (0.7), (0.8). This also anticipates the developments of the next section.

As in section 1 we consider the complex curve  $w_1w_2 = c$ , for small real  $c$ . It is invariant under both  $\rho_1$  and  $\rho_2$ , and

$$(4.18) \quad \{w_1w_2 = c\} \cap FP(\rho_i) = \{w_1w_2 = c, |w_1|^2 = c\lambda_i(c)\}.$$

If  $\epsilon = +1$ , we choose  $c$  so that  $c\lambda_1(c) > 0$ . Then

$$(4.19) \quad c\lambda_1(c) \leq |w_1|^2 \leq c\lambda_2(c), \quad w_2 = c/w_1,$$

defines an analytic annulus  $A_c$  on the curve with boundary on  $M_1 \cup M_2$ , which shrinks to the origin as  $c \rightarrow 0$ . The annular modulus (ratio of inner to outer radius) is  $|\lambda_1(c)|$ , which approaches  $|\lambda_1(0)| \neq 0$ . These annuli  $A_c$  sweep out a three dimensional manifold lying on the real analytic levi-flat set  $\text{Im}(w_1w_2) = 0, \text{Re}(w_1w_2) > 0$ .

In case (i) the transformation of  $\sigma$  into the normal form (3.16) exists if the linear part of  $\sigma$  is non-resonant, but may not converge [9]. In case (ii) it always exists and converges [9]. Combining this with the arguments just given yields the following.

**THEOREM 3.** *Let  $M_1$  and  $M_2$  be a pair of analytic real Lagrangian surfaces in  $\mathbb{C}^2$ , intersecting transversely at the origin. Let  $\rho_1$  and  $\rho_2$  be the associated anti-holomorphic involutions, with linear parts satisfying (1.13) in case (i), or (1.14) in case (ii). Suppose the linear part of  $\sigma = \rho_1\rho_2$  is non-resonant. Then in case (i) there exists a formal symplectic transformation taking the pair  $\rho_i$  into the form (4.5). In case (ii) there exists a holomorphic symplectic transformation taking the  $\rho_i$  into the form (4.12).*

By combining proposition (2) and theorem (3), we may carry some of the above results over to the higher dimensional case. For example, we have the following result.

**THEOREM 4.** *Let  $M_1$  and  $M_2$  be a pair of analytic real Lagrangian surfaces in  $\mathbb{C}^{2n}$ , with symplectic form (0.5), intersecting transversely at the origin. Suppose that*

$\sigma$  satisfies the conditions of proposition (2) with  $\mu_1 > 0$ . Suppose that  $\rho_1, \rho_2$  satisfy the conditions of proposition (2a). Then there exists a real analytic one-parameter family  $A_c, c_0 > c > 0$ , of analytic annuli in  $\mathbb{C}^{2n}$  bounding on  $M_1 \cup M_2$ , and shrinking to the intersection point as  $c \rightarrow 0$ .

This family of analytic annuli contributes to the local holomorphic hull of  $M_1 \cup M_2$ .

**5. Real Lagrangians and generating functions.** As in [10] we work on the holomorphic cotangent bundle with its canonical structure,

$$(5.1) \quad T^*(\mathbb{C}^n) \cong \mathbb{C}^{2n} \ni (z, p), \quad \theta = p \cdot dz = \sum_{\alpha=1}^n p_\alpha dz_\alpha, \quad \omega = d\theta.$$

Every real-valued function  $r(z)$  on  $\mathbb{C}^n$  gives a real Lagrangian  $M \subset T^*(\mathbb{C}^n)$ , namely the graph of  $\partial r$ ,

$$(5.2) \quad M : R_\alpha \equiv p_\alpha - \partial_\alpha r = 0, \quad \partial_\alpha = \partial / \partial z_\alpha, \quad 1 \leq \alpha \leq n,$$

since restricting to  $M$  gives  $Re(\omega) = dRe(\theta) = d(\partial r + \bar{\partial} r) / 2 = 0$ . A (1,0)-vector  $(dz, dp)$  is tangent to  $M$ , if it satisfies

$$(5.3) \quad \partial R_\alpha = 0, \quad \partial \bar{R}_\alpha = - \sum_{\beta} \partial_\beta \partial_{\bar{\alpha}} r dz_\beta = 0.$$

This means that its projection  $dz$  is in the nullspace of the Levi form, or (1,1)-hessian of  $r$ . In particular,  $M$  is totally real if and only if  $r$  has non-degenerate Levi form. All real analytic, totally real, real Lagrangians  $M$  are locally equivalent. This follows from applying the real analytic Darboux theorem to  $(M, Im(\omega))$  and then complexifying back to the ambient space. For a single smooth  $M$ , we may take  $r = z \cdot \bar{z}$ , to arbitrarily high order.

Every holomorphic function  $h(z)$  generates the symplectic map  $(z_\alpha, p_\alpha) \mapsto (z_\alpha, p_\alpha + \partial_\alpha h)$ . This transforms  $p_\alpha = \partial_\alpha r$  into  $p_\alpha = \partial_\alpha (r - h - \bar{h})$ , and allows us to remove any purely holomorphic and anti-holomorphic terms in  $r$ . If  $M_1$  and  $M_2$  are two real Lagrangians intersecting over  $z = 0$ , then their functions  $r_1$  and  $r_2$  have the same linear parts. By so transforming  $M_1$  and  $M_2$ , we may assume

$$(5.4) \quad \begin{aligned} r_1 &= \sum_{\alpha, \beta=1}^n b_{\alpha\bar{\beta}} z_\alpha \bar{z}_\beta + 2Re(a_{\alpha\beta} z_\alpha z_\beta) + \dots, \\ r_2 &= z \cdot \bar{z} + \dots, \quad a_{\alpha\beta} = a_{\beta\alpha}, \quad b_{\alpha\bar{\beta}} = \bar{b}_{\beta\bar{\alpha}}. \end{aligned}$$

One may apply a linear symplectic transformation to simplify further the quadratic terms in (5.4), and then use a generating function to construct a symplectic transformation to simplify the higher order terms (see [10]). However, this seems to be rather complicated, so we return to the approach of sections 1 and 2 focusing on the involutions  $\rho_i$  and map  $\sigma$ .

We restrict to the case,  $n = 1$ , so that (5.4) gives (0.6). We drop the higher order terms in (0.6) and work with quadratic parts,

$$(5.5) \quad \begin{aligned} r_1 &= az^2 + bz\bar{z} + \bar{a}\bar{z}^2, \quad b \neq 0, \\ r_2 &= z\bar{z}, \quad (b, a) \neq (1, 0). \end{aligned}$$

This gives the linear surfaces

$$(5.6) \quad \begin{aligned} M_1 &: p = r_{1z} = 2az + b\bar{z}, \\ M_2 &: p = r_{2z} = \bar{z}. \end{aligned}$$

By solving these equations for  $(z, p)$  in terms of  $(\bar{z}, \bar{p})$ , we get the matrices as in (2.1) for  $\rho_1, \rho_2$ , and  $\sigma$ ,

$$(5.7) \quad L_1 = b^{-1} \begin{bmatrix} -2\bar{a} & 1 \\ \Delta & 2a \end{bmatrix}, \quad L_2 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad M = L_1 \bar{L}_2 = b^{-1} \begin{bmatrix} 1 & -2\bar{a} \\ 2a & \Delta \end{bmatrix},$$

$$(5.8) \quad \det(M - \mu I) = \mu^2 - 2\delta\mu + 1, \quad 2b\delta = \Delta + 1, \quad \Delta = b^2 - 4|a|^2.$$

For the two cases (1.12) we have: (i)  $\mu\bar{\mu} = 1$  if and only if  $\delta^2 < 1$ ; and (ii)  $\mu = \bar{\mu}$  if and only if  $\delta^2 > 1$ . The case  $\epsilon = +1$  in case (ii) is equivalent to  $\delta > 1$ , since the trace of  $M$  must be positive. From this and the results of the last section we readily derive the theorem stated in the introduction.

#### REFERENCES

- [1] E. BISHOP, *Differentiable manifolds in complex Euclidean space*, Duke Math. J., 32 (1965), pp. 1–22.
- [2] A. D. BRJUNO, *Analytic form of differential equations*, Trans. Moscow Math. Soc., 25 (1971), pp. 131–288, 26 (1972), pp. 199–239.
- [3] X. GONG, *Anti-holomorphically reversible holomorphic maps that are not holomorphically reversible*, to appear.
- [4] W. KLINGENBERG, *Asymptotic curves on real analytic surfaces in  $\mathbf{C}^2$* , Math. Ann., 273 (1985), pp. 149–162.
- [5] J. MOSER AND S. WEBSTER, *Normal forms for real surfaces in  $\mathbf{C}^2$  near complex tangents and hyperbolic surface transformations*, Acta Math., 150 (1983), pp. 255–296.
- [6] G. PFEIFFER, *On the conformal mapping of curvilinear angles*, Trans. AMS, 18 (1917), pp. 185–198.
- [7] J. PÖSCHEL, *Invariant manifolds of complex analytic mappings near fixed points*, Les Houches, Session XLIII, vol. II (1984), pp. 949–964.
- [8] N. SIBONY, personal communication.
- [9] C. SIEGEL AND J. MOSER, *Lectures on Celestial Mechanics*, Springer-Verlag (1971).
- [10] S. WEBSTER, *Holomorphic symplectic normalization of a real function*, Ann. Sc. Norm. Sup. Pisa, vol. XIX (1992), pp. 69–86.
- [11] S. WEBSTER, *A note on extremal discs and double valued reflection*, AMS. Contemp. Math., 205 (1997), pp. 271–276.
- [12] B. WEINSTOCK, *On the polynomial convexity of the union of two maximal totally real subspaces of  $\mathbf{C}^n$* , Math. Ann., 282 (1988), pp. 131–138.
- [13] H. WEYL, *The Classical Groups*, Princeton University Press (1946).
- [14] J. YOCCOZ, *Théorème de Siegel, nombres de Bruno et polynômes quadriques*, Astérisque, 231 (1995), pp. 3–88.