

On the dimension of contact loci and the identifiability of tensors

Edoardo Ballico, Alessandra Bernardi and Luca Chiantini

Abstract. Let $X \subset \mathbb{P}^r$ be an integral and non-degenerate variety. Set $n := \dim(X)$. We prove that if the $(k+n-1)$ -secant variety of X has (the expected) dimension $(k+n-1)(n+1)-1 < r$ and X is not uniruled by lines, then X is not k -weakly defective and hence the k -secant variety satisfies identifiability, i.e. a general element of it is in the linear span of a unique $S \subset X$ with $\#(S)=k$. We apply this result to many Segre-Veronese varieties and to the identifiability of Gaussian mixtures $\mathcal{G}_{1,d}$. If X is the Segre embedding of a multiprojective space we prove identifiability for the k -secant variety (assuming that the $(k+n-1)$ -secant variety has dimension $(k+n-1)(n+1)-1 < r$, this is a known result in many cases), beating several bounds on the identifiability of tensors.

1. Introduction

The computation of the decompositions of tensors and the related evaluation of their ranks recently received great interest from people working in algebraic geometry. Indeed, a geometric analysis on projective spaces of tensors, Segre embeddings and their secant varieties produced currently several new results in the field.

From the geometric point of view, the study of decompositions and ranks can be turned to a general problem. For any (irreducible) algebraic variety X which spans a projective space \mathbb{P}^r , one can define the X -rank $r_X(q)$ of a point $q \in \mathbb{P}^r$ as the minimal cardinality of a finite set $S \subset X$ such that $q \in \langle S \rangle$, where $\langle \cdot \rangle$ denotes the linear span. Any set S for which the minimum is attained is a *minimal X -decomposition* of q .

In tensor analysis, one interprets \mathbb{P}^r as a space of tensors of given type $(n_1+1) \times \dots \times (n_s+1)$ (mod scalars), and X as the locus of tensors of rank 1. In geometric terms, X is the multiprojective product $\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_s}$, naturally embedded in \mathbb{P}^r via the Segre map. Then, the notions of X -rank and minimal decompositions match with the usual multilinear algebra notion. The set of tensors of fixed rank

k corresponds to an open subset of the k -th secant variety $\sigma_k(X)$ of X . Properties of the rank decompositions can be derived from geometric properties of X , which determine some features of $\sigma_k(X)$. Examples of such analysis are contained in several papers, e.g. see [8], [10], [25], [26], [36], [41], and the book [38].

In this paper, we study the problem of the *identifiability* of generic points $q \in \mathbb{P}^r$ of given X -rank k . A point q is X -identifiable when the set $\mathcal{S}(q, X)$ of minimal X -decompositions of q is a singleton; in any case, the cardinality of $\mathcal{S}(q, X)$ is called the *secant degree of q* or the k -th secant degree of q . In tensor analysis the identifiability of a tensor is fundamental for algorithms of data compression and for the analysis of mixture models (see [33]–[35], [43]). We will say that X is k -*identifiable* if a generic point of $\sigma_k(X)$ is X -identifiable.

We first work over an algebraically closed field of characteristic 0. Our investigation is based on the notion of weak defectivity introduced by the last author and C. Ciliberto in [25]. A projective variety X is *weakly k -defective* if hyperplanes of \mathbb{P}^r , which are tangent to X at k general points, are indeed tangent along a positive dimensional subvariety. It turns out (see [26, Theorem 2.5]) that when $r > k(n+1) - 1$ then X is k -identifiable, unless it is weakly k -defective (unfortunately the reverse is not true). Therefore, the knowledge on the non- k weak defectivity provides an effective way to determine geometrically the identifiability of generic points. This approach has been applied to tensors and linear systems of tensors in a series of papers ([1], [16], [18], [19], [29]–[31]), where identifiability problems are partially or totally solved.

A challenging task is therefore to get criterions that can establish identifiability (or not) in the cases when weak defectivity arises. To this purpose there is a related notion that is very helpful, meaning the *tangential weak defectivity*. Our new analysis starts with the following observation (see [27]). If X is a weakly k -defective variety of dimension n , then a general choice of a set A of k points on X determines a positive dimensional *tangential k -contact locus* $\Gamma_k(A)$: the tangency locus of the span of the tangent spaces to X at the points of A . If for a general choice of A we have that $\dim \Gamma_k(A) > 0$ then X is *tangential k -weakly defective*. It turns out that the k -th secant degree of a general element of X is related with the k -th secant degree of the contact locus $\Gamma_k(A)$ for a general choice of A (cf. Lemma 4.2, Corollary 4.3, [26, Theorem 2.4] and part (iii) of [27, Proposition 3.9]). We will exploit in more details the relation in Section 4.

Moreover, when $r \geq (k+1)(n+1) - 1$, then the weak k -defectivity of X implies the weak $(k+1)$ -defectivity. Moreover, we prove that, under some geometric assumption on X , if we take a generic subset A as above and let A' be obtained by adding to A a generic point of X , then the dimension of the tangential contact locus $\Gamma_{k+1}(A')$ is strictly bigger than the dimension of $\Gamma_k(A)$. As the dimension of

contact loci is bounded by $n-1$, by repeating the previous construction we obtain the following result.

Theorem (see Theorem 5.1) *Let $X \subset \mathbb{P}^r$ be an integral and non-degenerate n -dimensional variety and let k be a positive integer. Assume that X is not uniruled by lines and that $\dim \sigma_{k+n-1}(X) = (k+n-1)(n+1) - 1 < r$. Then X is not weakly j -defective for every $j \leq k$.*

It follows from the previous analysis that, with the assumptions of Theorem 5.1, X is j -identifiable for every $j \leq k$.

The result can be directly applied e.g. to Segre-Veronese embeddings of products of projective spaces, provided that the multidegree (d_1, \dots, d_s) satisfies $d_i > 1$ for all i . When some degree d_i is 1, as in the case of Segre embeddings (where $d_i = 1$ for all i), then Theorem 5.1 does not apply directly. However, for the Segre product we can modify the argument of Theorem 5.1, and prove that the same conclusion holds when for each $i \in \{1, \dots, s\}$ there is $j \in \{1, \dots, s\} \setminus \{i\}$ such that $n_j = n_i$ (see Theorem 5.3).

Combining Theorem 5.3 with the main result in [23], for the Segre embedding X of $s \geq 5$ copies of \mathbb{P}^1 , we find that X is not weakly k -defective, hence it is k -identifiable, whenever $(k+s-1)(s+1) < 2^s - 1$ (see Corollary 5.4).

Corollary 5.4 improves our previous knowledge on the weak defectivity of Segre embeddings of many copies of \mathbb{P}^1 , hence on the identifiability of generic binary tensors. Indeed it extends the range where we know that k -identifiability holds up to $k \leq \frac{s^2-1}{s+1} - s$, which beats all previously known bounds (see Section 4 of [20]) as s grows. Since identifiability cannot hold, for trivial dimensional reasons, when $k > s^2/(s+1) - 1$, then we are quite close to a complete analysis of the generic identifiability of binary tensors. We also get almost optimal results for the Segre embedding of $(\mathbb{P}^m)^s$, $m > 1$, see Corollary 5.5 (we use the very strong [9, Theorem 3.1]).

See Example 5.8 for the case of Gaussian mixtures $\mathcal{G}_{1,d}$.

When the assumptions of Theorem 5.1 does not hold, one cannot conclude the non weak k -defectivity of X . Yet, even if X is weakly k -defective, it can still be k -identifiable. We prove the following result.

Theorem (see Theorem 5.9) *Let X be the Segre embedding of a product $\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_a}$ in \mathbb{P}^r , with $n = \sum n_i = \dim(X)$. Let k be a positive integer. Assume that $\dim \sigma_{k+n-1}(X) = (k+n-1)(n+1) - 1 < r$ (in particular X is not $(k+n-1)$ -defective). Then X is j -identifiable for every $j \leq k$.*

It would be interesting to overcome the assumption “ X not uniruled by lines” of Theorem 5.3 in other cases, apart the Segre varieties covered by Theorem 5.9,

because it would expand the known cases of identifiability (e.g. for Grassmannians see [6], [17], [21], [22]; for the tangential variety of the Veronese varieties see [7]). We just recall that conjecturally only four Grassmannians have some defective secant variety.

We stress that, in particular, Remark 5.10 suggests that if one is interested in the k -identifiability of some specific Segre product then the computational procedure induced by Theorem 5.9 can be faster than previously known direct procedures.

In the last section of the paper we work over an algebraically closed base field \mathbb{K} of arbitrary characteristic. To include varieties X defined in positive characteristic, we will assume that X is *not very strange* (see Definition 6.1). Under this mild assumption, we prove the following.

Theorem 1.1. *Let M be an integral projective variety and \mathcal{L}, \mathcal{R} line bundles on M with \mathcal{L} very ample. Set $n := \dim M$. Fix a linear subspace $V \subseteq H^0(\mathcal{L})$ such that the associated map $j: M \rightarrow \mathbb{P}^r$ is an embedding with image not very strange. Fix an integer $k \leq \min\{h^0(\mathcal{R}), \dim V - n - 2\}$. Let $W \subseteq H^0(\mathcal{L} \otimes \mathcal{R})$ be the image of the multiplication map $V \otimes H^0(\mathcal{R}) \rightarrow H^0(\mathcal{L} \otimes \mathcal{R})$ and let $X \subset \mathbb{P}^r$, $r = \dim W - 1$, be the embedding induced by W . Then $\sigma_k(X)$ has dimension $kn + k - 1$ and $\sharp(\mathcal{S}(q, X)) = 1$ for a general $q \in \sigma_k(X)$.*

We will apply Theorem 1.1 to the Segre-Veronese embeddings of a multiprojective space in arbitrary characteristic (see Example 6.3), where we do not have an analogue of [26, Theorem 2.5], hence the analysis of the previous section does not apply. Notice that Theorem 1.1 does not apply to Segre embeddings, because it requires \mathcal{L} very ample (or at least birationally very ample) and $h^0(X, \mathcal{R}) > 1$.

The authors wish to thank C. Bocci, for his help to understand the case of almost unbalanced tensors (see Example 4.10 below). The authors also thank the anonymous referee for pointing out an improvement in the formula after Corollary 5.4.

2. General notation

In this section, we work with projective spaces \mathbb{P}^r defined over an algebraically closed field \mathbb{K} .

Let $X \subset \mathbb{P}^r$ be an integral and non-degenerate n -dimensional variety. We will denote with X_{reg} the set of regular points in X . For any $x \in X_{\text{reg}}$ we will denote with $T_x X$ the tangent space to X at x .

For any integer $k \geq 2$ we will denote with $\sigma_k(X)$ the k -th secant variety of X which is an irreducible projective variety. If $k(n+1) - 1 \leq r$, then $k(n+1) - 1$ is the expected dimension of $\sigma_k(X)$ and X is k -defective if $\dim \sigma_k(X) < k(n+1) - 1$.

We will often use the fact that if X is not k -defective, then, for all $k' \leq k$, X is not k' -defective.

Definition 2.1. For $q \in \mathbb{P}^r$ general, the X -rank $r_X(q)$ of a point $q \in \mathbb{P}^r$ is the minimal cardinality of a finite set $A \subset X$ such that $q \in \langle A \rangle$.

We define $\mathcal{S}_X(q)$ as the collection of subsets $A \subset X$ of cardinality $r_X(q)$ such that $q \in \langle A \rangle$. When $\mathcal{S}_X(q)$ is finite, we will write $\alpha_X(q)$ for the cardinality of $\mathcal{S}_X(q)$.

Remark 2.2. Since X is irreducible, then the secant varieties $\sigma_k(X)$ are irreducible. Moreover, a general point of $\sigma_k(X)$ has X -rank k .

By definition, the set $\mathcal{S}_X(q)$ is finite for $q \in \sigma_k(X)$ general if and only if $\dim \sigma_k(X) = k(n+1) - 1 \leq r$. In this case, we will write simply α_k for the cardinality of $\mathcal{S}_X(q)$, where $q \in \sigma_k(X)$ is generic. Such an α_k is often called the k -th secant degree of X .

The following result holds (with the same elementary proof) in arbitrary characteristic.

Proposition 2.3. *Assume $\dim \sigma_{k+1}(X) = (k+1)(n+1) - 1$ and $\alpha_k > 1$. Fix a general $q \in \sigma_{k+1}(X)$, any $A \in \mathcal{S}(q)$ and any $a \in A$. There are at least α_k elements $B_1, \dots, B_{\alpha_k} \in \mathcal{S}(q)$ (one of them being A) such that $a \in B_i$ for all $i = 1, \dots, \alpha_k$.*

Proof. Since $\dim \sigma_{k+1}(X) = (k+1)(n+1) - 1$, we have $\dim \sigma_k(X) = k(n+1) - 1$ and hence α_k is a well-defined finite positive integer. Fix a general $q \in \sigma_{k+1}(X)$ and take any $A \in \mathcal{S}(q)$ and any $a \in A$. Set $E := A \setminus \{a\}$. Since q is general in $\sigma_{k+1}(X)$ and $\dim \sigma_{k+1}(X) = (k+1)(n+1) - 1$, a dimensional count gives that for each $D \in \mathcal{S}(q)$ the set D is a general element of the $(k+1)$ -symmetric product of X . Thus E is a general subset of X with cardinality k . Since $\#(A) = r_X(q)$, we have $q \notin \langle A' \rangle$ for any $A' \subsetneq A$; in particular $q \notin \langle E \rangle$. Thus $\langle \{a, q\} \rangle$ is a line, which meets $\langle E \rangle$ in a unique point q' (because $\langle E \rangle + \langle a, q \rangle = \langle A \rangle$) and $q' \notin \langle E' \rangle$ for any $E' \subsetneq E$. For a general $q \in \langle A \rangle$, q' is a general element of $\langle E \rangle$. Since E is general, q' is general in $\sigma_k(X)$. Thus $r_X(q') = k$, $E \in \mathcal{S}(q')$ and $\#(\mathcal{S}(q')) = \alpha_k > 1$. Fix any $F \in \mathcal{S}(q')$. We have $q \in \langle \{F, a\} \rangle$. Since $r_X(q) = k+1$, we have $a \notin F$ and $F \cup \{a\} \in \mathcal{S}(q)$. Now we have exactly α_k such F 's, say F_1, \dots, F_{α_k} , the B_i of the statement are $B_i = \{F_i, a\}$ for $i = 1, \dots, \alpha_k$ concluding the proof. \square

3. Weak defectivity and identifiability

In this section, as well as in the next two sections, we assume that the algebraically closed base field \mathbb{K} has characteristic 0.

For a subset A of cardinality k contained in the regular locus X_{reg} we set:

$$M_A := \langle \cup_{x \in A} T_x X \rangle.$$

By Terracini's lemma ([8, Corollary 1.11]), for a general choice of A the space M_A is the tangent space to $\sigma_k(X)$ at a general point $u \in \langle A \rangle$. Thus $\dim M_A = \min\{k(n+1)-1, \dim \langle X \rangle\}$ for A general, unless X is k defective.

Definition 3.1. The *tangential k -contact locus* $\Gamma_k(A)$ is the closure in X of the union of all the irreducible components which contain at least one point of A , of the locus of points of X_{reg} where M_A is tangent to X .

We will write $t_k := \dim \Gamma_k(A)$ for a general choice of A .

We say that X is *weakly k -defective* if $t_k > 0$.

We will write just Γ_k instead of $\Gamma_k(A)$, when A is general.

With the terminology just introduced we have the following remark:

Remark 3.2. [25, Remark 3.1(iii) at p. 159] If $\sigma_{k+1}(X) \not\subseteq \mathbb{P}^r$ has the expected dimension $(k+1)\dim X + (k+1) - 1 < r$ and X is weakly k -defective, then X is also weakly $(k+1)$ -defective.

We will need the following result of [27].

Proposition 3.3. *For a general choice of $A \subset X_{\text{reg}}$, the algebraic set $\Gamma_k := \Gamma_k(A)$ is equidimensional and either it is irreducible or it has exactly k irreducible components, each of them containing a different point of A .*

Proof. It is Proposition 3.9 of [27]. (Notice that our $\sigma_k(X)$ is called the $(k-1)$ -th secant variety in [27], so that all the k 's there should be here replaced by $k+1$, in order to apply the results of [27] to our terminology). \square

Definition 3.4. Following [26] and [27], we say that Γ_k is of the *type I* (resp. *type II*) if Γ_k is irreducible (resp. it is not irreducible) for a general choice of A .

The fundamental technical results of the present paper is the following lemma. It shows how one can control the growth of the dimension of contact loci. Mainly, one must take care of the case of contact loci of type II, which are not irreducible.

Lemma 3.5. *Let $X \subset \mathbb{P}^r$ be an irreducible projective variety such that $\dim \sigma_{k+1}(X) = (k+1)(n+1) - 1 < r$ and assume that X is weakly k -defective. Then X is weakly $(k+1)$ -defective. Let B (resp. A) be a general subset of k (resp. $k+1$) points of X . Call Γ_k (resp. Γ_{k+1}) the tangential contact locus of X at the points of B (resp. A). Then either:*

- $\dim \Gamma_k < \dim \Gamma_{k+1}$; or
- both Γ_k and Γ_{k+1} have type II, each of the irreducible components of Γ_k and Γ_{k+1} is a linear space and these components are linearly independent, i.e.

$$\dim(\Gamma_{k+1}) = \dim(\Gamma_k) + \dim \Gamma_{k+1} = (k+1)(\dim \Gamma_k + 1) - 1.$$

Proof. Since $\dim \sigma_{k+1}(X) = (k+1)(n+1) - 1$, we have $\dim \sigma_k(X) = k(n+1) - 1$. By Remark 3.2, X is weakly $(k+1)$ -defective. By [25, first two lines after Definition 1.2], X is not i -defective for $i=1, \dots, k+1$.

By [27, Proposition 3.9] we know that both Γ_k and Γ_{k+1} are equidimensional, that A is in the smooth locus of Γ_{k+1} , that B is in the smooth locus of Γ_k and that either Γ_k (resp. Γ_{k+1}) is irreducible or it has exactly k (resp. $k+1$) irreducible components, each of them containing exactly one point of B (resp. A).

Since B is general, we may assume that $B \subset A$. Thus $\Gamma_k \subseteq \Gamma_{k+1}$. Since Γ_k is a proper subvariety of X and Γ_{k+1} contains B plus a general point of X , then $\Gamma_k \neq \Gamma_{k+1}$. Thus if either Γ_k or Γ_{k+1} is irreducible, then $t_k < t_{k+1}$ (where t_k is defined in Definition 3.1).

Now assume that both Γ_k and Γ_{k+1} are reducible, hence of type II, and that $t_k = t_{k+1} =: t$. Let $B = \{P_1, \dots, P_k\}$ and $A = \{P_{k+1}\} \cup B$. Call Γ^i the component of Γ_{k+1} passing through P_i , so that also $\Gamma_k = \Gamma^1 \cup \dots \cup \Gamma^k$. Call Π^i the linear span of Γ^i . Of course, by the generality of A , all the Π^i 's have the same dimension. Assume that $\dim \Pi^i > \dim \Gamma^i$ and let us show that we get a contradiction.

For $i=2, \dots, k+1$ call L_i the span of $\Pi^1 \cup \dots \cup \Pi^i$, which is also the span of $\Gamma^1 \cup \dots \cup \Gamma^i$. By [26, Proposition 2.5] we know that, for all i , $\dim(L_i) \geq it + i - 1$, since X is not i -defective. Moreover, by [27, Proposition 3.9] we know that $\dim(L_k) = kt + k - 1$ and $\dim(L_{k+1}) = (k+1)t + k$. If $\dim(L_{k-1}) > (k-1)t + k - 2$, then by Grassmann formula, the intersection of L_{k-1} and Π_k has dimension greater than $(k-1)t + k - 2 + \dim(\Pi_k) - kt - k + 1 = \dim(\Pi_k) - t - 1$, so that, by Grassmann formula again,

$$\dim(L_{k+1}) < \dim(L_k) + \dim(\Pi_{k+1}) - \dim(\Pi_i) + t + 1 = (k+1)t + k,$$

a contradiction. Thus $\dim(L_{k-1}) = (k-1)t + k - 2$. Arguing by induction, we obtain that $\dim(L_i) = it + i - 1$ for $i=2, \dots, k+1$. In particular, $\dim(\Pi^1 \cap \Pi^2) = 2 \dim(\Pi^1) - 2t - 1 \geq 0$. By the generality of A again, $\dim(\Pi^i \cap \Pi^j) = \dim(\Pi^1 \cap \Pi^2)$ for all i, j . Thus Π^3 intersects L_2 in dimension at least $\dim \Pi^1 \cap \Pi^2$. It follows that L_3 has dimension

$$\begin{aligned} \dim(L_3) &\leq \dim(L_2) + \dim(\Pi^3) - 2 \dim(\Pi^1) + 2t + 1 \\ &= (2t + 1) - \dim(\Pi^1) + 2t + 1 < 3t + 2, \end{aligned}$$

which yields the claimed contradiction. \square

4. Remarks on the secant degree of contact loci

We collect in this sections some remarks on the general *tangential* contact loci Γ_k of X , which clarify the relations between the k -th secant degree of X and the k -th secant degree of a general Γ_k . To do that, we need to distinguish the case where Γ_k is of type I (i.e. irreducible) from the case where Γ_k is of type II.

Notation 4.1. *For the rest of the section, let $X \subset \mathbb{P}^r$ be an integral and non-degenerate n -dimensional variety and let k be a positive integer. Assume that X is weakly k -defective with*

$$\dim \sigma_{k+n-1}(X) = (k+n-1)(n+1) - 1 < r.$$

Call Γ_k the tangential k -contact locus of X at a general set $A = \{P_1, \dots, P_k\}$ of k points of X , and call Π_k the linear span of Γ_k .

The following technical result is the analogue of [26, Theorem 2.4] for the tangential contact locus Γ_k . It is implicitly contained in part (iii) of [27, Proposition 3.9], but we prefer to make it explicit here.

Lemma 4.2. *For a general $Q \in \langle A \rangle \subset \Pi_k$, all subsets $B \subset X$ of cardinality k such that $Q \in \langle B \rangle$ are contained in the span of Γ_k .*

Proof. Notice that, by the generality of A , Q is also a general point in the secant variety $\sigma_k(X)$. Since Q is general, the tangent space to $\sigma_k(X)$ at Q is also the span of the tangent spaces to X at the points of B , by the Terracini's Lemma. The claim now follows from [27, proposition 3.9 (iii)]. \square

Corollary 4.3. *The k -secant degree of a projective variety X is equal to the k -secant degree of the intersection of X with the span of its tangential k -contact locus.*

When the general contact locus of X is of type I, we can give a more explicit statement.

Proposition 4.4. *Assume that Γ_k is irreducible (i.e. type I). Assume the existence of a component W of $\Pi_k \cap X$, different from Γ_k , such that Π_k is tangent to X along W . Then for all a, b with $a > 0$ and $a + b = k$, the join of the a -th secant variety of W with the b -th secant variety of Γ_k cannot cover Π_k .*

Proof. Assume on the contrary that the join covers Π_k . Since $\sigma_k(X)$ has the expected dimension and it is equal to the union of the spaces Π_k 's, when the k -th points P_1, \dots, P_k vary on X , then for dimensional reasons there cannot be a subvariety $Y \subsetneq X$ which contains all the respective components W 's. Thus, for P_1, \dots, P_k

general, a general point Q of W is a general point of X . Hence the contact variety of Q, P_2, \dots, P_k has the same properties of Γ_k . In particular, Π_k is tangent to X along an irreducible variety of the same dimension of Γ_k , which contains P_2, \dots, P_k and Q . Since $Q \notin \Gamma_k$, this is excluded by the generality of P_1, \dots, P_k . \square

Corollary 4.5. *Assume that Γ_k is irreducible (i.e. type I). Then the secant degree of X is equal to the secant degree of the general contact locus Γ_k .*

The situation becomes completely different when the general Γ_k is of type II.

Remark 4.6. Assume that $\dim \sigma_{k+n}(X) = (k+n)(n+1) - 1 < r$, so that X is also weakly $(k+1)$ -defective. Assume that a general tangential contact locus Γ_{k+1} is formed by $k+1$ linearly independent linear spaces. Then also the general tangential contact locus Γ_k is formed by k linearly independent linear spaces.

Indeed, with no loss of generality, as in the proof of Lemma 3.5, we may assume $\Gamma_k \subset \Gamma_{k+1}$. Thus Γ_k is of type II, and all its components are contained in one component of Γ_{k+1} . But clearly if one takes irreducible varieties $\Gamma^i, i=1, \dots, k$, of the same dimension t_k , such that each Γ^i spans a linear space Π^i of dimension $> t_k$, and the linear spaces Π^i 's are linearly independent, then the span of the union of the Γ^i 's cannot have dimension $kd+k-1$.

We point out the following example.

Example 4.7. ([18, Proposition 1.7]) Let X be the Grassmannian $G := \mathbb{G}(\mathbb{P}^2, \mathbb{P}^7)$ and let $k=3$. Then X is weakly 3-defective and, for a general choice of $P_1, P_2, P_3 \in X$, Γ_3 is the union of three disjoint linear spaces L_1, L_2, L_3 of dimension 3, such that L_i contains P_i . The span Π_3 of Γ_3 is also tangent to X along a linear space $W = \mathbb{P}^5$, which misses the P_i 's and intersects each L_i along a line.

One computes soon that the join of L_i, L_j, W cannot fill Π_3 , for obvious dimensional reasons. It follows that the secant degree of X equals the secant degree of Γ_3 , which is 1. Hence the generic 3-identifiability of G holds.

We can derive, as in the previous example a general statement for the case where the tangential contact variety Γ_k is of type II.

Proposition 4.8. *Assume that the general k -tangential contact locus Γ_k is of type II, i.e. it is the union of components $\Gamma^1, \dots, \Gamma^k$. Assume that Π_k is tangent to X along $\Gamma_k \cup \Gamma$, where $\Gamma \not\subset \Gamma_k$. If $\langle \Gamma^1 \cup \dots \cup \widehat{\Gamma^I} \cup \dots \cup \Gamma^k \cup \Gamma \rangle \neq \Pi_k$, where with $\widehat{\Gamma^I}$ we indicate any non empty subset of $\{\Gamma^1, \dots, \Gamma^k\}$, then the k -th secant degree of X is equal to the k -th secant degree of Γ_k .*

Corollary 4.9. *In the assumptions of the previous proposition, assume the Γ^i 's are linear subspaces of \mathbb{P}^r , and linearly independent. Then X is j -identifiable for any $j \leq k$.*

Proof. Clearly, if X is k -identifiable then it is also j -identifiable for every $j \leq k$, hence we need simply to show that X is k -identifiable. This last follows since the union of linearly independent linear spaces is identifiable. \square

On the other hand, there are cases (where X is an *almost unbalanced* products of projective spaces) in which Γ_k has type II and the previous proposition does not apply, moreover the k -th secant degree of Γ_k is different from the k -th secant degree of X .

A first example is the following.

Example 4.10. Consider the Segre variety X of $\mathbb{P}^2 \times \mathbb{P}^2 \times \mathbb{P}^5$ and let $k=5$. One computes that the general tangential contact locus Γ_5 consist of 5 linearly independent linear subspaces $\Gamma^1 \cup \dots \cup \Gamma^5$, where each Γ_i is a \mathbb{P}^4 which passes through one of the points P_i .

The span Π_5 of Γ_5 is a \mathbb{P}^{24} which is tangent to X along an extra $\Gamma = \mathbb{P}^4$. Γ is linearly independent from any subset of 4 among the Γ^i 's. Thus the 5-th secant degree of Γ_5 is 1, but the secant degree of X equals the secant degree of $\Gamma_5 \cup \Gamma$, hence it is 6.

A similar situation holds for the almost unbalanced cases defined in Section 8 of [20].

5. Applications to concrete examples

We show below how Lemma 3.5 applies, thanks to the observations contained in Section 4, to conclude the identifiability of some relevant cases of Segre products, Veronese varieties, Segre-Veronese varieties and moment varieties.

Non-trivial Segre products $X \subset \mathbb{P}^r$, X image of $\mathbb{P}^{a_1} \times \dots \times \mathbb{P}^{a_s}$, contain, for each $j=1, \dots, s$, linear spaces L_1, \dots, L_k , of dimension a_j passing through k general points, which are linearly independent, provided that $k(a_j+1)$ is smaller than $(r+1)$. We do not know examples where these spaces are contact loci of weakly defective Segre products. In any case, the previous lemma, together with the analysis of weakly defective varieties contained in [26] and [29], yield the following.

Theorem 5.1. *Let $X \subset \mathbb{P}^r$ be an integral and non-degenerate n -dimensional variety and let k be a positive integer. Assume that X is not uniruled by lines and that $\dim \sigma_{k+n-1}(X) = (k+n-1)(n+1) - 1 < r$ (in particular X is not $(k+n-$*

1)-defective). Then X is not weakly j -defective (hence it is j -identifiable) for every $j \leq k$.

Proof. If X is not weakly j -defective, then the claim follows by [26]. Assume that X is weakly j -defective. Hence X is also weakly $(j+i)$ -defective for $i=1, \dots, n-1$. Since X is not uniruled, then Γ_{j+i} cannot be a union of linear spaces of positive dimension, so by Lemma 3.5 we have:

$$1 \leq \dim \Gamma_j < \dots < \dim(\Gamma_{j+n-1}).$$

It follows that $\dim(\Gamma_{j+n-1}) \geq n$, a contradiction. \square

Theorem 5.1 is applicable to any embedding $X \subset \mathbb{P}^r$ with $\mathcal{O}_X(1) \cong \mathcal{L}^{\otimes 2} \otimes \mathcal{M}$ with \mathcal{L} an ample line bundle and \mathcal{M} a nef line bundle. In particular, it is applicable to any Veronese embedding of \mathbb{P}^n , since in this case $\dim \sigma_i(X)$ is known for all i by a famous theorem of Alexander and Hirschowitz ([10]). However, the result that we get from Theorem 5.1 for symmetric tensors is weaker than the results of [31], [36], where all the non-identifiable Veronese varieties are classified.

Let us turn to the study of Segre varieties. Fix integer $s \geq 2$, $n_i > 0$, $1 \leq i \leq s$. Set $n := n_1 + \dots + n_s$ and $r := -1 + \prod_{i=1}^s (n_i + 1)$ and let $X \subset \mathbb{P}^r$ be the (n -dimensional) Segre embedding of the multiprojective space $\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_s}$. Since X is uniruled by lines, we cannot apply Theorem 5.1 directly. On the other hand, the structure of linear spaces contained in Segre embeddings of multiprojective spaces is well known.

Remark 5.2. Let $\pi_i: X \rightarrow \mathbb{P}^{n_i}$ denote the projection of the Segre variety X into its i -th factor. For any $i=1, \dots, s$ let $\mathcal{O}_X(\varepsilon_i)$ be the line bundle $\pi_i^*(\mathcal{O}_{\mathbb{P}^{n_i}}(1))$. Each $\mathcal{O}_X(\varepsilon_i)$ is a spanned line bundle and $\mathcal{O}_X(1) = \otimes_{i=1}^s \mathcal{O}_X(\varepsilon_i)$. Let $L \subset X$ be any linear space of positive dimension. Since $\mathcal{O}_L(1) \cong \oplus_{i=1}^s \mathcal{O}_X(\varepsilon_i)|_L$, there is $j \in \{1, \dots, s\}$ such that $\mathcal{O}_X(\varepsilon_j)|_L \cong \mathcal{O}_L$ for all $i \neq j$, i.e. L is contained in a fiber of π_j . Fix an integer $k \geq 1$ such that $\dim \sigma_{k+1}(X) = (k+1)(n+1) - 1 < r$ and assume that X is weakly $(k+1)$ -defective, with general tangential $(k+1)$ -contact locus Γ_{k+1} equal to a union of linear spaces, say L_1, \dots, L_k, L_{k+1} , of dimension $\varepsilon_k > 0$. Thus for each $i=1, \dots, k+1$ there is a unique integer $j(i) \in \{1, \dots, s\}$ such that $\pi_{j(i)}(L_i)$ is a point. We get a function $\phi_{k+1}: \{1, \dots, k+1\} \rightarrow \{1, \dots, s\}$ defined by the formula $\phi_{k+1}(i) := j(i)$. The function ϕ_{k+1} cannot depend on the choice of $k+1$ general points. Then, moving the points independently, by monodromy we get that this function ϕ_{k+1} must be invariant for the action of the full symmetric group on $\{1, \dots, k+1\}$. Thus $j(i) = j(1)$ for all i .

Using the previous remark, we can prove the following.

Theorem 5.3. *Let X be the Segre embedding of $\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_k}$ in \mathbb{P}^r as above. Assume that for each $i \in \{1, \dots, s\}$ there is $j \in \{1, \dots, s\} \setminus \{i\}$ such that $n_j = n_i$. Assume also that $\dim \sigma_{k+n-1}(X) = (k+n-1)(n+1) - 1 < r$. Then X is not weakly j -defective (hence it is j -identifiable) for every $j \leq k$.*

Proof. Assume that X is weakly j -defective. Thus it is also weakly $(j+i)$ -defective, for $i=0, \dots, n-1$.

Each general contact locus Γ_{j+i} has dimension $t_{j+i} \leq n-1$ and so there cannot be a strictly increasing sequence of integer t_j, \dots, t_{j+n-1} . By Lemma 3.5 there is an integer $h \in \{j, \dots, j+n-1\}$ such that Γ_h is a union $L_1 \cup \dots \cup L_h$ of h independent linear spaces of dimension t_h . Thus, by Remark 5.2, there is an integer $m \in \{1, \dots, s\}$ such that, for all $i=1, \dots, h$, $\pi_m(L_i)$ is a linear space of dimension $\dim L_i$, while $\pi_x(L_i)$ is a point for all $x \in \{1, \dots, s\} \setminus \{m\}$.

Notice that the general contact locus Γ_h is invariant for linear automorphisms of \mathbb{P}^r preserving X , in the sense that for any automorphism $\phi: \mathbb{P}^r \rightarrow \mathbb{P}^r$ with $\phi(X) = X$ the set $\phi(\Gamma_h)$ may be taken as a general contact locus. Thus $\pi_x(\phi(L_i))$ is a point for each $x \in \{1, \dots, s\} \setminus \{m\}$, while $\pi_m(\phi(L_i))$ is a linear space of dimension $\dim L_i$. By assumption there is $z \in \{1, \dots, s\} \setminus \{j\}$ with $n_z = n_j$. Hence there is an automorphism ψ of X exchanging the m -th factor and the z -th factor. This automorphism extends to a linear automorphism $\phi: \mathbb{P}^r \rightarrow \mathbb{P}^r$ with $\phi(X) = X$, because the embedding $X \hookrightarrow \mathbb{P}^r$ is induced by the complete linear system $|\mathcal{O}_X(1, \dots, 1)|$ and $\psi^*(\mathcal{O}_X(1, \dots, 1)) \cong \mathcal{O}_X(1, \dots, 1)$. We would have that $\pi_z(L_i)$ is a linear space of dimension $\dim L_i$, which yields a contradiction. \square

The condition that for each $i \in \{1, \dots, s\}$ there is $j \in \{1, \dots, s\} \setminus \{i\}$ such that $n_j = n_i$ restricts the range of application of Theorem 5.3. However, the theorem applies to the products of many copies of a given \mathbb{P}^q , i.e. to the case of cubic tensors (provided that one knows the non-defectivity of the corresponding Segre embedding).

In particular, in the case $n_i = 1$ for all i , as an immediate corollary of [23] and Theorem 5.3, we obtain the following result.

Corollary 5.4. *Fix an integer $s \geq 5$ and an integer $k > 0$ such that $(k+s-1)(s+1) \leq 2^s - 1$. Let $X \subset \mathbb{P}^{2^s-1}$ be the Segre embedding of $(\mathbb{P}^1)^s$. Then X is not weakly k -defective, hence it is k -identifiable.*

Thus, we obtain that binary tensors, i.e. tensors of type $2 \times \dots \times 2$ (s times), are k -identifiable for

$$k \leq \frac{2^s - 1}{s + 1} - s + 1.$$

This bound is very near to $k_0 = \lceil 2^s / (s + 1) \rceil$ is the maximal integer for which k -identifiability can hold.

As an immediate corollary of [9, Theorem 3.1] we get the following result.

Corollary 5.5. *Take $X = (\mathbb{P}^m)^s \subset \mathbb{P}^r$, $m > 1$, $r + 1 = (m + 1)^s$. Set $a := \lfloor (m + 1)^s / (ms + 1) \rfloor$ and let ε be the only integer such that $0 \leq \varepsilon \leq m$ and $a \equiv \varepsilon \pmod{m + 1}$. If either $\varepsilon > 0$ or $(k + ms)(sm + 1) \leq (m + 1)^s - 1$, set $a' := a - \varepsilon$. If $\varepsilon = 0$ and $(ms + 1)a = (m + 1)^s$ set $a' := a - 1$. For any positive integer $k + ms - 1 \leq a'$ X is not weakly k -defective and hence it is k -identifiable.*

Take X as in Corollary 5.5. Note that if $k \geq a + 2$, then $\sigma_{a+2}(X)$ has expected dimension $> r$ and hence identifiability fails for $\sigma_{a+2}(X)$. In the special case $s = 3$, in Corollary 3.10 it is sufficient to assume $3m(k + 3m - 1) < (m + 1)^3$, because the secant varieties of $(\mathbb{P}^m)^3$ are not defective, by [39].

Complete results on the defectivity of Segre products with more than 3 factors, except for the case of many copies of \mathbb{P}^1 and for $(\mathbb{P}^m)^3$ ([39]) and an almost complete for [9], are not available. As far as we know, the best procedure to find theoretically the non defectivity of Segre products is the inductive method described in [5].

For Segre-Veronese varieties, which correspond to rank 1 partially symmetric tensors, we can join Theorem 5.1 and Theorem 5.3 to get:

Theorem 5.6. *Fix integer $s \geq 2$, $n_i > 0$, $d_i > 0$, $1 \leq i \leq s$. Set $n := n_1 + \dots + n_s$ and $r = -1 + \prod \binom{d_i + n_i}{d_i}$. Let $X \subset \mathbb{P}^r$ be the (n -dimensional) Segre-Veronese embedding of the multiprojective space $\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_s}$, via the linear system $L_1^{d_1} \otimes \dots \otimes L_s^{d_s}$, where each L_i is the pull-back of $\mathcal{O}_{\mathbb{P}^{n_i}}(1)$ in the projection onto the i -th factor.*

Assume that $\dim \sigma_{k+n-1}(X) = (k + n - 1)(n + 1) - 1 < r$. Assume also that either:

- $d_i > 1$ for all i ; or
- for all i with $d_i = 1$ there exists $j \neq i$ such that $d_j = 1$ and $n_i = n_j$.

Then X is not weakly j -defective (hence it is j -identifiable) for every $j \leq k$.

Proof. Notice that X is not uniruled by lines, unless $d_i = 1$ for some i . When some d_i is 1, one can still apply the trick introduced in the proof of Theorem 5.3. \square

Example 5.7. C. Araujo, A. Massarenti and R. Rischter proved that if $h \leq a^{\lfloor \log_2(d-1) \rfloor}$, then $\sigma_{h+1}(X)$ has the expected dimension (see [13, Theorem 1.1] for a more precise result).

For the case $s = 2$ there is a very strong conjecture and several results supporting it ([1], [2], [4]). When $s = 2$ H. Abo and M. C. Brambilla give a list of 9 classes of secant varieties, which are known to be defective and they conjecture that they are the only defective ones for $s = 2$ ([4, Conjecture 5.5]. See [8, Table 2] for the list of known (in 2010) non-defective cases when $s = 2$.

H. Abo and M. C. Brambilla also gave many examples of defective secant varieties when $s=3, 4$ ([3]). See [4, §5] for a general conjecture for the non-defectivity of Segre-Veronese varieties. If $d_i \geq 2$ for all i , then X is not uniruled by lines, so we may apply Theorem 5.1 to obtain the non weak defectivity.

Example 5.8. C. Améndola, J.-C. Faugère, K. Ranestad and B. Sturmfels ([11], [12]) studied the Gaussian moment variety $\mathcal{G}_{n,d} \subset \mathbb{P}^N$, $N = \binom{n+d}{n} - 1$, whose points are the vectors of all moments of degree $\leq d$ of an n -dimensional Gaussian distribution. We have $\dim \mathcal{G}_{n,d} = n(n+3)/2$. They proved that all the secant varieties of $\mathcal{G}_{1,d}$ are non-defective, but $\sigma_2(\mathcal{G}_{n,d})$ is defective for all $n \geq 3$ ([12, Theorem 13 and Remark 15]). To apply Theorem 5.1 to $\mathcal{G}_{1,d}$ we need prove that $\mathcal{G}_{1,d}$ is not uniruled by lines. If so, Theorem 5.1 may be applied to $\sigma_k(\mathcal{G}_{1,d})$ for all k with $2(k+1) < d$. We may assume $d \geq 4$. The variety $\mathcal{G}_{1,d} \subset \mathbb{P}^d$ is described in [11] and [12]; it is an arithmetically Cohen-Macaulay surface of degree $\binom{d}{2}$ ([11, Proposition 1 and the last part of its proof and Corollary 2]) and its singular locus is a line $L \subset \mathcal{G}_{1,d}$ ([12, Lemma 4]). Let $\ell_L: \mathbb{P}^d \setminus L \rightarrow \mathbb{P}^{d-2}$ the note the linear projection of from L and let $\pi: \mathcal{G}_{1,d} \setminus L \rightarrow \mathbb{P}^{d-2}$ be the restriction of ℓ_L . Assume that $\mathcal{G}_{1,d}$ is ruled by line and call Δ the one-dimensional family of lines giving the ruling. Since the Grassmannian are complete, Δ covers $\mathcal{G}_{1,d}$ and hence the closure of the image of π would be an integral and non-degenerate curve $Y \subset \mathbb{P}^{d-2}$ such that a general $L \in \Delta$ is mapped to a general point of Y . The surface $\mathcal{G}_{1,d}$ is covered by a family of degree d rational normal curves of \mathbb{P}^d , each of them tangent to L at a different point. Hence Y must be the rational normal curve of degree $d-2$ and X would be contained in a degree $d-2$ 3-dimensional cone $T \subset \mathbb{P}^d$ with vertex L . Since $\deg(\mathcal{G}_{1,d}) = \binom{d}{2} > 3(d-2)$, Bezout theorem gives that every cubic hypersurface containing $\mathcal{G}_{1,d}$ contains T , contradicting the fact that $\mathcal{G}_{1,d}$ is cut out by cubics ([11, Proposition 1]).

For Segre products, the most general result that we can obtain with our techniques is resumed in the following theorem. It can be applied to a wide class of Segre products, not covered by Theorem 5.3.

Theorem 5.9. *Let X be the Segre embedding of a product $\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_q}$ in \mathbb{P}^r , with $n = \sum n_i = \dim(X)$. Let k be a positive integer. Assume that $\dim \sigma_{k+n-1}(X) = (k+n-1)(n+1) - 1 < r$ (in particular X is not $(k+n-1)$ -defective). Then X is j -identifiable for every $j \leq k$.*

Proof. Assume that X is weakly k -defective. Then, as in the proof of Theorem 5.1, there exists some j , $k \leq j \leq k+n-2$, such that the general tangential contact loci Γ_j and Γ_{j+1} have the same dimension. Thus, by Lemma 3.5 and its proof, the contact loci Γ_j and Γ_{j+1} are formed by a union of linearly independent linear spaces L_i , say of dimension s . By Remark 5.2, the L_i 's are contained in some component of the product, which, as in the proof of Theorem 5.6, is the same for all i 's. Say

for simplicity that the L_i 's sits in the first component. Call Y the Segre embedding of $\mathbb{P}^{n_2} \times \dots \times \mathbb{P}^{n_q}$ and call $Q_i \in Y$ the point to which L_i maps in the projection $X \rightarrow Y$. The Q_i 's are general points of Y . Since X is not $(j+1)$ -defective, the span M of the L_i 's has dimension $(j+1)(s+1) - 1$. If M meets X only in the union of the L_i 's, which are linearly independent subspaces, then by [19, Example 2.2] and Lemma 4.2 X is $(j+1)$ -identifiable. Otherwise, there is a point $P \in (M \cap X) \setminus \bigcup L_i$. The point P cannot belong to the same fiber over Y of some L_i , for otherwise M contains the span of L_i and P , and its dimension cannot be $(j+1)(s+1) - 1$. Thus the projection of $M \cap X$ to Y contains a point Q different from the Q_i 's. It follows that for a general choice of j points Q_1, \dots, Q_j in Y the span of the Q_i 's meets Y in a point Q different from the Q_i 's. By the trisecant lemma ([14, p. 109], [24, Example 1.8], [25, Proposition 2.6]), this means that the codimension of Y in its span is at most $j - 1$. Since the codimension of Y is $\prod(n_i + 1) - \sum n_i - 1$ (where the product and the sum range from 2 to q), and moreover $(j+1)(n+1) < \prod_{i=1}^q (a_i + 1)$, after a short computation we get that:

$$a_1 > \prod_{i=2}^q (a_i + 1) - \sum_{i=2}^q a_i,$$

i.e. Y is *unbalanced* in the sense of [20, Section 8]. Since we also have $j \geq a_1$, it follows by [5, Theorem 4.4] that either $\sigma_{j+1}(X)$ covers \mathbb{P}^r , or X is $(j+1)$ -defective. In both cases we have a contradiction. \square

We should remark that the inductive method of [5] is extended to the study of weak defectivity in [20]. The results contained in [20] are, in several cases, improved by the results that one can get via Theorem 5.9.

From the computational point of view, let us point out the following:

Remark 5.10. For specific examples of Segre products X , the *stacked hessian* method introduced in [30] can provide a positive answer to the k -identifiability of X . The stacked hessian method requires the computations of the derivatives of the span of k tangent spaces at general points, when the points move.

In order to prove the non $(k+n)$ -defectivity of X , by the Terracini's lemma, one must compute the dimension of the span of $k+n$ general tangent spaces (see e.g. [32, Section 6]). When n is not too big with respect to k , this second method, together with Theorem 5.9, can provide the k -identifiability of X with a faster procedure.

6. Positive characteristic

In this section, we work over an algebraically closed field \mathbb{K} of arbitrary characteristic.

Definition 6.1. Let $X \subset \mathbb{P}^r$ be an integral and non-degenerate variety. Set $n := \dim(X)$. X is said to be *very strange* if for a general codimension n linear subspace $M \subset \mathbb{P}^r$ the set $X \cap M$ is not in linearly general position in M , i.e. there is a hyperplane of M containing at least $r - n + 2$ points of $M \cap X$ ([42]).

We recall that $X \subset \mathbb{P}^r$ is not very strange if a general 0-dimensional section of X is in uniform position ([14, Ch. 2], [42, Introduction]). Hence $X \subset \mathbb{P}^r$ is not very strange if either $\text{char}(\mathbb{K}) = 0$ or X is non-singular in codimension 1 and $r \geq n + 3$ ([42, Corollaries 1.6 and 2.2 and Theorem 0.1]). Thus none of the examples of the last section is very strange. X is not very strange if a general curve section of X is not strange ([42, Lemma 1.1]) and in particular X is not very strange if $\text{char}(\mathbb{K}) > \text{deg}(X)$.

Lemma 6.2. *Let $Y \subset \mathbb{P}^r$ be an integral and non-degenerate variety, which is not very strange. Set $n := \dim(Y)$. Fix a general $S \subset Y$ with $\sharp(S) \leq r - n - 1$. Then S is the set-theoretic base locus of the linear system on Y induced by $H^0(\mathbb{P}^r, \mathcal{I}_S(1))$.*

Proof. Let $N \subset \mathbb{P}^r$ be a general linear space of dimension $n - r - 1$. By Bertini’s theorem the scheme $N \cap Y$ is a finite set of $\text{deg}(Y)$ points. Since Y is not very strange, the set $N \cap Y$ is in linearly general position in N . Hence for any $E \subset N \cap Y$ with $\sharp(E) = \sharp(S) \leq n - r - 1$, the restriction of $U := H^0(\mathbb{P}^r, \mathcal{I}_E(1))$ to $N \cap Y$ has E as its set-theoretic base locus. Since N is a linear space the restriction of U to Y has base locus contained in $N \cap Y$. Thus E is the base locus of the restriction of U to T . Since Y is integral and non-degenerate, a general subset $A \subset Y$ with cardinality at most $r - n$ spans a general subspace of \mathbb{P}^r with dimension $\sharp(A) - 1$. Hence E is a general subset of Y with cardinality $\sharp(S)$. \square

Proof of Theorem 1.1. Fix a general $q \in \sigma_k(X)$. If $\dim \sigma_k(X) < kn + k - 1$, then a dimensional count gives that $\mathcal{S}(q, X)$ is infinite.

Hence to prove Theorem 1.1 it is sufficient to find a contradiction to the existence of $A, B \in \mathcal{S}(q, X)$ with $A \neq B$. Set $E := B \setminus A \cup B$ and $Z := A \cup E$. We have $h^1(\mathbb{P}^r, \mathcal{I}_Z(1)) > 0$, because $q \in \langle A \rangle \cap \langle B \rangle$ and $q \notin \langle A \cap B \rangle$ ([15, Lemma 1]), i.e. Z does not impose $\sharp(Z)$ independent conditions to W . We may also assume that A and B are general in their irreducible component of the constructible set $\mathcal{S}(q, X)$. Since q is general, we may assume that A (resp. B) is a general subset of X with cardinality s , but of course $A \cup B$ is far from being general (in general). We identify X with M , so that \mathcal{L} and \mathcal{R} are line bundles on X and $\mathcal{O}_X(1) \cong \mathcal{L} \otimes \mathcal{R}$. Since A is general and $\sharp(A) \leq \dim V$, the set of all $f \in V$ vanishing at all points of A is a linear subspace of V with dimension $\dim(V) - s$. Hence $h^1(\mathbb{P}^r, \mathcal{I}_A(1)) = 0$. Since A is general, by Lemma 6.2 there is a hypersurface $T \in |V|$ with $A \subset T$ and $E \cap T = \emptyset$. Let $f \in V \setminus \{0\}$ be an equation of T . The multiplication by f induces an isomorphism between $H^0(X, \mathcal{R})$ and

the linear subspace $W(-T):=H^0(X, \mathcal{I}_T \otimes \mathcal{L} \otimes \mathcal{R}) \cap W$ of all $g \in W$ vanishing on T . Moreover, for any finite subset $S \subset X \setminus T$ this isomorphism sends $H^0(X, \mathcal{I}_S \otimes \mathcal{R})$ onto $W(-T-S) = \{h \in W : h|_{T \cup S} \equiv 0\}$. Since $\#(E) \leq \#(B) \leq h^0(\mathcal{R})$ and B is general in X , we have $h^0(X, \mathcal{I}_E \otimes \mathcal{R}) = h^0(X, \mathcal{R}) - \#(E)$, i.e. E impose s independent conditions to $H^0(X, \mathcal{R})$. Since $T \cap E = \emptyset$, we get $\dim(W(-T-E)) = \dim(W(-T)) - \#(E)$. Hence $H^0(\mathbb{P}^r, \mathcal{I}_Z(1)) = \dim W - \#(Z)$, i.e. $h^1(\mathbb{P}^r, \mathcal{I}_Z(1)) = 0$, a contradiction. \square

We give an example of application of Theorem 1.1.

Example 6.3. Take $X = \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_s}$ and positive integers $d_i > 0$ and let $X \subset \mathbb{P}^r$, $r = -1 + \prod_{i=1}^s \binom{n_i + d_i}{n_i}$. Write $d_i = b_i + c_i$ with $b_i > 0$, $c_i \geq 0$ and $c_j > 0$ for some j . Set $n := n_1 + \dots + n_s$. If we fix any integer $k > 0$ such that

$$k \leq \min \left\{ -2 - n + \prod_{i=1}^s \binom{n_i + b_i}{n_i}, \prod_{i=1}^s \binom{n_i + c_i}{n_i} \right\},$$

then Theorem 1.1 applies and proves the identifiability of a general $q \in \mathbb{P}^r$ of X -rank k .

References

1. ABO, H., On non-defectivity of certain Segre-Veronese varieties, *J. Symbolic Comput.* **45** (2010), 1254–1269.
2. ABO, H. and BRAMBILLA, M. C., Secant varieties of Segre-Veronese varieties $\mathbb{P}^m \times \mathbb{P}^n$ embedded by $\mathcal{O}(1, 2)$, *Exp. Math.* **18** (2009), 369–384.
3. ABO, H. and BRAMBILLA, M. C., New examples of defective secant varieties of Segre-Veronese varieties, *Collect. Math.* **63** (2012), 287–297.
4. ABO, H. and BRAMBILLA, M. C., On the dimensions of secant varieties of Segre-Veronese varieties, *Ann. Mat. Pura Appl.* **192** (2013), 61–92.
5. ABO, H., OTTAVIANI, G. and PETERSON, C., Induction for secant varieties of Segre varieties, *Trans. Amer. Math. Soc.* **361** (2009), 767–792.
6. ABO, H., OTTAVIANI, G. and PETERSON, C., Non-defectivity of Grassmannians of planes, *J. Algebraic Geom.* **21** (2012), 1–20.
7. ABO, H. and VANNIEUWENHOVEN, N., Most secant varieties of tangential varieties to Veronese varieties are nondefective, *Trans. Amer. Math. Soc.* **370** (2018), 393–420.
8. ÅDLANDSVIK, B., Joins and higher secant varieties, *Math. Scand.* **62** (1987), 213–222.
9. ALADPOOSH, T. and HAGHIGHI, H., On the dimension of higher secant varieties of Segre varieties $\mathbb{P}^n \times \dots \times \mathbb{P}^n$, *J. Pure Appl. Algebra* **215** (2011), 1040–1052.
10. ALEXANDER, J. and HIRSCHOWITZ, A., Polynomial interpolation in several variables, *J. Algebraic Geom.* **4** (1995), 201–222.
11. AMÉNDOLA, C., FAUGÈRE, J.-C. and STURMFELS, B., Moment varieties of Gaussian mixtures, *J. Algebr. Stat.* **7** (2016), 14–28.

12. AMÉNDOLA, C., RANESTAD, K. and STURMFELS, B., Algebraic Identifiability of Gaussian Mixtures, *Int. Math. Res. Not. IMRN*, DOI: 10.1093/imrn/rnx090 (2016, to appear), *Preprint*, [arXiv:1612.01129](https://arxiv.org/abs/1612.01129).
13. ARAUJO, C., MASSARENTI, A. and RISCHTER, R., On nonsecant defectivity of Segre-Veronese varieties, *Preprint*, 2016. [arXiv:1611.01674](https://arxiv.org/abs/1611.01674).
14. ARBARELLO, E., CORNALBA, M., GRIFFITHS, P. and HARRIS, J., *Geometry of Algebraic Curves, Vol. I*, Springer, Berlin, 1985.
15. BALLICO, E. and BERNARDI, A., Decomposition of homogeneous polynomials with low rank, *Math. Z.* **271** (2012), 1141–1149.
16. BALLICO, E., BERNARDI, A., CATALISANO, M. V. and CHIANTINI, L., Grassman secants, identifiability, and linear systems of tensors, *Linear Algebra Appl.* **438** (2013), 121–135.
17. BAUR, K., DRAISMA, J. and de GRAAF, W., Secant dimensions of minimal orbits: computations and conjectures, *Exp. Math.* **16** (2007), 239–250.
18. BERNARDI, A. and VANZO, D., A new class of non-identifiable skew symmetric tensors, *Preprint*, 2016, [arXiv:1606.04158](https://arxiv.org/abs/1606.04158).
19. BOCCI, C. and CHIANTINI, L., On the identifiability of binary Segre products, *J. Algebraic Geom.* **22** (2013), 1–11.
20. BOCCI, C., CHIANTINI, L. and OTTAVIANI, G., Refined methods for the identifiability of tensors, *Ann. Mat. Pura Appl.* **193** (2014), 1691–1702.
21. BORALEVI, A., A note on secants of Grassmannians, *Rend. Istit. Mat. Univ. Trieste* **45** (2013), 67–72.
22. CATALISANO, M. V., GERAMITA, A. V. and GIMIGLIANO, A., Secant varieties of Grassmann varieties, *Proc. Amer. Math. Soc.* **183** (2004), 633–642.
23. CATALISANO, M. V., GERAMITA, A. V. and GIMIGLIANO, A., Secant varieties of $\mathbb{P}^1 \times \dots \times \mathbb{P}^1$ (n -times) are NOT defective for $n \geq 5$, *J. Algebraic Geom.* **20** (2011), 295–327.
24. CHIANTINI, L. and CILIBERTO, C., A few remarks on the lifting problem, *Astérisque* **218** (1993), 95–109.
25. CHIANTINI, L. and CILIBERTO, C., Weakly defective varieties, *Trans. Amer. Math. Soc.* **454** (2002), 151–178.
26. CHIANTINI, L. and CILIBERTO, C., On the concept of k -secant order of a variety, *J. Lond. Math. Soc.* **73** (2006), 436–454.
27. CHIANTINI, L. and CILIBERTO, C., On the dimension of secant varieties, *J. Eur. Math. Soc. (JEMS)* **73** (2006), 436–454.
28. CHIANTINI, L., MELLA, M. and OTTAVIANI, G., One example of general unidentifiable tensors, *J. Algebr. Stat.* **5** (2014), 64–71.
29. CHIANTINI, L. and OTTAVIANI, G., On generic identifiability of 3-tensors of small rank, *SIAM J. Matrix Anal. Appl.* **33** (2012), 1018–1037.
30. CHIANTINI, L., OTTAVIANI, G. and VANNIUWENHOVEN, N., An algorithm for generic and low-rank specific identifiability of complex tensors, *SIAM J. Matrix Anal. Appl.* **35** (2014), 1265–1287.
31. CHIANTINI, L., OTTAVIANI, G. and VANNIUWENHOVEN, N., On identifiability of symmetric tensors of subgeneric rank, *Trans. Amer. Math. Soc.* **369** (2017), 4021–4042.
32. CHIANTINI, L., OTTAVIANI, G. and VANNIUWENHOVEN, N., Effective criteria for specific identifiability of tensors and forms, *SIAM J. Matrix Anal. Appl.* **38** (2017), 656–681.

33. DOMANOV, I. and De LATHAUWER, L., On the uniqueness of the canonical polyadic decomposition of third-order tensors—Part I: Basic results and uniqueness of one factor matrix, *SIAM J. Matrix Anal. Appl.* **34** (2013), 855–875.
34. DOMANOV, I. and De LATHAUWER, L., On the uniqueness of the canonical polyadic decomposition of third-order tensors—Part II: Uniqueness of the overall decomposition, *SIAM J. Matrix Anal. Appl.* **34** (2013), 876–903.
35. DOMANOV, I. and De LATHAUWER, L., Generic uniqueness conditions for the canonical polyadic decomposition and INDSCAL, *SIAM J. Matrix Anal. Appl.* **36** (2015), 1567–1589.
36. GALUPPI, F. and MELLA, M., Identifiability of homogeneous polynomials and Cremona Transformations, *Preprint*, 2016, [arXiv:1606.06895](https://arxiv.org/abs/1606.06895).
37. KRUSKAL, J. B., Three-way arrays: rank and uniqueness of trilinear decompositions, with application to arithmetic complexity and statistics, *Linear Algebra Appl.* **18** (1977), 95–138.
38. LANDSBERG, J. M., *Tensors: Geometry and Applications*, Graduate Studies in Mathematics **128**, Amer. Math. Soc., Providence, 2012.
39. LICKTEIG, T., Typical tensorial rank, *Linear Algebra Appl.* **69** (1985), 95–120.
40. MASSARENTI, A. and RISCHTER, R., Non-secant defectivity via osculating projections, *Ann. Sc. Norm. Super.* DOI: 10.2422/2036-2145.201611_012 (2016, to appear), *Preprint*, [arXiv:1610.09332v1](https://arxiv.org/abs/1610.09332v1).
41. MELLA, M., Singularities of linear systems and the Waring problem, *Trans. Amer. Math. Soc.* **358** (2006), 5523–5538.
42. RATHMANN, J., The uniform position principle for curves in characteristic p , *Math. Ann.* **276** (1987), 565–579.
43. SIDIROPOULOS, N. D. and BRO, R., On the uniqueness of multilinear decomposition of N -way arrays, *J. Chemom.* **14** (2000), 229–239.

Edoardo Ballico
Dipartimento di Matematica
Univ. Trento
Trento
Italy
edoardo.ballico@unitn.it

Luca Chiantini
Dipartimento di Ingegneria dell'Informazione e Scienze Matematiche
Univ. Siena
Siena
Italy
luca.chiantini@unisi.it

Alessandra Bernardi
Dipartimento di Matematica
Univ. Trento
Trento
Italy
alessandra.bernardi@unitn.it

*Received July 10, 2017
in revised form December 1, 2017*