

On a remark by Ohsawa related to the Berndtsson-Lempert method for L^2 -holomorphic extension

Tai Terje Huu Nguyen and Xu Wang

Abstract. In [15, Remark 4.1], Ohsawa asked whether it is possible to prove Theorem 4.1 and Theorem 0.1 in [15] using the Berndtsson-Lempert method. We shall answer Ohsawa’s question affirmatively in this paper. Our approach also suggests to introduce the Legendre-Fenchel theory and weak psh-geodesics into the Berndtsson-Lempert method.

Contents

1	Introduction	173
2	The Berndtsson-Lempert method in a simple setting	175
3	A solution to Ohsawa’s question	176
4	Appendix: weak geodesics in the space of toric plurisubharmonic functions	179
	References	181

1. Introduction

In [15] Ohsawa gave a new proof of two theorems of Guan-Zhou (see [11], [12], [18] for further details and related results), Theorem 1.1 and 1.2 below, and asked in a remark (remark 4.1 in [15]) whether a proof using the Berndtsson-Lempert method in [9] can be had.

Theorem 1.1. (Theorem 4.1 in [15] and Corollary 1.8 in [13]) *Let Ω be a pseudoconvex domain in \mathbb{C}^n , $\Omega' := \{(z', z_n) \in \Omega : z_n = 0\}$, ϕ a plurisubharmonic function on Ω , $f \in \mathcal{O}(\Omega')$ a holomorphic function on Ω' , and let $\alpha > 0$. Then there exists*

$\tilde{f} \in \mathcal{O}(\Omega)$ a holomorphic function on Ω such that $\tilde{f}|_{\Omega'} = f$ and

$$\int_{\Omega} |\tilde{f}|^2 e^{-\phi - \alpha |z_n|^2} \leq \frac{\pi}{\alpha} \int_{\Omega'} |f|^2 e^{-\phi},$$

where integration is with respect to the Lebesgue measure.

Theorem 1.2. (Theorem 0.1 in [15]) *With the same assumptions and notation as in Theorem 1.1, there exists $\tilde{f} \in \mathcal{O}(\Omega)$ such that $\tilde{f}|_{\Omega'} = f$ and satisfying the estimate*

$$\int_{\Omega} \frac{|\tilde{f}|^2 e^{-\phi}}{(1 + |z_n|^2)^{1+\alpha}} \leq \frac{\pi}{\alpha} \int_{\Omega'} |f|^2 e^{-\phi}.$$

In this paper we answer Ohsawa’s question affirmatively by proving an extension theorem with estimate using the Berndtsson-Lempert method. This is Theorem 1.3 below and the main theorem of this paper. A key ingredient in our proof is the construction of a weight function θ using the Legendre-Fenchel transform and weak geodesics for plurisubharmonic functions (see the appendix in Section 4) to which we can apply the Berndtsson-Lempert technique.

Below we use the following terminology. Let $\sigma : U \rightarrow \mathbb{R}, x \mapsto \sigma(x)$ be a real-valued function where $U \subseteq \mathbb{R}^n$ is some subset and we write $x := (x_1, \dots, x_n)$. We will say that σ is increasing if σ is separately increasing in each argument. That is, more precisely, if for each $j \in \{1, 2, \dots, n\}$, the one-variable function $t \mapsto \sigma(x_1, \dots, x_{j-1}, t, x_{j+1}, \dots, x_n)$ is increasing for all $(x_1, \dots, x_{j-1}, t, x_{j+1}, \dots, x_n) \in U$.

Theorem 1.3. (Main Theorem) *Let Ω be a pseudoconvex domain in \mathbb{C}^n , $1 \leq k \leq n$ an integer,*

$$\Omega' := \{(z', z'') \in \Omega : z'' = 0\}, \quad z'' := (z_{n-k+1}, \dots, z_n),$$

ϕ a plurisubharmonic function on Ω , $f \in \mathcal{O}(\Omega')$ a holomorphic function on Ω' , and σ a convex increasing function on \mathbb{R}^k . Then there exists $\tilde{f} \in \mathcal{O}(\Omega)$ a holomorphic function on Ω such that $\tilde{f}|_{\Omega'} = f$ and

$$\int_{\Omega} |\tilde{f}|^2 e^{-\phi - \sigma(\ln |z''|)} \leq \int_{\Omega'} e^{-\sigma(\ln |z''|)} \int_{\Omega'} |f|^2 e^{-\phi},$$

where we write $\ln |z''| := (\ln |z_{n-k+1}|, \dots, \ln |z_n|)$, $\Omega'' := \{z'' \in \mathbb{C}^k : \ln |z''| \in \Omega''_{\mathbb{R}}\}$, $\Omega''_{\mathbb{R}}$ denotes the convex hull of $\{\ln |z''| : z \in \Omega\}$, and integration is with respect to the Lebesgue measure.

In the case that σ depends only on $|z''|^2$, Theorem 1.3 reduces to [18, Theorem 1.7]. It is also very likely that the main theorem in [12] implies Theorem 1.3. So our contribution is not the theorem itself, but rather a method of proof using the Berndtsson-Lempert method.

2. The Berndtsson-Lempert method in a simple setting

We start by recalling quickly the Berndtsson-Lempert method for L^2 -holomorphic extension ([9]) in the simple setting of domains in \mathbb{C}^n as it will apply to us. Specifically, we consider the following set-up. Let $\Omega \subseteq \mathbb{C}^n$ be a pseudoconvex domain and let ϕ be a plurisubharmonic function on Ω . For $z \in \Omega$, we write $z = (z', z_n)$. Let $\Omega' := \{z \in \Omega : z_n = 0\}$, suppose that $f \in \mathcal{O}(\Omega')$ is a given holomorphic function on Ω' , and suppose that $|z_n| < 1$ on Ω . We are interested in finding a holomorphic extension $\tilde{f} \in \mathcal{O}(\Omega)$ of f defined on all of Ω for which we have a good weighted L^2 estimate of the form ([16])

$$\int_{\Omega} |\tilde{f}|^2 e^{-\phi} \leq C \int_{\Omega'} |f|^2 e^{-\phi},$$

where integration is with respect to the Lebesgue measure, and C is some universal constant. The Berndtsson-Lempert method demonstrated in [9] is the following. Fix a positive constant $j > 1$ and consider the following plurisubharmonic function

$$(2.1) \quad \theta : \mathbb{C} \times \Omega \longrightarrow \mathbb{R}, \quad (\tau, z) \longmapsto \theta(\tau, z) := \theta^{\operatorname{Re} \tau}(z) := \phi + j \max\{\ln |z_n|^2 - \operatorname{Re} \tau, 0\}.$$

For each $t := \operatorname{Re} \tau \leq 0$, θ^t provides a weight function for a weighted L^2 inner product $\langle \cdot, \cdot \rangle_t$ and the corresponding induced weighted L^2 norm $\|\cdot\|_t$, given by

$$\langle u, v \rangle_t := \int_{\Omega} u \bar{v} e^{-\theta^t}, \quad \|u\|_t^2 := \langle u, u \rangle_t.$$

Since $|z_n| < 1$ on Ω , we have $\theta^0 = \phi$. Let f_t denote the holomorphic extension of f with minimal $\|\cdot\|_t$ -norm. Then we can write [9, section 3]

$$\|f_t\|_t^2 = \sup_{g \in C_0^\infty(\Omega')} \frac{|\xi_g(F)|^2}{\|\xi_g\|_t^2}, \quad \xi_g(F) := \int_{\Omega'} F \bar{g} e^{-\phi},$$

where F denotes *any* holomorphic extension of f , and where $\|\xi_g\|_t^2$ denotes the (squared) dual norm of the continuous linear functional ξ_g , given by

$$\|\xi_g\|_t^2 = \sup_{\|F\|_t=1} |\xi_g(F)|^2.$$

The crux of the argument is that by complex Brunn-Minkowski theory (see e.g. [2], [3], [6], [7]), under the assumption that θ is plurisubharmonic, we have that $t \mapsto \|\xi_g\|_t^2$ is log-convex. Hence $t \mapsto e^t \|\xi_g\|_t^2$ is log-convex. Suppose that this is also bounded as $t \rightarrow -\infty$. Then $t \mapsto t + \ln(\|\xi_g\|_t^2)$ is convex and bounded as $t \rightarrow -\infty$, and

hence increasing. Taking the exponential, we infer that $t \mapsto e^t \|\xi_g\|_t^2$ is increasing, which implies that $t \mapsto e^{-t} \|f_t\|_t^2$ is decreasing. Thus, we get the inequality

$$\|f_0\|_0^2 \leq \lim_{t \rightarrow -\infty} e^{-t} \|f_t\|_t^2.$$

Since f_t is the extension of f with *minimal* $\|\cdot\|_t$ -norm, the right-hand side limit is bounded by the limit of $e^{-t} \|F\|_t^2$ as $t \rightarrow -\infty$ with F any extension as above. Writing $f_0 := \tilde{f}$ and recalling that we are assuming $\theta^0 = \phi$, we therefore find

$$\int_{\Omega} |\tilde{f}|^2 e^{-\phi} \leq \liminf_{t \rightarrow -\infty} e^{-t} \|F\|_t^2 = \liminf_{t \rightarrow -\infty} e^{-t} \int_{\Omega} |F|^2 e^{-\theta^t}.$$

By [9, Lemma 3.2, Lemma 3.3], as $j \rightarrow \infty$, the right-hand side limit converges to $\pi \int_{\Omega'} |f|^2 e^{-\phi}$. Thus, we get an extension \tilde{f} with the following estimate:

$$\int_{\Omega} |\tilde{f}|^2 e^{-\phi} \leq \pi \int_{\Omega'} |f|^2 e^{-\phi}.$$

This estimate is known to be optimal (we have equality in the case $\Omega = \Omega' \times \{|z_n| < 1\}$ and ϕ does not depend on z_n) and was first proved in [10], [12].

3. A solution to Ohsawa’s question

In this section we apply the Berndtsson-Lempert method in Section 2 to a weight function θ that is different from the one in (2.1) to prove Theorem 3.1 below. Notice that taking $\sigma = \alpha e^{2x}$ or $\sigma = (1 + \alpha) \ln(1 + e^{2x})$ we get $L = \pi/\alpha$ in Theorem 3.1. Hence Theorem 1.1 and 1.2 follow. Our construction of θ is based on a Legendre-Fenchel transform approach to weak geodesics for plurisubharmonic functions (see the appendix in Section 4).

Theorem 3.1. *Let $\Omega \subseteq \mathbb{C}^n$ be pseudoconvex and write for $z \in \Omega$, $z = (z', z_n)$. Let ϕ be a plurisubharmonic function in Ω , $\Omega' := \{z \in \Omega : z_n = 0\}$, and $f \in \mathcal{O}(\Omega')$ a holomorphic function on Ω' . Put $x := \ln |z_n|$ and let $\sigma = \sigma(x)$ be a convex and increasing function in x with the property that $L := \int_{\mathbb{C}} e^{-\sigma(\ln |w|)} < \infty$. Then there exists $\tilde{f} \in \mathcal{O}(\Omega)$ a holomorphic function on Ω such that $\tilde{f}|_{\Omega'} = f$ and*

$$(3.1) \quad \int_{\Omega} |\tilde{f}|^2 e^{-\phi - \sigma(\ln |z_n|)} \leq L \int_{\Omega'} |f|^2 e^{-\phi}.$$

Proof. Fix $c \in \mathbb{R}$ and consider the weight function θ defined by

$$\theta(\tau, z) := \theta^{\operatorname{Re} \tau}(z) := \phi(z) + \psi^{\operatorname{Re} \tau}(\ln |z_n|), \quad t = \operatorname{Re} \tau > 0,$$

where

$$(3.2) \quad \psi^t(x) := \psi(t, x) := t\sigma \left(\frac{x-c}{t} + c \right).$$

One may directly verify that (see also the appendix in Section 4 for another approach) ψ is convex in (t, x) and increasing with respect to x . Since ϕ is assumed to be plurisubharmonic, we know that θ defined above is plurisubharmonic in (τ, z) . By the Berndtsson-Lempert method, using the same notation as in Section 2, it follows that

$$\rho_c(t) := \ln \|\xi_g\|_t^2 - 2c(t-1)$$

is convex as a function of t . Hence for all $t \in [1/2, 1)$,

$$\frac{\rho_c(1) - \rho_c(t)}{1-t} \geq \frac{\rho_c(1) - \rho_c(1/2)}{1-1/2}.$$

Observe that $\rho_c(1) = \ln \|\xi_g\|_1^2$ does not depend on c . Thus,

$$\ln \|\xi_g\|_1^2 \geq \rho_c(t) + 2(1-t) (\ln \|\xi_g\|_1^2 - \rho_c(1/2)),$$

for all $t \in [1/2, 1)$ and for all $c \in \mathbb{R}$. Now let

$$c := \frac{-1}{(1-t)^2},$$

and let $t \uparrow 1$. Lemma 3.2 below implies that

$$(3.3) \quad \ln \|\xi_g\|_1^2 \geq \limsup_{t \uparrow 1} \rho_c(t).$$

Hence, by (3.3), we have a holomorphic extension \tilde{f} with the following estimate

$$\begin{aligned} \int_{\Omega} |\tilde{f}|^2 e^{-\phi - \sigma(\ln |z_n|)} &\leq \liminf_{t \uparrow 1} e^{2c(t-1)} \int_{\Omega} |F|^2 e^{-\phi - \psi^t(\ln |z_n|)} \\ &= \liminf_{t \uparrow 1} e^{2/(1-t)} \int_{\Omega} |F|^2 e^{-\phi - t\sigma \left(\frac{\ln |z_n| + \frac{1}{1-t}}{t} \right)}, \end{aligned}$$

where F is any arbitrary holomorphic extension. From the change of variables

$$z_n = e^{\frac{1}{t-1}} w, \quad \Omega_t := \{(z', w) \in \mathbb{C}^n : (z', e^{\frac{1}{t-1}} w) \in \Omega\},$$

and the definition of L , we find that letting $t \uparrow 1$,

$$e^{2/(1-t)} \int_{\Omega} |F|^2 e^{-\phi - t\sigma \left(\frac{\ln |z_n| + \frac{1}{1-t}}{t} \right)} = \int_{\Omega_t} |F(z', e^{\frac{1}{t-1}} w)|^2 e^{-\phi \left(z', e^{\frac{1}{t-1}} w \right) - t\sigma \left(\frac{\ln |w|}{t} \right)}$$

converges to the right hand side of (3.1). Hence the theorem follows. \square

Lemma 3.2. *Let $c := \frac{-1}{(1-t)^2}$. Then $\lim_{t \uparrow 1} (1-t)\rho_c(1/2) = 0$.*

Proof. It suffices to show that $\limsup_{c \rightarrow -\infty} \rho_c(1/2) < \infty$. That is,

$$\limsup_{c \rightarrow \infty} \sup_{F \in \mathcal{O}(\Omega)} \frac{\left| \int_{\Omega'} F \bar{g} e^{-\phi} \right|^2}{e^{-c} \int_{\Omega} |F|^2 e^{-\phi - \frac{1}{2}\sigma(2 \ln |z_n| - c)}} < \infty.$$

This now follows from the change of variables $z_n = e^{c/2} w$ and the definition of L. \square

Remark. The change of variables argument in the proof of Theorem 3.1 also suggests to use another weight function, with

$$(3.4) \quad \psi^t(x) = \sigma(x-t), \quad t \in \mathbb{R}.$$

In fact, as we shall see next, the proof of Theorem 3.1 is simpler if we use this new weight function. Nevertheless, we still wish to include the proof given above since it is our first example of the use of weak geodesics in the Berndtsson-Lempert method.

3.1. Proof of the main theorem (Theorem 1.3)

Since $\Omega \subset \mathbb{C}^{n-k} \times \Omega''$, one may assume that $\sigma = \infty$ outside $\Omega''_{\mathbb{R}}$. Also we may assume that the right-hand side in the estimate in the theorem is finite, else there is nothing to prove. Replace ψ^t in (3.2) by

$$(3.5) \quad \psi^t(x) = \sigma(x-t), \quad t \in \mathbb{R}, \quad x-t := (x_1-t, \dots, x_k-t).$$

It is clear that $\psi^t(x)$ is convex in (t, x) and increasing in x . Hence, the corresponding weight function

$$\theta(\tau, z) = \phi(z) + \psi^{\operatorname{Re} \tau}(\ln |z_n|)$$

is plurisubharmonic in (τ, z) , and the Berndtsson-Lempert method applies. Thus, we know that $t \mapsto \ln \|\xi_g\|_t^2 + 2kt$ is convex. Moreover, the change of variables $z'' = e^t w$ gives,

$$(3.6) \quad \int_{\Omega} |F|^2 e^{-\phi - \sigma(\ln |z''| - t)} = e^{2kt} \int_{\{(z', e^t w) \in \Omega\}} |F(z', e^t w)|^2 e^{-\phi(z', e^t w) - \sigma(\ln |w|)},$$

where F , as before, is any arbitrary fixed extension of f . From (3.6) we infer that $\ln \|\xi_g\|_t^2 + 2kt$ is also bounded near $t = -\infty$. Hence, $t \mapsto \ln \|\xi_g\|_t^2 + 2kt$ is increasing and there exists a holomorphic extension \tilde{f} with

$$\int_{\Omega} |\tilde{f}|^2 e^{-\phi - \sigma(\ln |z''|)} \leq \liminf_{t \rightarrow -\infty} \int_{\{(z', e^t w) \in \Omega\}} |F(z', e^t w)|^2 e^{-\phi(z', e^t w) - \sigma(\ln |w|)}$$

$$\begin{aligned} &\leq \int_{\mathbb{C}^k} e^{-\sigma(\ln |w|)} \int_{\Omega'} |f|^2 e^{-\phi} \\ &= \int_{\Omega''} e^{-\sigma(\ln |w|)} \int_{\Omega'} |f|^2 e^{-\phi}, \end{aligned}$$

where in the last equality we have used that $\sigma = \infty$ outside $\Omega''_{\mathbb{R}}$. This completes the proof. \square

Remark. From (3.6), in the case that Ω is horizontally balanced (that is, $(z', z'') \in \Omega \Rightarrow (z', \tau z'') \in \Omega$ for all $\tau \in \mathbb{C}$ with $|\tau| < 1$), the method above is equivalent to the one associated to the weight function

$$\theta(\tau, z) := \phi(z', \tau z'') + \sigma(\ln |z''|).$$

This construction has already been used in [8, Section 2]. The advantage of (3.5) is that it also applies to general Ω .

4. Appendix: weak geodesics in the space of toric plurisubharmonic functions

By a toric plurisubharmonic function we mean a function of the following form

$$\psi(z) := \psi(\log |z_1|, \dots, \log |z_n|)$$

on \mathbb{C}^n , where ψ is a convex increasing function on \mathbb{R}^n .

Definition 4.1. We call a family, say $\{\psi^t\}_{0 < t < 1}$, of toric plurisubharmonic functions a weak geodesic⁽¹⁾ if

$$\Psi : (\tau, z) \mapsto \psi^{\operatorname{Re} \tau}(z), \quad \tau \in \mathbb{C}_{(0,1)} := \{\tau \in \mathbb{C} : 0 < \operatorname{Re} \tau < 1\}$$

is plurisubharmonic on $\mathbb{C}_{(0,1)} \times \mathbb{C}^n$ and $(i\partial\bar{\partial}\Psi)^{n+1} = 0$ on $\mathbb{C}_{(0,1)} \times (\mathbb{C} \setminus \{0\})^n$.

Remark. Noticing that Ψ is locally bounded on $\mathbb{C}_{(0,1)} \times (\mathbb{C} \setminus \{0\})^n$, we know that $(i\partial\bar{\partial}\Psi)^{n+1}$ is well defined. Moreover, by a local change of variables $z_j = e^{w_j}$, we know that $\{\psi^t\}_{0 < t < 1}$ is a weak geodesic if and only if

$$\psi(t, x) := \psi^t(x)$$

is convex on $(0, 1) \times \mathbb{R}^n$ and

$$MA(\psi) := \det(D_{(t,x)}^2 \psi) = 0$$

⁽¹⁾ For usual weak psh geodesics on compact Kähler manifolds, see [1, Section 2]. The weak geodesic is also called generalized geodesic in [5, Section 2.2], and maximal psh segment or psh geodesic segment in [17, page 5]. See [14] for the background.

on $(0, 1) \times \mathbb{R}^n$, where $D_{(t,x)}^2\psi$ denotes the real Hessian matrix of ψ with respect to (t, x) . Hence, by Corollary 2.5 in [4], $\{\psi^t\}_{0 < t < 1}$ is a weak geodesic if and only if the Legendre-Fenchel transform of $\psi(t, x)$ with respect to x for fixed t , say

$$(\psi^t)^*(\xi) := \sup_{x \in \mathbb{R}^n} x \cdot \xi - \psi^t(x),$$

is an affine function in t . This suggest the following definition.

Definition 4.2. We call a family, say $\{\psi^t\}_{0 < t < 1}$, of toric plurisubharmonic functions, a weak geodesic segment if

$$(\psi^t)^* = t(\sigma^1)^* + (1-t)(\sigma^0)^*$$

with σ^1, σ^0 convex increasing functions on \mathbb{R}^n .

Now let us consider the case $n=1$. Let $\sigma^1 := \sigma$ in Theorem 3.1. For σ^0 we take

$$\sigma^0 := \mathbf{1}_{(-\infty, c]}(x) = \begin{cases} 0, & x \leq c \\ \infty, & x > c. \end{cases}$$

Then we have

$$(\sigma^0)^*(\xi) = \sup_{x \in \mathbb{R}} x\xi - \sigma^0(x) = \sup_{x \leq c} x\xi = c\xi + \sup_{x \leq 0} x\xi = \begin{cases} c\xi & \xi \geq 0 \\ \infty & \xi < 0. \end{cases}$$

Since $\sigma^1 = \sigma$ is also increasing, we know that $(\sigma^1)^*(\xi) = \sigma^*(\xi) = \infty$ when $\xi < 0$. Hence ψ^t can be written as

$$\psi^t(x) = \sup_{\xi \in \mathbb{R}} x\xi - t\sigma^*(\xi) - (1-t)c\xi = t \left(\sup_{\xi \in \mathbb{R}} \frac{x - (1-t)c}{t} \xi - \sigma^*(\xi) \right),$$

and $\sigma^{**} = \sigma$ gives $\psi(t, x) = t\sigma\left(\frac{x-c}{t} + c\right)$. This is precisely (3.2). Note that (3.4) also gives a weak geodesic since

$$\sup_{x \in \mathbb{R}} x\xi - \sigma(x-t) = t\xi + \sigma^*(\xi)$$

is affine in t .

References

1. BERMAN, R. and BERNDTSSON, B., Convexity of the K-energy on the space of Kähler metrics and uniqueness of extremal metrics, *J. Am. Math. Soc.* **30** (2017), 1165–1196. [MR3671939](#)
2. BERNDTSSON, B., Subharmonicity properties of the Bergman kernel and some other functions associated to pseudoconvex domains, *Ann. Inst. Fourier (Grenoble)* **56** (2006), 1633–1662. [MR2282671](#)
3. BERNDTSSON, B., Curvature of vector bundles associated to holomorphic fibrations, *Ann. Math.* **169** (2009), 531–560. [MR2480611](#)
4. BERNDTSSON, B., aConvexity on the space of Kähler metrics, *Ann. Fac. Sci. Toulouse* **22** (2013), 713–746. [MR3137249](#)
5. BERNDTSSON, B., A Brunn-Minkowski type inequality for Fano manifolds and some uniqueness theorems in Kähler geometry, *Invent. Math.* **200** (2015), 149–200. [MR3323577](#)
6. BERNDTSSON, B., *aReal and complex Brunn-Minkowski Theory*, AMS Contemporary Mathematics **681**, 2017. [MR3603880](#)
7. BERNDTSSON, B., aComplex Brunn-Minkowski theory and positivity of vector bundles. [arXiv:1807.05844v1](#). [MR3966792](#)
8. BERNDTSSON, B., Bergman kernels for Paley-Wiener spaces and Nazarov’s proof of the Bourgain-Milman theorem. [arXiv:2008.00837](#). [MR2985302](#)
9. BERNDTSSON, B. and LEMPERT, L., aA Proof of the Ohsawa-Takegoshi theorem with sharp estimates, *J. Math. Soc. Japan* **68** (2016), 1461–1472. [MR3564439](#)
10. BLOCKI, Z., Suita conjecture and the Ohsawa-Takegoshi extension theorem, *Invent. Math.* **193** (2013), 149–158. [MR3069114](#)
11. GUAN, Q., A remark on the extension of L^2 holomorphic functions. [arXiv:1809.05876](#). [MR4083241](#)
12. GUAN, Q. and ZHOU, X., A solution of an L^2 extension problem with an optimal estimate and applications, *Ann. of Math. (2)* **181** (2015), 1139–1208. [MR3296822](#)
13. GUAN, Q. and ZHOU, X. Y., Strong openness of multiplier ideal sheaves and optimal L^2 extension, *Sci. China Math.* **60** (2017), 967–976. [MR3647126](#)
14. MABUCHI, T., Some symplectic geometry on compact Kähler manifolds, *Osaka J. Math.* **24** (1987), 227–252. [MR0909015](#)
15. OHSAWA, T., aOn the extension of L^2 holomorphic functions VIII - a remark on a theorem of Guan and Zhou, *Internat. J. Math.* **28**, no. 9, 1740005, 12 pp. (2017). [MR3690414](#)
16. OHSAWA, T. and TAKEGOSHI, K., On the extension of L^2 -holomorphic functions, *Math. Z.* **195** (1987), 197–204. [MR0892051](#)
17. RÉMI, R., Plurisubharmonic geodesics in spaces of non-Archimedean metrics of finite energy. [arXiv:2012.07972](#).
18. YAO, S., LI, Z. and ZHOU, X., On the optimal extension theorem and a question of Ohsawa, *Nagoya Mathematical Journal* **1**, <https://doi.org/10.1017/nmj.2020.34>.

Tai Terje Huu Nguyen
Department of Mathematical Sciences
Norwegian University of
Science and Technology
NO-7491
Trondheim
Norway
tai.t.h.nguyen@ntnu.no

Xu Wang
Department of Mathematical Sciences
Norwegian University of
Science and Technology
NO-7491
Trondheim
Norway
xu.wang@ntnu.no

*Received June 21, 2021
in revised form July 20, 2021*