

Stable functors and cohomology theory in abelian categories

Shoutao Guo and Li Liang

Abstract. In this paper, we first introduce stable functors with respect to a preenveloping/precovering subcategory and investigate some of their properties. Using that we then introduce and study a relative complete cohomology theory in abelian categories. Some properties of the cohomology including vanishing are given. As applications, we give some characterizations of objects of finite homological dimensions including the flat dimension, cotorsion dimension, Gorenstein injective/flat dimension and projectively coresolved Gorenstein flat dimension.

Introduction

Tate cohomology was initially defined for representations of finite groups. Avramov and Martsinkovsky [2] extended the definition so that it can work well for finitely generated modules of finite Gorenstein dimension over a noetherian ring. Sather-Wagstaff, Sharif and White [27] further investigated Tate cohomology for objects in abelian categories with enough projectives and injectives. As a broad generalization of Tate cohomology to the realm of infinite group algebras or even associative rings, complete cohomology was introduced by Vogel and Goichot [15], Mislin [22] and Benson and Carlson [4] independently, and was further treated by Avramov and Veliche [3] and Nucinkis [23]. The main purpose of this paper is to introduce and study a relative complete cohomology theory in abelian categories. Much of our motivation comes from the theory on stabilization of functors developed by Martsinkovsky and Russell recently in [18]–[20], which is very useful for studying complete homology theory.

Key words and phrases: stable functor, complete/Tate cohomology, special preenveloping/precovering subcategory.

2010 Mathematics Subject Classification: 18G25, 18G10, 18G15.

The definition of projective/injective stabilization of functors was first given by Auslander and Bridger in [1]. It is known that the Hom modulo projectives is actually the projective stabilization of the covariant Hom functor, which plays an important role in the field of representation theory. The applications of injective stabilization of covariant functors are displayed in the complete homology theory [20] and in the (co)torsion theory [19]. In this paper we introduce a relative version of stabilization of functors with respect to a preenveloping/precovering subcategory to extend the complete cohomology theory.

Let \mathbf{A} be an abelian category, and \mathbf{V} a special preenveloping subcategory of \mathbf{A} . For a contravariant additive functor F from \mathbf{A} to the category \mathbf{Ab} of abelian groups, the \mathbf{V} -stable functor $\overline{F}^{\mathbf{V}}$ of F is defined as the cokernel of the natural transformation $\rho: L_0^{\mathbf{V}}F \rightarrow F$; see 2.1. Here $L_0^{\mathbf{V}}F$ is the 0th left derived functor of F with respect to \mathbf{V} . Let M and N be objects in \mathbf{A} . As a special case where $F = \text{Hom}_{\mathbf{A}}(-, N)$, we denote $\overline{F}^{\mathbf{V}}(M)$ by $\overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N)$. Then there is an equality $\overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N) = \text{Hom}_{\mathbf{A}}(M, N) / \mathbf{V}\text{Hom}_{\mathbf{A}}(M, N)$; see Proposition 2.4. Here $\mathbf{V}\text{Hom}_{\mathbf{A}}(M, N)$ is the subgroup of $\text{Hom}_{\mathbf{A}}(M, N)$ consisting of those morphisms $f: M \rightarrow N$ that factor through an object in \mathbf{V} . In Section 2, we mainly discuss the exactness of \mathbf{V} -stable functors. The following result is from Corollary 2.12, Lemma 2.14 and Proposition 2.15.

Theorem A. *Let F be a contravariant additive functor from \mathbf{A} to \mathbf{Ab} , and \mathbf{V} a special preenveloping subcategory of \mathbf{A} . If F is half $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact, then so is $\overline{F}^{\mathbf{V}}$, and the following statements are equivalent.*

- (i) $\overline{F}^{\mathbf{V}}$ is right $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact.
- (ii) $\overline{F}^{\mathbf{V}} = 0$.
- (iii) F is right $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact.

Moreover, if F is left $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact, then the following statements are equivalent.

- (i) $\overline{F}^{\mathbf{V}}$ is left $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact.
- (ii) $\overline{F}^{\mathbf{V}}(\Theta_{\mathbf{V}}^i M) = 0$ for each object M in \mathbf{A} and all $i \geq 1$.
- (iii) $\overline{F}^{\mathbf{V}}(\Theta_{\mathbf{V}} M) = 0$ for each object M in \mathbf{A} .

In the above theorem, $\Theta_{\mathbf{V}}^i M$ denotes the i th \mathbf{V} -cosyzygy of M ; see 1.3. A contravariant functor F is said to be half $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact if for each $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact short exact sequence $0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0$ of objects in \mathbf{A} , the sequence $F(A'') \rightarrow F(A) \rightarrow F(A')$ is exact. If furthermore the sequence $0 \rightarrow F(A'') \rightarrow F(A) \rightarrow F(A')$ is exact, then we call F left $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact. Right $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact functors are defined dually; see 2.10. We mention that the dual result of Theorem A is given in Corollary 2.13, Lemma 2.14 and Proposition 2.17.

As an immediate consequence of Theorem A one gets that the functor $\overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(-, N)$ is left $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact if and only if $\overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(\Theta_{\mathbf{V}}^{\geq 1} M, N) = 0$ for each object M in \mathbf{A} if and only if $\overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(\Theta_{\mathbf{V}} M, N) = 0$ for each object M in \mathbf{A} , and if \mathbf{V} is closed under direct summands then $\overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(-, N)$ is right $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact if and only if $\overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N) = 0$ for each object M in \mathbf{A} if and only if $N \in \mathbf{V}$; see Corollary 2.16.

Using the tools developed in Section 2, we study a relative complete cohomology theory with respect to a special precovering/preenveloping subcategory in Sections 3 and 4.

Let \mathbf{V} be a special preenveloping subcategory of \mathbf{A} . For two objects M and N in \mathbf{A} and $n \in \mathbb{Z}$, the n th complete cohomology of M and N with respect to \mathbf{V} is defined as $\widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N) = \text{colim}_i \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(\Theta_{\mathbf{V}}^i M, \Theta_{\mathbf{V}}^{i+n} N)$; see Definition 3.4. The following result is from Theorem 3.10 and Propositions 3.12, 4.5 and 4.14.

Theorem B. *Let \mathbf{V} be a special preenveloping subcategory of \mathbf{A} , and let M and N be objects in \mathbf{A} with $M \xrightarrow{\simeq} I$ and $N \xrightarrow{\simeq} J$ proper \mathbf{V} -coresolutions. For each $n \in \mathbb{Z}$ there exist natural isomorphisms*

$$\widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N) \cong \text{colim}_i \text{Ext}_{\mathbf{A}}^1(\Theta_{\mathbf{V}}^{i+1} M, \Theta_{\mathbf{V}}^{i+n} N) \cong \text{H}^n(\widetilde{\text{Hom}}_{\mathbf{A}}(I, J)).$$

Moreover, if \mathbf{V} is closed under direct summands, and N has a Tate \mathbf{V} -coresolution $N \xrightarrow{\simeq} I \xrightarrow{\alpha} T$, then for each object M in \mathbf{A} and each $n \in \mathbb{Z}$, there is a natural isomorphism

$$\widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N) \cong \text{H}^n(\text{Hom}_{\mathbf{A}}(M, T)).$$

If furthermore $\text{Ext}_{\mathbf{A}}^{\geq 1}({}^{\perp} \mathbf{V}, \mathbf{V}) = 0$, then for each $n \in \mathbb{Z}$ there is a natural isomorphism

$$\widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N) \cong \text{colim}_i \text{S}_{\mathbf{V}}^{-i} \text{Ext}_{\mathbf{A}}^{n+i}(M, N).$$

In the above theorem, $\widetilde{\text{Hom}}_{\mathbf{A}}$ is the stable Hom functor (see 4.1), and $\text{S}_{\mathbf{V}}^{-i} \text{Ext}$ is the left satellite functor of contravariant Ext functor; see 1.5 for more details. We mention that the dual result of Theorem B for $\widetilde{\text{Ext}}_{\mathbf{W}}^n(M, N)$ is also true; see Theorem 3.10 and Propositions 3.12, 4.5 and 4.15.

The relative complete cohomology has expected properties including vanishing. The next result is from Theorem 4.9.

Theorem C. *Let \mathbf{V} be a special preenveloping subcategory of \mathbf{A} closed under direct summands. Then for each object N in \mathbf{A} the following statements are equivalent.*

- (i) $\mathbf{V}\text{-id}_{\mathbf{A}} N < \infty$.
- (ii) $\widetilde{\text{Ext}}_{\mathbf{V}}^n(N, -) = 0 = \widetilde{\text{Ext}}_{\mathbf{V}}^n(-, N)$ for all $n \in \mathbb{Z}$.

$$(iii) \widetilde{\text{Ext}}_{\mathcal{V}}^0(N, N) = 0.$$

The definition of \mathcal{V} -injective dimension is given in 1.7. The dual version of Theorem C can be found in Theorem 4.11.

In the final section, we give some applications of the above vanishing results, and characterize objects of finite homological dimensions including the flat dimension, cotorsion dimension, Gorenstein injective/flat dimension and projectively coresolved Gorenstein flat dimension.

1. Preliminaries

Throughout this paper, \mathcal{A} denotes an abelian category. We use the term “subcategory” to mean a “full and additive subcategory that is closed under isomorphisms”.

1.1 Special preenveloping/precovering subcategories. Given a subcategory \mathcal{X} of \mathcal{A} , we write

$$\begin{aligned} {}^{\perp}\mathcal{X} &= \{M \mid \text{Ext}_{\mathcal{A}}^1(M, X) = 0 \text{ for all } X \in \mathcal{X}\} \\ \mathcal{X}^{\perp} &= \{N \mid \text{Ext}_{\mathcal{A}}^1(X, N) = 0 \text{ for all } X \in \mathcal{X}\}. \end{aligned}$$

Here $\text{Ext}_{\mathcal{A}}^1(-, -)$ is the 1st Yoneda Ext group. A *special \mathcal{X} -preenvelope* of an object M in \mathcal{A} is an exact sequence $0 \rightarrow M \rightarrow X \rightarrow C \rightarrow 0$ with $X \in \mathcal{X}$ and $C \in {}^{\perp}\mathcal{X}$. Dually, a *special \mathcal{X} -precover* of M is an exact sequence $0 \rightarrow K \rightarrow X' \rightarrow M \rightarrow 0$ with $K \in \mathcal{X}^{\perp}$ and $X' \in \mathcal{X}$. Recall that a subcategory \mathcal{X} of \mathcal{A} is *special preenveloping* if each object in \mathcal{A} has a special \mathcal{X} -preenvelope. Dually a subcategory \mathcal{X} of \mathcal{A} is called *special precovering* if each object in \mathcal{A} has a special \mathcal{X} -precover.

Setup. Throughout this paper, the symbol \mathcal{V} denotes a special preenveloping subcategory of \mathcal{A} , and the symbol \mathcal{W} denotes a special precovering subcategory of \mathcal{A} .

1.2 Proper (co)resolutions. Let M be an object in \mathcal{A} . A proper \mathcal{V} -*coresolution* of M is a complex I of objects in \mathcal{V} such that $I^{-n} = 0 = H^n(I)$ for all $n > 0$ and $H^0(I) \cong M$, and the associated exact sequence $I^+ \cong 0 \rightarrow M \rightarrow I^0 \rightarrow I^1 \rightarrow \dots$ is $\text{Hom}_{\mathcal{A}}(-, \mathcal{V})$ -exact (that is, it remains exact after applying the functor $\text{Hom}_{\mathcal{A}}(-, V)$ to it for each $V \in \mathcal{V}$), which is always denoted $M \xrightarrow{\cong} I$. The *proper \mathcal{W} -resolutions* $P \xrightarrow{\cong} M$ of M are defined dually.

1.3 (Co)Syzygies. A proper \mathcal{V} -coresolution $M \xrightarrow{\cong} I$ of an object M in \mathcal{A} is called *special* if each $\text{Ker}(I^i \rightarrow I^{i+1})$ is in ${}^{\perp}\mathcal{V}$ for $i \geq 1$. Since \mathcal{V} is a special preenveloping subcategory, every object in \mathcal{A} has a special proper \mathcal{V} -coresolution. We

let $\Theta_V^i M$ denote the kernel $\text{Ker}(I^i \rightarrow I^{i+1})$ for some special proper V -coresolution $M \xrightarrow{\cong} I$; it is always called the i th V -cosyzygy of M . We always set $\Theta_V M = \Theta_V^1 M$.

Dually, a proper W -resolution $P \xrightarrow{\cong} M$ of an object M in \mathbf{A} is called *special* if each $\text{Coker}(P_{i+1} \rightarrow P_i)$ is in W^\perp for $i \geq 1$. Since W is a special precovering subcategory, every object in \mathbf{A} has a special proper W -resolution. We let $\Omega_i^W M$ denote the cokernel $\text{Coker}(P_{i+1} \rightarrow P_i)$ for some special proper W -resolution $P \xrightarrow{\cong} M$; it is always called the i th W -syzygy of M . We always set $\Omega^W M = \Omega_1^W M$.

1.4 Remark. We mention that $\Theta_V^i M$ is in ${}^\perp V$, and $\Omega_i^W M$ is in W^\perp for each $i \geq 1$, which are used frequently in the paper.

1.5 Left satellite functors. Let F be a contravariant additive functor from \mathbf{A} to the category \mathbf{Ab} of abelian groups. For an object M in \mathbf{A} , there is a special V -preenvelope $0 \rightarrow M \rightarrow I \xrightarrow{\pi} \Theta_V M \rightarrow 0$ of M with $I \in V$ and $\Theta_V M \in {}^\perp V$. Following Cartan and Eilenberg [6], the 1st *left satellite* of F with respect to V , denoted $S_V^{-1} F$, is defined as $S_V^{-1} F(M) = \text{Ker } F(\pi)$. Then $S_V^{-1} F$ is a contravariant additive functor from \mathbf{A} to \mathbf{Ab} , and it is independent of the choices of special V -preenvelopes. We set $S_V^{-n} F = S_V^{-1}(S_V^{-n+1} F)$ for each $n > 0$, and set $S_V^0 F = F$.

Let G be a covariant additive functor from \mathbf{A} to the category \mathbf{Ab} of abelian groups. For each object M in \mathbf{A} , there is a special W -precover $0 \rightarrow \Omega^W M \xrightarrow{\varepsilon} P \rightarrow M \rightarrow 0$ of M with $P \in W$ and $\Omega^W M \in W^\perp$. The 1st *left satellite* of G with respect to W , denoted $S_W^{-1} G$, is defined as $S_W^{-1} G(M) = \text{Ker } G(\varepsilon)$. Then $S_W^{-1} G$ is a covariant additive functor from \mathbf{A} to \mathbf{Ab} , and it is independent of the choices of special W -precovers. We set $S_W^{-n} G = S_W^{-1}(S_W^{-n+1} G)$ for each $n > 0$, and set $S_W^0 G = G$.

Let M and N be objects in \mathbf{A} . For the contravariant functor $F = \text{Ext}_A^i(-, N)$, the value of the left satellite functor $S_V^{-n} F$ at M , $S_V^{-n} F(M)$, is always denoted $S_V^{-n} \text{Ext}_A^i(M, N)$. For the covariant functor $G = \text{Ext}_A^i(M, -)$, the value of the left satellite functor $S_W^{-n} G$ at N , $S_W^{-n} G(N)$, is always denoted $S_W^{-n} \text{Ext}_A^i(M, N)$.

1.6 Remark. Since $S_V^{-n} F(I) = 0$ for each $I \in V$ and any $n > 0$, one has

$$S_V^{-n} F(M) = S_V^{-n+k} F(\Theta_V^k M)$$

for $n > k \geq 0$. Similarly, since $S_W^{-n} G(P) = 0$ for each $P \in W$ and any $n > 0$, one has $S_W^{-n} G(M) = S_W^{-n+k} G(\Omega_k^W M)$ for $n > k \geq 0$.

1.7 Dimensions and relative cohomology. The V -injective dimension of an object N in \mathbf{A} is the quantity

$$V\text{-id}_A N = \inf\{\sup\{n \geq 0 \mid I_n \neq 0\} \mid N \xrightarrow{\cong} I \text{ is a proper } V\text{-coresolution of } N\}.$$

Let N be an object in \mathbf{A} with $N \xrightarrow{\simeq} I$ a proper \mathbf{V} -coresolution. Then for each object M in \mathbf{A} and every $i \in \mathbb{Z}$, the i th *relative \mathbf{V} -cohomology* of N with coefficients in M is defined as

$$\mathrm{Ext}_{\mathbf{AV}}^i(M, N) = \mathrm{H}^i(\mathrm{Hom}_{\mathbf{A}}(M, I)).$$

Specially, if \mathbf{A} has enough injectives and \mathbf{V} is the subcategory of injectives, then $\mathbf{V}\text{-id}_{\mathbf{A}}N$ is the classical injective dimension, and $\mathrm{Ext}_{\mathbf{AV}}^i(M, N)$ is actually the group $\mathrm{Ext}_{\mathbf{A}}^i(M, N)$.

Dually, one has the definition of *\mathbf{W} -projective dimension*, $\mathbf{W}\text{-pd}_{\mathbf{A}}M$, of an object M in \mathbf{A} . Also, for objects M and N in \mathbf{A} with $P \xrightarrow{\simeq} M$ a proper \mathbf{W} -resolution, the i th *relative \mathbf{W} -cohomology* of M with coefficients in N is defined as

$$\mathrm{Ext}_{\mathbf{WA}}^i(M, N) = \mathrm{H}^i(\mathrm{Hom}_{\mathbf{A}}(P, N)).$$

1.8 Remark. The relation between $\mathrm{Ext}_{\mathbf{AV}}^i(M, N)$ and $\mathrm{Ext}_{\mathbf{WA}}^i(M, N)$ may be derived from balanced pairs given by Chen [9, Definition 1.1]. Specially, if (\mathbf{W}, \mathbf{V}) is a balanced pair in \mathbf{A} , then by [9, Lemma 2.1] there is a natural isomorphism $\mathrm{Ext}_{\mathbf{AV}}^i(M, N) \cong \mathrm{Ext}_{\mathbf{WA}}^i(M, N)$ for all objects M and N in \mathbf{A} , and each $i \geq 0$. In the following we will recall the definition of balanced pairs.

1.9 Balanced pairs. Recall that a pair (\mathbf{X}, \mathbf{Y}) of subcategories of \mathbf{A} is called a *balanced pair* if the following conditions hold:

- \mathbf{X} is precovering and \mathbf{Y} is preenveloping.
- For each object M in \mathbf{A} , there is a proper \mathbf{X} -resolution $X \rightarrow M$ such that the associated exact sequence X^+ is $\mathrm{Hom}_{\mathbf{A}}(-, \mathbf{Y})$ -exact.
- For each object N in \mathbf{A} , there is a proper \mathbf{Y} -coresolution $N \rightarrow Y$ such that the associated exact sequence Y^+ is $\mathrm{Hom}_{\mathbf{A}}(\mathbf{X}, -)$ -exact.

Balanced pairs arise naturally from cotorsion triplets. Recall from [9] that a triplet $(\mathbf{X}, \mathbf{Z}, \mathbf{Y})$ of subcategories of \mathbf{A} is a complete hereditary cotorsion triplet if both (\mathbf{X}, \mathbf{Z}) and (\mathbf{Z}, \mathbf{Y}) are complete hereditary cotorsion pairs, see [13] for the definition of complete hereditary cotorsion pairs. It follows from Estrada, Pérez and Zhu [14, Proposition 4.2] that if $(\mathbf{X}, \mathbf{Z}, \mathbf{Y})$ is a complete hereditary cotorsion triplet then (\mathbf{X}, \mathbf{Y}) is a balanced pair⁽¹⁾. As an immediate consequence one gets that if \mathbf{A} has enough projectives and injectives then $(\mathrm{Prj}, \mathrm{Inj})$ is a balanced pair, where Prj is the subcategory of projectives and Inj is the subcategory of injectives. For more examples of balanced pairs one refers to [9] and [14].

The next result can be found in [9, Lemma 2.4].

⁽¹⁾ This result was first proved by Chen in [9, Proposition 2.6] under the assumption that \mathbf{A} has enough projectives and injectives.

1.10 Lemma. *Let M be an object in \mathbf{A} . Then for each $n \geq 0$, the following statements are equivalent.*

- (i) $\mathbf{W}\text{-pd}_{\mathbf{A}} M \leq n$.
- (ii) $\text{Ext}_{\mathbf{W}\mathbf{A}}^i(M, -) = 0$ for all $i > n$.
- (iii) For each proper \mathbf{W} -resolution $P \xrightarrow{\simeq} M$, $\text{Coker}(P_{n+1} \rightarrow P_n)$ is in \mathbf{W} .

Dually, one has the following result.

1.11 Lemma. *Let N be an object in \mathbf{A} . Then for each $n \geq 0$, the following statements are equivalent.*

- (i) $\mathbf{V}\text{-id}_{\mathbf{A}} N \leq n$.
- (ii) $\text{Ext}_{\mathbf{A}\mathbf{V}}^i(-, N) = 0$ for all $i > n$.
- (iii) For each proper \mathbf{V} -coresolution $N \xrightarrow{\simeq} I$, $\text{Ker}(I^n \rightarrow I^{n+1})$ is in \mathbf{V} .

2. Stable functors of additive functors

In this section, we introduce relative stable functors of additive functors and investigate some properties including the exactness of these functors.

Setup. Throughout this section, we let \mathbf{F} (resp., \mathbf{G}) be a contravariant (resp., covariant) additive functor from \mathbf{A} to the category \mathbf{Ab} of abelian groups.

2.1 Stable functors. For each object M in \mathbf{A} , there is a proper \mathbf{V} -coresolution $M \xrightarrow{\simeq} I$. The i th left derived functor of \mathbf{F} with respect to \mathbf{V} , denoted $\mathbf{L}_i^{\mathbf{V}}\mathbf{F}$, is defined as $\mathbf{L}_i^{\mathbf{V}}\mathbf{F}(M) = \mathbf{H}_i(\mathbf{F}(I))$. It is known that $\mathbf{L}_i^{\mathbf{V}}\mathbf{F}(M)$ is independent of the choices of proper \mathbf{V} -coresolutions of M ; see Enochs and Jenda [13, Section 8.2].

According to the universal property of cokernels there is a natural transformation $\rho: \mathbf{L}_0^{\mathbf{V}}\mathbf{F} \rightarrow \mathbf{F}$. The cokernel of ρ , denoted by $\overline{\mathbf{F}}^{\mathbf{V}}$, is called the \mathbf{V} -stable functor of \mathbf{F} . For an object M in \mathbf{A} , $\overline{\mathbf{F}}^{\mathbf{V}}(M)$ is independent of the choices of proper \mathbf{V} -coresolutions of M . It is clear that $\overline{\mathbf{F}}^{\mathbf{V}}$ is a contravariant additive functor from \mathbf{A} to \mathbf{Ab} .

Dually, for each object N in \mathbf{A} , there is a proper \mathbf{W} -resolution $P \xrightarrow{\simeq} N$. The i th left derived functor of \mathbf{G} with respect to \mathbf{W} , denoted $\mathbf{L}_i^{\mathbf{W}}\mathbf{G}$, is defined as $\mathbf{L}_i^{\mathbf{W}}\mathbf{G}(M) = \mathbf{H}_i(\mathbf{G}(P))$. It is known that $\mathbf{L}_i^{\mathbf{W}}\mathbf{G}(N)$ is independent of the choices of proper \mathbf{W} -resolutions of N ; see [13, Section 8.2].

It follows from the universal property of cokernels that there is a natural transformation $\varrho: \mathbf{L}_0^{\mathbf{W}}\mathbf{G} \rightarrow \mathbf{G}$. The cokernel of ϱ , denoted by $\underline{\mathbf{G}}^{\mathbf{W}}$, is called the \mathbf{W} -stable functor of \mathbf{G} . For an object N in \mathbf{A} , $\underline{\mathbf{G}}^{\mathbf{W}}(N)$ is independent of the choices of proper \mathbf{W} -resolutions of N . Clearly, $\underline{\mathbf{G}}^{\mathbf{W}}$ is a covariant additive functor from \mathbf{A} to \mathbf{Ab} .

The following result is used frequently in the paper.

2.2 Theorem. *Let M and N be objects in \mathbf{A} . Then the following statements hold.*

(a) *Let $0 \rightarrow M \xrightarrow{d} I^0 \rightarrow \Theta_{\mathbf{V}} M \rightarrow 0$ be a special \mathbf{V} -preenvelope of M . Then there is a natural isomorphism $\overline{\mathbf{F}}^{\mathbf{V}}(M) \cong \text{Coker } \mathbf{F}(d)$.*

(b) *Let $0 \rightarrow \Omega^{\mathbf{W}} N \rightarrow P_0 \xrightarrow{f} N \rightarrow 0$ be a special \mathbf{W} -precover of N . Then there is a natural isomorphism $\underline{\mathbf{G}}^{\mathbf{W}}(N) \cong \text{Coker } \mathbf{G}(f)$.*

Proof. We only prove (a); the statement (b) is proved dually.

Fix a proper \mathbf{V} -coresolution $0 \rightarrow \Theta_{\mathbf{V}} M \rightarrow I^1 \rightarrow I^2 \rightarrow \dots$. Then one gets a proper \mathbf{V} -coresolution $0 \rightarrow M \xrightarrow{d} I^0 \rightarrow I^1 \rightarrow \dots$. This yields the following commutative diagram with exact rows and columns:

$$\begin{array}{ccccccc}
 \mathbf{F}(I^1) & \longrightarrow & \mathbf{F}(I^0) & \longrightarrow & \mathbf{L}_0^{\mathbf{V}} \mathbf{F}(M) & \longrightarrow & 0 \\
 \downarrow & & \mathbf{F}(d) \downarrow & & \rho_M \downarrow & & \\
 0 & \longrightarrow & \mathbf{F}(M) & \xlongequal{\quad} & \mathbf{F}(M) & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \\
 & & \text{Coker } \mathbf{F}(d) & \xrightarrow{\quad \eta \quad} & \overline{\mathbf{F}}^{\mathbf{V}}(M), & &
 \end{array}$$

where the dotted morphism η is induced by the universal property of cokernels. Then η is an isomorphism by the Snake Lemma. \square

2.3 Corollary. *If M is in \mathbf{V} and N is in \mathbf{W} , then $\overline{\mathbf{F}}^{\mathbf{V}}(M) = 0 = \underline{\mathbf{G}}^{\mathbf{W}}(N)$.*

Let M and N be objects in \mathbf{A} , and let $\mathbf{F} = \text{Hom}_{\mathbf{A}}(-, N)$ and $\mathbf{G} = \text{Hom}_{\mathbf{A}}(M, -)$. We always denote $\overline{\mathbf{F}}^{\mathbf{V}}(M)$ by $\overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N)$, and denote $\underline{\mathbf{G}}^{\mathbf{W}}(N)$ by $\underline{\text{Hom}}_{\mathbf{A}}^{\mathbf{W}}(M, N)$.

2.4 Proposition. *For objects M and N in \mathbf{A} , there are two equalities*

$$\begin{aligned}
 \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N) &= \text{Hom}_{\mathbf{A}}(M, N) / \mathbf{V}\text{Hom}_{\mathbf{A}}(M, N), \text{ and} \\
 \underline{\text{Hom}}_{\mathbf{A}}^{\mathbf{W}}(M, N) &= \text{Hom}_{\mathbf{A}}(M, N) / \mathbf{W}\text{Hom}_{\mathbf{A}}(M, N).
 \end{aligned}$$

Here $\mathbf{V}\text{Hom}_{\mathbf{A}}(M, N)$ (resp., $\mathbf{W}\text{Hom}_{\mathbf{A}}(M, N)$) is the subgroup of $\text{Hom}_{\mathbf{A}}(M, N)$ consisting of the morphisms that factor through an object in \mathbf{V} (resp., \mathbf{W}).

Proof. We only prove the first equality; the second one is proved dually.

For an object M in \mathbf{A} , there is a proper \mathbf{V} -coresolution $M \xrightarrow{\cong} I$. Applying the functor $\text{Hom}_{\mathbf{A}}(-, N)$ to the exact sequence $0 \rightarrow M \xrightarrow{d} I^0 \rightarrow I^1 \rightarrow \dots$, one gets the following commutative diagram:

$$\begin{array}{ccccccc}
 \text{Hom}_{\mathbf{A}}(I^1, N) & \longrightarrow & \text{Hom}_{\mathbf{A}}(I^0, N) & \xrightarrow{\pi} & \mathbf{L}_0^{\mathbf{V}} \text{Hom}_{\mathbf{A}}(M, N) & \longrightarrow & 0 \\
 & & d^* \downarrow & \swarrow \rho_M & & & \\
 & & \text{Hom}_{\mathbf{A}}(M, N) & & & &
 \end{array}$$

By the definition of \mathbf{V} -stable functor, one has

$$\overline{\mathrm{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N) = \mathrm{Hom}_{\mathbf{A}}(M, N) / \mathrm{Im} \rho_M = \mathrm{Hom}_{\mathbf{A}}(M, N) / \mathrm{Im} d^*.$$

So it is sufficient to show $\mathrm{Im} d^* = \mathbf{V}\mathrm{Hom}_{\mathbf{A}}(M, N)$. Clearly, $\mathrm{Im} d^* \subseteq \mathbf{V}\mathrm{Hom}_{\mathbf{A}}(M, N)$. Conversely, let f be in $\mathbf{V}\mathrm{Hom}_{\mathbf{A}}(M, N)$. Then there exists an object $C \in \mathbf{V}$ and morphisms $g: M \rightarrow C$ and $h: C \rightarrow N$ such that $f = hg$. Since the sequence $\mathrm{Hom}_{\mathbf{A}}(I^0, C) \rightarrow \mathrm{Hom}_{\mathbf{A}}(M, C) \rightarrow 0$ is exact, there exists a morphism $\lambda: I^0 \rightarrow C$ such that $g = \lambda d$. Then $f = hg = h\lambda d = d^*(h\lambda) \in \mathrm{Im} d^*$, as desired. \square

2.5 Remark. Assume that \mathbf{A} has enough injectives and \mathbf{V} is the subcategory of injectives in \mathbf{A} . Then by Proposition 2.4, $\overline{\mathrm{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N)$ is the stable Hom group based on injectives given by Nucinkis [23]. Dually, if \mathbf{A} has enough projectives and \mathbf{W} is the subcategory of projectives in \mathbf{A} , then by Proposition 2.4, $\underline{\mathrm{Hom}}_{\mathbf{A}}^{\mathbf{W}}(M, N)$ is the classical stable Hom group.

In the following we collect some results on exactness of \mathbf{V} -stable functors and \mathbf{W} -stable functors.

2.6. Let M be an object in \mathbf{A} with $0 \rightarrow M \rightarrow I \rightarrow \Theta_{\mathbf{V}}M \rightarrow 0$ and $0 \rightarrow M \rightarrow I' \rightarrow \Theta'_{\mathbf{V}}M \rightarrow 0$ special \mathbf{V} -preenvelopes. Then one gets the following commutative diagram with exact rows:

$$(2.6.1) \quad \begin{array}{ccccccc} 0 & \longrightarrow & M & \longrightarrow & I & \xrightarrow{g} & \Theta_{\mathbf{V}}M & \longrightarrow & 0 \\ & & \parallel & & \beta \downarrow & & \alpha \downarrow & & \\ 0 & \longrightarrow & M & \longrightarrow & I' & \xrightarrow{g'} & \Theta'_{\mathbf{V}}M & \longrightarrow & 0. \end{array}$$

Dually, let N be an object in \mathbf{A} with $0 \rightarrow \Omega^{\mathbf{W}}N \rightarrow P \rightarrow N \rightarrow 0$ and $0 \rightarrow \Omega'^{\mathbf{W}}N \rightarrow P' \rightarrow N \rightarrow 0$ special \mathbf{W} -precovers. Then one gets the next commutative diagram with exact rows:

$$(2.6.2) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \Omega^{\mathbf{W}}N & \longrightarrow & P & \longrightarrow & N & \longrightarrow & 0 \\ & & \varphi \downarrow & & \downarrow & & \parallel & & \\ 0 & \longrightarrow & \Omega'^{\mathbf{W}}N & \longrightarrow & P' & \longrightarrow & N & \longrightarrow & 0. \end{array}$$

2.7 Proposition. *Adopt the notation from 2.6. $\overline{\mathbf{F}}^{\mathbf{V}}(\alpha)$ and $\underline{\mathbf{G}}^{\mathbf{W}}(\varphi)$ are isomorphisms.*

Proof. We prove that $\overline{\mathbf{F}}^{\mathbf{V}}(\alpha)$ is an isomorphism; the next statement is proved dually.

We adopt the setup and notation in 2.6. One gets an exact sequence

$$0 \longrightarrow I \xrightarrow{\begin{pmatrix} \beta \\ g \end{pmatrix}} I' \oplus \Theta_{\mathbb{V}} M \xrightarrow{(g', -\alpha)} \Theta'_{\mathbb{V}} M \longrightarrow 0;$$

it is split as $I \in \mathbb{V}$ and $\Theta'_{\mathbb{V}} M \in {}^{\perp}\mathbb{V}$ (see Remark 1.4). Then there exists a morphism $(\tau, \lambda): I' \oplus \Theta_{\mathbb{V}} M \rightarrow I$ satisfying $\text{id}_I = (\tau, \lambda) \begin{pmatrix} \beta \\ g \end{pmatrix} = \tau\beta + \lambda g$. Hence one obtains the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & I & \xrightarrow{\begin{pmatrix} \beta \\ g \end{pmatrix}} & I' \oplus \Theta_{\mathbb{V}} M & \xrightarrow{(g', -\alpha)} & \Theta'_{\mathbb{V}} M \longrightarrow 0 \\ & & \parallel & & \downarrow \begin{pmatrix} \tau & \lambda \\ g' & -\alpha \end{pmatrix} & & \parallel \\ 0 & \longrightarrow & I & \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} & I \oplus \Theta'_{\mathbb{V}} M & \xrightarrow{(0, 1)} & \Theta'_{\mathbb{V}} M \longrightarrow 0. \end{array}$$

It follows from the Snake Lemma that $\begin{pmatrix} \tau & \lambda \\ g' & -\alpha \end{pmatrix}: I' \oplus \Theta_{\mathbb{V}} M \rightarrow I \oplus \Theta'_{\mathbb{V}} M$ is an isomorphism. Since $\bar{\mathbb{F}}^{\mathbb{V}}$ is an additive functor and $\bar{\mathbb{F}}^{\mathbb{V}}(I) = 0 = \bar{\mathbb{F}}^{\mathbb{V}}(I')$ by Corollary 2.3, it is obvious that $\bar{\mathbb{F}}^{\mathbb{V}}(\alpha)$ is an isomorphism. \square

2.8. Let $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ be an exact sequence of objects in \mathbf{A} which is $\text{Hom}_{\mathbf{A}}(-, \mathbb{V})$ -exact. Fix special \mathbb{V} -preenvelopes $0 \rightarrow M' \rightarrow I^0 \rightarrow \Theta_{\mathbb{V}} M' \rightarrow 0$ and $0 \rightarrow M'' \rightarrow H^0 \rightarrow \Theta_{\mathbb{V}} M'' \rightarrow 0$. One gets the following commutative diagram with exact rows and columns (see [13, Remark 8.2.2]):

$$(2.8.1) \quad \begin{array}{ccccccc} & & 0 & & 0 & & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & M' & \longrightarrow & M & \longrightarrow & M'' \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & I^0 & \longrightarrow & I^0 \oplus H^0 & \longrightarrow & H^0 \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \Theta_{\mathbb{V}} M' & \longrightarrow & S & \longrightarrow & \Theta_{\mathbb{V}} M'' \longrightarrow 0. \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 0 & & 0 & & 0 \end{array}$$

Since $\Theta_{\mathbb{V}} M'$ and $\Theta_{\mathbb{V}} M''$ are in ${}^{\perp}\mathbb{V}$ (see Remark 1.4), so is S . Thus the middle column is a special \mathbb{V} -preenvelope of M , and the third non-zero row is $\text{Hom}_{\mathbf{A}}(-, \mathbb{V})$ -exact; in this case we write S as $\Theta_{\mathbb{V}} M$. Inductively, one gets a $\text{Hom}_{\mathbf{A}}(-, \mathbb{V})$ -exact short exact sequence

$$0 \longrightarrow \Theta_{\mathbb{V}}^i M' \longrightarrow \Theta_{\mathbb{V}}^i M \longrightarrow \Theta_{\mathbb{V}}^i M'' \longrightarrow 0$$

for each $i \geq 1$.

Dually, let $0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$ be an exact sequence of objects in \mathbf{A} which is $\text{Hom}_{\mathbf{A}}(\mathbf{W}, -)$ -exact. Fix special \mathbf{W} -precovers $0 \rightarrow \Omega^{\mathbf{W}}N' \rightarrow P_0 \rightarrow N' \rightarrow 0$ and $0 \rightarrow \Omega^{\mathbf{W}}N'' \rightarrow Q_0 \rightarrow N'' \rightarrow 0$. One gets the following commutative diagram with exact rows and columns (see [13, Lemma 8.2.1]):

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \Omega^{\mathbf{W}}N' & \longrightarrow & T & \longrightarrow & \Omega^{\mathbf{W}}N'' \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & P_0 & \longrightarrow & P_0 \oplus Q_0 & \longrightarrow & Q_0 \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & N' & \longrightarrow & N & \longrightarrow & N'' \longrightarrow 0. \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array}$$

Since $\Omega^{\mathbf{W}}N'$ and $\Omega^{\mathbf{W}}N''$ are in \mathbf{W}^{\perp} (see Remark 1.4), so is T . Thus the middle column is a special \mathbf{W} -precover of N , and the first non-zero row is $\text{Hom}_{\mathbf{A}}(\mathbf{W}, -)$ -exact; in this case we write T as $\Omega^{\mathbf{W}}N$. Inductively, one gets a $\text{Hom}_{\mathbf{A}}(\mathbf{W}, -)$ -exact short exact sequence

$$0 \longrightarrow \Omega_i^{\mathbf{W}}N' \longrightarrow \Omega_i^{\mathbf{W}}N \longrightarrow \Omega_i^{\mathbf{W}}N'' \longrightarrow 0$$

for each $i \geq 1$.

2.9 Remark. In 2.8, one sees that $\Theta_{\mathbf{V}}^i M$ is actually based on the special proper \mathbf{V} -coresolution $M \xrightarrow{\simeq} I \oplus H$, where $M' \xrightarrow{\simeq} I$ and $M'' \xrightarrow{\simeq} H$ are special proper \mathbf{V} -coresolutions of M' and M'' , respectively. Similarly, $\Omega_i^{\mathbf{W}}N$ is actually based on the special proper \mathbf{W} -resolution $P \oplus Q \xrightarrow{\simeq} N$, where $P \xrightarrow{\simeq} N'$ and $Q \xrightarrow{\simeq} N''$ are special proper \mathbf{W} -resolutions of N' and N'' , respectively. Proposition 2.7 above asserts that $\overline{\mathbf{F}}^{\mathbf{V}}(\Theta_{\mathbf{V}}^i M)$ and $\underline{\mathbf{G}}^{\mathbf{W}}(\Omega_i^{\mathbf{W}}N)$ are independent of the choices of special proper \mathbf{V} -coresolutions and special proper \mathbf{W} -resolutions, respectively.

2.10. The contravariant functor \mathbf{F} is said to be *half* $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -*exact* if for each $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact short exact sequence $0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0$ of objects in \mathbf{A} , the sequence $\mathbf{F}(A'') \rightarrow \mathbf{F}(A) \rightarrow \mathbf{F}(A')$ is exact. If furthermore the sequence $0 \rightarrow \mathbf{F}(A'') \rightarrow \mathbf{F}(A) \rightarrow \mathbf{F}(A')$ is exact, then we call \mathbf{F} *left* $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -*exact*. *Right* $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -*exact* functors are defined dually.

The covariant functor \mathbf{G} is *half* $\text{Hom}_{\mathbf{A}}(\mathbf{W}, -)$ -*exact* if for each $\text{Hom}_{\mathbf{A}}(\mathbf{W}, -)$ -exact short exact sequence $0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0$ of objects in \mathbf{A} , the sequence $\mathbf{G}(A') \rightarrow \mathbf{G}(A) \rightarrow \mathbf{G}(A'')$ is exact. If furthermore the sequence $0 \rightarrow \mathbf{G}(A') \rightarrow \mathbf{G}(A) \rightarrow \mathbf{G}(A'')$ is exact, then we call \mathbf{G} *left* $\text{Hom}_{\mathbf{A}}(\mathbf{W}, -)$ -*exact*. *Right* $\text{Hom}_{\mathbf{A}}(\mathbf{W}, -)$ -*exact* functors are defined dually.

2.11 Theorem. *Let F be a left $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact functor and G a left $\text{Hom}_{\mathbf{A}}(\mathbf{W}, -)$ -exact functor. Then the following statements hold.*

(a) *For each $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact short exact sequence $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ in \mathbf{A} , there exists an exact sequence*

$$\dots \rightarrow \bar{F}^{\mathbf{V}}(\Theta_{\mathbf{V}}M) \rightarrow \bar{F}^{\mathbf{V}}(\Theta_{\mathbf{V}}M') \rightarrow \bar{F}^{\mathbf{V}}(M'') \rightarrow \bar{F}^{\mathbf{V}}(M) \rightarrow \bar{F}^{\mathbf{V}}(M').$$

(b) *For each $\text{Hom}_{\mathbf{A}}(\mathbf{W}, -)$ -exact short exact sequence $0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$ in \mathbf{A} , there exists an exact sequence*

$$\dots \rightarrow \underline{G}^{\mathbf{W}}(\Omega^{\mathbf{W}}N) \rightarrow \underline{G}^{\mathbf{W}}(\Omega^{\mathbf{W}}N'') \rightarrow \underline{G}^{\mathbf{W}}(N') \rightarrow \underline{G}^{\mathbf{W}}(N) \rightarrow \underline{G}^{\mathbf{W}}(N'').$$

Proof. We prove (a); the statement (b) is proved dually.

Applying the functor F to the diagram (2.8.1), one gets the following commutative diagram:

$$\begin{array}{ccccccc} & & 0 & & 0 & & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & F(\Theta_{\mathbf{V}}M'') & & F(\Theta_{\mathbf{V}}M) & & F(\Theta_{\mathbf{V}}M') \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & F(H^0) & \longrightarrow & F(I^0 \oplus H^0) & \longrightarrow & F(I^0) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & F(M'') & \longrightarrow & F(M) & \longrightarrow & F(M') \\ & & \downarrow & & \downarrow & & \downarrow \\ & & \bar{F}^{\mathbf{V}}(M'') & & \bar{F}^{\mathbf{V}}(M) & & \bar{F}^{\mathbf{V}}(M'). \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 0 & & 0 & & 0 \end{array}$$

Since F is left $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact, all rows and columns are exact by Theorem 2.2. Thus by the Snake Lemma one has an exact sequence

(2.11.1)

$$0 \rightarrow F(\Theta_{\mathbf{V}}M'') \rightarrow F(\Theta_{\mathbf{V}}M) \rightarrow F(\Theta_{\mathbf{V}}M') \rightarrow \bar{F}^{\mathbf{V}}(M'') \rightarrow \bar{F}^{\mathbf{V}}(M) \rightarrow \bar{F}^{\mathbf{V}}(M').$$

The sequence $0 \rightarrow \Theta_{\mathbf{V}}M' \rightarrow \Theta_{\mathbf{V}}M \rightarrow \Theta_{\mathbf{V}}M'' \rightarrow 0$ is a $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact short exact sequence as $\Theta_{\mathbf{V}}M'' \in {}^{\perp}\mathbf{V}$. Fix special \mathbf{V} -preenvelopes $0 \rightarrow \Theta_{\mathbf{V}}M' \rightarrow I^1 \rightarrow \Theta_{\mathbf{V}}^2M' \rightarrow 0$ and $0 \rightarrow \Theta_{\mathbf{V}}M'' \rightarrow H^1 \rightarrow \Theta_{\mathbf{V}}^2M'' \rightarrow 0$. There is a commutative diagram with exact

rows and columns:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \Theta_V M' & \longrightarrow & \Theta_V M & \longrightarrow & \Theta_V M'' \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & I^1 & \longrightarrow & I^1 \oplus H^1 & \longrightarrow & H^1 \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \Theta_V^2 M' & \longrightarrow & \Theta_V^2 M & \longrightarrow & \Theta_V^2 M'' \longrightarrow 0. \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array}$$

Since $\Theta_V^2 M'$ and $\Theta_V^2 M''$ are in ${}^\perp V$, so is $\Theta_V^2 M$. Thus the middle column is a special V -preenvelope of $\Theta_V M$. Applying the functor F to the above diagram and combining the exact sequence (2.11.1), one gets the following commutative diagram with exact columns:

$$\begin{array}{ccccccccccc}
 & & 0 & & 0 & & 0 & & & & \\
 & & \downarrow & & \downarrow & & \downarrow & & & & \\
 & & F(\Theta_V^2 M'') & \longrightarrow & F(\Theta_V^2 M) & \longrightarrow & F(\Theta_V^2 M') & & & & \\
 & & \downarrow & & \downarrow & & \downarrow & & & & \\
 0 & \longrightarrow & F(H^1) & \longrightarrow & F(I^1 \oplus H^1) & \longrightarrow & F(I^1) & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & F(\Theta_V M'') & \longrightarrow & F(\Theta_V M) & \longrightarrow & F(\Theta_V M') & \longrightarrow & \bar{F}^V(M'') & \longrightarrow & \bar{F}^V(M) & \longrightarrow & \bar{F}^V(M') \\
 & & \downarrow & & \downarrow & & \downarrow & & \parallel & & \parallel & & \parallel \\
 & & \bar{F}^V(\Theta_V M'') & \longrightarrow & \bar{F}^V(\Theta_V M) & \longrightarrow & \bar{F}^V(\Theta_V M') & \xrightarrow{\delta} & \bar{F}^V(M'') & \longrightarrow & \bar{F}^V(M) & \longrightarrow & \bar{F}^V(M'). \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0 & & 0 & & 0 & & 0
 \end{array}$$

Here the first three non-zero rows are exact, and $\delta: \bar{F}^V(\Theta_V M') \rightarrow \bar{F}^V(M'')$ is obtained by the universal property of cokernels. Taking out a part from the above commutative diagram, one gets the following commutative diagram with the first two rows and all columns exact:

$$\begin{array}{ccccccc}
 F(I^1 \oplus H^1) & \longrightarrow & F(I^1) & \longrightarrow & 0 \\
 \downarrow & & \downarrow & & \downarrow \\
 F(\Theta_V M) & \longrightarrow & F(\Theta_V M') & \longrightarrow & \bar{F}^V(M'') \\
 \downarrow & & \downarrow & & \parallel \\
 \bar{F}^V(\Theta_V M) & \longrightarrow & \bar{F}^V(\Theta_V M') & \xrightarrow{\delta} & \bar{F}^V(M''). \\
 \downarrow & & \downarrow & & \downarrow \\
 0 & & 0 & & 0
 \end{array}$$

It follows from [6, III. Lemma 3.2] that the sequence

$$\bar{F}^{\mathbf{V}}(\Theta_{\mathbf{V}}M) \longrightarrow \bar{F}^{\mathbf{V}}(\Theta_{\mathbf{V}}M') \xrightarrow{\delta} \bar{F}^{\mathbf{V}}(M'')$$

is exact. The exactness of the sequence $\bar{F}^{\mathbf{V}}(\Theta_{\mathbf{V}}M') \xrightarrow{\delta} \bar{F}^{\mathbf{V}}(M'') \rightarrow \bar{F}^{\mathbf{V}}(M)$ can be obtained similarly. Hence one gets that the sequence

$$\bar{F}^{\mathbf{V}}(\Theta_{\mathbf{V}}M'') \longrightarrow \bar{F}^{\mathbf{V}}(\Theta_{\mathbf{V}}M) \longrightarrow \bar{F}^{\mathbf{V}}(\Theta_{\mathbf{V}}M') \longrightarrow \bar{F}^{\mathbf{V}}(M'') \longrightarrow \bar{F}^{\mathbf{V}}(M) \longrightarrow \bar{F}^{\mathbf{V}}(M')$$

is exact. Continuing this process, one gets the desired exact sequence in the statement. \square

2.12 Corollary. *Let F be a left $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact functor. Then the following statements are equivalent.*

- (i) $\bar{F}^{\mathbf{V}}$ is left $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact.
- (ii) $\bar{F}^{\mathbf{V}}(\Theta_{\mathbf{V}}^i M) = 0$ for each object M in \mathbf{A} and all $i \geq 1$.
- (iii) $\bar{F}^{\mathbf{V}}(\Theta_{\mathbf{V}} M) = 0$ for each object M in \mathbf{A} .

Proof. The implication (ii) \Rightarrow (iii) is clear, and (iii) \Rightarrow (i) holds by Theorem 2.11.

(i) \Rightarrow (ii): For each $i \geq 1$, consider a special \mathbf{V} -preenvelope

$$0 \longrightarrow \Theta_{\mathbf{V}}^{i-1} M \longrightarrow I^{i-1} \longrightarrow \Theta_{\mathbf{V}}^i M \longrightarrow 0.$$

By (i) one gets an exact sequence $0 \rightarrow \bar{F}^{\mathbf{V}}(\Theta_{\mathbf{V}}^i M) \rightarrow \bar{F}^{\mathbf{V}}(I^{i-1})$. Since I^{i-1} is in \mathbf{V} , one gets $\bar{F}^{\mathbf{V}}(I^{i-1}) = 0$ by Corollary 2.3, and so $\bar{F}^{\mathbf{V}}(\Theta_{\mathbf{V}}^i M) = 0$. \square

Dually, we have the next result; one refers to Martsinkovsky and Zangurashvili [21, Theorem 4.10] for some more equivalent conditions in the case where \mathbf{W} is the subcategory of projectives and $\mathbf{G} = \text{Hom}_{\mathbf{A}}(M, -)$.

2.13 Corollary. *Let G be a left $\text{Hom}_{\mathbf{A}}(\mathbf{W}, -)$ -exact functor. Then the following statements are equivalent.*

- (i) $\underline{G}^{\mathbf{W}}$ is left $\text{Hom}_{\mathbf{A}}(\mathbf{W}, -)$ -exact.
- (ii) $\underline{G}^{\mathbf{W}}(\Omega_i^{\mathbf{W}} N) = 0$ for each object N in \mathbf{A} and all $i \geq 1$.
- (iii) $\underline{G}^{\mathbf{W}}(\Omega^{\mathbf{W}} N) = 0$ for each object N in \mathbf{A} .

An absolute analog of the following lemma can be found in [21, Proposition 4.3].

2.14 Lemma. *If F is a half $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact functor, then so is the functor $\bar{F}^{\mathbf{V}}$. If G is a half $\text{Hom}_{\mathbf{A}}(\mathbf{W}, -)$ -exact functor, then so is the functor $\underline{G}^{\mathbf{W}}$.*

Proof. We prove the first statement; the second one is proved dually.

Let $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ be a $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact short exact sequence in \mathbf{A} . Applying the functor F to the diagram (2.8.1), one gets the following commutative diagram with exact columns by Theorem 2.2:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & F(H^0) & \longrightarrow & F(I^0 \oplus H^0) & \longrightarrow & F(I^0) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & F(M'') & \longrightarrow & F(M) & \longrightarrow & F(M') \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & \bar{F}^{\mathbf{V}}(M'') & \longrightarrow & \bar{F}^{\mathbf{V}}(M) & \longrightarrow & \bar{F}^{\mathbf{V}}(M') \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array}$$

The first two rows are exact as F is a half $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact additive functor, then so is the third one by [6, III. Lemma 3.2]. Thus $\bar{F}^{\mathbf{V}}$ is half $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact. \square

2.15 Proposition. *Let F be a half $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact functor. Then the following statements are equivalent.*

- (i) $\bar{F}^{\mathbf{V}}$ is right $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact.
- (ii) $\bar{F}^{\mathbf{V}} = 0$.
- (iii) F is right $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact.

Proof. (i) \implies (ii): For an object M in \mathbf{A} , consider a special \mathbf{V} -preenvelope

$$0 \rightarrow M \rightarrow I^0 \rightarrow \Theta_{\mathbf{V}}M \rightarrow 0.$$

Then from (i) one has an exact sequence $\bar{F}^{\mathbf{V}}(I^0) \rightarrow \bar{F}^{\mathbf{V}}(M) \rightarrow 0$. Since $I^0 \in \mathbf{V}$, one gets $\bar{F}^{\mathbf{V}}(I^0) = 0$ by Corollary 2.3, and so $\bar{F}^{\mathbf{V}}(M) = 0$.

(ii) \implies (iii): By the definition of \mathbf{V} -stable functor, there exists an exact sequence of functors $L_0^{\mathbf{V}}F \xrightarrow{\rho} F \rightarrow \bar{F}^{\mathbf{V}} \rightarrow 0$. So ρ is an epimorphism by (ii). Since the functor $L_0^{\mathbf{V}}F$ is right $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact by [13, Theorem 8.2.5(2)], the statement (iii) holds.

(iii) \implies (i): Let $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ be a $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact short exact sequence in \mathbf{A} . By Lemma 2.14, the sequence $\bar{F}^{\mathbf{V}}(M'') \rightarrow \bar{F}^{\mathbf{V}}(M) \rightarrow \bar{F}^{\mathbf{V}}(M')$ is exact. Then by Theorem 2.2 one gets the following commutative diagram with exact

rows and columns:

$$\begin{array}{ccccccc}
 F(M'') & \longrightarrow & F(M) & \longrightarrow & F(M') & \longrightarrow & 0 \\
 \downarrow & & \downarrow & & \downarrow & & \\
 \overline{F}^{\mathbf{V}}(M'') & \longrightarrow & \overline{F}^{\mathbf{V}}(M) & \longrightarrow & \overline{F}^{\mathbf{V}}(M') & & \\
 \downarrow & & \downarrow & & \downarrow & & \\
 0 & & 0 & & 0 & &
 \end{array}$$

So the sequence $\overline{F}^{\mathbf{V}}(M'') \longrightarrow \overline{F}^{\mathbf{V}}(M) \longrightarrow \overline{F}^{\mathbf{V}}(M') \longrightarrow 0$ is exact. \square

Let N be an object in \mathbf{A} . We notice that if \mathbf{V} is closed under direct summands, then N is in \mathbf{V} if and only if the functor $F = \text{Hom}_{\mathbf{A}}(-, N)$ is right $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact. Thus the following result is immediate by Proposition 2.15.

2.16 Corollary. *Suppose that \mathbf{V} is closed under direct summands. Then the following statements are equivalent for an object N in \mathbf{A} .*

- (i) $\overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(-, N)$ is right $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact.
- (ii) $\overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N) = 0$ for each object M in \mathbf{A} .
- (iii) $N \in \mathbf{V}$.

Dually, we have the following two results.

2.17 Proposition. *Let \mathbf{G} be a half $\text{Hom}_{\mathbf{A}}(\mathbf{W}, -)$ -exact functor. Then the following statements are equivalent.*

- (i) $\underline{\mathbf{G}}^{\mathbf{W}}$ is right $\text{Hom}_{\mathbf{A}}(\mathbf{W}, -)$ -exact.
- (ii) $\underline{\mathbf{G}}^{\mathbf{W}} = 0$.
- (iii) \mathbf{G} is right $\text{Hom}_{\mathbf{A}}(\mathbf{W}, -)$ -exact.

2.18 Corollary. *Suppose that \mathbf{W} is closed under direct summands. Then the following statements are equivalent for an object M in \mathbf{A} .*

- (i) $\underline{\text{Hom}}_{\mathbf{A}}^{\mathbf{W}}(M, -)$ is right $\text{Hom}_{\mathbf{A}}(\mathbf{W}, -)$ -exact.
- (ii) $\underline{\text{Hom}}_{\mathbf{A}}^{\mathbf{W}}(M, N) = 0$ for each object N in \mathbf{A} .
- (iii) $M \in \mathbf{W}$.

3. A cohomology theory based on stable functors

In this section we focus on the stable functors $\overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(-, N)$ and $\underline{\text{Hom}}_{\mathbf{A}}^{\mathbf{W}}(M, -)$, and introduce a cohomology theory based on these functors. Note that there are equalities $\overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N) = \text{Hom}_{\mathbf{A}}(M, N) / \mathbf{V}\text{Hom}_{\mathbf{A}}(M, N)$ and $\underline{\text{Hom}}_{\mathbf{A}}^{\mathbf{W}}(M, N) = \text{Hom}_{\mathbf{A}}(M, N) / \mathbf{W}\text{Hom}_{\mathbf{A}}(M, N)$ for all objects M and N in \mathbf{A} by Proposition 2.4.

3.1 Lemma. *The following statements hold:*

(a) *Let N and N' be objects in \mathbf{A} . Then for each object $M \in \mathbf{A}$, there is a natural isomorphism $\overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N \oplus N') \cong \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N) \oplus \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N')$.*

(b) *Let M and M' be objects in \mathbf{A} . Then for each object $N \in \mathbf{A}$, there is a natural isomorphism $\underline{\text{Hom}}_{\mathbf{A}}^{\mathbf{W}}(M \oplus M', N) \cong \underline{\text{Hom}}_{\mathbf{A}}^{\mathbf{W}}(M, N) \oplus \underline{\text{Hom}}_{\mathbf{A}}^{\mathbf{W}}(M', N)$.*

Proof. We only prove (a); the statement (b) is proved dually.

Let $0 \rightarrow M \rightarrow I \rightarrow \Theta_{\mathbf{V}}M \rightarrow 0$ be a special \mathbf{V} -preenvelope of M . Then by Theorem 2.2, one obtains the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} \text{Hom}_{\mathbf{A}}(I, N \oplus N') & \longrightarrow & \text{Hom}_{\mathbf{A}}(M, N \oplus N') & \longrightarrow & \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N \oplus N') & \longrightarrow & 0 \\ \cong \downarrow & & \cong \downarrow & & \eta \downarrow & & \\ \text{Hom}_{\mathbf{A}}(I, N) \oplus \text{Hom}_{\mathbf{A}}(I, N') & \longrightarrow & \text{Hom}_{\mathbf{A}}(M, N) \oplus \text{Hom}_{\mathbf{A}}(M, N') & \longrightarrow & \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N) \oplus \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N') & \longrightarrow & 0, \end{array}$$

where η is obtained by the universal property of cokernels. Moreover, η is an isomorphism by the Five Lemma. \square

3.2. Let M and N be objects in \mathbf{A} . Adopt the notation from 2.6. There is an induced morphism $\alpha: \Theta_{\mathbf{V}}M \rightarrow \Theta'_{\mathbf{V}}M$, and one has an isomorphism

$$\begin{pmatrix} \tau & \lambda \\ g' & -\alpha \end{pmatrix}: I' \oplus \Theta_{\mathbf{V}}M \longrightarrow I \oplus \Theta'_{\mathbf{V}}M$$

by the proof of Proposition 2.7. Fix a proper \mathbf{V} -coresolution

$$0 \longrightarrow N \longrightarrow J^0 \longrightarrow J^1 \longrightarrow \dots$$

of N . Then one gets the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} \text{Hom}_{\mathbf{A}}(J^1, I' \oplus \Theta_{\mathbf{V}}M) & \longrightarrow & \text{Hom}_{\mathbf{A}}(J^0, I' \oplus \Theta_{\mathbf{V}}M) & \longrightarrow & L_0^{\mathbf{V}}\text{Hom}_{\mathbf{A}}(N, I' \oplus \Theta_{\mathbf{V}}M) & \longrightarrow & 0 \\ \downarrow \cong & & \downarrow \cong & & \phi \downarrow & & \\ \text{Hom}_{\mathbf{A}}(J^1, I \oplus \Theta'_{\mathbf{V}}M) & \longrightarrow & \text{Hom}_{\mathbf{A}}(J^0, I \oplus \Theta'_{\mathbf{V}}M) & \longrightarrow & L_0^{\mathbf{V}}\text{Hom}_{\mathbf{A}}(N, I \oplus \Theta'_{\mathbf{V}}M) & \longrightarrow & 0, \end{array}$$

where ϕ is obtained by the universal property of cokernels. Moreover, ϕ is an isomorphism by the Five Lemma. Furthermore, one obtains the following commutative diagram with exact rows by the definition of \mathbf{V} -stable functors:

$$\begin{array}{ccccccc} L_0^{\mathbf{V}}\text{Hom}_{\mathbf{A}}(N, I' \oplus \Theta_{\mathbf{V}}M) & \longrightarrow & \text{Hom}_{\mathbf{A}}(N, I' \oplus \Theta_{\mathbf{V}}M) & \longrightarrow & \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(N, I' \oplus \Theta_{\mathbf{V}}M) & \longrightarrow & 0 \\ \phi \downarrow \cong & & \downarrow \cong & & \psi \downarrow & & \\ L_0^{\mathbf{V}}\text{Hom}_{\mathbf{A}}(N, I \oplus \Theta'_{\mathbf{V}}M) & \longrightarrow & \text{Hom}_{\mathbf{A}}(N, I \oplus \Theta'_{\mathbf{V}}M) & \longrightarrow & \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(N, I \oplus \Theta'_{\mathbf{V}}M) & \longrightarrow & 0, \end{array}$$

where ψ is obtained by the universal property of cokernels. Again by the Five Lemma, ψ is an isomorphism. It follows from Corollary 2.16 and Lemma 3.1 that

the morphism from $\overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(N, \Theta_{\mathbf{V}}M)$ to $\overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(N, \Theta'_{\mathbf{V}}M)$ obtained by composition with ψ is an isomorphism.

Dually, one gets that the morphism from $\underline{\text{Hom}}_{\mathbf{A}}^{\mathbf{W}}(\Omega^{\mathbf{W}}N, M)$ to $\underline{\text{Hom}}_{\mathbf{A}}^{\mathbf{W}}(\Omega^{\mathbf{W}}N, M)$ is an isomorphism.

3.3 Lemma. *Let M and N be objects in \mathbf{A} . Then there is a natural morphism*

$$\Delta_1 : \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N) \longrightarrow \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(\Theta_{\mathbf{V}}M, \Theta_{\mathbf{V}}N).$$

Moreover, it is independent of the choices of \mathbf{V} -coszygies, that is, if $0 \rightarrow M \rightarrow I' \rightarrow \Theta'_{\mathbf{V}}M \rightarrow 0$ and $0 \rightarrow N \rightarrow J' \rightarrow \Theta'_{\mathbf{V}}N \rightarrow 0$ are another special \mathbf{V} -preenvelopes, then there exists a commutative diagram with columns isomorphisms

$$\begin{array}{ccc} \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N) & \xrightarrow{\Delta_1} & \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(\Theta_{\mathbf{V}}M, \Theta_{\mathbf{V}}N) \\ \parallel & & \downarrow \cong \\ \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N) & \xrightarrow{\Delta'_1} & \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(\Theta'_{\mathbf{V}}M, \Theta'_{\mathbf{V}}N). \end{array}$$

Proof. Fix a special \mathbf{V} -preenvelope $0 \rightarrow M \xrightarrow{\sigma} I^0 \rightarrow \Theta_{\mathbf{V}}M \rightarrow 0$ of M . Applying the functor $\text{Hom}_{\mathbf{A}}(-, N)$ to the above sequence and using Theorem 2.2, one gets the next diagram with the row exact and $\delta_1 = \overline{\delta}_1 \pi_1$:

(3.3.1)

$$\begin{array}{ccccc} \text{Hom}_{\mathbf{A}}(I^0, N) & \xrightarrow{\sigma^*} & \text{Hom}_{\mathbf{A}}(M, N) & \xrightarrow{\delta_1} & \text{Ext}_{\mathbf{A}}^1(\Theta_{\mathbf{V}}M, N) \rightarrow \text{Ext}_{\mathbf{A}}^1(I^0, N). \\ & & \searrow \pi_1 & & \swarrow \overline{\delta}_1 \\ & & & \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N) & \end{array}$$

For objects $\Theta_{\mathbf{V}}M$ and N , fix special \mathbf{V} -preenvelopes $0 \rightarrow \Theta_{\mathbf{V}}M \xrightarrow{\tau} I^1 \rightarrow \Theta_{\mathbf{V}}^2M \rightarrow 0$ and $0 \rightarrow N \rightarrow J^0 \xrightarrow{\varsigma} \Theta_{\mathbf{V}}N \rightarrow 0$. Consider the following commutative diagram with exact rows and columns:

$$(3.3.2) \quad \begin{array}{ccccc} \text{Hom}_{\mathbf{A}}(I^1, J^0) & \xrightarrow{\tau^*} & \text{Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}M, J^0) & & \\ \downarrow & & \downarrow \varsigma_* & & \\ \text{Hom}_{\mathbf{A}}(I^1, \Theta_{\mathbf{V}}N) & \longrightarrow & \text{Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}M, \Theta_{\mathbf{V}}N) & \xrightarrow{\pi_2} & \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(\Theta_{\mathbf{V}}M, \Theta_{\mathbf{V}}N). \\ & & \downarrow \partial_1 & \dashrightarrow \beta_1 & \\ & & \text{Ext}_{\mathbf{A}}^1(\Theta_{\mathbf{V}}M, N) & & \end{array}$$

Since both $\Theta_{\mathbf{V}}M$ and $\Theta_{\mathbf{V}}^2M$ are in ${}^{\perp}\mathbf{V}$ (see Remark 1.4), and J^0 is in \mathbf{V} , one has

$$\text{Ext}_{\mathbf{A}}^1(\Theta_{\mathbf{V}}M, J^0) = 0 = \text{Ext}_{\mathbf{A}}^1(\Theta_{\mathbf{V}}^2M, J^0),$$

and hence ∂_1 and τ^* are epimorphisms. One has $\pi_2\zeta_*\tau^*=0$, so $\pi_2\zeta_*=0$. Thus by the universal property of cokernels, one gets an epimorphism $\beta_1:\text{Ext}_A^1(\Theta_V M, N) \rightarrow \overline{\text{Hom}}_A^V(\Theta_V M, \Theta_V N)$ such that $\beta_1\partial_1=\pi_2$. Thus $\Delta_1=\beta_1\overline{\delta_1}$ is the desired morphism from $\overline{\text{Hom}}_A^V(M, N)$ to $\overline{\text{Hom}}_A^V(\Theta_V M, \Theta_V N)$, which is natural as the connecting morphisms δ_1 and ∂_1 are natural.

Finally the existence of the commutative diagram in the statement follows a standard argument, and the vertical arrow is an isomorphism; see Proposition 2.7 and 3.2. \square

Continuing the construction in Lemma 3.3, one gets a sequence

$$\overline{\text{Hom}}_A^V(M, \Theta_V^n N) \xrightarrow{\Delta_1} \overline{\text{Hom}}_A^V(\Theta_V M, \Theta_V^{1+n} N) \xrightarrow{\Delta_2} \overline{\text{Hom}}_A^V(\Theta_V^2 M, \Theta_V^{2+n} N) \longrightarrow \dots$$

So we have the next definition.

3.4 Definition. Let M and N be objects in A . For each $n \in \mathbb{Z}$, the n th *complete cohomology* of M and N with respect to V is defined as

$$\widetilde{\text{Ext}}_V^n(M, N) = \text{colim}_i \overline{\text{Hom}}_A^V(\Theta_V^i M, \Theta_V^{i+n} N).$$

3.5 Remark. It is easy to see that $\widetilde{\text{Ext}}_V^n(-, N)$ is a contravariant additive functor from A to Ab . Specially, if A has enough injectives and V is the subcategory of injectives, then $\widetilde{\text{Ext}}_V^n(M, N)$ is the cohomology group given by Nucinkis in [23].

The following result is dual to Lemma 3.3.

3.6 Lemma. *Let M and N be objects in A . Then there is a natural morphism*

$$\Lambda_1 : \underline{\text{Hom}}_A^W(M, N) \longrightarrow \underline{\text{Hom}}_A^W(\Omega^W M, \Omega^W N).$$

Moreover, it is independent of the choices of W -syzygies, that is, if $0 \rightarrow \Omega'^W M \rightarrow Q' \rightarrow M \rightarrow 0$ and $0 \rightarrow \Omega'^W N \rightarrow P' \rightarrow N \rightarrow 0$ are both special W -precovers, then there exists a commutative diagram with columns isomorphisms

$$\begin{array}{ccc} \underline{\text{Hom}}_A^W(M, N) & \xrightarrow{\Lambda_1} & \underline{\text{Hom}}_A^W(\Omega^W M, \Omega^W N) \\ \parallel & & \downarrow \cong \\ \underline{\text{Hom}}_A^W(M, N) & \xrightarrow{\Lambda'_1} & \underline{\text{Hom}}_A^W(\Omega'^W M, \Omega'^W N). \end{array}$$

Continuing the construction in Lemma 3.6, one gets a sequence

$$\underline{\text{Hom}}_A^W(\Omega_n^W M, N) \xrightarrow{\Lambda_1} \underline{\text{Hom}}_A^W(\Omega_{1+n}^W M, \Omega^W N) \xrightarrow{\Lambda_2} \underline{\text{Hom}}_A^W(\Omega_{2+n}^W M, \Omega_2^W N) \longrightarrow \dots$$

So we have the next definition.

3.7 Definition. Let M and N be objects in \mathbf{A} . For each $n \in \mathbb{Z}$, the n th complete cohomology of M and N with respect to \mathbf{W} is defined as

$$\widehat{\text{Ext}}_{\mathbf{W}}^n(M, N) = \text{colim}_i \underline{\text{Hom}}_{\mathbf{A}}^{\mathbf{W}}(\Omega_{i+n}^{\mathbf{W}} M, \Omega_i^{\mathbf{W}} N).$$

3.8 Remark. It is easy to see that $\widehat{\text{Ext}}_{\mathbf{W}}^n(M, -)$ is a covariant additive functor from \mathbf{A} to \mathbf{Ab} . Specially, if \mathbf{A} has enough projectives and \mathbf{W} is the subcategory of projectives, then $\widehat{\text{Ext}}_{\mathbf{W}}^n(M, N)$ is the cohomology group given in [4].

3.9 Construction. For objects M and N in \mathbf{A} , we construct the following commutative diagrams

(3.9.1)

$$\begin{array}{ccccccc} \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(M, N) & \xrightarrow{\Delta_1} & \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(\Theta_{\mathbf{V}} M, \Theta_{\mathbf{V}} N) & \xrightarrow{\Delta_2} & \overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(\Theta_{\mathbf{V}}^2 M, \Theta_{\mathbf{V}}^2 N) & \longrightarrow & \dots \\ \cong \downarrow & \nearrow \overline{\delta}_1 & \nearrow \beta_1 & \searrow \overline{\delta}_2 & \nearrow \beta_2 & \searrow \overline{\delta}_3 & \\ \text{Ext}_{\mathbf{A}}^1(\Theta_{\mathbf{V}} M, N) & \xrightarrow{\mu_1} & \text{Ext}_{\mathbf{A}}^1(\Theta_{\mathbf{V}} M, \Theta_{\mathbf{V}} N) & \xrightarrow{\mu_2} & \text{Ext}_{\mathbf{A}}^1(\Theta_{\mathbf{V}}^2 M, \Theta_{\mathbf{V}}^2 N) & \longrightarrow & \dots \\ \downarrow \iota_1 & \nearrow \mu_1 & \downarrow \iota_2 & \nearrow \mu_2 & \downarrow \iota_3 & & \\ \text{S}_{\mathbf{V}}^{-1} \text{Ext}_{\mathbf{A}}^1(M, N) & \xrightarrow{\Phi_1} & \text{S}_{\mathbf{V}}^{-1} \text{Ext}_{\mathbf{A}}^1(\Theta_{\mathbf{V}} M, \Theta_{\mathbf{V}} N) & \xrightarrow{\Phi_2} & \text{S}_{\mathbf{V}}^{-1} \text{Ext}_{\mathbf{A}}^1(\Theta_{\mathbf{V}}^2 M, \Theta_{\mathbf{V}}^2 N) & \longrightarrow & \dots \end{array}$$

(3.9.2)

$$\begin{array}{ccccccc} \underline{\text{Hom}}_{\mathbf{A}}^{\mathbf{W}}(M, N) & \longrightarrow & \underline{\text{Hom}}_{\mathbf{A}}^{\mathbf{W}}(\Omega^{\mathbf{W}} M, \Omega^{\mathbf{W}} N) & \longrightarrow & \underline{\text{Hom}}_{\mathbf{A}}^{\mathbf{W}}(\Omega_2^{\mathbf{W}} M, \Omega_2^{\mathbf{W}} N) & \longrightarrow & \dots \\ \cong \downarrow & \nearrow & \nearrow & \searrow & \nearrow & \searrow & \\ \text{Ext}_{\mathbf{A}}^1(M, \Omega^{\mathbf{W}} N) & \longrightarrow & \text{Ext}_{\mathbf{A}}^1(\Omega^{\mathbf{W}} M, \Omega_2^{\mathbf{W}} N) & \longrightarrow & \text{Ext}_{\mathbf{A}}^1(\Omega_2^{\mathbf{W}} M, \Omega_3^{\mathbf{W}} N) & \longrightarrow & \dots \\ \downarrow & \nearrow & \downarrow & \nearrow & \downarrow & \nearrow & \\ \text{S}_{\mathbf{W}}^{-1} \text{Ext}_{\mathbf{A}}^1(M, N) & \longrightarrow & \text{S}_{\mathbf{W}}^{-1} \text{Ext}_{\mathbf{A}}^1(\Omega^{\mathbf{W}} M, \Omega_2^{\mathbf{W}} N) & \longrightarrow & \text{S}_{\mathbf{W}}^{-1} \text{Ext}_{\mathbf{A}}^1(\Omega_2^{\mathbf{W}} M, \Omega_2^{\mathbf{W}} N) & \longrightarrow & \dots \end{array}$$

We construct the diagram (3.9.1); the diagram (3.9.2) is constructed similarly.

Adopt the setup and the notation from the proof of Lemma 3.3. One gets the monomorphism $\overline{\delta}_1$ and epimorphism β_1 with $\Delta_1 = \beta_1 \overline{\delta}_1$. Similarly, one gets the monomorphism $\overline{\delta}_2$ and epimorphism β_2 with $\Delta_2 = \beta_2 \overline{\delta}_2$.

Fix special \mathbf{V} -preenvelopes $0 \rightarrow \Theta_{\mathbf{V}} M \xrightarrow{\tau} I^1 \rightarrow \Theta_{\mathbf{V}}^2 M \rightarrow 0$ and $0 \rightarrow N \rightarrow J^0 \xrightarrow{\varsigma} \Theta_{\mathbf{V}} N \rightarrow 0$. Consider the next commutative diagram with exact rows and columns:

$$\begin{array}{ccc} \text{Hom}_{\mathbf{A}}(I^1, J^0) & \longrightarrow & \text{Hom}_{\mathbf{A}}(I^1, \Theta_{\mathbf{V}} N) \\ \tau^* \downarrow & & \downarrow \\ \text{Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}} M, J^0) & \xrightarrow{\varsigma^*} & \text{Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}} M, \Theta_{\mathbf{V}} N) \xrightarrow{\partial_1} \text{Ext}_{\mathbf{A}}^1(\Theta_{\mathbf{V}} M, N) \\ & & \delta_2 \downarrow \\ & & \text{Ext}_{\mathbf{A}}^1(\Theta_{\mathbf{V}}^2 M, \Theta_{\mathbf{V}} N) \\ & & \lambda \downarrow \\ & & \text{Ext}_{\mathbf{A}}^1(I^1, \Theta_{\mathbf{V}} N). \end{array}$$

Here both ∂_1 and τ^* are epimorphisms, as $\Theta_V M$ and $\Theta_V^2 M$ are in ${}^\perp V$ and $J^0 \in V$. It is easy to see $\delta_{2\zeta_*} = 0$, so one gets a morphism

$$\Psi_2 : \text{Ext}_A^1(\Theta_V M, N) \longrightarrow \text{Ext}_A^1(\Theta_V^2 M, \Theta_V N)$$

such that $\Psi_2 \partial_1 = \delta_2$. By the diagram (3.3.2), one has $\overline{\delta_2} \beta_1 \partial_1 = \overline{\delta_2} \pi_2 = \delta_2$. So one gets $\Psi_2 = \overline{\delta_2} \beta_1$, as ∂_1 is an epimorphism. Similar as above, one obtains $\Psi_3 = \overline{\delta_3} \beta_2$.

By the definition of left satellite functors, $S_V^{-1} \text{Ext}_A^1(M, N)$ is the kernel of the morphism from $\text{Ext}_A^1(\Theta_V M, N)$ to $\text{Ext}_A^1(I^0, N)$, so the morphism ι_1 in (3.9.1) is the natural embedding, and so are ι_2 and ι_3 . Consider the next diagram

$$\begin{array}{ccccc} & & \text{Ext}_A^1(\Theta_V M, N) & & \\ & \swarrow \mu_1 \text{ (dashed)} & \downarrow \Psi_2 & & \\ S_V^{-1} \text{Ext}_A^1(\Theta_V M, \Theta_V N) & \xrightarrow{\iota_2} & \text{Ext}_A^1(\Theta_V^2 M, \Theta_V N) & \xrightarrow{\lambda} & \text{Ext}_A^1(I^1, \Theta_V N). \end{array}$$

Since $\lambda \Psi_2 \partial_1 = \lambda \delta_2 = 0$ and ∂_1 is an epimorphism, one has $\lambda \Psi_2 = 0$. Thus there is a morphism $\mu_1 : \text{Ext}_A^1(\Theta_V M, N) \rightarrow S_V^{-1} \text{Ext}_A^1(\Theta_V M, \Theta_V N)$ such that $\iota_2 \mu_1 = \Psi_2$. Set $\Phi_1 = \mu_1 \iota_1$. Similarly, one gets the morphism μ_2 in (3.9.1) with $\iota_3 \mu_2 = \Psi_3$. Set $\Phi_2 = \mu_2 \iota_2$.

Finally, consider the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \overline{\text{Hom}}_A^V(M, N) & \xrightarrow{\overline{\delta_1}} & \text{Ext}_A^1(\Theta_V M, N) & \longrightarrow & \text{Ext}_A^1(I^0, N) \\ & & \eta_1 \downarrow \text{(dotted)} & & \parallel & & \parallel \\ 0 & \longrightarrow & S_V^{-1} \text{Ext}_A^1(M, N) & \xrightarrow{\iota_1} & \text{Ext}_A^1(\Theta_V M, N) & \longrightarrow & \text{Ext}_A^1(I^0, N), \end{array}$$

where the first exact sequence is given in the diagram (3.3.1), the morphism η_1 is obtained by the universal property of kernels. So one has $\iota_1 \eta_1 = \overline{\delta_1}$, moreover, it follows from the Five Lemma that η_1 is an isomorphism. Similarly, one gets the second and third vertical isomorphisms $\eta_2 : \overline{\text{Hom}}_A^V(\Theta_V M, \Theta_V N) \rightarrow S_V^{-1} \text{Ext}_A^1(\Theta_V M, \Theta_V N)$ and $\eta_3 : \overline{\text{Hom}}_A^V(\Theta_V^2 M, \Theta_V^2 N) \rightarrow S_V^{-1} \text{Ext}_A^1(\Theta_V^2 M, \Theta_V^2 N)$ in (3.9.1) such that $\iota_2 \eta_2 = \overline{\delta_2}$ and $\iota_3 \eta_3 = \overline{\delta_3}$. It implies that the triangles on the right side of the first, second and third vertical isomorphisms in (3.9.1) are commutative, respectively.

Since $\iota_2 \eta_2 \beta_1 = \overline{\delta_2} \beta_1 = \Psi_2 = \iota_2 \mu_1$ and ι_2 is a monomorphism, one has $\eta_2 \beta_1 = \mu_1$, which implies that the triangle on the left side of the second vertical isomorphism in (3.9.1) is commutative. Similarly, one obtains the triangle on the left side of the third vertical isomorphism in (3.9.1) is commutative, that is $\eta_3 \beta_2 = \mu_2$. Hence all the squares in (3.9.1) are commutative.

Now continuing this construction one gets the commutative diagram (3.9.1).

3.10 Theorem. *Let M and N be objects in \mathbf{A} . For each $n \in \mathbb{Z}$ there exist natural isomorphisms*

$$\widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N) \cong \text{colim}_i \text{Ext}_{\mathbf{A}}^1(\Theta_{\mathbf{V}}^{i+1}M, \Theta_{\mathbf{V}}^{i+n}N) \cong \text{colim}_i S_{\mathbf{V}}^{-1} \text{Ext}_{\mathbf{A}}^1(\Theta_{\mathbf{V}}^iM, \Theta_{\mathbf{V}}^{i+n}N)$$

and

$$\widehat{\text{Ext}}_{\mathbf{W}}^n(M, N) \cong \text{colim}_i \text{Ext}_{\mathbf{A}}^1(\Omega_{i+n}^{\mathbf{W}}M, \Omega_{i+1}^{\mathbf{W}}N) \cong \text{colim}_i S_{\mathbf{W}}^{-1} \text{Ext}_{\mathbf{A}}^1(\Omega_{i+n}^{\mathbf{W}}M, \Omega_i^{\mathbf{W}}N).$$

Proof. Since the functor colim_i is exact, one gets the first isomorphism by the commutative diagram (3.9.1). The isomorphism

$$\widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N) \cong \text{colim}_i S_{\mathbf{V}}^{-1} \text{Ext}_{\mathbf{A}}^1(\Theta_{\mathbf{V}}^iM, \Theta_{\mathbf{V}}^{i+n}N)$$

holds again by (3.9.1). The remaining isomorphisms can be proved similarly. \square

3.11 Construction. Let M and N be objects in \mathbf{A} and $n \in \mathbb{Z}$. For each $k \geq 1$, the exact sequence $0 \rightarrow \Theta_{\mathbf{V}}^k M \rightarrow I^k \rightarrow \Theta_{\mathbf{V}}^{k+1} M \rightarrow 0$ yields an exact sequence

$$\text{Ext}_{\mathbf{A}}^{n+k}(\Theta_{\mathbf{V}}^k M, N) \xrightarrow{\delta} \text{Ext}_{\mathbf{A}}^{n+k+1}(\Theta_{\mathbf{V}}^{k+1} M, N) \longrightarrow \text{Ext}_{\mathbf{A}}^{n+k+1}(I^k, N).$$

The connecting morphism δ induces a morphism from $\text{Ext}_{\mathbf{A}}^{n+k}(\Theta_{\mathbf{V}}^k M, N)$ to the kernel $S_{\mathbf{V}}^{-1} \text{Ext}_{\mathbf{A}}^{n+k+1}(\Theta_{\mathbf{V}}^k M, N) \cong S_{\mathbf{V}}^{-(k+1)} \text{Ext}_{\mathbf{A}}^{n+k+1}(M, N)$; see Remark 1.6. Composed with the natural embedding from $S_{\mathbf{V}}^{-k} \text{Ext}_{\mathbf{A}}^{n+k}(M, N) \cong S_{\mathbf{V}}^{-1} \text{Ext}_{\mathbf{A}}^{n+k}(\Theta_{\mathbf{V}}^{k-1} M, N)$ to $\text{Ext}_{\mathbf{A}}^{n+k}(\Theta_{\mathbf{V}}^k M, N)$ one gets a morphism

$$\delta: S_{\mathbf{V}}^{-k} \text{Ext}_{\mathbf{A}}^{n+k}(M, N) \longrightarrow S_{\mathbf{V}}^{-(k+1)} \text{Ext}_{\mathbf{A}}^{n+k+1}(M, N).$$

Similarly, consider the connected sequence $\text{Ext}_{\mathbf{A}}^*(M, -)$ of covariant functors, one gets a morphism

$$\partial: S_{\mathbf{W}}^{-k} \text{Ext}_{\mathbf{A}}^{n+k}(M, N) \longrightarrow S_{\mathbf{W}}^{-(k+1)} \text{Ext}_{\mathbf{A}}^{n+k+1}(M, N).$$

As constructed in 3.11, $\{S_{\mathbf{V}}^{-i} \text{Ext}_{\mathbf{A}}^{n+i}(M, N)\}_{i \geq 1}$ and $\{S_{\mathbf{W}}^{-j} \text{Ext}_{\mathbf{A}}^{n+j}(M, N)\}_{j \geq 1}$ are direct systems. Then we have the next result.

3.12 Proposition. *Suppose that $\text{Ext}_{\mathbf{A}}^{\geq 1}({}^{\perp}\mathbf{V}, \mathbf{V}) = 0 = \text{Ext}_{\mathbf{A}}^{\geq 1}(\mathbf{W}, \mathbf{W}^{\perp})$. Let M and N be objects in \mathbf{A} . Then for each $n \in \mathbb{Z}$, there are natural isomorphisms*

$$\widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N) \cong \text{colim}_i S_{\mathbf{V}}^{-i} \text{Ext}_{\mathbf{A}}^{n+i}(M, N)$$

and

$$\widehat{\text{Ext}}_{\mathbf{W}}^n(M, N) \cong \text{colim}_j S_{\mathbf{W}}^{-j} \text{Ext}_{\mathbf{A}}^{n+j}(M, N).$$

Proof. We prove the first isomorphism; the second one is proved dually. Fix an integer n . For each $i \geq \max\{1, 1-n\}$, one gets a natural isomorphism

$$\mathrm{Ext}_{\mathbf{A}}^{n+i}(\Theta_{\mathbf{V}}^i M, N) \cong \mathrm{Ext}_{\mathbf{A}}^1(\Theta_{\mathbf{V}}^i M, \Theta_{\mathbf{V}}^{i+n-1} N)$$

by dimension shifting as $\mathrm{Ext}_{\mathbf{A}}^{\geq 1}(\perp \mathbf{V}, \mathbf{V})=0$. Then it follows from (3.3.2) that there exists an epimorphism

$$\beta_i : \mathrm{Ext}_{\mathbf{A}}^{n+i}(\Theta_{\mathbf{V}}^i M, N) \longrightarrow \overline{\mathrm{Hom}}_{\mathbf{A}}^{\mathbf{V}}(\Theta_{\mathbf{V}}^i M, \Theta_{\mathbf{V}}^{i+n} N).$$

Applying functor $\mathrm{Hom}_{\mathbf{A}}(-, N)$ to the special \mathbf{V} -preenvelope

$$0 \longrightarrow \Theta_{\mathbf{V}}^i M \xrightarrow{\iota} I^i \xrightarrow{\pi} \Theta_{\mathbf{V}}^{i+1} M \longrightarrow 0,$$

one gets a long exact sequence

(3.12.1)

$$\dots \longrightarrow \mathrm{Ext}_{\mathbf{A}}^{n+i}(\Theta_{\mathbf{V}}^i M, N) \xrightarrow{\delta_i} \mathrm{Ext}_{\mathbf{A}}^{n+i+1}(\Theta_{\mathbf{V}}^{i+1} M, N) \longrightarrow \mathrm{Ext}_{\mathbf{A}}^{n+i+1}(I^i, N) \longrightarrow \dots$$

This yields that $\{\mathrm{Ext}_{\mathbf{A}}^{n+i}(\Theta_{\mathbf{V}}^i M, N)\}_{i \geq \max\{1, 1-n\}}$ is a direct system. The functor colim_i is exact, so one gets an epimorphism

$$\beta_{M,N} : \mathrm{colim}_i \mathrm{Ext}_{\mathbf{A}}^{n+i}(\Theta_{\mathbf{V}}^i M, N) \longrightarrow \mathrm{colim}_i \overline{\mathrm{Hom}}_{\mathbf{A}}^{\mathbf{V}}(\Theta_{\mathbf{V}}^i M, \Theta_{\mathbf{V}}^{i+n} N).$$

We next clarify $\beta_{M,N}$ is a monomorphism. Applying functor $\mathrm{Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}^i M, -)$ to the special \mathbf{V} -preenvelope

$$0 \longrightarrow \Theta_{\mathbf{V}}^{i+n-1} N \xrightarrow{\tau} J^{i+n-1} \xrightarrow{\sigma} \Theta_{\mathbf{V}}^{i+n} N \longrightarrow 0,$$

one gets an exact sequence

$$\mathrm{Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}^i M, J^{i+n-1}) \longrightarrow \mathrm{Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}^i M, \Theta_{\mathbf{V}}^{i+n} N) \longrightarrow \mathrm{Ext}_{\mathbf{A}}^1(\Theta_{\mathbf{V}}^i M, \Theta_{\mathbf{V}}^{i+n-1} N) \longrightarrow 0,$$

as $\Theta_{\mathbf{V}}^i M \in \perp \mathbf{V}$ and $J^{i+n-1} \in \mathbf{V}$. An element in $\mathrm{Ker}(\beta_{M,N})$ can be represented by an element $\overline{\varphi}$ in

$$\begin{aligned} \mathrm{Ext}_{\mathbf{A}}^{n+i}(\Theta_{\mathbf{V}}^i M, N) &\cong \mathrm{Ext}_{\mathbf{A}}^1(\Theta_{\mathbf{V}}^i M, \Theta_{\mathbf{V}}^{i+n-1} N) \\ &\cong \mathrm{Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}^i M, \Theta_{\mathbf{V}}^{i+n} N) / \mathrm{Im} \mathrm{Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}^i M, \sigma) \end{aligned}$$

for some i ; one writes $\overline{\varphi} = \varphi + \mathrm{Im} \mathrm{Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}^i M, \sigma)$ with $\varphi \in \mathrm{Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}^i M, \Theta_{\mathbf{V}}^{i+n} N)$. Then it suffices to prove $\delta_i(\overline{\varphi})=0$. It follows from (3.3.2) that

$$\begin{aligned} \beta_{M,N}(\overline{\varphi}) &= \beta_{M,N}(\varphi + \mathrm{Im} \mathrm{Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}^i M, \sigma)) \\ &= \varphi + \mathbf{V} \mathrm{Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}^i M, \Theta_{\mathbf{V}}^{i+n} N) \end{aligned}$$

$$= 0$$

in $\overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(\Theta_{\mathbf{V}}^i M, \Theta_{\mathbf{V}}^{i+n} N)$. This yields that φ factors through an object in \mathbf{V} , and hence through I^i such that $\varphi = \lambda\iota$ for $\lambda \in \text{Hom}_{\mathbf{A}}(I^i, \Theta_{\mathbf{V}}^{i+n} N)$. Consider the following commutative diagram with exact rows:

$$\begin{array}{ccccc} \Theta_{\mathbf{V}}^i M & \xrightarrow{\iota} & I^i & \xrightarrow{\pi} & \Theta_{\mathbf{V}}^{i+1} M \\ \varphi \downarrow & \swarrow \lambda & \psi \downarrow & \swarrow \phi & \omega \downarrow \\ \Theta_{\mathbf{V}}^{i+n} N & \xrightarrow{\tau} & J^{i+n} & \xrightarrow{\sigma} & \Theta_{\mathbf{V}}^{i+n+1} N. \end{array}$$

Since $(\psi - \tau\lambda)\iota = 0$ and the sequence

$$0 \longrightarrow \text{Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}^{i+1} M, J^{i+n}) \xrightarrow{\pi^*} \text{Hom}_{\mathbf{A}}(I^i, J^{i+n}) \xrightarrow{\iota^*} \text{Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}^i M, J^{i+n}) \longrightarrow 0$$

is exact, one has $\psi - \tau\lambda \in \text{Ker } \iota^* = \text{Im } \pi^*$. Thus there is a morphism ϕ in $\text{Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}^{i+1} M, J^{i+n})$ satisfying $\psi - \tau\lambda = \pi^*(\phi) = \phi\pi$. So $\sigma\phi\pi = \sigma(\psi - \tau\lambda) = \sigma\psi = \omega\pi$, which yields that $\sigma\phi = \omega$ as π is an epimorphism. Hence

$$\delta_i(\overline{\varphi}) = \delta_i(\varphi + \text{Im Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}^i M, \sigma)) = \omega + \text{Im Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}^{i+1} M, \sigma).$$

Since $\omega = \sigma\phi = \text{Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}^{i+1} M, \sigma)(\phi) \in \text{Im Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}^{i+1} M, \sigma)$, it follows that $\delta_i(\overline{\varphi}) = 0$. This implies that $\beta_{M,N}$ is a monomorphism, and hence an isomorphism.

Finally, we prove that there is an isomorphism

$$\text{colim}_i \text{Ext}_{\mathbf{A}}^{n+i}(\Theta_{\mathbf{V}}^i M, N) \cong \text{colim}_i \text{S}_{\mathbf{V}}^{-i} \text{Ext}_{\mathbf{A}}^{n+i}(M, N).$$

Consider the exact sequence (3.12.1). Then one gets the next equalities:

$$\begin{aligned} \text{Im } \delta_i &= \text{Ker}(\text{Ext}_{\mathbf{A}}^{n+i+1}(\Theta_{\mathbf{V}}^{i+1} M, N) \longrightarrow \text{Ext}_{\mathbf{A}}^{n+i+1}(I^i, N)) \\ &= \text{S}_{\mathbf{V}}^{-1} \text{Ext}_{\mathbf{A}}^{n+i+1}(\Theta_{\mathbf{V}}^i M, N) \\ &= \text{S}_{\mathbf{V}}^{-i-1} \text{Ext}_{\mathbf{A}}^{n+i+1}(M, N), \end{aligned}$$

where the last equality follows from Remark 1.6. Passing onto direct limits one gets

$$\text{colim}_i \text{Ext}_{\mathbf{A}}^{n+i}(\Theta_{\mathbf{V}}^i M, N) \cong \text{colim}_i \text{Im } \delta_i = \text{colim}_i \text{S}_{\mathbf{V}}^{-i} \text{Ext}_{\mathbf{A}}^{n+i}(M, N).$$

This completes the proof. \square

3.13 Remark. The first isomorphism in Proposition 3.12 was proved by Nucinkis in [23, Theorem 3.6] for $\mathbf{V} = \text{Inj}$ in the category of R -modules, and the second one was proved by Kropholler in [17, Section 3.3] for $\mathbf{W} = \text{Prj}$ in the category of R -modules; see also Celikbas, Christensen, Liang and Piepmeyer [8, Appendix B].

4. Stable cohomology

In this section we consider the stable cohomology with respect to a preenveloping/precovering subcategory (not necessarily special). Throughout this section all complexes are cochain complexes of objects in \mathbf{A} . We start by recalling the definition of stable cohomology that was first introduced by Goichot [15].

4.1. For complexes X and Y of objects in \mathbf{A} , the symbol $\mathrm{Hom}_{\mathbf{A}}(X, Y)$ denotes the complex of abelian groups with the degree- n term

$$\mathrm{Hom}_{\mathbf{A}}(X, Y)^n = \prod_{i \in \mathbb{Z}} \mathrm{Hom}(X^i, Y^{n+i})$$

and the differential given by $\partial(\alpha) = \partial_Y \alpha - (-1)^{|\alpha|} \alpha \partial_X$ for a homogeneous element α . The *bounded Hom-complex* $\overline{\mathrm{Hom}}_{\mathbf{A}}(X, Y)$ is the subcomplex of $\mathrm{Hom}_{\mathbf{A}}(X, Y)$ with degree- n term

$$\overline{\mathrm{Hom}}_{\mathbf{A}}(X, Y)^n = \prod_{i \in \mathbb{Z}} \mathrm{Hom}_{\mathbf{A}}(X^i, Y^{n+i}).$$

We denote by $\widetilde{\mathrm{Hom}}_{\mathbf{A}}(X, Y)$ the quotient complex $\mathrm{Hom}_{\mathbf{A}}(X, Y) / \overline{\mathrm{Hom}}_{\mathbf{A}}(X, Y)$, which is called *stable Hom-complex*.

4.2 Definition. Let M and N be objects in \mathbf{A} with $M \xrightarrow{\simeq} I$ and $N \xrightarrow{\simeq} J$ proper \mathbf{V} -coresolutions of M and N , respectively. For each $n \in \mathbb{Z}$, the *nth bounded cohomology* of M and N with respect to \mathbf{V} is

$$\overline{\mathrm{Ext}}_{\mathbf{V}}^n(M, N) = \mathrm{H}^n(\overline{\mathrm{Hom}}_{\mathbf{A}}(I, J)),$$

and the *nth stable cohomology* of M and N with respect to \mathbf{V} is

$$\widetilde{\mathrm{Ext}}_{\mathbf{V}}^n(M, N) = \mathrm{H}^n(\widetilde{\mathrm{Hom}}_{\mathbf{A}}(I, J)).$$

Dually, let $P \xrightarrow{\simeq} M$ and $Q \xrightarrow{\simeq} N$ be proper \mathbf{W} -resolutions of M and N , respectively. For each $n \in \mathbb{Z}$, the *nth bounded cohomology* of M and N with respect to \mathbf{W} is

$$\overline{\mathrm{Ext}}_{\mathbf{W}}^n(M, N) = \mathrm{H}^n(\overline{\mathrm{Hom}}_{\mathbf{A}}(P, Q)),$$

and the *nth stable cohomology* of M and N with respect to \mathbf{W} is

$$\widetilde{\mathrm{Ext}}_{\mathbf{W}}^n(M, N) = \mathrm{H}^n(\widetilde{\mathrm{Hom}}_{\mathbf{A}}(P, Q)).$$

4.3 Remark. Any two proper \mathbf{V} -coresolutions and proper \mathbf{W} -resolutions of M are homotopy equivalent, respectively; see [13, Section 8.2]. Thus the above definitions of bounded cohomology and stable cohomology are independent of the choices of proper \mathbf{V} -coresolutions and proper \mathbf{W} -resolutions, respectively.

4.4. For a complex $X = \dots \rightarrow X^{n-1} \xrightarrow{\partial^{n-1}} X^n \rightarrow X^{n+1} \rightarrow X^{n+2} \rightarrow \dots$ of objects in \mathbf{A} , the symbol $X^{\supset n}$ denotes the quotient complex

$$\dots \rightarrow 0 \rightarrow \text{Coker } \partial^{n-1} \rightarrow X^{n+1} \rightarrow X^{n+2} \rightarrow \dots,$$

the symbol $X^{\geq n}$ denotes the subcomplex

$$\dots \rightarrow 0 \rightarrow X^n \rightarrow X^{n+1} \rightarrow \dots,$$

and the symbol $\Sigma^n(-)$ denotes the shift functor.

In view of Proposition 2.4, it can be proved similarly as in [23, Theorem 4.4] (see also [8, Appendix B]) that stable cohomology is actually the cohomology given in Section 3. In the following we give the proof for the convenience of the reader.

4.5 Proposition. *Let M and N be objects in \mathbf{A} . Then for all $n \in \mathbb{Z}$, there are natural isomorphisms*

$$\widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N) \cong \widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N) \text{ and } \widetilde{\text{Ext}}_{\mathbf{W}}^n(M, N) \cong \widehat{\text{Ext}}_{\mathbf{W}}^n(M, N).$$

Proof. We prove the first isomorphism; the second one is proved dually.

Let $M \xrightarrow{\sim} I$ and $N \xrightarrow{\sim} J$ be proper \mathbf{V} -coresolutions of M and N , respectively. Suppose that $\tilde{\mu}$ is an element of $\widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N)$ represented by a morphism μ of degree n , which is a chain map in high degrees, i.e., the following diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Theta_{\mathbf{V}}^i M & \longrightarrow & I^i & \longrightarrow & I^{i+1} \longrightarrow \dots \\ & & \hat{\mu}^i \downarrow & & \mu^i \downarrow & & \mu^{i+1} \downarrow \\ 0 & \longrightarrow & \Theta_{\mathbf{V}}^{i+n} N & \longrightarrow & J^{i+n} & \longrightarrow & J^{i+n+1} \longrightarrow \dots \end{array}$$

is commutative up to a sign $(-1)^n$ for $i \gg 0$. Thus μ^i induces a unique element $\hat{\mu}^i \in \text{Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}^i M, \Theta_{\mathbf{V}}^{i+n} N)$. In this way, μ defines an element $\hat{\mu} \in \widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N)$ in view of Proposition 2.4. In order to show this yields a morphism

$$\phi^n : \widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N) \longrightarrow \widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N),$$

it must be verified that $\hat{\mu}$ is independent of the choices of representative μ of $\tilde{\mu}$ in $\widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N)$. If $\tilde{\mu} = \tilde{\nu}$ in $\widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N)$, then $\mu - \nu$ is 0-homotopic in high degrees, so one has the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Theta_{\mathbf{V}}^i M & \xrightarrow{\iota} & I^i & \xrightarrow{d^i} & I^{i+1} \longrightarrow \dots \\ & & \hat{\mu}^i - \hat{\nu}^i \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \Theta_{\mathbf{V}}^{i+n} N & \xrightarrow{\iota'} & J^{i+n} & \longrightarrow & J^{i+n+1} \longrightarrow \dots \end{array}$$

δ^i (dashed arrow from $\Theta_{\mathbf{V}}^i M$ to J^{i+n})
 δ^{i+1} (dashed arrow from I^{i+1} to J^{i+n+1})
 $\mu^i - \nu^i$ (dashed arrow from I^i to J^{i+n})

for all $i \gg 0$. Since $\iota'(\hat{\mu}^i - \hat{\nu}^i) = (\mu^i - \nu^i)\iota = (\iota'\delta^i + \delta^{i+1}d^i)\iota = \iota'\delta^i\iota$ and ι' is a monomorphism, one has $\hat{\mu}^i - \hat{\nu}^i = \delta^i\iota$. Then $\hat{\mu}^i - \hat{\nu}^i$ factors through $I^i \in \mathbf{V}$, whence it is zero in $\overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(\Theta_{\mathbf{V}}^i M, \Theta_{\mathbf{V}}^{i+n} N)$.

We will continue to claim that ϕ^n is an isomorphism. On one hand, let $\tilde{\mu}$ be an element of $\widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N)$ such that $\phi^n(\tilde{\mu}) = 0$ in $\widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N)$. Then for all $i \gg 0$, the induced morphism $\hat{\mu}^i: \Theta_{\mathbf{V}}^i M \rightarrow \Theta_{\mathbf{V}}^{i+n} N$ factors through an object $C \in \mathbf{V}$ and hence through $\iota: \Theta_{\mathbf{V}}^i M \rightarrow I^i$. So one can construct a morphism $\delta^{k+1}: I^{k+1} \rightarrow J^{k+n}$ for each $k \geq i$, so μ is 0-homotopic in high degrees, i.e., $\tilde{\mu} = 0$ in $\widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N)$. On the other hand, let $\hat{\mu} \in \widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N)$ be a family of elements in the direct system of $\overline{\text{Hom}}_{\mathbf{A}}^{\mathbf{V}}(\Theta_{\mathbf{V}}^i M, \Theta_{\mathbf{V}}^{i+n} N)$. Such a family is represented by an element $\bar{\mu} = \mu + \mathbf{V}\text{Hom}_{\mathbf{A}}(\Theta_{\mathbf{V}}^i M, \Theta_{\mathbf{V}}^{i+n} N)$ for some $i \gg 0$. Extending μ , one gets a morphism $I^{\geq i} \rightarrow \Sigma^n(J^{\geq i+n})$, which yields an element $\tilde{\mu}$ in $\widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N)$ with $\phi^n(\tilde{\mu}) = \hat{\mu}$. \square

Since $\widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N)$ and $\widetilde{\text{Ext}}_{\mathbf{W}}^n(M, N)$ (resp., $\widetilde{\text{Ext}}_{\mathbf{W}}^n(M, N)$ and $\widehat{\text{Ext}}_{\mathbf{W}}^n(M, N)$) are naturally isomorphic, we will not distinguish these two notations; we use the notation $\widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N)$ (resp., $\widetilde{\text{Ext}}_{\mathbf{W}}^n(M, N)$).

4.6 Proposition. *Let M and N be objects in \mathbf{A} . Then there are exact sequences*

$$\dots \longrightarrow \overline{\text{Ext}}_{\mathbf{V}}^i(M, N) \longrightarrow \text{Ext}_{\mathbf{AV}}^i(M, N) \longrightarrow \widetilde{\text{Ext}}_{\mathbf{V}}^i(M, N) \longrightarrow \overline{\text{Ext}}_{\mathbf{V}}^{i+1}(M, N) \longrightarrow \dots$$

and

$$\dots \longrightarrow \overline{\text{Ext}}_{\mathbf{W}}^i(M, N) \longrightarrow \text{Ext}_{\mathbf{WA}}^i(M, N) \longrightarrow \widehat{\text{Ext}}_{\mathbf{W}}^i(M, N) \longrightarrow \overline{\text{Ext}}_{\mathbf{W}}^{i+1}(M, N) \longrightarrow \dots$$

Proof. We prove the first one; the second one is proved dually.

Fix proper \mathbf{V} -coresolutions $M \xrightarrow{\simeq} I$ and $N \xrightarrow{\simeq} J$. Then one gets an exact sequence

$$(4.6.1) \quad 0 \longrightarrow \overline{\text{Hom}}_{\mathbf{A}}(I, J) \longrightarrow \text{Hom}_{\mathbf{A}}(I, J) \longrightarrow \widehat{\text{Hom}}_{\mathbf{A}}(I, J) \longrightarrow 0.$$

It follows from a result by Christensen, Frankild and Holm [10, Proposition 2.7] that $\text{Ext}_{\mathbf{AV}}^n(M, N) \cong \mathbf{H}^n(\text{Hom}_{\mathbf{A}}(I, J))$ for all $n \in \mathbb{Z}$. Thus (4.6.1) yields the exact sequence in the statement. \square

4.7 Proposition. *The following statements hold:*

(a) *Let $0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$ be a $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact short exact sequence in \mathbf{A} . Then for each object M in \mathbf{A} there is an exact sequence*

$$\dots \longrightarrow \widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N') \longrightarrow \widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N) \longrightarrow \widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N'') \longrightarrow \widetilde{\text{Ext}}_{\mathbf{V}}^{n+1}(M, N') \longrightarrow \dots$$

(b) Let $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ be a $\text{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact short exact sequence in \mathbf{A} . Then for each object N in \mathbf{A} there is an exact sequence

$$\dots \rightarrow \widetilde{\text{Ext}}_{\mathbf{V}}^n(M'', N) \rightarrow \widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N) \rightarrow \widetilde{\text{Ext}}_{\mathbf{V}}^n(M', N) \rightarrow \widetilde{\text{Ext}}_{\mathbf{V}}^{n+1}(M'', N) \rightarrow \dots$$

Proof. We only prove (a); the statement (b) is proved similarly.

Let $N' \xrightarrow{\simeq} I$ and $N'' \xrightarrow{\simeq} H$ be proper \mathbf{V} -coresolutions of N' and N'' , respectively. Then N has a proper \mathbf{V} -coresolution $N \xrightarrow{\simeq} J$ such that there is a degree-wise split exact sequence $0 \rightarrow I \rightarrow J \rightarrow H \rightarrow 0$; see [13, Remark 8.2.2]. Let $M \xrightarrow{\simeq} L$ be a proper \mathbf{V} -coresolution of M . Then the sequence

$$0 \rightarrow \widetilde{\text{Hom}}_{\mathbf{A}}(L, I) \rightarrow \widetilde{\text{Hom}}_{\mathbf{A}}(L, J) \rightarrow \widetilde{\text{Hom}}_{\mathbf{A}}(L, H) \rightarrow 0$$

is exact, which yields the exact sequence in the statement. \square

The next result is proved dually.

4.8 Proposition. *The following statements hold:*

(a) Let $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ be a $\text{Hom}_{\mathbf{A}}(\mathbf{W}, -)$ -exact short exact sequence in \mathbf{A} . Then for each object N in \mathbf{A} there is an exact sequence

$$\dots \rightarrow \widehat{\text{Ext}}_{\mathbf{W}}^n(M'', N) \rightarrow \widehat{\text{Ext}}_{\mathbf{W}}^n(M, N) \rightarrow \widehat{\text{Ext}}_{\mathbf{W}}^n(M', N) \rightarrow \widehat{\text{Ext}}_{\mathbf{W}}^{n+1}(M'', N) \rightarrow \dots$$

(b) Let $0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$ be a $\text{Hom}_{\mathbf{A}}(\mathbf{W}, -)$ -exact short exact sequence in \mathbf{A} . Then for each object M in \mathbf{A} there is an exact sequence

$$\dots \rightarrow \widehat{\text{Ext}}_{\mathbf{W}}^n(M, N') \rightarrow \widehat{\text{Ext}}_{\mathbf{W}}^n(M, N) \rightarrow \widehat{\text{Ext}}_{\mathbf{W}}^n(M, N'') \rightarrow \widehat{\text{Ext}}_{\mathbf{W}}^{n+1}(M, N') \rightarrow \dots$$

The following are the vanishing results that were advertised in the introduction; a special case where \mathbf{V} is the subcategory of injectives was first proved by Nucinkis in [23, Theorem 3.7].

4.9 Theorem. *Suppose that \mathbf{V} is closed under direct summands. Then for each object N in \mathbf{A} , the following statements are equivalent.*

- (i) $\mathbf{V}\text{-id}_{\mathbf{A}} N < \infty$.
- (ii) $\widetilde{\text{Ext}}_{\mathbf{V}}^n(N, -) = 0 = \widetilde{\text{Ext}}_{\mathbf{V}}^n(-, N)$ for all $n \in \mathbb{Z}$.
- (iii) $\widetilde{\text{Ext}}_{\mathbf{V}}^0(N, N) = 0$.

Proof. The implication (ii) \implies (iii) is clear.

(i) \implies (ii): Let M be an object in \mathbf{A} with $M \xrightarrow{\simeq} J$ a proper \mathbf{V} -coresolution. Since $\mathbf{V}\text{-id}_{\mathbf{A}} N$ is finite, it follows from Lemma 1.11 that there is a proper \mathbf{V} -coresolution $N \xrightarrow{\simeq} I$ with I bounded. So one has $\text{Hom}_{\mathbf{A}}(I, J) = \widetilde{\text{Hom}}_{\mathbf{A}}(I, J)$. This implies

$$\widetilde{\text{Ext}}_{\mathbf{V}}^n(N, M) = 0 = \widetilde{\text{Ext}}_{\mathbf{V}}^n(M, N)$$

for all $n \in \mathbb{Z}$.

(iii) \implies (i): Let $N \xrightarrow{\sim} I$ be a proper \mathbf{V} -coresolution of N . Then one gets $H^0(\widetilde{\text{Hom}}_{\mathbf{A}}(I, I)) = 0$, and so for $\text{id}_I \in \text{Hom}_{\mathbf{A}}(I, I)^0$ one has

$$\text{id}_I + \overline{\text{Hom}}_{\mathbf{A}}(I, I)^0 \in Z^0(\widetilde{\text{Hom}}_{\mathbf{A}}(I, I)) = B^0(\widetilde{\text{Hom}}_{\mathbf{A}}(I, I)).$$

Thus there is a morphism $\varphi \in \text{Hom}_{\mathbf{A}}(I, I)^{-1}$ such that $\text{id}_I - \partial(\varphi) \in \overline{\text{Hom}}_{\mathbf{A}}(I, I)^0$ is bounded. So there is an integer $i \gg 0$ such that $\partial_I^{i-1} \varphi^i + \varphi^{i+1} \partial_I^i = \text{id}_{I^i}$. Thus one has $\partial_I^{i-1} \varphi^i \partial_I^{i-1} = \partial_I^{i-1}$, which yields that the epimorphism $\partial_I^{i-1}: I^{i-1} \rightarrow \text{Im } \partial_I^{i-1}$ is split. \mathbf{V} is closed under direct summands, so one has $\text{Im } \partial_I^{i-1} \in \mathbf{V}$. Thus $\mathbf{V}\text{-id}_{\mathbf{A}} N$ is finite. \square

4.10 Corollary. *Suppose that \mathbf{V} is closed under direct summands. Then for each object N in \mathbf{A} and $n \in \mathbb{Z}$, the following statements are equivalent.*

- (i) $\text{Ext}_{\mathbf{AV}}^i(C, N) = 0$ for each object $C \in \mathbf{V}$ and all $i \geq n$.
- (ii) $\overline{\text{Ext}}_{\mathbf{V}}^i(-, N) = 0$ for all $i \geq n$.

Proof. (i) \implies (ii): Let M be an object with $M \xrightarrow{\sim} I$ a proper \mathbf{V} -coresolution, and let $\alpha: N \xrightarrow{\sim} J$ be a proper \mathbf{V} -coresolution of N . For each $C \in \mathbf{V}$, the complex $\text{Hom}_{\mathbf{A}}(C, (\text{Cone } \alpha)^{\triangleright n-2})$ is acyclic by (i), where $\text{Cone } \alpha$ denotes the mapping cone of α . Thus by a result by Celikbas, Christensen, Liang and Piepmeyer [7, Proposition A.2], the complex $\overline{\text{Hom}}_{\mathbf{A}}(I, (\text{Cone } \alpha)^{\triangleright n-2})$ is acyclic. So for each $i \geq n$ one has

$$\overline{\text{Ext}}_{\mathbf{V}}^i(M, N) = H^i(\overline{\text{Hom}}_{\mathbf{A}}(I, J)) = H^i(\overline{\text{Hom}}_{\mathbf{A}}(I, (\text{Cone } \alpha)^{\triangleright n-2})) = 0.$$

(ii) \implies (i): For each $C \in \mathbf{V}$ and each $i \in \mathbb{Z}$, one has $\widetilde{\text{Ext}}_{\mathbf{V}}^i(C, N) = 0$ by Theorem 4.9. So $\text{Ext}_{\mathbf{AV}}^i(C, N) = 0$ for each $i \geq n$ by Proposition 4.6. \square

Dually, we have the following two results.

4.11 Theorem. *Suppose that \mathbf{W} is closed under direct summands. Then for each object M in \mathbf{A} , the following statements are equivalent.*

- (i) $\text{W-pd}_{\mathbf{A}} M < \infty$.
- (ii) $\widehat{\text{Ext}}_{\mathbf{W}}^n(-, M) = 0 = \widehat{\text{Ext}}_{\mathbf{W}}^n(M, -)$ for all $n \in \mathbb{Z}$.
- (iii) $\widehat{\text{Ext}}_{\mathbf{W}}^0(M, M) = 0$.

4.12 Corollary. *Suppose that \mathbf{W} is closed under direct summands. Then for each object N in \mathbf{A} and $n \in \mathbb{Z}$, the following statements are equivalent.*

- (i) $\text{Ext}_{\mathbf{WA}}^i(M, D) = 0$ for each object $D \in \mathbf{W}$ and all $i \geq n$.
- (ii) $\overline{\text{Ext}}_{\mathbf{W}}^i(M, -) = 0$ for all $i \geq n$.

We end this section with a new computation of stable cohomology $\widetilde{\text{Ext}}_{\mathbf{V}}^n(-, -)$ (resp., $\widehat{\text{Ext}}_{\mathbf{W}}^n(-, -)$) via Tate \mathbf{V} -coresolutions (resp., Tate \mathbf{W} -resolutions).

4.13 Tate (co)resolutions. Recall from [27] that a complex T of objects in \mathbf{V} is *totally \mathbf{V} -acyclic* if it is acyclic and the complex $\mathrm{Hom}_{\mathbf{A}}(C, T)$ and $\mathrm{Hom}_{\mathbf{A}}(T, C)$ are acyclic for each object $C \in \mathbf{V}$. Let N be an object in \mathbf{A} . A *Tate \mathbf{V} -coresolution* of N is a diagram $N \xrightarrow{\simeq} I \xrightarrow{\alpha} T$ wherein T is a totally \mathbf{V} -acyclic complex of objects in \mathbf{V} , $N \xrightarrow{\simeq} I$ is a proper \mathbf{V} -coresolution of N , and α^n is an isomorphism for $n \gg 0$.

Dually, one has the definitions of a *totally \mathbf{W} -acyclic* complex H and a *Tate \mathbf{W} -resolution* $H \xrightarrow{\gamma} P \xrightarrow{\simeq} M$ of M .

The next result generalizes [23, Theorem 7.3].

4.14 Proposition. *Suppose that \mathbf{V} is closed under direct summands. Let N be an object in \mathbf{A} that has a Tate \mathbf{V} -coresolution $N \xrightarrow{\simeq} I \xrightarrow{\alpha} T$. Then for each object M in \mathbf{A} and all $n \in \mathbb{Z}$, there is a natural isomorphism*

$$\widetilde{\mathrm{Ext}}_{\mathbf{V}}^n(M, N) \cong \mathrm{H}^n(\mathrm{Hom}_{\mathbf{A}}(M, T)).$$

Proof. Fix $n \in \mathbb{Z}$, and let $p \geq n$ such that α^i is an isomorphism for each $i \geq p$. Set $L = \mathrm{Ker}(\partial_I^p)$ and $K = \mathrm{Ker}(\partial_T^{n-1})$. Since the complex T is $\mathrm{Hom}_{\mathbf{A}}(-, \mathbf{V})$ -exact, $K \xrightarrow{\simeq} \Sigma^{n-1} T^{\geq n-1}$ is a proper \mathbf{V} -coresolution of K . Thus for each object $C \in \mathbf{V}$ and $i \geq 1$, one has

$$\begin{aligned} \mathrm{Ext}_{\mathbf{AV}}^i(C, K) &= \mathrm{H}^i(\mathrm{Hom}_{\mathbf{A}}(C, \Sigma^{n-1} T^{\geq n-1})) \\ &= \mathrm{H}^{i+n-1}(\mathrm{Hom}_{\mathbf{A}}(C, T^{\geq n-1})) \\ &= \mathrm{H}^{i+n-1}(\mathrm{Hom}_{\mathbf{A}}(C, T)) \\ &= 0, \end{aligned}$$

where the last equality holds as T is $\mathrm{Hom}_{\mathbf{A}}(\mathbf{V}, -)$ -exact. This yields $\overline{\mathrm{Ext}}_{\mathbf{V}}^i(M, K) = 0$ for all $i \geq 1$ by Corollary 4.10. Thus from Proposition 4.6 one gets $\overline{\mathrm{Ext}}_{\mathbf{V}}^1(M, K) \cong \mathrm{Ext}_{\mathbf{AV}}^1(M, K)$. This is the third isomorphism in the next computation

$$\begin{aligned} \widetilde{\mathrm{Ext}}_{\mathbf{V}}^n(M, N) &\cong \widetilde{\mathrm{Ext}}_{\mathbf{V}}^{n-p}(M, L) \\ &\cong \widetilde{\mathrm{Ext}}_{\mathbf{V}}^1(M, K) \\ &\cong \mathrm{Ext}_{\mathbf{AV}}^1(M, K) \\ &\cong \mathrm{H}^1(\mathrm{Hom}_{\mathbf{A}}(M, \Sigma^{n-1} T^{\geq n-1})) \\ &= \mathrm{H}^n(\mathrm{Hom}_{\mathbf{A}}(M, T^{\geq n-1})) \\ &= \mathrm{H}^n(\mathrm{Hom}_{\mathbf{A}}(M, T)). \end{aligned}$$

The first two isomorphisms hold by Proposition 4.7 and Theorem 4.9; the fourth one holds as $K \xrightarrow{\simeq} \Sigma^{n-1} T^{\geq n-1}$ is a proper \mathbf{V} -coresolution of K . \square

Dually, we have the following result.

4.15 Proposition. *Suppose that \mathcal{W} is closed under direct summands. Let M be an object in \mathcal{A} that has a Tate \mathcal{W} -resolution $H \rightarrow P \xrightarrow{\simeq} M$. Then for each object N in \mathcal{A} and all $n \in \mathbb{Z}$, there is a natural isomorphism*

$$\widehat{\text{Ext}}_{\mathcal{W}}^n(M, N) \cong H^n(\text{Hom}_{\mathcal{A}}(H, N)).$$

4.16 Remark. We will study the relation between $\widetilde{\text{Ext}}_{\mathcal{V}}^n(M, N)$ and $\widehat{\text{Ext}}_{\mathcal{W}}^n(M, N)$ when $(\mathcal{W}, \mathcal{V})$ is a balanced pair, and give a balance result in [16]; see 1.9 for the definition of balanced pairs.

5. Applications of the vanishing results

Let \mathcal{A} be the category $R\text{-Mod}$ of left R -modules, where R is an associative ring. In this section, we give some applications of the vanishing results showed in the above section, and give some characterizations of modules of finite homological dimension including the flat dimension, cotorsion dimension, Gorenstein injective (flat) dimension and projectively coresolved Gorenstein flat dimension.

5.1 Gorenstein injective objects. Recall from Enochs and Jenda [12] that a left R -module M is called *Gorenstein injective* if there is a $\text{Hom}_R(\text{Inj}, -)$ -exact acyclic complex $\dots \rightarrow E_1 \rightarrow E_0 \rightarrow E^0 \rightarrow E^1 \rightarrow \dots$ of injective left R -modules such that $M \cong \text{Ker}(E^0 \rightarrow E^1)$, where Inj is the subcategory of injective left R -modules. Let $\mathcal{V} = \text{GInj}$ be the subcategory of Gorenstein injective left R -modules. It was proved by Šaroch and Št'ovíček that \mathcal{V} is a special preenveloping subcategory; see [25, Theorem 5.6]. In this case, the \mathcal{V} -injective dimension $\mathcal{V}\text{-id}N$ of a left R -module N is actually the Gorenstein injective dimension $\text{Gid}_R N$.

The next result gives a characterization of modules of finite Gorenstein injective dimension; it follows from Theorem 4.9.

5.2 Proposition. *The following are equivalent for a left R -module N .*

- (i) $\text{Gid}_R N < \infty$.
- (ii) $\widetilde{\text{Ext}}_{\text{GInj}}^n(N, -) = 0 = \widetilde{\text{Ext}}_{\text{GInj}}^n(-, N)$ for all $n \in \mathbb{Z}$.
- (iii) $\widetilde{\text{Ext}}_{\text{GInj}}^0(N, N) = 0$.

5.3 Flat and cotorsion objects. A left R -module M is called *cotorsion* (see Enochs [11]) if $\text{Ext}_R^1(F, M) = 0$ for each flat left R -module F . Let $\mathcal{V} = \text{Cot}$ be the subcategory of cotorsion left R -modules and $\mathcal{W} = \text{Flat}$ the subcategory of flat left R -modules. Then \mathcal{V} is a special preenveloping subcategory and \mathcal{W} is a special precovering subcategory; see [5, Proposition 2]. In this case, the \mathcal{V} -injective dimension $\mathcal{V}\text{-id}N$ of a left R -module N is actually the cotorsion dimension $\text{cd}_R N$; see Mao and Ding [24]. Meanwhile, the \mathcal{W} -projective dimension $\mathcal{W}\text{-pd}M$ of a left R -module M is actually the flat dimension $\text{fd}_R M$.

We have the following two results by applying Theorem 4.9 and Theorem 4.11 respectively.

5.4 Proposition. *The following are equivalent for a left R -module N .*

- (i) $\text{cd}_R N < \infty$.
- (ii) $\widehat{\text{Ext}}_{\text{Cot}}^n(N, -) = 0 = \widetilde{\text{Ext}}_{\text{Cot}}^n(-, N)$ for all $n \in \mathbb{Z}$.
- (iii) $\widehat{\text{Ext}}_{\text{Cot}}^0(N, N) = 0$.

5.5 Proposition. *The following are equivalent for a left R -module M .*

- (i) $\text{fd}_R M < \infty$.
- (ii) $\widehat{\text{Ext}}_{\text{Flat}}^n(-, M) = 0 = \widehat{\text{Ext}}_{\text{Flat}}^n(M, -)$ for all $n \in \mathbb{Z}$.
- (iii) $\widehat{\text{Ext}}_{\text{Flat}}^0(M, M) = 0$.

5.6 Gorenstein flat modules. Recall from [13] that a left R -module M is called *Gorenstein flat* if there exists an acyclic complex $\dots \rightarrow F_1 \rightarrow F_0 \rightarrow F^0 \rightarrow F^1 \rightarrow \dots$ of flat left R -modules with $M \cong \text{Ker}(F^0 \rightarrow F^1)$ such that it remains exact after applying the functor $E \otimes_R -$ for each injective right R -module E . Let $\mathbf{W} = \mathbf{GFlat}$ be the subcategory of Gorenstein flat left R -modules. Then \mathbf{W} is a special precovering subcategory by [25, Corollary 4.12], and the \mathbf{W} -projective dimension $\mathbf{W}\text{-pd}M$ of a left R -module M is actually the Gorenstein flat dimension $\text{Gfd}_R M$.

The next result is immediate by Theorem 4.11.

5.7 Proposition. *The following are equivalent for a left R -module M .*

- (i) $\text{Gfd}_R M < \infty$.
- (ii) $\widehat{\text{Ext}}_{\mathbf{GFlat}}^n(-, M) = 0 = \widehat{\text{Ext}}_{\mathbf{GFlat}}^n(M, -)$ for all $n \in \mathbb{Z}$.
- (iii) $\widehat{\text{Ext}}_{\mathbf{GFlat}}^0(M, M) = 0$.

5.8 Projectively coresolving Gorenstein flat modules. Recall from [25] that a left R -module M is called *projectively coresolved Gorenstein flat* if there exists an acyclic complex $\dots \rightarrow P_1 \rightarrow P_0 \rightarrow P^0 \rightarrow P^1 \rightarrow \dots$ of projective left R -modules with $M \cong \text{Ker}(P^0 \rightarrow P^1)$ such that it remains exact after applying the functor $E \otimes_R -$ for each injective right R -module E . Let $\mathbf{W} = \mathbf{PGF}$ be the subcategory of projectively coresolved Gorenstein flat left R -modules. Then \mathbf{W} is a special precovering subcategory by [25, Theorem 4.9], and the \mathbf{W} -projective dimension $\mathbf{W}\text{-pd}M$ of a left R -module M is actually the projectively coresolved Gorenstein flat dimension $\text{PGF-dim}_R M$.

The next result is immediate by Theorem 4.11.

5.9 Proposition. *The following are equivalent for a left R -module M .*

- (i) $\text{PGF-dim}_R M < \infty$.
- (ii) $\widehat{\text{Ext}}_{\mathbf{PGF}}^n(-, M) = 0 = \widehat{\text{Ext}}_{\mathbf{PGF}}^n(M, -)$ for all $n \in \mathbb{Z}$.
- (iii) $\widehat{\text{Ext}}_{\mathbf{PGF}}^0(M, M) = 0$.

Acknowledgments. The authors would like to express sincere thanks to the referee for valuable comments, suggestions and corrections which resulted in a significant improvement of the paper. The authors also thank Xiaoyan Yang for helpful discussions related to this work. This research was partly supported by NSF of China (Grant No. 12271230), NSF of Gansu Province (Grant No. 22JR5RA375), and the Foundation for Innovative Fundamental Research Group Project of Gansu Province (Grant No. 23JRRA684).

References

1. AUSLANDER, L. and BRIDGER, M., *Stable Module Theory*, Memories of the American Mathematical Society **94**, Am. Math. Soc., Providence, 1969.
2. AVRAMOV, L. L. and MARTSINKOVSKY, A., Absolute, relative, and Tate cohomology of modules of finite Gorenstein dimension, *Proc. Lond. Math. Soc.* **85** (2002), 393–440.
3. AVRAMOV, L. L. and VELICHE, O., Stable cohomology over local rings, *Adv. Math.* **213** (2007), 93–139.
4. BENSON, D. J. and CARLSON, J. F., Products in negative cohomology, *J. Pure Appl. Algebra* **82** (1992), 107–129.
5. BICAN, L., BASHIR, R. E. I. and ENOCHS, E. E., All modules have flat covers, *Bull. Lond. Math. Soc.* **33** (2001), 385–390.
6. CARTAN, H. and EILENBERG, S., *Homological Algebra*, Princeton University Press, Princeton, 1956.
7. CELIKBAS, O., CHRISTENSEN, L. W., LIANG, L. and PIEPMEYER, G., Stable homology over associative rings, *Trans. Amer. Math. Soc.* **369** (2017), 8061–8086.
8. CELIKBAS, O., CHRISTENSEN, L. W., LIANG, L. and PIEPMEYER, G., Complete homology over associative rings, *Israel J. Math.* **221** (2017), 1–24.
9. CHEN, X.-W., Homotopy equivalences induced by balanced pairs, *J. Algebra* **324** (2010), 2718–2731.
10. CHRISTENSEN, L. W., FRANKILD, A. and HOLM, H., On Gorenstein projective, injective and flat dimensions – a functorial description with applications, *J. Algebra* **302** (2006), 231–279.
11. ENOCHS, E. E., Flat covers and flat cotorsion modules, *Proc. Amer. Math. Soc.* **92** (1984), 179–184.
12. ENOCHS, E. E. and JENDA, O. M. G., Gorenstein injective and projective modules, *Math. Z.* **220** (1995), 611–633.
13. ENOCHS, E. E. and JENDA, O. M. G., *Relative Homological Algebra*, de Gruyter Expositions in Mathematics **30**, de Gruyter, Berlin, 2000.
14. ESTRADA, S., PÉREZ, M. A. and ZHU, H., Balanced pairs, cotorsion triplets and quiver representations, *Proc. Edinb. Math. Soc.* **63** (2020), 67–90.
15. GOICHOT, F., Homologie de Tate-Vogel équivariante, *J. Pure Appl. Algebra* **82** (1992), 39–64.
16. GUO, S. T., LIANG, L. and YANG, X. Y., Balancedness of complete cohomology with respect to subcategories, preprint.

17. KROPHOLLER, P. H., Hierarchical decompositions, generalized Tate cohomology, and groups of type $(FP)_{\infty}$, in *Combinatorial and Geometric Group Theory*, London Math. Soc. Lecture Note Ser. **204**, *Edinburgh, 1993*, pp. 190–216, Cambridge University Press, Cambridge, 1995.
18. MARTSINKOVSKY, A. and RUSSELL, J., Injective stabilization of additive functors, I. Preliminaries, *J. Algebra* **530** (2019), 429–469.
19. MARTSINKOVSKY, A. and RUSSELL, J., Injective stabilization of additive functors, II. Co(torsion) and the Auslander-Gruson-Jensen functor, *J. Algebra* **548** (2020), 53–95.
20. MARTSINKOVSKY, A. and RUSSELL, J., Injective stabilization of additive functors, III. Asymptotic stabilization of the tensor product, *Algebra Discrete Math.* **31** (2021), 120–151.
21. MARTSINKOVSKY, A. and ZANGURASHVILI, D., The stable category of a left hereditary ring, *J. Pure Appl. Algebra* **219** (2015), 4061–4089.
22. MISLIN, G., Tate cohomology for arbitrary groups via satellites, *Topology Appl.* **56** (1994), 293–300.
23. NUCINKIS, B. E. A., Complete cohomology for arbitrary rings using injectives, *J. Pure Appl. Algebra* **131** (1998), 297–318.
24. MAO, L. X. and DING, N. Q., The cotorsion dimension of modules and rings, in *Abelian Groups, Rings, Modules, and Homological Algebra*, Lecture Notes in Pure and Applied Mathematics **249**, pp. 217–233, CRC Press, Boca Raton, 2006.
25. ŠAROCH, J. and ŠT’OVÍČEK, J., Singular compactness and definability for Σ -cotorsion and Gorenstein modules, *Selecta Math. (N.S.)* **26** (2020), Paper No. 23.
26. SALCE, L., Cotorsion theories for abelian groups, *Sympos. Math.* **23** (1978), 11–32.
27. SATHER-WAGSTAFF, S., SHARIF, T. and WHITE, D., Tate cohomology with respect to semidualizing modules, *J. Algebra* **324** (2010), 2336–2368.

Shoutao Guo
 Department of Mathematics
 Lanzhou Jiaotong University
 Lanzhou CN-730070
 China
guoshoutao9022@163.com

Li Liang
 (Corresponding author)
 Department of Mathematics
 Lanzhou Jiaotong University
 Gansu Provincial Research Center for Basic Disciplines of Mathematics and Statistics
 Lanzhou CN-730070
 China
liliangnju@gmail.com
<https://sites.google.com/site/liliangnju>

Received January 16, 2023
in revised form June 4, 2023