

Squaring the magic

SERGIO L. CACCIATORI, BIANCA L. CERCHIAI AND ALESSIO MARRANI

We construct and classify all possible Magic Squares (MS's) related to Euclidean or Lorentzian rank-3 simple Jordan algebras, both on normed division algebras and split composition algebras. Besides the known Freudenthal-Rozenfeld-Tits MS, the single-split Günaydin-Sierra-Townsend MS, and the double-split Barton-Sudbery MS, we obtain other 7 Euclidean and 10 Lorentzian novel MS's.

We elucidate the role and the meaning of the various non-compact real forms of Lie algebras, entering the MS's as symmetries of theories of Einstein-Maxwell gravity coupled to nonlinear sigma models of scalar fields, possibly endowed with local supersymmetry, in $D = 3, 4$ and 5 space-time dimensions. In particular, such symmetries can be recognized as the U -dualities or the stabilizers of scalar manifolds within space-time with standard Lorentzian signature or with other, more exotic signatures, also relevant to suitable compactifications of the so-called M^* - and M' - theories. Symmetries pertaining to some attractor U -orbits of magic supergravities in Lorentzian space-time also arise in this framework.

1. Introduction

Magic Squares (MS's), arrays of Lie algebras enjoying remarkable symmetry properties under reflection with respect to their main diagonal, were discovered long time ago by Freudenthal, Rozenfeld and Tits [1–3], and their structure and fascinating properties have been studied extensively in mathematics and mathematical physics, especially in relation to exceptional Lie algebras (see *e.g.* [4–12]).

Following the seminal papers by Günaydin, Sierra and Townsend [13, 14], MS's have been related to the generalized electric-magnetic (U -)duality¹ symmetries of particular classes of Maxwell-Einstein supergravity theories

¹Here U -duality is referred to as the “continuous” symmetries of [15]. Their discrete versions are the U -duality non-perturbative string theory symmetries introduced by Hull and Townsend [16].

(MESGT's), called *magic* (see also [17–21]). In particular, non-compact, real forms of Lie algebras, corresponding to non-compact symmetries of (super)gravity theories, have become relevant as symmetries of the corresponding rank-3 *simple* Jordan algebras [22], defined over normed division ($\mathbb{A} = \mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{O}$) or split ($\mathbb{A}_S = \mathbb{R}, \mathbb{C}_S, \mathbb{H}_S, \mathbb{O}_S$) composition algebras [23].

Later on, some other MS's have been constructed in literature through the exploitation of Tits' formula [2] (*cfr.* (2.1) below). On the other hand, the role of *Lorentzian* rank-3 simple Jordan algebras in constructing unified MESGT's in $D = 5$ and 4 Lorentzian space-time dimensions (through the determination of the cubic Chern-Simons *FFA* coupling in the Lagrangian density) has been investigated in [24–26].

In the present paper, we focus on Tits' formula (and its trialitarian reformulation, namely Vinberg's formula [4]; *cfr.* (2.17) below), and construct and classify all possible MS structures consistent with Euclidean or Lorentzian rank-3 simple Jordan algebras. We also elucidate the MS structure, in terms of maximal and symmetric embeddings on their rows and columns.

It should be remarked that most of the MS's which we determine (classified according to the sequences of algebras entering their rows and columns) are new and never appeared in literature. Indeed, as mentioned above, before the present survey only particular types of MS's, exclusively related to Euclidean Jordan algebras, were known, namely the original Freudenthal-Rozenfeld-Tits (FRT) MS $\mathcal{L}_3(\mathbb{A}, \mathbb{B})$ [1–3], the single-split supergravity Günaydin-Sierra-Townsend (GST) MS $\mathcal{L}_3(\mathbb{A}_S, \mathbb{B})$ [13], and the double-split Barton-Sudbery (BS) MS $\mathcal{L}_3(\mathbb{A}_S, \mathbb{B}_S)$ [8] (which also appeared in [27]). Besides these ones, only a particular “mixed” MS (denoted as $\mathcal{L}_3(\tilde{\mathbb{A}}, \mathbb{B})$ in our classification; see below) recently appeared in [21], in the framework of an explicit construction of a manifestly maximally covariant symplectic frame for the special Kähler geometry of the scalar fields of $D = 4$ magic MESGT's. The entries of the last row/ column of the magic squares have been computed also in [28], depending on the norm of the composition algebras involved.

Furthermore, we elucidate the role and the meaning of the various non-compact, real forms of Lie algebras as symmetries of Einstein-Maxwell gravity theories coupled to non-linear sigma models of scalar fields, possibly endowed with local supersymmetry. We consider U -dualities in $D = 3, 4$ and 5 space-time dimensions, with the standard Lorentzian signature or with other, more exotic signatures, such as the Euclidean one and others with two timelike dimensions. Interestingly, symmetries pertaining to particular compactifications of 11-dimensional theories alternative to M -theory,

namely to the so-called M^* -theory and M' -theory [29, 30], appear in this framework.

Frequently, the Lie algebras entering the MS's also enjoy an interpretation as stabilizers of certain orbits of an irreducible representation of the U -duality itself, in which the (Abelian) field strengths of the theory sit (possibly, along with their duals). The *stratification* of the related representation spaces under U -duality has been extensively studied in the supergravity literature, starting from [31, 32] (see *e.g.* [33] for a brief introduction), in relation to extremal black hole solutions and their attractor behaviour (see *e.g.* [34] for a comprehensive review).

A remarkable role is played by exceptional Lie algebras. It is worth observing that the particular non-compact real forms² $\mathfrak{f}_{4(-20)}$ and $\mathfrak{e}_{6(-14)}$, occurring as particular symmetries of flux configurations supporting non-supersymmetric attractors in magic MESGT's, can be obtained in the framework of MS's *only* by considering *Lorentzian* rank-3 Jordan algebras on division or split algebras.

Thus, the present investigation not only classifies all MS's based on rank-3 Euclidean or Lorentzian simple Jordan algebras, but also clarifies their role in generating non-compact symmetries of the corresponding (possibly, locally supersymmetric) theories of gravity in various dimensions and signatures of space-time.

The plan of the paper is as follows.

In Sec. 2, we recall some basic facts and definitions on rank-3 (*alias* cubic) Jordan algebras and MS's, and present Tits' and Vinberg's formulæ, which will be crucial for our classification.

Then, in Sec. 3 we compute and classify all 4×4 MS's based on rank-3 simple (generic) Jordan algebras of *Euclidean* type. We recover the known FRT, GST and BS MS's, and other 7 independent MS arrays, and we analyze the role of the corresponding symmetries in (super)gravity theories.

Sec. 4 deals with rank-3 simple (generic) Jordan algebras of *Lorentzian* type, and with the corresponding MS structures, all previously unknown. In particular, the Lorentzian FRT MS (Table 11), which is symmetric and contains only non-compact Lie algebras, is relevant to certain (non-supersymmetric) attractors in the corresponding theory.

²For simplicity's sake, in the following treatment, we will not distinguish between *algebra* level and *group* level. In the present investigation, indeed, we are not interested in dealing with various *discrete* factors \mathbb{Z}_n possibly arising at group level [10].

A detailed analysis of the MS structure, and further group-theoretical and physical considerations, are given in the concluding Sec. 5.

2. Magic squares and Jordan algebras

We start by briefly recalling the definition of a magic square: A *Magic Square* (MS) is an array of Lie algebras $\mathcal{L}(\mathbb{A}, \mathbb{B})$, where \mathbb{A} and \mathbb{B} are normed division or split composition algebras which label the rows and columns, respectively. The entries of $\mathcal{L}(\mathbb{A}, \mathbb{B})$ are determined by *Tits' formula* [2]:

$$(2.1) \quad \mathcal{L}(\mathbb{A}, \mathbb{B}) = \text{Der}(\mathbb{A}) \oplus \text{Der}(\mathfrak{J}^{\mathbb{B}}) \dot{+} (\mathbb{A}' \otimes \mathfrak{J}'^{\mathbb{B}}).$$

The symbol \oplus denotes direct sum of algebras, whereas $\dot{+}$ stands for direct sum of vector spaces. Moreover, Der are the linear derivations, with $\mathfrak{J}^{\mathbb{B}}$ we indicate the Jordan algebra on \mathbb{B} , and the prime amounts to considering only traceless elements.

In order to understand all these ingredients of the Tits' formula (2.1), it is necessary to introduce some notation first. The octonions are defined through the isomorphism $\mathbb{O} \cong \langle 1, e_1, \dots, e_7 \rangle_{\mathbb{R}}$, where $\langle \cdot \rangle_{\mathbb{R}}$ means the real span. The multiplication rule of the octonions is described by the Fano plane:




Figure 1: The Fano plane and the octonionic product

Let (e_i, e_j, e_k) be an ordered triple of points lying on a given line with the order specified by the direction of the arrow. Then the multiplication is

given by:

$$e_i e_j = e_k, \quad \text{and} \quad e_j e_i = -e_k,$$

together with:

$$e_i^2 = -1, \quad \text{and} \quad 1 e_i = e_i 1 = e_i.$$

\mathbb{O}' denotes the imaginary octonions. The split octonions \mathbb{O}_S can be obtained e.g. by substituting the imaginary units $e_i \rightarrow \tilde{e}_i$, $i = 4, 5, 6, 7$, so that they satisfy $\tilde{e}_i^2 = 1$ instead of $e_i^2 = -1$ (see e.g. [35]).

If the quaternions \mathbb{H} and the complex numbers \mathbb{C} are represented e.g. by the isomorphisms: $\mathbb{H}_S \cong \langle 1, e_1, e_5, e_6 \rangle_{\mathbb{R}}$, and $\mathbb{C}_S \cong \langle 1, e_4 \rangle_{\mathbb{R}}$, the split quaternions \mathbb{H}_S and the split complex numbers \mathbb{C}_S can be represented by the isomorphisms:

$$(2.2) \quad \mathbb{H}_S \cong \langle 1, e_1, \tilde{e}_5, \tilde{e}_6 \rangle_{\mathbb{R}}, \quad \mathbb{C}_S \cong \langle 1, \tilde{e}_4 \rangle_{\mathbb{R}}.$$

As for the octonions, the prime denotes the purely imaginary quaternions \mathbb{H}' and complex numbers \mathbb{C}' , respectively.

An inner product can be defined on any of the above division algebras \mathbb{A} as:

$$(2.3) \quad \langle x_1, x_2 \rangle := \text{Re}(\bar{x}_1 x_2), \quad x_1, x_2 \in \mathbb{A},$$

where the conjugation “ $\bar{\cdot}$ ” changes the sign of the imaginary part.

The algebra of derivations $\text{Der}(\mathbb{A})$ is given by:

$$(2.4) \quad \text{Der}(\mathbb{A}) := \{D \in \text{End}(\mathbb{A}) \mid D(x_1 x_2) = D(x_1) x_2 + x_1 D(x_2) \\ \forall x_1, x_2 \in \mathbb{A}\},$$

i.e. by the maps satisfying the Leibniz rule. Then, if L and R respectively are the left and right translation in \mathbb{A} , a derivation $D_{x_1, x_2} \in \text{Der}(\mathbb{A})$ can be constructed from $x_1, x_2 \in \mathbb{A}$ as:

$$(2.5) \quad D_{x_1, x_2} := [L_{x_1}, L_{x_2}] + [R_{x_1}, R_{x_2}] + [L_{x_1}, R_{x_2}],$$

which, when applied to an element $x_3 \in \mathbb{A}$, becomes:

$$D_{x_1, x_2}(x_3) = [[x_1, x_2], x_3] - 3((x_1 x_2) x_3 - x_1 (x_2 x_3)).$$

The main ingredient entering in the Tits’ formula (2.1) is the Jordan algebra \mathfrak{J} [22, 23], which is defined in the following way: A *Jordan algebra*

\mathfrak{J} is a vector space defined over a *ground field* \mathbb{F} , equipped with a bilinear product \circ satisfying:

$$(2.6) \quad X \circ Y = Y \circ X;$$

$$(2.7) \quad X^2 \circ (X \circ Y) = X \circ (X^2 \circ Y), \quad \forall X, Y \in \mathfrak{J}.$$

The Jordan algebras relevant for the present investigation are *rank-3* Jordan algebras \mathfrak{J}_3 over $\mathbb{F} = \mathbb{R}$, which also come equipped with a cubic norm:

$$(2.8) \quad N : \mathfrak{J} \rightarrow \mathbb{R},$$

$$N(\lambda X) = \lambda^3 N(X), \quad \forall \lambda \in \mathbb{R}, X \in \mathfrak{J}.$$

There is a general prescription for constructing rank-3 Jordan algebras, due to Freudenthal, Springer and Tits [36–38], for which all the properties of the Jordan algebra are essentially determined by the cubic norm N (for a sketch of the construction see also [39]).

In the present investigation, we realize a rank-3 Jordan algebra $\mathfrak{J}^{\mathbb{B}}$ over the division or split algebra \mathbb{B} as the set of all 3×3 matrices J with entries in \mathbb{B} satisfying:

$$(2.9) \quad \eta J^\dagger \eta = J,$$

where $\eta = \text{diag}\{\epsilon, 1, 1\}$, with $\epsilon = 1$ for the *Euclidean* Jordan algebra $\mathfrak{J}_3^{\mathbb{B}}$, and $\epsilon = -1$ for the *Lorentzian* Jordan algebra³ $\mathfrak{J}_{1,2}^{\mathbb{B}}$ (see *e.g.* [24]), *i.e.* J is of the form:

$$(2.10) \quad J = \begin{pmatrix} a_1 & x_1 & x_2 \\ \epsilon \bar{x}_1 & a_2 & x_3 \\ \epsilon \bar{x}_2 & \bar{x}_3 & a_3 \end{pmatrix},$$

with $a_i \in \mathbb{R}$, and $x_i \in \mathbb{B}$, $i = 1, 2, 3$. Thus, out of the all the Jordan algebras from the classification in [23], we are restricting ourselves to the consideration of all the *simple* rank-3 Jordan algebras except for the *non-generic* case

³The following *Jordan algebraic isomorphism* holds:

$$\mathfrak{J}_{1,2}^{\mathbb{B}} \sim \mathfrak{J}_{2,1}^{\mathbb{B}},$$

and in general:

$$\mathfrak{J}_{M,N}^{\mathbb{B}} \sim \mathfrak{J}_{N,M}^{\mathbb{B}}.$$

of $\mathfrak{J} = \mathbb{R}$ itself⁴. The (*commutative*) Jordan product \circ (2.6)–(2.7) is realized as the *symmetrized* matrix multiplication:

$$(2.11) \quad j_1 \circ j_2 := \frac{1}{2}(j_1 j_2 + j_2 j_1), \quad j_1, j_2 \in \mathfrak{J}_3^{\mathbb{B}}.$$

It is then possible to introduce an inner product on the Jordan algebra:

$$(2.12) \quad \langle j_1, j_2 \rangle := Tr(j_1 \circ j_2).$$

As an example, for both the rank-3 Jordan algebras $\mathfrak{J}_3^{\mathbb{O}}$ and $\mathfrak{J}_3^{\mathbb{O}^s}$, the relevant vector space is the representation space **27** pertaining to the fundamental irrep. of $E_{6(-26)}$ resp. $E_{6(6)}$, and the cubic norm N is realized in terms of the completely symmetric invariant rank-3 tensor d_{IJK} in the **27** ($I, J, K = 1, \dots, 27$):

$$(2.13) \quad (\mathbf{27} \times \mathbf{27} \times \mathbf{27})_s \ni \exists! \mathbf{1} \equiv d_{IJK};$$

$$(2.14) \quad N(X) \equiv d_{IJK} X^I X^J X^K.$$

A detailed study of the rank-3 totally symmetric invariant d -tensor of Lorentzian rank-3 Jordan algebras can be found in [24].

The last important ingredient entering Eq. (2.1) is the *Lie product* $[\cdot, \cdot]$, which extends the multiplication structure also to $\mathbb{A}' \otimes \mathfrak{J}'^{\mathbb{B}}$, thus endowing $\mathcal{L}(\mathbb{A}, \mathbb{B})$ with the structure of a (Lie) algebra. Its general explicit expression can be found *e.g.* in Eq. (2.5) of [12]:

$$(2.15) \quad [h_1 \otimes j_1, h_2 \otimes j_2] := \frac{1}{12} \langle j_1, j_2 \rangle D_{h_1, h_2} - \langle h_1, h_2 \rangle [L_{j_1}, L_{j_2}] + \frac{1}{2} [h_1, h_2] \otimes (j_1 \circ j_2 - \frac{1}{3} \langle j_1, j_2 \rangle I_3).$$

Tits' formula (2.1) can be rewritten in a more symmetric way in \mathbb{A} and \mathbb{B} by generalizing the concept of derivations to that of *triality* (see *e.g.* [4, 11, 35]):

⁴The MS row which can be associated to $\mathfrak{J} = \mathbb{R}$ and to the *semi-simple* rank-3 Jordan algebras $\mathfrak{J} = \mathbb{R} \oplus \Gamma_{m,n}$ [23] is known (see *e.g.* Table 1 of [40], as well as Table 1 of [27]).

By their very definition, these algebras already have a signature, and, therefore, it would not make sense to treat them here.

$$(2.16) \quad \text{Tri}(\mathbb{A}) = \{(A, B, C) \text{ with } A, B, C \in \text{End}(\mathbb{A}) \\ | A(x_1x_2) = B(x_1)x_2 + x_1C(x_2)\}.$$

This leads to *Vinberg's formula* [4]:

$$(2.17) \quad \mathcal{L}(\mathbb{A}, \mathbb{B}) = \text{tri}(\mathbb{A}) \oplus \text{tri}(\mathbb{B}) \dot{+} 3\mathbb{A} \otimes \mathbb{B},$$

which implies:

$$(2.18) \quad \mathcal{L}(\mathbb{A}, \mathbb{B}) = \mathcal{L}(\mathbb{B}, \mathbb{A}),$$

a relation which will be useful in subsequent treatment.

A remarkable property of Jordan algebras is that they have various symmetry groups, which are relevant to supergravity theories and appear as entries in the MS's.

The *derivations algebra* $\text{Der}(\mathfrak{J}^{\mathbb{B}})$ generates the *automorphisms group* $\text{Aut}(\mathfrak{J}^{\mathbb{B}})$ of the Jordan algebra.

The *structure algebra* $\text{Str}(\mathbb{A})$, which for a general algebra \mathbb{A} is defined to be the Lie algebra generated by the left and right multiplication maps, in the case of a Jordan algebra can be expressed as [8]:

$$(2.19) \quad \text{Str}(\mathfrak{J}^{\mathbb{B}}) := \text{Der}(\mathfrak{J}^{\mathbb{B}}) \dot{+} L(\mathfrak{J}^{\mathbb{B}}) \text{ with } L(\mathfrak{J}^{\mathbb{B}}) := \{L_j | j \in \mathfrak{J}^{\mathbb{B}}\},$$

and its Lie algebra structure follows from $[D, L_j] = L_{Dj}$ for $D \in \text{Der}(\mathfrak{J}^{\mathbb{B}})$, $j \in \mathfrak{J}^{\mathbb{B}}$ and $[L_{j_1}, L_{j_2}] \in \text{Der}(\mathfrak{J}^{\mathbb{B}})$ for $j_1, j_2 \in \mathfrak{J}^{\mathbb{B}}$.

The *reduced structure algebra* $\text{Str}_0(\mathfrak{J}^{\mathbb{B}})$ is then defined as the quotient of $\text{Str}(\mathfrak{J}^{\mathbb{B}})$ by the subspace of multiples of L_1 , with 1 the identity of $\mathfrak{J}^{\mathbb{B}}$. It can be verified that $\text{Str}_0(\mathfrak{J}^{\mathbb{B}}) = \mathcal{L}(\mathbb{C}_S, \mathbb{B})$.

The *conformal algebra* $\text{Conf}(\mathfrak{J}^{\mathbb{B}})$ is the vector space [41, 42]:

$$(2.20) \quad \text{Conf}(\mathfrak{J}^{\mathbb{B}}) := \text{Str}(\mathfrak{J}^{\mathbb{B}}) \dot{+} 2\mathfrak{J}^{\mathbb{B}},$$

and its Lie algebra structure is defined by the brackets $[(x, 0), (y, 0)] = 0 = [(0, x), (0, y)]$ and $[(x, 0), (0, y)] = \frac{1}{2}(L_{xy} + [L_x, L_y])$ for $(x, y) \in \text{Conf}(\mathfrak{J}^{\mathbb{B}})$. It turns out that $\text{Conf}(\mathfrak{J}^{\mathbb{B}}) = \mathcal{L}(\mathbb{H}_S, \mathbb{B})$.

Finally, for the *quasi-conformal algebra* $\text{QConf}(\mathfrak{J}^{\mathbb{B}})$ [27, 41–43] (see also *e.g.* Sec. 3.5 of [44]), it can be seen that $\text{QConf}(\mathfrak{J}^{\mathbb{B}}) = \mathcal{L}(\mathbb{O}_S, \mathbb{B})$.

3. Magic squares \mathcal{L}_3 over rank-3 *Euclidean* Jordan algebras

By exploiting Tits' formula (2.1), we can now construct all possible MS's \mathcal{L}_3 based on rank-3 *Euclidean* Jordan algebras over the division algebras $\mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{O}, \mathbb{C}_S, \mathbb{H}_S$ and \mathbb{O}_S , by taking into account that $\mathbb{C} \subset \mathbb{H}, \mathbb{H}_S$ and $\mathbb{H} \subset \mathbb{O}, \mathbb{O}_S$, while $\mathbb{C}_S \subset \mathbb{H}_S$ and $\mathbb{H}_S \subset \mathbb{O}_S$. Thus, the possible sequences to be specified on the rows and columns of \mathcal{L}_3 are only four:

$$(3.1) \quad \begin{aligned} \mathbb{A} &= \mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{O}; \\ \widehat{\mathbb{A}} &= \mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{O}_S; \\ \widetilde{\mathbb{A}} &= \mathbb{R}, \mathbb{C}, \mathbb{H}_S, \mathbb{O}_S; \\ \mathbb{A}_S &= \mathbb{R}, \mathbb{C}_S, \mathbb{H}_S, \mathbb{O}_S, \end{aligned}$$

giving rise a priori to sixteen possible structures of Euclidean MS \mathcal{L}_3 .

However, by virtue of (2.17) and (2.18), it is enough to explicitly list only the magic squares for which the number of split division algebras labeling the rows is bigger or equal to that of the columns. This yields only ten different structures of Euclidean MS \mathcal{L}_3 , which we list and analyze below.

1. The *Freudenthal-Rozenfeld-Tits* (FRT) MS⁵ $\mathcal{L}_3(\mathbb{A}, \mathbb{B})$ [1–3]

	\mathbb{R}	\mathbb{C}	\mathbb{H}	\mathbb{O}
\mathbb{R}	$SO(3)$	$SU(3)$	$USp(6)$	$F_{4(-52)}$
\mathbb{C}	$SU(3)$	$SU(3) \times SU(3)$	$SU(6)$	$E_{6(-78)}$
\mathbb{H}	$USp(6)$	$SU(6)$	$SO(12)$	$E_{7(-133)}$
\mathbb{O}	$F_{4(-52)}$	$E_{6(-78)}$	$E_{7(-133)}$	$E_{8(-248)}$

Table 1: The *Freudenthal-Rozenfeld-Tits* (FRT) MS $\mathcal{L}_3(\mathbb{A}, \mathbb{B})$

This is a symmetric MS ($\mathcal{L}_3(\mathbb{A}, \mathbb{B}) = \mathcal{L}_3(\mathbb{A}, \mathbb{B})^T$), and it contains only compact (real) Lie algebras.

⁵The subscript in brackets denotes the *character* χ of the real form under consideration, namely the difference between the number of non-compact and compact generators [45]. Thus, in the case of *compact* real forms (as for all entries of FRT MS), the character is nothing but the opposite of the dimension of the algebra/group itself.

2. The *Günaydin-Sierra-Townsend* (GST) *single-split* MS $\mathcal{L}_3(\mathbb{A}_S, \mathbb{B})$ [13]

	\mathbb{R}	\mathbb{C}	\mathbb{H}	\mathbb{O}
\mathbb{R}	$SO(3)$	$SU(3)$	$USp(6)$	$F_{4(-52)}$
\mathbb{C}_S	$SL(3, \mathbb{R})$	$SL(3, \mathbb{C})$	$SU^*(6)$	$E_{6(-26)}$
\mathbb{H}_S	$Sp(6, \mathbb{R})$	$SU(3, 3)$	$SO^*(12)$	$E_{7(-25)}$
\mathbb{O}_S	$F_{4(4)}$	$E_{6(2)}$	$E_{7(-5)}$	$E_{8(-24)}$

Table 2: The *Günaydin-Sierra-Townsend* (GST) *single-split* MS $\mathcal{L}_3(\mathbb{A}_S, \mathbb{B})$

This is a non-symmetric MS ($\mathcal{L}_3(\mathbb{A}_S, \mathbb{B}) \neq \mathcal{L}_3(\mathbb{A}_S, \mathbb{B})^T$), and it displays symmetries relevant to (quarter-maximal) Maxwell-Einstein supergravity theories (MESGT's) with 8 supersymmetries, in various space-time signatures and dimensions.

The **fourth row** displays $QConf(\mathfrak{J}_3^{\mathbb{B}})$, the *quasi-conformal* symmetries of $\mathfrak{J}_3^{\mathbb{B}}$ [41, 42], which are the U -duality symmetries of $\mathcal{N} = 4$ *magic* theories in⁶ $D = (2, 1)$ (*i.e.* Lorentzian) space-time dimensions [13, 46], based on the *extended Freudenthal triple system* (EFTS) $\mathfrak{T}(\mathfrak{J}_3^{\mathbb{B}})$.

The **third row** displays $Conf(\mathfrak{J}_3^{\mathbb{B}})$, the *conformal* symmetries of $\mathfrak{J}_3^{\mathbb{B}}$ (2.20) [41, 42]:

- They are the U -duality symmetries of $\mathcal{N} = 2$, $D = (3, 1)$ *magic* MESGT's [13, 14] based on the *Freudenthal triple system* (FTS) $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{B}})$ [47].
- Up to a commuting Ehlers $SL(2, \mathbb{R})$ factor, they are the stabilizers of the extended scalar manifold of the $\mathfrak{T}(\mathfrak{J}_3^{\mathbb{B}})$ -based *magic* theories in $D = (3, 0)$ (*i.e.* Euclidean) space-time dimensions [48, 49].
- However, other (exotic) supergravity theories can be considered, obtained from suitable compactifications of theories in 11 dimensions alternative to the usual $D = (10, 1)$ M -theory, but still consistent with the existence of a *real* 32-dimensional spinor, namely M^* -theory in $D = (9, 2)$ and M' -theory in $D = (6, 5)$ [29]. By exploiting the analysis of [30], $Conf(\mathfrak{J}_3^{\mathbb{B}})$ (up to the Ehlers $SL(2, \mathbb{R})$) factor can also be regarded as the stabilizers of the the extended scalar manifold of the $\mathfrak{T}(\mathfrak{J}_3^{\mathbb{B}})$ -based *magic* theories in $D = (3, 0)_{M^*}$, $D = (3, 0)_{M'}$ and $D = (0, 3)_{M'}$ dimensions, where the subscript denotes the 11-dimensional origin throughout. For instance, for the theories based on $\mathfrak{T}(\mathfrak{J}_3^{\mathbb{H}})$,

⁶The first and second entries in the pair $D = (s, t)$ are to be read as the number of spacelike (s) and timelike (t) dimensions.

$\mathfrak{T}(\mathfrak{J}_3^{\mathbb{O}})$ and $\mathfrak{T}(\mathfrak{J}_3^{\mathbb{O}_S})$, the following embedding of symmetric cosets holds:

$$(3.2) \quad \frac{E_{7(-5)}}{SO^*(12) \times SL(2, \mathbb{R})} \Big|_{\mathfrak{T}(\mathfrak{J}_3^{\mathbb{H}}, H^*)} \subset \left[\frac{E_{8(-24)}}{E_{7(-25)} \times SL(2, \mathbb{R})} \Big|_{\mathfrak{T}(\mathfrak{J}_3^{\mathbb{O}}, H^*)} \cap \frac{E_{8(8)}}{SO^*(16)} \Big|_{\mathfrak{T}(\mathfrak{J}_3^{\mathbb{O}_S})} \right],$$

where “ H^* ” denotes the para-quaternionic structure of the corresponding spaces, which have vanishing character ($\chi = 0$; see *e.g.* [50] for a recent study of such manifolds).

The **second row** displays $Str_0(\mathfrak{J}_3^{\mathbb{B}})$, the *reduced structure* symmetries of $\mathfrak{J}_3^{\mathbb{B}}$ [8]:

- They are the U -duality symmetries of $\mathcal{N}=2, D=(4,1)$ *magic* MESGT’s [13, 14] based on $\mathfrak{J}_3^{\mathbb{B}}$.
- They are the stabilizers of the non-BPS $Z_H \neq 0$ “large” U -orbit of the corresponding MESGT in $D=(3,1)$ [31, 51].
- They are the stabilizers (up to a Kaluza-Klein $SO(1,1)$ commuting factor) of the scalar manifolds of $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{B}})$ -based $\mathcal{N}=2, magic$ MESGT’s in $D=(4,0)$.
- Considering more exotic theories, they are the stabilizers (up to a Kaluza-Klein $SO(1,1)$ commuting factor) of the scalar manifolds of $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{B}})$ -based $\mathcal{N}=2, magic$ MESGT’s in $D=(4,0)_{M^*}, D=(4,0)_{M'}$ and $D=(0,4)_{M'}$ dimensions. For instance, for the theories based on $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{H}}), \mathfrak{M}(\mathfrak{J}_3^{\mathbb{O}})$ and $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{O}_S})$, the following embedding of symmetric cosets holds:

$$(3.3) \quad \frac{SO^*(12)}{SU^*(6) \times SO(1,1)} \Big|_{\mathfrak{M}(\mathfrak{J}_3^{\mathbb{H}}, K^*)} \subset \left[\frac{E_{7(-25)}}{E_{6(-26)} \times SO(1,1)} \Big|_{\mathfrak{M}(\mathfrak{J}_3^{\mathbb{O}}, K^*)} \cap \frac{E_{7(7)}}{SU^*(8)} \Big|_{\mathfrak{M}(\mathfrak{J}_3^{\mathbb{O}_S})} \right],$$

where “ K^* ” denotes the (special) pseudo-Kähler structure of the corresponding spaces, which also have vanishing character ($\chi = 0$).

The **first row** displays $Aut(\mathfrak{J}_3^{\mathbb{B}}) = mcs(Str_0(\mathfrak{J}_3^{\mathbb{B}}))$, namely the *automorphisms* of $\mathfrak{J}_3^{\mathbb{B}}$:

- They are the stabilizers of the scalar manifolds of $\mathcal{N} = 2$, $D = (4, 1)$ *magic* MESGTs [13, 14] based on $\mathfrak{J}_3^{\mathbb{B}}$.
- They are stabilizers of the (1/2-)BPS “large” U -orbit in the same theory [52, 53].
- Considering more exotic theories, $Aut(\mathfrak{J}_3^{\mathbb{B}})$ can also be regarded as the stabilizers of the scalar manifolds of the same $\mathfrak{J}_3^{\mathbb{B}}$ -based theory in $D = (0, 5)_{M'}$ dimensions.

3. The *Barton-Sudbery* (BS) *double-split* MS $\mathcal{L}_3(\mathbb{A}_S, \mathbb{B}_S)$ [8], which also appeared more recently in [27]

	\mathbb{R}	\mathbb{C}_S	\mathbb{H}_S	\mathbb{O}_S
\mathbb{R}	$SO(3)$	$SL(3, \mathbb{R})$	$Sp(6, \mathbb{R})$	$F_{4(4)}$
\mathbb{C}_S	$SL(3, \mathbb{R})$	$SL(3, \mathbb{R}) \times SL(3, \mathbb{R})$	$SL(6, \mathbb{R})$	$E_{6(6)}$
\mathbb{H}_S	$Sp(6, \mathbb{R})$	$SL(6, \mathbb{R})$	$SO(6, 6)$	$E_{7(7)}$
\mathbb{O}_S	$F_{4(4)}$	$E_{6(6)}$	$E_{7(7)}$	$E_{8(8)}$

Table 3: The *Barton-Sudbery* (BS) *double-split* MS $\mathcal{L}_3(\mathbb{A}_S, \mathbb{B}_S)$

This is a symmetric MS ($\mathcal{L}_3(\mathbb{A}_S, \mathbb{B}_S) = \mathcal{L}_3(\mathbb{A}_S, \mathbb{B}_S)^T$), and it displays symmetries relevant to Maxwell-Einstein theories of gravity with 8 (*quarter-maximal*, $\mathbb{B}_S = \mathbb{R}$) or 32 (*maximal*, $\mathbb{B}_S = \mathbb{O}_S$) supersymmetries, or *without* ($\mathbb{B}_S = \mathbb{C}_S, \mathbb{H}_S$) any supersymmetry at all (see e.g. [27]).

The **fourth row** displays $QConf(\mathfrak{J}_3^{\mathbb{B}_S})$, the *quasi-conformal* symmetries of $\mathfrak{J}_3^{\mathbb{B}_S}$, which are the U -duality symmetries of ME(S)GT’s in $D = (2, 1)$ dimensions, based on the EFTS $\mathfrak{T}(\mathfrak{J}_3^{\mathbb{B}_S})$ [41, 42].

The **third row** displays $Conf(\mathfrak{J}_3^{\mathbb{B}_S})$, the *conformal* symmetries of $\mathfrak{J}_3^{\mathbb{B}_S}$ [41, 42]:

- They are the U -duality symmetries of $D = (3, 1)$ ME(S)GT’s based on the FTS $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{B}_S})$ [47].
- By extending the analysis of [30] to non-maximally supersymmetric theories of gravity, they also are (up to a commuting Ehlers $SL(2, \mathbb{R})$ factor) the stabilizers of the extended scalar manifold of the $\mathfrak{T}(\mathfrak{J}_3^{\mathbb{B}_S})$ -based magic theories in $D = (2, 1)_{M^*}$, $D = (1, 2)_{M^*}$, $D = (2, 1)_{M'}$, and

$D = (1, 2)_{M'}$ dimensions. This holds with the exclusion of the case $\mathbb{B}_S = \mathbb{O}_S$, in which maximal supersymmetry constrains the stabilizer to match the \mathcal{R} -symmetry, namely $SO(8, 8)$. For instance, for the theories based on $\mathfrak{T}(\mathfrak{J}_3^{\mathbb{H}_S})$ and $\mathfrak{T}(\mathfrak{J}_3^{\mathbb{O}_S})$, the following embedding of symmetric cosets holds:

$$(3.4) \quad \frac{E_{7(7)}}{SO(6, 6) \times SL(2, \mathbb{R})} \subset \left[\frac{E_{8(8)}}{SO(8, 8)} \cap \frac{E_{8(8)}}{E_{7(7)} \times SL(2, \mathbb{R})} \right],$$

$\mathfrak{T}(\mathfrak{J}_3^{\mathbb{H}_S}), H^*$ $\mathfrak{T}(\mathfrak{J}_3^{\mathbb{O}_S})$ H^*

where the para-quaternionic spaces also have vanishing character ($\chi = 0$). Note that for the $\mathfrak{T}(\mathfrak{J}_3^{\mathbb{O}_S})$ -based theory, $\frac{E_{8(8)}}{SO(8, 8)}$ is the enlarged scalar manifold, whereas $\frac{E_{8(8)}}{E_{7(7)} \times SL(2, \mathbb{R})}$ can be regarded as a particular, non-compact pseudo-Riemannian version of the rank-4 quaternionic symmetric manifold $\frac{E_{8(-24)}}{E_{7(-25)} \times SU(2)}$, the c -map [54] of the rank-3 special Kähler space $\frac{E_{7(-25)}}{E_{6(-78)} \times U(1)}$ (scalar manifold of the $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{O}})$ -based MESGT in $D = (3, 1)$ [13, 14]; for a recent treatment, see *e.g.* [21]).

The **second row** displays $Str_0(\mathfrak{J}_3^{\mathbb{B}_S})$, the *reduced structure* symmetries of $\mathfrak{J}_3^{\mathbb{B}_S}$ [8]:

- They are the U -duality symmetries of $D = (4, 1)$ ME(S)GT's based on $\mathfrak{J}_3^{\mathbb{B}_S}$.
- They are the stabilizers of a certain “large” U -orbit of the corresponding ME(S)GT in $D = (3, 1)$ (which, in presence of local supersymmetry, is the non-BPS one [31, 51]).
- They are the stabilizers (up to a Kaluza-Klein $SO(1, 1)$ commuting factor) of the scalar manifolds of $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{B}})$ -based ME(S)GT's in $D = (2, 2)_{M^*}$ and $D = (2, 2)_{M'}$ dimensions. This holds with the exclusion of the case $\mathbb{B}_S = \mathbb{O}_S$, in which maximal supersymmetry constrains the stabilizer to match the \mathcal{R} -symmetry, namely $SL(8, \mathbb{R})$. For instance, for the theories based on $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{H}_S})$ and $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{O}_S})$, the following embedding of symmetric cosets holds:

$$(3.5) \quad \frac{SO(6, 6)}{SL(6, \mathbb{R}) \times SO(1, 1)} \subset \left[\frac{E_{7(7)}}{SL(8, \mathbb{R})} \cap \frac{E_{7(7)}}{E_{6(6)} \times SO(1, 1)} \right],$$

$\mathfrak{M}(\mathfrak{J}_3^{\mathbb{H}_S}), K^*$ $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{O}_S})$ K^*

where the (special) pseudo-Kähler spaces also have vanishing character ($\chi = 0$). Note that for the $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{O}_S})$ -based theory, $\frac{E_{7(7)}}{SL(8, \mathbb{R})}$ is the scalar manifold, whereas $\frac{E_{7(7)}}{E_{6(6)} \times SO(1,1)}$ can be regarded as a particular, non-compact pseudo-Riemannian version of the rank-3 special Kähler symmetric manifold $\frac{E_{7(-25)}}{E_{6(-78)} \times SU(2)}$, the R -map [26, 55] of the rank-2 real special space $\frac{E_{6(-26)}}{F_{4(-52)}}$ (scalar manifold of the $\mathfrak{J}_3^{\mathbb{O}}$ -based MESGT in $D = (4, 1)$ [13, 14]).

The **first row** displays $Aut(\mathfrak{J}_3^{\mathbb{B}_S})$, namely the *automorphisms* of $\mathfrak{J}_3^{\mathbb{B}_S}$, which can be regarded as the stabilizers of the scalar manifolds of $\mathfrak{J}_3^{\mathbb{B}}$ -based ME(S)GTs in $D = (3, 2)_{M^*}$, $D = (3, 2)_{M'}$ and $D = (2, 3)_{M'}$ dimensions. This holds with the exclusion of the case $\mathbb{B}_S = \mathbb{O}_S$, in which maximal supersymmetry constrains the stabilizer to match the \mathcal{R} -symmetry, namely $Sp(8, \mathbb{R})$. For instance, for the theories based on $\mathfrak{J}_3^{\mathbb{H}_S}$ and $\mathfrak{J}_3^{\mathbb{O}_S}$, the following embedding of symmetric cosets holds:

$$(3.6) \quad \frac{SL(6, \mathbb{R})}{Sp(6, \mathbb{R})_{\mathfrak{J}_3^{\mathbb{H}_S}}} \subset \left[\frac{E_{6(6)}}{SL(8, \mathbb{R})_{\mathfrak{J}_3^{\mathbb{O}_S}}} \cap \frac{E_{6(6)}}{F_{4(4)}} \right].$$

Note that for the $\mathfrak{J}_3^{\mathbb{O}_S}$ -based theory, $\frac{E_{6(6)}}{SL(8, \mathbb{R})}$ is the scalar manifold, whereas $\frac{E_{6(6)}}{F_{4(4)}}$ can be regarded as a particular, non-compact pseudo-Riemannian version of the rank-2 real special symmetric manifold $\frac{E_{6(-26)}}{F_{4(-52)}}$ (scalar manifold of the $\mathfrak{J}_3^{\mathbb{O}}$ -based MESGT in $D = (4, 1)$ [13, 14]). Moreover, $\frac{E_{6(6)}}{F_{4(4)}}$ can be regarded as the “large” $\frac{1}{8}$ -BPS U -orbit of the $\mathfrak{J}_3^{\mathbb{O}_S}$ -based maximal supergravity theory in $D = (4, 1)$ [31, 32, 56].

4. The first “mixed” MS $\mathcal{L}_3(\tilde{\mathbb{A}}, \mathbb{B})$ [21] This is a non-symmetric MS

	\mathbb{R}	\mathbb{C}	\mathbb{H}	\mathbb{O}
\mathbb{R}	$SO(3)$	$SU(3)$	$USp(6)$	$F_{4(-52)}$
\mathbb{C}	$SU(3)$	$SU(3) \times SU(3)$	$SU(6)$	$E_{6(-78)}$
\mathbb{H}_S	$Sp(6, \mathbb{R})$	$SU(3, 3)$	$SO^*(12)$	$E_{7(-25)}$
\mathbb{O}_S	$F_{4(4)}$	$E_{6(2)}$	$E_{7(-5)}$	$E_{8(-24)}$

Table 4: The first “mixed” MS $\mathcal{L}_3(\tilde{\mathbb{A}}, \mathbb{B})$ [21]

$(\mathcal{L}_3(\tilde{\mathbb{A}}, \mathbb{B}) \neq \mathcal{L}_3(\tilde{\mathbb{A}}, \mathbb{B})^T)$. It displays symmetries relevant to the construction of maximally manifestly covariant parametrizations (as well as Iwasawa decompositions) of the scalar manifolds of $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{B}})$ -based MESGT's in $D = (3, 1)$ (the case $\mathbb{B} = \mathbb{O}$ has been studied in detail in [21]).

5. – 10. All the other Euclidean MS's \mathcal{L}_3 can be computed (as to our knowledge, they never appeared in the literature), and we report them in Tables 5 – 10.

	\mathbb{R}	\mathbb{C}	\mathbb{H}	\mathbb{O}
\mathbb{R}	$SO(3)$	$SU(3)$	$USp(6)$	$F_{4(-52)}$
\mathbb{C}	$SU(3)$	$SU(3) \times SU(3)$	$SU(6)$	$E_{6(-78)}$
\mathbb{H}	$USp(6)$	$SU(6)$	$SO(12)$	$E_{7(-133)}$
\mathbb{O}_S	$F_{4(4)}$	$E_{6(2)}$	$E_{7(-5)}$	$E_{8(-24)}$

Table 5: MS $\mathcal{L}_3(\hat{\mathbb{A}}, \mathbb{B})$

	\mathbb{R}	\mathbb{C}	\mathbb{H}	\mathbb{O}_S
\mathbb{R}	$SO(3)$	$SU(3)$	$USp(6)$	$F_{4(4)}$
\mathbb{C}	$SU(3)$	$SU(3) \times SU(3)$	$SU(6)$	$E_{6(2)}$
\mathbb{H}	$USp(6)$	$SU(6)$	$SO(12)$	$E_{7(-5)}$
\mathbb{O}_S	$F_{4(4)}$	$E_{6(2)}$	$E_{7(-5)}$	$E_{8(8)}$

Table 6: MS $\mathcal{L}_3(\hat{\mathbb{A}}, \hat{\mathbb{B}})$

	\mathbb{R}	\mathbb{C}	\mathbb{H}	\mathbb{O}_S
\mathbb{R}	$SO(3)$	$SU(3)$	$USp(6)$	$F_{4(4)}$
\mathbb{C}	$SU(3)$	$SU(3) \times SU(3)$	$SU(6)$	$E_{6(2)}$
\mathbb{H}_S	$Sp(6, \mathbb{R})$	$SU(3, 3)$	$SO^*(12)$	$E_{7(7)}$
\mathbb{O}_S	$F_{4(4)}$	$E_{6(2)}$	$E_{7(-5)}$	$E_{8(8)}$

Table 7: MS $\mathcal{L}_3(\tilde{\mathbb{A}}, \hat{\mathbb{B}})$

It can be noticed that $\mathcal{L}_3(\hat{\mathbb{A}}, \hat{\mathbb{B}})$, given by Table 6, and $\mathcal{L}_3(\tilde{\mathbb{A}}, \tilde{\mathbb{B}})$, given by Table 9, are symmetric, while all the other ones are non-symmetric. By suitably generalizing the approach of [21] to non-compact spaces, these

	\mathbb{R}	\mathbb{C}	\mathbb{H}	\mathbb{O}_S
\mathbb{R}	$SO(3)$	$SU(3)$	$USp(6)$	$F_{4(4)}$
\mathbb{C}_S	$SL(3, \mathbb{R})$	$SL(3, \mathbb{C})$	$SU^*(6)$	$E_{6(6)}$
\mathbb{H}_S	$Sp(6, \mathbb{R})$	$SU(3, 3)$	$SO^*(12)$	$E_{7(7)}$
\mathbb{O}_S	$F_{4(4)}$	$E_{6(2)}$	$E_{7(-5)}$	$E_{8(8)}$

Table 8: MS $\mathcal{L}_3(\mathbb{A}_S, \widehat{\mathbb{B}})$

	\mathbb{R}	\mathbb{C}	\mathbb{H}_S	\mathbb{O}_S
\mathbb{R}	$SO(3)$	$SU(3)$	$Sp(6, \mathbb{R})$	$F_{4(4)}$
\mathbb{C}	$SU(3)$	$SU(3) \times SU(3)$	$SU(3, 3)$	$E_{6(2)}$
\mathbb{H}_S	$Sp(6, \mathbb{R})$	$SU(3, 3)$	$SO(6, 6)$	$E_{7(7)}$
\mathbb{O}_S	$F_{4(4)}$	$E_{6(2)}$	$E_{7(7)}$	$E_{8(8)}$

Table 9: MS $\mathcal{L}_3(\widetilde{\mathbb{A}}, \widetilde{\mathbb{B}})$

	\mathbb{R}	\mathbb{C}	\mathbb{H}_S	\mathbb{O}_S
\mathbb{R}	$SO(3)$	$SU(3)$	$Sp(6, \mathbb{R})$	$F_{4(4)}$
\mathbb{C}_S	$SL(3, \mathbb{R})$	$SL(3, \mathbb{C})$	$SL(6, \mathbb{R})$	$E_{6(6)}$
\mathbb{H}_S	$Sp(6, \mathbb{R})$	$SU(3, 3)$	$SO(6, 6)$	$E_{7(7)}$
\mathbb{O}_S	$F_{4(4)}$	$E_{6(2)}$	$E_{7(7)}$	$E_{8(8)}$

Table 10: MS $\mathcal{L}_3(\mathbb{A}_S, \widetilde{\mathbb{B}})$

MS's may be used to explicitly construct pseudo-Riemannian scalar manifolds of theories of Maxwell-Einstein (super)gravity in non-Lorentzian spacetimes, also obtained from compactifications of M^* -theory or M' -theory. For instance, the symmetric MS $\mathcal{L}_3(\widetilde{\mathbb{A}}, \widetilde{\mathbb{B}})$ can be used to determine a (maximally) manifestly $(E_{6(2)} \times U(1))$ -covariant construction of the rank-3 pseudo-Riemannian special Kähler manifold $\frac{E_{7(7)}}{E_{6(2)} \times U(1)}$, which is a non-compact version of the aforementioned Riemannian special Kähler symmetric coset $\frac{E_{7(-25)}}{E_{6(-78)} \times U(1)}$ (scalar manifold of the $\mathfrak{N}(\mathfrak{J}_3^0)$ -based MESGT in $D = (3, 1)$ [13, 14]).

4. Magic squares $\mathcal{L}_{1,2}$ over rank-3 Lorentzian Jordan algebras

We will now exploit Tits' formula (2.1) in order to construct all possible MS's $\mathcal{L}_{1,2}$ based on rank-3 Lorentzian Jordan algebras over the division algebras $\mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{O}, \mathbb{C}_S, \mathbb{H}_S$ and \mathbb{O}_S . As discussed at the start of Sec. 3, by virtue of (2.17) and (2.18), it is enough to explicitly list only the magic squares for which the number of split division algebras labeling the rows is bigger or equal to that of the columns.

We would like to point out that, as to our knowledge, these MS's never appeared in literature. Interestingly, their study has been motivated also by the investigation of the stabilizers of the class of "large" non-BPS $Z = 0$ U -orbits in magic MESGT's in $D = (3, 1)$ dimensions [51], which indeed provide the third row of $\mathcal{L}_{1,2}(\mathbb{A}, \mathbb{B})$, the Lorentzian counterpart of the FRT MS $\mathcal{L}_3(\mathbb{A}, \mathbb{B})$ [1–3] given in Table 1.

Moreover, it should be remarked that the two non-compact real forms $F_{4(-20)}$ and $E_{6(-14)}$, which do not occur Euclidean MS's \mathcal{L}_3 , can instead be obtained from Tits' formula (2.1) or the Vinberg's formula (2.17) by considering Lorentzian MS's $\mathcal{L}_{1,2}$. It holds that [10, 28]:

$$(4.1) \quad \mathfrak{f}_{4(-20)} = \text{Der} \left(\mathfrak{J}_{1,2}^{\mathbb{O}} \right) \\ = \text{Der}(\mathbb{O}) \oplus \text{Der} \left(\mathfrak{J}_{1,2}^{\mathbb{R}} \right) \dot{+} \left(\mathbb{O}' \otimes \mathfrak{J}'^{\mathbb{R}}_{1,2} \right);$$

$$(4.2) \quad \mathfrak{e}_{6(-14)} = \text{Der} \left(\mathfrak{J}_{1,2}^{\mathbb{O}} \right) \dot{+} \left(e_4 \otimes \mathfrak{J}'^{\mathbb{O}}_{1,2} \right) \\ = \text{Der}(\mathbb{O}) \oplus \text{Der} \left(\mathfrak{J}_{1,2}^{\mathbb{C}} \right) \dot{+} \left(\mathbb{O}' \otimes \mathfrak{J}'^{\mathbb{C}}_{1,2} \right).$$

The ten possible different structures of Lorentzian MS $\mathcal{L}_{1,2}$ are listed and analyzed below.

1. The Lorentzian FRT MS $\mathcal{L}_{1,2}(\mathbb{A}, \mathbb{B})$

	\mathbb{R}	\mathbb{C}	\mathbb{H}	\mathbb{O}
\mathbb{R}	$SL(2, \mathbb{R})$	$SU(2, 1)$	$USp(4, 2)$	$F_{4(-20)}$
\mathbb{C}	$SU(2, 1)$	$SU(2, 1) \times SU(2, 1)$	$SU(4, 2)$	$E_{6(-14)}$
\mathbb{H}	$USp(4, 2)$	$SU(4, 2)$	$SO(8, 4)$	$E_{7(-5)}$
\mathbb{O}	$F_{4(-20)}$	$E_{6(-14)}$	$E_{7(-5)}$	$E_{8(8)}$

Table 11: The Lorentzian FRT MS $\mathcal{L}_{1,2}(\mathbb{A}, \mathbb{B})$

This is a symmetric MS ($\mathcal{L}_{1,2}(\mathbb{A}, \mathbb{B}) = \mathcal{L}_{1,2}(\mathbb{A}, \mathbb{B})^T$), and it contains only non-compact (real) Lie algebras.

As mentioned above, the **first row** displays:

- the stabilizer of the “large” non-BPS U -orbit (with $Z_H \neq 0$) of the $\mathfrak{J}_3^{\mathbb{B}}$ -based magic MESGT in $D = (4, 1)$ dimensions [52, 53].
- the stabilizer of the scalar manifold of the same theory in $D = (5, 0)$ [48, 49].
- Considering more exotic theories, the stabilizer of the scalar manifold of the same theory in $D = (4, 1)_{M^*}$, $D = (5, 0)_{M^*}$, $D = (4, 1)_{M'}$, $D = (1, 4)_{M'}$ and $D = (5, 0)_{M'}$ dimensions. For instance, for the theories based on $\mathfrak{J}_3^{\mathbb{H}}$, $\mathfrak{J}_3^{\mathbb{O}}$ and $\mathfrak{J}_3^{\mathbb{O}_S}$, the following embedding of symmetric cosets holds:

$$(4.3) \quad \frac{SU^*(6)}{USp(4, 2)_{\mathfrak{J}_3^{\mathbb{H}}}} \subset \left[\frac{E_{6(-26)}}{F_{4(-20)}_{\mathfrak{J}_3^{\mathbb{O}}}} \cap \frac{E_{6(6)}}{USp(4, 4)_{\mathfrak{J}_3^{\mathbb{O}_S}}} \right].$$

The **second row** displays:

- the stabilizer of the “large” non-BPS U -orbit (with $Z_H = 0$) of the $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{B}})$ -based magic MESGT’s in $D = (3, 1)$ dimensions [31, 51].
- Considering more exotic theories, the stabilizer (up to a commuting $U(1)$ factor) of the scalar manifold of the same theory in $D = (3, 1)_{M^*}$, $D = (3, 1)_{M'}$ and $D = (1, 3)_{M'}$ dimensions. For instance, for the theories based on $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{H}})$, $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{O}})$ and $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{O}_S})$, the following embedding of symmetric cosets holds:

$$(4.4) \quad \frac{SO^*(12)}{SU(4, 2) \times U(1)_{\mathfrak{J}_3^{\mathbb{H}}, K}} \subset \left[\frac{E_{7(-25)}}{E_{6(-14)} \times U(1)_{\mathfrak{J}_3^{\mathbb{O}}, K}} \cap \frac{E_{7(7)}}{SU(4, 4)_{\mathfrak{J}_3^{\mathbb{O}_S}}} \right],$$

where “ K ” denotes the (special) Kähler structure of the corresponding spaces. Note that $\frac{SO^*(12)}{SU(4, 2) \times U(1)}$ and $\frac{E_{7(-25)}}{E_{6(-14)} \times U(1)}$ are particular pseudo-Riemannian non-compact forms of the rank-3 special Kähler Riemannian symmetric cosets $\frac{SO^*(12)}{U(6)}$ and $\frac{E_{7(-25)}}{E_{6(-78)} \times U(1)}$ (scalar manifolds of the $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{H}})$ - and $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{O}})$ - based magic MESGT’s in $D = (3, 1)$ dimensions).

The **third row** displays the stabilizer (up to $SU(2)$ factor) of the scalar manifold of the $\mathfrak{T}(\mathfrak{J}_3^{\mathbb{B}})$ -based magic theories in $D = (2, 1)_{M^*}$, $D = (1, 2)_{M^*}$, $D = (2, 1)_{M'}$ and $D = (1, 2)_{M'}$ dimensions. For instance, for the theories based on $\mathfrak{T}(\mathfrak{J}_3^{\mathbb{H}})$, $\mathfrak{T}(\mathfrak{J}_3^{\mathbb{O}})$ and $\mathfrak{T}(\mathfrak{J}_3^{\mathbb{O}_S})$, the following embedding of symmetric cosets holds:

$$(4.5) \quad \frac{E_{7(-5)}}{SO(8,4) \times SU(2)} \Big|_{\mathfrak{T}_3^{\mathbb{H}}, H} \subset \left[\frac{E_{8(-24)}}{E_{7(-5)} \times SU(2)} \Big|_{\mathfrak{T}_3^{\mathbb{O}}, H} \cap \frac{E_{8(8)}}{SO(8,8)} \Big|_{\mathfrak{T}_3^{\mathbb{O}_S}} \right],$$

where “ H ” denotes the quaternionic structure of the corresponding spaces. Note that $\frac{E_{7(-5)}}{SO(8,4) \times SU(2)}$ and $\frac{E_{8(-24)}}{E_{7(-5)} \times SU(2)}$ are particular pseudo-Riemannian non-compact forms of the rank-4 quaternionic Riemannian symmetric cosets $\frac{E_{7(-5)}}{SO(12) \times SU(2)}$ and $\frac{E_{8(-24)}}{E_{7(-133)} \times SU(2)}$ (extended scalar manifolds of the $\mathfrak{T}(\mathfrak{J}_3^{\mathbb{H}})$ - and $\mathfrak{T}(\mathfrak{J}_3^{\mathbb{O}})$ - based magic theories in $D = (2, 1)$ dimensions).

Finally, the **fourth row** can be characterized as displaying the non-compact real forms which (besides $QConf(\mathfrak{J}_3^{\mathbb{B}})$; *cfr.* the fourth row of the GST MS $\mathcal{L}_3(\mathbb{A}_S, \mathbb{B})$ in Table 2) embed maximally (by an $SU(2)$ factor) the non-compact real forms in the third row.

2. The Lorentzian GST single-split MS $\mathcal{L}_{1,2}(\mathbb{A}_S, \mathbb{B})$

	\mathbb{R}	\mathbb{C}	\mathbb{H}	\mathbb{O}
\mathbb{R}	$SL(2, \mathbb{R})$	$SU(2, 1)$	$USp(4, 2)$	$F_{4(-20)}$
\mathbb{C}_S	$SL(3, \mathbb{R})$	$SL(3, \mathbb{C})$	$SU^*(6)$	$E_{6(-26)}$
\mathbb{H}_S	$Sp(6, \mathbb{R})$	$SU(3, 3)$	$SO^*(12)$	$E_{7(-25)}$
\mathbb{O}_S	$F_{4(4)}$	$E_{6(2)}$	$E_{7(-5)}$	$E_{8(-24)}$

Table 12: Lorentzian GST MS $\mathcal{L}_{1,2}(\mathbb{A}_S, \mathbb{B})$

This is a non-symmetric MS ($\mathcal{L}_{1,2}(\mathbb{A}_S, \mathbb{B}) \neq \mathcal{L}_{1,2}(\mathbb{A}_S, \mathbb{B})^T$).

The **second**, **third** and **fourth rows** match the corresponding rows of its Euclidean counterpart, namely of the GST MS $\mathcal{L}_3(\mathbb{A}_S, \mathbb{B})$ given in Table 2.

On the other hand, the **first row** coincides with the first row of the Lorentzian FRT MS $\mathcal{L}_{1,2}(\mathbb{A}, \mathbb{B})$ given in Table 11.

3. The Lorentzian BS *double-split* MS $\mathcal{L}_{1,2}(\mathbb{A}_S, \mathbb{B}_S)$

	\mathbb{R}	\mathbb{C}_S	\mathbb{H}_S	\mathbb{O}_S
\mathbb{R}	$SL(2, \mathbb{R})$	$SL(3, \mathbb{R})$	$Sp(6, \mathbb{R})$	$F_{4(4)}$
\mathbb{C}_S	$SL(3, \mathbb{R})$	$SL(3, \mathbb{R}) \times SL(3, \mathbb{R})$	$SL(6, \mathbb{R})$	$E_{6(6)}$
\mathbb{H}_S	$Sp(6, \mathbb{R})$	$SL(6, \mathbb{R})$	$SO(6, 6)$	$E_{7(7)}$
\mathbb{O}_S	$F_{4(4)}$	$E_{6(6)}$	$E_{7(7)}$	$E_{8(8)}$

Table 13: Lorentzian BS MS $\mathcal{L}_{1,2}(\mathbb{A}_S, \mathbb{B}_S)$

This is a symmetric MS ($\mathcal{L}_{1,2}(\mathbb{A}_S, \mathbb{B}_S) = \mathcal{L}_{1,2}(\mathbb{A}_S, \mathbb{B}_S)^T$). It matches its Euclidean counterpart, namely the BS *double-split* MS $\mathcal{L}_3(\mathbb{A}_S, \mathbb{B}_S)$ given in Table 3, up to the first entry (from the left) in the first row, which reads:

$$(4.6) \quad SL(2, \mathbb{R}) = \mathcal{L}_{1,2}(\mathbb{R}, \mathbb{R}) \neq \mathcal{L}_3(\mathbb{R}, \mathbb{R}) = SO(3).$$

4. The Lorentzian counterpart $\mathcal{L}_{1,2}(\tilde{\mathbb{A}}, \mathbb{B})$ of the first “mixed” MS $\mathcal{L}_3(\tilde{\mathbb{A}}, \mathbb{B})$ reads:

	\mathbb{R}	\mathbb{C}	\mathbb{H}	\mathbb{O}
\mathbb{R}	$SL(2, \mathbb{R})$	$SU(2, 1)$	$USp(4, 2)$	$F_{4(-20)}$
\mathbb{C}	$SU(2, 1)$	$SU(2, 1) \times SU(2, 1)$	$SU(4, 2)$	$E_{6(-14)}$
\mathbb{H}_S	$Sp(6, \mathbb{R})$	$SU(3, 3)$	$SO^*(12)$	$E_{7(-25)}$
\mathbb{O}_S	$F_{4(4)}$	$E_{6(2)}$	$E_{7(-5)}$	$E_{8(-24)}$

Table 14: The Lorentzian first “mixed” MS $\mathcal{L}_{1,2}(\tilde{\mathbb{A}}, \mathbb{B})$

This is a non-symmetric MS ($\mathcal{L}_{1,2}(\tilde{\mathbb{A}}, \mathbb{B}) \neq \mathcal{L}_{1,2}(\tilde{\mathbb{A}}, \mathbb{B})^T$). Its **third** and **fourth rows** coincide with those of its Euclidean counterpart, namely of first “mixed” MS $\mathcal{L}_3(\tilde{\mathbb{A}}, \mathbb{B})$, given in Table 4. On the other hand, its **first** and **second rows** match those of the Lorentzian FRT MS $\mathcal{L}_{1,2}(\mathbb{A}, \mathbb{B})$, given in Table 11.

5. – 10. All the other Lorentzian MS’s $\mathcal{L}_{1,2}$ can be computed, and we report them in Tables 15 – 20. It can be noticed that $\mathcal{L}_{1,2}(\tilde{\mathbb{A}}, \widehat{\mathbb{B}})$, given by Table 16, and $\mathcal{L}_{1,2}(\tilde{\mathbb{A}}, \widetilde{\mathbb{B}})$, given by Table 19, are symmetric, while all the other ones are non-symmetric. By suitably generalizing the approach of [21] to non-compact spaces, also these MS’s may be used to explicitly construct pseudo-Riemannian scalar manifolds of theories of Maxwell-Einstein

(super)gravity in non-Lorentzian space-times, also obtained from compactifications of M^* -theory or M' -theory.

	\mathbb{R}	\mathbb{C}	\mathbb{H}	\mathbb{O}
\mathbb{R}	$SL(2, \mathbb{R})$	$SU(2, 1)$	$USp(4, 2)$	$F_{4(-20)}$
\mathbb{C}	$SU(2, 1)$	$SU(2, 1) \times SU(2, 1)$	$SU(4, 2)$	$E_{6(-14)}$
\mathbb{H}	$USp(4, 2)$	$SU(4, 2)$	$SO(8, 4)$	$E_{7(-5)}$
\mathbb{O}_S	$F_{4(4)}$	$E_{6(2)}$	$E_{7(-5)}$	$E_{8(-24)}$

Table 15: Lorentzian MS $\mathcal{L}_{1,2}(\widehat{\mathbb{A}}, \mathbb{B})$

	\mathbb{R}	\mathbb{C}	\mathbb{H}	\mathbb{O}_S
\mathbb{R}	$SL(2, \mathbb{R})$	$SU(2, 1)$	$USp(4, 2)$	$F_{4(4)}$
\mathbb{C}	$SU(2, 1)$	$SU(2, 1) \times SU(2, 1)$	$SU(4, 2)$	$E_{6(2)}$
\mathbb{H}	$USp(4, 2)$	$SU(4, 2)$	$SO(8, 4)$	$E_{7(-5)}$
\mathbb{O}_S	$F_{4(4)}$	$E_{6(2)}$	$E_{7(-5)}$	$E_{8(8)}$

Table 16: Lorentzian MS $\mathcal{L}_{1,2}(\widehat{\mathbb{A}}, \widehat{\mathbb{B}})$

	\mathbb{R}	\mathbb{C}	\mathbb{H}	\mathbb{O}_S
\mathbb{R}	$SL(2, \mathbb{R})$	$SU(2, 1)$	$USp(4, 2)$	$F_{4(4)}$
\mathbb{C}	$SU(2, 1)$	$SU(2, 1) \times SU(2, 1)$	$SU(4, 2)$	$E_{6(2)}$
\mathbb{H}_S	$Sp(6, \mathbb{R})$	$SU(3, 3)$	$SO^*(12)$	$E_{7(7)}$
\mathbb{O}_S	$F_{4(4)}$	$E_{6(2)}$	$E_{7(-5)}$	$E_{8(8)}$

Table 17: Lorentzian MS $\mathcal{L}_{1,2}(\widetilde{\mathbb{A}}, \widehat{\mathbb{B}})$

5. Analysis

Below we list some observations on common properties, as well as on differences, among the two sets of 4×4 MS's over rank-3 *Euclidean* (Tables 1 – 10) and *Lorentzian* (Tables 11 – 20) rank-3 (*simple*, generic) Jordan algebras.

- 1) For $\mathcal{L}_3(\mathbb{A}, \mathbb{B})$ and $\mathcal{L}_{1,2}(\mathbb{A}, \mathbb{B})$ (namely for the FRT MS - Table 1 - and its Lorentzian analogue - Table 11 -), the symmetries in the second row/column are embedded into the symmetries in the third one with a factor $U(1)$ or $SO(2)$, while the symmetries in the third row/column

	\mathbb{R}	\mathbb{C}	\mathbb{H}	\mathbb{O}_S
\mathbb{R}	$SL(2, \mathbb{R})$	$SU(2, 1)$	$USp(4, 2)$	$F_{4(4)}$
\mathbb{C}_S	$SL(3, \mathbb{R})$	$SL(3, \mathbb{C})$	$SU^*(6)$	$E_{6(6)}$
\mathbb{H}_S	$Sp(6, \mathbb{R})$	$SU(3, 3)$	$SO^*(12)$	$E_{7(7)}$
\mathbb{O}_S	$F_{4(4)}$	$E_{6(2)}$	$E_{7(-5)}$	$E_{8(8)}$

Table 18: Lorentzian MS $\mathcal{L}_{1,2}(\mathbb{A}_S, \widehat{\mathbb{B}})$

	\mathbb{R}	\mathbb{C}	\mathbb{H}_S	\mathbb{O}_S
\mathbb{R}	$SL(2, \mathbb{R})$	$SU(2, 1)$	$Sp(6, \mathbb{R})$	$F_{4(4)}$
\mathbb{C}	$SU(2, 1)$	$SU(2, 1) \times SU(2, 1)$	$SU(3, 3)$	$E_{6(2)}$
\mathbb{H}_S	$Sp(6, \mathbb{R})$	$SU(3, 3)$	$SO(6, 6)$	$E_{7(7)}$
\mathbb{O}_S	$F_{4(4)}$	$E_{6(2)}$	$E_{7(7)}$	$E_{8(8)}$

Table 19: Lorentzian MS $\mathcal{L}_{1,2}(\widetilde{\mathbb{A}}, \widetilde{\mathbb{B}})$

	\mathbb{R}	\mathbb{C}	\mathbb{H}_S	\mathbb{O}_S
\mathbb{R}	$SL(2, \mathbb{R})$	$SU(2, 1)$	$Sp(6, \mathbb{R})$	$F_{4(4)}$
\mathbb{C}_S	$SL(3, \mathbb{R})$	$SL(3, \mathbb{C})$	$SL(6, \mathbb{R})$	$E_{6(6)}$
\mathbb{H}_S	$Sp(6, \mathbb{R})$	$SU(3, 3)$	$SO(6, 6)$	$E_{7(7)}$
\mathbb{O}_S	$F_{4(4)}$	$E_{6(2)}$	$E_{7(7)}$	$E_{8(8)}$

Table 20: Lorentzian MS $\mathcal{L}_{1,2}(\mathbb{A}_S, \widetilde{\mathbb{B}})$

are embedded into the symmetries in the fourth one with a factor $SO(3)$ or $SU(2)$. Examples of such maximal and symmetric embeddings from $\mathcal{L}_{1,2}(\mathbb{A}, \mathbb{B})$ read

$$(5.1) \quad \begin{aligned} E_{7(-5)} &\supset E_{6(-14)} \times U(1); \\ E_{6(-14)} &\supset SU(4, 2) \times SU(2). \end{aligned}$$

Analogously, for $\mathcal{L}_3(\mathbb{A}_S, \mathbb{B})$ and $\mathcal{L}_{1,2}(\mathbb{A}_S, \mathbb{B})$ (namely for the *single-split* GST MS - Table 2 - and its Lorentzian analogue - Table 12 -), the symmetries in the second column (row) are embedded into the symmetries

in the third column (row) with a factor $U(1)$ ($SO(1, 1)$), whereas the symmetries in the third column (row) are embedded into the symmetries in the fourth column (row) with a factor $SU(2)$ ($SU(1, 1)$). And, similarly, for $\mathcal{L}_3(\mathbb{A}_S, \mathbb{B}_S)$ and $\mathcal{L}_{1,2}(\mathbb{A}_S, \mathbb{B}_S)$ (namely for the *double-split* BS MS - Table 3 - and its Lorentzian analogue - Table 13 -), the symmetries in the second row/column are embedded into the symmetries in the third one with a factor $SO(1, 1)$, while the symmetries in the third row/column are embedded into the symmetries in the fourth one with a factor $SU(1, 1)$. Analogous results holds for all other Euclidean (Tables 4 – 10) and Lorentzian (Tables 11 – 20) MS's. The *rationale* of all this is the following. When the embedding of H into G in the next row/ column of the MS contains an extra factor $T = U(1)$, $SO(1, 1)$, $SU(2)$ or $SU(1, 1)$, this reflects the structure of the symmetric coset $\frac{G}{H \times T}$, which then carries a complex (special Kähler), (special) pseudo-Kähler, quaternionic or para-quaternionic structure, respectively.

- 2) When all the aforementioned commuting factors are taken into account, all the embeddings in the MS's are maximal and symmetric [45].
- 3) From Tits' formula (2.1), it can be realized that the factor $SO(2)$ or $SO(1, 1)$, needed to maximally embed the symmetries in the second row into the symmetries in the third one, is in turn embedded respectively into $\text{Aut}(\mathbb{H}) = SO(3)$ or $\text{Aut}(\mathbb{H}_S) = SL(2, \mathbb{R})$; on the other hand, the factor $SU(2)$ or $SU(1, 1)$, needed to maximally embed the symmetries in the third row into the symmetries in the fourth one, is in turn embedded respectively into $\text{Aut}(\mathbb{O}) = G_{2(-14)}$ or $\text{Aut}(\mathbb{O}_S) = G_{2(2)}$. The relevant (maximal and symmetric) embeddings read:

$$\begin{aligned}
 &G_{2(-14)} \supset SU(2) \times SU(2); \\
 &G_{2(2)} \supset SU(1, 1) \times SU(1, 1); \\
 (5.2) \quad &SU(2) \supset SO(2); \\
 &SU(1, 1) \supset SO(1, 1).
 \end{aligned}$$

Analogous considerations can be made for the embeddings of the columns. The factor $U(1)$ or $SO(1, 1)$, needed to maximally embed the symmetries in the second column into the symmetries is in turn embedded respectively into $\text{Aut}(\mathfrak{J}_{1,2}^{\mathbb{H}}) = USp(4, 2)$ or $\text{Aut}(\mathfrak{J}_{1,2}^{\mathbb{H}_S}) = Sp(6, \mathbb{R})$; on the other hand, the factor $SU(2)$ or $SU(1, 1)$, needed to maximally embed the symmetries in the third column into the symmetries in the

fourth one, is in turn embedded respectively into $\text{Aut}(\mathfrak{J}_{1,2}^{\mathbb{O}}) = F_{4(-20)}$ or $\text{Aut}(\mathfrak{J}_{1,2}^{\mathbb{O}_S}) = F_{4(4)}$. The relevant (maximal and symmetric) embeddings read:

$$\begin{aligned}
 & F_{4(-20)} \supset USp(4, 2) \times SU(2); \\
 & F_{4(4)} \supset Sp(6, \mathbb{R}) \times SU(1, 1); \\
 (5.3) \quad & USp(4, 2) \supset SU(2, 1) \times U(1); \\
 & Sp(6, \mathbb{R}) \supset SL(3, \mathbb{R}) \times SO(1, 1).
 \end{aligned}$$

Therefore, for each of the embeddings of a row/column in the next, these generators always have the same origin.

- 4) The symmetries of Euclidean and Lorentzian rank-3 Jordan algebras over *division* algebras can be read from the rows of the corresponding single-split MS, namely from the GST MS $\mathcal{L}_3(\mathbb{A}_S, \mathbb{B})$ (Table 2) and from its Lorentzian counterpart, i.e. the MS $\mathcal{L}_{1,2}(\mathbb{A}_S, \mathbb{B})$ (Table 12). For Euclidean rank-3 Jordan algebras, it holds:

	Row 1: Automorphism	$\text{Aut}(\mathfrak{J}_3^{\mathbb{B}}) = \mathcal{L}_3(\mathbb{R}, \mathbb{B});$
(5.4)	Row 2: Reduced Structure	$\text{Str}_0(\mathfrak{J}_3^{\mathbb{B}}) = \mathcal{L}_3(\mathbb{C}_S, \mathbb{B});$
	Row 3: Conformal	$\text{Conf}(\mathfrak{J}_3^{\mathbb{B}}) = \mathcal{L}_3(\mathbb{H}_S, \mathbb{B});$
	Row 4: QuasiConformal	$\text{QConf}(\mathfrak{J}_3^{\mathbb{B}}) = \mathcal{L}_3(\mathbb{O}_S, \mathbb{B}).$

Since the second, third and fourth rows of $\mathcal{L}_3(\mathbb{A}_S, \mathbb{B})$ and $\mathcal{L}_{1,2}(\mathbb{A}_S, \mathbb{B})$ match, this implies that the reduced structure, conformal and quasi-conformal symmetries of Euclidean and Lorentzian rank-3 Jordan algebras over *division* algebras coincide:

$$\begin{aligned}
 & \text{Str}_0(\mathfrak{J}_{1,2}^{\mathbb{A}}) = \text{Str}_0(\mathfrak{J}_3^{\mathbb{A}}); \\
 (5.5) \quad & \text{Conf}(\mathfrak{J}_{1,2}^{\mathbb{A}}) = \text{Conf}(\mathfrak{J}_3^{\mathbb{A}}); \\
 & \text{QConf}(\mathfrak{J}_{1,2}^{\mathbb{A}}) = \text{QConf}(\mathfrak{J}_3^{\mathbb{A}}),
 \end{aligned}$$

whereas their automorphisms differ:

$$(5.6) \quad \text{Aut}(\mathfrak{J}_3^{\mathbb{B}}) = \mathcal{L}_3(\mathbb{R}, \mathbb{B}) \neq \mathcal{L}_{1,2}(\mathbb{R}, \mathbb{B}) = \text{Aut}(\mathfrak{J}_{1,2}^{\mathbb{B}}).$$

This is consistent with the analysis of [24, 25].

5) Analogously, the symmetries of Euclidean and Lorentzian rank-3 Jordan algebras $\mathfrak{J}_3^{\mathbb{B}_S}$ over *split* algebras can be read from the rows of the corresponding double-split MS, namely from the BS MS $\mathcal{L}_3(\mathbb{A}_S, \mathbb{B}_S)$ (Table 3) and from its Lorentzian counterpart, i.e. the MS $\mathcal{L}_{1,2}(\mathbb{A}_S, \mathbb{B}_S)$ (Table 13). For Euclidean rank-3 Jordan algebras, it holds:

$$\begin{aligned}
 \text{Row 1: Automorphism} & \quad \text{Aut} \left(\mathfrak{J}_3^{\mathbb{B}_S} \right) = \mathcal{L}_3 \left(\mathbb{R}, \mathbb{B}_S \right); \\
 \text{Row 2: Reduced Structure} & \quad \text{Str}_0 \left(\mathfrak{J}_3^{\mathbb{B}_S} \right) = \mathcal{L}_3 \left(\mathbb{C}_S, \mathbb{B}_S \right); \\
 \text{Row 3: Conformal} & \quad \text{Conf} \left(\mathfrak{J}_3^{\mathbb{B}_S} \right) = \mathcal{L}_3 \left(\mathbb{H}_S, \mathbb{B}_S \right); \\
 \text{Row 4: QuasiConformal} & \quad \text{QConf} \left(\mathfrak{J}_3^{\mathbb{B}_S} \right) = \mathcal{L}_3 \left(\mathbb{O}_S, \mathbb{B}_S \right).
 \end{aligned}
 \tag{5.7}$$

Since the second, third and fourth rows of $\mathcal{L}_3(\mathbb{A}_S, \mathbb{B}_S)$ and $\mathcal{L}_{1,2}(\mathbb{A}_S, \mathbb{B}_S)$ match, this implies that the reduced structure, conformal and quasi-conformal symmetries of Euclidean and Lorentzian rank-3 Jordan algebras over *split* algebras coincide:

$$\begin{aligned}
 \text{Str}_0 \left(\mathfrak{J}_{1,2}^{\mathbb{A}_S} \right) &= \text{Str}_0 \left(\mathfrak{J}_3^{\mathbb{A}_S} \right); \\
 \text{Conf} \left(\mathfrak{J}_{1,2}^{\mathbb{A}_S} \right) &= \text{Conf} \left(\mathfrak{J}_3^{\mathbb{A}_S} \right); \\
 \text{QConf} \left(\mathfrak{J}_{1,2}^{\mathbb{A}_S} \right) &= \text{QConf} \left(\mathfrak{J}_3^{\mathbb{A}_S} \right).
 \end{aligned}
 \tag{5.8}$$

On the other hand, since the first rows of $\mathcal{L}_3(\mathbb{A}_S, \mathbb{B}_S)$ and $\mathcal{L}_{1,2}(\mathbb{A}_S, \mathbb{B}_S)$ match (with the exception of the first entry from the left), it also follows that their automorphisms coincide:

$$\begin{aligned}
 \text{Aut} \left(\mathfrak{J}_3^{\mathbb{B}_S} \right) &= \mathcal{L}_3 \left(\mathbb{R}, \mathbb{B}_S \right) = \mathcal{L}_{1,2} \left(\mathbb{R}, \mathbb{B}_S \right) = \text{Aut} \left(\mathfrak{J}_{1,2}^{\mathbb{B}_S} \right), \\
 \mathbb{B}_S &= \mathbb{C}_S, \mathbb{H}_S, \mathbb{O}_S,
 \end{aligned}
 \tag{5.9}$$

whereas Eq. (4.6) can be interpreted as follows:

$$\begin{aligned}
 SL(2, \mathbb{R}) &= \text{Aut} \left(\mathfrak{J}_{1,2}^{\mathbb{R}} \right) = \mathcal{L}_{1,2}(\mathbb{R}, \mathbb{R}) \\
 &\neq \mathcal{L}_3(\mathbb{R}, \mathbb{R}) = \text{Aut} \left(\mathfrak{J}_3^{\mathbb{R}} \right) = SO(3).
 \end{aligned}
 \tag{5.10}$$

6) The complexification of the Jordan algebras $\mathfrak{J}_3^{\mathbb{A}}$ and $\mathfrak{J}_{1,2}^{\mathbb{A}}$ by means of a *Cayley-Dickson procedure* should in principle allow to recover all Euclidean and Lorentzian magic squares given in Tables 1 – 20, as

suitable sections of only two magic squares over the bi-octonions [10, 44].

- 7) In our treatment, we never mentioned unified MESGT's based on $\mathfrak{J}_{1,2}^{\mathbb{A}}$ (in $D = (4, 1)$) and on $\mathfrak{M}(\mathfrak{J}_{1,2}^{\mathbb{A}})$ (in $D = (3, 1)$), which are endowed with a *non-homogeneous* scalar manifold \mathcal{M} [24–26]. However, it respectively holds [24, 25]

$$(5.11) \quad D = (4, 1) : \mathcal{M}(\mathfrak{J}_{1,2}^{\mathbb{A}}) \subset \frac{Str_0(\mathfrak{J}_{1,2}^{\mathbb{A}})}{Aut(\mathfrak{J}_{1,2}^{\mathbb{A}})} = \frac{Str_0(\mathfrak{J}_3^{\mathbb{A}})}{Aut(\mathfrak{J}_{1,2}^{\mathbb{A}})};$$

$$(5.12) \quad D = (3, 1) : \mathcal{M}(\mathfrak{M}(\mathfrak{J}_{1,2}^{\mathbb{A}})) \subset \frac{Conf(\mathfrak{J}_{1,2}^{\mathbb{A}})}{K(\mathfrak{J}_{1,2}^{\mathbb{A}})} = \frac{Conf(\mathfrak{J}_3^{\mathbb{A}})}{K(\mathfrak{J}_{1,2}^{\mathbb{A}})}.$$

$\frac{Str_0(\mathfrak{J}_{1,2}^{\mathbb{A}})}{Aut(\mathfrak{J}_{1,2}^{\mathbb{A}})}$ (5.11) can also be regarded as the scalar manifold of the $\mathfrak{J}_3^{\mathbb{A}}$ -based magic MESGT in $D = (5, 0)$ dimensions, as well as in $D = (4, 1)_{M^*}$, $D = (5, 0)_{M^*}$, $D = (4, 1)_{M'}$, $D = (1, 4)_{M'}$ and $D = (5, 0)_{M'}$ dimensions (see Sec. 4). Moreover, $\frac{Str_0(\mathfrak{J}_{1,2}^{\mathbb{A}})}{Aut(\mathfrak{J}_{1,2}^{\mathbb{A}})}$ can be identified also with the “large” non-BPS U -orbit (with $Z_H \neq 0$) of the $\mathfrak{J}_3^{\mathbb{A}}$ -based magic MESGT in $D = (4, 1)$ dimensions [52, 53]. On the other hand, $\frac{Conf(\mathfrak{J}_{1,2}^{\mathbb{A}})}{K(\mathfrak{J}_{1,2}^{\mathbb{A}})}$ (5.12), whose stabilizer is given (up to a $U(1)$ factor) by the second row of the Lorentzian FRT MS $\mathcal{L}_{1,2}(\mathbb{A}, \mathbb{B})$ (Table 11), is the *Koecher upper half plane* of $\mathfrak{J}_{1,2}^{\mathbb{A}}$ [25], which can be identified also with the “large” non-BPS U -orbit (with $Z_H = 0$) of the $\mathfrak{M}(\mathfrak{J}_3^{\mathbb{B}})$ -based magic MESGT's in $D = (3, 1)$ dimensions [31, 51]. Moreover, by adding an additional $U(1)$ factor in the stabilizer, $\frac{Conf(\mathfrak{J}_{1,2}^{\mathbb{A}})}{K(\mathfrak{J}_{1,2}^{\mathbb{A}}) \times U(1)}$ can also be regarded as the scalar manifold of the $\mathfrak{J}_3^{\mathbb{A}}$ -based magic MESGT in $D = (3, 1)_{M^*}$, $D = (3, 1)_{M'}$ and $D = (1, 3)_{M'}$ dimensions (see Sec. 4).

Acknowledgments

The work of B.L.C. has been supported in part by the European Commission under the FP7-PEOPLE-IRG-2008 Grant No. PIRG04-GA-2008-239412 “*String Theory and Noncommutative Geometry*” (STRING).

References

- [1] H. Freudenthal, *Lie groups in the foundations of geometry*. Adv. Math., **1**:145, (1963).
- [2] J. Tits, *Algèbres alternatives, algèbres de Jordan et algèbres de Lie exceptionnelles. I. Construction* (French). Nederl. Akad. Wetensch. Proc. Ser., **A 69**:223, (1966).
- [3] B. A. Rozenfeld, *Geometrical interpretation of the compact simple Lie groups of the class E* (in Russian). Dokl. Akad. Nauk. SSSR, **106**:600, (1956).
- [4] A. L. Onishchik and E. B. Vinberg, *Lie Groups and Lie Algebras III*. Springer, Berlin, 1991.
- [5] R. C. Arcuri, *Conformal and Critical Embeddings, Infinite Magic Square and a new Clifford Product*. J. Math. Phys., **32**:1890, (1991).
- [6] M. Santander and F. J. Herranz, “Cayley-Klein” Schemes for Real Lie Algebras and Freudenthal Magic Squares. [arXiv:physics/9702031](https://arxiv.org/abs/physics/9702031).
- [7] J. M. Landsberg and L. Manivel, *The Projective Geometry of Freudenthal’s Magic Square*. J. of Algebra, **239**:477, (2001), [arXiv:math/9908039](https://arxiv.org/abs/math/9908039).
- [8] C. H. Barton and A. Sudbery, *Magic Squares of Lie Algebras*. [arXiv:math/0001083](https://arxiv.org/abs/math/0001083). C. H. Barton and A. Sudbery, *Magic squares and matrix models of Lie algebras*. Adv. in Math., **180**:596, (2003), [arxiv:math/0203010](https://arxiv.org/abs/math/0203010) [math.RA].
- [9] I. Cunha and A. Elduque, *The Extended Freudenthal Magic Square and Jordan Algebras*. Manuscripta Math. **123**:325, (2007), [arXiv:math/0608191](https://arxiv.org/abs/math/0608191).
- [10] I. Yokota, *Exceptional Lie Groups*. [arXiv:0902.0431](https://arxiv.org/abs/0902.0431) [math.DG].
- [11] J. M. Evans, *Trialities and Exceptional Lie Algebras: Deconstructing the Magic Square*. [arXiv:0910.1828](https://arxiv.org/abs/0910.1828) [hep-th].
- [12] S. L. Cacciatori, F. D. Piazza and A. Scotti, *E_7 groups from octonionic magic square*. Adv. Theor. Math. Phys. **15**:1605, (2011), [arXiv:1007.4758](https://arxiv.org/abs/1007.4758) [math-ph].
- [13] M. Günaydin, G. Sierra and P. K. Townsend, *Exceptional Supergravity Theories and the Magic Square*. Phys. Lett., **B133B**:72, (1983).

- [14] M. Günaydin, G. Sierra and P. K. Townsend, *The Geometry of $\mathcal{N}=2$ Maxwell-Einstein Supergravity and Jordan Algebras*. Nucl. Phys. **B242**: 244 (1984).
- [15] E. Cremmer and B. Julia, *The $\mathcal{N}=8$ Supergravity Theory. 1. The Lagrangian*. Phys. Lett., **B80**:48 (1978). E. Cremmer and B. Julia, *The $SO(8)$ Supergravity*. Nucl. Phys., **B159**:141, (1979).
- [16] C. Hull and P. K. Townsend, *Unity of Superstring Dualities*. Nucl. Phys., **B438**:109, (1995), [arXiv:hep-th/9410167](#).
- [17] P. Truini, G. Olivieri and L. C. Biedenharn, *The Jordan Pair Content of the Magic Square and the Geometry of the Scalars in $\mathcal{N}=2$ Supergravity*. Lett. Math. Phys., **9**:255, (1985).
- [18] K. Dasgupta, V. Hussin and A. Wissanji, *Quaternionic Kähler Manifolds, Constrained Instantons and the Magic Square I*. Nucl. Phys., **B793**:34, (2008), [arXiv:0708.1023 \[hep-th\]](#).
- [19] J. Gutt, *Special Riemannian Geometries and the Magic Square of Lie Algebras*. [arXiv:0810.2138](#).
- [20] S. Ferrara and A. Marrani, *Symmetric Spaces in Supergravity* Proceedings of the Conference “Symmetry in Mathematics and Physics: Celebrating V.S. Varadarajan’s 70th Birthday”, [arXiv:0808.3567 \[hep-th\]](#); L. J. Boya, *Symmetric spaces of exceptional groups*. Phys. Atom. Nucl., **73**:209, (2010), [arXiv:0811.0554 \[math-ph\]](#).
- [21] S. L. Cacciatori, B. L. Cerchiai and A. Marrani, *Magic Coset Decompositions*. Adv. Theor. Math. Phys. **17**: 1077, (2013), [arXiv:1201.6314 \[hep-th\]](#).
- [22] P. Jordan, *Über die multiplikation quanten-mechanischer grossen*. Zschr. f. Phys., **80**:285, (1933); P. Jordan, *Über verallgemeinerungsmöglichkeiten des formalismus der quantenmechanik*. Nachr. Ges. Wiss. Gottingen, (1933) 209–214; N. Jacobson, *Some groups of transformations defined by Jordan algebras*. J. Reine Angew. Math., **207**, (1961) 61–85; N. Jacobson, “*Structure and Representations of Jordan Algebras*”, Vol. 39, American Mathematical Society Colloquium Publications, 1968.
- [23] P. Jordan, J. von Neumann and E. P. Wigner, *On an Algebraic generalization of the quantum mechanical formalism*. Annals Math., **35**:29, (1934).

- [24] M. Günaydin and M. Zagermann, *Unified Maxwell-Einstein and Yang-Mills-Einstein Supergravity Theories in Five Dimensions*. JHEP, **0307**:023, (2003), [arXiv:hep-th/0304109](#).
- [25] M. Günaydin, S. McReynolds and M. Zagermann, *Unified $\mathcal{N}=2$ Maxwell-Einstein and Yang-Mills-Einstein supergravity theories in four dimensions*. JHEP, **0509**:026, (2005), [arXiv:hep-th/0507227](#).
- [26] M. Günaydin, S. McReynolds and M. Zagermann, *The R-map and the Coupling of $N=2$ Tensor Multiplets in 5 and 4 Dimensions*. JHEP, **0601**:168, (2006), [arXiv:hep-th/0511025](#).
- [27] M. Günaydin and O. Pavlyk, *Quasiconformal Realizations of $E_{6(6)}$, $E_{7(7)}$, $E_{8(8)}$ and $SO(n+3, m+3)$, $\mathcal{N} \geq 4$ Supergravity and Spherical Vectors* [arXiv:0904.0784 \[hep-th\]](#).
- [28] A. Elduque, *A new look at Freudenthal's Magic Square*. In: "Non Associative Algebra and Its Applications" (L. Sabinin, L. Sbitneva and I.P. Shestakov, eds.), Lect. Notes in Pure and Applied Mathematics **246**, (2006), 149–165.
- [29] C. Hull, *Duality and the Signature of Space-Time*. JHEP, **9807**:021, (1998), [arXiv:hep-th/9807127](#).
- [30] S. Ferrara, *Spinors, superalgebras and the signature of space-time*. [arXiv:hep-th/0101123](#).
- [31] S. Ferrara and M. Günaydin, *Orbits of Exceptional Groups, Duality and BPS States in String Theory*. Int. J. Mod. Phys., **A13**:2075, (1998), [hep-th/9708025](#).
- [32] H. Lu, C. N. Pope and K. S. Stelle, *Multiplet structures of BPS solitons*. Class. Quant. Grav., **15**:537, (1998), [arXiv:hep-th/9708109](#).
- [33] A. Marrani, *Charge Orbits and Moduli Spaces of Black Hole Attractors*. Lect. Notes Math., **2027**:155, (2011), [arXiv:1012.3559 \[hep-th\]](#).
- [34] S. Bellucci, S. Ferrara, R. Kallosh and A. Marrani, *Extremal Black Hole and Flux Vacua Attractors*. Lect. Notes Phys., **755**:115, (2008), [arXiv:0711.4547](#).
- [35] J. C. Baez, *The Octonions*. Bull. Am. Math. Soc., **39**:145, (2002), [arXiv:math/0105155 \[math-ra\]](#).
- [36] T. A. Springer, *Characterization of a class of cubic forms*. Nederl. Akad. Wetensch. Proc. Ser., **A24**, (1962), 259–265.

- [37] K. McCrimmon, *The Freudenthal-Springer-Tits construction of exceptional Jordan algebras*. Trans. Amer. Math. Soc., **139**, (1969), 495–510.
- [38] K. McCrimmon, *A Taste of Jordan Algebras*. Springer-Verlag New York Inc., New York, 2004.
- [39] L. Borsten, D. Dahanayake, M. J. Duff and W. Rubens, *Black Holes Admitting a Freudenthal Dual*. Phys. Rev., **D80**:026003, (2009), [arXiv:0903.5517 \[hep-th\]](#).
- [40] B. Pioline, *Lectures on Black Holes, Topological Strings and Quantum Attractors*. Class. Quant. Grav., **23**:S981, (2006), [arXiv:hep-th/0607227](#).
- [41] M. Günaydin, *Generalized conformal and superconformal group actions and Jordan algebras*. Mod. Phys. Lett., **A8**:1407, (1993), [arXiv:hep-th/9301050](#); M. Günaydin, K. Koepsell and H. Nicolai, *Conformal and quasiconformal realizations of exceptional Lie groups*. Commun. Math. Phys., **221**:57, (2001), [arXiv:hep-th/0008063](#).
- [42] M. Günaydin and O. Pavlyk, *Spectrum Generating Conformal and Quasiconformal U-Duality Groups, Supergravity and Spherical Vectors*. JHEP, **1004**:070, (2010), [arXiv:0901.1646 \[hep-th\]](#).
- [43] M. Günaydin, *Unitary Realizations of U-duality Groups as Conformal and Quasiconformal Groups and Extremal Black Holes of Supergravity Theories*. AIP Conf. Proc., **767**:268, (2005), [arXiv:hep-th/0502235](#).
- [44] M. Rios, *Jordan C*-Algebras and Supergravity*. Preprint, [arXiv:1005.3514 \[hep-th\]](#).
- [45] R. Gilmore, *Lie Groups, Lie Algebras, and Some of Their Applications*. Dover, New York, 2006.
- [46] B. de Wit, A.K. Tollsten and H. Nicolai, *Locally supersymmetric D = 3 nonlinear sigma models*. Nucl. Phys., **B392**:3, (1993), [arXiv:hep-th/9208074](#).
- [47] R. B. Brown, *Groups of type E₇*. Jour. Reine Angew. Math., **236**:79, (1969); J. R. Faulkner, *A construction of Lie algebras from a class of ternary algebras*. Trans. Amer. Math. Soc., **155**:397, (1971); K. Meyberg, *Eine Theorie der Freudenthalschen Tripelsysteme, I, II*. Ned. Akad. Wetenschap., **71**:162, (1968).
- [48] P. Breitenlohner, D. Maison and G. Gibbons, *4-Dimensional black holes from Kaluza-Klein theories*. Commun. Math. Phys., **120**:295, (1988).

- [49] E. Bergshoeff, W. Chemissany, A. Ploegh, M. Trigiante and T. Van Riet, *Generating Geodesic Flows and Supergravity Solutions*. Nucl. Phys., **B812**:343, (2009), [arXiv:0806.2310 \[hep-th\]](#).
- [50] S. Ferrara, A. Marrani and M. Trigiante, *Super-Ehlers in Any Dimension*. JHEP, **1211**:068, (2012), [arXiv:1206.1255 \[hep-th\]](#).
- [51] S. Bellucci, S. Ferrara, M. Günaydin and A. Marrani, *Charge Orbits of Symmetric Special Geometries and Attractors*. Int. J. Mod. Phys., **A21**:5043, (2006), [arXiv:hep-th/0606209](#).
- [52] S. Ferrara and M. Günaydin, *Orbits and Attractors for $\mathcal{N}=2$ Maxwell-Einstein Supergravity Theories in Five Dimensions*. Nucl. Phys., **B759**:1, (2006), [arXiv:hep-th/0606108](#).
- [53] B. L. Cerchiai, S. Ferrara, A. Marrani and B. Zumino, *Charge Orbits of Extremal Black Holes in Five Dimensional Supergravity*. Phys. Rev., **D82**:085010, (2010), [arXiv:1006.3101 \[hep-th\]](#).
- [54] S. Cecotti, S. Ferrara and L. Girardello, *Geometry of type II superstrings and the moduli of superconformal field theories*. Int. J. Mod. Phys., **A4**:2475, (1989).
- [55] B. de Wit, F. Vanderseypen and A. Van Proeyen, *Symmetry Structure of Special Geometries* Nucl. Phys. **B400**:463 (1993), [arXiv:hep-th/9210068](#).
- [56] L. Borsten, D. Dahanayake, M. J. Duff, S. Ferrara, A. Marrani and W. Rubens, *Observations on Integral and Continuous U-duality Orbits in $\mathcal{N}=8$ Supergravity*. Class. Quant. Grav., **27**:185003, (2010), [arXiv:1002.4223 \[hep-th\]](#).

DIPARTIMENTO DI SCIENZE ED ALTA TECNOLOGIA
UNIVERSITÀ DEGLI STUDI DELL'INSUBRIA
VIA VALLEGGIO 11, 22100 COMO, ITALY
AND INFN, SEZIONE DI MILANO, VIA CELORIA 16, 20133 MILANO, ITALY
E-mail address: sergio.cacciatori@uninsubria.it

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI MILANO
VIA SALDINI 50, 20133 MILANO, ITALY
AND INFN, SEZIONE DI MILANO, VIA CELORIA 16, 20133 MILANO, ITALY
E-mail address: bianca.letizia.cerchiai@cern.ch

PHYSICS DEPARTMENT, THEORY UNIT, CERN
CH 1211, GENEVA 23, SWITZERLAND
E-mail address: alessio.marrani@cern.ch