

On topological approach to local theory of surfaces in Calabi-Yau threefolds

SERGEI GUKOV, CHIU-CHU MELISSA LIU,
ARTAN SHESHMANI, AND SHING-TUNG YAU

We study the web of dualities relating various enumerative invariants, notably Gromov-Witten invariants and invariants that arise in topological gauge theory. In particular, we study Donaldson-Thomas gauge theory and its reductions to $D = 4$ and $D = 2$ which are relevant to the local theory of surfaces in Calabi-Yau threefolds.

1	Introduction	1679
2	Preliminaries	1681
3	Dimensional reduction of DT theory to 4d Vafa-Witten theory	1692
4	Dimensional reduction of DT theory to 2d σ-model	1707
5	Dimensional reduction of torsion DT theory	1715
	References	1720

1. Introduction

Not so long ago, enumerative invariants were completely unfamiliar to physicists and were regarded more as a curiosity rather than serious mathematics in the math world. At present, enumerative invariants are heavily used in theoretical physics and play a major role in many branches of pure mathematics: symplectic geometry, algebraic geometry, mirror symmetry, and gauge theory.

In gauge theory, the prominent examples of enumerative invariants are Donaldson polynomials and Seiberg-Witten invariants, which help to distinguish different smooth structures on 4-manifolds. In recent years, other 4-manifold invariants have been introduced by changing the gauge theory (the PDE's and the "counting" problem) or, by changing dimension, similar gauge theory invariants were defined on higher-dimensional manifolds. Notable examples include Donaldson-Thomas (DT) invariants which, roughly speaking, can be viewed as six-dimensional analogues of Donaldson invariants on 4-manifolds [25].

In symplectic geometry and mirror symmetry, enumerative invariants play an equally important role; it is almost impossible to imagine these fields without enumerative invariants. Most of the original versions of such invariants were based on the problem of curve "counting" of some form. More recently, however, this was generalized to a much larger framework where the role of curves is less special and, in the same time, the borderline between gauge theory and symplectic/algebraic geometry started to blur. For example, Donaldson-Thomas invariants that we mentioned earlier and that originally were introduced via gauge theory, nowadays are actually more widely regarded as part of algebraic geometry and mirror symmetry, since they are defined as the intersection numbers over the moduli space of sheaves (say for example the ideal sheaves of one dimensional subschemes of the ambient variety).

String theory also stimulated the process of erasing borders. It led to many dualities which relate enumerative invariants in different dimensions and even across the fields. For example, gauge theory itself appears in string theory through *open* strings, whereas curve counting (Gromov-Witten theory) is naturally a part of the *closed* string theory. Therefore, string dualities which relate open and closed strings (also known as gauge/gravity dualities) provide a fundamental, conceptual reason why one might hope to have a relation between gauge theory enumerative invariants and, say, Gromov-Witten invariants.

In this paper, we explore a rich web of inter-relations and dualities between different enumerative invariants.

Acknowledgements

We would like to thank Jun Li, Davesh Maulik, and Richard Thomas for helpful conversations. The work of S. Gukov is supported in part by the U.S. Department of Energy, Office of Science, Office of High Energy Physics, under Award Number DE-SC0011632 and in part by the ERC Starting

Grant no. 335739 “Quantum fields and knot homologies” funded by the European Research Council under the European Union Seventh Framework Programme. The work of C.-C. Liu is partially supported by NSF grants DMS-1206667 and DMS-1159416. A. Sheshmani would like to thank Kavli IPMU, MIT, Harvard and the Institute Henri Poincaré (IHP) for creating the opportunity of initiating the discussions about the current article. The work of A. Sheshmani was partially supported by the World Premier International Research Center Initiative (WPI Initiative), MEXT, Japan and Laboratory of Mirror Symmetry NRU HSE, RF Government grant, ag. No 14.641.31.0001. S.-T. Y. was partially supported by NSF DMS-0804454, NSF PHY-1306313, and Simons 38558. The work of S.-T. Yau is partially supported by NSF grants DMS 1308244, DMS-159412, PHY-1306313, and PHY-0937443.

2. Preliminaries

2.1. Gromov-Witten theory

2.1.1. Gromov-Witten invariants and Taubes’s Gromov invariants of symplectic 4-manifolds. In this subsection, we briefly review certain Gromov-Witten invariants of compact symplectic 4-manifolds and Taubes’s Gromov invariants [43, 85].

Let (S, ω) be a compact symplectic 4-manifold with an almost complex structure J compatible with the symplectic structure ω , so that (S, ω, J) is an almost Kähler manifold. If J is integrable then (S, ω, J) is a Kähler manifold.

Let $\overline{\mathcal{M}}_{g,n}(S, \beta)$ be the moduli space of genus g , degree $\beta \in H_2(S; \mathbb{Z})$ stable maps to (S, ω, J) . The virtual dimension of $\overline{\mathcal{M}}_{g,n}(S, \beta)$ is given by

$$\int_{\beta} c_1(S) + g - 1 + n = -K_S \cdot \beta + g - 1 + n,$$

where K_S is the canonical class of S .

Let $\nu \in H^4(S; \mathbb{Z})$ be the Poincaré dual of the class of a point. We define

$$(1) \quad N_{g,\beta}^S := \begin{cases} \int_{[\overline{\mathcal{M}}_{g,n}(S,\beta)]^{\text{vir}}} \prod_{i=1}^n \text{ev}_i^*(\nu), & \text{if } n = -K_S \cdot \beta + g - 1 \geq 0, \\ 0, & \text{if } -K_S \cdot \beta + g - 1 < 0, \end{cases}$$

where $\text{ev}_i : \overline{\mathcal{M}}_{g,n}(S, \beta) \rightarrow S$ is the evaluation at the i -th marked point. By the divisor equation, any genus g , degree β primary Gromov-Witten invariants of S can be reduced to $N_{g,\beta}^S$. In general, Gromov-Witten invariants

count connected parametrized curves, and are rational numbers instead of integers.

By adjunction formula, the genus of a connected, embedded smooth J -holomorphic curve in class $\beta \in H_2(S; \mathbb{Z})$ is given by

$$1 + \frac{1}{2}(\beta \cdot \beta + K_S \cdot \beta).$$

Gromov invariant $N_{g,\beta}^S$ counts connected embedded curves of genus g in class β passing through n generic points if the following three conditions holds:

- 1) $g = 1 + \frac{1}{2}(\beta \cdot \beta + K_S \cdot \beta)$.
- 2) $n = \frac{1}{2}(\beta \cdot \beta - K_S \cdot \beta) \geq 0$.
- 3) $\beta \notin T$, where $T = \{\beta \in H_2(S, \mathbb{Z}) \mid \beta \cdot K_S = \beta \cdot \beta = 0\}$ is the set of *toroidal cases*.

When the above conditions hold, the moduli space of J -holomorphic curves of genus g in class β passing through n points contains no multiply covered curves for generic J (see Ruan [81], Taubes [85]). By [43, Section 4],

$$N_{g,\beta}^S = \widehat{Gr}_S(\beta)$$

where $\widehat{Gr}_S(\beta)$ is the connected version of Taubes's Gromov invariant in class β . Taubes's Gromov invariant $Gr_S(\beta)$, which counts possibly disconnected curves, is equivalent to certain Seiberg-Witten invariant by Taubes's famous theorem.

In this paper, we will assume J is integrable, so that the pair (ω, J) is a Kähler structure on S .

2.1.2. Non-equivariant local Gromov-Witten invariants of surfaces in a Calabi-Yau threefold. Let S be a projective Fano surface, and let X_S be the total space of the canonical line bundle K_S over S . Then X_S is a noncompact Calabi-Yau 3-fold. Given any nonzero $\beta \in H_2(S; \mathbb{Z}) \cong H_2(X_S; \mathbb{Z})$, we have

$$\overline{\mathcal{M}}_{g,0}(X_S, \beta) = \overline{\mathcal{M}}_{g,0}(S, \beta)$$

as Deligne-Mumford stacks. The virtual dimensions of the perfect obstruction theories on $\overline{\mathcal{M}}_{g,0}(X_S, \beta)$ and on $\overline{\mathcal{M}}_{g,0}(S, \beta)$ are 0 and $-K_S \cdot \beta + g - 1$,

respectively, and

$$N_{g,\beta}^{X_S} = \int_{[\overline{\mathcal{M}}_{g,0}(X_S,\beta)]^{\text{vir}}} 1 = \int_{[\overline{\mathcal{M}}_{g,0}(S,\beta)]^{\text{vir}}} e(V_{g,\beta})$$

is the genus g , degree β Gromov-Witten invariant of X_S , where $V_{g,\beta}$ is a rank $-K_S \cdot \beta + g - 1$ vector bundle over $\overline{\mathcal{M}}_{g,0}(S, \beta)$ whose fiber over $[f : C \rightarrow S]$ is $H^1(C, f^*K_S)$. Note that because β is nonzero and $-K_S$ is ample, we have $\deg(f^*K_S) < 0$, so $H^0(C, f^*K_S) = 0$. (See [17, Section 5.2].) The invariants $N_{g,\beta}^{X_S}$ can be viewed as local invariants of the Fano surfaces S in a Calabi-Yau 3-fold.

When S is projective but not Fano, the Gromov-Witten invariants of X_S are not necessarily defined, due to the non-compactness of moduli spaces $\overline{\mathcal{M}}_{g,0}(X_S, \beta)$. Observe that, when the class β satisfies $-K_S \cdot \beta > 0$, $V_{g,\beta}$ is a rank $-K_S \cdot \beta + g - 1$ vector bundle over $\overline{\mathcal{M}}_{g,0}(S, \beta)$. This allows us to define the genus g , degree β Gromov-Witten invariant of X_S by

$$(2) \quad N_{g,\beta}^{X_S} := \int_{[\overline{\mathcal{M}}_{g,0}(S,\beta)]^{\text{vir}}} e(V_{g,\beta}).$$

Note that $e(V_{g,\beta}) = 1$ if and only if $H^1(C, f^*K_S) = 0$ for all $[f : C \rightarrow X_S] \in \overline{\mathcal{M}}_{g,0}(S, \beta)$. In this case

$$\int_{[\overline{\mathcal{M}}_{g,0}(X_S,\beta)]^{\text{vir}}} 1 = \int_{[\overline{\mathcal{M}}_{g,0}(S,\beta)]^{\text{vir}}} 1$$

is also the genus g , degree β Gromov-Witten invariant of the surface S . We have

$$-K_S \cdot \beta + g - 1 = 0$$

where $-K_S \cdot \beta > 0$, so we must have $g = 0$ and $K_S \cdot \beta = -1$. We summarize the above discussion in the following lemma.

Lemma 1. *Let S be a projective surface, and suppose $\beta \in H_2(S, \mathbb{Z})$ satisfies $K_S \cdot \beta = -1$. Then*

$$N_{0,\beta}^{X_S} = N_{0,\beta}^S$$

where the left hand side is defined by (2) and the right hand side is defined as in Section 2.1.1.

In the remainder of this subsection, we assume

$$S \text{ is Fano, } g = 0, \quad K_S \cdot \beta = -1.$$

Then a curve in class β cannot be multiply covered. We claim that there is no bubbling. Indeed if $\beta = \beta_1 + \beta_2$ where β_1, β_2 are nonzero effective class, we have $-K_S \cdot \beta_i > 0$ for $i = 1, 2$ so $-K_S \cdot \beta \geq 2$, which is a contradiction. Similarly, β cannot be represented by a disconnected holomorphic curve. Combining the above discussion and the discussion Section 2.1.1, we conclude:

Lemma 2. *Let S be a Fano surface, and suppose that $\beta \in H_2(S, \mathbb{Z})$ satisfies*

$$K_S \cdot \beta = \beta \cdot \beta = -1.$$

Then

$$N_{0,\beta}^{X_S} = N_{0,\beta}^S = \widehat{Gr}_S(\beta) = Gr_S(\beta).$$

Let B_k denote \mathbb{P}^2 blowup at k generic points. By classification of surfaces, Fano surfaces are: \mathbb{P}^2 , $\mathbb{P}^1 \times \mathbb{P}^1$, and del Pezzo surfaces B_k where $1 \leq k \leq 8$. If $S = \mathbb{P}^2$ or $\mathbb{P}^1 \times \mathbb{P}^1$ then there is no class β satisfying $K_S \cdot \beta = -1$. If $S = B_k$ and $e_1, \dots, e_k \in H_2(S, \mathbb{Z})$ are classes of exceptional divisors, then $K_S \cdot e_i = e_i \cdot e_i = -1$.

2.1.3. Equivariant local Gromov-Witten invariants of surfaces in a Calabi-Yau threefold. It is possible to define *residue invariants* in the presence of a torus action [15, Section 2.1]. We first consider the special case of toric surfaces. We have

$$N_{g,\beta}^{X_S} = \int_{[\overline{\mathcal{M}}_{g,0}(X_S, \beta)]^{\text{vir}}} 1 = \int_{[\overline{\mathcal{M}}_{g,0}(X_S, \beta)^T]^{\text{vir}}} \frac{1}{e_T(N^{\text{vir}})}.$$

where T acts on X_S , and $e_T(N^{\text{vir}})$ is the T -equivariant Euler class of the virtual normal bundle of $\overline{\mathcal{M}}_{g,0}(X_S, \beta)^T$ in $\overline{\mathcal{M}}_{g,0}(X_S, \beta)$. When the T -action on X_S induced from a T -action on the base S , so that the T -action on X_S acts trivially on the canonical line bundle of the toric threefold X_S , these invariants can be evaluated by the algorithm of the topological vertex [2]. Indeed, for local toric surfaces, only 2-leg vertex is needed; the algorithm in this case is described in [4]. The topological vertex is proved in the 1-leg case in [60, 78] and the 2-leg case in [57, 61]. The full 3-leg case is a consequence of the proof of the GW/DT correspondence of toric 3-folds in [67].

In general, if there is a torus action on X_S such that X_S^T is contained in the zero section S and $\overline{\mathcal{M}}_{g,0}(X_S, \beta)^T = \overline{\mathcal{M}}_{g,0}(S, \beta)^T$ is compact, we define

$$(3) \quad N_{g,\beta}^{X_S}(t_1, \dots, t_r) = \int_{[\overline{\mathcal{M}}_{g,0}(X_S, \beta)^T]^{\text{vir}}} \frac{1}{e_T(N^{\text{vir}})} \in \mathbb{Q}(t_1, \dots, t_r)$$

where t_1, \dots, t_r are the generators of the $T = (\mathbb{C}^*)^r$ equivariant cohomology of a point. Alternatively,

$$(4) \quad N_{g,\beta}^{X_S}(t_1, \dots, t_r) = \int_{[\overline{\mathcal{M}}_{g,0}(S, \beta)^T]^{\text{vir}}} \frac{e_T(H^1)/e_T(H^0)}{e_T(N^{\text{vir}})} \in \mathbb{Q}(t_1, \dots, t_r)$$

where $e_T(N^{\text{vir}})$ is the T -equivariant Euler class of the virtual normal bundle of $\overline{\mathcal{M}}_{g,0}(S, \beta)^T$ in $\overline{\mathcal{M}}_{g,0}(S, \beta)$. Here H^1 and H^0 are vector bundles on each connected component of $\overline{\mathcal{M}}_{g,0}(S, \beta)^T$, and their fibres over $[f : C \rightarrow S]$ are $H^1(C, f^*K_S)$ and $H^0(C, f^*K_S)$, respectively.

2.2. Donaldson-Thomas theory

We consider a general gauge group G which can be any compact connected Lie group. The complexification $G^{\mathbb{C}}$ of G is a connected reductive complex algebraic group. For example, when $G = U(N)$, we have $G^{\mathbb{C}} = GL(N, \mathbb{C})$. The space of connections on a principal G -bundle P over a Kähler manifold can be identified with the space of $\bar{\partial}$ -operators on the principal $G^{\mathbb{C}}$ bundle $P^{\mathbb{C}} = P \times_G G^{\mathbb{C}}$.

The Donaldson-Thomas theory is a topological gauge theory defined on a complex Kahler 3-fold X which, roughly speaking, “counts” solutions to the Hermitian Yang-Mills (HYM) equations (see e.g. [1, Definition 3.2]):

$$(5) \quad F_A^{0,2} = 0, \quad \text{Tr}_{k_0} F_A^{1,1} = \tau.$$

where $A \in \Omega_X^1(\text{ad}P)$ is the gauge connection, k_0 is the Kähler form on X , and τ is an element in the center of the Lie algebra \mathfrak{g} of G . For example, $\tau = 0$ if G is semisimple, and τ is a multiple of the identity element if $G = U(N)$.

This six-dimensional TQFT can be obtained by a topological twist of the dimensional reduction of the ten-dimensional super-Yang-Mills [3, 6, 10, 12, 25, 45]. Under the reduction,

$$SO(10) \rightarrow SO(6) \times SO(4)$$

and bosonic/fermionic fields decompose as

$$(6) \quad \begin{array}{ll} \text{bosons :} & \mathbf{10} \rightarrow (\mathbf{6}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{4}) \\ \text{fermions :} & \mathbf{16} \rightarrow (\mathbf{4}, \mathbf{2}) \oplus (\bar{\mathbf{4}}, \mathbf{2}) \end{array}$$

In addition, if X is Kähler, we have $SO(6) \rightarrow U(3) \cong SU(3) \times U(1)$. The twisted six-dimensional gauge theory is obtained by mixing the last $U(1)$ factor with $U(1) \subset SO(4)$. As a result, one finds a theory with $\mathcal{N}_T = 2$ topological supersymmetries (nilpotent BRST symmetries) and the following field content:

bosonic:	1-form	A	
	(3, 0)-form	φ	
		complex scalar	$\phi, \bar{\phi}$
fermionic:	1-form	$\psi^{1,0}, \psi^{0,1}$	3+3
	2-form	$\chi^{2,0}, \chi^{0,2}$	3+3
	3-form	$\psi^{3,0}, \psi^{0,3}$	1+1
	two scalars	$\eta, \bar{\eta}$	1+1

where all fields transform in the adjoint representation of the gauge group G and in the last column we counted the number of real components. Notice that, altogether, the fermionic fields contain 16 real components. If X is a Calabi-Yau space, the resulting theory has $\mathcal{N}_T = 4$ topological supersymmetries.

The partition function of the six-dimensional topological gauge theory localizes on the solutions to the following equations

$$(7) \quad \begin{aligned} F_A^{0,2} &= \bar{\partial}_A^\dagger \bar{\varphi} \\ \text{Tr}_{k_0} F_A^{1,1} + *[\varphi, \bar{\varphi}] &= \tau \\ d_A \phi &= 0 \end{aligned}$$

where the first two equations can be recognized as perturbations of the HYM equations (5). The Bianchi identity and the first equation imply $\bar{\partial}_A \bar{\partial}_A^\dagger \bar{\varphi} = 0$, so when X is compact, the first equation is decoupled into two equations $\bar{\partial}_A^\dagger \bar{\varphi} = 0$ and $F_A^{0,2} = 0$, and the path integral localizes on the solutions to

$$(8) \quad \bar{\partial}_A^\dagger \bar{\varphi} = 0, \quad F_A^{0,2} = 0, \quad \text{Tr}_{k_0} F_A^{1,1} + *[\varphi, \bar{\varphi}] = \tau$$

Note that the solutions to the HYM equations (5) are also solutions to (8). In [38], the moduli space defined by (8) is called the *extended* moduli space

of Einstein-Hermitian (i.e. HYM) connections. We have

$$\mathcal{M}_{\text{HYM}}(X, G) \subset \mathcal{M}_{\text{DT}}(X, G),$$

where $\mathcal{M}_{\text{HYM}}(X, G)$ and $\mathcal{M}_{\text{DT}}(X, G)$ are moduli spaces of solutions to (5) and (8), respectively. Given a holomorphic structure, the space of solutions to $\bar{\partial}_A^\dagger \bar{\varphi} = 0$ can be identified with $H^0(X, K_X \otimes \text{ad}(P)) = H^3(X, \text{ad}(P))^\vee$.

When X is Calabi-Yau, φ can be viewed as an element of $H^0(X, \text{ad}(P))$. If P is stable, then φ lies in the center $Z(\mathfrak{g})$ of \mathfrak{g} , so $[\varphi, \bar{\varphi}] = 0$, so the second and third equations in (8) become the HYM equations (5). For example, $Z(\mathfrak{su}(N))$ is trivial, and $Z(\mathfrak{u}(N))$ is spanned by the identity matrix. When X is Fano, i.e., $-K_X > 0$, we have $H^0(X, K_X \otimes \text{ad}(P)) = H^3(X, \text{ad}(P))^\vee = 0$.

Let \mathcal{M} denote the moduli space of gauge equivalence classes of solutions to (7). The tangent space of \mathcal{M} at a point (A, φ) can be identified with H^1 of the following deformation complex¹

$$(9) \quad 0 \rightarrow \Omega_X^{0,0}(\text{ad}P_{\mathbb{C}}) \xrightarrow{D_1} \Omega_X^{0,1}(\text{ad}P_{\mathbb{C}}) \oplus \Omega_X^{0,3}(\text{ad}P_{\mathbb{C}}) \xrightarrow{D_2} \Omega_X^{0,2}(\text{ad}P_{\mathbb{C}}) \rightarrow 0.$$

When X is Calabi-Yau, the index of the above deformation complex is zero. We have

$$(10) \quad \dim T_{(A,\varphi)}\mathcal{M} = \dim \mathcal{N}_{(A,\varphi)}$$

where $\mathcal{N}_{(A,\varphi)}$ is H^2 of the above deformation complex. In this case, from 1-loop determinants one finds that the partition function computes the Euler characteristic of the obstruction bundle \mathcal{N} . We have

$$\begin{aligned} Z_{DT}^G &= \int DAD\varphi D(\dots) \exp(S(A, \varphi, \dots)) \\ &= \int_{\mathcal{M}} e(\mathcal{N}) \exp \int_X \text{Tr} \left(\frac{g_s}{3!} F \wedge F \wedge F + \frac{1}{2} F \wedge F \wedge k_0 + F \wedge \vartheta \right) \end{aligned}$$

where $e(\mathcal{N})$ is the Euler class of \mathcal{N} . The integral over X is topological in the sense that it is constant on each connected component of \mathcal{M} . For example,

¹Using $\bar{\partial}_A^2 \eta = [F_A^{0,2}, \eta]$ and $F_A^{0,2} = \bar{\partial}_A^\dagger \bar{\varphi}$, one can check that this is actually a complex:

$$D_2 \circ D_1(\eta) = D_2(\bar{\partial}_A \eta, [\bar{\varphi}, \eta]) = \bar{\partial}_A^2 \eta - \bar{\partial}_A^\dagger [\bar{\varphi}, \eta] = [\bar{\partial}_A^\dagger \bar{\varphi}, \eta] - \bar{\partial}_A^\dagger [\bar{\varphi}, \eta] = 0$$

when $G = U(N)$, let $\mathcal{M}_{ch_1, ch_2, ch_3}$ be the union of connected components of \mathcal{M} with fixed Chern characters ch_1, ch_2, ch_3 . Then

$$(11) \quad Z_{DT}^{U(N)}(q, t, T) = \sum_{ch_1, ch_2, ch_3} \int_{\mathcal{M}_{ch_1, ch_2, ch_3}} e(\mathcal{N}) q^{ch_3(\mathcal{E})} t^{ch_2(\mathcal{E})} T^{ch_1(\mathcal{E})}$$

where $q = e^{g_s}$, t , and T are formal variables that encode dependence of the Donaldson-Thomas partition function on the Chern classes. More generally, when X is not a Calabi-Yau manifold, the index of the deformation complex (9) is

$$(12) \quad \int_X c_1(T_X) \left(ch_0 ch_2 - \frac{ch_1^2}{2} + \frac{ch_0^2 - 1}{24} c_2(T_X) \right) = \text{rank}_{\mathbb{C}} \mathcal{N} - \dim_{\mathbb{C}} \mathcal{M}$$

and the correlation functions are of the form

$$(13) \quad \int_{\mathcal{M}_{ch_1, ch_2, ch_3}} e(\mathcal{N}) \mathcal{O}_1 \cdots \mathcal{O}_k$$

where $\mathcal{O}_i \in H^*(\mathcal{M}_{ch_1, ch_2, ch_3})$ are topological observables, invariant under the BRST symmetry.

In order to define (13) mathematically, one needs to compactify the moduli space \mathcal{M} . By Uhlenbeck and Yau’s result [91], the moduli space of gauge equivalence classes of HYM $U(N)$ -connections can be identified with the moduli space of isomorphism classes of polystable vector bundles of rank N . The later can be compactified by the moduli space of rank N semistable sheaves. In the algebro-geometric framework, the deformation complex (7) is replaced by the tangent-obstruction complex which encodes the deformation theory of the moduli problem. When higher obstruction groups vanish, one obtains virtual fundamental cycle $[\mathcal{M}]^{\text{vir}}$, and the mathematical definition of the integral (13) is given by

$$(14) \quad \int_{[\mathcal{M}_{ch_1, ch_2, ch_3}]^{\text{vir}}} \mathcal{O}_1 \cdots \mathcal{O}_k \in \mathbb{Z}.$$

The deformation theory and the virtual fundamental cycle have been established for smooth projective Calabi-Yau and Fano 3-folds in [90]. In string theory, the Donaldson-Thomas theory with gauge group $G = U(N)$ can be interpreted as a gauge theory on N D6-branes wrapped on X . In this interpretation, the integer coefficients of the partition function Z_{DT} count BPS states with D0, D2, D4, and D6-brane changes. We remind that D-brane charges are described by the Mukai vector, $Q \in H^0(X) \oplus H^2(X) \oplus$

$H^4(X) \oplus H^6(X)$, given by

$$Q = ch(\mathcal{E})\sqrt{Td(X)}$$

These invariants are in fact the generalized DT invariants which were constructed and computed by the wall crossing techniques of Kontsevich-Soibelman [49] and Joyce-Song [46]. In connection with Gromov-Witten invariants which count holomorphic curves, one considers moduli space of ideal sheaves of curves. These are torsion free, rank 1 sheaves with trivial determinant, so they correspond to abelian theory with $ch_1 = 0$. The conjectural GW/DT correspondence [65, 66] (proved for toric 3-folds [67] and local curves [15, 80]) states that

$$Z'_{DT}(q, t) = Z'_{GW}(g_s, t)$$

where $Z'_{DT}(q, t)$ and $Z'_{GW}(g_s, t)$ are the *reduced* DT and GW partition functions. Roughly speaking, $Z'_{DT}(q, t)$ is obtained from $Z_{DT}(q, t)$, by removing the contribution of point-like instantons, and $Z'_{GW}(g_s, t)$ is obtained from $Z_{GW}(g_s, t)$ by removing the contribution of constant maps.

In certain cases, the Donaldson-Thomas theory on X can be related to topological theories in lower dimensions. In this paper, we describe two such relations which, in a sense, are analogues of the corresponding relations in four-dimensional topological gauge theory studied, respectively, in [9] and [70].

2.3. Seiberg-Witten theory

In this subsection, we briefly review the definition of the Seiberg-Witten invariants [95]

$$(15) \quad SW_S : Spin^c(S) \rightarrow \mathbb{Z}$$

and their relation [85] with the enumerative invariants of S .

Up to the finite group $H^1(S, \mathbb{Z}_2)$, the set of $Spin^c$ structures on a 4-manifold S is parameterized by integral cohomology classes² which reduce

²Sometimes, we parametrize the set of $Spin^c$ structures by $\lambda = \frac{c}{2} \in H^2(S, \mathbb{Z}) + w_2(S)/2$, which is widely used in the physics literature.

to $w_2(X) \bmod 2$

$$\text{Spin}^c(S) = \{c \in H^2(S, \mathbb{Z}) \mid c \equiv w_2(S) \bmod 2\}$$

Given a Spin^c structure $c \in \text{Spin}^c(S)$, let L be the corresponding Hermitian line bundle, and S_L^\pm the corresponding spinor bundles. The Seiberg-Witten monopole equations are equations for a pair (A, Ψ) , where A is a unitary connection on L and Ψ is a smooth section of S_L^+ . In order to write the equations, we need to introduce the Dirac operator

$$D_A : S_L^+ \rightarrow S_L^-$$

and a map

$$\Omega^0(S_L^+) \rightarrow \Omega^0(\text{ad}_0 S_L^+)$$

$$\Psi \mapsto i(\Psi \otimes \Psi^*)_0$$

where Ψ^* is the adjoint of Ψ and $\text{ad}_0 S_L^+$ is the subbundle of the adjoint bundle of S_L^+ consisting of the traceless skew-Hermitian endomorphisms. Then, the Seiberg-Witten equations take the form [95]

$$D_A \Psi = 0$$

$$F_A^+ - i(\Psi \otimes \Psi^*)_0 = 0$$

where in the second equation we used the identification between the space of self-dual 2-forms and skew-Hermitian automorphisms of the positive spin representation, $\text{ad}_0 S_L^+ \cong \Lambda_+^2$.

Let \mathcal{M}_c be the moduli space of solutions to the Seiberg-Witten equations. It has virtual dimension

$$d_c = \frac{1}{4}(c^2 - 2\chi(S) - 3\sigma(S))$$

The Seiberg-Witten invariants are defined as³

$$SW_S(c) = \int_{\mathcal{M}_c} a_D^{d_c/2}$$

where a_D is a 2-form which represents the first Chern class of the universal line bundle on the moduli space \mathcal{M}_c . Notice, that the dimension of the

³Even though some of our examples below include 4-manifolds with $b_1 > 0$, we will not consider classes in $H^1(S; \mathbb{R})$. In particular, we will not worry about the orientation of $H^1(S; \mathbb{R})$.

Seiberg-Witten moduli space and $b_2^+ + b_1$ have opposite parity. This fact can be used to establish vanishing of Seiberg-Witten invariants when $b_2^+ + b_1$ is even. Another useful fact is a vanishing theorem [95], which says that $SW_S(c) = 0$ whenever S admits a metric of positive scalar curvature. This theorem is especially helpful in some of our examples, where the existence of a positive curvature Kähler-Einstein metric on S is directly related to the existence of a complete Ricci flat Kähler metric (Calabi-Yau metric) on X_S [88, 89, 97].

Among the examples that we consider, there are 4-manifolds with $b_2^+ = 1$. In such cases, Seiberg-Witten invariants are not quite topological invariants; they are only piece-wise constant and exhibit discontinuous behavior under wall crossing that we briefly review below.

Let H be the hyperbolic space

$$H = \{h \in H_{DR}^2(S) \mid h^2 = 1\}$$

This space has two connected components and the choice of one of them orients the lines $H_g^{2,+}(S)$ for all metrics g . Having fixed a component H_0 of H , every metric defines a unique self-dual 2-form ω of length 1 with $[\omega] \in H_0$, called the period point, $\omega^2 = 1$. We have $H^{2,+}(S; \mathbb{R}) \cong \mathbb{R}\omega$ and c can be written as $c = c_+ + c_-$ where $c_+ = (c \cdot \omega)\omega$.

The space H is divided into a set of chambers C , and for every metric g whose self-dual 2-form ω lies in $\mathbb{R}_+ \cdot C$ one can define the corresponding Seiberg-Witten invariants $SW_{S,g} = SW_{S,C}$. Notice, that changing ω to $-\omega$ corresponds to changing the orientation of the Seiberg-Witten moduli space, so that

$$(16) \quad SW_{S,-C} = -SW_{S,C}$$

The Seiberg-Witten invariants $SW_{S,C}$ can be defined using the perturbed (or “twisted”) monopole equations⁴

$$D_A \Psi = 0$$

$$(F_A + 2\pi i\beta)^+ = i(\Psi \otimes \Psi^*)_0$$

where β is a closed 2-form with $b = [\beta] \in H^2(S; \mathbb{R})$. Equivalently, we consider a Spin^c structure given by $(c - b)$. Each characteristic class $c \in H^2(S, \mathbb{Z})$,

⁴In topological gauge theory, the perturbation by the self-dual 2-form β^+ corresponds to perturbation by a mass term [96].

that is an integral lift of $w_2(S)$, defines four chambers of type c :

$$C_{H_0, \pm} = \{(h, b) \in H_0 \times H_{DR}^2(S) \mid \pm (c - b) \cdot h < 0\}$$

where H_0 is one of the components of H . Following the standard conventions, we denote the corresponding Seiberg-Witten invariants by $SW_{S, H_0}^{\pm}(c)$ (or simply by $SW_S^{\pm}(c)$). Notice, according to (16), we have $SW_{S, -H_0}^{\pm}(c) = -SW_{S, H_0}^{\mp}(c)$.

The difference between the Seiberg-Witten invariants across a wall is given by a simple formula, which for a simply-connected surface S takes the form

$$SW_S^+(c) - SW_S^-(c) = 1$$

For example, for $S = \mathbb{P}^2$, we have $\text{Spin}^c(\mathbb{P}^2) \cong (2\mathbb{Z} + 1)h$, where h denotes the first Chern class of $\mathcal{O}(1)$, $h^2 = 1$. And for every $c \equiv h \pmod{2}$, there are two chambers of type c :

$$C_{H_0, \pm} = \{(h, b) \in H_0 \times H_{DR}^2(S) \mid \pm (c - b) \cdot h < 0\}$$

where $H_0 = \{h\}$ and $d_c = \frac{1}{4}(c^2 - 9)$. The corresponding Seiberg-Witten invariants are

$$(17) \quad \begin{aligned} SW_S^+(c) &= 1 & c \cdot h &\geq 3 \\ SW_S^+(c) &= 0 & c \cdot h &< 3 \\ SW_S^-(c) &= -1 & c \cdot h &\leq -3 \\ SW_S^-(c) &= 0 & c \cdot h &> -3 \end{aligned}$$

More generally, for Kähler surfaces with $b_2^+ = 1$ and $b_1 = 0$, we have $SW_S^{\pm}(c) = \{0, 1\}$ or $SW_S^{\pm}(c) = \{0, -1\}$ as soon as $d_c \geq 0$ [77].

3. Dimensional reduction of DT theory to 4d Vafa-Witten theory

3.1. Dimensional reduction of DT theory on an elliptic fibration

Suppose that the 3-fold X is an elliptic fibration over a surface S . In particular, one can consider a trivial fibration, $X = E \times S$. Then, we can interpret the partition function of the Donaldson-Thomas theory on X as counting BPS states in the T-dual system. Specifically, under a T-duality along E , the D6-branes on X turn into D4-branes on S , the D0-brane charge becomes the charge of the D2-brane on E , and so on.

For general gauge group G , D-brane charges can be viewed as homology classes which are Poincaré duals of the obstruction classes $o_i \in H^i(X, \pi_1(G))$. We consider compact connected Lie groups, so $\pi_0(G)$ is trivial and the first obstruction class $o_1 = 0$. The second obstruction class $o_2 \in H^2(X, \pi_1(G))$ is the magnetic flux m . When $G = U(N)$, $m = c_1 = ch_1 \in H^2(X, \mathbb{Z})$ is the first Chern class (which is also the first Chern character); when $G = SO(N)$ ($N > 2$), $m = w_2 \in H^2(X, \mathbb{Z}/2\mathbb{Z})$, is the second Steifel-Whitney class. In the rest of Section 3, we will consider $G = U(N)$, and the D-brane charges are described by the Mukai vector as recalled in Section 2.2.

In general, the resulting configuration of D-branes is similar to the one we started with. However, there are several cases when this duality can be very useful and can relate Donaldson-Thomas invariants to some invariants in the four-dimensional topological gauge theory. The four-dimensional gauge theory is a twisted gauge theory on N D4-branes wrapped around the base S , obtained from N D6-branes in the original setup. This gauge theory is a topological twist of $\mathcal{N} = 4$ super-Yang-Mills studied by Vafa and Witten. It has the following spectrum [93]:

bosonic:	1-form	A	
	self-dual 2-form	B_+	3
	scalar	C	1
fermionic:	complex scalar	$\phi, \bar{\phi}$	2
	self-dual 2-forms	$\chi, \tilde{\psi}$	3+3
	vectors	$\psi, \tilde{\chi}$	4 + 4
	scalars	η, ζ	1+1

where in the last column we again summarized the number of real components.⁵ Under a favorable set of conditions⁶, the partition function in this theory localizes on anti-self-dual gauge fields (instantons):

$$(18) \quad F_A^+ = 0$$

and has the form similar to (11).

In general, the partition function of this four-dimensional topological gauge theory can be interpreted as counting BPS states with D2, D4, and D6 brane charges on X , where all of these branes are wrapped on the elliptic fiber E . Specifically, the dependence on the D2 and D4 brane charges is through the gauge coupling $q = \exp(2\pi i\tau)$ and the 't Hooft magnetic fluxes m , while D6-brane charge corresponds to the rank of the gauge group.

⁵Notice, that the total number of fermions is 16.

⁶More specifically, when vanishing theorems hold.

In our case, $G = U(N)$ and D2, D4, D6 brane charges are cycles dual to ch_2, ch_1, ch_0 , respectively, and the partition function of the Vafa-Witten theory can be interpreted as a partition function of the Donaldson-Thomas theory,

$$(19) \quad Z_{DT}(X)_{ch_3=0} = Z_{VW}(S) = \sum_{k,m} \chi(\mathcal{M}_{m,k}) t^m q^k$$

where $\mathcal{M}_{m,k}$ is the moduli space of solutions to (18) with

$$k = \int_S ch_2(\mathcal{E}), \quad m = ch_1(\mathcal{E}), \quad N = ch_0(\mathcal{E}).$$

Notice that we have $c_1(T_X) = c_1(T_S)$ so the index (12) becomes

$$\int_X c_1(T_S) \left(ch_0 ch_2 - \frac{ch_1^2}{2} + \frac{ch_0^2 - 1}{24} c_2(T_S) \right)$$

which is zero when ch_0, ch_1, ch_2 are classes from $H^*(S, \mathbb{Z})$, so this part of the theory is balanced (in the sense of [20]). This is consistent with the fact that the Vafa-Witten theory is balanced, even when $c_1(T_S) \neq 0$.

The partition function $Z_{DT}(X) = Z_{DT}(E \times S)$ is defined mathematically when S is a K3 surfaces. In this case, let

$$\tilde{N}_{ch_1, ch_2, ch_3} = \int_{[\mathcal{M}_{ch_1, ch_2, ch_3}]^{vir}} 1$$

be the Donaldson-Thomas (holomorphic Casson) invariants. R. Thomas verified that in some cases $\tilde{N}_{m,k,0} = N^2 \chi(\mathcal{M}_{m,k})$, where the factor N^2 comes from the N -th root of unity in the Jacobian of E [90].

3.2. Abelian Vafa-Witten theory

In the abelian gauge theory, we have $ch_0(\mathcal{E}) = 1$. In order for $ch_2(\mathcal{E})$ to be non-trivial, we need to enlarge the space of gauge connections to include singular gauge fields (point-like instantons) which mathematically can be described as ideal sheaves. The partition function of the abelian gauge theory on S is

$$(20) \quad Z(q, t) = \frac{\vartheta_\Gamma(t, \tau, \bar{\tau})}{\eta(\tau)^{\chi(S)}}$$

where the factor $\eta^{-\chi(S)}$ is the contribution from the point-like instantons ($m = 0$) and the theta-function comes from the lattice sum over magnetic fluxes $m \in \Gamma$.

The moduli space of point-like instantons can be identified with Hilbert scheme of points on S . According to Hirzebruch-Höfer [37], the orbifold Euler number of the symmetric product $\text{Sym}^k S$ is equal to the Euler number of the Hilbert scheme $\text{Hilb}^k S$ of k points in S , which is all we need.

Göttsche [33] has shown:

$$\sum_{k=0}^{\infty} q^k P_t(\text{Hilb}^k S) = \prod_{m=1}^{\infty} \prod_{i=1}^4 (1 - (-t)^{2m-2+1} q^m)^{(-1)^{i+1} b_i(S)}$$

In particular,

$$\sum_{k \geq 0} \chi(\text{Hilb}^k S) q^{k-\chi(S)/24} = \eta(\tau)^{-\chi(S)}$$

or equivalently,

$$(21) \quad \sum_{k \geq 0} \chi(\text{Hilb}^k S) q^k = \prod_{n > 0} \frac{1}{(1 - q^n)^{\chi(M)}} = q^{\frac{\chi(S)}{24}} \eta(\tau)^{-\chi(S)}$$

This formula can be also obtained as a special case of the elliptic genus for the symmetric product CFT with target space $\text{Sym}^k S$ [21].

3.3. Examples and applications

Below we consider two groups of examples: 1) abelian gauge theory on S , and 2) non-abelian $U(N)$ gauge theory with $m = 0$. The former is more relevant to the DT/GW correspondence, while the latter gives us partition function of the DT theory on elliptic fibration with $ch_1(\mathcal{E}) = 0$ and $ch_3(\mathcal{E}) = 0$.

3.4. Abelian DT theory and GW theory on a trivial elliptic fibration

Let $X = E \times S \rightarrow S$ be a trivial elliptic fibration over a smooth projective surface S , where E is an elliptic curve. In this subsection, we follow Edidin and Qin's computations of DT and GW invariants of $X = E \times S$ in [26].

3.4.1. Abelian DT theory. Following [65, 66], given $n \in \mathbb{Z}$ and $\beta \in H_2(X; \mathbb{Z})$, let $I_n(X, \beta)$ be the moduli space of ideal sheaves I_Z of 1-dimensional closed subschemes Z of X satisfying $\chi(\mathcal{O}_Z) = n$ and $[Z] = \beta$.

The virtual dimension of $I_n(X, \beta)$ is $-\int_{\beta} K_S$. Let $\beta_0 = [E] \in H_2(X; \mathbb{Z})$ be the fiber class. The virtual dimension of $I_n(X, k\beta_0)$ is zero. Define

$$\tilde{N}_{n,k} = \int_{[I_n(X, k\beta_0)]^{\text{vir}}} 1.$$

The integral can be calculated by localization. The elliptic curve E is a group (compact torus). Let E act on itself by group multiplication and act on S trivially. This gives a torus action on $X = E \times S$ and torus actions on moduli spaces $I_n(X, k\beta_0)$. The action has no fixed point unless $n = 0$. So

$$\tilde{N}_{n,k} = 0, \quad n \neq 0.$$

When $n = 0$, the moduli space $I_0(X, k\beta_0)$ can be identified with $\text{Hilb}^k S$, Hilbert scheme of k points in S , which is a smooth variety of complex dimension $2k$. The virtual dimension of $I_0(X, k[E])$ is zero. Edidin and Qin proved the following dimensional reduction of Donaldson-Thomas theory on X to Vafa-Witten theory on S in algebraic geometry:

Fact 1. [26, Lemma 3.4 (i)] *The obstruction bundle over $I_0(X, d[E]) \cong \text{Hilb}^k S$ is isomorphic to the tangent bundle of $\text{Hilb}^k S$.*

As a consequence [26, Lemma 3.4 (ii)],

$$\tilde{N}_{0,k} = \int_{[I_0(X, k\beta_0)]^{\text{vir}}} 1 = \int_{\text{Hilb}^k S} e(T_{\text{Hilb}^k S}) = \chi(\text{Hilb}^k S).$$

We have

$$\sum_{k=0}^{\infty} \tilde{N}_{0,k} v^k = \sum_{k=0}^{\infty} \chi(\text{Hilb}^k S) v^k = \prod_{m>0} \frac{1}{(1 - v^m)^{\chi(S)}}.$$

3.4.2. GW theory. For any $g, k \geq 0$ the moduli space $\overline{\mathcal{M}}_{g,0}(X, k\beta_0)$ of genus g , degree $k\beta_0$ stable maps to X has virtual dimension zero. Define

$$N_{g,k} = \int_{[\overline{\mathcal{M}}_{g,0}(X, k\beta_0)]^{\text{vir}}} 1.$$

The moduli space $\overline{\mathcal{M}}_{g,0}(X, k\beta_0)$ can be identified with $\overline{\mathcal{M}}_{g,0}(E, k[E]) \times S$. The virtual dimension of $\overline{\mathcal{M}}_{g,0}(X, k\beta_0)$ is zero, and the virtual dimension of

$\overline{\mathcal{M}}_{g,0}(E, k[E])$ is $2g - 2$. Let

$$\pi_1 : \overline{\mathcal{M}}_{g,0}(E, k[E]) \times S \rightarrow \overline{\mathcal{M}}_{g,0}(E, k[E]), \quad \pi_2 : \overline{\mathcal{M}}_{g,0}(E, k[E]) \times S \rightarrow S$$

be projections to the first and second factors, respectively. Let

$$\mathbb{E} \rightarrow \overline{\mathcal{M}}_{g,0}(E, k[E])$$

be the Hodge bundle, and let \mathbb{E}^\vee denote its dual. Then

$$N_{g,k} = \int_{[\overline{\mathcal{M}}_{g,0}(X, k\beta_0)]^{\text{vir}}} 1 = \int_{[\overline{\mathcal{M}}_{g,0}(E, k[E])]^{\text{vir}} \times [S]} e(\pi_1^* \mathbb{E}^\vee \otimes \pi_2^* T_S)$$

Again, E acts on moduli spaces $\overline{\mathcal{M}}_{g,0}(X, k\beta_0)$, and the action has no fixed points unless $g = 1$. In this case, we have

$$N_{1,k} = \int_{[\overline{\mathcal{M}}_{1,0}(E, k[E])]^{\text{vir}}} 1 \cdot \int_S e(T_S) = \chi(S) N_{1,k}^E$$

where

$$N_{1,k}^E = \int_{[\overline{\mathcal{M}}_{1,0}(E, k[E])]^{\text{vir}}} 1$$

We have

$$\sum_{k=1}^{\infty} N_{1,k} v^k = \chi(S) \sum_{k=1}^{\infty} N_{1,k}^E v^k.$$

By [8, 19],

$$\exp \left(\sum_{k=1}^{\infty} N_{1,k}^E v^k \right) = \frac{1}{\prod_{m>0} (1 - v^m)}.$$

so

$$\exp \left(\sum_{k=1}^{\infty} N_{1,k} v^k \right) = \prod_{m>0} \frac{1}{(1 - v^m)^{\chi(S)}}.$$

3.4.3. Non-compact surfaces. The identity

$$(22) \quad \sum_{k=0}^{\infty} \tilde{N}_{0,k} v^k = \exp \left(\sum_{k=1}^{\infty} N_{1,k} v^k \right) = \prod_{m>0} \frac{1}{(1 - v^m)^{\chi(S)}}.$$

is also valid for certain noncompact surfaces. For example, suppose that S admits a \mathbb{C}^* action such that the fixed point set $S^{\mathbb{C}^*}$ is compact. Then

one can use the torus action to define residue DT invariants $\tilde{N}_{0,k}$ and GW invariants $N_{1,k}$. We have

$$N_{1,k} = \int_{[\overline{\mathcal{M}}_{1,0}(E,k[E])]^{\text{vir}}} 1 \cdot \int_{S^{\mathbb{C}^*}} \frac{e_{\mathbb{C}^*}(T_S)}{e_{\mathbb{C}^*}(N_{S^T/S})} = N_{1,k}^E \chi(S^{\mathbb{C}^*}) = N_{1,k}^E \chi(S).$$

For example, S can be the total space of a line bundle over a curve, or a resolution of ADE singularities.

3.4.4. Trivial elliptic fibration over a K3 surface. Let $X = S \times E$, where S be a projective K3 surface. Then any effective curve class $\beta \in H_2(X, \mathbb{Z})$ is of the form

$$\beta = \beta_1 + k\beta_0$$

where $\beta_0 = [E]$ as before, and β_1 is in the image of the injective map $H_2(S, \mathbb{Z}) \rightarrow H_2(X, \mathbb{Z})$ induced by the inclusion

$$S = S \times \{p_0\} \hookrightarrow S \times E = X$$

where p_0 is some point in E .

$\beta_1 = 0$: by Section 3.4.2, $N_{g,k\beta_0}^{S \times E} = 0$ unless $g = 1$, and

$$\exp\left(\sum_{k=0}^{\infty} N_{g=1,k\beta_0}^{S \times E} v^k\right) = \prod_{m>0} \frac{1}{(1-v^m)^{24}}.$$

$\beta_1 > 0$ (viewed as a class in $H_2(S, \mathbb{Z})$): we have (see e.g. [69, 75])

$$[\overline{\mathcal{M}}_{g,0}(S \times E, \beta_1 + k\beta_0)]^{\text{vir}} = 0,$$

so

$$N_{g,\beta_1+k\beta_0}^{S \times E} = 0.$$

Therefore, Gromov-Witten theory of $S \times E$ is almost trivial. However, $S \times E$ has very interesting *reduced* Gromov-Witten theory [75, 76].

3.5. Half K3

3.5.1. Construction. Let $[X_0, X_1, X_2]$ be homogeneous coordinates on \mathbb{P}^2 . Let $F_0(X_0, X_1, X_2)$ and $F_1(X_0, X_1, X_2)$ be two homogeneous polynomials

of degree 3 such that

$$D_0 = \{[X_0, X_1, X_2] \in \mathbb{P}^2 \mid F_0(X_0, X_1, X_2) = 0\},$$

$$D_1 = \{[X_0, X_1, X_2] \in \mathbb{P}^2 \mid F_1(X_0, X_1, X_2) = 0\}$$

are two smooth cubic curves intersecting transversally at 9 points q_1, \dots, q_9 . D_0, D_1 are elliptic curves.

Consider a pencil of elliptic curves in \mathbb{P}^2 containing D_0 and D_1 :

$$B_9 = \{([Z_0, Z_1], [X_0, X_1, X_2]) \in \mathbb{P}^1 \times \mathbb{P}^2 \mid Z_0 F_0(X_0, X_1, X_2) + Z_1 F_1(X_0, X_1, X_2) = 0\}.$$

Define $p_1 : \mathbb{P}^1 \times \mathbb{P}^2 \rightarrow \mathbb{P}^1$ and $p_2 : \mathbb{P}^1 \times \mathbb{P}^2 \rightarrow \mathbb{P}^2$ to be the projections to the first and second factors, respectively. And, let $\pi_1 : B_9 \rightarrow \mathbb{P}^1$ and $\pi_2 : B_9 \rightarrow \mathbb{P}^2$ be the restrictions of these projections. Then $\pi_1 : B_9 \rightarrow \mathbb{P}^1$ is an elliptic fibration, and $\pi_2 : B_9 \rightarrow \mathbb{P}^2$ is blowup of \mathbb{P}^2 at q_1, \dots, q_9 . The exceptional divisor $E_i = \pi_2^{-1}(q_i)$ is a section of $\pi_1 : B_9 \rightarrow \mathbb{P}^1$ and can be identified with $\mathbb{P}^1 \times \{q_i\} \subset \mathbb{P}^1 \times \mathbb{P}^2$.

3.5.2. Basic topology. Let $H_1, H_2 \in H^2(\mathbb{P}^1 \times \mathbb{P}^2; \mathbb{Z})$ be the pull back of the hyperplane class under p_1, p_2 , respectively. Then

$$H^*(\mathbb{P}^1 \times \mathbb{P}^2; \mathbb{Z}) = \mathbb{Z}[H_1, H_2]/(H_1^2, H_2^3).$$

Let $h_1, h_2 \in H^2(B_9; \mathbb{Z})$ be the restriction of H_1, H_2 respectively. Then h_1 is the Poincaré dual of the homology class of a fiber F of $\pi_1 : S \rightarrow \mathbb{P}^1$, and h_2 is the Poincaré dual of the homology class of a curve $E_0 \subset S$ such that $\pi_1|_{E_0} : E_0 \rightarrow \mathbb{P}^1$ is a degree 3 cover. We have $E_0 \cdot E_0 = 1$ and the genus of E_0 is zero. By adjunction formula, we have $c_1(T_{B_9}) = h_1$. So the pair (B_9, F) is relative Calabi-Yau in the sense that $K_{B_9} + F = 0$. In particular, $h^{2,0}(B_9) = h^{0,2}(B_9) = 0$.

We have

$$\chi(B_9) = \int_{B_9} c_2(T_{B_9}) = \int_{\mathbb{P}^1 \times \mathbb{P}^2} \frac{(1 + H_1)^2(1 + H_2)^3}{1 + H_1 + 3H_2} (H_1 + 3H_2) = 12.$$

Alternatively,

$$\chi(B_9) = \chi(\mathbb{P}^2) - 9\chi(\text{point}) + 9\chi(\mathbb{P}^1) = 3 - 9 + 18 = 12$$

By the Lefschetz hyperplane theorem, $H_1(B_9; \mathbb{Z}) = 0$. So the Hodge diamond of B_9 is given by:

$$\begin{array}{ccccc}
 & & & & 1 \\
 & & & 0 & & 0 \\
 & 0 & & 10 & & 0 \\
 & & 0 & & 0 & \\
 & & & & & 1
 \end{array}$$

For $i = 0, \dots, 9$, let $e_i \in H_2(B_9; \mathbb{Z})$ be the homology class represented by the curve E_i , respectively. Then

$$H_2(B_9; \mathbb{Z}) = \bigoplus_{i=0}^9 \mathbb{Z}e_i,$$

and the intersection form is given by

$$e_i \cdot e_j = g_{ij}, \quad g_{ij} = \text{diag}(1, -1, \dots, -1).$$

so that

$$H_2(B_9, \mathbb{Z}) = \Gamma^{1,9}$$

is the unique 10 dimensional odd unimodular lattice of signature $(1, 9)$.

In terms of e_i , the fiber class $[F]$ and the base class $[B]$ have the form

$$[F] = 3e_0 - \sum_{i=1}^9 e_i, \quad [B] = e_9$$

such that

$$[F] \cdot [B] = 1 \quad [B] \cdot [B] = -1 \quad [F] \cdot [F] = 0$$

We have an orthogonal decomposition

$$\Gamma^{1,9} = \Gamma^{1,1} \oplus E_8(-1)$$

where

- $\Gamma^{1,1}$ is the sublattice generated by $[F]$ and $[B]$, which is the unique 2 dimensional odd unimodular lattice of signature $(1, 1)$, and
- $E_8(-1)$ is the sublattice generated by $-e_0 + e_1 + e_2 + e_3, e_1 - e_2, e_2 - e_3, \dots, e_7 - e_8$, which is the negative E_8 lattice.

3.5.3. Local GW invariants from topological YM theory. Recall that $K_{B_9} = -F$. Suppose that

$$\beta \in H_2(X_{B_9}; \mathbb{Z}) \cong H_2(B_9; \mathbb{Z})$$

and $\beta \cdot [F] > 0$. Then

$$\overline{\mathcal{M}}_{g,0}(X_{B_9}, \beta) = \overline{\mathcal{M}}_{g,0}(B_9, \beta).$$

The virtual dimension of $\overline{\mathcal{M}}_{g,0}(X_{B_9}; \beta)$ is zero, while the virtual dimension of $\overline{\mathcal{M}}_{g,0}(B_9; \beta)$ is $\beta \cdot [F] + g - 1$. Define

$$(23) \quad N_{g,\beta}^{X_{B_9}} = \int_{[\overline{\mathcal{M}}_{g,0}(X_{B_9}, \beta)]^{\text{vir}}} 1 = \int_{[\overline{\mathcal{M}}_{g,0}(B_9, \beta)]^{\text{vir}}} e(V_{g,\beta})$$

where $V_{g,\beta}$ is a vector bundle over $\overline{\mathcal{M}}_{g,0}(B_9; \beta)$ whose fiber over $[f : C \rightarrow B_9]$ is

$$H^1(C, f^* \mathcal{O}_{B_9}(-F)).$$

Note that $H^0(C, f^* \mathcal{O}_S(-F)) = 0$ since $[C] \cdot (-[F]) < 0$. By Riemann-Roch, the rank of $V_{g,\beta}$ is $\beta \cdot [F] + g - 1$.

Introduce the Gromov-Witten potential,

$$\mathcal{F}(\lambda; t) := \sum_g \lambda^{2g-2} \mathcal{F}_g(t), \quad \mathcal{F}_g(t) = \sum_{\beta} N_{g,\beta}^{X_{B_9}} e^{2\pi i t \cdot \beta}$$

where $t = (\phi, \tau, \vec{m})$ are the Kähler moduli, such that ϕ and τ correspond to the base and fiber moduli, and $m_i, i = 1, \dots, 8$ denote the remaining E_8 moduli.

Since B_9 is elliptic, one can use the fiberwise T-duality to relate the counting of holomorphic curves on B_9 to the counting of instantons on B_9 . As a result, one finds a relation

$$(24) \quad \mathcal{F}_0(t) = \sum_{n=1}^{\infty} q^{n/2} Z_n(m_i; \tau) e^{2\pi i n \phi}$$

where $q = e^{2\pi i \tau}$ and $Z_n(m_i; \tau)$ is the partition function of the topological $\mathcal{N} = 4$ $U(n)$ Yang-Mills on B_9 . It is defined as a generating function

$$Z_n := \sum_k \chi(\mathcal{M}_k) q^k$$

where \mathcal{M}_k is the moduli space of instantons on S with instanton number k . Thus, (24) relates the counting of holomorphic curves in the class $\beta =$

$n[B] + k[F]$ to the counting of $U(n)$ instantons with instanton number k . Let us consider some explicit examples.

$n = 0$: In this case, the rank of the gauge group is zero and there are no Yang-Mills instantons. Therefore,

$$N_{g,\beta}^{X_{B_9}} = 0$$

for $\beta = k[F]$.

$n = 1$: In the abelian Yang-Mills theory, the only non-trivial instantons are point-like. The moduli space of point-like instantons of charge k can be identified with $\text{Hilb}^k B_9$, the Hilbert scheme of k points on the surface B_9 . This suggests $Z_1 \sim \eta(q)^{-1/12}$.

In general,

$$q^{1/2} Z_1(m_i; \tau) = \frac{\theta_{E_8}(\tau; m_i) q^{1/2}}{\eta(q)^{12}} = \theta_{E_8}(\tau; m_i) \prod_{m>0} \frac{1}{(1 - q^m)^{12}}.$$

The precise definition of $\theta_{E_8}(\tau; m_1, \dots, m_8)$ can be found in [41, section 2.6]. It can be written as

$$\theta_{E_8}(\tau; m_1, \dots, m_8) = \sum_{k_i \in \frac{1}{2}\mathbb{Z}} A_{k_1, \dots, k_8}(q) y_1^{k_1} \dots y_8^{k_8}$$

where $y_i = e^{2\pi i m_i}$, $A_{0, \dots, 0}(q) = 1$. Hence, one finds

$$(25) \quad \sum_{k=0}^{\infty} N_{0, [B]+k[F]}^{X_{B_9}} q^k = \prod_{m>0} \frac{1}{(1 - q^m)^{12}} = 1 + 12q + 90q^2 + 520q^3 + 2535q^4 + 10908q^5 + \dots$$

In fact, the duality prediction (25) has been verified mathematically. The rank of $V_{0, [B]+k[F]}$ in (23) is zero, so

$$N_{0, [B]+k[F]}^{X_{B_9}} = \int_{[\overline{\mathcal{M}}_{0,0}(B_9; [B]+k[F])]^{\text{vir}}} 1 = N_{0, [B]+k[F]}^{B_9}$$

Let $N_{g,n} = N_{g, [B]+(g+n)[F]}^{B_9}$, where $N_{g,\beta}^{B_9}$ is defined by (1). Bryan and Leung [14] showed that

$$\sum_{n=0}^{\infty} N_{g,n} q^n = \left(\sum_{k=1}^{\infty} k \left(\sum_{d|k} d \right) q^{k-1} \right)^g \prod_{m>0} \frac{1}{(1 - q^m)^{12}}.$$

The case $g = 0$ gives exactly the formula (25).

$n \geq 2$: In general, Z_n have the following structure [70]:

$$(26) \quad Z_n(0; \tau) = \eta^{-12n} P_{6n-2}(E_2(\tau), E_4(\tau), E_6(\tau))$$

where P_{6n-2} is a quasi-homogeneous polynomial of weight $6n - 2$, and E_{2m} are the Eisenstein modular forms of weight $2m$.

In general $q^{n/2} Z_n(0; \tau)$ gives us

$$(27) \quad \sum_k \left(\sum_{\beta \cdot [B]=k, \beta \cdot [F]=n} N_{0, \beta}^{X_{B_9}} \right) q^k.$$

instead of

$$(28) \quad \sum_{k=0}^{\infty} N_{0, n[B]+k[F]}^{X_{B_9}} q^k$$

For example,

$$(29) \quad q^{1/2} Z_1(0; \tau) = \frac{\theta_{E_8}(\tau; 0) q^{1/2}}{\eta(q)^{12}} = \frac{E_4(\tau)}{\prod_{m>0} (1 - q^m)}$$

$$= 1 + 12q + 330q^2 + 3400q^3 + 26295q^4 + 161628q^5 + \dots$$

which is different from (25).

We have

$$(30) \quad qZ_2(0; \tau) = -\frac{1}{8} + \frac{18441}{2}q + 673760q^2 + \frac{82133595}{4}q^3 + \dots$$

Gromov-Witten invariants of X_{B_9} , i.e. local Gromov-Witten invariants of the half K3, are studied in [39], [82], [47], etc.

3.6. Dimensional reduction of DT theory on a local surface

A closely related problem appears when we study the total space, X_S , of the canonical line bundle of a complex algebraic surface S . We will be mainly interested in enumerative invariants of X_S when S is a Fano surface. In this case, any non-constant holomorphic map to X_S factors through the zero section $S \rightarrow X_S$, so the Gromov-Witten invariants of X_S can be viewed as local Gromov-Witten invariants of S in a Calabi-Yau three-fold. Similarly, Donaldson-Thomas gauge theory on X_S effectively reduces to a four-dimensional gauge theory on S .

3.6.1. The reduced deformation complex. In order to see this more explicitly, let us consider the deformation complex (9) for the total space X_S , of the canonical line bundle of S . Let v be a local complex coordinate on the fiber of X_S . Then, a (p, q) -form on X_S which is constant along the fiber locally can be written as

$$\omega^{(p,q)} = \tilde{\omega}^{(p,q)} + \tilde{\omega}^{(p,q-1)} \wedge d\bar{v} + \tilde{\omega}^{(p-1,q)} \wedge dv + \tilde{\omega}^{(p-1,q-1)} \wedge dv \wedge d\bar{v}$$

where we use $\tilde{\omega}^{(p,q)}$ to denote a (pull-back) of a (p, q) -form on S . Let $\hat{\Omega}^{0,k}$ be the space of $(0, k)$ -forms on X_S which are constant along the fiber. Then it decomposes into the following spaces of anti-holomorphic forms on S :

$$(31) \quad \begin{aligned} \hat{\Omega}^{0,3} &= \Omega_S^{0,2} \\ \hat{\Omega}^{0,k} &= \Omega_S^{0,k} \oplus \Omega_S^{0,k-1} \quad k = 1, 2 \\ \hat{\Omega}^{0,0} &= \Omega_S^{0,0} \end{aligned}$$

Hence, in terms of $(0, k)$ -forms on S , the complex (9) reads

$$(32) \quad 0 \rightarrow \Omega^{0,0} \rightarrow \Omega^{0,0} \oplus \Omega^{0,1} \oplus \Omega^{0,2} \rightarrow \Omega^{0,1} \oplus \Omega^{0,2} \rightarrow 0$$

We now describe the differentials in (32). We assume that the fields are constant along the fiber of $X_S \rightarrow S$. Then $A \in \Omega^1_{X_S}(\text{End}\mathcal{E})$ can be decomposed as

$$(33) \quad A = A_2 + Cdv + \bar{C}d\bar{v},$$

where v is the local coordinate in the fiber of $X \rightarrow S$,

$$A_2 \in \Omega^1_S(\text{End}\mathcal{E}), \quad C \in \Omega^0_S(\text{End}\mathcal{E} \otimes K_S^{-1}), \quad \bar{C} \in \Omega^0_S(\text{End}\mathcal{E} \otimes K_S).$$

and $\bar{\varphi}$ can be written as

$$(34) \quad \bar{\varphi} = \bar{\varphi}_2 \wedge d\bar{v}$$

where $\bar{\varphi}_2 \in \Omega^{0,2}_S(\text{End}\mathcal{E} \otimes K_S)$.

Equations (7) can be written as

$$(35) \quad F_{A_2}^{0,2} = [C, \bar{\varphi}_2]$$

$$(36) \quad \bar{\partial}_{A_2} \bar{C} = \bar{\partial}^\dagger_{A_2} \bar{\varphi}_2$$

$$(37) \quad \text{Tr}_\omega F_{A_2}^{1,1} + [C, \bar{C}] + *[\varphi_2, \bar{\varphi}_2] = \ell I.$$

where ω is the Kähler form on S .

Let $V = \text{End}\mathcal{E}$. The deformation complex at (C, A_2, φ_2) is given by

$$0 \rightarrow \Omega_S^{0,0}(V) \xrightarrow{D_1} \Omega_S^{0,0}(V \otimes K_S) \oplus \Omega_S^{0,1}(V) \oplus \Omega_S^{0,2}(V \otimes K_S) \\ \xrightarrow{D_2} \Omega_S^{0,1}(V \otimes K_S) \oplus \Omega_S^{0,2}(V) \rightarrow 0$$

where

$$D_1(\eta) = ([\bar{C}, \eta], \bar{\partial}_{A_2}\eta, [\bar{\varphi}_2, \eta]) \\ D_2(\delta\bar{C}, \delta A_2^{0,1}, \delta\bar{\varphi}_2) = \left([\bar{C}, \delta A_2^{0,1}] + \bar{\partial}_{A_2}(\delta\bar{C}) - \bar{\partial}_{A_2}^\dagger(\delta\bar{\varphi}_2), \bar{\partial}_{A_2}(\delta A_2^{0,1}) - [C, \delta\bar{\varphi}_2]\right)$$

This complex has to be identified with a deformation complex in a suitable four-dimensional gauge theory.

3.6.2. Three twists of $\mathcal{N} = 4$ super-Yang-Mills. There are three different twists of $\mathcal{N} = 4$ super-Yang-Mills in four dimensions, which can be conveniently characterized by the following homomorphisms [93]:

- 1) $SU(2)_L \times SU(2)_R \times SU(4)_I \rightarrow SU(2)'_L \times SU(2)_R \times SU(2)_F$, under which $\mathbf{4} \rightarrow (\mathbf{2}, \mathbf{1}) \oplus (\mathbf{2}, \mathbf{1})$
- 2) $SU(2)_L \times SU(2)_R \times SU(4)_I \rightarrow SU(2)'_L \times SU(2)_R \times SU(2)_F \times U(1)$, so that $\mathbf{4} \rightarrow (\mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{1})$
- 3) $SU(2)_L \times SU(2)_R \times SU(4)_I \rightarrow SU(2)'_L \times SU(2)'_R \times U(1)$, so that $\mathbf{4} \rightarrow (\mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2})$

and which relate the rotation symmetry and R-symmetry in the original theory and its twisted (topological) version. The first twist leads to a TQFT known as the Vafa-Witten theory [93]. The second is the Donaldson-Witten twist that leads to non-abelian monopole equations [52]. And the last twist is sometimes called Marcus twist [63] or GL twist, for its connection to the geometric Langlands program [51].

These three TQFTs involve (though in a somewhat different way) certain moduli spaces: 1) moduli space of self-dual gauge connections; 2) moduli space of adjoint non-abelian monopoles; 3) moduli space of complexified flat gauge connections. The second theory based on non-abelian monopole equations has only one topological supersymmetry, $\mathcal{N}_T = 1$. On the other hand, theories 1) and 3) have $\mathcal{N}_T = 2$ topological supersymmetry.

For example, in the Vafa-Witten $\mathcal{N} = 4$ twisted gauge theory, the deformation complex is [93]:

$$(38) \quad 0 \rightarrow \Omega_S^0(\text{ad}P) \rightarrow \Omega_S^0(\text{ad}P) \oplus \Omega_S^1(\text{ad}P) \oplus \Omega_S^{2,+}(\text{ad}P) \\ \rightarrow \Omega_S^1(\text{ad}P) \oplus \Omega_S^{2,+}(\text{ad}P) \rightarrow 0$$

For comparison, let us write the deformation complex in the generalization of $\mathcal{N} = 2$ Seiberg-Witten theory, where the monopole fields take values in $\Gamma(S, S^+ \otimes E)$,

$$0 \rightarrow \Omega^0(g_E) \rightarrow \Omega^1(g_E) \oplus \Gamma(S, S^+ \otimes E) \rightarrow \Omega^{2,+}(g_E) \oplus \Gamma(S, S^- \otimes E) \rightarrow 0$$

Here, g_E denotes the representation of the Lie algebra $g = \text{Lie}(G)$ associated to a representation R . If $P \rightarrow S$ is the principal G -bundle over S , and V is a vector space associated with representation R , we can form the associated vector bundle $E = P \times_G V$.

In order to compare the deformation complex of the six-dimensional gauge theory on X_S with deformation complexes in familiar four-dimensional topological gauge theories (1)-(3), we now assume that S is Kähler, and use the famous result of Donaldson. Namely, for a bundle E over a Kahler surface S , we can identify the moduli space of irreducible ASD connections with the set of equivalence classes of stable holomorphic bundles \mathcal{E} which are topologically equivalent to E . We fix the holomorphic line bundle $\det \mathcal{E}$ so that $G = SU(n)$. There are natural isomorphisms

$$(39) \quad \Omega^1(g_E) \cong \Omega^{0,1}(\text{End}_0 \mathcal{E}) \\ \Omega^0(g_E) \oplus \Omega^{2,+}(g_E) \cong \Omega^0(\text{End}_0 \mathcal{E}) \oplus \Omega^{0,2}(\text{End}_0 \mathcal{E})$$

where $\text{End}_0 \mathcal{E}$ denotes the trace-free endomorphisms of the stable holomorphic vector bundle \mathcal{E} . Using such isomorphisms, we can write the deformation complex (32) as

$$0 \rightarrow \Omega^0 \rightarrow \Omega^0 \oplus \Omega^1 \oplus \Omega^{2,+} \rightarrow \Omega^1 \oplus \Omega^{2,+} \rightarrow 0$$

where elements of all these spaces are valued in g_E associated with $\text{End}_0(\mathcal{E})$. Now, this complex does look like the deformation complex (38) in Vafa-Witten theory!

On a Kähler surface S , we further have isomorphisms

$$\begin{aligned} S^+ &= \Omega^{0,0} \oplus \Omega^{0,2} \\ S^- &= \Omega^{0,1} \\ \Omega_{\mathbb{C}}^{2,+} &= \Omega^{2,0} \oplus \Omega^0 \cdot \omega \oplus \Omega^{0,2} \end{aligned}$$

As pointed out in [90], there is a simple way to relate the Vafa-Witten theory on S and the Donaldson-Thomas theory on X_S . Given a stable bundle \mathcal{E} on S , let $i_*\mathcal{E}$ be the pushforward torsion sheaf on X_S supported on S . This gives an inclusion $i_* : \mathcal{M}(S) \rightarrow \mathcal{M}(X_S)$, whose image is a connected component of $\mathcal{M}(X_S)$. The restriction of the deformation theory on $\mathcal{M}(X_S)$ to $i_*\mathcal{M}(S)$ coincides with that on $\mathcal{M}(S)$.

4. Dimensional reduction of DT theory to 2d σ -model

4.1. Topological reduction

The second duality we consider is similar to the topological reduction of four-dimensional gauge theory studied in [9]. Namely, we consider X to be of the form $X = \Sigma_\ell \times S$, where Σ_ℓ is a Riemann surface of genus ℓ and S is a Kähler surface. Let g_Σ and g_S be the metrics of Σ and of S , respectively. We first consider a general rescaling

$$g_\Sigma \rightarrow t^p g_\Sigma, \quad g_S \rightarrow t^q g_S$$

where t is a positive number which tends to zero. Then the Lagrangian density of the Yang-Mills action scales as

$$\begin{aligned} F_X \wedge *F_X &\rightarrow t^p F_S \wedge *F_S + 2t^q (d_S A_\Sigma - D_\Sigma A_S) \wedge *(d_S A_\Sigma - D_\Sigma A_S) \\ &\quad + t^{-p+2q} F_\Sigma \wedge *F_\Sigma. \end{aligned}$$

When $(p, q) = (0, 1)$, we have $g_\Sigma \rightarrow g_\Sigma$, $g_S \rightarrow t g_S$ (S is “small”), and

$$\begin{aligned} F_X \wedge *F_X &\rightarrow F_S \wedge *F_S + 2t (d_S A_\Sigma - D_\Sigma A_S) \wedge *(d_S A_\Sigma - D_\Sigma A_S) \\ &\quad + t^2 F_\Sigma \wedge *F_\Sigma. \end{aligned}$$

The resulting theory is the 4d Vafa-Witten theory discussed in Section 3.

This section concerns the case $(p, q) = (-1, 0)$. In this case, we have

$$\begin{aligned}
 g_\Sigma &\rightarrow \frac{1}{t}g_\Sigma, & g_S &\rightarrow g_S \quad (\Sigma \text{ is "large"}), \\
 F_X \wedge *F_X &\rightarrow \frac{1}{t}F_S \wedge *F_S + 2(d_S A_\Sigma - D_\Sigma A_S) \wedge *(d_S A_\Sigma - D_\Sigma A_S) \\
 &+ tF_\Sigma \wedge *F_\Sigma.
 \end{aligned}$$

As in [9], the resulting theory on Σ_ℓ is $\mathcal{N} = 2$ topological sigma-model, whose target space is the moduli space of solutions to the equations in the *four-dimensional* gauge theory on S . As we explained in Section 3, these are anti-self-dual gauge fields on S (when vanishing theorems hold). Therefore, after the topological reduction we obtain $\mathcal{N} = 2$ sigma-model on Σ_ℓ with the target space $\mathcal{M}_{ASD}(S)$.

Since the six-dimensional theory is topological, it should not matter whether we reduce on S or on Σ_ℓ . This means that the partition function $Z_{DT}(X)$ should be equal to the partition function of the four-dimensional gauge theory on S , as well as to the partition function of the $\mathcal{N} = 2$ sigma-model on Σ_ℓ ,

$$Z_{DT}(X) = Z_{4D}(S) = Z_{2D}(\Sigma_\ell)$$

4.2. Moduli spaces of instantons and correlation functions

We consider a general gauge group G which can be any compact connected Lie group. Let $\mathcal{M}_{ASD}(S, G)$ denote the moduli space of gauge equivalence classes of ASD G -connections on S , which are solutions to (18). Let $\mathcal{M}_{HYM}(\Sigma_\ell \times S, G)$ denote the moduli space of gauge equivalence classes of HYM G -connections on $\Sigma_\ell \times S$, which are solutions to (5). Each of the spaces $\mathcal{M}_{ASD}(S, G)$ and $\mathcal{M}_{HYM}(\Sigma_\ell \times S, G)$ can be represented as a union of components labelled by the topological type of the underlying topological principal G -bundle.

When G is semi-simple, the ASD equation (18) is equivalent to the 2-dimensional version of the HYM equations (5). When G is not semi-simple, there is a topological obstruction to the existence of ASD connections. Let H be the connected component of $Z(G)$, the center of G . Then H is a compact torus. Let $G_{ss} = [G, G]$ be the commutator subgroup, which is also the maximal connected semisimple subgroup of G . There is a finite cover $H \times G_{ss} \rightarrow G$ given by $(h, g) \mapsto hg$. This is a group homomorphism. The kernel is isomorphic to $D = H \cap G_{ss}$ which is a finite subgroup of H . Note that G_{ss} is a normal subgroup of G . Consider the exact sequence of abelian

groups

$$1 \rightarrow \pi_1(G_{ss}) \rightarrow \pi_1(G) \rightarrow \pi_1(G/G_{ss}) \rightarrow 1$$

where $G/G_{ss} = H/D \cong \mathbb{Z}^{\oplus \dim_{\mathbb{R}} H}$. We have

$$\dots \rightarrow H^2(S, \pi_1(G_{ss})) \rightarrow H^2(S, \pi_1(G)) \xrightarrow{\deg} H^2(S, \mathbb{Z})^{\oplus \dim_{\mathbb{R}} H} \rightarrow \dots$$

Let $m = o_2(P) \in H^2(S, \pi_1(G))$ be the magnetic flux of a principal G -bundle P , so that $\deg(m) = (m_1, \dots, m_r)$, where $m_i \in H^2(S; \mathbb{Z})$ and $r = \dim_{\mathbb{R}} H$. Then P admits an ASD connection only if $m_i \cdot [k_0] = 0$, where k_0 is the Kähler class. Given $c \in H^2(S; \mathbb{Z}) \setminus \{0\}$, let

$$\Gamma_c = \{k \in H^{1,1}(S) \mid c \wedge k \wedge k = 0\}.$$

Then Γ_c is a real codimension-1 subspace of $H^{1,1}(S)$. Choose k_0 such that

$$[k_0] \notin \bigcup_{c \in H^2(S; \mathbb{Z}), c \neq 0} \Gamma_c$$

Then $m_i \cdot [k_0] = 0$ implies $m_i = 0$. In this case P admits an ASD connection only if

$$\deg(P) \stackrel{\text{def}}{=} \deg(o_2(P)) = 0 \in H^2(S; \mathbb{Z})^{\oplus r}.$$

For example, when $G = U(N)$, we have $G_{ss} = SU(N)$ and $o_2(P) = \deg(P) = c_1(P)$. When $G = SO(N)$ ($N > 2$), we have $G_{ss} = SO(N)$, $o_2(P) = w_2(P)$, and $\deg(P) = 0$.

The complexification $G^{\mathbb{C}}$ of G is a connected reductive algebraic group over \mathbb{C} . The moduli space $\mathcal{M}_{\text{ASD}}(S, G)$ can be identified with the moduli space $\mathcal{M}(S, G^{\mathbb{C}})$ of S -equivalence classes of semi-stable holomorphic principal $G^{\mathbb{C}}$ -bundles of degree zero on S [24], and the moduli space $\mathcal{M}_{\text{HYM}}(\Sigma_{\ell} \times S, G)$ can be identified with the moduli space $\mathcal{M}(\Sigma_{\ell} \times S, G^{\mathbb{C}})$ of S -equivalence classes of semi-stable holomorphic principal $G^{\mathbb{C}}$ -bundles over $\Sigma_{\ell} \times S$ [1, 91].

The partition function of the $\mathcal{N} = 2$ sigma-model on Σ_{ℓ} localizes on $\text{Mor}(\Sigma_{\ell}, \mathcal{M}(S, G^{\mathbb{C}}))$, the space of holomorphic maps from Σ_{ℓ} to $\mathcal{M}(S, G^{\mathbb{C}})$. The partition function of the Donaldson-Thomas theory on $\Sigma_{\ell} \times S$ localizes on $\mathcal{M}(\Sigma_{\ell} \times S, G^{\mathbb{C}})$. We now construct a map between the two moduli spaces $\text{Mor}(\Sigma_{\ell}, \mathcal{M}(S, G^{\mathbb{C}}))$ and $\mathcal{M}(\Sigma_{\ell} \times S, G^{\mathbb{C}})$. Let $\mathcal{P} \rightarrow \mathcal{M}(S, G^{\mathbb{C}}) \times S$ be the universal principal $G^{\mathbb{C}}$ -bundle. (It does not exist as a usual principal bundle, see e.g. [51, Section 7.1].) Given a holomorphic map $u : \Sigma_{\ell} \rightarrow \mathcal{M}(S, G^{\mathbb{C}})$, we get a holomorphic principal $G^{\mathbb{C}}$ -bundle $(u \times \text{id}_S)^* \mathcal{P}$ over $\Sigma_{\ell} \times S$. We get a

map

$$(40) \quad \Phi : \text{Mor}(\Sigma_\ell, \mathcal{M}(S, G^{\mathbb{C}})) \longrightarrow \mathcal{M}(\Sigma_\ell \times S, G^{\mathbb{C}}), \quad u \mapsto (u \times \text{id}_S)^*\mathcal{P}$$

which is birational.

Let $\mathcal{O}_1, \dots, \mathcal{O}_k$ be observables in the σ -model on Σ_ℓ . Our topological reduction implies that there are corresponding observables $\tilde{\mathcal{O}}_1, \dots, \tilde{\mathcal{O}}_k$ in the DT theory on $\Sigma_\ell \times S$ such that

$$(41) \quad \langle \mathcal{O}_1 \cdots \mathcal{O}_k \rangle = \langle \tilde{\mathcal{O}}_1 \cdots \tilde{\mathcal{O}}_k \rangle$$

Geometrically, these correlation functions should be interpreted as intersection numbers on moduli spaces $\text{Mor}(\Sigma_\ell, \mathcal{M}(S, G^{\mathbb{C}}))$ and $\mathcal{M}(\Sigma_\ell \times S, G^{\mathbb{C}})$. To define these intersection numbers, we need to compactify these moduli spaces.

When S is a projective surface, the moduli spaces $\mathcal{M}(S, G^{\mathbb{C}})$ and $\mathcal{M}(\Sigma_\ell \times S, G^{\mathbb{C}})$ can be compactified using algebraic geometry. When $G = U(N)$, $G^{\mathbb{C}} = GL(N, \mathbb{C})$, these moduli spaces can be identified with moduli spaces of isomorphism classes of polystable holomorphic vector bundles, and the compactified moduli spaces are obtained by including semistable torsion free sheaves. For general G , the compactified moduli spaces $\overline{\mathcal{M}}(S, G^{\mathbb{C}})$ and $\overline{\mathcal{M}}(\Sigma_\ell \times S, G^{\mathbb{C}})$ are obtained by including semistable *principal sheaves* [32] or *singular principal bundles* [83]. These compactified moduli spaces are projective but may have singularities.

To compactify $\text{Mor}(\Sigma_\ell, \mathcal{M}(S, G^{\mathbb{C}}))$, we first compactify the target and get a partial compactification $\text{Mor}(\Sigma_\ell, \overline{\mathcal{M}}(S, G^{\mathbb{C}}))$ which consists of morphisms from Σ_ℓ to $\overline{\mathcal{M}}(S, G^{\mathbb{C}})$. The final compactification $\overline{\text{Mor}}(\Sigma_\ell, \overline{\mathcal{M}}(S, G^{\mathbb{C}}))$ is obtained by including stable maps from a curve with a root component isomorphic to Σ_ℓ and bubble components which are spheres. Ideally, (40) extends to a morphism $\Phi : \overline{\text{Mor}}(\Sigma_\ell, \overline{\mathcal{M}}(S, G^{\mathbb{C}})) \longrightarrow \overline{\mathcal{M}}(\Sigma_\ell \times S, G^{\mathbb{C}})$. The inverse is a rational map: a semi-stable $G^{\mathbb{C}}$ -bundle $P \rightarrow \Sigma_\ell \times S$ representing an element in $\mathcal{M}(\Sigma_\ell \times S, G^{\mathbb{C}})$ defines a morphism $\Sigma_\ell \rightarrow \overline{\mathcal{M}}(S, G^{\mathbb{C}})$.

$$\tilde{\mathcal{O}}_i \in H^*(\overline{\mathcal{M}}(\Sigma_\ell \times S, G^{\mathbb{C}})), \quad \mathcal{O}_i = \Phi^* \tilde{\mathcal{O}}_i \in H^*(\overline{\text{Mor}}(\Sigma_\ell, \overline{\mathcal{M}}(S, G^{\mathbb{C}}))).$$

The correlation functions (41) can be interpreted as intersection numbers of the cohomology classes \mathcal{O}_i 's and $\tilde{\mathcal{O}}_i$'s. For a singular moduli space one needs a virtual fundamental class to define intersection theory. The topological reduction suggests that there exist virtual fundamental classes such that

$$\Phi_* [\overline{\text{Mor}}(\Sigma_\ell, \overline{\mathcal{M}}(S, G^{\mathbb{C}}))]^{\text{vir}} = [\overline{\mathcal{M}}(\Sigma_\ell \times S, G^{\mathbb{C}})]^{\text{vir}}.$$

Then

$$(42) \quad \int_{[\overline{\text{Mor}}(\Sigma_\ell, \overline{\mathcal{M}}(S, G^{\mathbb{C}}))]^{\text{vir}}} \mathcal{O}_1 \cdots \mathcal{O}_k = \int_{[\overline{\mathcal{M}}(\Sigma_\ell \times S, G^{\mathbb{C}})]^{\text{vir}}} \tilde{\mathcal{O}}_1 \cdots \tilde{\mathcal{O}}_k.$$

The left and right hand sides of (42) can be viewed as the mathematical definition of the left and right hand sides of (41), respectively.

Here we propose some natural observables such that (42) is expected to hold. Given a point $p \in \Sigma_\ell$, define

$$\begin{aligned} \text{ev}_p : \overline{\text{Mor}}(\Sigma_\ell, \overline{\mathcal{M}}(S, G^{\mathbb{C}})) &\longrightarrow \overline{\mathcal{M}}(S, G^{\mathbb{C}}) \\ &u \mapsto u(p) \\ r_p : \overline{\mathcal{M}}(\Sigma_\ell \times S, G^{\mathbb{C}}) &\longrightarrow \overline{\mathcal{M}}(S, G^{\mathbb{C}}) \\ &P \mapsto P|_{\{p\} \times S} \end{aligned}$$

Then $\text{ev}_p = r_p \circ \Phi$. Given $\alpha_1, \dots, \alpha_k \in H^*(\overline{\mathcal{M}}(S, G^{\mathbb{C}}))$ and k distinct points $p_1, \dots, p_k \in \Sigma_\ell$, define

$$\begin{aligned} \langle \alpha_1 \cdots \alpha_k \rangle_{\sigma|_{ASD}}^{\Sigma_\ell|S} &= \int_{[\overline{\text{Mor}}(\Sigma_\ell, \overline{\mathcal{M}}(S, G^{\mathbb{C}}))]^{\text{vir}}} \text{ev}_{p_1}^* \alpha_1 \cup \cdots \cup \text{ev}_{p_k}^* \alpha_k \\ \langle \alpha_1 \cdots \alpha_k \rangle_{DT}^{\Sigma_\ell \times S} &= \int_{[\overline{\mathcal{M}}(\Sigma_\ell \times S, G^{\mathbb{C}})]^{\text{vir}}} r_{p_1}^* \alpha_1 \cup \cdots \cup r_{p_k}^* \alpha_k \end{aligned}$$

The definition is independent of choices of p_1, \dots, p_k . The correlation functions $\langle \alpha_1 \cdots \alpha_k \rangle_{\sigma|_{ASD}}^{\Sigma_\ell|S}$ are k -point *mixed*⁷ Gromov-Witten invariants of $\overline{\mathcal{M}}(S, G^{\mathbb{C}})$. We should have

$$(43) \quad \langle \alpha_1 \cdots \alpha_k \rangle_{\sigma|_{ASD}}^{\Sigma_\ell|S} = \langle \alpha_1 \cdots \alpha_k \rangle_{DT}^{\Sigma_\ell \times S}$$

4.3. Abelian theory

In the abelian theory,

$$\mathcal{M}_{ASD}(S) \cong \text{Hilb}^k S$$

which is a resolution of the symmetric product orbifold, $\text{Sym}^k S$. In this case, DT theory on $\Sigma_\ell \times S$ is related to GW theory on $\Sigma_\ell \times S$ by the GW/DT correspondence conjectured in [65, 66]. These lead to the equivalence of the following theories:

- S1 Gromov-Witten theory on $\Sigma_\ell \times S$.

⁷since we fix the complex structure of the domain

S2 Donaldson-Thomas theory on $\Sigma_\ell \times S$.

S3 Mixed Gromov-Witten invariants with domain Σ_ℓ and target $\text{Hilb}^k S$.

We now give a more precise conjecture relating S2 and S3. We first introduce correlation functions in S3. Classical cohomology ring of $\text{Hilb}^k S$ is determined first for \mathbb{C}^2 in [54, 94] and then for K3 [55]. The general case is treated in [18]. As vector spaces, there is an isomorphism:

$$\mathcal{F} = \text{Sym}^* \left(t^{-1} \mathbb{Q}[t^{-1}] \otimes H^*(S; \mathbb{Q}) \right) \cong \bigoplus_{k \geq 0} H^*(\text{Hilb}^k S; \mathbb{Q}).$$

A basis of \mathcal{F} is given by *cohomology weighted partitions*

$$\alpha_{-\mu_1}(\gamma_{i_1}) \cdots \alpha_{-\mu_n}(\gamma_{i_n} \mid 0).$$

where $(\mu_1, \mu_2, \dots, \mu_\ell)$ is a partition, and $\{\gamma_i\}$ is a basis of the cohomology group $H^*(S; \mathbb{Q})$.

Assume that $3\ell - 3 + n \geq 0$, so that the moduli space $\overline{\mathcal{M}}_{\ell,n}$ of genus ℓ , n -pointed stable curves is nonempty. We fix n distinct points x_1, \dots, x_n on Σ_ℓ , and let $\xi \in H^{6\ell-6+2n}(\overline{\mathcal{M}}_{\ell,n}; \mathbb{Q})$ be the Poincaré dual of the point class $[(\Sigma_\ell, x_1, \dots, x_n)] \in \overline{\mathcal{M}}_{\ell,n}$. Given $\beta \in H_2(\text{Hilb}^k S, \mathbb{Z})$, let $\pi: \overline{\mathcal{M}}_{\ell,n}(\text{Hilb}^k S, \beta) \rightarrow \overline{\mathcal{M}}_{\ell,n}$ be the forgetful map. We have the following genus ℓ , n -point, mixed Gromov-Witten invariants:

$$\langle (\mu^1, \delta^1), \dots, (\mu^n, \delta^n) \rangle_{\xi, \beta} = \int_{[\overline{\mathcal{M}}_{\ell,n}(\text{Hilb}^k S, \beta)]^{\text{vir}}} \pi^* \xi \cup \prod_{i=1}^n \text{ev}_i^*(\mu^i, \delta^i)$$

where each μ^i is a partition of k , and $\delta^i \in H^*(S; \mathbb{Q})^{\otimes \ell(\mu^i)}$.

Let $\{D_1, \dots, D_m\}$ be a basis of $H^2(S; \mathbb{Z})$. Then

$$\begin{aligned} \tilde{D}_0 &= \alpha_{-2}(1) (\alpha_{-1}(1))^{k-1} \mid 0, \\ \tilde{D}_i &= \alpha_{-1}(D_i) (\alpha_{-1}(1))^{k-1} \mid 0, \quad i = 1, \dots, m \end{aligned}$$

form a basis of $H^2(\text{Hilb}^k(S); \mathbb{Z})$. Define

$$\begin{aligned} (44) \quad & \langle (\mu^1, \delta^1), \dots, (\mu^k, \delta^n) \rangle_{\xi}^{\text{Hilb}^k(S)} \\ &= \sum_{\beta \in H_2(\text{Hilb}^k(S); \mathbb{Z})} \sum_{\beta} \langle (\mu^1, \delta^1), \dots, (\mu^k, \delta^n) \rangle_{\xi, \beta} \tilde{q}_0^{\beta \cdot \tilde{D}_0} \prod_{j=1}^m \tilde{q}_j^{\beta \cdot \tilde{D}_j} \end{aligned}$$

We next introduce correlation functions in S2. We fix n distinct points x_1, \dots, x_n on Σ_ℓ as before. Given $\beta \in H_2(S; \mathbb{Z})$, there are maps

$$\begin{aligned} \epsilon_i &: I_s(\Sigma_\ell \times S / \{x_1, \dots, x_n\} \times S, k[\Sigma_\ell] + \beta) \\ &\rightarrow \text{Hilb}^k(\{x_i\} \times S) \cong \text{Hilb}^k(S, \mathbb{Z}), \quad i = 1, \dots, n. \end{aligned}$$

Let (μ^i, δ^i) be cohomology weighted partitions as above. Define

$$\begin{aligned} &\langle (\mu^1, \delta^1), \dots, (\mu^n, \delta^n) \rangle_{s, \beta}^{DT(\xi, S)} \\ &= \int_{[I_s(\Sigma_\ell \times S / \{x_1, \dots, x_n\} \times S, k[\Sigma_\ell] + \beta)]^{\text{vir}}} \prod_{i=1}^n \epsilon_i^*(\mu^i, \delta^i). \\ (45) \quad &\langle (\mu^1, \delta^1), \dots, (\mu^n, \delta^n) \rangle_k^{DT(\xi, S)} \\ &= q^{-k} \sum_{s \in \mathbb{Z}} \sum_{\beta \in H_2(S, \mathbb{Z})} \langle (\mu^1, \delta^1), \dots, (\mu^n, \delta^n) \rangle_{s, \beta}^{DT(\xi, S)} q^s \prod_{j=1}^m q_j^{\beta \cdot D_j} \end{aligned}$$

Conjecture 1 (Maulik-Oblomkov). *Let S be a hyper-Kähler surface. Then*

$$(46) \quad \langle (\mu^1, \delta^1), \dots, (\mu^n, \delta^n) \rangle_\xi^{\text{Hilb}^k(S)} = \langle (\mu^1, \delta^1), \dots, (\mu^n, \delta^n) \rangle_k^{DT(\xi, S)}$$

Maulik and Oblomkov’s original conjecture concerns the genus zero case (i.e. $\Sigma_\ell = \mathbb{P}^1$), but the general case should follow from the genus 0 case by degenerating Σ_ℓ to a union of \mathbb{P}^1 ’s. Maulik and Oblomkov have checked that Conjecture 1 fails for Enriques surfaces.

Conjecture 2 (Maulik-Oblomkov). *Let S be a Fano surface, and let $\Sigma_\ell = \mathbb{P}^1$. Let $c_1(S) = \sum_{j=1}^m c_j D_j$. Then*

$$(47) \quad \langle (\mu^1, \delta^1), \dots, (\mu^n, \delta^n) \rangle_\xi^{\text{Hilb}^k(S)} = \langle (\mu^1, \delta^1), \dots, (\mu^n, \delta^n) \rangle_k^{DT(\xi, S)}$$

with $\tilde{q}_0 = q$, $\tilde{q}_j = (1 - q^{-1})^{c_j} q_j$.

Maulik and Oblomkov have checked Conjecture 2 for $\mathbb{P}^1 \times \mathbb{C}$ and $\mathcal{O}_{\mathbb{P}^1}(-1) \rightarrow \mathbb{P}^1$.

We summarize some known results which motivated the above conjectures. When the genus $\ell = 0$, then 3-point correlation functions in $Z_{2D}(\Sigma_0) = Z_{2D}(\mathbb{P}^1)$ are the structure constants of the quantum cohomology ring of $\mathcal{M}_{ASD}(S)$. Here we consider small quantum cohomology, so the structure

constants of the quantum product involves 3-point genus zero Gromov-Witten invariants. The big quantum cohomology involves n -point genus zero Gromov-Witten invariants, which are not the same as the n -point mixed Gromov-Witten invariants. The crepant resolution conjecture asserts that the quantum cohomology ring of $\text{Hilb}^k S$ is isomorphic to the orbifold quantum cohomology ring of $\text{Sym}^k S$ [16, 92, 99]

When $S = \mathbb{C}^2$, $\Sigma_0 \times S = \mathbb{P}^1 \times \mathbb{C}^2$ is non-compact, so a priori neither GW theory nor DT theory is defined. Using the torus action on \mathbb{C}^2 , one can define equivariant GW and DT theories.

- C1 Equivariant Gromov-Witten theory on $\mathbb{P}^1 \times \mathbb{C}^2$.
- C2 Equivariant Donaldson-Thomas theory on $\mathbb{P}^1 \times \mathbb{C}^2$.
- C3 Equivariant quantum cohomology of $\text{Hilb}_k \mathbb{C}^2$.
- C4 Equivariant orbifold quantum cohomology of $\text{Sym}^k \mathbb{C}^2$.

The equivalences among C1, C2, C3 are proved by Bryan, Okounkov, Pandharipande [15, 79, 80], whereas the equivalence between C3 and C4 is studied in [13].

Maulik and Oblomkov established the equivalence of the following theories when S is an ADE resolution.

- A1 Equivariant Gromov-Witten theory on $\mathbb{P}^1 \times S$.
- A2 Equivariant Donaldson-Thomas theory on $\mathbb{P}^1 \times S$.
- A3 Equivariant quantum cohomology of $\text{Hilb}_k S$.

In particular, $C2 \Leftrightarrow C3$ and $A2 \Leftrightarrow A3$ are special cases of the equivariant version of (46).

We may consider the case where S is a orbifold.

- O1 Equivariant orbifold Gromov-Witten theory on $\mathbb{P}^1 \times [\mathbb{C}^2/\mathbb{Z}_{n+1}]$.
- O2 Equivariant Donaldson Thomas theory on $\mathbb{P}^1 \times [\mathbb{C}^2/\mathbb{Z}_{n+1}]$.
- O3 Equivariant quantum cohomology of $\text{Hilb}_k[\mathbb{C}^2/\mathbb{Z}_{n+1}]$.

By Maulik-Okounkov [68], when $S = \mathcal{A}_n$ (resolution of A_n -singularity), A3 is equivalent of O3, which can be viewed as an example of Ruan's Crepant Transformation Conjecture. Z. Zhou [100] established the equivalence between O2 and O3.

Finally, let S be a smooth projective K3 surface. The Gromov-Witten theories of $K3$ and $K3 \times \mathbb{P}^1$ are trivial, so one considers reduced Gromov-Witten theory defined by reduced virtual class. The compact case is much more difficult than the non-compact case.

R1 Reduced Gromov-Witten theory of $\mathbb{P}^1 \times S$

R2 Reduced Donaldson-Thomas theory on $\mathbb{P}^1 \times S$.

R3 Reduced quantum cohomology of $\text{Hilb}_k S$.

G. Oberdieck studies R1 and R3 in [74] and [73], respectively. In [73], Oberdieck conjectures a formula for the quantum multiplication with divisor classes on $\text{Hilb}_k S$, and prove the conjecture in the first non-trivial case $\text{Hilb}_2 S$.

5. Dimensional reduction of torsion DT theory

5.1. Seiberg-Witten invariants via Mochizuki's wall crossing

Let S be a smooth projective surface and h an ample divisor on it. We assume that $H^1(\mathcal{O}_S) = 0$ and the arithmetic genus $p_g = \dim H^2(\mathcal{O}_S) > 0$. (For instance, any smooth hypersurface in a quintic 3-fold satisfies this assumption.) Let us take an element

$$v = (2, a, n) \in H^0(S) \oplus H^2(S) \oplus H^4(S)$$

such that $h \cdot a$ is an odd number. Then by this choice, any h -semistable sheaf $E \in \text{Coh}(S)$ with $\text{ch}(E) = v$ is h -stable. Now define $\mathcal{M}_h(v)$ to be the moduli space of h -stable sheaves $E \in \text{Coh}(S)$ with $\text{ch}(E) = v$. For simplicity, we assume that there exists a universal sheaf $\mathcal{E} \in \text{Coh}(S \times \mathcal{M}_h(v))$. Let p_M be the projection from $S \times \mathcal{M}_h(v)$ to $\mathcal{M}_h(v)$. We have the decomposition

$$\mathbf{R}p_{M*} \mathbf{R}\mathcal{H}om(\mathcal{E}, \mathcal{E}) = \mathbf{R}p_{M*} \mathbf{R}\mathcal{H}om(\mathcal{E}, \mathcal{E})_0 \oplus \mathbf{R}p_{M*} \mathcal{O}_{S \times \mathcal{M}_h(v)}$$

and the perfect obstruction theory

$$E_{\mathcal{M}_h(v)}^\bullet := \mathbf{R}p_{M*} \mathbf{R}\mathcal{H}om(\mathcal{E}, \mathcal{E})_0^\vee \rightarrow \mathbf{L}_{\mathcal{M}_h(v)}.$$

We have the associated virtual cycle $[\mathcal{M}_h(v)]^{\text{vir}}$ whose virtual dimension d is

$$d = a^2 - 4n - 3\chi(\mathcal{O}_S).$$

Now let $P(\mathcal{E})$ be a polynomial in the slant products $\text{ch}_i(\mathcal{E})/b$ for the elements $b \in H^*(S)$ and $i \in \mathbb{Z}_{\geq 0}$. By the wall crossing argument using the master space, Mochizuki described the invariant

$$\int_{[M_h(v)]^{\text{vir}}} P(\mathcal{E})$$

in terms of the Seiberg-Witten invariants and certain integration over the Hilbert schemes of points on S .

The SW invariant is then defined as follows: for any $c \in \text{NS}(S)$, let L be the line bundle on S such that $c_1(L) = c$, which is uniquely determined following the assumption that $H^1(\mathcal{O}_S) = 0$. Let $M(c)$ be the moduli space of non-zero morphisms $\mathcal{O}_S \rightarrow L$, which is isomorphic to $\mathbb{P}(H^0(S, L))$. The natural deformation theory of pairs $\mathcal{O}_S \rightarrow L$ induces an obstruction bundle $O(c)$ on $M(c)$, which fits into the exact sequence

$$\begin{aligned} 0 \rightarrow H^1(S, L) \otimes \mathcal{O}_{M(c)}(1) \rightarrow O(c) \rightarrow H^2(S, \mathcal{O}_S) \otimes \mathcal{O}_{M(c)} \\ \rightarrow H^2(S, L) \otimes \mathcal{O}_{M(c)}(1) \rightarrow 0. \end{aligned}$$

The virtual cycle $[M(c)]^{\text{vir}}$ is defined to be the Euler class of $O(c)$. If it is non-zero, then the virtual dimension is zero⁸, and

$$\text{SW}(c) = \int_{[M(c)]^{\text{vir}}} 1.$$

By setting $d(c) = h^0(S, L) - 1$, the SW invariant is computed as (cf. [71, Proposition 6.3.1])

$$\text{SW}(c) = (-1)^{d(c)} \binom{p_g - 1}{d(c)}.$$

Let $S^{[n]}$ be the Hilbert scheme of n -points in S . For $j = 1, 2$, let $\mathcal{Z}_i \subset S \times S^{[n_i]}$ be the universal subscheme and $\mathcal{I}_i \subset \mathcal{O}_{S \times S^{[n_i]}}$ be its ideal sheaf. Below we consider the decomposition

$$a_1 + a_2 = a, \quad a_i \in \text{NS}(S).$$

⁸The $p_g > 0$ assumption is required here in [71, Proposition 6.3.1].

We denote by e^{a_i} the line bundle on S whose first Chern class equals to a_i . We define $Q(\mathcal{I}_1 e^{a_1-s}, \mathcal{I}_2 e^{a_2+s})$ to be the Euler class of the following \mathbb{C}^* -equivariant virtual vector bundle on $S^{[n_1]} \times S^{[n_2]}$:

$$-\mathbf{R}p_* \mathbf{R}Hom(\mathcal{I}_1 e^{a_1-s}, \mathcal{I}_2 e^{a_2+s}) - \mathbf{R}p_* \mathbf{R}Hom(\mathcal{I}_2 e^{a_2+s}, \mathcal{I}_1 e^{a_1-s}).$$

Here s is the equivariant parameter with respect to the trivial \mathbb{C}^* -action, and p is the projection from $S \times S^{[n_1]} \times S^{[n_2]}$ to $S^{[n_1]} \times S^{[n_2]}$. All the equivariant sheaves in the derived inner Hom's are pulled back to $S \times S^{[n_1]} \times S^{[n_2]}$.

We also consider the rank n_i vector bundle on $S^{[n_i]}$, given by

$$\mathcal{V}_i = p_*(\mathcal{O}_{\mathbb{Z}_i} \otimes e^{a_i})$$

We define $\mathcal{A}(a_1, a_2, v)$ to be

$$(48) \quad \begin{aligned} &\mathcal{A}(a_1, a_2, v) \\ &= \sum_{\substack{n_1+n_2=n-a_1a_2 \\ n_1>n_2}} \int_{S^{[n_1]} \times S^{[n_2]}} \text{Res}_{s=0} \left(\frac{P(\mathcal{I}_1 e^{a_1-s} \oplus \mathcal{I}_2 e^{a_2+s})}{Q(\mathcal{I}_1 e^{a_1-s}, \mathcal{I}_2 e^{a_2+s})} \frac{e(\mathcal{V}_1) \cdot e(\mathcal{V}_2 e^{2s})}{(2s)^{n_1+n_2-p_g}} \right). \end{aligned}$$

The following result was obtained by Mochizuki:

Theorem 1. (Mochizuki [71, Theorem 1.4.6]) *Assume that $ah > 2K_S h$ and $\chi(v) = \int_S v \cdot \text{td}_S \geq 1$. Then we have the following formula:*

$$\frac{1}{2} \int_{\mathcal{M}_h(v)} P(\mathcal{E}) = - \sum_{\substack{a_1+a_2=a \\ a_1 h < a_2 h}} \text{SW}(a_1) \cdot 2^{1-\chi(v)} \cdot \mathcal{A}(a_1, a_2, v).$$

Remark 1. *Note that the factor 1/2 in the LHS comes from the difference between Mochizuki's convention and ours. Mochizuki used the moduli space of oriented stable sheaves, which is a μ_2 -gerb over our moduli space $M_h(v)$.*

Remark 2. *The assumptions $ah > 2K_S h$ and $\chi(v) \geq 1$ are satisfied if we replace v by $v \cdot e^{kh}$ for $k \gg 0$.*

5.2. Torsion DT invariants and Seiberg-Witten theory

Let (S, h) and $v \in H^*(S, \mathbb{Q})$ be as in the previous subsection, and consider

$$(49) \quad \pi: X_S = \omega_S \rightarrow S$$

the total space of the canonical line bundle on S . Note that X_S is a non-compact Calabi-Yau 3-fold, i.e. $\omega_{X_S} \cong \mathcal{O}_{X_S}$. Denote

$$(50) \quad \text{Coh}_c(X_S) \subset \text{Coh}(X_S)$$

to be the abelian category of coherent sheaves on X_S whose supports are compact. The slope function μ_h on $\text{Coh}_c(X_S) \setminus \{0\}$ defined as

$$\mu_h(E) = \frac{c_1(\pi_*E) \cdot h}{\text{rank}(\pi_*E)} \in \mathbb{Q} \cup \{\infty\}$$

determines a slope stability condition on $\text{Coh}_c(X_S)$ in the usual way. Let $\overline{\mathcal{M}}_h(v)$ be the moduli space of μ_h -stable sheaves $E \in \text{Coh}_c(X_S)$ with $\text{ch}(\pi_*E) = v = \langle r, \gamma, n \rangle$. The DT invariant for the local surface X is then defined by

$$(51) \quad \text{DT}_h(v) = \int_{\overline{\mathcal{M}}_h(v)} \nu_M \, d\chi.$$

Here ν_M is Behrend’s constructible function [7] on $\overline{\mathcal{M}}_h(v)$.

The one-dimensional complex torus \mathbb{C}^* acts on X_S by re-scaling on the fibers of π . By the localization, the DT invariant (51) coincides with the integration of ν_M over the \mathbb{C}^* -fixed locus $\overline{\mathcal{M}}_h(v)^{\mathbb{C}^*}$. Note that the moduli space $\mathcal{M}_h(v)$ is an open and closed subscheme of $\overline{\mathcal{M}}_h(v)^{\mathbb{C}^*}$.

Now let $\overline{\mathcal{E}} \in \text{Coh}(X_S \times \overline{\mathcal{M}}_h(v))$ be the universal family, and $p_{\overline{\mathcal{M}}}$ the projection from $X_S \times \overline{\mathcal{M}}_h(v)$ to $\overline{\mathcal{M}}_h(v)$. Let $\mathbf{R}\mathcal{H}om(\overline{\mathcal{E}}, \overline{\mathcal{E}})_0$ be the cone of the composition

$$\mathbf{R}\mathcal{H}om_{\overline{\mathcal{M}}_h(v) \times X_S}(\overline{\mathcal{E}}, \overline{\mathcal{E}}) \xrightarrow{\pi_*} \mathbf{R}\mathcal{H}om_{\overline{\mathcal{M}}_h(v) \times S}(\pi_*\overline{\mathcal{E}}, \pi_*\overline{\mathcal{E}}) \xrightarrow{\text{tr}} \mathcal{O}_{\overline{\mathcal{M}}_h(v) \times S}.$$

We obtain the \mathbb{C}^* -fixed trace-free perfect⁹ obstruction theory

$$(52) \quad E_{\overline{\mathcal{M}}_h(v)}^\bullet := (\mathbf{R}p_{\overline{\mathcal{M}}}^* \mathbf{R}\mathcal{H}om(\overline{\mathcal{E}}, \overline{\mathcal{E}})_0[1])^{\vee \mathbb{C}^*} \rightarrow \mathbf{L}_{\overline{\mathcal{M}}_h(v)^{\mathbb{C}^*}}.$$

Let $[\overline{\mathcal{M}}_h(v)^{\mathbb{C}^*}]^{\text{vir}}$ be the associated virtual fundamental class. Instead of working with the invariant (51), we can consider the invariant

$$(53) \quad \overline{\overline{\text{DT}}}_h(v) = \int_{[\overline{\mathcal{M}}_h(v)^{\mathbb{C}^*}]^{\text{vir}}} c \left((\mathbf{R}p_{\overline{\mathcal{M}}}^* \mathbf{R}\mathcal{H}om(\overline{\mathcal{E}}, \overline{\mathcal{E}})_0[1])^{\vee \mathbb{C}^*} \right).$$

⁹By the Serre duality and the non-trivial \mathbb{C}^* -weight on ω_S , the higher obstruction space vanishes after taking the \mathbb{C}^* -fixed part.

Remark 3. Here the definition of the invariant (53) is motivated by Fantechi-Göttsche’s virtual Euler numbers [27, Section 4]. In fact, it coincides with the virtual Euler number of $\overline{\mathcal{M}}_h(v)$ up to a sign. In particular, when $\overline{\mathcal{M}}_h(v)^{\mathbb{C}^*}$ is non-singular with the expected dimension, then the above invariant coincides with the topological Euler number of $\overline{\mathcal{M}}_h(v)$ up to a sign.

Now let $r = 2$. The \mathbb{C}^* -fixed locus $\overline{\mathcal{M}}_h(v)^{\mathbb{C}^*}$ decomposes into two components

$$\overline{\mathcal{M}}_h(v)^{\mathbb{C}^*} = \mathcal{M}_h(v) \coprod \widehat{\mathcal{M}}_h(v)$$

where $\mathcal{M}_h(v)$ is, roughly speaking, the moduli space of μ_h -stable torsion-free sheaves of rank 2 on S , and $\widehat{\mathcal{M}}_h(v)$ is the moduli space of \mathbb{C}^* -fixed μ_h -stable sheaves with rank one on the “fat” surface $2S$. Now we can define the DT invariants associated to each component, by restricting the obstruction theory in (52) to each component and integrating against the corresponding induced virtual cycle. Let

$$(54) \quad \widehat{\text{DT}}_h(v) = \int_{[\widehat{\mathcal{M}}_h(v)]^{\text{vir}}} c\left(\left(\mathbf{R}p_{\overline{\mathcal{M}}^*} \mathbf{R}\mathcal{H}om(\overline{\mathcal{E}}, \overline{\mathcal{E}})_0[1]\right)^{\vee \mathbb{C}^*}\right)$$

be the contribution from the “fat” surface $2S$ and

$$(55) \quad \text{DT}_h(v) = \int_{[\mathcal{M}_h(v)]^{\text{vir}}} c\left(\left(\mathbf{R}p_{\overline{\mathcal{M}}^*} \mathbf{R}\mathcal{H}om(\overline{\mathcal{E}}, \overline{\mathcal{E}})_0[1]\right)^{\vee \mathbb{C}^*}\right)$$

be the contribution from rank 2 sheaves on S . Note that, as we saw in the last section, the latter contributions have been computed in some cases by Mochizuki [71] (look at Theorem 1). On the other hand, it can be shown [31, Proposition 3.13, 3.14, Corollary 3.15, and Proposition 3.21] that the contribution of $\widehat{\mathcal{M}}_h(v)$ to $\widehat{\text{DT}}_h(v)$ is given by the invariants of “nested Hilbert schemes” on S , denoted by $S_\beta^{[n_1 > n_2]}$, parametrizing 2-step flags $\mathcal{I}_{Z_1}(-C_1) \hookrightarrow \mathcal{I}_{Z_2}$ of (possibly twisted) ideal sheaves of subschemes, $(Z_1, Z_2), (C_1)$ of S , where $Z_i, i = 1, 2$ are zero dimensional subschemes of S with length n_1, n_2 respectively, $C_1 \subset S$ is a divisor with $[C_1] = \beta$ for some suitable β , such that Z_2 is a subscheme of $Z_1 \cup C_1$ (hence it induces the injective map of the corresponding ideal sheaves). We then prove the following identity in [31, Theorem 4] relating $\overline{\text{DT}}_h(v)$ to SW invariants and the invariants of nested Hilbert schemes contributing to $\widehat{\text{DT}}_h(v)$.

$$(56) \quad \overline{\text{DT}}_h(v) = - \sum_{\substack{a_1 + a_2 = a \\ a_1 h < a_2 h}} \text{SW}(a_1) \cdot 2^{1-\chi(v)} \cdot \mathcal{A}(a_1, a_2, v, P) + \widehat{\text{DT}}_h(v).$$

Here P is a universally defined explicit integrand [31, Proposition 4.4], which is defined independent of S and $\widehat{\text{DT}}_h(v)$ is, roughly speaking, given as a sum over the contribution of nested Hilbert schemes $S_\beta^{[n_1 > n_2]}$, for all allowed values of n_1, n_2, β induced by the choice of v . It must be pointed out that by S-duality consideration, the generating series of the invariants on the left-hand side of (56) is expected to be given by partition function of Vafa-Witten invariants [31, Remark 2.10] which are expected to have modular properties [84, 93], while the generating series of $\widehat{\text{DT}}_h(v)$ is also (in some cases) shown to be given by modular forms of certain weight [30, Section 6]. Therefore, equation (56) implies that the generating series of the Seiberg-Witten invariants (despite not being necessarily modular itself) can in some cases be written in terms of modular forms.

References

- [1] B. Anichouche and I. Biswas, *Einstein-Hermitian connections on polystable principal bundles over a compact Kähler manifold*, Amer. J. Math. **123** (2001), 207–228.
- [2] M. Aganagic, A. Klemm, M. Mariño, and C. Vafa, *The topological vertex*, Commun. Math. Phys. **254** (2005), 425–478.
- [3] B. S. Acharya, M. O’Loughlin, and B. J. Spence, *Higher-dimensional analogues of Donaldson-Witten theory*, Nucl. Phys. B **503** (1997) 657. [arXiv:hep-th/9705138](#).
- [4] M. Aganagic, M. Mariño, and C. Vafa, *All loop topological string Amplitudes from Chern-Simons theory*, Comm. Math. Phys. **247** (2004), no. 2, 467–512.
- [5] M. Aganagic, H. Ooguri, N. Saulina, and C. Vafa, *Black Holes, q -deformed 2d Yang-Mills and nonperturbative topological strings*, Nucl. Phys. B **715** (2005), no. 1-2, 23, 304–348.
- [6] L. Baulieu, H. Kanno, and I. M. Singer, *Special quantum field theories in eight and other dimensions*, Commun. Math. Phys. **194** (1998), no. 1, 149–175.
- [7] K. Behrend, *Donaldson-Thomas invariants via microlocal geometry*, Ann. of Math. (2) **170** (2009), no. 3, 1307–1338.

- [8] M. Bershadsky, S. Cecotti, H. Ooguri, and C. Vafa, *Holomorphic anomalies in topological field theories*, Nucl. Phys. B **405** (1993), no. 2-3, 279–304.
- [9] M. Bershadsky, A. Johansen, V. Sadov, and C. Vafa, *Topological reduction of 4D SYM to 2D σ -models*, Nuclear Phys. B **448** (1995), no. 1-2, 166–186. [arXiv:hep-th/9501096](#).
- [10] M. Bershadsky, C. Vafa, and V. Sadov, *D-branes and topological field theories*, Nucl. Phys. B **463** (1996) 420. [arXiv:hep-th/9511222](#).
- [11] M. Blau and G. Thompson, *Euclidean SYM theories by time reduction and special holonomy manifolds*, Phys. Lett. B **415** (1997) 242. [arXiv:hep-th/9706225](#).
- [12] M. Blau and G. Thompson, *Euclidean SYM theories by time reduction and special holonomy manifolds*, Phys. Lett. B **415** (1997), no. 3, 242–252.
- [13] J. Bryan and T. Graber, *The crepant resolution conjecture*, Algebraic geometry — Seattle 2005, Part 1, 23–42, Proc. Sympos. Pure Math. **80**, Part 1, Amer. Math. Soc., Providence, RI, 2009.
- [14] J. Bryan and N. C. Leung, *The enumerative geometry of K3 surfaces and modular forms*, J. Amer. Math. Soc. **13** (2000), no. 2, 371–410.
- [15] J. Bryan and R. Pandharipande, *The local Gromov-Witten theory of curves*, with an appendix by Bryan, C. Faber, A. Okounkov and Pandharipande, J. Amer. Math. Soc. **21** (2008), no. 1, 101–136.
- [16] W. Chen and Y. Ruan, *Orbifold Gromov-Witten theory*, in: Orbifolds in mathematics and physics (Madison, WI, 2001), pp. 25–85, Contemp. Math. **310**, Amer. Math. Soc., Providence, RI, 2002.
- [17] T.-M. Chiang, A. Klemm, S.-T. Yau, and E. Zaslow, *Local mirror symmetry: Calculations and interpretations*, Adv. Theor. Math. Phys. **3** (1999), no. 3, 495–565. [arXiv:hep-th/9903053](#).
- [18] K. Costello and I. Grojnowski, *Hilbert schemes, Hecke algebras and the Calogero-Sutherland system*, [arXiv:math.AG/0310189](#).
- [19] R. Dijkgraaf, *Mirror symmetry and elliptic curves*, in: The moduli space of curves (Texel Island, 1994), 149–163, Progr. Math. **129**, Birkhäuser Boston, Boston, MA, 1995.

- [20] R. Dijkgraaf and G. W. Moore, *Balanced topological field theories*, Commun. Math. Phys. **185** (1997), no. 2, 411–440.
- [21] R. Dijkgraaf, G. W. Moore, E. Verlinde, and H. Verlinde, *Elliptic genera of symmetric products and second quantized strings*, Commun. Math. Phys. **185** (1997) no. 1, 197–209.
- [22] D.-E. Diaconescu, B. Florea, *The ruled vertex and nontoric del Pezzo surfaces*, J. Hep **12**, (2006), no. 028.
- [23] D.-E. Diaconescu, B. Florea, and N. Saulina, *A vertex formalism for local ruled surfaces*, Comm. Math. Phys. **265**, (2006) no. 1, 201–226.
- [24] S. Donaldson, *Anti-self-dual Yang-Mills connections over complex algebraic surfaces and stable vector bundles*, Proc. Lond. Math. Soc. **50** (1985).
- [25] S. Donaldson and R. Thomas, *Gauge theory in higher dimensions*, in: The Geometric Universe: Science, Geometry, and the Work of Roger Penrose, S. Huggett et. al eds., Oxford Univ. Press, 1998.
- [26] D. Edidin and Z. Qin, *The Gromov-Witten and Donaldson-Thomas correspondence for trivial elliptic fibrations*, Internat. J. Math. **18** (2007), no. 7, 821–838.
- [27] B. Fantechi and L. Göttsche, *Riemann-Roch theorems and elliptic genus for virtually smooth schemes*, Geom. Topol. **14** (2010), no. 1, 83–115.
- [28] C. Faber and R. Pandharipande, *Hodge integrals and Gromov-Witten theory*, Invent. Math. **139** (2000), no. 1, 173–199.
- [29] R. Fintushel and R. Stern, *Immersed spheres in 4-manifolds and the immersed Thom conjecture*, Turkish Journal of Mathematics **19** (1995), 145–157.
- [30] A. Gholampour, A. Sheshmani, and S.-T. Yau, *Nested Hilbert Scheme on Surfaces: Virtual Fundamental Class*, arXiv:1701.08899.
- [31] A. Gholampour, A. Sheshmani, and S.-T. Yau, *Localized Donaldson-Thomas theory of surfaces*, arXiv:1701.08902.
- [32] T. L. Gómez and I. Sols, *Projective moduli space of semistable principal sheaves for a reductive group*, Comm. Alg. Alg. Geom., Catania, (2001).

- [33] L. Göttsche, *The Betti numbers of the Hilbert scheme of points on a smooth projective surface*, Math. Ann. **286** (1990), 193–207.
- [34] L. Göttsche and R. Pandharipande, *The quantum cohomology of blow-ups of P^2 and enumerative geometry*, J. Differential Geom. **48** (1998), no. 1, 61–90.
- [35] P. Griffiths and J. Harris, *Principles of algebraic geometry*, Pure and Applied Mathematics, Wiley-Interscience [John Wiley & Sons], New York, 1978. xii+813pp.
- [36] R. Gopakumar and C. Vafa, *M-theory and topological strings. I,II*, arXiv:hep-th/9809187; hep-th/9812127.
- [37] F. Hirzebruch, T. Höfer, *On the Euler number of an orbifold*, Math. Ann. **286** (1990) 255.
- [38] C. Hofman and J-S. Park, *Cohomological Yang-Mills theories on Kahler 3-folds*, Nuclear Phys. B **600** (2001), no. 1, 133–162.
- [39] S. Hosono, M.-H. Saito, and A. Takahashi, *Holomorphic anomaly equation and BPS state counting of rational elliptic surface*, Adv. Theor. Math. Phys. **3** (1999), 177–208.
- [40] S. Hosono, M.-H. Saito, and A. Takahashi, *Relative Lefschetz action and BPS state counting*, Internat. Math. Res. Notices, (2001), no. 15, 783–816.
- [41] S. Hosono, *Counting BPS states via holomorphic anomaly equations*, in: Calabi-Yau varieties and mirror symmetry (Toronto, ON, 2001), 57–86, Fields Inst. Commun. **38**, Amer. Math. Soc., Providence, RI, 2003.
- [42] J. Hu, *Gromov-Witten invariants of blow-ups along surfaces*, Compositio Math. **125** (2001), no. 3, 345–352.
- [43] E.-N. Ionel and T. Parker, *The Gromov invariants of Ruan-Tian and Taubes*, Math. Res. Lett. **4** (1997), no. 4, 521–532.
- [44] A. Iqbal, *All genus topological string amplitudes and 5-brane Webs as Feynman diagrams*, arXiv:hep-th/0207114.
- [45] A. Iqbal, N. Nekrasov, A. Okounkov, and C. Vafa, *Quantum foam and topological strings*, J. High Energy Phys. (2008), no. 4, 011, 47pp.
- [46] D. Joyce and Y. Song, *A theory of generalized Donaldson-Thomas invariants*, Mem. Amer. Math. Soc., Publication **217**, (2012), no. 1020.

- [47] Albrecht Klemm, Jan Manschot, and Thomas Wotschke, *Quantum geometry of elliptic Calabi-Yau manifolds*, Commun. Number Theory Phys. **6** (2012), no. 4, 849–917.
- [48] M. Kontsevich and Y. Manin, *Gromov-Witten classes, quantum cohomology, and enumerative geometry*, Comm. Math. Phys. **164** (1994), no. 3, 525–562.
- [49] M. Kontsevich and Y. Soibelman, *Stability structures, motivic Donaldson-Thomas invariants and cluster transformations*, arXiv: math.AG/0811.2435.
- [50] S. Katz, A. Klemm, and C. Vafa, *M-theory, topological strings, and spinning black holes*, Adv. Theor. Math. Phys. **3** (1999), 1445–1537.
- [51] A. Kapustin and E. Witten, *Electric-magnetic duality and the geometric Langlands program*, Commun. Number Theory Phys. **1** (2007), no. 1, 1–236.
- [52] J. M. F. Labastida and C. Lozano, *Mathai-Quillen formulation of twisted $N = 4$ supersymmetric gauge theories in four-dimensions*, Nuclear Phys. B **502** (1997), no. 3, 741–790.
- [53] C. Lozano and M. Mariño, *Donaldson invariants of product ruled surfaces and two-dimensional gauge theories*, Commun. Math. Phys. **220** (2001), no. 2, 231–261.
- [54] M. Lehn and C. Sorger, *Symmetric groups and the cup product on the cohomology of Hilbert schemes*, Duke Math. J. **110**, (2001), no. 2, 345–357.
- [55] M. Lehn and C. Sorger, *The cup product of the Hilbert scheme for $K3$ surfaces*, Invent. math. **152**, (2003), no. 2, 305–329.
- [56] A. M. Li and Y. B. Ruan, *Symplectic surgery and Gromov-Witten invariants of Calabi-Yau 3-folds*, Invent. Math. **145** (2001), no. 1, 151–218.
- [57] J. Li, C.-C. M. Liu, K. Liu, and J. Zhou, *A mathematical theory of the topological vertex*, Geom. Topol. **13** (2009), no. 1, 527–621.
- [58] J. Li, *A Degeneration formula of GW-invariants*, J. Diff. Geom. **60** (2002), no. 2, 177–354.
- [59] T.-J. Li and A.-K. Liu, *The equivalence between SW and Gr in the case where $b^+ = 1$* , Internat. Math. Res. Notices **7** (1999), 335–345;

- Uniqueness of symplectic canonical class, surface cone and symplectic cone of 4-manifolds with $b^+ = 1$* , J. Diff. Geom. **58** (2001), 331–370.
- [60] C.-C. M. Liu, K. Liu, and J. Zhou, *A proof of a conjecture of Mariño-Vafa on Hodge integrals*, J. Differential Geom. **65** (2003), no. 2, 289–340.
- [61] C.-C. M. Liu, K. Liu, and J. Zhou, *A formula of two-partition Hodge integrals*, J. Amer. Math. Soc. **20** (2007), no. 1, 149–184.
- [62] C. Lozano and M. Mariño, *Donaldson invariants of product ruled surfaces and two-dimensional gauge theories*, Commun. Math. Phys. **220** (2001), no. 2, 231–261.
- [63] N. Marcus, *The Other topological twisting of $N = 4$ Yang-Mills*, Nucl. Phys. B **452** (1995), no. 1-2, 331–345.
- [64] M. Mariño and G. W. Moore, *Donaldson invariants for non-simply connected manifolds*, Commun. Math. Phys. **203** (1999), no. 2, 249–267. [arXiv:hep-th/9804104](https://arxiv.org/abs/hep-th/9804104).
- [65] D. Maulik, N. Nekrasov, A. Okounkov, and R. Pandharipande, *Gromov-Witten theory and Donaldson-Thomas theory I*, Compos. Math. **142** (2006), no. 5, 1263–1285. .
- [66] D. Maulik, N. Nekrasov, A. Okounkov, and R. Pandharipande, *Gromov-Witten theory and Donaldson-Thomas theory II*, Compos. Math. **142**, (2006), no. 5, 1286–1304.
- [67] D. Maulik, A. Oblomkov, A. Okounkov, and R. Pandharipande, *Gromov-Witten/Donaldson-Thomas correspondence for toric 3-folds*, Invent. math. **186** (2011) no. 2, 435–479.
- [68] D. Maulik and A. Okounkov, *Quantum groups and quantum cohomology*, [arXiv:1211.1287](https://arxiv.org/abs/1211.1287).
- [69] D. Maulik and R. Pandharipande, *Gromov-Witten theory and Noether-Lefschetz theory*, in: A Celebration of Algebraic Geometry, pp. 469–507, Clay Math. Proc. **18**, Amer. Math. Soc., Providence, RI, 2013.
- [70] J. A. Minahan, D. Nemeschansky, C. Vafa, and N. P. Warner, *E-strings and $N = 4$ topological Yang-Mills theories*, Nucl. Phys. B **527** (1998), no. 3, 581–623.
- [71] T. Mochizuki, *Donaldson type invariants for algebraic surfaces*, Lect. Notes Math., Springer-Verlag, Berlin, Vol. 1972, 2009.

- [72] G. Moore and E. Witten, *Integration over the u -plane in Donaldson theory*, Adv. Theor. Math. Phys. **1** (1998), no. 2, 298–387.
- [73] G. Oberdieck, *Gromov-Witten invariants of the Hilbert schemes of points of a $K3$ surface*, Geom. Topol. **22** (2018), no. 1, 323–437.
- [74] G. Oberdieck, *Gromov-Witten theory of $K3 \times \mathbb{P}^1$ and quasi-Jacobi forms*, arXiv:1605.05238.
- [75] G. Oberdieck and R. Pandharipande, *Curve counting on $K3 \times E$, the Igusa cusp form χ_{10} , and descendent integration*, $K3$ surfaces and their moduli, 245–278, Progr. Math. **315**, Birkhäuser/Springer, [Cham], 2016.
- [76] G. Oberdieck and J. Shen, *Curve counting on elliptically fibered Calabi-Yau 3-folds*, arXiv:1608.07073.
- [77] C. Okonek and A. Teleman, *Seiberg-Witten invariants for manifolds with $b_+ = 1$, and the universal wall crossing formula*, Internat. J. Math. **7** (1996), no. 6, 811–832.
- [78] A. Okounkov and R. Pandharipande, *Hodge integrals and invariants of the unknot*, Geom. Topol. **8** (2004), 675–699.
- [79] A. Okounkov and R. Pandharipande, *Quantum cohomology for the Hilbert scheme of points in the plane*, Invent. math. **179** (2010), no. 3, 523–557.
- [80] A. Okounkov and R. Pandharipande, *The local Donaldson-Thomas theory of curves*, Geom. Topol. **14** (2010), 1503–1567.
- [81] Y. Ruan, *Symplectic topology and complex surfaces*, in: Geometry and Analysis on Complex Manifolds, pp. 171–197, World Sci. Publ., River Edge, NJ, 1994.
- [82] K. Sakai, *Topological string amplitudes for the local $\frac{1}{2}$ $K3$ surface*, PTEP. Prog. Theor. Exp. Phys. 2017, no. 3, 033B09, 29 pp.
- [83] A. Schmitt, *Singular principal bundles over higher-dimensional manifolds and their moduli spaces*, Int. Math. Res. Not. (2002), no. 23, 1183–1209.
- [84] Y. Tanaka and R. P. Thomas, *Vafa-Witten invariants for projective surfaces I: stable case*, arXiv:1702.08487.
- [85] C. Taubes, *Seiberg Witten and Gromov invariants for symplectic 4-manifolds*, International Press, Somerville, MA, 2000.

- [86] G. Tian and S.-T. Yau, *Kähler-Einstein metrics on complex surfaces with $c_1 > 0$* , *Comm. Math. Phys.* **112** (1987), no. 1, 175–203.
- [87] G. Tian, *On Calabi's conjecture for complex surfaces with positive first Chern class*, *Invent. Math.* **101** (1990), no. 1, 101–172.
- [88] G. Tian and S.-T. Yau, *Complete Kähler manifolds with zero Ricci curvature. I*, *J. Amer. Math. Soc.* **3** (1990), no. 3, 579–609.
- [89] G. Tian and S.-T. Yau, *Complete Kähler manifolds with zero Ricci curvature. II*, *Invent. Math.* **106** (1991), no. 1, 27–60.
- [90] R. P. Thomas, *A holomorphic Casson invariants for Calabi-Yau 3-folds, and bundles on K3 fibrations*, *J. Diff. Geom.* **54** (2000), no. 2, 367–438.
- [91] K. Uhlenbeck and S.-T. Yau, *On the existence of Hermitian-Yang-Mills connections in stable vector bundles*, *Comm. Pure Appl. Math.* **39** (1986), no. S, supp. S257–S293.
- [92] C. Vafa, *String vacua and orbifoldized LG models*, *Modern Phys. Lett. A* **4** (1989), no. 12, 1169–1185.
- [93] C. Vafa and E. Witten, *A Strong coupling test of S duality*, *Nucl. Phys. B* **431** (1994), no. 1-2, 3–77.
- [94] E. Vasserot, *Sur l'anneau de cohomologie du schéma de Hilbert de \mathbb{C}^2* , *C. R. Acad. Sci. Paris, Sé. I Math.* **332** (2001), no. 1, 7–12.
- [95] E. Witten, *Monopoles and four-manifolds*, *Math. Res. Lett.* **1** (1994), no. 6, 769–796.
- [96] E. Witten, *Supersymmetric Yang-Mills theory on a four manifold*, *J. Math. Phys.* **35** (1994), no. 10, 5101–5135.
- [97] S.-T. Yau, *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation. I*, *Comm. Pure Appl. Math.* **31** (1978), no. 3, 339–411.
- [98] S.-T. Yau and E. Zaslow, *BPS states, string duality, and nodal curves on K3*, *Nuclear Phys. B* **471** (1996), no. 3, 503–512.
- [99] E. Zaslow, *Topological orbifold models and quantum cohomology rings*, *Comm. Math. Phys.* **156** (1993), no. 2, 301–331.
- [100] Z. Zhou, *Donaldson-Thomas theory of $[\mathbb{C}^2/\mathbb{Z}_{n+1}] \times \mathbb{P}^1$* , [arXiv:1510.00871](https://arxiv.org/abs/1510.00871).

WALTER BURKE INSTITUTE FOR THEORETICAL PHYSICS
CALIFORNIA INSTITUTE OF TECHNOLOGY, PASADENA, CA 91125, USA
AND MAX-PLANCK-INSTITUT FÜR MATHEMATIK
VIVATSGASSE 7, D-53111 BONN, GERMANY
E-mail address: gukov@theory.caltech.edu

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY
2990 BROAD- way, NEW YORK, NY 10027, USA
E-mail address: ccliu@math.columbia.edu

CENTER FOR QUANTUM GEOMETRY AND MODULI SPACES
DEPARTMENT OF MATHEMATICS, AARHUS UNIVERSITY
NY MUNKEGADE 118, BUILDING 1530, 8000 AARHUS C, DENMARK
AND CENTER FOR MATHEMATICAL SCIENCES AND APPLICATIONS
HARVARD UNIVERSITY, CAMBRIDGE, MA 02138, USA
AND NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS, RUS-
SIAN FEDERATION, LABORATORY OF MIRROR SYMMETRY
NRU HSE, 6 USACHEVA STR., MOSCOW, RUSSIA, 119048
E-mail address: artan@cmsa.fas.harvard.edu

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY
CAMBRIDGE, MA 02138, USA
E-mail address: yau@math.harvard.edu