

T-Duality from super Lie n -algebra cocycles for super p -branes

DOMENICO FIORENZA, HISHAM SATI, AND URS SCHREIBER

We compute the L_∞ -theoretic double dimensional reduction of the F1/D p -brane super L_∞ -cocycles with coefficients in rationalized twisted K-theory from the 10d type IIA and type IIB super Lie algebras down to 9d. We show that the two resulting coefficient L_∞ -algebras are naturally related by an L_∞ -isomorphism which we find to act on the super p -brane cocycles by the infinitesimal version of the rules of topological T-duality and inducing an isomorphism between K^0 -cocycles in type IIA and K^1 -cocycles in type IIB, rationally. In particular this is a derivation of the Buscher rules for RR-fields (Hori’s formula) from first principles. Moreover, we show that these L_∞ -algebras are the homotopy quotients of the RR-charge coefficients by the “T-duality Lie 2-algebra”. We find that the induced L_∞ -extension is a gerby extension of a $9 + (1 + 1)$ dimensional (i.e. “doubled”) T-duality correspondence super-spacetime, which serves as a local model for T-folds. We observe that this still extends, via the D0-brane cocycle of its type IIA factor, to a $10 + (1 + 1)$ -dimensional super Lie algebra. Finally we show that this satisfies expected properties of a local model space for F-theory elliptic fibrations.

| | | |
|----------|--|-------------|
| 1 | Introduction | 1210 |
| 2 | Supersymmetry super Lie n-algebras | 1216 |
| 3 | Double dimensional reduction | 1224 |
| 4 | The brane supercocycles | 1236 |
| 5 | Super L_∞-algebraic T-duality | 1242 |
| 6 | T-Correspondence space and Doubled spacetimes | 1248 |

| | | |
|----------|--------------------------------|-------------|
| 7 | T-Duality Lie 2-algebra | 1255 |
| 8 | F-theory | 1259 |
| | References | 1265 |

1. Introduction

Understanding and constructing fields and branes in string theory and M-theory in a manner compatible with supersymmetry and with the various dualities is a fundamental problem, both from the theoretical as well as the phenomenological point of view. On the former, a complete solution to this problem would provide a solid ground for deriving the theory from firm (mathematical) principles. On the latter, it would help in the systematic classification of allowable vacua.

The fundamental super p -branes that have no gauge fields on their world-volume and which propagate on super-Minkowski spacetime are defined via Green-Schwarz type action functionals. These are higher-dimensional super coset WZW-type functionals [32], for super-Minkowski regarded as the super-coset of super-Poincaré by the Spin cover of the Lorentz group. Accordingly, these p -branes are classified by the invariant super Lie algebra cohomology of the supersymmetry algebras, a fact known as the “old brane scan” [1]. When super-Minkowski target spacetime is generalized to curved super-spacetimes, then this statement applies super-tangent-space wise: the bispinorial component of the field strength super $(p + 2)$ -form H_{p+2} to which the p -brane couples is constrained to coincide in each tangent space with the left-invariant form $\frac{1}{p!} (\bar{\psi} \wedge \Gamma_{a_1 \dots a_p} \psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_p}$ corresponding to the super-cocycles in the old brane scan [5, 6]. Notice that this is in direct analogy for instance to G_2 -structures on 7-manifolds, which are given by differential 3-forms that are constrained to coincide tangent-space wise with a fixed 3-cocycle on \mathbb{R}^7 . In particular, the bosonic part of the field strength H_{p+2} may vanish identically, and still its bispinorial component is constrained to coincide with the given cocycle super-tangent-space wise. In this way the super Lie algebra cocycles tightly control the structure of super p -brane charges.

We view the above as a powerful statement: While in general these differential forms H_{p+2} are just the image in real cohomology (“rationalization”) of components of some more refined cohomology theory, this says that at least the rational image of the charges of branes without gauge fields on

their worldvolume is classified by super Lie algebra cohomology. Notice that in certain instances rationalization of twisted generalized cohomology in the treatment of T-duality is even forced upon us (see [47]).

This phenomenon turns out to generalize also to those branes that do carry (higher) gauge fields on their worldvolume, such as the D-branes and the M5-brane — *if* one generalizes the Chevalley-Eilenberg algebras of super-Minkowski spacetimes to quasi-free differential-graded super-commutative algebras with generators also in higher degree [17, 61]. These DG-algebras are just the “FDAs” from the supergravity literature [15, 20]. For instance what in [17] is identified as the cocycle for the M5-brane earlier appeared as an algebraic ingredient in the construction of 11d supergravity in [20]; similarly the algebra for the Dp -brane charges found in [61] and [17] appears earlier as an ingredient for constructing type II supergravity in [14].

Now, while (super) Lie algebra cohomology is a respectable mathematical subject, what are these “extended super Minkowski algebras” that carry the D-brane and M-brane charges, really? In [27] we had pointed out, following [63], that these “FDA”s are naturally identified as the Chevalley-Eilenberg algebras of *super Lie n -algebras* also called *n -term super L_∞ -algebras*, for higher n ¹, and that under this identification the Lie theory that underlies the “old brane scan” turns into the “higher Lie theory” or homotopy theory of super Lie n -algebras that sees the entire super p -brane content [27].


This homotopy-theoretic perspective sheds further light on the classification of super p -branes. For instance, it identifies the extended super-Minkowski spacetimes of [17, 61] as being the higher central super Lie n -algebra extensions of super-Minkowski spacetime that are classified by the 3-cocycles for the superstring and by the 4-cocycle for the M2-brane. This is in higher analogy to how 2-cocycles classify ordinary central extensions. Namely, these extended spacetimes are the *homotopy fibers* of the corresponding cocycles [27, Prop. 3.5], see example 2.5 below.

This means that by embedding super Lie algebra theory into the larger context of homotopy super Lie algebra theory (super L_∞ -algebra theory)

¹Notice that these are Lie n -algebras in the sense of Stasheff [45, 46, 63], not “ n -Lie algebras” in the sense of Filippov. However, the two notions are not unrelated. At least the Filippov 3-algebras that appear in the BLG model of coincident solitonic M2-branes may naturally be understood as Stasheff-type Lie 2-algebras equipped with a metric form. This observation is due to [57, section 2], based on [24].

then *all* super p -branes are found by a sequence of consecutive higher invariant extensions, yielding a “bouquet of branes” growing out of the super-spacetimes [27]. This provides the generalization of the “old brane scan” that was argued to be needed in [49]. There, the multiplicity of elementary and solitonic p -brane solutions to supergravity theories was shown to cover many more values of (D, d) than the classic κ -symmetric points on the brane scan, suggesting that the original classification needs to be generalized. Indeed, our work pins down that the required generalization is from super Lie algebra cohomology to super L_∞ -algebra cohomology.

In fact this bouquet of invariant higher super L_∞ -cocycles is rooted in 0-dimensional super-space, the superpoint. It is a diagram of super Lie n -algebras of the following form:



Here every solid arrow denotes a central super L_∞ -algebra extension which is invariant with respect to the maximal semisimple part of the bosonic body of the external automorphisms (i.e. all automorphisms modulo R-symmetries) of the super L_∞ -algebra that is being extended. These turn out to be the respective Lorentz groups (their Spin-covers). Notice that the claim is that it is the maximality of these invariant extensions which implies that the extensions of the superpoint are super-Minkowski spacetimes, and that they are precisely of dimension increasing from 0 through 3,4, 6, 10 to 11. The top of this diagram is discussed in [27]. The proof of the “trunk” of the bouquet

is due to [37]. (For dimensions 0 to 3 the statement was observed earlier in [67].)

This shows that the core structure of string/M-theory follows from first principles in higher super Lie algebra theory, with no need of an external input from Lorentz geometry, or Spin geometry. Instead, Lorentzian geometry and Spin geometry is discovered (re-discovered) by analyzing the super-point in higher super Lie theory, as is the existence of all the super p -branes in their respective super-spacetimes.² This further suggests that higher super Lie theory knows much more about the inner working of string/M-theory, and that we may check conjectures on M-theory and discover its missing details by a systematic homotopy theoretic analysis of the superpoint. In the present article we are concerned with discovering and studying *T-duality* from this perspective (see e.g. [2]). We will show that this allows to systematically derive phenomena that have been proposed or conjectured but which seem to have been lacking a derivation from first principles, such as the rules of “topological T-duality” on supermanifolds, the nature of super-T-folds and the emergence of F-theory from M-theory.

In order to describe the action of T-duality on the F1/D p branes, we need the cocycles for the super D-branes not as cocycles with coefficients in \mathbb{R} on the type II extended super-Minkowski super Lie 2-algebras that are denoted **string**_{IIA} and **string**_{IIB} in the above diagram, but we need to descend them to cocycles on the type II super-Minkowski super Lie 1-algebras themselves, where they will take values in more complicated richer coefficients. This homotopical descent of the brane cocycles we have previously discovered and studied in [28] and [29]. There we showed that homotopy theory allows to descend the iterated \mathbb{R} -valued cocycles for separate p -branes, defined on an extended Minkowski spacetime, back to single cocycles on plain super-Minkowski spacetime, but now taking values in more complex coefficients. We showed in [29, section 4] that applying this homotopy-theoretic descent to the type IIA D p -brane cocycles gives that jointly they combine with the cocycle for the type IIA superstring to one single cocycle, now with coefficients in the L_∞ -algebra which is the image under Lie differentiation of the classifying space $KU/\mathbf{BU}(1)$ for twisted K-theory. Of course twisted K-theory has famously been argued earlier to be the correct cohomology

² Compare to [51, p.41]: “Perhaps we need to understand the nature of time itself better. [...] One natural way to approach that question would be to understand in what sense time itself is an emergent concept, and one natural way to make sense of such a notion is to understand how pseudo-Riemannian geometry can emerge from more fundamental and abstract notions such as categories of branes.”

theory in which F1/D p -brane (background) charges properly take value [76] [52] [42] [10]. Therefore, it is of interest to explore how much more the homotopy theory of super Lie- n algebra may teach us about string/M-theory.


In the present article, we first observe in section 3 that (double) dimensional reduction is naturally encoded on super L_∞ -algebras by *cyclification*, namely by the process which in terms of rational homotopy theory corresponds to passing to homotopy quotients of free loop spaces by the rotation action on loops. We find that L_∞ -theoretically this (double) dimensional reduction is an *isomorphism*, hence has an inverse (“oxidation”, see [49]) that completely reconstructs the higher dimensional situation (by the L_∞ -incarnation of D0-brane condensation [27, Remark 3.11, 4.6]). In this sense this process is non-perturbative, a fact that is important for our discussion of F-theory further below.

We first use this reduction isomorphism in section 3 to recall from [27] the form of the descended type IIA F1/D p -brane cocycles for $p \in \{0, 2, 4\}$ as the dimensional reduction of the M-brane cocycles from 11d. Then we observe that down in 10d these are enhanced to cocycles for p -branes for $p \in \{0, 2, 4, 8, 10\}$, using the Fierz identity analysis in [17]. We also state the corresponding type IIB F1/D p -brane cocycles. These may be similarly extracted from analysis of Fierz identities [61], but our main theorem below (Theorem 5.3) also implies the form of either (IIA or IIB) from the other.

In section 5 we first compute the dimensional reduction of the type IIA F1/D p -brane cocycles from the 10d super-Minkowski super Lie algebra to 9d. We observe that the two L_∞ -algebras of coefficients of the descended cocycles (for type IIA and type IIB) are manifestly related by an L_∞ -isomorphism (Proposition 5.1.) Then we show in Theorem 5.3 that the action of this isomorphism on the descended type II L_∞ -cocycles implements *T-duality* between the F1/D p brane charges of IIA and type IIB string theory: we discover that the super-cocycles follow the infinitesimal version of the rules of “topological T-duality”, originally proposed by [9] with precise formulation due to [12][13], this is the content of Remarks 5.4 and 6.3 below. Notice that even though the coefficients we obtain are just the rational image of the twisted K-theory that appears in topological T-duality, this is the first time (to the best of our knowledge) that the rules for topological T-duality are actually derived from string theoretic first principles, and that topological T-duality is connected to local spacetime supersymmetry. In fact our derivation shows that the existence and structure of (topological) T-duality acting of F1/D p -brane charges is entirely controlled by higher super Lie algebra theory.

Similarly, we derive the Buscher rules for RR-fields (*Hori's formula*) from first principles this way (Proposition 6.4, Remark 6.5).

Our main theorem (Theorem 5.3) shows that in the category of super L_∞ -algebras T-duality is incarnated as the right part of a diagram of the following form:



The bottom left part of this diagram we discover in section 6. Namely topological T-duality in the alternative formulation of [13] is controlled by the *correspondence space* which is the fiber product of the IIA spacetime with the T-dual IIB spacetime over their joint 9d base. In the string theory literature this is essentially what is known as the *doubled spacetime* [39]. We first show that the correspondence space axiom for topological T-duality due to [13] is satisfied by the relevant super Lie n -algebra extensions of super-Minkowski spacetimes (Proposition 6.2).

Then we show that after stripping off the RR-charge coefficients in 9d, the remaining coefficient L_∞ -algebra is the delooping of the “T-duality Lie 2-algebra” (Def. 7.1, Rem. 7.2). This is the homotopy fiber of the cup product of two universal first Chern classes, in direct analogy to how the “string Lie 2-algebra” that controls the anomaly cancellation in heterotic string theory is the homotopy fiber of the second Chern class/first Pontryagin class [64]. We show that the T-duality Lie 2-algebra has a natural ∞ -action on the direct sum of twisted K^0 and K^1 (Prop. 7.3). Finally we construct the T-duality 2-group-principal 2-bundle over 9d super-Minkowski spacetime that is classified by the descended F1/Dp-brane cocycles, and find that this is equivalent to the total space of the (either) gerbe extension of the T-duality correspondence spacetime (Proposition 7.5 below).

It has been shown in [53] that the principal 2-bundles of which this is the local model space are the right mathematical formulation of Hull's concept of *T-folds* [38] [39]. This we will further discuss elsewhere. Here we just remark that, as we already amplified for T-duality itself, while here we obtain just the infinitesimal/rational image of this structure, we connect (for the first time, to our knowledge) to spacetime super-symmetry and obtain what is in fact the local model for *super T-folds*.

Finally, in section 8 we observe that these doubled super T-correspondence spacetimes still carry the D0-brane L_∞ -cocycle inherited through their type IIA fiber factor. Hence there is a central super extension of this to a super Lie algebra of bosonic dimension 10+2. We show in this last section that this super Lie algebra has the correct properties to be expected of the local model space for an F-theory elliptic fibration, according to [40, 72]. This gives the top left part of the above diagram, below this is Prop. 8.3.

Related literature. Our T-duality takes place on superspaces, for a related discussion see [70]. There are various other approaches to T-duality in relation to branes. The brane worldvolume approach to T-duality transformations as transformations which mix the worldvolume field equations with Bianchi identities is discussed in [58] [69]. T-duality between D-branes is realized on the underlying p -brane solutions of type IIA and type IIB supergravity in [4]. The relation to worldsheet and spacetime supersymmetry is discussed in [3], for the Green-Schwarz superstring in [19] [44], and in the presence of RR fields in [30]. In [31], a superspace with manifest T-duality including Ramond-Ramond gauge fields is presented. The superspace is defined by the double nondegenerate super-Poincare algebras where Ramond-Ramond charges are introduced by central extension. In [56] an interpretation of T-dualization procedure of type II superstring theory in double space is given, taking into account compatibility between supersymmetry and T-duality. A geometry of superspace corresponding to double field theory for type II supergravity is introduced in [16] based on an orthosymplectic extension $\text{OSp}(d, d|2s)$ of the continuous T-duality group.

2. Supersymmetry super Lie n -algebras

Here we introduce what we need below on super Lie n -algebras associated with supersymmetry. Similarly to how L_∞ -algebras are extensions of Lie algebras to include higher brackets, super L_∞ -algebras are extensions of Lie superalgebras to include higher graded brackets. These turn out to be

defined via more familiar differential graded (DG) algebras when we restrict to finite-dimensional super-vector spaces.

Definition 2.1. Write

$$CE : \text{sL}_{\infty}^{\text{fin}} \text{Alg}_{\mathbb{R}} \hookrightarrow \text{dgAlg}_{\mathbb{R}}^{\text{op}}$$

for the full subcategory of the opposite of that of differential graded-algebras augmented over \mathbb{R} (i.e. DG \mathbb{R} -algebras equipped with a algebra homomorphism to \mathbb{R}) whose underlying graded algebra is freely generated as a graded super-commutative algebra on a \mathbb{Z} -graded super-vector space which is degreewise finite dimensional. This is the category of *super L_{∞} -algebras of finite type*.

If we consider just \mathbb{R} -algebras $\text{Alg}_{\mathbb{R}}$ instead of augmented \mathbb{R} -algebras $\text{Alg}_{/\mathbb{R}}$, then the analogous full inclusion

$$CE : \text{sL}_{\infty}^{\text{cvd,fin}} \text{Alg}_{\mathbb{R}} \hookrightarrow \text{dgAlg}_{\mathbb{R}}^{\text{op}}$$

is called that of possibly *curved L_{∞} -algebras of finite type*, with possibly *curved* homomorphism between them.

Under the functor $\text{dgAlg}_{/\mathbb{R}}^{\text{op}} \rightarrow \text{dgAlg}_{\mathbb{R}}^{\text{op}}$ which forgets the augmentation, then every L_{∞} -algebra is regarded as a curved L_{∞} -algebra with *vanishing curvature* and as such there are then possibly curved homomorphisms between non-curved L_{∞} -algebras.

Remark 2.2. This means that for $\mathfrak{g} \in \text{sL}_{\infty} \text{Alg}_{\mathbb{R}}$ a super L_{∞} -algebra, then every generator of its Chevalley-Eilenberg algebra $CE(\mathfrak{g})$ carries a bidegree $(n, \sigma) \in \mathbb{Z} \times (\mathbb{Z}/2)$ and for two such elements the following holds

$$\omega_1 \wedge \omega_2 = (-1)^{n_1 n_2 + \sigma_1 \sigma_2} \omega_2 \wedge \omega_1.$$

(these signs are as in [15, II.2.109] and [22, appendix §6]). These DG-algebras $CE(\mathfrak{g})$ are the “Cartan integrable systems” of [20] and the “free differential algebras” (FDAs) of [73] [15, III.6].³

By forming the linear dual of a differential graded algebra of finite type, it equivalently becomes a differential graded co-algebra. That every L_{∞} -algebra gives a differential co-algebra is originally due to [45] and that this

³It is however crucial that they are not in general free as differential algebras, but just as graded-commutative algebras. In rational homotopy theory one also speaks of “quasi-free” or “semi-free” dg-algebras.

faithfully reflects the original L_∞ -algebra is due to [46], see [63, around def. 13]. In more modern parlance this is due to the Koszul duality between the operads for Lie algebras and that for commutative algebras.

While differential co-algebras are less familiar in practice, they have the advantage that they immediately reflect also L_∞ -algebras not of finite type. This gives a full inclusion

$$\mathrm{sL}_\infty^{\mathrm{fin}} \mathrm{Alg}_{\mathbb{R}} \hookrightarrow \mathrm{sL}_\infty \mathrm{Alg}_{\mathbb{R}}$$

into the category of possibly degreewise infinite dimensional super L_∞ -algebras, and similarly for the curved case

$$\mathrm{sL}_\infty^{\mathrm{cvd}, \mathrm{fin}} \mathrm{Alg}_{\mathbb{R}} \hookrightarrow \mathrm{sL}_\infty^{\mathrm{cvd}} \mathrm{Alg}_{\mathbb{R}}.$$

Notice that there is a fully faithful inclusion

$$\mathrm{sLieAlg}_{\mathbb{R}} \hookrightarrow \mathrm{sL}_\infty \mathrm{Alg}_{\mathbb{R}}$$

of ordinary super Lie algebras into super L_∞ -algebras, whose image is those \mathfrak{g} for which all generators in $\mathrm{CE}(\mathfrak{g})$ are in degree $(1, \sigma)$, for some super-degree σ . Notice that for \mathfrak{g} a super L_∞ -algebra structure on a graded super vector space V , its CE-differential may be co-restricted to its co-unary piece (which sends single generators to single generators). This is the super cochain complex dual to the *underlying super chain complex* of the L_∞ -algebra.

Example 2.3. For $n \in \mathbb{N}$ we write $b^n \mathbb{R}$ for the super L_∞ -algebra for which $\mathrm{CE}(b^n \mathbb{R})$ has a single generator in bidegree $(n+1, \text{even})$, and vanishing differential (the “line Lie n -algebra”). For \mathfrak{g} any super L_∞ -algebra, we call a homomorphism

$$\mu : \mathfrak{g} \longrightarrow b^n \mathbb{R}$$

an L_∞ -cocycle of degree $(n+1)$ on \mathfrak{g} with coefficients in \mathbb{R} . Because dually this is, by definition, a closed element in $\mathrm{CE}(\mathfrak{g})$ of degree $(n+1)$ (see [63, section 6.3]).

We are going to use homotopy theory of super L_∞ -algebras. A standard method to present such are model categories ([35]), but for our purposes here a more lightweight structure is fully sufficient: that of a Brown category of fibrant objects. See [55, Def. 3.54] for review in a context that we are concerned with here.

Proposition 2.4 ([59]). *There is a model category whose category of fibrant objects is precisely the category $sL_\infty\text{Alg}_\mathbb{R}$ of super L_∞ -algebras, and such that on these the weak equivalences are the morphisms for which the underlying morphism of dual super chain complexes (Remark 2.2) is a quasi-isomorphism, and whose fibrations are the morphisms that induce a surjection on the underlying chain complexes. In terms of the dual Chevalley-Eilenberg algebras, this corresponds to an injection on the graded linear subspaces spanned by the generators.*

Proof. By [59, Prop 4.36, Prop. 4.42] there is a model category for ordinary (i.e. bosonic) L_∞ -algebras with these properties. By chasing through the proofs there, one finds that they immediately generalize to the super-algebraic situation. \square

Example 2.5 ([27, Prop. 3.5]). For \mathfrak{g} any super L_∞ -algebra and

$$\mu_{p+2} : \mathfrak{g} \longrightarrow b^{p+1}\mathbb{R}$$

any homomorphism into the line Lie $(p + 1)$ -algebra (a $(p + 2)$ -cocycle on \mathfrak{g} , Example 2.3), the CE-algebra $\text{CE}(\widehat{\mathfrak{g}})$ of its homotopy fiber $\widehat{\mathfrak{g}} \xrightarrow{\text{hofib}(\mu_{p+2})} \mathfrak{g}$ is given from that of $\text{CE}(\mathfrak{g})$ by adjoining a single generator b_{p+1} in degree $p + 1$ and extending the differential by

$$d_{\widehat{\mathfrak{g}}} b_{p+1} = \mu_{p+2},$$

i.e.

$$\text{CE}(\widehat{\mathfrak{g}}) \simeq \text{CE}(\mathfrak{g}) [b] / (db_{p+1} = \mu_{p+2}).$$

Notice that for \mathfrak{g} an ordinary super Lie algebra, and for $p = 0$, i.e. for the case of a 2-cocycle, then $\widehat{\mathfrak{g}}$ thus defined is simply the ordinary central extension of super Lie algebras classified by the 2-cocycle. Therefore in the general case we may think of the homotopy fiber $\widehat{\mathfrak{g}}$ as the *higher central extension* of super L_∞ -algebras classified by a super L_∞ -cocycle.

We are interested in *supersymmetry* super Lie algebras, and their extensions to super L_∞ -algebras. To be self-contained, we briefly collect now some basics on Majorana spinors and spacetime supersymmetry algebras, as well as their interrelation, as the spacetime dimension ranges from 11 down to 9. We use conventions as in [15, II.7.1], except for the first two points to follow, where we use the opposite signs. This means that our Clifford matrices behave as in [15, II.7.1], the only difference is in a sign when raising a spacetime index or lowering a spacetime index.

Definition 2.6.

- (i) The Lorentzian spacetime metric is $\eta := \text{diag}(-1, +1, +1, +1, \dots)$.
- (ii) The Clifford algebra relation is $\Gamma_a \Gamma_b + \Gamma_b \Gamma_a = -2\eta_{ab}$.
- (iii) The timelike index is $a = 0$, the spacelike indices range $a \in \{1, \dots, d - 1\}$.
- (iv) A unitary Dirac representation of $\text{Spin}(d - 1, 1)$ is on \mathbb{C}^{2^ν} where $d \in \{2\nu, 2\nu + 1\}$, via Clifford matrices such that $\Gamma_0^\dagger = \Gamma_0$ and $\Gamma_a^\dagger = -\Gamma_a^\dagger$ for $a \geq 1$.
- (v) For $\psi \in \text{Mat}_{\nu \times 1}(\mathbb{C})$ a complex spinor, we write $\bar{\psi} := \psi^\dagger \Gamma_0$ for its Dirac conjugate. If we have Majorana spinors forming a real sub-representation S then restricted to these the Dirac conjugate coincides with the Majorana conjugate $\psi^\dagger \Gamma_0 = \psi^T C$ (where C is the Charge conjugation matrix).

As usual we write

$$\Gamma_{a_1 \dots a_p} := \frac{1}{p!} \sum_{\text{permutations } \sigma} (-1)^{|\sigma|} \Gamma_{a_{\sigma(1)}} \dots \Gamma_{a_{\sigma(p)}}$$

for the anti-symmetrization of products of Clifford matrices. These conventions imply that all Γ_a are self-conjugate with respect to the pairing $\overline{(-)}(-)$, hence that

$$(\bar{\psi} \Gamma_{a_1 \dots a_p} \psi)^* = (-1)^{p(p-1)/2} \bar{\psi} \Gamma_{a_1 \dots a_p} \psi$$

holds for all ψ . This means that the following expressions are real numbers

$$\begin{aligned} \bar{\psi} \psi, \quad \bar{\psi} \Gamma_a \psi, \quad i \bar{\psi} \Gamma_{a_1 a_2} \psi, \quad i \bar{\psi} \Gamma_{a_1 a_2 a_3} \psi, \\ \bar{\psi} \Gamma_{a_1 \dots a_4} \psi, \quad \bar{\psi} \Gamma_{a_1 \dots a_5} \psi, \quad i \bar{\psi} \Gamma_{a_1 \dots a_6} \psi, \quad \dots \end{aligned}$$

Definition 2.7. Given $d \in \mathbb{N}$ and N a real $\text{Spin}(d - 1, 1)$ -representation (hence some direct sum of Majorana and Majorana-Weyl representations), the corresponding super-Minkowski super Lie algebra

$$\mathbb{R}^{d-1, 1|N} \in \text{sLieAlg}_{\mathbb{R}}$$

is the super Lie algebra defined by the fact that its Chevalley-Eilenberg algebra is the $(\mathbb{N}, \mathbb{Z}/2)$ -bigraded-commutative differential algebra generated from elements $\{e^a\}_{a=0}^{d-1}$ in bidegree $(1, \text{even})$ and from elements $\{\psi^\alpha\}_{\alpha=1}^{\dim N}$ in

bidegree (1, odd) with differential given by

$$d\psi^\alpha = 0, \quad de^a = \bar{\psi} \wedge \Gamma^a \psi.$$

Here on the right we use the spinor-to-vector bilinear pairing, regarded as a super 2-form, i.e. in terms of the charge conjugation matrix C this is

$$\bar{\psi} \wedge \Gamma^a \psi = (C\Gamma^a)_{\alpha\beta} \psi^\alpha \wedge \psi^\beta,$$

where summation over repeated indices is understood. Notice that we do not include a factor of $\frac{1}{2}$ in the definition of de^a .

Definition 2.8. Let $\{\gamma_a\}_{a=0}^{d-1}$ be a Dirac representation on \mathbb{C}^{16} of the Lorentzian $d = 9$ Clifford algebra as above. We obtain a Dirac representation of the $d = 10$ and $d = 11$ Clifford algebra by taking the following block matrices acting on $\mathbb{C}^{16} \oplus \mathbb{C}^{16}$

$$\Gamma_{a \leq 8} := \begin{pmatrix} 0 & \gamma^a \\ \gamma^a & 0 \end{pmatrix}, \quad \Gamma_9 := \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}, \quad \Gamma_{10} := \begin{pmatrix} iI & 0 \\ 0 & -iI \end{pmatrix},$$

where I is the identity matrix.

Remark 2.9. The unique irreducible Majorana representation of $\text{Spin}(10, 1)$ is of real dimension 32. Under the inclusions

$$\text{Spin}(8, 1) \hookrightarrow \text{Spin}(9, 1) \hookrightarrow \text{Spin}(10, 1)$$

this representation branches as

$$\mathbf{32} \mapsto \mathbf{16} \oplus \overline{\mathbf{16}} \mapsto \mathbf{16} \oplus \mathbf{16},$$

where in the middle $\mathbf{16}$ and $\overline{\mathbf{16}}$ are the left and right chiral Majorana-Weyl representations in 10d, while on the right the $\mathbf{16}$ is again the unique irreducible real representation in 9d. Under this branching we decompose a Majorana spinor $\psi \in \mathbf{32}$ as

$$\psi = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}$$

with $\psi_1 \in \mathbf{16}$ and $\psi_2 \in \overline{\mathbf{16}}$ or $\mathbf{16}$.

When we consider T-duality and S-duality below, the Clifford algebra generators will receive various re-interpretations. To make this transparent we introduce the following notation.

Definition 2.10. Define another set of matrices $\{\Gamma_a^B\}_{a=0}^9$ by

$$\Gamma_a^B := \begin{cases} \Gamma_a & | a \leq 8, \\ \begin{pmatrix} 0 & \mathbf{I} \\ \mathbf{I} & 0 \end{pmatrix} & | a = 9. \end{cases}$$

For emphasis we write the original matrices also as $\Gamma_a^A := \Gamma_a$, for $a \leq 9$.

Moreover we also write

$$\sigma_1 := \Gamma_9, \quad \sigma_2 := -\Gamma_9\Gamma_{10}, \quad \sigma_3 := \Gamma_{10}.$$

Remark 2.11. The matrices $\{\Gamma_a^B\}_{a=0}^9$ in Def. 2.10 do not represent a Clifford algebra, but the product of any even number of them represents the correct such product acting on $\mathbf{16} \oplus \mathbf{16}$. For instance $\exp(\omega^{ab}\Gamma_{ab}^B)$ are the elements of the $\text{Spin}(d-1, 1)$ -representation on $\mathbf{16} \oplus \mathbf{16}$. Also, for *odd* $p = 2k + 1$, each of the pairings

$$\overline{\psi}\Gamma_{a_1 \dots a_p}^B \psi = \psi^\dagger \Gamma_0^B \Gamma_{a_1 \dots a_p}^B \psi$$

is the sum of the corresponding pairings on two copies of $\mathbf{16}$.

Remark 2.12. (i) By Def. 2.10 and Def. 2.8 we have

$$\Gamma_9^B = i\Gamma_9\Gamma_{10} = \Gamma_9\Gamma_{11}$$

or, equivalently,

$$\Gamma_9 = i\Gamma_9^B\Gamma_{10}.$$

This simple relation is crucial in the proof of T-duality in Theorem 5.3.

(ii) This relation also makes it manifest that Γ_9^B commutes not only with all Γ_{ab}^B for $a, b \leq 8$, but also with all $\Gamma_a^B\Gamma_9^B$. Consequently, Γ_{10} as well as Γ_9 are invariant under the IIB Spin-action, in that (with the notation in Def. 2.10)

$$\exp(-\omega^{ab}\Gamma_{ab}^B) \sigma_i \exp(\omega^{ab}\Gamma_{ab}^B) = \sigma_i$$

for $i \in \{1, 2, 3\}$.

(iii) Conversely, rotation in the (9, 10)-plane leaves all the Γ_a^B invariant, in that

$$\exp(-\frac{\alpha}{4}\Gamma_9\Gamma_{10}) \Gamma_a^B \exp(\frac{\alpha}{4}\Gamma_9\Gamma_{10}) = \Gamma_a^B.$$


Definition 2.13. We write $\mathbb{R}^{10,1|32}$ (in M-theory) and $\mathbb{R}^{9,1|16+\overline{16}}$ (in type IIA) and $\mathbb{R}^{9,1|16+16}$ (in type IIB) and $\mathbb{R}^{8,1|16+16}$ (in common 9d) for the super-Minkowski super Lie algebras (Def. 2.7) given by the Spin representations of Remark 2.9.

We now start with an observation relating the algebraic structures of super-Minkowski spacetimes (Def. 2.7) in dimensions nine and ten.


Proposition 2.14. *The bilinear spinor-to-vector pairings $\psi \mapsto (\overline{\psi}\Gamma^a\psi)$ in dimensions 11 and 10 constitute 2-cocycles on the super-Minkowski super Lie algebras of one dimension lower (Def. 2.13):*

- (i) $c_2^M := \overline{\psi} \wedge \Gamma_{10}\psi \in \text{CE}(\mathbb{R}^{9,1|16+\overline{16}});$
- (ii) $c_2^{\text{IIA}} := \overline{\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}} \wedge \Gamma_9^{\text{IIA}} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \in \text{CE}(\mathbb{R}^{8,1|16+16});$
- (iii) $c_2^{\text{IIB}} := \overline{\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}} \wedge \Gamma_9^{\text{IIB}} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \in \text{CE}(\mathbb{R}^{8,1|16+16}).$

Moreover, these 2-cocycles classify consecutive central super Lie algebra extensions of super-Minkowski spacetime from 9d to 11d (Def. 2.13) in that we get the following diagram of super L_∞ -algebras



where each “hook”



corresponds to the central extension $\widehat{\mathfrak{g}}$ of \mathfrak{g} classified by the 2-cocycle ω_2 (i.e. is a homotopy fiber sequence of super L_∞ -algebras, according to Example 2.5).

Proof. To see that the given 2-forms are indeed cocycles: they are trivially closed (by Def. 2.7), and so all that matters is that we have a well-defined super-2-form in the first place. Since the ψ^α are in bidegree (1, odd), they all commute with each other. and hence the condition is that the spinor-to-vector pairing is symmetric. This is the case for Majorana spinors. (This is a simple but deep fact, highlighted before in [17, (2.4)], [27, Prop. 4.5]).

Now we consider the extensions. Notice that for \mathfrak{g} any super Lie algebra (of finite dimension), and for $\omega \in \wedge^2 \mathfrak{g}^*$ a Lie algebra 2-cocycle on it, the Lie algebra extension $\widehat{\mathfrak{g}}$ that this classifies is neatly characterized in terms of its dual Chevalley-Eilenberg algebra; it is simply the original CE algebra with one new generator e in degree (1, even) adjoined, and with the differential of e taken to be ω :

$$\text{CE}(\widehat{\mathfrak{g}}) = ((\text{CE}(\mathfrak{g}) \otimes \langle e \rangle), de = \omega).$$

Hence in the case of $\omega = c_2^{\text{IIA}}$ we identify the new generator with e^9 . Furthermore, we see from Prop. 2.12 that the equation $de^9 = c_2^{\text{IIA}}$ is precisely what distinguishes the CE-algebra of $\mathbb{R}^{8,1|\mathbf{16}+\mathbf{16}}$ from that of $\mathbb{R}^{9,1|\mathbf{16}+\mathbf{16}}$. This follows by Def. 2.7, Def. 2.8, and using the fact that the spinors all have the same underlying representation space by Remark 2.9.

The other two cases are directly analogous. \square

3. Double dimensional reduction

In this section we formalize the concept of *double dimensional reduction* of branes on a circle bundle (originally due to [23]) as an equivalence between higher homotopy algebra structures (Theorem 3.8 below), which is at the same time a formalization and a generalization of earlier treatments (as for instance [25]). This may be iterated to deal with reducing on successive circles, i.e. on tori. The free and cyclic loop space construction will play a major role in this formalization (see Remark 3.1 below), which in terms of L_∞ -cohomolgy translates into passage to Hochschild and cyclic L_∞ -homology (see Remark 3.2 below). We close this section below in Remark 3.9 with spelling out how the L_∞ -theoretic statement of Theorem 3.8 indeed reflects the informal idea of double dimensional reduction.

Here we make use of one of the main result of *rational homotopy theory* (see e.g. [33] for review); namely, that the rational homotopy type of a connected topological space X with nilpotent fundamental group is completely and faithfully encoded into a suitable L_∞ -algebra $\mathfrak{l}(X)$, which one may think of as being the ∞ -Lie algebra of the loop group ΩX (itself an “ ∞ -group”, namely a grouplike A_∞ -space). The *Sullivan model* dg-algebras of rational homotopy theory [71], are nothing but the CE-algebras of (particularly good representatives) of connective L_∞ -algebras:

| topological space | loop ∞ -group | L_∞ -algebra | Sullivan model |
|-------------------|----------------------|---------------------|----------------------------|
| X | ΩX | $\mathfrak{l}(X)$ | $\text{CE}(\mathfrak{l}X)$ |

The observation that traditional rational homotopy theory sits inside the homotopy theory of L_∞ -algebras was implicit already in the original [60], but was made fully explicit only in [34], on which the model [59] is based, which we use in Proposition 2.4. A review of rational homotopy theory from this modern perspective is in [11, section 2]. For more exposition in our context see also [29, appendix A].

We recall the following fact due to Vigué et al.:

Remark 3.1 (Sullivan model for free and cyclic loop spaces). The *cyclic loop space* of a topological space X is the homotopy quotient $\mathcal{L}X/S^1$, where $\mathcal{L}X := \text{Maps}(S^1, X)$ denotes the free loop space of X and the S^1 -action is given by rotation of loops. If X is simply connected, and $(\wedge^\bullet V, d_x)$ is a minimal Sullivan model for the rationalization of X , then a Sullivan model for the rationalization of the free loop space $\mathcal{L}X$ of X is given by [75]

$$\text{CE}(\mathfrak{l}(\mathcal{L}X)) = (\wedge^\bullet (V \oplus sV), d_{\mathcal{L}X}),$$

where sV is V with degrees shifted down by one, and with $d_{\mathcal{L}X}$ acting for $v \in V$ as $d_{\mathcal{L}X} v = d_x v$, $d_{\mathcal{L}X} sv = -s d_x v$ where on the right $s: V \rightarrow sV$ is extended uniquely as a graded derivation. A Sullivan model for the rationalization of the cyclic space $\mathcal{L}X/S^1$ is given by [74]

$$\text{CE}(\mathfrak{l}(\mathcal{L}X/S^1)) = (\wedge^\bullet (V \oplus sV \oplus \langle \omega_2 \rangle), d_{\mathcal{L}X/S^1})$$

with $d_{\mathcal{L}X/S^1} \omega_2 = 0$ and with $d_{\mathcal{L}X/S^1}$ acting on $w \in \wedge^1 V \oplus sV$ as $d_{\mathcal{L}X/S^1} w = d_{\mathcal{L}X} w + \omega_2 \wedge sw$. Moreover, the canonical sequence of L_∞ -homomorphisms

$$\mathfrak{l}(\mathcal{L}X) \longrightarrow \mathfrak{l}(\mathcal{L}X/S^1) \longrightarrow b\mathbb{R}$$

is a rational model for the homotopy fiber sequence $\mathcal{L}X \rightarrow \mathcal{L}X/S^1 \rightarrow BS^1$ that exhibits the homotopy quotient.

Remark 3.2 (Jones’ theorem). The Sullivan models for free and cyclic loop spaces as in Remark 3.1 will appear (in a super L_∞ -theoretic generalization to follow in Definition 3.3) below in Theorem 3.8 purely as part of an L_∞ -theoretic incarnation of the concept of double dimensional reduction. All the more does it seem interesting to briefly recall the close relation of these dg-algebras to Hochschild homology and to cyclic homology. This is known as *Jones’ theorem* [41] (see [48] for expository review): Let X be a simply connected topological space. Then the ordinary cohomology $H^\bullet(-)$ of its free loop space is isomorphic to the Hochschild homology HH_\bullet of its dg-algebra $C^\bullet(X)$ of singular cochains:

$$H^\bullet(\mathcal{L}X) \simeq \mathrm{HH}_\bullet(C^\bullet(X)).$$

Similarly, the S^1 -equivariant cohomology of the free loop space, hence the cohomology of the cyclic loop space, is isomorphic to the cyclic homology $\mathrm{HC}_\bullet(-)$ of the cochains:

$$H^\bullet(\mathcal{L}X/S^1) \simeq \mathrm{HC}_\bullet(C^\bullet(X)).$$

Specifically if X happens to carry the structure of a smooth manifold, then the dg-algebra of singular chains with real coefficients is quasi-isomorphic to the de-Rham dg-algebra of smooth differential forms $\Omega^\bullet(X)$, and hence in this case

$$H^\bullet(\mathcal{L}X, \mathbb{R}) \simeq \mathrm{HH}_\bullet(\Omega^\bullet(X)) \quad \text{and} \quad H^\bullet(\mathcal{L}X/S^1, \mathbb{R}) \simeq \mathrm{HC}_\bullet(\Omega^\bullet(X)).$$

Therefore in this case Remark 3.1 together with the central theorem of Quillen-Sullivan rational homotopy says that the cochain cohomology of the Chevalley-Eilenberg complexes of the L_∞ -algebras $\mathfrak{l}(\mathcal{L}X)$ and $\mathfrak{l}(\mathcal{L}X/S^1)$ coincide with the Hochschild homology and the cyclic homology, respectively, of the CE-algebra of the L_∞ -algebra $\mathfrak{l}(X)$:

$$\begin{aligned} H^\bullet(\mathrm{CE}(\mathfrak{l}(\mathcal{L}X))) &\simeq \mathrm{HH}_\bullet(\mathrm{CE}(\mathfrak{l}(X))) \\ \text{and } H^\bullet(\mathrm{CE}(\mathfrak{l}(\mathcal{L}X/S^1))) &\simeq \mathrm{HC}_\bullet(\mathrm{CE}(\mathfrak{l}(X))). \end{aligned}$$

Remark 3.1 motivates the following generalization:

Definition 3.3. Let \mathfrak{h} be a super L_∞ -algebra.

1) The *free loop algebra* of \mathfrak{h} is the super L_∞ -algebra defined by

$$\text{CE}(\mathfrak{L}\mathfrak{h}) = (\wedge^\bullet(\mathfrak{h}^* \oplus s\mathfrak{h}^*), d_{\mathfrak{L}\mathfrak{h}}),$$

where $s\mathfrak{h}^*$ is a copy of \mathfrak{h}^* with bi-degrees shifted by $(-1, \text{even})$, with $d_{\mathfrak{L}\mathfrak{h}}$ acting as

$$\begin{aligned} d_{\mathfrak{L}\mathfrak{h}}v &= d_{\mathfrak{h}}v \\ d_{\mathfrak{L}\mathfrak{h}}sv &= -sd_{\mathfrak{h}}v \end{aligned}$$

for all $v \in \mathfrak{h}^*$, where s is the graded derivation of degree $(-1, \text{even})$ which sends the original generators $v \in \mathfrak{h}^*$ to their shifted image $sv \in s\mathfrak{h}^*$.

2) The *cyclification* of \mathfrak{h} is the L_∞ -algebra $\mathfrak{L}\mathfrak{h}/\mathbb{R}$ defined by

$$\text{CE}(\mathfrak{L}\mathfrak{h}/\mathbb{R}) = (\wedge^\bullet(\mathfrak{h}^* \oplus s\mathfrak{h}^* \oplus \langle \omega_2 \rangle), d_{\mathfrak{L}\mathfrak{h}/\mathbb{R}})$$

with

$$d_{\mathfrak{L}\mathfrak{h}/\mathbb{R}} \omega_2 = 0$$

and with $d_{\mathfrak{L}\mathfrak{h}/\mathbb{R}}$ acting on $w \in \mathfrak{h} \oplus s\mathfrak{h}$ as

$$d_{\mathfrak{L}\mathfrak{h}/\mathbb{R}} w = d_{\mathfrak{L}\mathfrak{h}} w + \omega_2 \wedge sw.$$

Lemma 3.4. *The free loop algebra construction from Definition 3.3 extends to a functor on the category of super L_∞ -algebras*

$$\mathfrak{L}(-): sL_\infty\text{Alg}_{\mathbb{R}} \longrightarrow sL_\infty\text{Alg}_{\mathbb{R}}$$

by taking a super L_∞ -homomorphism $f: \mathfrak{h}_1 \rightarrow \mathfrak{h}_2$ to the homomorphism $\mathfrak{L}f$ whose dual $(\mathfrak{L}f)^*$ is given on generators $v \in \wedge^1\mathfrak{h}_2^*$ by

$$\begin{aligned} (\mathfrak{L}f)^*(v) &= f^*(v) \\ (\mathfrak{L}f)^*(s_2v) &= s_1f^*(v). \end{aligned}$$

Similarly the cyclification operation from Definition 3.3 extends to a functor

$$\mathfrak{L}(-)/\mathbb{R}: sL_\infty\text{Alg}_{\mathbb{R}} \longrightarrow sL_\infty\text{Alg}_{\mathbb{R}}$$

by taking a homomorphism f to the homomorphism $\mathfrak{L}f/\mathbb{R}$ whose dual is given on v and sv by $(\mathfrak{L}f)^*$ as before, and which sends the copy of the generator ω_2 in $\text{CE}(\mathfrak{L}\mathfrak{h}_2/\mathbb{R})$ to the generator of the same name in $\text{CE}(\mathfrak{L}\mathfrak{h}_1/\mathbb{R})$.

Proof. To see that $\mathfrak{L}(-)$ is functorial on homomorphisms of graded algebras it is sufficient to observe that the relation $(\mathfrak{L}f)^*(s_2v) = s_1(\mathfrak{L}f)^*(v)$ on generators implies that $(\mathfrak{L}f)^*$ commutes with s generally. For instance on binary wedge products of generators we get

$$\begin{aligned} (\mathfrak{L}f)^*(s(v_1 \wedge v_2)) &= (\mathfrak{L}f)^*((sv_1) \wedge v_2 + (-1)^{\deg(v_1+1)}(sv_2)) \\ &= (\mathfrak{L}f)^*(sv_1) \wedge (\mathfrak{L}f)^*(v_2) \\ &\quad + (-1)^{\deg(v_1)+1}(\mathfrak{L}f)^*(v_1) \wedge (\mathfrak{L}f)^*(sv_2) \\ &= s((\mathfrak{L}f)^*(v_1)) \wedge (\mathfrak{L}f)^*(v_2) \\ &\quad + (-1)^{\deg((\mathfrak{L}f)^*(v_1))+1}(\mathfrak{L}f)^*(v_1) \wedge s((\mathfrak{L}f)^*(v_2)) \\ &= s(\mathfrak{L}f)^*(v_1 \wedge v_2). \end{aligned}$$

It remains to see that $(\mathfrak{L}f)^*$ respects the differential. On the unshifted generators v this is so because f^* does respect the differential. For shifted generators it follows by this computation:

$$\begin{aligned} (\mathfrak{L}f)^*(d_{\mathfrak{L}\mathfrak{h}_2}s_2v) &= -(\mathfrak{L}f)^*(s_2d_{\mathfrak{L}\mathfrak{h}_2}v) \\ &= -s_1(\mathfrak{L}f)^*(d_{\mathfrak{h}_2}v) \\ &= -s_1f^*(d_{\mathfrak{h}_2}v) \\ &= -s_1d_{\mathfrak{h}_1}f^*(v) \\ &= -s_1d_{\mathfrak{L}\mathfrak{h}_1}f^*(v) \\ &= d_{\mathfrak{L}\mathfrak{h}_1}s_1(\mathfrak{L}f)^*(v) \\ &= d_{\mathfrak{L}\mathfrak{h}_1}(\mathfrak{L}f)^*(s_2v) \end{aligned}$$

Finally, that also $(\mathfrak{L}f/\mathbb{R})^*$ respects the differential follows by this computation:

$$\begin{aligned} (\mathfrak{L}f/\mathbb{R})^*(d_{\mathfrak{L}\mathfrak{h}_2/\mathbb{R}}w) &= (\mathfrak{L}f/\mathbb{R})^*(d_{\mathfrak{L}\mathfrak{h}_2}w + \omega_2 \wedge sw) \\ &= (\mathfrak{L}f)^*(d_{\mathfrak{L}\mathfrak{h}_2}w) + (\mathfrak{L}f/\mathbb{R})^*(\omega_2 \wedge sw) \\ &= d_{\mathfrak{L}\mathfrak{h}_2}(\mathfrak{L}f)^*(w) + \omega_2 \wedge (\mathfrak{L}f)^*(sw) \\ &= d_{\mathfrak{L}\mathfrak{h}_2}(\mathfrak{L}f)^*(w) + \omega_2 \wedge s(\mathfrak{L}f)^*(w). \end{aligned}$$

where we used the previous statements about $(\mathfrak{L}f)^*$. □

The two super L_∞ -algebras from Definition 3.3 are related as follows.

Proposition 3.5. *For any super L_∞ -algebra \mathfrak{h} , its free loop algebra and its cyclification (Def. 3.3) sit in a homotopy fiber sequence of the form*

$$\mathfrak{L}\mathfrak{h} \longrightarrow \mathfrak{L}\mathfrak{h}/\mathbb{R} \xrightarrow{\omega_2} b\mathbb{R},$$

where ω_2 is the 2-cocycle given by the CE-element of the same name in Def. 3.3.

Proof. The second morphism is a fibration (according to Proposition 2.4), by the surjection evident from their definition. Moreover, from the form of the differentials, the first morphism is directly seen to be the ordinary fiber of the second. Hence it models the homotopy fiber. \square

The key observation now, noticed in a particular case in [29, Prop. 3.8], is that the cyclification of coefficients is what formalizes “double dimensional reduction” of branes [23], i.e. the joint process of reducing spacetime dimension and reducing in parallel the dimension of branes in this spacetime, or dually, the cohomological degrees of their charges. Or rather: those branes/charges that “wrap” the dimension being reduced are to reduce in parallel, while those that do not wrap should just descend. We formalize this by Theorem 3.8 below, see Remark 3.9 further below for discussion of the physical interpretation. To state the formalization, we first need to make explicit the following basic construction:

Definition 3.6. Given any super L_∞ -algebra $\mathfrak{b} \in sL_\infty\text{Alg}_{\mathbb{R}}$, then the *slice over \mathfrak{g}* is the category $(sL_\infty\text{Alg}_{\mathbb{R}})_{/\mathfrak{b}}$ whose objects are super L_∞ -homomorphisms $\mathfrak{g} \xrightarrow{\phi} \mathfrak{b}$ into \mathfrak{b} , and whose morphisms are super L_∞ -homomorphisms $\mathfrak{g}_1 \rightarrow \mathfrak{g}_2$ that respect the morphisms down to \mathfrak{b} in that they make the diagrams shown on the right commute:

$$\text{Hom}_{/\mathfrak{b}}((\mathfrak{g}_1, \phi_1), (\mathfrak{g}_2, \phi_2)) := \left\{ \begin{array}{ccc} \mathfrak{g}_1 & \text{---} & \mathfrak{g}_2 \\ & \searrow \phi_1 & \swarrow \phi_2 \\ & & \mathfrak{b} \end{array} \right\}.$$

Example 3.7. Consider the operation that sends any super L_∞ -homomorphism of the form

$$\mu_{p+2}: \mathfrak{g} \longrightarrow b^{p+1}\mathbb{R}$$

(i.e. a super L_∞ -cocycle, according to Example 2.3) to its homotopy fiber $\widehat{\mathfrak{g}}$ represented via Example 2.5

$$\text{CE}(\widehat{\mathfrak{g}}) := \text{CE}(\mathfrak{g})[b_{p+1}]/(db_{p+1} = \mu_{p+2}).$$

This operation extends to a functor on the slice category (Def. 3.6) over $b^{p+1}\mathbb{R}$

$$\text{hofib}: (sL_\infty\text{Alg}_{\mathbb{R}})_{/b^{p+1}\mathbb{R}} \longrightarrow sL_\infty\text{Alg}_{\mathbb{R}},$$

by taking any super L_∞ -homomorphism f in

$$\begin{array}{ccc} \mathfrak{g}_1 & \xrightarrow{f} & \mathfrak{g}_2 \\ & \searrow^{\mu_{p+2}^1} & \swarrow_{\mu_{p+2}^2} \\ & & b^{p+1}\mathbb{R} \end{array}$$

to the homomorphism $\text{hofib}(f)$ whose dual $(\text{hofib}(f))^*$ is given on $\text{CE}(\mathfrak{g}_2)$ by f^* and sends the generator b_{p+1}^2 to b_{p+1}^1 . This respects the differential on the original generators because f^* does, and it respects the differential on the new generator because

$$\begin{aligned} (\text{hofib}(f))^*(d_{\widehat{\mathfrak{g}}_2} b_{p+1}^2) &= (\text{hofib}(f))^*(\mu_{p+2}^2) \\ &= f^*(\mu_{p+2}^2) \\ &= \mu_{p+1}^1 \\ &= d_{\widehat{\mathfrak{g}}_1} b_{p+1}^1 \\ &= d_{\widehat{\mathfrak{g}}_1} (\text{hofib}(f))^* b_{p+1}^2, \end{aligned}$$

where the third equality used is equivalent to the commutativity of the triangular diagram above.

Theorem 3.8 (L_∞ -theoretic double dimensional reduction and oxidation). *Let \mathfrak{g} and \mathfrak{h} be super L_∞ -algebras, such that $\text{CE}(\mathfrak{h})$ has no generators in degree 1. Let moreover $c_2 : \mathfrak{g} \rightarrow b\mathbb{R}$ be a 2-cocycle (Definition 2.3) and $\pi : \widehat{\mathfrak{g}} \rightarrow \mathfrak{g}$ be the central extension classified by c_2 , according to Example 2.5. Then there is a bijection*

$$\text{Hom}(\widehat{\mathfrak{g}}, \mathfrak{h}) \begin{array}{c} \xleftarrow{\text{oxidation}} \\ \xrightarrow[\text{reduction}]{\simeq} \end{array} \text{Hom}_{/b\mathbb{R}}(\mathfrak{g}, \mathfrak{L}\mathfrak{h}/\mathbb{R})$$

between super L_∞ -homomorphisms out of $\widehat{\mathfrak{g}}$ into \mathfrak{h} and super L_∞ -homomorphism over $b\mathbb{R}$ from \mathfrak{g} (in the sense of Definition 3.6) to the cyclification

$\mathfrak{L}\mathfrak{h}/\mathbb{R}$ of \mathfrak{h} (Def. 3.3):

$$\left(\widehat{\mathfrak{g}} \longrightarrow \mathfrak{h} \right) \longleftrightarrow \left(\begin{array}{ccc} \mathfrak{g} & \longrightarrow & \mathfrak{L}\mathfrak{h}/\mathbb{R} \\ & \searrow c_2 & \swarrow \omega_2 \\ & & b\mathbb{R} \end{array} \right).$$

More generally, if we consider \mathfrak{g} and \mathfrak{h} as curved L_∞ -algebras with vanishing curvature, according to Definition 2.1, then the bijection holds without any condition on \mathfrak{h} as a bijection between hom-sets of possibly curved super L_∞ -homomorphism. Moreover, in this case the bijection is natural in its arguments (i.e. compatible with pre- and postcomposition with curved super L_∞ -homomorphisms). In other words, the functors from Example 3.3 (Lemma 3.4) and Example 3.7 then form an adjoint pair (e.g. [8, chapter 3]) with hofib left adjoint to the cyclification functor $\mathfrak{L}(-)/\mathbb{R}$ from Def. 3.3:

$$sL_\infty \text{Alg}_{\mathbb{R}}^{\text{cvd}} \xleftarrow{\text{hofib}} (sL_\infty \text{Alg}_{\mathbb{R}}^{\text{cvd}})_{/b\mathbb{R}} \xrightarrow[\mathfrak{L}(-)/\mathbb{R}]{\perp}$$

Finally, the unit $\eta_{(\mathfrak{g}),\omega_2}$ of this adjunction,

$$\begin{array}{ccc} \mathfrak{g} & \xrightarrow{\eta_{\mathfrak{g},c_2}} & \mathfrak{L}\widehat{\mathfrak{g}}/\mathbb{R} \\ & \searrow c_2 & \swarrow \omega_2 \\ & & b\mathbb{R} \end{array}$$

(i.e. the image under the above correspondence of the identity on some $\widehat{\mathfrak{g}}$) is dually given by the map that

- 1) sends unshifted generators to themselves, except for e (the extra generator with $d_{\widehat{\mathfrak{g}}}e = c_2$ according to example 2.5) which is sent to zero,
- 2) sends all shifted generators to zero, except for the shift $s(e)$ of the extra generator, that instead is sent to minus the algebra unit $s(e) \mapsto -1$ (this makes it a curved L_∞ -morphism).

Proof. We discuss the adjunction for curved L_∞ -homomorphism. The statement for the non-curved case then follows immediately as a special case.

First consider the bijection as such. Dually we need to show that there is an identification

$$\text{Hom}(\text{CE}(\mathfrak{h}), \text{CE}(\widehat{\mathfrak{g}})) \simeq \text{Hom}^{\text{CE}(b\mathbb{R})/}(\text{CE}(\mathfrak{L}\mathfrak{h}/\mathbb{R}), \text{CE}(\mathfrak{g})).$$

between homomorphisms of dg-algebras (not augmented) out of $\text{CE}(\mathfrak{h})$ into $\text{CE}(\widehat{\mathfrak{g}})$ and dg-algebra homomorphisms under $\text{CE}(b\mathbb{R})$ out of $\text{CE}(\mathfrak{L}\mathfrak{h}/\mathbb{R})$ into $\text{CE}(\mathfrak{g})$

$$\left(\text{CE}(\mathfrak{h}) \longrightarrow \text{CE}(\widehat{\mathfrak{g}}) \right) \longleftrightarrow \left(\begin{array}{ccc} & \text{CE}(b\mathbb{R}) & \\ \omega_2 \swarrow & & \searrow c_2 \\ \text{CE}(\mathfrak{L}\mathfrak{h}/\mathbb{R}) & \longrightarrow & \text{CE}(\mathfrak{g}) \end{array} \right).$$

Since, after forgetting the differential, $\text{CE}(\mathfrak{h})$ is a free graded polynomial algebra, we may write

$$\text{CE}(\mathfrak{h}) = (\mathbb{R}[\{x_p\}], d_{\text{CE}(\mathfrak{h})}x_i = P_i(\{x_p\}))$$

for suitable polynomials P_i in the variables $\{x_p\}$. Then a DGCA homomorphism from $\text{CE}(\mathfrak{h})$ to $\text{CE}(\widehat{\mathfrak{g}})$ is just a collection of elements α_p in $\text{CE}(\widehat{\mathfrak{g}})$, one for each generator of $\text{CE}(\mathfrak{h})$, such that

$$d_{\text{CE}(\widehat{\mathfrak{g}})}\alpha_i = P_i(\{\alpha_p\}).$$

Since $\widehat{\mathfrak{g}}$ is the the central extension of \mathfrak{g} classified by ω_2 , we have

$$\text{CE}(\widehat{\mathfrak{g}}) = (\text{CE}(\mathfrak{g})[e], d_{\text{CE}(\widehat{\mathfrak{g}})}e = \gamma_2),$$

where γ_2 is the image of the generator ω_2 of $\mathbb{R}[\omega_2] \simeq \text{CE}(b\mathbb{R})$ under the map $c_2 : \mathbb{R}[\omega_2] \rightarrow \text{CE}(\mathfrak{g})$. Since e has degree 1, we then see that there is a vector space decomposition

$$\text{CE}(\widehat{\mathfrak{g}}) = \text{CE}(\mathfrak{g}) \oplus (e \wedge \text{CE}(\mathfrak{g})).$$

Consequently, we can uniquely decompose all the elements α_p as

$$(1) \quad \alpha_p = \beta_p - e \wedge \tilde{\alpha}_{p-1},$$

with $\tilde{\alpha}_{p-1}, \beta_p \in \text{CE}(\mathfrak{g})$. We denote this by

$$(2) \quad \pi_*(\alpha_p) := \tilde{\alpha}_{p-1} \quad \text{and} \quad \alpha_p|_{\mathfrak{g}} := \beta_p$$

in order to amplify the geometric interpretation. Forgetting the differential, we have

$$\text{CE}(\mathfrak{L}\mathfrak{h}/\mathbb{R}) = \mathbb{R}[\{x_p, y_{p-1}\}, \omega_2],$$

so that the assignment

$$(3) \quad x_p \mapsto \beta_p; \quad y_{p-1} \mapsto \tilde{\alpha}_{p-1}$$

precisely defines a morphism of graded commutative algebras under $\mathbb{R}[\omega_2]$ from $\text{CE}(\mathfrak{L}\mathfrak{h}/\mathbb{R})$ to $\text{CE}(\mathfrak{g})$. That is, the image of the additional generator ω_2 of $\text{CE}(\mathfrak{L}\mathfrak{h}/\mathbb{R})$ is prescribed to be γ_2 by the requirement of having a morphism under $\mathbb{R}[\omega_2]$. We are therefore left with checking that this is indeed a morphism of DGCA's. In terms of the generators $\{x_p, y_{p-1}\}$ and ω_2 , the differential in $\text{CE}(\mathfrak{L}\mathfrak{h}/\mathbb{R})$ reads

$$\begin{aligned} d_{\text{CE}(\mathfrak{L}\mathfrak{h}/\mathbb{R})}x_i &= P_i(\{x_p\}) + \omega_2 \wedge y_{p-1}, \\ d_{\text{CE}(\mathfrak{L}\mathfrak{h}/\mathbb{R})}y_{i-1} &= - \sum_j y_{j-1} \frac{\partial P_i}{\partial x_j}(\{x_p\}), \\ d_{\text{CE}(\mathfrak{L}\mathfrak{h}/\mathbb{R})}\omega_2 &= 0. \end{aligned}$$

The fact that the above assignment is a morphism of DGCA's then follows from the matching

$$\begin{aligned} P_i(\{\beta_p\}) - e \wedge \sum_j \tilde{\alpha}_{j-1} \frac{\partial P_i}{\partial x_j}(\{\beta_p\}) &= P_i(\{\beta_p - e \wedge \tilde{\alpha}_p\}) \\ &= P_i(\{\alpha_p\}) \\ &= d_{\text{CE}(\hat{\mathfrak{g}})}\alpha_i \\ &= d_{\text{CE}(\hat{\mathfrak{g}})}(\beta_i - e \wedge \tilde{\alpha}_{i-1}) \\ &= d_{\text{CE}(\mathfrak{g})}\beta_i - \gamma_2 \wedge \tilde{\alpha}_{i-1} + e \wedge d_{\text{CE}(\mathfrak{g})}\tilde{\alpha}_{i-1}. \end{aligned}$$

This establishes the bijection for any fixed $\hat{\mathfrak{g}}$ and \mathfrak{h} . For this bijection to be *natural* we need to show that for every morphism

$$\begin{array}{ccc} \mathfrak{g}_2 & \xrightarrow{f} & \mathfrak{g}_1 \\ & \searrow (\omega_2)_2 & \swarrow (\omega_2)_1 \\ & & b\mathbb{R} \end{array}$$

in $(sL_\infty\text{Alg}_{\mathbb{R}})_{/b\mathbb{R}}$ (Def. 3.6) and every morphism

$$\mathfrak{h}_1 \xrightarrow{g} \mathfrak{h}_2$$

in $sL_\infty\text{Alg}_\mathbb{R}$ the following diagram (of functions between hom-sets) commutes:

$$\begin{array}{ccc} \text{Hom}(\widehat{\mathfrak{g}}_1, \mathfrak{h}_1) & \xrightarrow[\text{reduction}]{\simeq} & \text{Hom}_{/b\mathbb{R}}(\mathfrak{g}_1, \mathfrak{L}\mathfrak{h}_1/\mathbb{R}) \\ g \circ (-) \circ \text{hofib}(f) \downarrow & & \downarrow \mathfrak{L}g/\mathbb{R} \circ (-) \circ f \\ \text{Hom}(\widehat{\mathfrak{g}}_2, \mathfrak{h}_2) & \xrightarrow[\text{reduction}]{\simeq} & \text{Hom}_{/b\mathbb{R}}(\mathfrak{g}_2, \mathfrak{L}\mathfrak{h}_2/\mathbb{R}), \end{array}$$

where the vertical maps are those given by pre- and postcomposition, as indicated.

By unwinding the definition, one finds that indeed both ways of going around this square take a homomorphism $\phi: \widehat{\mathfrak{g}}_1 \rightarrow \mathfrak{h}_1$ to the homomorphism $\mathfrak{g}_2 \rightarrow \mathfrak{L}\mathfrak{h}_2/\mathbb{R}$ which is given, dually, for any generator $v \in \wedge^1 \mathfrak{h}_1^*$ by

$$\begin{aligned} v &\mapsto f^*((\phi^*(g^*(v)))|_{\mathfrak{g}_1}) \\ sv &\mapsto f^*((\pi_1)_*(\phi^*(g^*(v)))) \end{aligned}$$

where we are using (2).

This establishes the adjunction. Finally we show that the adjunction unit η is as claimed. One way to see this is to apply the above correspondence to the identity morphism on $\widehat{\mathfrak{g}}$. But it is instructive to spell this out: First observe that the map as claimed is indeed well defined: Its dual incarnation respects the differential on e because

$$d_{\mathfrak{L}\widehat{\mathfrak{g}}/\mathbb{R}}e = c_2 + \omega_2 \wedge s(e) \mapsto c_2 + c_2 \wedge (-1) = 0,$$

Showing that this is indeed the adjunction unit is equivalent to showing that image of any $\widehat{\mathfrak{g}} \xrightarrow{\phi} \mathfrak{h}$ under the adjunction correspondence is equal to the composite

$$\mathfrak{g} \xrightarrow{\eta_{\mathfrak{g}, c_2}} \mathfrak{L}\widehat{\mathfrak{g}}/\mathbb{R} \xrightarrow{\mathfrak{L}\phi/\mathbb{R}} \mathfrak{L}\mathfrak{h}/\mathbb{R} .$$

This now follows by direct inspection. By the above, the dual of this composite sends any unshifted generator c_p to

$$c_p \mapsto \alpha_p = \beta_p - e \wedge \tilde{\alpha}_{p-1} \mapsto \beta_p$$

and any shifted generator $s(c_p)$ to

$$s(c_p) \mapsto s(\alpha_p) = s(\beta_p) - s(e) \wedge \tilde{\alpha}_{p-1} - e \wedge s(\tilde{\alpha}_{p-1}) \mapsto -(-1) \wedge \tilde{\alpha}_{p-1} = \tilde{\alpha}_{p-1}.$$

This is indeed as in assignment (3) above. □

Remark 3.9 (Geometric interpretation of L_∞ -algebraic dimensional reduction). The general theory of adjoint functors (e.g. [8, chapter 3]) provides insight as to the geometric nature of the super L_∞ -algebraic formalization of dimensional reduction from Theorem 3.8. Namely given any pair of adjoint functors $L \dashv R$, then the unit of the adjunction $\eta_x : x \rightarrow RLx$ is such that the natural bijection between hom-sets

$$\text{Hom}(Lx, y) \xrightarrow{\cong} \text{Hom}(x, Rx)$$

is given by sending any morphism of the form $\phi : Lx \rightarrow y$ to the composite

$$x \xrightarrow{\eta_x} RLx \xrightarrow{R\phi} Ry.$$

Specified to the situation in Theorem 3.8, this means that the L_∞ -theoretic double dimensional reduction of a super L_∞ -homomorphism

$$\widehat{\mathfrak{g}} \xrightarrow{\phi} \mathfrak{h}$$

on a central \mathbb{R} -extension $\widehat{\mathfrak{g}}$ of some super L_∞ -algebra \mathfrak{g} is the following composite:

$$\mathfrak{g} \xrightarrow{\eta_{\mathfrak{g}}} \mathfrak{L}\widehat{\mathfrak{g}}/\mathbb{R} \xrightarrow{\mathfrak{L}(\phi)/\mathbb{R}} \mathfrak{L}\mathfrak{h}/\mathbb{R}.$$

In terms of the geometric interpretation via rational homotopy theory from Remark 3.1 the morphism $\eta_{\mathfrak{g}}$ here has the following interpretation:

Let $\widehat{X} \rightarrow X$ be a principal circle bundle. Then there is a map

$$X \longrightarrow \mathfrak{L}\widehat{X}/S^1$$

which sends each point of X to the loop that winds around the circle fiber over that point, at unit parameter speed. As a map to the free loop space $\mathfrak{L}\widehat{X}$ this would not be well defined unless the circle bundle were trivial, because by definition of principal circle bundles its fibers are identified with the typical fiber (the circle) only up to rigid rotation of that circle. But this is precisely the relation that is divided out by passing to the cyclified space $\mathfrak{L}\widehat{X}/S^1$, which makes the assignment of points to the loops that wind around their fibers be well defined.

Hence given any map of spaces $\widehat{X} \xrightarrow{f} H$, then we may pass to the induced map on loops in \widehat{X} modulo rigid rotation, and then precompose with

the above fiber-assigning map

$$X \longrightarrow \mathcal{L}\widehat{X}/S^1 \xrightarrow{\mathcal{L}f/S^1} \mathcal{L}H/S^1 .$$

Under the Quillen-Sullivan functor from spaces to their associated L_∞ -algebras in rational homotopy theory, this is the L_∞ -algebraic construction above.

And this shows just how this formalizes the intuitive picture of double dimensional reduction: For let Σ_{p+1} be a manifold of dimension $p + 1$ and let $\Sigma_{p+1} \rightarrow X$ be the worldvolume of some p -brane in X . Thinking of $f : \widehat{X} \rightarrow H$ as classifying a background field for $(p + 1)$ -branes on \widehat{X} (for instance for $H = K(\mathbb{Z}, p + 3)$, the classifying space for ordinary cohomology) then the dimensionally reduced coupling term is given by the composite

$$\Sigma_{p+1} \longrightarrow X \longrightarrow \mathcal{L}\widehat{X}/S^1 \xrightarrow{\mathcal{L}f/S^1} \mathcal{L}H/S^1 .$$

To see what this does, consider what happens locally over some chart U on which the circle extension $\widehat{X} \rightarrow X$ is topologically trivial. Then by the ordinary product/hom adjunction $S^1 \times (-) \dashv [S^1, -] = \mathcal{L}(-)$ this is equivalently the composite

$$\Sigma_{p+1} \times S^1 \longrightarrow \widehat{U} \xrightarrow{f|_U} H .$$

But this is nothing than the value of the background field f not on the p -brane worldvolume Σ_{p+1} , but on the worldvolume $\Sigma_{p+1} \times S^1$ of a $(p + 1)$ -brane, which “wraps” the circle fiber in $\widehat{U} = U \times S^1$. This is precisely the physical picture of double dimensional reduction, originally due to [23].

4. The brane supercocycles

We now naturally associate to systems of super p -brane species certain supercocycles taking values in super L_∞ -algebras arising from spheres and related topological spaces/spectra. This follows the geometric approach to cocycles in supergravity, as in [20] [15]. The algebras for type IIA and type IIB that we obtain may also be found in [14] [17] [61] [21].

Definition 4.1. Write $\mathfrak{IS}^4 \in \mathfrak{sL}_\infty\text{Alg}$ for the Chevalley-Eilenberg algebra which is the minimal Sullivan model of the 4-sphere according to [71]:

$$\text{CE}(\mathfrak{IS}^4) = \left\{ \begin{array}{l} dg_4 = 0 \\ dg_7 = -\frac{1}{2}g_4 \wedge g_4 \end{array} \right\} .$$

Definition 4.2. On the super Lie algebra $\mathbb{R}^{10,1|\mathbf{32}}$ (Def. 2.7, Remark 2.9) define the cochains

$$\begin{aligned} \mu_{M_2} &:= \frac{i}{2} (\bar{\psi} \wedge \Gamma_{a_1 a_2} \psi) \wedge e^{a_1} \wedge e^{a_2}, \\ \mu_{M_5} &:= \frac{1}{5!} (\bar{\psi} \wedge \Gamma_{a_1 \dots a_5} \psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_5}, \end{aligned}$$

where the indices run in the set $\{0, 1, \dots, 10\}$.

Proposition 4.3. *The elements in Def. 4.2 satisfy*

$$d\mu_{M_5} = \frac{1}{2} \mu_{M_2} \wedge \mu_{M_2},$$

hence constitute a super L_∞ -algebra homomorphism to the 4-sphere (Def. 4.1):

$$(\mu_{M_2}, \mu_{M_5}) : \mathbb{R}^{10,1|\mathbf{32}} \longrightarrow \mathbb{S}^4.$$

Proof. Using the differential relations $d\psi^\alpha = 0$ and $de^a = \bar{\psi} \Gamma^a \psi$ from Def. 2.7, we have

$$\begin{aligned} d\mu_{M_5} &= \frac{1}{5!} d \left((\bar{\psi} \Gamma_{a_1 \dots a_5} \psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_5} \right) \\ &= \frac{1}{4!} \underbrace{(\bar{\psi} \Gamma_{[a_1 \dots a_4 a]} \psi) (\bar{\psi} \Gamma^a \psi)}_{=3(\bar{\psi} \Gamma_{[a_1 a_2]} \psi) (\bar{\psi} \Gamma_{a_3 a_4]} \psi)} \wedge e^{a_1} \wedge \dots \wedge e^{a_4} \\ &= -\frac{1}{2} \left(\frac{i}{2} (\bar{\psi} \wedge \Gamma_{a_1 a_2} \psi) \wedge e^{a_1} \wedge e^{a_2} \right) \wedge \left(\frac{i}{2} (\bar{\psi} \wedge \Gamma_{a_3 a_4} \psi) \wedge e^{a_3} \wedge e^{a_4} \right) \\ &= -\frac{1}{2} \mu_{M_2} \wedge \mu_{M_2}, \end{aligned}$$

where the equality under the brace is the Fierz identity from [15, (3.27a)]. □

Remark 4.4. In fact there is a stronger statement: A priori the WZW term for the M5-brane is a 7-cocycle not on $\mathbb{R}^{10,1|\mathbf{32}}$, but on the super Lie 3-algebra **m2brane** which is its higher extension (according to example 2.5) by the 4-cocycle μ_{M_2} [27]:

$$\begin{array}{ccc} \mathbf{m2brane} & \xrightarrow{\mu'_{M_5}} & b^6 \mathbb{R} \\ \text{hofib}(\mu_{M_2}) \downarrow & & \\ \mathbb{R}^{10,1|\mathbf{32}} & \searrow \mu_{M_2} & \\ & & b^3 \mathbb{R} \end{array}$$

Given such a situation, **m2brane** is exhibited as a $b^2\mathbb{R}$ -principal ∞ -bundle over $\mathbb{R}^{10,1|\mathbf{32}}$ [54] and one may ask for the $b^2\mathbb{R}$ -equivariant homotopy descent of μ'_{M5} down to the base space $\mathbb{R}^{9,1|\mathbf{16}+\mathbf{16}}$. As discussed in [28] this exists and is given, up to equivalence of L_∞ -cocycles, by the S^4 -valued cocycle (μ_{M2}, μ_{M5}) from Proposition 4.3.

Example 4.5 ([29, prop. 3.8]). The double dimensional reduction according to Theorem 3.8 of the M2/M5-brane cocycle from Definition 4.3 and Remark 4.4 is a cocycle of the form

$$\begin{array}{ccc}
 \mathbb{R}^{10,1|\mathbf{32}} & \xrightarrow{\{\mu_{F1}^A, \mu_{NS5}^A, \mu_{D2}, \mu_{D4}\}} & \mathcal{L}S^4/b\mathbb{R} \\
 & \searrow \mu_{D0}^A & \swarrow \omega_2 \\
 & & b\mathbb{R}
 \end{array}$$

However, we highlight that there is gauge enhancement: more cocycles appear after dimensional reduction, notably the D6-brane cocycle, with all cocycles being assembled together in a model for rational twisted complex topological K-theory in cohomological degrees 0 and 1. As a precise statement this is Proposition 4.8 below, for which we need the following definitions:

Definition 4.6 ([29, section 4]). Write $\mathfrak{l}(KU/BU(1))$ for the L_∞ -algebra with CE-algebra

$$\text{CE}(\mathfrak{l}(KU/BU(1))) := (\mathbb{R}[h_3, \{\omega_{2p}\}_{p \in \mathbb{Z}}], d\omega_{2p+2} = h_3 \wedge \omega_{2p})$$

and $\mathfrak{l}(\Sigma KU/BU(1))$ for the L_∞ -algebra with CE-algebra

$$\text{CE}(\mathfrak{l}(\Sigma KU/BU(1))) := (\mathbb{R}[h_3, \{\omega_{2p+1}\}_{p \in \mathbb{Z}}], d\omega_{2p+3} = h_3 \wedge \omega_{2p+1}).$$

Starting with cocycles in M-theory on $\mathbb{R}^{9,1|\mathbf{32}}$ we get cocycles in type IIA on $\mathbb{R}^{8,1|\mathbf{16}+\mathbf{16}}$ by integration over the fiber $(\pi_{10})_*$ of the rational circle bundle $S^1_{\mathbb{R}} \rightarrow \mathbb{R}^{9,1|\mathbf{32}} \xrightarrow{\pi_{10}} \mathbb{R}^{8,1|\mathbf{16}+\mathbf{16}}$, established in [27, Prop. 4.5].

Definition 4.7. We denote the components of the double dimensional reduction of the M-brane cocycles as follows, using the notation in Eq. (2):

$$\begin{aligned} \mu_{D_0} &:= \bar{\psi}\Gamma^{10}\psi \\ &= \bar{\psi}\Gamma_{10}\psi, \\ \mu_{F_1}^A &:= (\pi_{10})_*(\mu_{M_2}) \\ &= i(\bar{\psi} \wedge \Gamma_a \Gamma_{10}\psi) \wedge e^a, \\ \mu_{D_2} &:= (\mu_{M_2})|_{8+1} \\ &= \frac{i}{2}(\bar{\psi} \wedge \Gamma_{a_1 a_2}\psi) \wedge e^{a_1} \wedge e^{a_2}, \\ \mu_{D_4} &:= (\pi_{10})_*(\mu_{M_5}) \\ &= +\frac{1}{4!}(\bar{\psi} \wedge \Gamma_{a_1 \dots a_4}\Gamma_{10}\psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_4}. \end{aligned}$$

Furthermore, consider the following elements in $CE(\mathbb{R}^{9,1|\mathbf{16}+\overline{\mathbf{16}}})$:

$$\begin{aligned} \mu_{D_6} &:= \frac{i}{6!}(\bar{\psi}\Gamma_{a_1 \dots a_6}\psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_6}, \\ \mu_{D_8} &:= \frac{1}{8!}(\bar{\psi}\Gamma_{a_1 \dots a_8}\Gamma_{10}\psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_8}, \\ \mu_{D_{10}} &:= \frac{i}{10!}(\bar{\psi}\Gamma_{a_1 \dots a_{10}}\psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_{10}}, \end{aligned}$$

where the indices run through $\{0, 1, \dots, 9\}$

Proposition 4.8. *The elements in Def. 4.7 satisfy the following differential conditions*

$$\begin{aligned} d\mu_{F_1}^A &= 0, \\ d\mu_{D_0} &= 0, \quad d\mu_{D_{10}} = \mu_{F_1}^A \wedge \mu_{D_8} = 0, \\ d\mu_{D_2} &= \mu_{F_1}^A \wedge \mu_{D_0}, \quad d\mu_{D_4} = \mu_{F_1}^A \wedge \mu_{D_2}, \\ d\mu_{D_6} &= \mu_{F_1}^A \wedge \mu_{D_4}, \quad d\mu_{D_8} = \mu_{F_1}^A \wedge \mu_{D_6}, \end{aligned}$$

hence they constitute an L_∞ -cocycle shown as the top morphism of the following diagram:

$$\begin{array}{ccc} & \{\mu_{F_1}^A, \mu_{D_0}, \mu_{D_2}, \mu_{D_4}, \mu_{D_6}, \mu_{D_8}, \mu_{D_{10}}\} & \\ & \curvearrowright & \\ \mathbb{R}^{9,1|\mathbf{16}+\overline{\mathbf{16}}} & \xrightarrow{\mathfrak{L}(\mu_{M_2}, \mu_{M_5})/b\mathbb{R}} \mathfrak{L}S^4/b\mathbb{R} & \xrightarrow{\quad} \mathfrak{l}(ku/BU(1)) \\ & & \downarrow \\ & & \mathfrak{l}(ku_{\leq 6}/BU(1)). \end{array}$$

Here for emphasis we also displayed the double dimensional reduction of the M-brane cocycle from example 4.5, and indicated that these coincide with the IIA cocycles on the F1, the D0, D2, and D4. Note that the M-brane cocycles also produce the NS5, but not the D6 and higher, which appear only in 10d (“gauge enhancement”).

Proof. The first equation follows for instance from $d\mu_{M2} = 0$ under dimensional reduction. The two equations in the second row follow trivially, by $d\psi^\alpha = 0$ and since there is no bosonic 11-form on $\mathbb{R}^{9,1}$. Regarding the equations in the third row: Using $d\psi^\alpha = 0$ and $de^a = \bar{\psi}\Gamma^a\psi$ (Def. 2.7) we find that they are equivalently rewritten as follows:

$$\begin{aligned} & d\mu_{D2} = \mu_{F1}^A \wedge \mu_{D0} \\ \Leftrightarrow & -i (\bar{\psi}\Gamma_{a_1 a} \psi) \wedge (\bar{\psi}\Gamma^a \psi) = +i (\bar{\psi}\Gamma_{a_1} \Gamma_{10} \psi) \wedge \bar{\psi}\Gamma_{10} \psi \\ & d\mu_{D4} = \mu_{F1}^A \wedge \mu_{D2} \\ \Leftrightarrow & -\frac{1}{3!} (\bar{\psi}\Gamma_{[a_1 a_2 a_3] a} \Gamma_{10} \psi) (\bar{\psi}\Gamma^a \psi) = -\frac{1}{2} (\bar{\psi}\Gamma_{[a_1 a_2} \psi) (\bar{\psi}\Gamma_{a_3]} \Gamma_{10} \psi) \\ & d\mu_{D6} = \mu_{F1}^A \wedge \mu_{D4} \\ \Leftrightarrow & -\frac{i}{5!} (\bar{\psi}\Gamma_{[a_1 \dots a_5] a} \psi) (\bar{\psi}\Gamma^a \psi) = \frac{i}{4!} (\bar{\psi}\Gamma_{[a_1 \dots a_4} \Gamma_{10} \psi) (\bar{\psi}\Gamma_{a_5]} \Gamma_{10} \psi) \\ & d\mu_{D8} = \mu_{F1}^A \wedge \mu_{D6} \\ \Leftrightarrow & -\frac{1}{7!} (\bar{\psi}\Gamma_{[a_1 \dots a_7] a} \Gamma_{10} \psi) (\bar{\psi}\Gamma^a \psi) = -\frac{1}{6!} (\bar{\psi}\Gamma_{[a_1 \dots a_6} \psi) (\bar{\psi}\Gamma_{a_7]} \Gamma_{10} \psi) . \end{aligned}$$

That these conditions hold may be checked to be equivalent to the statement of [17, expressions (6.8) with coefficients as found above (6.9)].

Alternatively, our main Theorem 5.3 below implies that the F1/Dp-cochains for type IIA are cocycles precisely if those for type IIB are, which we state below as def. 4.9. This implies that the above Fierz identities in type IIA hold precisely if those in type IIB hold. The latter has been checked independently in [61, section 2], see prop. 4.10 below. □

We now consider a similar construction for the type IIB theory.

Definition 4.9. Define the following elements in the Chevalley-Eilenberg algebra of the type IIB super-Minkowski spacetime $\mathbb{R}^{9,1|16+16}$ (Def. 2.13):

$$\begin{aligned}
 c_2^B &:= \bar{\psi}\Gamma_9^B\psi = \bar{\psi}\Gamma_B^9\psi, \\
 \mu_{F1}^B &:= i(\bar{\psi}\Gamma_a^B\Gamma_{10}\psi) \wedge e^a \\
 \mu_{D1} &:= i(\bar{\psi}\Gamma_a^B\Gamma_9\psi) \wedge e^a, \\
 \mu_{D3} &:= \frac{1}{3!}(\bar{\psi}\Gamma_{a_1\dots a_3}^B(\Gamma_9\Gamma_{10})\psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_3}, \\
 \mu_{D5} &:= \frac{i}{5!}(\bar{\psi}\Gamma_{a_1\dots a_5}^B\Gamma_9\psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_5}, \\
 \mu_{D7} &:= \frac{1}{7!}(\bar{\psi}\Gamma_{a_1\dots a_7}^B(\Gamma_9\Gamma_{10})\psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_7}, \\
 \mu_{D9} &:= \frac{i}{9!}(\bar{\psi}\Gamma_{a_1\dots a_9}^B\Gamma_9\psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_9}.
 \end{aligned}$$

Proposition 4.10. *The elements in Def. 4.9 constitute an L_∞ -cocycle in IIA super-Minkowski spacetime with coefficients in the model for twisted K^1 from Def. 4.6:*

$$\mathbb{R}^{9,1|\mathbf{16}+\mathbf{16}} \longrightarrow \mathfrak{l}(\Sigma\text{KU}/\text{BU}(1)).$$

Proof. By matching Clifford algebra conventions via Remark 2.10, one finds that this is the statement in [61, section 2]. But we may also re-derive this as a consequence of Theorem 5.3 below, which says that, under the dimensional reduction isomorphism from Theorem 3.8, the IIA cocycles of Prop. 4.8 are sent to the IIB elements from Def. 4.9 by the L_∞ -isomorphism of Prop. 5.1. Since L_∞ -homomorphisms preserve cocycles, the above claim follows via Theorem 3.8 and Theorem 5.3 from Proposition 4.8. \square

In summary, Proposition 4.8 and Proposition 4.10 say that the homotopical descent of the cocycles for the WZW-terms of the super p -branes from extended super-Minkowski spacetime down to the actual super-Minkowski spacetime is of the following form:

| String theory | Unified brane cocycles | Rational image of |
|---------------|--|-----------------------|
| type IIA | $\mathbb{R}^{9,1 \mathbf{16}+\overline{\mathbf{16}}} \xrightarrow{\mu_{F1/Dp}^A} \mathfrak{l}(\text{KU}/\text{BU}(1))$ | twisted K^0 -theory |
| type IIB | $\mathbb{R}^{9,1 \mathbf{16}+\mathbf{16}} \xrightarrow{\mu_{F1/Dp}^B} \mathfrak{l}(\Sigma\text{KU}/\text{BU}(1))$ | twisted K^1 -theory |

Notice that in this article, for ease of terminology we say “brane charge”, for the force field (“flux”) that the given brane feels, sourced by the (“magnetic”) background charge. For D-branes these are the *RR-field strengths*, while in the literature it is usual to say “D-brane charge” for the (“electric”) charge carried by the D-brane itself. That the classification of the RR-fields

is in K^0/K^1 for type II A/B was first argued in [52, p. 6] for the untwisted case. An explicit extension to the twisted case is indicated in [9] [50], which corresponds to the fields found above.

5. Super L_∞ -algebraic T-duality

The goal of this section is to describe how T-duality appears as an isomorphism between the F1/Dp-brane L_∞ -cocycles on the type IIA and type IIB supersymmetry super Lie algebra in 10d, after double dimensional reduction, in the sense of Theorem 3.8, down to 9d.

We have seen (Sec. 3) how the double dimensional reduction crucially involves cyclification. We will apply this for the corresponding spectra, i.e. rational twisted K-theory, to connect to the cocycles that we have just encountered in Sec. 4.

Proposition 5.1. *The cyclification $\mathfrak{L}(\mathrm{KU}/\mathrm{BU}(1))/\mathbb{R}$ (Definition 3.3) of $\mathfrak{l}(\mathrm{KU}/\mathrm{BU}(1))$ (Definition 4.6) has CE-algebra*

$$\mathrm{CE}(\mathfrak{L}(\mathrm{KU}/\mathrm{BU}(1))/\mathbb{R}) = \left\{ \begin{array}{l} dc_2 = 0, \quad d\tilde{c}_2 = 0 \\ dh_3 = -c_2 \wedge \tilde{c}_2 \\ d\omega_{2p+2} = h_3 \wedge \omega_{2p} + c_2 \wedge \omega_{2p+1} \\ d\omega_{2p+1} = h_3 \wedge \omega_{2p-1} + \tilde{c}_2 \wedge \omega_{2p} \end{array} \right\}.$$

The cyclification $\mathfrak{L}(\Sigma\mathrm{KU}/\mathrm{BU}(1))/\mathbb{R}$ of $\mathfrak{l}(\Sigma\mathrm{KU}/\mathrm{BU}(1))$ has CE-algebra

$$\mathrm{CE}(\mathfrak{L}(\Sigma\mathrm{KU}/\mathrm{BU}(1))/\mathbb{R}) = \left\{ \begin{array}{l} dc_2 = 0, \quad d\tilde{c}_2 = 0 \\ dh_3 = -c_2 \wedge \tilde{c}_2 \\ d\omega_{2p+2} = h_3 \wedge \omega_{2p} + \tilde{c}_2 \wedge \omega_{2p+1} \\ d\omega_{2p+1} = h_3 \wedge \omega_{2p-1} + c_2 \wedge \omega_{2p} \end{array} \right\}.$$

Hence there is an L_∞ -isomorphism of the form

$$\phi_T : \mathfrak{l}(\mathcal{L}(\mathrm{KU}/\mathrm{BU}(1))/S^1) \xrightarrow[\simeq]{\phi_T} \mathfrak{l}(\mathcal{L}(\Sigma\mathrm{KU}/\mathrm{BU}(1))/S^1)$$

relating the cyclifications of the rational twisted KU-coefficients, which is given by

$$c_2 \longleftrightarrow \tilde{c}_2, \quad h_3 \longmapsto h_3, \quad \omega_p \longmapsto \omega_p.$$

Proof. By Definition 3.3, as a polynomial algebra the CE-algebra of $\mathfrak{L}(\mathrm{KU}/\mathrm{BU}(1))$ is obtained from the CE-algebra of $\mathfrak{l}(\mathrm{KU}/\mathrm{BU}(1))$ by adding

a shifted copy of each generator. We denote by ω_{2p-1} the shifted copy of ω_{2p} and by $-\tilde{c}_2$ the shifted copy of h_3 . The differential is then defined by

$$d\omega_{2p+2} = h_3 \wedge \omega_{2p}, \quad d\omega_{2p+1} = h_3 \wedge \omega_{2p-1} + \tilde{c}_2 \wedge \omega_{2p}, \quad dh_3 = 0, \quad d\tilde{c}_2 = 0.$$

Next, again by Definition 3.3, the CE-algebra of $\mathfrak{L}(\mathrm{KU}/\mathrm{BU}(1))/\mathbb{R}$ is obtained by adding a further degree 2 generator c_2 and defining the differential as

$$\begin{aligned} d\omega_{2p+2} &= h_3 \wedge \omega_{2p} + c_2 \wedge \omega_{2p+1}, & d\omega_{2p+1} &= h_3 \wedge \omega_{2p-1} + \tilde{c}_2 \wedge \omega_{2p}, \\ dc_2 &= 0, & d\tilde{c}_2 &= 0, & dh_3 &= -c_2 \wedge \tilde{c}_2. \end{aligned}$$

The proof for $\mathfrak{L}(\Sigma\mathrm{KU}/\mathrm{BU}(1))/\mathbb{R}$ is completely analogous. □

Example 5.2. By Prop. 4.8, the combined F1/D p -brane cocycles on type IIA super-Minkowski spacetime constitute an L_∞ -homomorphism (corresponding to rational twisted K^0) of the form

$$\mu_{F1/Dp}^{\mathrm{IIA}} : \mathbb{R}^{9,1|\mathbf{16}+\overline{\mathbf{16}}} \xrightarrow{\{\mu_{F1}^{\mathrm{IIA}}, \{\mu_{D2p}\}\}} \mathfrak{l}(\mathrm{KU}/\mathrm{BU}(1))$$

with the coefficient L_∞ -algebra on the right from Def. 4.6. By Prop. 2.14 and Theorem 3.8 this is naturally identified with a dimensionally reduced cocycle of the form

$$\begin{array}{ccc} \mathfrak{L}(\mu_{F1/Dp}^{\mathrm{IIA}})/\mathbb{R} : \mathbb{R}^{8,1|\mathbf{16}+\mathbf{16}} & \xrightarrow{\left\{ \begin{array}{l} \mu_{F1}^{\mathrm{IIA}}, \{\mu_{D2p}\} \\ (\pi_9^{\mathrm{IIA}})_* \mu_{F1}^{\mathrm{IIA}}, \{(\pi_9^{\mathrm{IIA}})_* \mu_{D2p}\} \\ c_2^{\mathrm{IIA}} \end{array} \right\}} & \mathfrak{L}(\mathfrak{l}(\mathrm{KU}/\mathrm{BU}(1)))/\mathbb{R} \\ & \searrow c_2^{\mathrm{IIA}} & \swarrow \omega_2 \\ & & b\mathbb{R} \end{array}$$

with coefficients now given by Prop. 5.1, where in braces on top we see the original cocycles without the pieces containing e^9 , below that we see the e^9 -components, and in the last line the cocycle that classifies the IIA extension. Similarly, the combined F1/D p -brane cocycles on type IIB super-Minkowski spacetime constitute an L_∞ -homomorphism (corresponding to rational twisted K^1) of the form

$$\mu_{F1/Dp}^{\mathrm{IIB}} : \mathbb{R}^{9,1|\mathbf{16}+\mathbf{16}} \xrightarrow{\{\mu_{F1}^{\mathrm{IIB}}, \{\mu_{D(2p+1)}\}\}} \mathfrak{l}(\mathrm{KU}/\mathrm{BU}(1))$$

and by Prop. 2.14 and Theorem 3.8 this is naturally identified with a dimensionally reduced cocycle of the form

$$\mathfrak{L}(\mu_{F_1/Dp}^{\text{IIB}})/\mathbb{R} : \mathbb{R}^{8,1|\mathbf{16}+\mathbf{16}} \xrightarrow{\left\{ \begin{array}{l} \mu_{F_1}^{\text{IIB}}, \{\mu_{D(2p+1)}\} \\ (\pi_9^{\text{IIB}})_* \mu_{F_1}^{\text{IIB}}, \{(\pi_9^{\text{IIB}})_* \mu_{D(2p+1)}\} \end{array} \right\}} \mathfrak{L}(\mathfrak{I}(\Sigma\text{KU}/\text{BU}(1)))/\mathbb{R} .$$

$$\begin{array}{ccc} & & \\ & \searrow^{c_2^{\text{IIB}}} & \swarrow_{\omega_2} \\ & & b\mathbb{R} \end{array}$$

Now for the original F_1/Dp cocycles $\mu_{F_1/Dp}^{\text{IIA}}$ and $\mu_{F_1/Dp}^{\text{IIB}}$ from [29, Sec. 4] it does not make sense to ask whether they are equivalent, since their domains are not. However, their double dimensional reductions in Example 5.2, to which by Theorem 3.8 they bijectively correspond, do have the same domain $\mathbb{R}^{8,1|\mathbf{16}+\mathbf{16}}$. This is ultimately due to the fact that the two inequivalent chiral $\text{Spin}(9, 1)$ -representations become isomorphic as $\text{Spin}(8, 1)$ -representations — see Remark 2.9. Hence for these it makes sense to ask whether they are equivalent L_∞ -homomorphisms. We now establish that indeed they are:

Theorem 5.3. *The L_∞ -algebra isomorphism of Proposition 5.1 takes the dimensionally reduced type IIA F_1/Dp -cocycle $\mathfrak{L}(\mu_{F_1/Dp}^{\text{IIA}})/\mathbb{R}$ of Example 5.2 to the dimensionally reduced IIB cocycle $\mathfrak{L}(\mu_{F_1/Dp}^{\text{IIB}})/\mathbb{R}$, making the following diagram of super L_∞ -algebras commute:*

$$\begin{array}{ccccc} & & b\mathbb{R} & \xleftarrow{\omega_2} & \mathfrak{L}(\Sigma\text{KU}/\text{BU}(1))/\mathbb{R} \\ & \nearrow^{c_2^{\text{IIB}}} & & & \uparrow \simeq \phi_T \\ \mathbb{R}^{8,1|\mathbf{16}+\mathbf{16}} & \xrightarrow{\mathfrak{L}(\mu_{F_1/Dp}^{\text{IIB}})/\mathbb{R}} & & & \mathfrak{L}(\text{KU}/\text{BU}(1))/\mathbb{R} \\ & \searrow_{\mathfrak{L}(\mu_{F_1/Dp}^{\text{IIA}})/\mathbb{R}} & & & \downarrow \\ & & b\mathbb{R} & \xleftarrow{\omega_2} & \end{array}$$

Proof. First of all we need to check that

$$\omega_2(\phi_T(\mathfrak{L}(\mu_{F_1/Dp}^{\text{IIA}})/\mathbb{R})) = c_2^{\text{IIB}}.$$

By Theorem 3.8 and Prop. 5.1 the left hand side here is the integration over the fiber $-(\pi_9^{\text{IIA}})_*(\mu_{F_1}^{\text{IIA}})$. Hence we need to check that

$$(4) \quad -(\pi_9^{\text{IIA}})_*(\mu_{F_1}^{\text{IIA}}) = c_2^{\text{IIB}}.$$

In components this equality says that $i\bar{\psi}\Gamma_9\Gamma_{10}\psi = \bar{\psi}\Gamma_9^B\psi$. This holds by direct inspection — see Remark 2.12. Similarly, one has

$$(5) \quad -(\pi_9^{\text{IIB}})_*(\mu_{F_1}^{\text{IIB}}) = c_2^{\text{IIA}},$$

due to the identity $i\bar{\psi}\Gamma_9^B\Gamma_{10}\psi = \bar{\psi}\Gamma_9\psi$. Moreover, we also have

$$\mu_{F_1}^{\text{IIA}}|_{8+1} = \mu_{F_1}^{\text{IIB}}|_{8+1}$$

due to the identity $\Gamma_a^B = \Gamma_a$ for $0 \leq a \leq 8$. In view of this, to see that $\mu_{F_1}^{\text{IIA}}$ from Def. 4.7 is sent to $\mu_{F_1}^{\text{IIB}}$ from Def. 4.9, use that ϕ_T swaps c_2^{IIA} with c_2^{IIB} , while keeping the restriction $\mu_{F_1}|_{8+1}$ intact. Now we need to check that the D-brane charges are sent to each other. Unwinding the definitions, this means that interchanging the components of fiber integration $(\pi_9^A)_*(-)$ and restriction $(-)|_{8+1}$ then the process in expression (2) turns the IIA-brane elements from Def. 4.7 to the IIB-brane elements from Def. 4.9. This is indeed the case, as the following explicit computations show (where under the braces we keep using Remark 2.12):

$$\begin{aligned} (\boxed{\text{D1}}) \quad & (\pi_9^A)_*(\mu_{D_2}) - e^9 \wedge (\mu_{D_0}|_{8+1}) \\ &= i \sum_{a=0}^8 \bar{\psi}\Gamma_a\Gamma_9\psi \wedge e^a - \bar{\psi} \underbrace{\Gamma_{10}}_{=-i\Gamma_9^B\Gamma_9} \psi \wedge e^9 \\ &= i(\bar{\psi}\Gamma_a^B\Gamma_9\psi) \wedge e^a \\ &= \mu_{D_1}. \end{aligned}$$

$$\begin{aligned} (\boxed{\text{D3}}) \quad & (\pi_9^A)_*(\mu_{D_4}) - e^9 \wedge (\mu_{D_2}|_{8+1}) \\ &= \frac{1}{3!} \sum_{a_i=0}^8 \bar{\psi}\Gamma_{a_1 a_2 a_3} \Gamma_9 \Gamma_{10} \psi \wedge e^{a_1} \wedge e^{a_2} \wedge e^{a_3} \\ &\quad - \frac{i}{2} \sum_{a_i=0}^8 \bar{\psi}\Gamma_{a_1 a_2} \psi \wedge e^{a_1} \wedge e^{a_2} \wedge e^9 \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{3!} \sum_{a_i=0}^8 \bar{\psi} \Gamma_{a_1 a_2 a_3} \Gamma_9 \Gamma_{10} \psi \wedge e^{a_1} \wedge e^{a_2} \wedge e^{a_3} \\
&\quad + \frac{1}{3!} 3 \sum_{a_i=0}^8 \bar{\psi} \Gamma_{a_1 a_2} \underbrace{\Gamma_9^B(\Gamma_9 \Gamma_{10})}_{=-i} \psi \wedge e^{a_1} \wedge e^{a_2} \wedge e^9 \\
&= \frac{1}{3!} (\bar{\psi} \Gamma_{a_1 a_2 a_3}^B(\Gamma_9 \Gamma_{10}) \psi) \wedge e^{a_1} \wedge e^{a_2} \wedge e^{a_3} \\
&= \mu_{D_3}.
\end{aligned}$$


$$\begin{aligned}
\boxed{\text{D5}} \quad &(\pi_9)_*(\mu_{D_6}) - e^9 \wedge (\mu_{D_4}|_{8+1}) \\
&= \frac{i}{5!} \sum_{a_i=0}^8 (\bar{\psi} \Gamma_{a_1 \dots a_5} \Gamma_9 \psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_5} \\
&\quad - \frac{1}{4!} \sum_{a_i=0}^8 (\bar{\psi} \Gamma_{a_1 \dots a_4} \Gamma_{10} \psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_4} \wedge e^9 \\
&= \frac{i}{5!} \sum_{a_i=0}^8 (\bar{\psi} \Gamma_{a_1 \dots a_5} \Gamma_9 \psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_5} \\
&\quad - \frac{1}{5!} 5 \sum_{a_i=0}^8 \left(\bar{\psi} \Gamma_{a_1 \dots a_4} \underbrace{\Gamma_{10}}_{=-i \Gamma_9^B \Gamma_9} \psi \right) \wedge e^{a_1} \wedge \dots \wedge e^{a_4} \wedge e^9 \\
&= \frac{i}{5!} (\bar{\psi} \Gamma_{a_1 \dots a_5}^B \Gamma_9 \psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_5} \\
&= \mu_{D_5}.
\end{aligned}$$

$$\begin{aligned}
\boxed{\text{D7}} \quad &(\pi_9)_*(\mu_{D_8}) - e^9 \wedge (\mu_{D_6}|_{8+1}) \\
&= \frac{1}{7!} \sum_{a_i=0}^8 (\bar{\psi} \Gamma_{a_1 \dots a_7} \Gamma_9 \Gamma_{10} \psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_7} \\
&\quad - \frac{i}{6!} \sum_{a_i=0}^8 (\bar{\psi} \Gamma_{a_1 \dots a_6} \psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_6} \wedge e^9 \\
&= \frac{1}{7!} \sum_{a_i=0}^8 (\bar{\psi} \Gamma_{a_1 \dots a_7} \Gamma_9 \Gamma_{10} \psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_7} \\
&\quad + \frac{1}{7!} 7 \sum_{a_i=0}^8 \left(\bar{\psi} \Gamma_{a_1 \dots a_6} \underbrace{\Gamma_9^B(\Gamma_9 \Gamma_{10})}_{=-i} \psi \right) \wedge e^{a_1} \wedge \dots \wedge e^{a_6} \wedge e^9 \\
&= \frac{1}{7!} (\bar{\psi} \Gamma_{a_1 \dots a_7}(\Gamma_9 \Gamma_{10}) \psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_7} \\
&= \mu_{D_7}.
\end{aligned}$$


$$\begin{aligned}
 \boxed{\text{D9}} \quad & (\pi_9)_*(\mu_{D10}) - e^9 \wedge (\mu_{D8}|_{8+1}) \\
 &= \frac{i}{9!} \sum_{a_i=0}^8 (\bar{\psi} \Gamma_{a_1 \dots a_9} \Gamma_9 \psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_9} \\
 &\quad - \frac{1}{8!} \left(\bar{\psi} \Gamma_{a_1 \dots a_8} \underbrace{\Gamma_{10}}_{=-i\Gamma_9^B \Gamma_9} \psi \right) \wedge e^{a_1} \wedge \dots \wedge e^{a_8} \wedge e^9 \\
 &= \frac{i}{9!} \sum_{a_i=0}^8 (\bar{\psi} \Gamma_{a_1 \dots a_9} \Gamma_9 \psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_9} \\
 &\quad + \frac{i}{9!} 9 \sum_{a_i=0}^8 (\bar{\psi} \Gamma_{a_1 \dots a_8} \Gamma_9^B \Gamma_9 \psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_8} \wedge e^9 \\
 &= \frac{i}{9!} (\bar{\psi} \Gamma_{a_1 \dots a_9} \Gamma_9 \psi) \wedge e^{a_1} \wedge \dots \wedge e^{a_9} \\
 &= \mu_{D9}.
 \end{aligned}$$

□

Remark 5.4 (Topological T-duality I). (i) The interpretation of our super-Minkowski spacetimes as tangent spaces of spacetime manifolds X , and extensions of them as corresponding fiber bundles $X_{10} \rightarrow X_9$ of spacetime manifolds, means that the diagram of super L_∞ algebras from Proposition 2.14



together with 3-cocycles $\mu_{F1}^{\text{IIA/B}}$ globalizes to a diagram of manifolds of the form



which carry closed differential super 3-forms $H_3^{A/B} \in \Omega_{\text{cl}}^3(X_{10}^{A/B})$.

Indeed, the consistency of the Green-Schwarz sigma-model for the type II superstring on $X_{10}^{A/B}$ requires that the bispinorial component of the super-3-form H here is constrained to coincide on each tangent spacetime with our cocycle $\mu_{F_1}^{IIA/B}$ (this follows from [5, equation (2.11)]). In particular, if H happens to have vanishing bosonic component, then it is entirely fixed by restricting on each tangent space to our cocycle $\mu_{F_1}^{IIA/B}$ (this is amplified in [5, equation (2.15)]). Similarly, these circle bundles have first Chern classes whose representing forms $C_2^{A/B}$ need to have bispinorial super-components that super-tangent-space wise coincide with the cocycles $c_2^{IIA/B}$.

(ii) Hence the globalization of the super-tangent-space wise equivalence that we see in equations (4-5) in the proof of Theorem 5.3 imposes the global condition

$$C_2^{IIA/B} = -(\pi_9^{IIB/A})_*(H_3^{IIB/A}).$$

This relation is precisely what is used as an axiom for “topological T-duality” in [9, (1.8)], [12, lemma 2.12, lemma 2.33].

6. T-Correspondence space and Doubled spacetimes

Above we considered T-duality as an equivalence of classifying maps (moduli) of fields. Here we consider the incarnation of this equivalence in terms of the higher extended super-Minkowski spaces that are classified thereby.

With every L_∞ -cocycle on some superalgebra \mathfrak{g} , we get some L_∞ -extension that it classifies, namely its homotopy fiber $\widehat{\mathfrak{g}} \rightarrow \mathfrak{g}$, according to Example 2.5. If \mathfrak{g} is some super-Minkowski super Lie algebra (Sec. 2), then this extension is the higher extended super-Minkowski spacetime which may be thought of as containing condensates of those brane species that the cocycle classifies [27, Remark 3.11]. In particular a 2-cocycle corresponds to a 0-brane and the corresponding extension is just an ordinary central extension, hence grows one extra dimension of spacetime, as befits a 0-brane condensate [27, Remark 4.6].

Here we analyze this phenomenon for the cocycles that classify the type II branes on $9d$ $N = 2$ super-Minkowski spacetime, according to Example 5.2. By Theorem 5.3 there is only one such cocycle, up to equivalence, since the dimensional reduction $\mathfrak{L}(\mu_{F_1/Dp}^{IIA})/\mathbb{R}$ of the IIA branes and the dimensional reduction $\mathfrak{L}(\mu_{F_1/Dp}^{IIB})/\mathbb{R}$ of the IIB branes agrees in 9d, by L_∞ -algebraic T-duality. Hence we just write $\mathfrak{L}(\mu_{F_1/Dp}^{II})/\mathbb{R}$ for either of them. Now, as made explicit in the diagram in Theorem 5.3, this cocycle contains contributions from *two* 0-brane species. One of these is the D0-brane

of type IIA in 10d, descended down to 9d, and the other is the double dimensional reduction of the type IIB string 3-cocycle from 10d to 9d. Hence the condensation of these [27, Remark 3.11, 4.6] grows *two* extra spacetime dimensions. By Theorem 5.3, these are the infinitesimal version of what in finite T-duality are the two circle fibers S_A^1 and S_B^1 of Remark 5.4. Hence in the notation of that remark, we obtain the fiber product spacetime

$$\begin{array}{ccccc}
 & & S_A^1 \times S_B^1 \hookrightarrow & X_{10}^{\text{IIA}} \times_{X_9} & X_{10}^{\text{IIB}} \\
 & & \swarrow & & \searrow \\
 S_A^1 \longrightarrow & X_{10}^{\text{IIA}} & & (pb) & X_{10}^{\text{IIB}} \longleftarrow S_B^1 \\
 & \searrow^{\pi_9^{\text{IIA}}} & & & \swarrow_{\pi_9^{\text{IIB}}} \\
 & & X_9 & &
 \end{array}$$

which is an $S_A^1 \times S_B^1$ -fiber bundle over X_9 .


Definition 6.1. Write

$$\mathbb{R}^{8+(1+1),1|32} := \mathbb{R}^{9,1|\mathbf{16}+\overline{\mathbf{16}}} \times_{\mathbb{R}^{8,1|\mathbf{16}+\mathbf{16}}} \mathbb{R}^{9,1|\mathbf{16}+\mathbf{16}}$$

as shorthand for the fiber product of the type IIA super-Minkowski spacetime with its IIB version, over their common 9d base, according to Prop. 2.14, hence for the super Lie algebra fitting into the following fiber product diagram

$$\begin{array}{ccccc}
 & & \mathbb{R}^{8+(1+1),1|32} & & \\
 & & \swarrow^{p_B} & & \searrow^{p_A} \\
 \mathbb{R}^{9,1|\mathbf{16}+\mathbf{16}} & & & (pb) & \mathbb{R}^{9,1|\mathbf{16}+\overline{\mathbf{16}}} \\
 & \searrow & & & \swarrow \\
 & & \mathbb{R}^{8,1|\mathbf{16}+\mathbf{16}} & &
 \end{array}$$

Proposition 6.2. *We have a diagram of super L_∞ -algebras of the following form*



where

- $\widehat{\mathbb{R}^{9,1|\mathbf{16}+\overline{\mathbf{16}}}} \simeq \mathbf{string}_{IIA}$ is the super Lie 2-algebra extension of type IIA super-Minkowski spacetime by the 3-cocycle for the type IIA super-string; i.e. the infinitesimal model of the (super-)gerbe underlying the type IIA B-field;
- $\widehat{\mathbb{R}^{9,1|\mathbf{16}+\mathbf{16}}} \simeq \mathbf{string}_{IIB}$ is the super Lie 2-algebra extension of type IIB super-Minkowski spacetime by the 3-cocycle for the type IIB super-string; i.e. the infinitesimal model of the (super-)gerbe underlying the type IIB B-field;
- by slight abuse of notation, $\pi_9^{A/B}$ always denotes the map that projects out the 9th dimension of type A/B, respectively, hence dually, on underlying graded algebras, the canonical map $(-) \rightarrow (-)[e_9^{A/B}]$ induced from adjoining the vielbein generator $e_9^{A/B}$

such that

- 1) a horizontal isomorphism ν exists as shown, between the pullback of these two extensions to the correspondence space, making the diagram commute;
- 2) on the classifying 3-cocycles it is given by the Poincaré form

$$\mathcal{P} := e_9^A \wedge e_9^B$$

as

$$p_B^*(\mu_{F1}^B) - p_A^*(\mu_{F1}^A) = d\mathcal{P}.$$

Proof. By Example 2.5, the extended super-Minkowski super Lie 2-algebra on the far left is given by

$$\text{CE}(\widehat{\mathbb{R}^{9,1|\mathbf{16}+\overline{\mathbf{16}}}}) = \text{CE}(\mathbb{R}^{9,1|\mathbf{16}+\overline{\mathbf{16}}})[f_2^A]/(df_2^A = \mu_{F_1}^A)$$

and that on the far right by

$$\text{CE}(\widehat{\mathbb{R}^{9,1|\mathbf{16}+\mathbf{16}}}) = \text{CE}(\mathbb{R}^{9,1|\mathbf{16}+\mathbf{16}})[f_2^B]/(df_2^B = \mu_{F_1}^B).$$

Their pullbacks along the projections p_A and p_B are directly seen to be given by further adjoining the generator e_9^B or e_9^A , respectively.

Then define ν by defining its dual ν^* by sending all generators in $p_A^* \text{CE}(\widehat{\mathbb{R}^{9,1|\mathbf{16}+\overline{\mathbf{16}}}})$ to the generator of the same name in $p_B^* \text{CE}(\widehat{\mathbb{R}^{9,1|\mathbf{16}+\mathbf{16}}})$ except for f_2^A , for which we set

$$(6) \quad \nu^* : f_2^A \mapsto f_2^B - \mathcal{P}.$$

To see that this indeed gives a homomorphism, recall equation (4) which followed explicitly in components via Remark 2.12

$$i \bar{\psi} \Gamma_9^{A/B} \Gamma_{10} \psi = \bar{\psi} \Gamma_9^{B/A} \psi.$$

Here this means that on the doubled correspondence space the type IIA and IIB superstring cocycles $\mu_{F_1}^{A/B}$ (Definition 4.7 and Definition 4.9) may be expressed in terms of each other as follows (as above in (??))

$$(7) \quad \mu_{F_1}^{A/B} = \mu_{F_1}^9 + e_9^{A/B} \wedge (\bar{\psi} \wedge \Gamma_9^{B/A} \psi),$$

where $\mu_{F_1}^9$ denotes their common summand, the one which involves only the generators e_a for $a \leq 8$. This implies first of all the claimed coboundary between the 3-cocycles:

$$\begin{aligned} \mu_{F_1}^A + d\mathcal{P} &= \underbrace{\mu_{F_1}^9 + e_9^A \wedge \bar{\psi} \Gamma_9^B \psi}_{\mu_{F_1}^A} + \underbrace{e_9^B \wedge \bar{\psi} \Gamma_9^A \psi - e_9^A \wedge \bar{\psi} \Gamma_9^B \psi}_{d(e_9^A \wedge e_9^B)} \\ &= \mu_{F_1}^9 + e_9^B \wedge \bar{\psi} \Gamma_9^A \psi \\ &= \mu_{F_1}^B. \end{aligned}$$

In turn, this implies now that ν^* is indeed a dg-algebra homomorphism, because in the extended type IIA dg-algebra $p_B^* \text{CE}(\widehat{\mathbb{R}^{9,1|\mathbf{16}+\mathbf{16}}})$ we have the


following relations:

$$\begin{aligned}
 d(\nu^*(f_2^A)) &= d(f_2^B - \mathcal{P}) \\
 &= \mu_{F1}^B - d\mathcal{P} \\
 &= \mu_{F1}^A \\
 &= \nu^*(d(f_2^A)).
 \end{aligned}$$

Finally, ν clearly has an inverse ν^{-1} . Its linear dual is the identity on all generators except f_2^B , where it is

$$(\nu^{-1})^* : f_2^B \mapsto f_2^A + \mathcal{P}. \quad \square$$

Remark 6.3 (Topological T-duality II). Proposition 6.2 is evidently the infinitesimal and supergeometric picture of topological T-duality as considered in [13, def. 2.8]. There, one considers two circle bundles X_{10}^{IIA} and X_{10}^{IIB} , carrying a $U(1)$ -gerbe \mathcal{G}_{IIA} and \mathcal{G}_{IIB} , respectively, such that there is an equivalence between these gerbes after pullback to the fiber product:



and such that this equivalence is fiberwise exhibited by the 2-cocycle which is the cup-product of the canonical 1-classes on the two circle fibers.

Therefore, we find both perspectives on topological T-duality (Remark 5.4 and Remark 6.3), from the analysis of the super-tangent-space wise super L_∞ -cocycles (Prop. 5.3 and Prop. 6.2, respectively). (Previously these rules had been guessed, not derived from string theory.) Of course these two perspectives are supposed to be equivalent, rationally. In section 7 we see how this equivalence arises within the homotopy theory of super L_∞ -algebras.

Before we turn to that, we observe that the correspondence super-space-time $\mathbb{R}^{9+(1,1),1}$ ³² also serves to clarify the operation of T-duality on the

RR-fields: Consider the standard fact (see also [29, section 4]) that a degree-3 twisted KU-valued cocycle on any super L_∞ -algebra \mathfrak{g}

$$\left\{ \begin{array}{ccc} \mathfrak{g} & \xrightarrow{\{\omega_{2p}|p \in \mathbb{N}\}} & \mathfrak{l}(\text{KU}/\text{BU}(1)) \\ \mu_3 \searrow & & \nearrow \\ & b^2\mathbb{R} & \end{array} \right\} \leftrightarrow \{\omega_{2p} \in \text{CE}(\mathfrak{g}) \mid d_{\mathfrak{g}}\omega_{2p} = \mu_3 \wedge \omega_{2p-2}\}$$

is naturally identified with an untwisted ku-cocycle on the higher central extension $\widehat{\mathfrak{g}}$. The latter is classified by the twist μ_3 according to Example 2.5

$$\left\{ \begin{array}{ccc} \widehat{\mathfrak{g}} & \xrightarrow{\{\tilde{\omega}_{2p}|p \in \mathbb{N}\}} & \mathfrak{l}(\text{KU}) \\ \text{hofib}(\mu_3) \downarrow & & \\ \mathfrak{g} & & \end{array} \right\} \leftrightarrow \{\tilde{\omega}_{2p} \in \text{CE}(\widehat{\mathfrak{g}}) \mid d\tilde{\omega}_{2p} = 0\}$$

under the relation

$$\tilde{\omega}_{2p} = [\exp(-f_2) \wedge C]_{2p}, \quad C := \sum_{p \in \mathbb{N}} \omega_{2p},$$

where f_2 is the new generator of $\text{CE}(\widehat{\mathfrak{g}}) = \text{CE}(\mathfrak{g})[f_2, df_2 = \mu_3]$ according to Example 2.5, and where $[-]_{2p}$ denotes taking the summand of homogeneous degree $2p$. Under this identification, the top three morphisms in the diagram of Proposition 6.2 define a linear map

$$(\pi_{\mathfrak{g}}^A)_* \circ \nu^* \circ (\pi_{\mathfrak{g}}^B)^*$$

from the twisted cocycle of the type IIA super-Minkowski spacetime to that of type IIB. Here the two pullback operations are just the dual morphisms on CE-algebras, while the pushforward operation $(\pi_{\mathfrak{g}}^A)_*$ is defined as in expressions (1) and (2).

Proposition 6.4. *This operation is a well defined morphism on twisted cohomology groups*

$$(\pi_{\mathfrak{g}}^A)_* \circ \nu^* \circ (\pi_{\mathfrak{g}}^B)^* : H_{\mu_{F_1}^A}(\mathbb{R}^{9,1|\mathbf{16}+\overline{\mathbf{16}}}, \mathfrak{l}(\text{KU})) \longrightarrow H_{\mu_{F_1}^B}(\mathbb{R}^{9,1|\mathbf{16}+\mathbf{16}}, \mathfrak{l}(\Sigma\text{KU}))$$

and is in fact an isomorphism. Moreover, it identifies the type IIA D-brane cocycles with those of type IIB, as in Theorem 5.3:

$$(8) \quad \exp(-f_2^B) \wedge C^B = (\pi_{\mathfrak{g}}^A)_* \circ \nu^* \circ (\pi_{\mathfrak{g}}^B)^* (\exp(-f_2^A) \wedge C^A),$$

where

$$C^A := \mu_{D_0} + \mu_{D_2} + \cdots + \mu_{D_{10}}, \quad C^B := \mu_{D_1} + \mu_{D_3} + \cdots + \mu_{D_9}.$$

Proof. First, the pullback operation $(\pi_9^B)^*$ leaves the form of the cocycles unaffected, and simply regards them inside the larger CE-algebra which has the generator e_9^B adjoined. Next, by Proposition 6.2, equation (6), the pullback along ν amounts to substituting f_2^A by $f_2^B - \mathcal{P} = f_2^B - e_9^A \wedge e_9^B$. This means that

$$(9) \quad \nu^* : \exp(-f_2^A)C^A \mapsto \exp(-f_2^B + \mathcal{P}) = \exp(-f_2^B) \exp(\mathcal{P})C^A.$$

Since $\mathcal{P} = e_9^A \wedge e_9^B$ squares to zero, this in turn amounts to sending

$$C^A \mapsto C^A + e_9^A \wedge e_9^B \wedge C^A.$$

Observe then that fiber integration $(\pi_9^A)_*$ applied to wedge products with $e_9^A \wedge e_9^B$ amounts to wedge product with e_9^B of the part that does not contain a factor e_9^A :

$$(\pi_9^A)_* (e_9^B \wedge e_9^A \wedge (\cdots)) = -e_9^B \wedge (\cdots)|_{8+1}$$

(since e_9^A squares to zero and using equations (1) and (2)). Summing up, it follows that acting on elements of the form $\exp(f_2^A)C^A$ we have

$$(10) \quad (\pi_9^A)_* \circ \nu^* \circ (\pi_9^B)^* = (\pi_9^A)_*(-) - e_9^B \wedge (-)|_{8+1}.$$

Comparison of the right hand side with the proof of Theorem 5.3 (equations [D1](#) through [D9](#)) shows that this is precisely what establishes the T-duality isomorphism between twisted cocycles in general, and which identifies the type IIA D-brane cocycles with those of type IIB in particular. □

Remark 6.5 (Topological T-duality III). In view of Remark 6.3, the isomorphism in Proposition 6.4 is seen to be the super L_∞ -algebraic analog of the key result of classical topological T-duality [12, Theorem 3.13]. Moreover, via equation (9) the identification from equation (8) is

$$\exp(-f_2^B)C^B = (\pi_9^A)_* (\exp(\mathcal{P}) \wedge (\pi_9^B)^* (\exp(-f_2^A)C^A)).$$

This is precisely the form usually known as the *Buscher rules for RR-fields*, or the *Hori formula* [36, equation (1.1)]. Here we obtained this from just the form of the super-cocycles, to which the bifermionic summand of more general RR-fields are constrained to be equal at the level of super-tangent spaces.

7. T-Duality Lie 2-algebra

It was proposed in [38] [39] that there ought to be a concept of “T-folds” which generalizes that of manifolds to a situation where diffeomorphisms may be accompanied by T-duality transformations. In [53] it is shown that the correct mathematical formulation of this concept is by spaces associated to principal 2-bundles ([54]) for structure 2-group a certain “T-duality 2-group”. We show now that the extended supergeometry implied by the cocycle $\mathfrak{L}(\mu_{F1/Dp}^{\text{II}})/\mathbb{R}$ from Example 5.2 provides a systematic derivation of this structure, infinitesimally, but including the supergeometric aspects.

First of all, we consider the following sub- L_∞ -algebra of $\mathfrak{L}(\text{KU}/\text{BU}(1))/\mathbb{R}$:

Definition 7.1. The *delooped T-duality Lie 2-algebra* is given by

$$\text{CE}(b\mathcal{T}_1) = \left\{ \mathbb{R}[c_2, \tilde{c}_2, h_3]; \quad \begin{array}{l} dc_2 = 0, \quad d\tilde{c}_2 = 0 \\ dh_3 = -c_2 \wedge \tilde{c}_2 \end{array} \right\}.$$

Remark 7.2 (T-duality Lie 2-group). The delooped T-duality Lie 2-algebra $b\mathcal{T}(1)$ from Definition 7.1 is the L_∞ -algebra corresponding to a smooth 2-group $T(1)$, the *T-duality 2-group*, with smooth delooping 2-stack $\mathbf{BT}(1)$. The construction of the latter follows from [26, section 3.2]: Consider the homotopy fiber

$$\begin{array}{ccc} \mathbf{BT}(1) & \longrightarrow & * \\ \text{hofib}(\mathbf{c}_1 \cup \mathbf{c}'_1) \downarrow & & \downarrow \\ \mathbf{BU}(1) \times \mathbf{BU}(1) & \xrightarrow{\mathbf{c}_1 \cup \mathbf{c}'_1} & \mathbf{B}^3U(1), \end{array}$$

where $\mathbf{BU}(1)$ is the smooth stack of principal $U(1)$ -bundles, $\mathbf{B}^3U(1)$ is the smooth stack of $\mathbf{B}^2U(1)$ -principal 2-bundles (bundle 2-gerbes) and $\mathbf{c}_1 \cup \mathbf{c}'_1$ is external cup product of the universal first Chern class with itself, regarded as a morphism of smooth stacks. Here $\mathbf{c}_1 \cup \mathbf{c}'_1$ is a homomorphism of smooth higher group stacks, and so its homotopy fiber inherits group structure, too. This means that a $T(1)$ -principal bundle over a smooth (super-)manifold X is the datum of two principal $U(1)$ -bundles on X together with a trivialization of the product of their first Chern classes. One manifestly sees that, translated in terms of Chevalley-Eilenberg algebras, this is precisely the content of Definition 7.1.

Finally, forgetting the smooth structure and passing to topological geometric realization $|-|$ (as in [65]), then the above homotopy fiber of smooth

higher stacks becomes the ordinary homotopy fiber of topological spaces

$$|\mathbf{BT}(1)| \xrightarrow{\text{hofib}(c_1 \cup c'_1)} K(\mathbb{Z}, 2) \times K(\mathbb{Z}, 2) \xrightarrow{c_1 \cup c'_1} K(\mathbb{Z}, 4),$$

where now c_1 and c'_1 are the ordinary (non-stacky) universal first Chern classes and we form their ordinary cup product. This identifies the geometric realization $|\mathbf{BT}(1)|$ with the space that was identified as the classifying space for topological T-duality pairs in [12, thm. 2.17].

Indeed, we now discuss how this is reflected on the level of super L_∞ -algebras, seeing that the T-duality Lie 2-algebra $b\mathcal{T}_1$ sits inside the coefficients $\mathcal{L}(\text{KU}/\text{BU}(1))/S^1$ as the classifying L_∞ -algebra for the ‘‘T-duality’’ pairs consisting of circle fibrations and degree-3 classes, while the rest of $\mathcal{L}(\text{KU}/\text{BU}(1))/S^1$ encodes the K-theory classes on this background.

Proposition 7.3. *There is a homotopy fiber sequence*

$$\begin{array}{ccc} \mathfrak{I}(\text{KU} \oplus \Sigma\text{KU}) & \longrightarrow & \mathfrak{I}(\mathcal{L}(\text{KU}/\text{BU}(1))/S^1) \\ & & \downarrow \\ & & b\mathcal{T}_1 \end{array}$$

which exhibits the cyclified L_∞ -algebras in Prop. 5.1 as homotopy quotients of $\mathfrak{I}(\text{KU})$ and $\mathfrak{I}(\Sigma\text{KU})$, respectively, by the Lie 2-algebra of the T-duality 2-group (Prop. 7.1)

$$\mathfrak{I}(\mathcal{L}(\text{KU}/\text{BU}(1))/\mathbb{R}) \simeq \mathfrak{I}((\text{KU} \oplus \Sigma\text{KU})/\mathcal{T}_1).$$


Proof. Consider the corresponding dual diagram of CE-algebras:

$$\begin{array}{ccccc} \text{CE}(\mathfrak{I}(\text{KU} \oplus \Sigma\text{KU})) & \longleftarrow & \text{CE}(\mathfrak{I}(\mathcal{L}(\text{KU}/\text{BU}(1))/S^1)) & \longleftarrow & \text{CE}(b\mathcal{T}_1) \\ \\ \left\{ \begin{array}{l} d\omega_{2p+2} = 0 \\ d\omega_{2p+3} = 0 \end{array} \right\} & \longleftarrow & \left\{ \begin{array}{l} dc_2 = 0, \quad d\tilde{c}_2 = 0 \\ dh_3 = c_2 \wedge \tilde{c}_2 \\ d\omega_{2p+2} = h_3 \wedge \omega_{2p} + c_2 \wedge \omega_{2p+1} \\ d\omega_{2p+3} = h_3 \wedge \omega_{2p+1} + \tilde{c}_2 \wedge \omega_{2p+2} \end{array} \right\} & \longleftarrow & \left\{ \begin{array}{l} dc_2 = 0, \quad d\tilde{c}_2 = 0 \\ dh_3 = c_2 \wedge \tilde{c}_2 \end{array} \right\}. \end{array}$$

The morphism on the right is the dual of a fibration, according to Prop. 2.4, since it is an inclusion of generators, and the morphism on the left is clearly its ordinary cofiber, hence is a model for its homotopy cofiber. \square

By the construction from Section 5 one sees that the triple $(c_2^A, c_2^B, \mu_{F_1}^9)$ defines a \mathcal{T}_1 -valued cocycle on $\mathbb{R}^{8,1|16+16}$. This leads to the following interrelations between models for 9-dimensional spacetime, their T-folds, and the dimensionally reduced twisted K-theory.

Definition 7.4. Define the 9-dimensional *T-fold super Lie 2-algebra* $\mathbb{R}_{\text{dbt}^A}^{8,1|16+16}$ to be the homotopy fiber of dimensional reduced IIA-fields, and define $\mathbb{R}_{\text{dbt}^B}^{8,1|16+16}$ to be the homotopy fiber of the dimensional reduced IIB-fields according to Example 5.2



Of course by Theorem 5.3 these two homotopy fibers are going to be equivalent, but for our purposes it is interesting to make explicit how they are equivalent:

Proposition 7.5. *The canonical model for the Chevalley-Eilenberg algebra of the super L_∞ -algebras $\mathbb{R}_{\text{dbt}^{A/B}}^{8,1|16+16}$ from Def. 7.4 is*

$$\text{CE}(\mathbb{R}_{\text{dbt}^A}^{8,1|16+16}) = \left\{ \begin{array}{l} d\psi^\alpha = 0 \\ de^a = \bar{\psi}\Gamma_a\psi \mid a \leq 8 \\ de_9^A = \bar{\psi}\Gamma_9^A\psi \\ de_9^B = \bar{\psi}\Gamma_9^B\psi \\ df_2 = \mu_{F_1}^{\text{IIA}} \end{array} \right\}$$

$$\text{CE}(\mathbb{R}_{\text{dbt}^B}^{8,1|16+16}) = \left\{ \begin{array}{l} d\psi^\alpha = 0 \\ de^a = \bar{\psi}\Gamma_a\psi \mid a \leq 8 \\ de_9^A = \bar{\psi}\Gamma_9^A\psi \\ de_9^B = \bar{\psi}\Gamma_9^B\psi \\ df_2 = \mu_{F_1}^{\text{IIB}} \end{array} \right\}.$$

These are just the objects in the top left and top right of the “topological” T -duality diagram in Prop. 6.2:

$$\mathbb{R}_{\text{dbl}^A}^{8,1|\mathbf{16}+\mathbf{16}} = p_A^* \widehat{\mathbb{R}^{9,1|\mathbf{16}+\overline{\mathbf{16}}}}, \quad \mathbb{R}_{\text{dbl}^B}^{8,1|\mathbf{16}+\mathbf{16}} = p_B^* \widehat{\mathbb{R}^{9,1|\mathbf{16}+\overline{\mathbf{16}}}}.$$

Proof. We consider the case of type IIA, the case for IIB is of course directly analogous. To find the homotopy fiber, we may replace the point inclusion by a fibration and then take the ordinary pullback. Dually, by Prop. 2.4 we need to find an inclusion of $\mathfrak{L}(\text{KU}/\text{BU}(1))/\mathbb{R}$ into CE-algebra whose underlying co-unary cochain complex is null, and then take the ordinary pushout of $(c_a^A, c_2^B, \mu_{F1}^9)$ along that,

$$\begin{array}{ccc} \text{CE}(\mathbb{R}_{\text{dbl}^A}^{8,1|\mathbf{16}+\mathbf{16}}) & \longleftarrow & A \xrightarrow{\simeq} 0 \\ \uparrow & \text{(p.o)} & \uparrow \\ \text{CE}(\mathbb{R}^{8,1|\mathbf{16}+\mathbf{16}}) & \longleftarrow & \text{CE}(b\mathcal{T}_1). \\ & \text{(} c_2^{\text{IIA}}, c_2^{\text{IIB}}, \mu_{F1}^9 \text{)} & \end{array}$$

To build such an A , we first need to adjoin generators e_9^A and e_9^B to $\text{CE}(b\mathcal{T}_1)$ to render the cocycles c_2 and \tilde{c}_2 trivial, by setting

$$de_9^A := c_2, \quad de_9^B := \tilde{c}_2.$$

Furthermore, we need to kill h_3 , which is a cocycle in the underlying dual chain complex of $b\mathcal{T}_1$. However, adjoining a generator f_2 with $df_2 = h_3$ would fail the condition that $d^2\theta = 0$. However, to remedy that we may set

$$df_2 = h_3 + e_9^A \wedge \tilde{c}_2,$$

which is consistent with $d^2\theta_2 = 0$ (recall that $dh_3 = -c_2 \wedge \tilde{c}_2$ according to Prop. 5.1) yet still removes h_3 in the cohomology of the underlying dual chain complex (since the co-unary restriction of d on θ_2 is h_3). In conclusion, a possible choice for A is the quotient

$$A := \text{CE}(b\mathcal{T}_1)[\{e_9^A, e_9^B, \theta_2\}] \Big/ \left(\begin{array}{l} de_9^A = c_2, \quad de_9^B = \tilde{c}_2 \\ db_2 = h_3 + e_9^A \wedge \tilde{c}_2 \end{array} \right).$$


Now the pushout in question is directly read off, using the values of $(c_2^{\text{IIA}}, c_2^{\text{IIB}}, h_3)$ from section 5. So c_2 gets identified with $\bar{\psi}\Gamma_9\psi$, \tilde{c}_2 gets identified with $\bar{\psi}\Gamma_9^B\psi$ and h_3 gets identified with $\mu_{F1}^A|_{8+1}$ in the pushout while \tilde{c}

is identified with $-(\pi_9^A)_*(\mu_{F_1}^A)$. Hence $h_3 + e_9^A \wedge \tilde{C}$ gets identified with

$$\mu_{F_1}^A|_{8+1} - e_9^A \wedge (\pi_9^A)_*(\mu_{F_1}^A) = \mu_{F_1}$$

as claimed. □

As a consequence, this shows how the equivalence in Prop. 6.2 follows from Theorem 5.3, since the operation of forming homotopy fibers sends weak equivalences to weak equivalences.



8. F-theory

We have discussed above in section 6 the “doubled” correspondence super Lie algebra

$$\mathbb{R}^{8+(1+1),1|32} := \mathbb{R}^{9,1|16+16} \times_{\mathbb{R}^{8,1|16+16}} \mathbb{R}^{9,1|16+\overline{16}}$$

of bosonic dimension $9 + 2$ that serves to interpolate between type IIA and type IIB super-spacetime via T-duality. It is immediate to see that this inherits the D0-brane 2-cocycle of its type IIA factor and hence extends to a super Lie algebra of bosonic dimension $10+2$:

Definition 8.1. Write $\mathbb{R}^{9+(1+1),1|32}$ for the central super Lie algebra extension of $\mathbb{R}^{8+(1+1),1|32}$ (def. 6.1) classified by the 2-cocycle $p_A^* c_2^M$, i.e. fitting

into a homotopy fiber sequence of the following form:

$$\begin{array}{ccccc} \mathbb{R}^{9+(1+1),1|32} & & & & \\ \text{hofib}(p_A^* c_2^M) \downarrow & & & & \\ \mathbb{R}^{8+(1+1),1|32} & \xrightarrow{p_A} & \mathbb{R}^{9,1|\mathbf{16}+\overline{\mathbf{16}}} & \xrightarrow{c_2^M} & b\mathbb{R}. \end{array}$$

where c_2^M is from Def. 2.14.

This super Lie algebra reflects both the relation between the M-theoretic 11d spacetime and the type IIA super-spacetime, as well as the correspondence of the latter to the type IIB super-spacetime.

Remark 8.2. Unwinding the definition, the CE-algebra of the super Lie algebra $\mathbb{R}^{9+(1+1),1|32}$ of Def. 8.1 has the following generating relations

$$\begin{aligned} de_0 &= \bar{\psi}\Gamma_0\psi, & de_1 &= \bar{\psi}\Gamma_1\psi, & \dots, & & de_8 &= \bar{\psi}\Gamma_8\psi, \\ de_9^B &= \bar{\psi}\Gamma_9^B\psi, & de_9 &= \underbrace{\bar{\psi}\Gamma_9\psi}_{\sigma_1}, & & & de_{10} &= \underbrace{\bar{\psi}\Gamma_{10}\psi}_{\sigma_3}, \end{aligned}$$

where on the right we have the 12 algebra elements from Def. 2.10.

Since the homotopy fiber of c_2^M alone is the local model for M-theoretic spacetime, hence for non-perturbative type IIA string theory, this suggests that the above homotopy fiber $\mathbb{R}^{9+(1+1),1|32}$ of $p_A^* c_2^M$ is similarly related to a non-perturbative description of type IIB string theory. That type IIB string theory ought to have a non-perturbative description in terms of 10+2 dimensional fibrations over 10-dimensional super-spacetime is known as the *F-theory* conjecture, due to [72]. In order to formalize aspects of this, we recall how this conjecture is motivated, see also for instance [40]:

The motivation of the F-theory conjecture from the M-theory conjecture.

- 1) Assume that a non-perturbative completion of type IIA string theory exists, given by a geometric theory on a 10+1 dimensional Riemannian circle fibration (M-theory).

- 2) Pass to the limit that the radius R_M of the circle fiber is infinitesimal to obtain perturbative type IIA string theory with coupling constant

$$g_{\text{IIA}} = R_M / \ell_s,$$

where ℓ_s is the string scale.

- 3) Consider the situation when the 10d IIA spacetime is itself a 9+1 dimensional circle fibration with circle fiber S_A^1 , so that in total the original 11d spacetime is a Riemannian torus fibration over 9d with fiber $S_M^1 \times S_A^1$.
- 4) Invoke perturbative T-duality to find an equivalent perturbative type IIB string theory on a 9+1-dimensional fibration with dual circle fiber S_B^1 of radius

$$R_B = \ell_s^2 / R_A$$

By the rules of perturbative T-duality, the coupling constant of the IIB theory is

$$\begin{aligned} g_{\text{IIB}} &= g_{\text{IIA}} \frac{\ell_s}{R_A} \\ &= R_M / R_A. \end{aligned}$$


Note that the last two steps can be combined [68]: M-theory compactified on a torus is supposed to be equivalent to type IIB superstring theory compactified on a circle in the limit of small volume of the torus.

- 5) While this was “derived” for infinitesimal R_M , observe that the resulting formula for the IIB coupling evidently extrapolates to finite R_M and then says that the IIB theory remains weakly coupled for finite (large) R_M if only R_A is suitably scaled along. Regard this as evidence for a non-perturbative version of T-duality which relates the non-perturbative IIA theory (M-theory) with some non-perturbative completion of IIB string theory, to be called *F-theory*.
- 6) Collect the geometric data that went into this construction: Retain information both of the S_A^1 -fiber, hence of the doubled correspondence space, and of the S_M^1 -fiber to obtain in total a 10+2-dimensional $S_M^1 \times S_A^1$ -fiber bundle over 10d type IIB spacetime.
- 7) Since the coupling constant g_{IIB} depends only on the ratio R_M / R_A , hence only on the complex structure of this torus, conclude that this is to be regarded as a 10+2-dimensional elliptic fibration.

- 8) Check that the action of S-duality on type IIB fields corresponds to the automorphisms of the elliptic fiber.

We now observe that the super Lie algebra from Def. 8.1 has just the right structure to be the model for the super-tangent space of this F-theory elliptic fibration.

Proposition 8.3. *The super Lie algebra $\mathbb{R}^{9+(1+1),1|32}$ from Def. 8.1 fits into a diagram of super L_∞ -algebras of the following form*



where each square is a (homotopy) pullback square (homotopy Cartesian).

Proof. This follows immediately from the pasting law for homotopy pullbacks of super L_∞ -algebras. It is also directly checked explicitly. \square

In particular, this says that the composite diagonal morphism

$$\mathbb{R}^{8+(1+1),1|32} \longrightarrow \mathbb{R}^{8,1|16+16}$$

exhibits its domain as a $\text{Lie}(S_M^1 \times S_A^1)$ -fibration over the type IIB super-spacetime.

Remark 8.4. Remark 8.2 says that the bosonically 12-dimensional $\mathbb{R}^{9+(1+1),1|32}$ is a super Lie algebra, but not a super-Minkowski Lie algebra,

hence not a super-symmetry algebra in the sense of spacetime supersymmetry. Instead, the diagram in Prop. 8.3 shows how it projects onto various genuine super-spacetime algebras.

This is consistent with the observations and assertions in the literature that F-theory does not have a straightforward spacetime interpretation and, furthermore, that the alternative seems to emanate from superalgebras, but not of the usual type. In [7], it is shown how the algebras in 10, 11, and 12 dimensions can be described by a web of dualities as different faces of the ortho-symplectic superalgebra $\text{OSp}(1|32)$. In particular, for F-theory the corresponding algebra has no vector operator, so that there is no generator for translations. This implies that F-theory, in contrast to M-theory, has no straightforward spacetime interpretation. It was noted in [18, end of Sec. 6] that the fact that the type IIA and IIB theories have $\text{SL}(2, \mathbb{R})$ and $\text{SL}(1|1)$ global symmetry, respectively, highlights the similarity between the symmetries of the two, but more importantly point to a possibly fermionic twelve-dimensional origin.

Hence, if we regard $\mathbb{R}^{9+(1+1),1|32}$ as a \mathbb{R}^2 -fibration over the IIB-spacetime according to Prop. 8.3, then the last check in the list of F-theory desiderata from above is that its fiber automorphisms induce S-duality transformation on the type IIB fields:

Definition 8.5. Write

$$\mu_{F1}^F, \mu_{Dp}^F \in \text{CE}(\mathbb{R}^{9+(1+1),1|32})$$

for the pullback of the type IIB F1/D p -brane cocycles from Def. 4.9 along the projection

$$\mathbb{R}^{9+(1+1),1|32} \longrightarrow \mathbb{R}^{9,1|16+16}$$

from Prop. 8.3.

Hence in terms of generators, via Remark 8.2, these CE-elements have the same form as in Def. 4.9, but with all generators e^a renamed as e_B^a . Notably the cocycles for the F1- and D1-string on the F-theory space read

$$\mu_{F1}^F = i (\bar{\psi} \Gamma_a^B \Gamma_{10} \psi) \wedge e_B^a \quad \text{and} \quad \mu_{D1}^F = i (\bar{\psi} \Gamma_a^B \Gamma_9 \psi) \wedge e_B^a.$$

Proposition 8.6. *Rotation in the (9, 10)-plane of the fiber \mathbb{R}^2 of the F-theory super Lie algebra from Def. 8.1 is a super Lie algebra automorphism*

$$\begin{aligned}
 \phi_\alpha : \mathbb{R}^{9+(1+1),1|32} &\longrightarrow & \mathbb{R}^{9+(1+1),1|32} \\
 e_B^a &\mapsto & e_B^a \\
 \begin{pmatrix} e^9 \\ e^{10} \end{pmatrix} &\mapsto & \begin{pmatrix} \cos(\alpha) & \sin(\alpha) \\ -\sin(\alpha) & \cos(\alpha) \end{pmatrix} \cdot \begin{pmatrix} e^9 \\ e^{10} \end{pmatrix} \\
 \psi &\mapsto & \exp\left(\frac{\alpha}{4}\Gamma_9\Gamma_{10}\right)\psi.
 \end{aligned}$$

Under this automorphism the F1-string and the D1-string are turned into superpositions of each other, in that their F-theoretic cocycles from Def. 8.5 satisfy:

$$\phi_\alpha^* \begin{pmatrix} \mu_{D1}^F \\ \mu_{F1}^F \end{pmatrix} = \begin{pmatrix} \cos(\alpha) & \sin(\alpha) \\ -\sin(\alpha) & \cos(\alpha) \end{pmatrix} \cdot \begin{pmatrix} \mu_{D1}^F \\ \mu_{F1}^F \end{pmatrix}.$$

Proof. That ϕ_α is indeed a super Lie algebra automorphism follows with Remark 2.12. Hence acting with ϕ_α on μ_{F1}^F and μ_{D1}^F leaves all the factors in the explicit formula in Def. 8.5 invariant, except for Γ_9 and Γ_{10} , which are rotated into each other. □

Under the identification $\sigma_2 = -\Gamma_9\Gamma_{10}$ from Def. 2.10 this is the action of S-duality on the F1/D1 cocycles according to [27, Remark 4.9].

Acknowledgement

We thank Jonathan Pridham for discussion of his model structure for L_∞ -algebras. U.S. thanks Thomas Nikolaus for discussion of T-folds.

D.F. thanks New York University Abu Dhabi (NYUAD) for kind hospitality during the writing of this paper.

H.S. would like to thank Ralph Kaufmann and Alexander Voronov, the organizers of the AMS special sessions on Topology and Physics in Minneapolis in October 2016, for the opportunity to present the results of this work.

U.S. thanks the Max Planck Institute for Mathematics (MPI) in Bonn for kind hospitality during the writing of this paper. U.S. was supported by RVO:67985840.

References

- [1] A. Achúcarro, J. Evans, P. Townsend, and D. Wiltshire, *Super p -branes*, Phys. Lett. **B 198** (1987), 441–446.
- [2] E. Alvarez, L. Alvarez-Gaumé, and Y. Lozano, *An introduction to T-duality in string theory*, Nucl. Phys. Proc. Suppl. **41** (1995), 1–20.
- [3] I. Bachas and K. Sfetsos, *T-duality and world-sheet supersymmetry*, Phys. Lett. **B 349** (1995), 448–457.
- [4] E. Bergshoeff and M. de Roo, *D-branes and T-duality*, Phys. Lett. **B 380** (1996), 265–272.
- [5] E. Bergshoeff, E. Sezgin, and P. K. Townsend, *Superstring actions in $D = 3, 4, 6, 10$ curved superspace*, Phys. Lett. **B 169** (1986), 191–196.
- [6] E. Bergshoeff, E. Sezgin, and P. K. Townsend, *Supermembranes and eleven dimensional supergravity*, Phys. Lett. **B 189** (1987), 75–78.
- [7] E. Bergshoeff and A. Van Proeyen, *The many faces of $\text{OSp}(1|32)$* , Class. Quantum Grav. **17** (2000), 3277.
- [8] F. Borceaux, *Handbook of Categorical Algebra I: Basic Category Theory*, Cambridge University Press (1994)
- [9] P. Bouwknegt, J. Evslin, and V. Mathai, *T-duality: Topology change from H-flux*, Commun. Math. Phys. **249** (2004), 383–415.
- [10] P. Bouwknegt and V. Mathai, *D-branes, B-fields and twisted K-theory*, J. High Energy Phys. **03** (2000), 007.
- [11] U. Buijs, Y. Félix, and A. Murillo, *L_∞ -models of based mapping spaces*, J. Math. Soc. Japan **63** (2011), no. 2, 503–524.
- [12] U. Bunke and T. Schick, *On the topology of T-duality*, Rev. Math. Phys. **17** (2005), 77–112.
- [13] U. Bunke, P. Rumpf, and T. Schick, *The topology of T-duality for T^n -bundles*, Rev. Math. Phys. **18** (2006), 1103.
- [14] L. Castellani, *Chiral $D = 10, N = 2$ supergravity on the group manifold I: Free differential algebra and solution of Bianchi identities*, Nucl. Phys. **B 294** (1987), no. 4, 877–889.
- [15] L. Castellani, R. D’Auria, and P. Fré, *Supergravity and Superstrings – A Geometric Perspective*, World Scientific, Singapore, 1991.

- [16] M. Cederwall, *Double supergeometry*, J. High Energy Phys. **06** (2016), 155.
- [17] C. Chryssomalakos, J. de Azcárraga, J. Izquierdo, and C. Pérez Bueno, *The geometry of branes and extended superspaces*, Nucl. Phys. **B 567** (2000), 293–330.
- [18] E. Cremmer, B. Julia, H. Lu, and C. N. Pope, *Dualisation of Dualities, II: Twisted self-duality of doubled fields and superdualities*, Nucl. Phys. **B 535** (1998), 242–292.
- [19] M. Cvetič, H. Lu, C. N. Pope, and K. S. Stelle, *T-duality in the Green-Schwarz formalism, and the massless/massive IIA duality map*, Nucl. Phys. **B 573** (2000), 149i–176.
- [20] R. D’Auria and P. Fré, *Geometric supergravity in $D = 11$ and its hidden supergroup*, Nucl. Phys. **B 201** (1982), 101–140.
- [21] R. D’Auria, P. Fré, P. Grassi, and M. Trigiante, *Pure spinor superstrings on generic type IIA supergravity backgrounds*, J. High Energy Phys. **0807** (2008), 059.
- [22] P. Deligne and D. Freed, *Supersolutions*, in P. Deligne et al. Quantum Fields and Strings, Amer. Math. Soc., Providence, RI, 1999,
- [23] M. Duff, P. Howe, T. Inami, and K. Stelle, *Superstrings in $D = 10$ from Supermembranes in $D = 11$* , Phys. Lett. **B 191** (1987), 70.
- [24] P. de Medeiros, J. Figueroa-O’Farrill, E. Méndez-Escobar, and P. Ritter, *On the Lie-algebraic origin of metric 3-algebras*, Commun. Math. Phys. **290** (2009), 871–902.
- [25] J. Figueroa-O’Farrill and J. Simón, *Supersymmetric Kaluza-Klein reductions of M2 and M5-branes*, Adv. Theor. Math. Phys. **6** (2003), 703–793.
- [26] D. Fiorenza, H. Sati, and U. Schreiber, *Extended higher cup-product Chern-Simons theories*, J. Geom. Phys. **74** (2013), 130–163.
- [27] D. Fiorenza, H. Sati, and U. Schreiber, *Super Lie n -algebra extensions, higher WZW models, and super p -branes with tensor multiplet fields*, Intern. J. Geom. Meth. Mod. Phys. **12** (2015), 1550018.
- [28] D. Fiorenza, H. Sati, and U. Schreiber, *The WZW term of the M5-brane and differential cohomotopy*, J. Math. Phys. **56** (2015), 102301.

- [29] D. Fiorenza, H. Sati, and U. Schreiber, *Rational sphere valued supercocycles in M-theory and type IIA string theory*, Journal of Geometry and Physics **114** (April 2017).
- [30] S. F. Hassan, *$SO(d, d)$ transformations of Ramond-Ramond fields and space-time spinors*, Nucl. Phys. **B 583** (2000), 431–453.
- [31] M. Hatsuda, K. Kamimura, and W. Siegel, *Superspace with manifest T-duality from type II superstring*, J. High Energy Phys. **06** (2014), 039.
- [32] M. Henneaux and L. Mezincescu, *A Sigma model interpretation of Green-Schwarz covariant superstring action*, Phys. Lett. **B 152** (1985), 340–342.
- [33] K. Hess, *Rational homotopy theory: a brief introduction*, in: Interactions between Homotopy Theory and Algebra, pp. 175–202, Contemp. Math. **436**.
- [34] V. Hinich, *DG coalgebras as formal stacks* J. Pure and Applied Algebra **162** (2001), 209–250.
- [35] P. Hirschhorn, *Model Categories and Their Localizations*, Amer. Math. Soc., Providence, RI, 2003.
- [36] K. Hori, *D-branes, T-duality and index theory*, Adv. Theor. Math. Phys. **3** (1999), 281.
- [37] J. Huerta and U. Schreiber, *M-Theory from the superpoint*, Letters in Mathematical Physics (2018).
- [38] C. Hull, *A geometry for non-geometric string backgrounds*, J. High Energy Phys. **0510** (2005), 065.
- [39] C. Hull, *Doubled geometry and T-folds*, J. High Energy Phys. **0707** (2007), 080.
- [40] C. Johnson, *From M-theory to F-theory, with branes*, Nucl. Phys. **B 507** (1997), 227–244.
- [41] J. D. S. Jones, *Cyclic homology and equivariant homology*, Invent. Math. **87** (1987), 403–423.
- [42] A. Kapustin, *D-branes in a topologically nontrivial B-field*, Adv. Theor. Math. Phys. **4** (2000), 127.

- [43] A. Konechny and A. Schwarz, *On $(k \oplus l|q)$ -dimensional supermanifolds*, in J. Wess and V. Akulov (eds.), *Supersymmetry and Quantum Field Theory*, Lecture Notes in Physics **509**, Springer (1998).
- [44] B. Kulik and R. Roiban, *T-duality of the Green-Schwarz superstring*, *J. High Energy Phys.* **0209** (2002), 007.
- [45] T. Lada and J. Stasheff, *Introduction to sh Lie algebras for physicists*, *Int. J. Theo. Phys.* **32** (1993), 1087–1103.
- [46] T. Lada and M. Markl, *Strongly homotopy Lie algebras*, *Commun. in Algebra* **23** (1995), 2147–2161.
- [47] J. A. Lind, H. Sati, and C. Westerland, *A higher categorical analogue of topological T-duality for sphere bundles*, [arXiv:1601.06285](https://arxiv.org/abs/1601.06285).
- [48] J.-L. Loday, *Free loop space and homology*, [arXiv:1110.0405](https://arxiv.org/abs/1110.0405).
- [49] H. Lu, C. N. Pope, E. Sezgin, and K. S. Stelle, *Stainless super p-branes*, *Nucl. Phys. B* **456** (1995), 669–698.
- [50] V. Mathai and H. Sati, *Some relations between twisted K-theory and E_8 gauge theory*, *J. High Energy Phys.* **0403** (2004), 016.
- [51] G. Moore, *Physical mathematics and the future*, talk at Strings 2014. <http://www.physics.rutgers.edu/~gmoore/PhysicalMathematicsAndFuture.pdf>
- [52] G. Moore and E. Witten, *Self-duality, Ramond-Ramond Fields, and K-theory*, *JHEP* 0005 (2000), 032.
- [53] T. Nikolaus and K. Waldorf, *Higher geometry for non-geometric T-duals*, [arXiv:1804.00677](https://arxiv.org/abs/1804.00677).
- [54] T. Nikolaus, U. Schreiber, and D. Stevenson, *Principal ∞ -bundles – General theory*, *J. Homotopy Related Structr.* **10** (2015), 749–801.
- [55] T. Nikolaus, U. Schreiber, and D. Stevenson, *Principal ∞ -bundles – Presentations*, *J. Homotopy and Related Structr.* **10** (2015), 565–622.
- [56] B. Nikolić and B. Sazdović, *T-dualization of type II superstring theory in double space*, [arXiv:1505.06044](https://arxiv.org/abs/1505.06044).
- [57] S. Palmer and C. Saemann, *M-brane models from non-abelian gerbes*, *J. High Energy Phys.* **1207** (2012), 010.
- [58] R. Percacci and E. Sezgin, *On target space duality in p-branes*, *Mod. Phys. Lett. A* **10** (1995), 441–450.

- [59] J. P. Pridham, *Unifying derived deformation theories*, Adv. Math. **224** (2010), 772–826.
- [60] D. Quillen, *Rational homotopy theory*, Ann. of Math., 2nd Series, **90** (1969), no. 2, 205–295.
- [61] M. Sakaguchi, *IIB-branes and new spacetime superalgebras*, J. High Energy Phys. **0004** (2000), 019.
- [62] H. Sati and U. Schreiber, *Lie n -Algebras of BPS charges*, J. High Energy Phys. **2017** (2017), 87.
- [63] H. Sati, U. Schreiber, and J. Stasheff, *L_∞ -algebra connections and applications to String- and Chern-Simons n -transport*, Quantum Field Theory, pp. 303–424, Birkhäuser, Basel (2009).
- [64] H. Sati, U. Schreiber, and J. Stasheff, *Twisted differential String- and Fivebrane structures*, Commun. Math. Phys. **315** (2012), 169–213.
- [65] U. Schreiber, *Differential cohomology in a cohesive ∞ -topos*, <https://ncatlab.org/schreiber/show/differential+cohomology+in+a+cohesive+topos>.
- [66] U. Schreiber, *Structure theory for higher WZW terms*, lectures at Higher Structures in String Theory and Quantum Field Theory, ESI Vienna 2015. <https://ncatlab.org/schreiber/show/Structure+Theory+for+Higher+WZW+Terms>.
- [67] U. Schreiber, *Modern physics Formalized in modal homotopy type theory*, talk at J. Ladyman (org.), Applying Homotopy Type Theory to Physics, Bristol, April 2015. <https://ncatlab.org/schreiber/show/Modern+Physics+formalized+in+Modal+Homotopy+Type+Theory>.
- [68] J. H. Schwarz, *M theory extensions of T-duality*, Frontiers in Quantum Field Theory (1995).
- [69] A. Sen, *T-duality of p -branes*, Mod. Phys. Lett. **A11** (1996), 827–834.
- [70] W. Siegel, *Superspace duality in low-energy superstrings*, Phys. Rev. **D48** (1993), 2826–2837.
- [71] D. Sullivan, *Infinitesimal computations in topology*, Publ. Math. de I.H.É.S. **47** (1977), 269–331.
- [72] C. Vafa, *Evidence for F-theory*, Nucl. Phys. **B 469** (1996), 403–418.

- [73] P. van Nieuwenhuizen, *Free graded differential superalgebras*, Group Theoretical Methods in Physics, pp. 228–247, Lecture Notes in Physics **180**, M. Serdaroglu and E. Inonu (eds.), Springer-Verlag, Berlin, Germany, 1983.
- [74] M. Vigué-Poirrier and D. Burghelea, *A model for cyclic homology and algebraic K-theory of 1-connected topological spaces*, J. Differential Geom. **22** (1985), 243–253.
- [75] M. Vigué-Poirrier and D. Sullivan, *The homology theory of the closed geodesic problem*, J. Differential Geom. **11** (1976), 633–644.
- [76] E. Witten, *D-branes and K-theory*, J. High Energy Phys. **12** (1998), 019.

DIPARTIMENTO DI MATEMATICA, LA SAPIENZA UNIVERSITÀ DI ROMA
PIAZZALE ALDO MORO 2, 00185 ROME, ITALY
E-mail address: `fiorenza@mat.uniroma1.it`

UNIVERSITY OF PITTSBURGH, PITTSBURGH, PA 15260, USA
AND NEW YORK UNIVERSITY, ABU DHABI, UAE
E-mail address: `hsati@nyu.edu`

MATHEMATICS INSTITUTE OF THE ACADEMY
ŽITNA 25, 115 67 PRAHA 1, CZECH REPUBLIC
E-mail address: `schreiber@math.cas.cz`, `urs.schreiber@gmail.com`