

On non-degeneracy of Riemannian Schwarzschild-anti de Sitter metrics

PIOTR T. CHRUSCIEL, ERWANN DELAY, AND PAUL KLINGER

We prove that the TT -gauge-fixed linearised Einstein operator is non-degenerate for Riemannian Kottler (“Schwarzschild-anti de Sitter”) metrics with dimension- and topology-dependent ranges of mass parameter. We provide evidence that this remains true for all such metrics except the spherical ones with a critical mass.

| | | |
|------------|--|-------------|
| 1 | Introduction | 1216 |
| 2 | Riemannian Kottler metrics | 1218 |
| 3 | Master functions | 1224 |
| 4 | Gauge vectors for vanishing master functions | 1235 |
| 5 | Triviality of the kernel | 1239 |
| Appendix A | Divergence of a symmetric two-tensor in coordinates | 1241 |
| Appendix B | Divergence & double divergence of h_{ij}^S and h_{aj}^S | 1244 |
| Appendix C | The decomposition of h | 1245 |
| Appendix D | The asymptotics of scalar modes | 1246 |
| Appendix E | The asymptotics of vector modes | 1249 |
| Appendix F | The linearised Birkhoff theorem | 1250 |
| Appendix G | The $l = 1$ modes for $K = 1, n = 2$ | 1252 |

| | | |
|-------------------|--|-------------|
| Appendix H | A linearised Birkhoff-type theorem with $K = 0$ | 1258 |
| Appendix I | A linearised Birkhoff-type theorem with $K = -1$ | 1260 |
| Appendix J | The Riemannian Kerr anti-de Sitter metrics | 1261 |
| Appendix K | The master equation for vector perturbations | 1264 |
| References | | 1266 |

1. Introduction

There is currently considerable interest in the literature in spacetimes with a negative cosmological constant. In particular one is interested in existence of stationary black hole solutions of the Einstein equations with $\Lambda < 0$, with or without sources, and in properties thereof. Many such solutions have been constructed numerically, e.g. in [6, 7, 25, 26, 30].

In [2, 9] we showed how to construct infinite dimensional families of non-singular stationary black hole spacetimes, solutions of the Einstein equations with a negative cosmological constant in vacuum or with various matter sources, assuming that a suitable linearised operator was an isomorphism. In [2, Proposition D.2] we proved the isomorphism property at Kottler solutions with negatively curved sections of conformal infinity for some dimension-dependent explicit ranges of the mass parameter (the whole range of masses in space-time dimension four, and the whole range of negative masses in all dimensions). However, the case of flat or positively curved conformal infinity has been open so far. The aim of this work is to prove the optimal result for all topologies in spacetime dimension four, and to provide partial answers to this problem in higher dimensions. This extends immediately the applicability of the existence theorems of [2, 9] to the topologies and dimensions covered here.

Indeed, we prove:

Theorem A. *Let us denote by P_L the linearisation, at Riemannian Kottler metrics (2.1) with negative cosmological constant, of the TT-gauge-fixed Einstein operator. Then:*

- 1) P_L has no L^2 -kernel in spacetime dimension $n + 2 = 4$ except for spherical black holes with mass parameter

$$(1.1) \quad \mu = \frac{n}{n + 1} \left(\ell \sqrt{\frac{n - 1}{n + 1}} \right)^{n-1} .$$

- 2) P_L has no L^2 -kernel for toroidal and higher genus black holes for all mass parameters μ in dimension $n = 2$, as well as for open ranges of parameters $\mu \in (0, \mu(n))$ for $n > 2$ and $K = 0$, where $\mu(n) > 0$ solves a polynomial equation; cf. Table 1 in low dimensions.

Theorem A is a rewording of Theorem 5.1 below.

In order to prove an equivalent of part (2) of Theorem A for black holes in higher-dimensions with $K \in \{1, -1\}$ it remains to establish a higher-dimensional topology-independent linearised-Birkhoff-type theorem and to treat the $l = 1$ modes for $K = 1$ in dimension $n > 2$. This is done in the accompanying paper [17].

We also propose a scheme, supported by numerical evidence, that could establish an equivalent of part (1) of Theorem A for all topologies and space-time dimensions, except for the exceptional value (1.1) of the mass in the spherical case. We thus conjecture:

Conjecture B. P_L has no L^2 -kernel except if $K = 1$ and μ is given by (1.1).

In view of the results in [17], in order to establish the conjecture one would need to prove global existence for the ODE that we integrate numerically.

We emphasise that our four-dimensional result, namely point (1) of Theorem A holds regardless of the genus of the black hole. The proof in [2, Proposition D.2] of the higher-genus case $K = -1$ of (1) uses a completely different method, and leads to restricted ranges of masses in higher dimension. Our method here provides an alternative argument which, at the level of numerical tests, covers all masses.

Now, the idea in [2, 9] is to show that the construction of stationary Lorentzian solutions near a known static d -dimensional metric $\mathring{\mathbf{g}}$ can be carried-out using the implicit function theorem near a Riemannian d -dimensional partner metric $\mathring{\mathbf{g}}$. Consider then the operator P_L obtained by linearising the Einstein equations at the ‘‘Wick rotated’’ metric $\mathring{\mathbf{g}}$ associated to $\mathring{\mathbf{g}}$ and imposing the TT gauge. The construction of [2, 9] applies if P_L has

no L^2 -kernel; we then say that the Riemannian metric \mathring{g} is non-degenerate. We show here that non-degeneracy holds for Kottler metrics with spherical black hole horizons in four spacetime dimension, and for toroidal black hole horizons in all dimensions, for wide ranges of masses.

Some comments on the proof are in order. For this, let h be an element of the L^2 -kernel of the operator P_L defined above. Our proof relies heavily on the remarkable construction of master functions of Ishibashi and Kodama [18]. Indeed, h can be decomposed into a sum of eigentensors of relevant operators, which we will refer to as modes. The modes split into two families, which we will refer to as exceptional modes and master modes. The master modes of h are controlled by the Ishibashi-Kodama master functions, which we show to be zero under the conditions above. We use the vanishing of the master modes to show existence of a vector field *with controlled asymptotic behaviour* which can be used to gauge-away the relevant part of h . Likewise we show that the special modes are pure gauge in the last, controlled, sense. Adding the associated vector fields and using the TT-gauge condition, together with suitable Birkhoff-type linearised theorems, we establish that the kernel is trivial in the cases listed in Theorem A.

We note that a non-trivial kernel of P_L implies existence of growing linear modes in the corresponding spacetime, except perhaps for variations in the direction of nearby stationary metrics. So proving that P_L has no L^2 -kernel has implications for the dynamics of the associated Lorentzian solutions. However, non-degeneracy does not imply linear dynamical stability on the Lorentzian side, since non-degeneracy only excludes modes with a specific spectrum of frequencies.

2. Riemannian Kottler metrics

We start with a review of the Riemannian counterparts of the “generalised Kottler [22] metrics”, also known as “Schwarzschild-anti de Sitter metrics”, or “Birmingham metrics” [5].

In what follows we will be making extensive use of [18], in order to be consistent as much as possible with their notations we set

$$n = d - 2,$$

where d is the *dimension of spacetime*. (The reader is warned that this *does not* coincide with our notations in [2, 9], where n denotes space dimension $d - 1$.)

The metrics of interest read

$$(2.1) \quad \mathring{g} = \underbrace{\left(\frac{r^2}{\ell^2} + K - \frac{2\mu}{r^{n-1}} \right)}_{=:f(r)} dt^2 + \frac{dr^2}{\frac{r^2}{\ell^2} + K - \frac{2\mu}{r^{n-1}}} + r^2 h_K,$$

where ℓ is a constant related to the cosmological constant $\Lambda < 0$ by the formula

$$\ell = \sqrt{-\frac{n(n+1)}{2\Lambda}} > 0,$$

μ is a real constant, and $K \in \{0, \pm 1\}$. Furthermore, h_K is a metric on an n -dimensional Einstein manifold nN . The metric \mathring{g} is Einstein if the Ricci tensor of h_K equals $(n-1)Kh_K$. This will be assumed in what follows.

We require that f has positive zeros, we denote by $r_0 > 0$ the largest such zero, which we assume to be of first order (as will necessarily be the case if $K \geq 0$ and $\mu > 0$):

$$(2.2) \quad \frac{r_0^2}{\ell^2} + K - \frac{2\mu}{r_0^{n-1}} = 0.$$

After introducing a new coordinate \check{r} by the formula

$$(2.3) \quad \check{r}(r) = \int_{r_0}^r \frac{1}{\sqrt{\frac{s^2}{\ell^2} + K - \frac{2\mu}{s^{n-1}}}} ds,$$

one can rewrite the metric (2.1) as

$$(2.4) \quad \mathring{g} = d\check{r}^2 + \check{r}^2 H(\check{r}) dt^2 + r^2 h_K,$$

where H is obtained by dividing g_{tt} by \check{r}^2 . Elementary analysis, using the fact that r_0 is a simple zero of F , shows that

$$H(0) = \frac{f'(r_0)^2}{4} = \left(\frac{(n+1)\ell^{-2}r_0^2 + K(n-1)}{2r_0} \right)^2.$$

This implies that a periodic identification of t with period

$$(2.5) \quad T := \frac{4\pi}{f'(r_0)}$$

guarantees that $d\check{r}^2 + \check{r}^2 H(\check{r}) dt^2$ is a smooth metric on \mathbb{R}^2 with a rotation axis at $\check{r} = 0$. As a result, (2.4) defines a smooth Riemannian metric on

$$(2.6) \quad M := \mathbb{R}^2 \times {}^nN.$$

The metric (2.1) can be smoothly conformally compactified by introducing, for large r , a coordinate $\rho := 1/r$ and rescaling:

$$(2.7) \quad \rho^2 \mathring{g} = (\ell^{-2} + K\rho^2 - 2\mu\rho^{n+1})dt^2 + \frac{d\rho^2}{\ell^{-2} + K\rho^2 - 2\mu\rho^{n+1}} + h_K.$$

Hence, the $(n+1)$ -dimensional conformal boundary $\partial M := \{\rho = 0\}$ of M is diffeomorphic to $S^1 \times {}^n N$, with conformal metric

$$(2.8) \quad \ell^{-2}dt^2 + h_K.$$

As already pointed out, it has been shown in [2, Appendix D] that all such solutions with $n+1 = 3$ and $K = -1$ are non-degenerate. Furthermore, in that reference some non-degenerate families of higher-dimensional such solutions have been described explicitly. We thus conjecture that the ranges of mass parameters there are not sharp, and that the arguments proposed here can be used to establish non-degeneracy for all values of μ when $K \leq 0$.

In the analysis of the master equations below we assume that $({}^n N, h_K)$ is a (locally) maximally symmetric compact manifold. When $K \geq 0$ the hypothesis that $\Lambda < 0$ and that of existence of a black hole with $r_0 > 0$ implies

$$\mu > 0.$$

When $K = -1$ one needs instead

$$(2.9) \quad \mu > \mu_{\min} := -\frac{1}{n+1} \left(\frac{n+1}{\ell^2(n-1)} \right)^{\frac{1-n}{2}},$$

with the corresponding outermost-horizon radius $r_0 = 1/\sqrt{\frac{n+1}{\ell^2(n-1)}}$.

The idea is to reduce the question of non-degeneracy to the Riemannian equivalent of the “master equations” of Ishibashi and Kodama [12, 18] as follows:

- 1) rewrite the master equations in a Riemannian form;
- 2) work-out the asymptotic behaviour of the master fields corresponding to L^2 -elements of the kernel of the shifted Lichnerowicz operator P_L defined below;
- 3) prove that all associated solutions of the master equations are trivial;
- 4) prove that elements of the L^2 -kernel with trivial master fields vanish identically.

Point (1) above is straightforward once the impressive work in [18] has been carried out, but the remaining parts require some work. We note that point (4) captures the fact that the master equations contain the whole gauge-invariant information about the linearised gravitational field.

Consider, thus, an element $h = h_{\mu\nu}dx^\mu dx^\nu$ of the L^2 -kernel of

$$(2.10) \quad P_L := \Delta_L + 2(n + 1),$$

where Δ_L is the Lichnerowicz Laplacian, acting on symmetric two-tensor fields $h_{\mu\nu}$ as [4, § 1.143]

$$(2.11) \quad \Delta_L h_{\mu\nu} = -\nabla^\alpha \nabla_\alpha h_{\mu\nu} + R_{\mu\alpha} h^\alpha{}_\nu + R_{\nu\alpha} h^\alpha{}_\mu - 2R_{\mu\alpha\nu\beta} h^{\alpha\beta}.$$

It follows from [23] that $|h|_{\mathfrak{g}} = O(\rho^{n+1})$, or in local coordinates near the conformal boundary,

$$(2.12) \quad h_{\mu\nu} = O(\rho^{n-1}).$$

Let us show, first, that h satisfies the linearised Einstein equations. For this, recall that the Hodge Laplacian acting on one-forms is defined as

$$(2.13) \quad \Delta_H := d_g^* d + d d_g^* = \nabla^* \nabla + \text{Ric}.$$

If the Ricci tensor is covariantly constant, then [24]

$$\text{div} \circ \Delta_L = \Delta_H \circ \text{div}.$$

This implies that L^2 -elements of the kernel of P_L are divergence-free: Indeed, assume that h is in the L^2 -kernel of P_L and let $u = \text{div } h$. We have

$$0 = \int \langle \text{div}(\Delta_L + 2(n + 1))h, u \rangle = \int \langle (\Delta_H + 2(n + 1))u, u \rangle,$$

so

$$\int |du|^2 + |d_g^* u|^2 + 2(n + 1)|u|^2 = 0,$$

and $u \equiv 0$ follows. Note that there are no boundary terms in the integration-by-parts above by, e.g., [23].

Next, recall that we always have

$$\text{Tr} \circ \Delta_L = -\Delta \circ \text{Tr}$$

(we use the convention $\Delta = \nabla^\alpha \nabla_\alpha$). It follows that elements of the L^2 -kernel of P_L are trace-free.

The linearisation of the trace-shifted Ricci tensor reads

$$(2.14) \quad D(\text{Ric} + (n+1))h_{\mu\nu} = \frac{1}{2}\Delta_L h_{\mu\nu} + (n+1)h_{\mu\nu} - (\text{div}^* \text{div grav } h)_{\mu\nu},$$

where

$$(2.15) \quad \begin{aligned} \text{grav } h &= h - \frac{1}{2}\text{Tr}_g hg, & (\text{div } h)_\mu &= -\nabla^\nu h_{\mu\nu}, \\ (\text{div}^* w)_{\mu\nu} &= \frac{1}{2}(\nabla_\mu w_\nu + \nabla_\nu w_\mu), \end{aligned}$$

and where Tr denotes the trace (note the geometers' convention to include a negative sign in the definition of divergence).

We have just seen that tensors in the kernel of P_L are transverse and traceless. It follows from (2.14)–(2.15) that they are also in the kernel of the linearised vacuum Einstein operator.

Now, it follows immediately from the analysis in [18] (see also [21, Appendix B]) that, similarly to the Lorentzian case, the linearised Einstein operator for metrics (2.1) on manifolds as in (2.6) leaves invariant the subspaces of “scalar”, “vector”, and “tensor modes”.

The modes $l = 0, 1$ require separate attention. A detailed analysis of this case, in Lorentzian signature, four spacetime dimensions, and spherical black-hole topology, can be found in [11].¹ Indeed, it is shown there that the $l = 0, 1$ modes are, up to a gauge transformation, variations of the mass parameter of the Schwarzschild-anti de Sitter metrics ($l = 0$), or variations of the angular-momentum parameter when the Schwarzschild-anti de Sitter metric is viewed as a member of the Kerr-anti de Sitter family of metrics ($l = 1$). We show in Appendices F–H how to adapt the arguments of [11] to the cases of interest here.

As is well known, solutions of the linearised Riemannian Einstein equations corresponding to variations of the mass parameter are not in L^2 , except if μ is given by (2.21) below, which is seen as follows: Replacing t in (2.1) by a 2π -periodic variable φ and r by ρ , defined as

$$(2.16) \quad \rho^2 = \frac{4f}{(f'(r_0))^2},$$

¹Compare [14, Section 5]. The variable \mathbf{x} used there is actually awkward for the $l = 0$ modes because its definition involves an operator which becomes singular at $r = 3m$ precisely for this mode. The calculations there for spherically symmetric perturbations become clear if instead of \mathbf{x} one uses $\mathcal{B}^{-1}\mathbf{x} = (1 - \frac{3m}{r})\mathbf{x}$. The operator \mathcal{B} has been introduced there to obtain simple formulae for all remaining modes.

we find, in all spacetime dimensions $d = n + 2$,

$$(2.17) \quad \mathring{g} = \rho^2 d\varphi^2 + \frac{(f'(r_0))^2}{(f'(r(\rho)))^2} d\rho^2 + r(\rho)^2 h_K.$$

Variations of the mass parameter take the form

$$(2.18) \quad \frac{d\mathring{g}}{d\mu} = \left[\underbrace{\frac{d(f'(r_0))}{d\mu} \frac{2f'(r_0)}{(f'(r))^2}}_I - \frac{2(f'(r_0))^2}{(f'(r))^3} \left(\underbrace{\frac{\partial f'}{\partial \mu}(r)}_{II} + \underbrace{f''(r) \frac{dr}{d\mu}}_{III} \right) \right] d\rho^2 + \underbrace{2r \frac{dr}{d\mu}}_{IV} h_K.$$

The part marked I contributes an asymptotically constant term to the norm $|d\mathring{g}/d\mu|_{\mathring{g}}^2$, unless $d(f'(r_0))/d\mu$ vanishes. The term coming from II is of order r^{-2n-2} . For the parts III and IV we consider

$$(2.19) \quad \begin{aligned} \frac{dr}{d\mu} &= \frac{\partial f^{-1}}{\partial \mu} \left(\frac{(f'(r_0))^2 \rho^2}{4} \right) + (f^{-1})' \left(\frac{(f'(r_0))^2 \rho^2}{4} \right) \frac{f'(r_0) \rho^2}{2} \frac{d(f'(r_0))}{d\mu} \\ &= O(r^{-n}) + O(r^2) \frac{d(f'(r_0))}{d\mu}. \end{aligned}$$

This implies that the terms in $|d\mathring{g}/d\mu|_{\mathring{g}}^2$ coming from III and IV are of order r^{-2n-2} if $d(f'(r_0))/d\mu = 0$ and of order r^2 otherwise.

Therefore $d\mathring{g}/d\mu \in L^2$ if and only if

$$(2.20) \quad \frac{d(f'(r_0))}{d\mu} = 0.$$

This equation has no solution with $r_0 > 0$ when $K = 0$ or $K = -1$, while if $K = 1$ this leads to

$$(2.21) \quad r_0 = r_c := \ell \sqrt{\frac{n-1}{n+1}}, \quad \mu = \mu_c := \frac{n}{n+1} \left(\ell \sqrt{\frac{n-1}{n+1}} \right)^{n-1}.$$

In Appendix J we review the Riemannian Kerr-anti de Sitter metrics, and we prove there that variations of angular momentum lead to linearised solutions of the vacuum Einstein equations which are not in L^2 either.

3. Master functions

The master functions of Ishibashi and Kodama, which we denote by $\Phi_{S,I}$, $\Phi_{V,I}$, and $\Phi_{T,I}$, where the index I runs over all eigenfunctions of Δ_{h_K} on nN (i.e. $l = l(I)$), are solutions of a two-dimensional Schrödinger equation

$$(3.1) \quad \Delta_{2\mathring{g}}\Phi_{i,I} - V_{i,l}\Phi_{i,I} = 0, \quad i \in \{S, V, T\},$$

where

$$(3.2) \quad 2\mathring{g} = f dt^2 + \frac{dr^2}{f},$$

with $f = f(r)$ as in (2.1), while

$$(3.3) \quad V_{S,k} = \frac{1}{16r^2 (m + xn(n+1)/2)^2} \times \left\{ \begin{aligned} & [n^3(n+2)(n+1)^2x^2 - 12n^2(n+1)(n-2)mx \\ & + 4(n-2)(n-4)m^2]\ell^{-2}r^2 + n^4(n+1)^2x^3 \\ & + n(n+1)[4(2n^2 - 3n + 4)m \\ & + n(n-2)(n-4)(n+1)K]x^2 \\ & - 12n[(n-4)m + n(n+1)(n-2)K]mx \\ & + 16m^3 + 4Kn(n+2)m^2 \end{aligned} \right\}, \quad k^2 > nK,$$

$$(3.4) \quad V_{V,k_V} = \frac{1}{r^2} \left\{ \begin{aligned} & k_V^2 + K + \frac{n(n-2)}{4}K \\ & + \frac{n(n-2)}{4}\ell^{-2}r^2 - 3\frac{n^2\mu}{2r^{n-1}} \end{aligned} \right\}, \quad k_V^2 - (n-1)K > 0,$$

$$(3.5) \quad V_{T,k_T} = \frac{1}{r^2} \left\{ \begin{aligned} & k_T^2 + 2K + \frac{n(n-2)}{4}K \\ & + \frac{n(n+2)}{4}\ell^{-2}r^2 + \frac{n^2\mu}{2r^{n-1}} \end{aligned} \right\},$$

where

$$x = 2\mu r^{1-n}, \quad m = k^2 - nK.$$

Note that, compared to the equivalent equations in [18], our equation (3.1) omits a term $f(r)^{-1}$ in front of $V_{i,l}$. We have absorbed this factor into the definition of the potentials $V_{i,l}$. For a sphere there are no master functions for $l = 0, 1$; for a torus no master functions $\Phi_{S,I}$ and $\Phi_{V,I}$ for $k = 0 = k_V$;

and no master functions for $k = 0$ (“scalar master potential”) in the higher genus case.

For $K = 1$ we have $k^2 = l(l + n - 1)$ with $l \geq 0$, $k_V^2 = l(l + n - 1) - 1$ with $l \geq 1$, and $k_T^2 = l(l + n - 1) - 2$ with $l \geq 2$.

For $K = 0$ and a flat torus at infinity,

$$\mathbb{T}^n = \underbrace{S^1 \times \dots \times S^1}_{n \text{ factors}}$$

with each S^1 -coordinate of period 2π , and with $h_{K=0} \equiv \gamma_{ij} dx^i dx^j$ where $\partial_\mu \gamma_{ij} = 0$, we have

$$(3.6) \quad k^2, k_V^2, k_T^2 \in \{\gamma^{ij} k_i k_j\}_{k_i \in \mathbb{N}}.$$

Equation (3.6) is an immediate consequence of decompositions into Fourier series.

After a periodic identification of t with period T given by (3.2), the metric (3.2) becomes a smooth rotation-invariant conformally-compactifiable metric on \mathbb{R}^2 , with a smooth center of rotation at $\check{r} = 0$ (equivalently, at $r = r_0$).

Chasing through the definitions of [18] we show in Appendices D.1, D.2, and K, using the notation there, that for linearised solutions which are in L^2 we have $H_T = O(\rho^{n+1})$, $|f_a|_{\mathfrak{g}} = O(\rho^{n+1})$, $|f_{ab}|_{\mathfrak{g}} = O(\rho^{n+1})$, etc., resulting in

$$(3.7) \quad \Phi_{S,I} = \begin{cases} O(\rho^{n/2-1}), & n > 2; \\ o(1), & n = 2, \end{cases}$$

$$(3.8) \quad \Phi_{V,I} = O(\rho^{n/2}),$$

$$(3.9) \quad \Phi_{T,I} = O(\rho^{n/2+1}).$$

3.1. Vanishing via the maximum principle

We want to prove the vanishing of the master functions for L^2 -elements of the kernel. The simplest case occurs when the potentials in (3.3)–(3.5) are non-negative. Since all the $\Phi_{i,I}$ ’s tend to zero as r tends to infinity, the vanishing of the corresponding master function follows from the maximum principle. This leads to rigorous statements for restricted dimensions and masses. For the purposes of the proof of Theorem A, the main conclusions of the analysis that follows in this section are:

Proposition 3.1. *Let h be an element of the L^2 -kernel of the operator P_L defined in (2.10). Then:*

- 1) *The associated tensor master functions vanish.*
- 2) *The associated scalar master functions vanish if $n = 2$ and either $K = 0$ or $K = 1$ and $l \geq 2$.*

Remark 3.2. Note that the above concerns only the master modes, i.e. those which are controlled by the master functions $\Phi_{i,I}$. The scalar modes with $k = 0$ (in particular, the modes corresponding to the variation of mass), and the modes $K = 1$ and $l = 1$ (which include the variation of angular momentum) require separate attention.

Since we have already shown decay of the master functions, it remains to analyze positivity of the potentials occurring in the master equations. We have not attempted an exhaustive analysis of this, but we certainly have positivity under the following circumstances, keeping in mind that we are working in the region where $f > 0$:

- for V_{T,k_T} .
- When $n = 2$ the formula (3.3) for $V_{S,k}$ simplifies to

$$(3.10) \quad \begin{cases} V_{S,k} = \frac{1}{3} \left(\frac{6\mu}{r^3} + \frac{k^2-2}{r^2} + \frac{2\ell^2(k^6-3k^4+4)+216\mu^2}{(6\mu+(k^2-2)r)^2} \right), & K = 1; \\ V_{S,k} = \frac{1}{3} \left(\frac{6\mu}{r^3} + \frac{k^2}{r^2} + \frac{2\ell^2k^6+216\mu^2}{\ell^2(6\mu+k^2r)^2} \right), & K = 0; \\ V_{S,k} = \frac{1}{3} \left(\frac{6\mu}{r^3} + \frac{k^2+2}{r^2} + \frac{2\ell^2(k^6+3k^4-4)+216\mu^2}{\ell^2(6\mu+(k^2+2)r)^2} \right), & K = -1. \end{cases}$$

So $V_{S,k} \geq 0$ if $n = 2$ and either $K = 0$, or $K = -1$ and $\mu \geq 0$, or $K = 1$ and $l \geq 1$. Further, when $K = 0$ and $n > 4$ one also finds $V_{S,k} \geq 0$ when $0 < \mu < \mu(n)\ell^{n-1}$ for a function $\mu(n)$ which solves a polynomial equation. Indeed, by inserting $\mu = 0$ into (3.3), we see that for $n > 4$ the scalar potential is positive for small masses. As we also know that it is positive at large r , we find that the limiting value of μ is reached when $V_{S,k}$ has a minimum with value zero, i.e. when the resultant of the denominators of $V_{S,k}$ and $\partial_r V_{S,k}$, after dividing each by a suitable power of r corresponding to the root $r = 0$, has a positive solution. This resultant is a polynomial in μ and k . Now, $V_{S,k}$ is positive for all r and sufficiently large k at any fixed value of μ , so the problem is solved by choosing the smallest zero of a finite number of polynomials in μ parameterised by k . Similarly when $K = 1$ and $n \in \{4, 5\}$.

Note that

$$(3.11) \quad V_{S,k} \xrightarrow{r \rightarrow \infty} V_{S,k}(\infty) := \frac{(n-2)(n-4)}{4\ell^2},$$

so that the dimensions $n = 3, 4$ (thus, spacetime dimensions five and six) require different considerations in any case.

- It holds that $V_{V,k_V} \geq 0$ if

$$(3.12) \quad \begin{cases} \frac{n(n+1)}{2} (2\mu\ell^{1-n})^{2/(n+1)} \leq k_V^2, & K = 0; \\ l \geq 2 \text{ and } 0 < \mu < \ell, & K = 1 \text{ and } n = 2; \\ l \geq 2 \text{ and } 0 < \mu < \frac{2}{9}\ell^2, & K = 1 \text{ and } n = 3; \\ k_V \geq k_{V,1}(n) \text{ and } 0 < \mu < \mu_1(k_V, n)\ell^{n-1}, & K = 1 \text{ and } n \geq 4; \\ k_V \geq k_{V,-1}(n) \text{ and } \mu_{\min} < \mu < \mu_{-1}(k_V, n)\ell^{n-1}, & K = -1, \end{cases}$$

for some functions $k_{V,K}(n)$ and $\mu_K(k_V, n) > 0$, where μ_{\min} is given by (2.9).

3.2. Vanishing using the bottom of the spectrum

In this section we will prove further vanishing theorems for the master functions by studying the first L^2 -eigenvalue λ_1 , and more precisely the kernel, of Schrödinger operators $\nabla^* \nabla + V$ with a smooth potential V and an asymptotically hyperbolic metric ${}^2\mathring{g}$ on \mathbb{R}^2 ,

$${}^2\mathring{g} = d\check{r}^2 + \check{r}^2 H(\check{r}) dt^2 = \frac{dr^2}{f} + f dt^2,$$

where, as before,

$$f = \frac{r^2}{\ell^2} + K - \frac{2\mu}{r^{n-1}} > 0, \quad \frac{df}{dr} = 2 \left(\frac{r}{\ell^2} + (n-1) \frac{\mu}{r^n} \right) > 0,$$

with \check{r} is as in (2.3). Recall that r^{-s} is in L^2 if and only if $s > 1/2$. Using Theorem C of [23] we have the following:

Lemma 3.3. *Let V be a smooth potential on \mathbb{R}^2 . We assume that $-\Delta + V$ has a non-trivial indicial interval with smallest characteristic index s_- . If $-\Delta + V$ has no L^2 -kernel, then this last operator has no non trivial function of order $o(r^{-s_-})$ at infinity in its kernel.*

We would like to apply this lemma to (3.1). We note the following indicial exponents for the master equations, which turn-out to depend only on the type of the mode: in obvious notation,

$$(3.13) \quad s_{\pm}^S = \frac{1 \pm \sqrt{1 + (n - 2)(n - 4)}}{2},$$

$$(3.14) \quad s_{\pm}^V = \frac{1 \pm \sqrt{1 + n(n - 2)}}{2},$$

$$(3.15) \quad s_{\pm}^T = \frac{1 \pm \sqrt{1 + n(n + 2)}}{2}.$$

Remark 3.4. Note that for $n = 3$ we have $s_{\pm}^S = \frac{1}{2}$ so we can not use directly Theorem C of [23] as in Lemma 3.3, but this weight $1/2$ is the critical weight to be in L^2 so the conclusion of Lemma 3.3 remains true if we replace $o(r^{-s-})$ with $O(r^{-s--\varepsilon})$ for some $\varepsilon > 0$, which is the case in our applications.

We now study the L^2 -spectrum of our Schrödinger operators.

Lemma 3.5. *Let X be a vector field in $W_{loc}^{1,1}$. The first L^2 -eigenvalue of $\nabla^*\nabla + V$ acting on functions is bigger than or equal to the almost-everywhere-infimum of*

$$(3.16) \quad \tilde{V} := \nabla_i X^i - |X|^2 + V.$$

Proof. Let $\delta_X \in \mathbb{R} \cup \{-\infty\}$ be the a.e.-infimum of \tilde{V} . For any function u smooth with compact support we have $\int \nabla_i(u^2 X^i) = 0$, so

$$\int |\nabla u|^2 + \int |X|^2 u^2 \geq -2 \int u X^i \nabla_i u = \int u^2 \nabla_i X^i.$$

We thus obtain

$$\int u(\nabla^*\nabla + V)u \geq \int \tilde{V}u^2 \geq \delta_X \int u^2. \quad \square$$

In order to apply Lemma 3.5, we need to find a vector field $X \in W_{loc}^{1,1}$ such that we have, weakly,

$$\tilde{V} = \operatorname{div} X - |X|^2 + V \geq 0,$$

with \tilde{V} positive on an open set. Indeed, if such a vector field X exists, and if u is in the L^2 -kernel of $-\Delta + V$, then (see the last inequality in the

proof above) u has to vanish on the open set where $\tilde{V} > 0$, so by unique continuation $u = 0$ everywhere.

We look for X of the form $X = S\partial_r$ where S is a function on \mathbb{R}^2 , then

$$(3.17) \quad \tilde{V} = \operatorname{div} X - |X|^2 + V = \partial_r S - \frac{S^2}{f} + V.$$

This should be compared with the Ishibashi-Kodama modified potential \tilde{V} [12] which differs from ours by a multiplicative factor f (similarly to the potentials V_{\dots} entering the master equations).

So whenever their \tilde{V} is non negative with their choice of S , we obtain positivity of our \tilde{V} by taking $X = S\partial_r$. We can then apply Lemma 3.5 after verifying that X belongs to $W_{\text{loc}}^{1,1}$. This works very well for vector modes in any dimension, leading to:

Corollary 3.6. *Under the condition $k_V^2 > (n - 1)K$, the L^2 kernel of $-\Delta + V_{V,k_V}$ is trivial for all $n \geq 2$ and $K \in \{-1, 0, 1\}$. We deduce that any function in the kernel which is of order $o(r^{-s_V^-})$ at infinity, where s_V^- is given by (3.14), is trivial.*

Remark 3.7. Corollary 3.6 applies to $K = 1$ and $l \geq 2$, since then we have

$$k_V^2 - (n - 1)K = (l - 1)(l + n) > 0.$$

Proof of Corollary 3.6. In [12, Equation (2.21)] one takes $S = \frac{nf}{2r}$, so we take $X = \frac{nf}{2r}\partial_r$, which is in $W_{\text{loc}}^{1,1}$. With this choice of X , the potential in the vector master equation (as in [12, Equation (2.17)])

$$V_V = \frac{1}{r^2} \left[k_V^2 - (n - 1)K + \frac{n(n + 2)}{4}f - \frac{n}{2}r \frac{df}{dr} \right],$$

is transformed to

$$\tilde{V}_V = \frac{1}{r^2} [k_V^2 - (n - 1)K] > 0.$$

Whence the triviality of the kernel, as claimed. □

Let us turn our attention to the scalar potential. The following corollary of Lemma 3.5 gives an alternative proof of point (2) of Proposition 3.1, and extends that last proposition to the case $K = -1$ (cf., however, Remark 3.2):

Corollary 3.8. *Under the condition $k^2 > \max(0, 2K)$, the L^2 kernel of $-\Delta + V_{S,k}$ is trivial for $n = 2$ for any $K = -1, 0, 1$. We deduce that any function of order $o(1)$ at infinity and in the kernel is trivial.*

Proof. We use the function S chosen in [19, Equation (6.23)], where here we have $n = 2$, $Q = \delta = 0$, $H = h_- = H_- = m + \frac{6\mu}{r}$, $m = k^2 - 2K$, that is

$$S = \frac{f}{H} \frac{dH}{dr},$$

so $X = S\partial_r$ is in $W_{\text{loc}}^{1,1}$. Our scalar potential V_S becomes (see [19, Equation (6.24)] with a factor f removed)

$$\tilde{V}_S = \frac{k^2 m}{r^2 H},$$

which is positive if $k \neq 0$ and $m = k^2 - 2K > 0$. □

To continue, set

$$(3.18) \quad \lambda := \ell \inf_{r \geq r_0} \frac{d\sqrt{f}}{dr} \in \begin{cases} \{1\}, & K \in \{-1, 0\}; \\ (0, 1), & K = 1. \end{cases}$$

We have:

Corollary 3.9. *The L^2 first eigenvalue of $\nabla^* \nabla$ acting on functions is bigger than or equal to $\lambda^2 / (4\ell^2)$.*

Proof. We choose $X = c\sqrt{f}\partial_r \in W_{\text{loc}}^{1,1}$, where the positive constant c will be chosen later. It follows that $|X| = c$ and $\text{div } X = c\partial_r(\sqrt{f}) > 0$, with

$$(\text{div } X)^2 = c^2 \frac{(\partial_r f)^2}{4f} = c^2 \left(\frac{1}{\ell^2} + (n-1) \frac{\mu}{r^{n+1}} \right)^2 \left(\frac{1}{\ell^2} + K \frac{1}{r^2} - \frac{2\mu}{r^{n+1}} \right)^{-1},$$

where $2\mu = r_0^{n+1}/\ell^2 + Kr_0^{n-1}$, $r_0 > 0$, $r \geq r_0$. If $K = 0, -1$ the function $(\text{div } X)^2$ is decreasing (recall that we have assumed (2.9) when $K = -1$) and tends to $c^2\ell^{-2}$, but for $K = 1$, this function has a positive infimum less than $c^2\ell^{-2}$, say $\lambda^2 c^2 \ell^{-2}$. Thus

$$\text{div } X - |X|^2 \geq \lambda c \ell^{-1} - c^2.$$

The right-hand side is maximised by $c = \lambda \ell^{-1} / 2$, giving

$$\text{div } X - |X|^2 \geq \lambda^2 / (4\ell^2). \quad \square$$

To apply Corollary 3.9, note that the L^2 -kernel of $-\Delta + V$ will be trivial if $V + \lambda^2/(4\ell^2) \geq 0$ and is strictly positive somewhere. More generally, from the proof above with $X = \frac{\lambda}{2\ell}\sqrt{f}\partial_r$, this kernel will be trivial if

$$(3.19) \quad \tilde{V} = V + \frac{\lambda}{2\ell} \left(\frac{d\sqrt{f}}{dr} - \frac{\lambda}{2\ell} \right) \geq 0,$$

and positive on an open set. This leads to (compare Proposition 3.1 and Remark 3.2):

Proposition 3.10. *Let h be an element of the L^2 -kernel of the operator P_L defined in (2.10). Let $n \in \{4, 5\}$ if $K = 1$, and $n \in \mathbb{N}$ otherwise.*

- 1) *Let $K \in \{0, 1\}$. There exists a function $\mu(K, n) > 0$ such that for $0 < \mu < \mu(K, n)$ the corresponding scalar master functions vanish.*
- 2) *Let $K = -1$, and let $\lambda_1 > 0$ be the first non-zero eigenvalue of the scalar Laplacian of $({}^nN, h_K)$. There exists a function $\mu(\lambda_1, n) > \mu_{\min}$ and an $\varepsilon(n) > 0$ such that for $\lambda_1 \geq \varepsilon(n)$ and $\mu_{\min} < \mu < \mu(\lambda_1, n)$ the corresponding scalar master functions vanish.*

Remark 3.11. By a result of Schoen [28] we have, for the case $K = -1$, the lower bound $\lambda_1 \geq (n - 1)^2/4$ under the assumption that the volume of nN is sufficiently small, as described in detail there. We have checked in dimensions $2 \leq n \leq 10$ that Schoen’s estimate is sufficient to obtain $\mu(\lambda_1, n) > 0$, we give approximate values of an upper bound of the μ ’s allowed in Table 1. Thus the result is not empty.

The actual values of $\varepsilon(n)$ are lower than this estimate, we give approximate values for dimensions $2 \leq n \leq 10$ in Table 2. □

Similarly to the discussion in the paragraph following (3.10), Equation (3.19) can be used to determine $\mu(K, n)$ as a root of a high order polynomial. These polynomials grow with dimension, being extremely long already in $n = 4$, so that they cannot be usefully displayed here.

As an illustration, we list approximate numerical values of $\mu(K, n)$ for dimensions $n \leq 10$ in Table 1 and the corresponding values of r_0 in Table 3.

Let us return to the definition (3.17) of \tilde{V} . It is tempting to try to choose S so that $\tilde{V} \geq 0$ globally, thus solving the equation

$$(3.20) \quad \partial_r S - \frac{S^2}{f} + V = \tilde{V} \geq 0,$$

| n | $K = 0$ | $K = 1$ | $K = -1$ |
|-----|-----------------------|----------------------|----------|
| 2 | ∞ | ∞ | ∞ |
| 3 | 1/48 | – | 1/24 |
| 4 | 4.4×10^{-3} | 3.5×10^{-1} | 0.11 |
| 5 | 4.7×10^{-4} | 2.2×10^{-2} | 0.12 |
| 6 | 3.9×10^{-5} | – | 0.19 |
| 7 | 2.7×10^{-6} | – | 0.26 |
| 8 | 1.6×10^{-7} | – | 0.35 |
| 9 | 8.6×10^{-9} | – | 0.45 |
| 10 | 4.0×10^{-10} | – | 0.57 |

Table 1: For $\ell^{n-1}\mu$ between $\ell^{n-1}\mu_{\min}$ (where μ_{\min} is the lowest value giving positive r_0 , i.e. $\mu_{\min} = 0$ for $K \in \{0, 1\}$ and $\mu_{\min} < 0$ for $K = -1$, given by (2.9)) and the values shown here we obtain trivial L^2 kernel of $-\Delta + V_S$. For $K = -1$ we have used the fact that the first non-zero eigenvalue of the Laplacian on nN is $\geq (n-1)^2/4$, which holds by [28] under the assumption that the volume of nN is sufficiently small, cf. [28] for details. For the cases marked “–” there is no $\mu > 0$ such that the inequality (3.19) is satisfied for all relevant k (i.e. those not covered by the linearised Birkhoff theorem or the discussion of $l = 1$ modes in Appendix G).

| n | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|------------------|---|------|------|------|------|------|------|------|------|
| $\varepsilon(n)$ | 0 | 0.76 | 1.13 | 1.51 | 1.57 | 1.88 | 1.98 | 2.26 | 2.33 |

Table 2: This table shows approximate values of the function $\varepsilon(n)$, appearing in Proposition 3.10, obtained from the inequality (3.19).

for some given function $\tilde{V}(r)$ which is positive in an open set.² For this, we numerically solved the ODE (3.17) for S using MATHEMATICA, with \tilde{V}_S set to zero, with initial value $S(r_0) = 0$. The integration was performed using the `StiffnessSwitching` method and a `WorkingPrecision` of 30–100 depending on the value of μ . If the resulting solutions have a zero at a point $r_{S,0}$ such that the potential V_S is positive for all $r \geq r_{S,0}$, we extend S for $r \geq r_{S,0}$ by setting it to be equal to zero there, giving a non-negative \tilde{V}_S for all $r > r_0$ and $\tilde{V}_S > 0$ for $r \geq r_{S,0}$. This procedure works for all attempted

²Once this work was completed we noticed [16], where very similar ideas are used in a related context.

| n | $K = 0$ | $K = 1$ | $K = -1$ |
|-----|----------|----------|----------|
| 2 | ∞ | ∞ | ∞ |
| 3 | 0.45 | – | 1.01 |
| 4 | 0.38 | 0.76 | 1.08 |
| 5 | 0.31 | 0.43 | 1.11 |
| 6 | 0.25 | – | 1.13 |
| 7 | 0.22 | – | 1.12 |
| 8 | 0.19 | – | 1.13 |
| 9 | 0.16 | – | 1.14 |
| 10 | 0.14 | – | 1.15 |

Table 3: This table is similar to Table 1, except that we list the value of r_0 , the largest zero of f , rather than the value of μ . Thus, for r_0 smaller than the values shown here we obtain a trivial L^2 kernel of $-\Delta + V_S$ with $\ell = 1$.

mass parameters and dimensions $n \geq 5$ for $K = 0, 1$, and all positive masses for $K = -1$. Indeed, we tested masses in all orders of magnitude between the limits in Table 1 and $\mu = 10^{30}$ (with $\ell = 1$) for dimensions $n = 5, 6, 100$, and at least 5 values of the mass parameter within this range for each dimension $5 \leq n \leq 100$.

In dimension $n = 4$, for $K = 1$, this construction of S works for μ small enough, giving a larger mass range than that obtained from (3.19) (up to $\mu \approx 1.5$ for $\ell = 1$). A typical result for S and \tilde{V}_S is shown in Figure 1.




Figure 1: A typical numerical solution for S with $\tilde{V} = 0$, cut off at the third zero of S . Here $n = 7, K = 0, k = 1, \mu = 100, \ell = 1$.

In dimensions $n = 3$ and 4 for all K , and for $K = -1$ with negative mass parameter in all dimensions, the numerical solution S obtained in this way does not have a zero (except for $K = 1$ with low mass) but appears to exist globally. Since we need positivity of \tilde{V} somewhere to conclude, we instead numerically solved the ODE (3.17) with $\tilde{V} = \varepsilon r^{-2}$ for $\varepsilon > 0$ (for $\mu > 0$ we can choose $\varepsilon = 1$, for $\mu < 0$ the choice of ε depends on μ), with again initial value $S(r_0) = 0$. The resulting numerical solution S appears to grow asymptotically linearly for large r and we therefore cannot set it to zero at some $r_{S,0}$, as a negative jump in S would add a negative distributional component to \tilde{V} . Instead we just use the solution directly, without cutting off at finite distance. This works in fact for all masses, dimensions, and $K = 0, 1, -1$, but the results are less conclusive as one has then to rely on the behaviour of the solution up to $r = \infty$, while the numerical integration necessarily stops at some finite r . A typical result for S and \tilde{V}_S for this approach is shown in Figure 2.




Figure 2: A typical numerical solution for S with $\tilde{V} = 1/r^2$. Here $n = 3$, $K = 1$, $l = 2$, $\mu = 10$, $\ell = 1$.

In any case, our numerical experiments strongly hint at the following conjecture:

Conjecture 3.12. *There are no non-trivial scalar master modes associated with the L^2 -kernel of the operator P_L given by (2.10).*

4. Gauge vectors for vanishing master functions

In this section we study the consequences of the vanishing of the master functions. We consider perturbations of the $(n + 2)$ -dimensional metric

$$\mathring{g} = g_{ab}dx^a dx^b + r^2\gamma_{ij}dx^i dx^j,$$

where

$$g_{ab}dx^a dx^b = fdt^2 + f^{-1}dr^2,$$

and where γ is the metric of an n -dimensional manifold with constant sectional curvature K . The indices a, b take values t, r , while i, j are “angular” indices. We will denote by D the covariant derivative associated to g and by \widehat{D} that of γ .

As emphasised in [18], a general metric perturbation h can be split into “scalar”, “vector”, and “tensor” parts as

$$(4.1) \quad h = h^S + h^V + h^T.$$

An analytic description of the splitting (4.1), without referring to a mode-decomposition, is presented in Appendix C.

We show in Appendix A below that the TT conditions are consistent with the splitting above.

The components in (4.1) can be expanded as [18, Sections 2.1, 5.1 and 5.2]

$$(4.2) \quad \begin{aligned} h_{ab}^S &= \sum_I f_{ab,I}^S \mathbb{S}^I, & h_{ai}^S &= \sum_I r f_{a,I}^S \mathbb{S}_i^I, \\ h_{ij}^S &= \sum_I 2r^2 (H_{L,I}^S \gamma_{ij} \mathbb{S}^I + H_{T,I}^S \mathbb{S}_{ij}^I), \end{aligned}$$

$$(4.3) \quad h_{ab}^V = 0, \quad h_{ai}^V = \sum_I r f_{a,I}^V \mathbb{V}_i^I, \quad h_{ij}^V = \sum_I 2r^2 H_{T,I}^V \mathbb{V}_{ij}^I,$$

$$(4.4) \quad h_{ab}^T = 0, \quad h_{ai}^T = 0, \quad h_{ij}^T = \sum_I 2r^2 H_{T,I}^T \mathbb{T}_{ij}^I,$$

where the following holds: for $K = 0$, the above are obvious decompositions in Fourier series, with coordinate-independent tensor components leading to

zero eigenvalues. For $K = 1$ (see, e.g. [8, 27])

$$(4.5) \quad (\widehat{\Delta}_n + k^2)\mathbb{S}^I = 0, \quad k^2 = l(l + n - 1), \quad l = 0, 1, 2, \dots,$$

$$(4.6) \quad (\widehat{\Delta}_n + k_V^2)\mathbb{V}_i^I = 0, \quad k_V^2 = l(l + n - 1) - 1, \quad l = 1, 2, \dots,$$

$$(4.7) \quad (\widehat{\Delta}_n + k_T^2)\mathbb{T}_{ij}^I = 0, \quad k_T^2 = l(l + n - 1) - 2, \quad l = 2, \dots, \quad n > 2.$$

For $K = -1$ there is no example of compact quotient of \mathbb{H}^n with explicit values of the whole spectrum, but we have a countable set of increasing eigenvalues of finite multiplicity:

$$k^2 = (\lambda_0 = 0), \lambda_1, \dots, \quad k_V^2 = \lambda_{V,0}, \dots,$$

where $\lambda_{V,0} \geq (n - 1)$ by non-negativity of the Hodge Laplacian (2.13). The first non zero eigenvalue λ_1 is greater or equal than a computable constant [28] that can be chosen to be $(n - 1)^2/4$ for sufficiently small volumes.

Whatever the value of $K \in \{-1, 0, 1\}$ one sets

$$(4.8) \quad \mathbb{S}_i^I = -\frac{1}{k}\widehat{D}_i\mathbb{S}^I, \quad k \neq 0,$$

$$(4.9) \quad \mathbb{S}_{ij}^I = \frac{1}{k^2}\widehat{D}_i\widehat{D}_j\mathbb{S}^I + \frac{1}{n}\gamma_{ij}\mathbb{S}^I, \quad k \neq 0,$$

$$(4.10) \quad \mathbb{V}_{ij}^I = -\frac{1}{2k_V}(\widehat{D}_i\mathbb{V}_j^I + \widehat{D}_j\mathbb{V}_i^I) = -\frac{1}{2k_V}\mathcal{L}_{\mathbb{V}^I}\gamma_{ij}, \quad k_V \neq 0.$$

We define the sets \mathbb{I}_S^* and \mathbb{I}_V^* of indices I corresponding to modes governed by the scalar and vector master equations:

$$(4.11) \quad \mathbb{I}_S^* = \{I \mid k(I) > 0, k^2(I) > nK\},$$

$$(4.12) \quad \mathbb{I}_V^* = \{I \mid k_V(I) > 0, k_V^2(I) > (n - 1)K\},$$

and denote by \mathbb{I}_S^\diamond and \mathbb{I}_V^\diamond their complements.

Note that k_V never vanishes when $K \in \{\pm 1\}$, and that those vector fields \mathbb{V}_j^I for which $l(I)$ in (4.6) equals one, are Killing vector fields of γ_{ij} (cf., e.g., the eigenvalue λ_1^1 for the Hodge Laplacian in [8, Theorem 3.1]), so that the associated tensor \mathbb{V}_{ij}^I vanishes.

From now on we assume that the master functions Φ_S , Φ_V , and Φ_T vanish.

The vanishing of the tensor potential directly gives $h^T = 0$ [18, (5.4)].

The vanishing of the vector potential implies [18, (5.10)–(5.13)], for modes such that $k_V \neq 0$,

$$(4.13) \quad f_{a,I}^V = -\frac{r}{k_V} D_a H_{T,I}^V.$$

Let us define the vector field $Y^{V\star}$ as

$$(4.14) \quad Y^{V\star} := -r^2 \sum_{I \in \mathbb{I}_V^\star} \frac{H_{T,I}^V}{k_V(I)} \mathbb{V}_j^I \gamma^{ij} \partial_i.$$

The convergence of the series is justified in Appendix K.

Let us denote by $h_{ij}^{V\star}$ and $h_{ia}^{V\star}$ tensors in which we have collected all those modes in h_{ij}^V and h_{ia}^V which are governed by the master equations, and by $h_{ij}^{V\circ}$ and $h_{ia}^{V\circ}$ whatever remains:

$$(4.15) \quad h_{ai}^{V\star} = \sum_{I \in \mathbb{I}_V^\star} r f_{a,I}^V \mathbb{V}_i^I,$$

$$(4.16) \quad h_{ij}^{V\star} = \sum_{I \in \mathbb{I}_V^\star} 2r^2 H_{T,I}^V \mathbb{V}_{ij}^I,$$

$$(4.17) \quad h_{ai}^{V\circ} = \sum_{I \in \mathbb{I}_V^\circ} r f_{a,I}^V \mathbb{V}_i^I,$$

$$(4.18) \quad h_{ij}^{V\circ} = \sum_{I \in \mathbb{I}_V^\circ} 2r^2 H_{T,I}^V \mathbb{V}_{ij}^I.$$

From (K.10) and (K.15), Appendix K, we obtain

$$(4.19) \quad \|D_a(r^{-2} h_{ij}^{V\star}) dx^i dx^j\|_{H^k(n_N)} \leq 2r^{-2} \|h_{ai}^{V\star} dx^i\|_{H^{k+1}(n_N)},$$

$$(4.20) \quad h_{ij}^V = h_{ij}^{V\circ} + r^2 (\mathcal{L}_{Y^{V\star}} \gamma)_{ij} = h_{ij}^{V\circ} + (\mathcal{L}_{Y^{V\star}} \mathring{\mathbf{g}})_{ij}.$$

Using (4.13), the mixed components h_{ai}^V take the form

$$(4.21) \quad \begin{aligned} h_{ai}^V &= \sum_I r f_{a,I}^V \mathbb{V}_i^I = h_{ai}^{V\circ} - \sum_{I \in \mathbb{I}_V^\star} \frac{r^2}{k_V} \mathbb{V}_i^I D_a H_{T,I}^V \\ &= h_{ai}^{V\circ} + D_a Y_i^{V\star} = h_{ai}^{V\circ} + (\mathcal{L}_{Y^{V\star}} \mathring{\mathbf{g}})_{ai} \end{aligned}$$

(note that $\mathring{\mathbf{g}}_{ij} \gamma^{jk} = r^2 \delta_i^k$). We therefore have

$$(4.22) \quad h^V = h^{V\circ} + \mathcal{L}_{Y^{V\star}} \mathring{\mathbf{g}}.$$

We continue with the scalar variations. We define

$$(4.23) \quad h_{ab}^{S\circ} = \sum_{I \notin \mathbb{I}_S^*} f_{ab,I}^S \mathbb{S}^I, \quad h_{ab}^{S\star} = \sum_{I \in \mathbb{I}_S^*} f_{ab,I}^S \mathbb{S}^I,$$

with obvious similar definitions for $h_{ai}^{S\star}$, etc.

The vanishing of the scalar master function implies [18, (2.7), (2.21) and (3.9)]

$$(4.24) \quad H_{L,I}^S + \frac{1}{n} H_{T,I}^S + \frac{1}{r} D^a r X_{a,I} = 0,$$

$$(4.25) \quad 0 = f_{ab,I}^S + (\mathcal{L}_{X_I} g)_{ab}.$$

where we have assumed that $k \neq 0$ and

$$(4.26) \quad X_{a,I} = \frac{r}{k} (f_{a,I}^S + \frac{r}{k} D_a H_{T,I}^S).$$

Using (4.24), the angular part of the variation can be written as

$$(4.27) \quad \begin{aligned} h_{ij}^{S\star} &= \sum_{I \in \mathbb{I}_S^*} 2r^2 \left(\frac{H_{T,I}^S}{k^2} \widehat{D}_i \widehat{D}_j \mathbb{S}^I - \gamma_{ij} \mathbb{S}^I \frac{1}{r} X(r) \right) \\ &= r^2 \mathcal{L}_{Y^{S\star}} \gamma_{ij} + 2r Y^{S\star}(r) \gamma_{ij} \\ &= \mathcal{L}_{Y^{S\star}}(r^2 \gamma_{ij}), \end{aligned}$$

where

$$(4.28) \quad Y^{S\star} := \sum_{I \in \mathbb{I}_S^*} \left(\frac{1}{k^2} H_{T,I}^S \gamma^{ij} \widehat{D}_j \mathbb{S}^I \partial_i - \mathbb{S}^I X^a \partial_a \right).$$

The convergence of all series above can be justified in a way very similar to that of Appendix K, and will be omitted.

Using (4.26) and (4.8) the mixed part is given by

$$(4.29) \quad \begin{aligned} h_{ai}^S &= \sum_{I \in \mathbb{I}_S^*} \widehat{D}_i \mathbb{S}^I \left(-X_{a,I} + \frac{r^2}{k^2} D_a H_{T,I}^S \right) \\ &= \widehat{D}_i Y_{S,a} + D_a Y_{S,i} = (\mathcal{L}_{Y^{S\star}} \mathring{g})_{ai}. \end{aligned}$$

The r, t part of the variation is finally, using (4.25),

$$(4.30) \quad h_{ab}^S = \sum_{I \in \mathbb{I}_S^*} f_{ab,I}^S \mathbb{S}^I = - \sum_{I \in \mathbb{I}_S^*} \mathbb{S}^I (\mathcal{L}_{Y^{S\star}} g)_{ab} = (\mathcal{L}_{Y^{S\star}} g)_{ab}.$$

In conclusion, we have

$$(4.31) \quad h = h^\diamond + \mathcal{L}_{Y^\star} \mathring{\mathfrak{g}},$$

with

$$(4.32) \quad \begin{aligned} Y^\star &= Y^{S^\star} + Y^{V^\star} \\ &= \sum_{I \in \mathbb{I}_S^\star} \left(\frac{1}{k^2} H_{T,I}^S \widehat{D}^i S^I \partial_i - S^I X_I^a \partial_a \right) - \sum_{I \in \mathbb{I}_V^\star} \frac{H_{T,I}^V}{k_V} \mathbb{V}^I. \end{aligned}$$

Using the estimates in Appendix D.1 we obtain for the asymptotics of Y^\star

$$(4.33) \quad \begin{aligned} |Y^\star|^2 &= \mathring{\mathfrak{g}}_{tt}((Y^\star)^t)^2 + \mathring{\mathfrak{g}}_{rr}((Y^\star)^r)^2 + r^2 |(Y^\star)^i|_\gamma^2 \\ &= O(r^{-2-2n+4}) + O(r^{2-2-2n}) + O(r^{2-2-2n}) \\ &= O(r^{2-2n}). \end{aligned}$$

5. Triviality of the kernel

We are ready now to pass to the proof of non-degeneracy:

Theorem 5.1. *Consider an $(n + 2)$ -dimensional Riemannian Kottler metric $\mathring{\mathfrak{g}}$ as in (2.1), with $\mu > 0$ for $K \in \{0, 1\}$ or μ satisfying (2.9) for $K = -1$, and an axis of rotation at $r = r_0 > 0$, as described in Section 2. Suppose that*

- 1) $K = 0, n \geq 2, \mu$ as in Proposition 3.10; or
- 2) $K = 1, n = 2, \mu \neq \mu_c$ given by (2.21); or
- 3) $K = -1, n = 2$.

Then $(\mathbb{R}^2 \times {}^n N, \mathring{\mathfrak{g}})$ is non-degenerate in the sense described in the introduction.

Proof. Let $h \in L^2$ be in the kernel of the operator P_L . Consider the decomposition of h into master scalar, vectorial, and tensor modes, together with their non-master counterparts:

$$(5.1) \quad h = \underbrace{h^{S^\star} + h^{S^\diamond}}_{h^S} + \underbrace{h^{V^\star} + h^{V^\diamond}}_{h^V} + \underbrace{h^{T^\star}}_{h^T},$$

(note that all tensorial modes are controlled by master functions).

Suppose, first, that $K = 0$. We show in Appendix H below that all angle-independent modes h^\diamond of h (in particular, all non-master modes) are pure gauge:

$$(5.2) \quad h^\diamond := h^{S^\diamond} + h^{V^\diamond} = \mathcal{L}_{Y^\diamond} \mathring{\mathfrak{g}}, \quad |Y^\diamond|_{\mathring{\mathfrak{g}}} = O(r^{-3}).$$

It follows from Proposition 3.10 together with the analysis in Section D.2 that, for $n \geq 2$, all master modes are likewise pure gauge for a nontrivial range of mass parameters μ (for all $\mu > 0$ when $n = 2$):

$$(5.3) \quad h^\star := h^{S^\star} + h^{V^\star} + h^{T^\star} = \mathcal{L}_{Y^\star} \mathring{\mathfrak{g}}, \quad |Y^\star|_{\mathring{\mathfrak{g}}} = O(r^{1-n}).$$

Thus

$$(5.4) \quad h = \mathcal{L}_Y \mathring{\mathfrak{g}}, \quad Y := Y^\star + Y^\diamond, \quad |Y|_{\mathring{\mathfrak{g}}} = O(r^{-1}).$$

Now, h in (4.31) is in TT -gauge, and the operator obtained by composing the divergence and the trace-free part of the Lie derivative is precisely the operator L^*L considered in [23, Proposition G]. Keeping in mind that our n is shifted by one as compared to the parameter n used in [23], from [23, Proposition G] we find that the indicial radius of L^*L is $R = \frac{n+3}{2}$. From [23, Theorem C(c)], choosing δ there so that the Sobolev space $C_\delta^{k,\alpha}$ contains only decaying fields, we see that any element Y in the L^2 -kernel of L^*L has to decay at infinity at a rate as close to the lower end of the indicial interval as desired. (In fact a more careful analysis shows that an element of the L^2 -kernel must decay as

$$(5.5) \quad |Y|_{\mathring{\mathfrak{g}}} = O(r^{-(\frac{n+1}{2}+R)}) = O(r^{-2}).)$$

Since the L^2 -kernel of L^*L is the same as the L^2 -kernel of L , one can invoke [1, Proposition 6.2.2] to conclude that the L^2 -kernel of L^*L is always trivial.

Next, (5.4) shows that for our gauge field Y it holds that $Y\rho^\delta$ is in L^2 for $\delta > \frac{3-n}{2}$; choosing $\delta \in (\frac{3-n}{2}, \frac{n+3}{2})$ we can thus use [23, Theorem C(b)] to conclude that $Y \equiv 0$. Hence

$$(5.6) \quad h \equiv 0,$$

which concludes the proof when $K = 0$.

The argument for $K = \pm 1$ is very similar: when $n = 2$ there are no non-master tensor modes; the fact that non-master scalar modes are pure gauge

is the contents of the linearised Birkhoff theorems of Appendices F and I, while the pure-gauge character of the $l = 1, K = 1$ modes is established in Appendix G. \square

Appendix A. Divergence of a symmetric two-tensor in coordinates

The object of this appendix is to show that the TT condition does not mix the scalar, vector, and tensor modes.

Let us consider a warped product metric of the following form (note that this allows for metrics more general than the metric \mathring{g} of the main body of the paper):

$$\mathcal{G} = g_{ab}(y)dy^a dy^b + r^2(y)\gamma_{ij}(x)dx^i dx^j,$$

where the y^a 's are local coordinates on a m -dimensional manifold and the x^i 's are local coordinates in n -dimensions. The non trivial Christoffel symbols of \mathcal{G} are:

$$\begin{aligned} \Gamma_{ab}^c &= \Gamma_{ab}^c(g), & \Gamma_{ij}^k &= \Gamma_{ij}^k(\gamma), \\ \Gamma_{ij}^c &= -g^{cb}r\partial_b r \gamma_{ij}, & \Gamma_{ib}^k &= r^{-1}\partial_b r \delta_i^k. \end{aligned}$$

The divergence of a 2-tensor is

$$\nabla_\alpha T^{\alpha\beta} = \partial_\alpha T^{\alpha\beta} + \Gamma_{\alpha\sigma}^\alpha T^{\sigma\beta} + \Gamma_{\alpha\sigma}^\beta T^{\alpha\sigma},$$

or equivalently

$$\nabla_\alpha T^{\alpha\beta} = (\sqrt{|\mathcal{G}|})^{-1}\partial_\alpha(\sqrt{|\mathcal{G}|}T^{\alpha\beta}) + \Gamma_{\alpha\sigma}^\beta T^{\alpha\sigma},$$

where

$$\sqrt{|\mathcal{G}|} = \sqrt{|g|}\sqrt{|\gamma|}r^n.$$

We deduce

$$\nabla_\alpha T^{\alpha\beta} = (\sqrt{|g|})^{-1}r^{-n}\partial_\alpha(r^n\sqrt{|g|}T^{a\beta}) + (\sqrt{|\gamma|})^{-1}\partial_i(\sqrt{|\gamma|}T^{i\beta}) + \Gamma_{\alpha\sigma}^\beta T^{\alpha\sigma}.$$

In particular

$$\begin{aligned} \nabla_\alpha T^{ab} &= (\sqrt{|g|})^{-1}r^{-n}\partial_\alpha(r^n\sqrt{|g|}T^{ab}) \\ &+ (\sqrt{|\gamma|})^{-1}\partial_i(\sqrt{|\gamma|}T^{ib}) + \Gamma_{ac}^b T^{ac} - g^{bc}r\partial_c r \gamma_{ij}T^{ij}, \end{aligned}$$

so

$$(A.1) \quad \nabla_\alpha T^{\alpha b} = r^{-n} D_a(r^n T^{ab}) + \widehat{D}_i(T^{ib}) - g^{bc} r \partial_c r \gamma_{ij} T^{ij},$$

where \widehat{D} is the γ -connection and D is the g -connection. Similarly

$$\begin{aligned} \nabla_\alpha T^{\alpha j} &= (\sqrt{|g|})^{-1} r^{-n} \partial_a(r^n \sqrt{|g|} T^{aj}) \\ &\quad + (\sqrt{|\gamma|})^{-1} \partial_i(\sqrt{|\gamma|} T^{ij}) + \Gamma_{kl}^j T^{kl} + r^{-1} \partial_b r T^{bj}, \end{aligned}$$

so

$$(A.2) \quad \nabla_\alpha T^{\alpha j} = r^{-n} D_a(r^n T^{aj}) + \widehat{D}_i(T^{ij}) + r^{-1} \partial_b r T^{bj}.$$

A.1. Divergence of a scalar variation

Let us write, as in [18, Equation (2.4)],

$$T^{ab} = \mathbb{S} f^{ab}, \quad T^{ai} = r^{-1} f^a \mathbb{S}^i, \quad T^{ij} = 2r^{-2} (H_L \mathbb{S} \gamma^{ij} + H_T \mathbb{S}^{ij}),$$

Note that here $\mathbb{S}^i = \gamma^{ij} \mathbb{S}_j$ but $T^{ai} = \mathring{\mathbf{g}}^{ij} T^a_j$. From (A.1) and (A.2)

$$\begin{aligned} \nabla_\alpha T^{\alpha a} &= \mathbb{S} r^{-n} D_b(r^n f^{ab}) + r^{-1} f^a \widehat{D}_i \mathbb{S}^i - g^{ac} \partial_c r \gamma_{ij} 2r^{-1} (H_L \mathbb{S} \gamma^{ij} + H_T \mathbb{S}^{ij}), \\ \nabla_\alpha T^{\alpha j} &= r^{-n} D_b(r^{n-1} f^b) \mathbb{S}^j + 2r^{-2} (H_L \widehat{D}^j \mathbb{S} + H_T \widehat{D}_i \mathbb{S}^{ij}) + r^{-2} \partial_b r f^b \mathbb{S}^j. \end{aligned}$$

Assume moreover that (see [18], Equations (2.2) and (2.6))

$$\Delta_\gamma \mathbb{S} = -k^2 \mathbb{S}, \quad \mathbb{S}_i = -\frac{1}{k} \partial_i \mathbb{S}, \quad \mathbb{S}_{ij} = \frac{1}{k^2} \widehat{D}_i \partial_j \mathbb{S} + \frac{1}{n} \gamma_{ij} \mathbb{S}.$$

We obtain

$$\begin{aligned} \nabla_\alpha T^{\alpha a} &= \mathbb{S} r^{-n} D_b(r^n f^{ab}) + r^{-1} f^a k \mathbb{S} - g^{ac} \partial_c r \gamma_{ij} 2r^{-1} (H_L \mathbb{S} \gamma^{ij} + H_T \mathbb{S}^{ij}) \\ &= \mathbb{S} [r^{-n} D_b(r^n f^{ab}) + r^{-1} f^a k - 2r^{-1} g^{ac} \partial_c r n H_L], \\ \nabla_\alpha T^{\alpha j} &= r^{-n} D_b(r^{n-1} f^b) \mathbb{S}^j + 2r^{-2} (H_L \widehat{D}^j \mathbb{S} + H_T \widehat{D}_i \mathbb{S}^{ij}) + r^{-2} \partial_b r f^b \mathbb{S}^j \\ &= \widehat{D}^j \mathbb{S} \left[-\frac{1}{k} r^{-n} D_b(r^{n-1} f^b) \right. \\ &\quad \left. + 2r^{-2} \left(H_L + H_T \left(-1 + \frac{(n-1)K}{k^2} + \frac{1}{n} \right) \right) - \frac{1}{k} r^{-2} \partial_b r f^b \right]. \end{aligned}$$

A.2. Divergence of a vector variation

If we have

$$T^{ab} = 0, \quad T^{ai} = r^{-1} f^a \mathbb{V}^i, \quad T^{ij} = 2r^{-2} H_T \mathbb{V}^{ij},$$

then from (A.1) and (A.2)

$$\begin{aligned} \nabla_\alpha T^{\alpha a} &= r^{-1} f^a \widehat{D}_i \mathbb{V}^i - g^{ac} \partial_c r \gamma_{ij} 2r^{-1} H_T \mathbb{V}^{ij}, \\ \nabla_\alpha T^{\alpha j} &= r^{-n} D_b (r^{n-1} f^b) \mathbb{V}^j + 2r^{-2} H_T \widehat{D}_i \mathbb{V}^{ij} + r^{-2} \partial_b r f^b \mathbb{V}^j. \end{aligned}$$

Assume moreover that (see [18], Equations (5.7a), (5.7b) and (5.9))

$$\Delta_\gamma \mathbb{V} = -k_V^2 \mathbb{V}, \quad \mathbb{V}_{ij} = -\frac{1}{2k_V} (\widehat{D}_i \mathbb{V}_j + \widehat{D}_j \mathbb{V}_i), \quad \widehat{D}_i \mathbb{V}^i = 0,$$

where $k_V \neq 0$, then

$$\begin{aligned} \nabla_\alpha T^{\alpha a} &= 0, \\ \nabla_\alpha T^{\alpha j} &= \mathbb{V}^j \left(r^{-n} D_b (r^{n-1} f^b) - \frac{1}{k_V} r^{-2} H_T [-k_V^2 + (n-1)K] + r^{-2} \partial_b r f^b \right). \end{aligned}$$

A.3. Divergence of a tensor variation

If we assume that

$$T^{ab} = 0, \quad T^{ai} = 0, \quad T^{ij} = 2r^{-2} H_T \mathbb{T}^{ij},$$

then from (A.1) and (A.2)

$$\begin{aligned} \nabla_\alpha T^{\alpha a} &= -g^{ac} \partial_c r \gamma_{ij} 2r^{-1} H_T \mathbb{T}^{ij}, \\ \nabla_\alpha T^{\alpha j} &= 2r^{-2} H_T \widehat{D}_i \mathbb{T}^{ij}. \end{aligned}$$

Assume moreover that (see [18] equation (5.1a) and (5.1b))

$$\gamma_{ij} \mathbb{T}^{ij} = 0, \quad \widehat{D}_i \mathbb{T}^{ij} = 0,$$

then

$$\nabla_\alpha T^{\alpha \mu} = 0.$$

Appendix B. Divergence & double divergence of h_{ij}^S and h_{aj}^S

The aim of this appendix is to derive some divergence identities, as implicitly used in Appendix C.

We start by calculating the divergence of \mathbb{S}_{jk}^I . Assuming $k \neq 0$ it holds that

$$\begin{aligned} \widehat{D}_i(\gamma^{ij}\mathbb{S}_{jk}^I) &= \frac{1}{k^2}\gamma^{ij}\widehat{D}_i\widehat{D}_j\widehat{D}_k\mathbb{S}^I + \frac{1}{n}\widehat{D}_k\mathbb{S}^I \\ &= -\widehat{D}_k\mathbb{S}^I + \frac{1}{k^2}\gamma^{\ell m}\widehat{R}_{mk}\widehat{D}_\ell\mathbb{S}^I + \frac{1}{n}\widehat{D}_k\mathbb{S}^I \\ &= \left(\frac{1}{n} - 1 + \frac{(n-1)K}{k^2}\right)\widehat{D}_k\mathbb{S}^I, \end{aligned}$$

where we have used $\widehat{R}_{ij} = (n-1)K\gamma_{ij}$ and

$$\nabla_i\nabla_j\nabla_k f = \nabla_k\nabla_j\nabla_i f + R^\ell{}_{jki}\partial_\ell f.$$

The double divergence of \mathbb{S}_{jk}^I is then

$$\widehat{D}_\ell\widehat{D}_i(\gamma^{\ell k}\gamma^{ij}\mathbb{S}_{jk}^I) = -\left(k^2\left(\frac{1}{n} - 1\right) + (n-1)K\right)\mathbb{S}^I.$$

For $k \neq 0$ this gives for the divergence and double divergence of $\delta\mathring{g}_{ij}^S$

$$\begin{aligned} \text{(B.1)} \quad \widehat{D}_i(\gamma^{ij}\delta\mathring{g}_{jk}^S) &= \sum_I 2r^2 \left[H_{L,S}^I \widehat{D}_k\mathbb{S}^I + H_{T,S}^I (\widehat{D}_i\gamma^{ij}\mathbb{S}_{jk}^I) \right] \\ &= \sum_I 2r^2 \widehat{D}_k\mathbb{S}^I \left[H_{L,S}^I + H_{T,S}^I \left(\frac{1}{n} - 1 + \frac{(n-1)K}{k^2} \right) \right], \end{aligned}$$

and

$$\begin{aligned} \text{(B.2)} \quad \widehat{D}_\ell\widehat{D}_i(\gamma^{\ell k}\gamma^{ij}\delta\mathring{g}_{jk}^S) &= \sum_I -2r^2 k^2 \mathbb{S}^I \left[H_{L,S}^I + H_{T,S}^I \left(\frac{1}{n} - 1 + \frac{(n-1)K}{k^2} \right) \right]. \end{aligned}$$

Similarly, for $\mathbb{V}_{\ell m}^I$, assuming $k_V \neq 0$,

$$\begin{aligned} \widehat{D}_j(\gamma^{\ell j}\mathbb{V}_{\ell m}^I) &= -\frac{1}{2k_V}\gamma^{\ell j}\left(\widehat{D}_j\widehat{D}_\ell\mathbb{V}_m^I + \widehat{D}_j\widehat{D}_m\mathbb{V}_\ell^I\right) \\ &= -\frac{1}{2k_V}\left(-k_V^2\mathbb{V}_m^I + \widehat{D}_m\gamma^{\ell j}\widehat{D}_j\mathbb{V}_\ell^I + \widehat{R}_{\ell jm}^j\gamma^{\ell i}\mathbb{V}_i^I\right) \\ &= \frac{1}{2}\mathbb{V}_m^I(k_V + (n-1)K), \end{aligned}$$

where we have used $\widehat{D}_j(\gamma^{jk}\mathbb{V}_k) = 0$ [18, (5.7b)], and

$$\widehat{D}_i\widehat{D}_j(\gamma^{mi}\gamma^{\ell j}\mathbb{V}_{\ell m}^I) = (k_V + (n-1)K)\widehat{D}_i(\gamma^{mi}\mathbb{V}_m) = 0.$$

Therefore

$$(B.3) \quad \widehat{D}_i(\gamma^{ij}\delta\hat{g}_{jk}^V) = \sum_I r^2 H_{T,V}^I \mathbb{V}_m^I (k_V + (n-1)K),$$

and

$$(B.4) \quad \widehat{D}_\ell\widehat{D}_i(\gamma^{\ell k}\gamma^{ij}\delta\hat{g}_{jk}^V) = 0.$$

Appendix C. The decomposition of h

In this appendix we wish to justify the decomposition (4.1) of h .

Recall that we denote by x^a the coordinates t and r , and by x^i the coordinates on the compact boundaryless Riemannian manifold (N, γ) . The metric functions h_{ab} form obviously a family of t - and r -dependent scalar functions on N , similarly for the γ -trace $\gamma^{ij}h_{ij}$ of $h_{\mu\nu}$. Next, the fields $h_{ai}dx^i$ form a family of t - and r -dependent covectors on N , while $h_{ij}dx^i dx^j$ forms a similar family of tensor fields on N . Removing the γ -trace of $h_{ij}dx^i dx^j$, one obtains a family of trace-free tensor fields on N .

We have the standard two L^2 -orthogonal decompositions for vector or covector fields, and for trace-free symmetric two-tensor fields (see e.g. [3]):

$$\begin{aligned} C^\infty(N, TN) &= \text{Im } d \oplus \ker d^*, \\ C^\infty(N, \mathring{S}_2 N) &= \text{Im } \mathring{\mathcal{L}} \oplus \ker \text{div}, \end{aligned}$$

where $\mathring{\mathcal{L}}$ is the conformal Killing operator,

$$\mathring{\mathcal{L}}(W)_{ij} = \widehat{D}_i W_j + \widehat{D}_j W_i - \frac{2}{n}\widehat{D}^k W_k \gamma_{ij}.$$

This allows to write any covector field uniquely as

$$W_i = W_i^S + W_i^V, \quad W^S \in \text{Im } d, \quad W^V \in \ker d^*.$$

This decomposition can be applied to the fields $h_{ai}dx^i$.

Next, any trace free symmetric tensor field $u_{ij}dx^i dx^j$ can be written in unique way as

$$u_{ij} = u_{ij}^S + u_{ij}^V + u_{ij}^T, \quad u^S \in \mathring{\mathcal{L}}(\text{Im } d), \quad u^V \in \mathring{\mathcal{L}}(\ker d^*), \quad u^T \in \ker \text{div}.$$

We apply this last decomposition to the γ -trace-free part of $h_{ij}dx^i dx^j$ which, together with what been said above, results in (4.1).

Note that (minus) the divergence of u is also decomposed as

$$\widehat{D}^j u_{ij} = \widehat{D}^j u_{ij}^S + \widehat{D}^j u_{ij}^V.$$

Next, if we use (compare Appendix B)

$$\begin{aligned} \text{div } \mathring{\mathcal{L}} &= \widehat{D}^* \widehat{D}_g - \text{Ric} + \frac{n-2}{2} dd^* \\ &= \Delta_{\text{Hodge}} - 2 \text{Ric} + \frac{n-2}{2} dd^* \\ &= d^* d + \frac{n}{2} dd^* - 2 \text{Ric}, \end{aligned}$$

and if the metric γ is Einstein, we have

$$\text{div } u^S \in \text{Im } d, \quad \text{div } u^V \in \ker d^*.$$

More precisely, if $\text{Ric} = K(n-1)$ and $u^S = \mathring{\mathcal{L}}dS$ then

$$\text{div } u^S = d \left[\frac{n}{2} (d^* dS) - 2K(n-1)S \right],$$

and if $u^V = \mathring{\mathcal{L}}V$, with $d^*V = 0$ then

$$\text{div } u^V = d^* dV - 2K(n-1)V.$$

Appendix D. The asymptotics of scalar modes

Whatever the value of k , for large r the potentials $V_{S,k}$ tend to

$$V_{S,k}(\infty) = \frac{(n-2)(n-4)}{4}.$$

We find the characteristic exponents of the master equation by solving the equation

$$s(1 - s) + V_{S,k}(\infty) = 0,$$

giving

$$s_{\pm} = \frac{1 \pm |n - 3|}{2}.$$

This has to be compared with the asymptotics

$$(D.1) \quad \Phi_{S,I} = O(\rho^{(n-2)/2}),$$

as obtained by directly translating the asymptotic behaviour of elements of the L^2 -kernel of the linearised Einstein operator into the asymptotic behaviour of the master fields. For $n \geq 4$ the decay (D.1) corresponds to s_+ . For $n = 3$ there is only one index, implying logarithmic terms in an asymptotic expansion. For $n = 2$ the decay (D.1) corresponds to s_- , which is zero, but we will improve this asymptotics below.

D.1. Estimate of $\Phi_{S,I}$ in any dimension n

From $|h|_{\mathfrak{g}} = O(\rho^{n+1})$ we obtain for the components in (t, r, z^j) coordinates ($j = 2, \dots, n + 1$)

$$(D.2) \quad \begin{aligned} h_{tt} &= O(r^{1-n}), & h_{tr} &= O(r^{-1-n}), & h_{rr} &= O(r^{-3-n}), \\ h_{tj} &= O(r^{1-n}), & h_{rj} &= O(r^{-1-n}), & h_{jk} &= O(r^{1-n}). \end{aligned}$$

By (4.2)

$$\begin{aligned} f_{tt,I}^S &= O(r^{1-n}), & f_{tr,I}^S &= O(r^{-1-n}), & f_{rr,I}^S &= O(r^{-3-n}), \\ f_{t,I}^S &= O(r^{-n}), & f_{r,I}^S &= O(r^{-2-n}), & H_{L/T,I}^S &= O(r^{-1-n}). \end{aligned}$$

Then we have for $X_{a,I}$, defined in (4.26), assuming $k \neq 0$,

$$\begin{aligned} X_{t,I} &= O(r^{1-n}), \\ X_{r,I} &= \frac{r}{k}(f_r + \frac{r}{k}D_r H_T) \\ &= \frac{r}{k} \left(O(r^{-2-n}) + \frac{r}{k}O(r^{-2-n}) \right) = O(r^{-n}), \end{aligned}$$

and [18, Equation (2.7)]

$$\begin{aligned}
 F_I &:= H_{L,I}^S + \frac{H_{T,I}^S}{n} + r^{-1}D^a r X_{a,I} \\
 &= O(r^{-1-n}) + r^{-1}g^{rr} X_{r,I} \\
 &= O(r^{-1-n}) + O(r^{1-n}) = O(r^{1-n}), \\
 F_{ab,I} &:= f_{ab,I}^S + D_a X_{b,I} + D_b X_{a,I}, \\
 F_{tt,I} &= O(r^{1-n}), \quad F_{tr,I} = O(r^{-n}), \quad F_{rr,I} = O(r^{-n-1}).
 \end{aligned}$$

The rescaled quantities, defined as [18, (2.13)]

$$\tilde{F}_I := r^{n-2}F_I, \quad \tilde{F}_{ab,I} := r^{n-2}F_{ab,I},$$

are

$$\tilde{F}_I = O(r^{-1}), \quad \tilde{F}_{tt,I} = O(r^{-1}), \quad \tilde{F}_{tr,I} = O(r^{-2}), \quad \tilde{F}_{rr,I} = O(r^{-3}).$$

Then [18, (2.20)]

$$\begin{aligned}
 X_I &:= \tilde{F}_{t,I}^t - 2\tilde{F}_I = O(r^{-1}), \quad Y_I := \tilde{F}_{r,I}^r - 2\tilde{F}_I = O(r^{-1}), \\
 Z_I &:= \tilde{F}_{t,I}^r = O(1),
 \end{aligned}$$

and finally [18, (3.1)]

$$\Phi_{S,I} := H_I^{-1}r^{1-n/2}(n\tilde{Z}_I - r(X_I + Y_I)) = O(r^{1-n/2}) = O(\rho^{n/2-1}),$$

where

$$\begin{aligned}
 H_I &:= m + xn(n+1)/2 = O(1), \\
 m &:= k(I)^2 - nK, \quad x := 2\mu r^{1-n},
 \end{aligned}$$

and \tilde{Z} is of the same order as Z .

D.2. Estimate of Φ_S in dimension $n = 2$

We revisit the preceding equations when $n = 2$. From [18, (2.11),(2.13)] we see that

$$\tilde{F} = F, \quad F^a_a = 0.$$

Let us write

$$H_T = h_t r^{-3} + O(r^{-3-\varepsilon}), \quad D_r H_T = -3h_t r^{-4} + O(r^{-4-\varepsilon}),$$

where $\varepsilon > 0$. From the estimates in Section D.1,

$$\begin{aligned} X_r &= -3\frac{1}{k^2}h_t r^{-2} + O(r^{-2-\varepsilon}), & F &= -3\frac{1}{k^2}h_t r^{-1} + O(r^{-1-\varepsilon}) \\ F_{tt} &= O(r^{-1}), & F_{tr} &= O(r^{-2}), \end{aligned}$$

and

$$F_{rr} = 2D_r(r^2 D_r H_T) + O(r^{-3-\varepsilon}) = 12h_t r^{-3} + O(r^{-3-\varepsilon}).$$

From $F^a_a = 0$ we see that $h_t = 0$, so F and then $X + Y$ decay faster. We deduce from the definitions of X and Y that there exists $\varepsilon > 0$ such that

$$F, X, Y = O(r^{-1-\varepsilon}).$$

This information inserted into [18, (2.24d)] gives

$$\tilde{Z} = O(r^{-\varepsilon}).$$

In conclusion

$$\Phi_{S,I} = O(r^{-\varepsilon}),$$

tends to zero at infinity.

Appendix E. The asymptotics of vector modes

In order to estimate the rate of decay of $\Phi_{V,I}$ we start with [18, (5.10)]

$$\begin{aligned} F_r &= f_r + \frac{r}{k_V} D_r H_T = O(r^{-2-n}) + O(r^{-1-n}) = O(r^{-1-n}), \\ F_t &= f_t + \frac{r}{k_V} D_t H_T = O(r^{-n}). \end{aligned}$$

From [18, (5.12)] we have

$$r^{n-1} F_a = \varepsilon_a^b D_b \Omega,$$

where $\varepsilon_{ab} = O(1)$. Therefore

$$D_t \Omega = O(1), \quad D_r \Omega = O(r^{-3}),$$

and $\Omega = O(1)$. By [18, (5.13)] we have

$$\Phi_{V,I} = r^{-n/2} \Omega = O(r^{-n/2}) = O(\rho^{n/2}).$$

Appendix F. The linearised Birkhoff theorem

In the case $K = 1$, $n = 2$, a gauge transformation $h_{\mu\nu} \rightarrow h_{\mu\nu} + \mathcal{L}_Y \mathring{\mathfrak{g}}_{\mu\nu}$, with gauge vector Y , takes the form

$$(F.1) \quad h_{tt} \rightarrow h_{tt} + Y^r \partial_r f + 2f \partial_t Y^t,$$

$$(F.2) \quad h_{tr} \rightarrow h_{tr} + f^{-1} \partial_t Y^r + f \partial_r Y^t,$$

$$(F.3) \quad h_{rr} \rightarrow h_{rr} + Y^r \partial_r f^{-1} + 2f^{-1} \partial_r Y^r,$$

$$(F.4) \quad h_{\theta\varphi} \rightarrow h_{\theta\varphi} + r^2 \sin^2 \theta \partial_\theta Y^\varphi + r^2 \partial_\varphi Y^\theta,$$

$$(F.5) \quad h_{\varphi\varphi} \rightarrow h_{\varphi\varphi} + 2r^2 \sin^2 \theta (r^{-1} Y^r + Y^\theta \cot \theta + \partial_\varphi Y^\varphi),$$

$$(F.6) \quad h_{\theta\theta} \rightarrow h_{\theta\theta} + 2r Y^r + 2r^2 \partial_\theta Y^\theta,$$

$$(F.7) \quad h_{t\theta} \rightarrow h_{t\theta} + f \partial_\theta Y^t + r^2 \partial_t Y^\theta,$$

$$(F.8) \quad h_{t\varphi} \rightarrow h_{t\varphi} + f \partial_\varphi Y^t + r^2 \sin^2 \theta \partial_t Y^\varphi,$$

$$(F.9) \quad h_{r\theta} \rightarrow h_{r\theta} + f^{-1} \partial_\theta Y^r + r^2 \partial_r Y^\theta,$$

$$(F.10) \quad h_{r\varphi} \rightarrow h_{r\varphi} + f^{-1} \partial_\varphi Y^r + r^2 \sin^2 \theta \partial_r Y^\varphi.$$

We consider an $l = 0$ linearised solution $h_{\mu\nu}$ of the Einstein equations, i.e.

$$(F.11) \quad h_{ab} = h_{ab}(t, r), \quad h_{ia} \equiv 0, \quad h_{ij} = \psi(t, r) \mathring{\mathfrak{g}}_{ij}.$$

We assume that $h \in L^2$ and is in the kernel of the operator P_L . We set

$$(F.12) \quad Y^r = r\psi/2 = O(r^{-2}), \quad Y^i \equiv 0$$

(recall that $O(r^{-2})$ is understood for large r), which implies

$$h_{ij} = \mathcal{L}_Y \mathring{\mathfrak{g}}_{ij}.$$

Recall that $r_0 > 0$ has been defined as the largest zero of f . Let $\varepsilon > 0$. We define Y_ε^t by integrating (F.2) in r and cutting-off near r_0 :

$$(F.13) \quad Y_\varepsilon^t = -\chi_\varepsilon(r) \int_r^\infty f^{-1} (h_{rt} - \partial_t Y^r f^{-1}) dr = O(r^{-4}),$$

where χ_ε is a smooth function which is identically equal to one for $r \geq r_0 + \varepsilon$ and zero for $r \in [r_0, r_0 + \frac{1}{2}\varepsilon]$. (The vector field defined by the integral above without the cut-off might be singular at $r = r_0$; we will see at the end of the argument that the cut-off was unnecessary, but this is not clear from the outset. Note that if $\varepsilon_1 > \varepsilon_2$ then $Y_{\varepsilon_1}(r) = Y_{\varepsilon_2}(r)$ for $r > r_0 + \varepsilon_1$.)

Then, for $r > r_0 + \varepsilon$,

$$h_{tr} = \mathcal{L}_{Y_\varepsilon \mathring{\mathfrak{g}}}.$$

An analysis of the components of $\mathcal{L}_{Y_\varepsilon \mathring{\mathfrak{g}}}$, using (F.1)–(F.10), proves that

$$|\mathcal{L}_{Y_\varepsilon \mathring{\mathfrak{g}}}|_{\mathring{\mathfrak{g}}}^2 = O(r^{-6}),$$

so $\mathcal{L}_{Y_\varepsilon \mathring{\mathfrak{g}}} \in L^2$.

Set

$$(F.14) \quad \widehat{h}_{\mu\nu} = h_{\mu\nu} - \mathcal{L}_{Y_\varepsilon \mathring{\mathfrak{g}}}_{\mu\nu},$$

thus $\widehat{h}_{\mu\nu}$ is a solution of the linearised Einstein equations with all components vanishing for large r except possibly h_{tt} and h_{rr} . Now, a variation of the metric arising from a variation $\delta\mu$ of the mass in the coordinate system (2.1) takes the form

$$(F.15) \quad 2\frac{\delta\mu}{r}(-dt^2 + f^{-2}dr^2),$$

which suggests that it might be convenient to define new functions \widetilde{h}_{rr} and \widetilde{h}_{tt} as

$$(F.16) \quad \widetilde{h}_{rr} := r f^2 \widehat{h}_{rr}, \quad \widetilde{h}_{tt} := r(\widehat{h}_{tt} + f^2 \widehat{h}_{rr}).$$

Inserting (F.16) into the linearised Einstein tensor $G'[h]_{\mu\nu}$ one finds, again for $r > r_0 + \varepsilon$,

$$(F.17) \quad G'_{tr}[h] = \frac{\partial_t \widetilde{h}_{rr}}{r(-2\mu + r^3 + r)},$$

thus \widetilde{h}_{rr} depends at most upon r . One can now eliminate the second radial derivative of \widetilde{h}_{tt} between the G_{tt} and G_{rr} equations, obtaining

$$(F.18) \quad \partial_r \left(\frac{\widetilde{h}_{tt}}{rf} \right) = 0.$$

Hence,

$$(F.19) \quad \widetilde{h}_{tt} = C(t)rf.$$

for $r > r_0 + \varepsilon$, for some function C depending only upon t . Inserting all this into the equations $G_{ij} = 0$ gives $\partial_r \widetilde{h}_{rr} = 0$, and thus \widetilde{h}_{rr} is a constant, say $2\delta\mu$.

In terms of \widehat{h}_{rr} and \widehat{h}_{tt} we now have, for $r > r_0 + \varepsilon$,

$$(F.20) \quad \widehat{h}_{tt} = fC(t) - \frac{2\delta\mu}{r}, \quad \widehat{h}_{rr} = \frac{2\delta\mu}{rf^2}.$$

This is not in L^2 unless $C(t) \equiv 0$. Further, our gauge-transformed field with $C(t) = 0$ is a variation of the mass, and therefore, as shown in Section 2, it is in L^2 if and only if the mass takes the critical value.

For all masses except the critical one the tensor field h is therefore pure gauge for $r > r_0 + \varepsilon$. Since ε is arbitrary, we conclude that for $r > r_0$ we have

$$(F.21) \quad h_{\mu\nu} = \mathcal{L}_{Y_0} \mathring{\mathfrak{g}}_{\mu\nu},$$

where Y_0 is defined as in (F.13) with $\chi_0 \equiv 1$. As h has been assumed to be smooth, it is simple to show from (F.21) that Y_0 is smooth across the rotation axis, so that h is pure gauge everywhere, as desired.

If we denote by $h^{S\circ}$ the $l = 0$ -part of h , and by $Y^{S\circ}$ the gauge vector field Y_0 just constructed, we have, for $\mu \neq \mu_c$,

$$(F.22) \quad h^{S\circ} = \mathcal{L}_{Y^{S\circ}} \mathring{\mathfrak{g}}, \quad |Y^{S\circ}|_{\mathring{\mathfrak{g}}} = O(r^{-3}).$$

Appendix G. The $l = 1$ modes for $K = 1$, $n = 2$

In this section we analyze the $l = 1$ modes when $({}^nN, \gamma_{ij})$ is a two-dimensional round unit sphere. We follow the treatment of the Lorentzian case in [11], which requires only trivial modifications when addressing the Riemannian setting. We present the argument here because of the need of establishing estimates for the gauge vector fields.

As explained in [13, end of Section 2.2] (compare [11, Section II.A]), in the case at hand a general metric perturbation can be decomposed as $h_{\mu\nu} = h_{\mu\nu}^{(-)} + h_{\mu\nu}^{(+)}$, where $h_{\mu\nu}^{(-)}$ is the *odd* and $h_{\mu\nu}^{(+)}$ the *even* part. The linearised Einstein equations decouple into separate equations for the odd and even parts.

G.1. Odd perturbations

Odd perturbations take the form [11, Equation (35)]

$$(G.1) \quad h_{\mu\nu}^{(-)} = \begin{pmatrix} 0 & r^2 h_a^{(l,m)} \widehat{\varepsilon}_i^j \widehat{\nabla}_j S_{(l,m)} \\ r^2 h_a^{(l,m)} \widehat{\varepsilon}_i^j \widehat{\nabla}_j S_{(l,m)} & 2r^4 k^{(l,m)} \widehat{\varepsilon}_{(i}{}^m \widehat{\nabla}_{j)} \widehat{\nabla}_m S_{(l,m)} \end{pmatrix},$$

where $h_a^{(l,m)}$ and $k^{(l,m)}$ are functions of t and r , $\widehat{\nabla}$ is the covariant derivative on the sphere, $\widehat{\varepsilon}_{ij} = \sin\theta(\delta_i^\theta\delta_j^\phi - \delta_i^\phi\delta_j^\theta)$ and $S_{(l,m)}$ are the scalar spherical harmonics.

Restricting to $l = 1$ modes, the ij components of $h_{\mu\nu}^{(-)}$ vanish and the ai components take the form

$$(G.2) \quad h_{ai}^{(-)} = \sum_{m=-1}^1 \sqrt{\frac{3}{4\pi}} h_a^{(l=1,m)} J_{(m)i}$$

where the $J_{(m)}$'s form a basis of Killing vector fields on S^2 .

Gauge transformations defined by a gauge vector Y of the form [11, (28)]

$$Y^a = 0, \quad Y^i = r^4 \widehat{\varepsilon}^{ij} \widehat{\nabla}_j f_Y,$$

for some function f_Y , preserve the odd character of the perturbation and those with

$$f_Y = \sum_{m=-1}^1 f_Y^{(m)} S_{(l=1,m)},$$

stay within the $l = 1$ modes. The effect of such a gauge transformation on the perturbation is given by [11, (73)]

$$h_{ai}^{(-)} \rightarrow \sum_{m=-1}^1 \sqrt{\frac{3}{4\pi}} (h_a^{(l=1,m)} + r^2 \partial_a f_Y^{(m)}) r^{-2} J_{(m)i},$$

with all other components unaffected.

Defining $\widehat{h}^{(-)}$ by $h_{\mu\nu}^{(-)} = \widehat{h}_{\mu\nu}^{(-)} + \mathcal{L}_{Y^{\diamond-}} \mathring{\mathfrak{g}}$ with a gauge vector $Y^{\diamond-}$ defined as $(Y^{\diamond-})^a := 0$ and

$$(G.3) \quad (Y^{\diamond-})^i = -r^4 \widehat{\varepsilon}^{ij} \partial_j \left(\sum_{m=-1}^1 S_{(l=1,m)} \int h_r^{(l=1,m)} r^{-2} dr \right) = O(r^{-4}),$$

the components $\widehat{h}_{ri}^{(-)}$ vanish, leaving only $\widehat{h}_{ti}^{(-)}$. The norm of the gauge part is found to be

$$|\mathcal{L}_{Y^{\diamond-}} \mathring{\mathfrak{g}}|_{\mathring{\mathfrak{g}}}^2 = O(r^{-6}),$$

and, as $f_Y^{(m)}$ is regular at r_0 and therefore $f^{-1} \partial_t f_Y^{(m)}$ is as well, $\mathcal{L}_{Y^{\diamond-}} \mathring{\mathfrak{g}} \in L^2$.

As the $\widehat{h}_{ti}^{(-)}$ -components are of the form (G.2), we can set one of them to zero by a rotation, leaving $\widehat{h}_{t\varphi}^{(-)} = \sqrt{\frac{3}{4\pi}} \widehat{h}_t^{(l=1,0)} \sin^2\theta$, for some functions $\widehat{h}_t^{(l=1,0)}(t, r)$, as the only non-zero component of the perturbation.

The linearised Einstein equations give

$$(G.4) \quad r^2 \partial_r^2 \widehat{h}_t^{(l=1,0)} - 2\widehat{h}_t^{(l=1,0)} = 0,$$

$$(G.5) \quad 2\partial_t \widehat{h}_t^{(l=1,0)} - r\partial_r \partial_t \widehat{h}_t^{(l=1,0)} = 0.$$

Integrating (G.5) twice, we obtain

$$\widehat{h}_t^{(l=1,0)} = C_1 r^2 t + g(r),$$

where C_1 has to vanish because of the periodicity of the t coordinate. Inserting this into (G.4) leads to

$$\widehat{h}_t^{(l=1,0)} = C_2 r^2 + C_3 r^{-1}.$$

As the tensors $dt dx^i$ are not smooth at the axis of rotation $r = r_0$ we require $C_2 r_0^2 + C_3 r_0^{-1} = 0$, i.e.

$$\widehat{h}_t^{(l=1,0)} = C_2 \frac{r^3 - r_0^3}{r}.$$

Perturbations of this form are exactly the variations of angular momentum within the Riemannian Kerr anti-de Sitter family which are analyzed in Section J. As these variations are not in L^2 we have $\widehat{h} = 0$, and if we denote by $h^{\diamond 1-}$ the odd part of the $l = 1$ modes, we obtain

$$(G.6) \quad h^{\diamond 1-} = \mathcal{L}_{Y^{\diamond-} \mathring{g}}, \quad |Y^{\diamond-}|_{\mathring{g}} = O(r^{-3}).$$

G.2. Even perturbations

Even $l = 1$ solutions of the linearised Einstein equations can be parameterised as [11, Equation (36)]

$$(G.7) \quad (h_{\alpha\beta}^{(+)}) = \begin{pmatrix} h_{ab}^{(\ell=1)} & \widehat{D}_i q_a^{(\ell=1)} \\ \widehat{D}_i q_b^{(\ell=1)} & \frac{r^2}{2} J^{(\ell=1)} \gamma_{ij} \end{pmatrix}.$$

Under gauge transformations with gauge-vector Y of the form

$$(G.8) \quad (Y_\alpha) = (Y_a, r^2 \widehat{D}_i X),$$

$(h_{\alpha\beta}^{(+)})$ transforms to $(\widehat{h}_{\alpha\beta}^{(+)})$ given by [11, Equation (122)]

$$(G.9) \quad \begin{pmatrix} h_{ab}^{(\ell=1)} + \tilde{D}_a Y_b^{(\ell=1)} + \tilde{D}_b Y_a^{(\ell=1)} & \widehat{D}_i (q_a^{(\ell=1)} + r^2 \tilde{D}_a X^{(\ell=1)} + Y_a^{(\ell=1)}) \\ \widehat{D}_i (q_b^{(\ell=1)} + r^2 \tilde{D}_b X^{(\ell=1)} + Y_b^{(\ell=1)}) & \frac{r^2}{2} \gamma_{ij} (J^{(\ell=1)} - 4X^{(\ell=1)} + \frac{4}{r} Y_{(\ell=1)}^a \tilde{D}_{ar}) \end{pmatrix}.$$

According to [11, Section IV.A.2], it is convenient to define (X, Y_a) by solving the following system of equations:

$$(G.10) \quad r^2 D_t X^{(\ell=1)} + Y_t = -q_t^{(\ell=1)} = O(r^{-1}),$$

$$(G.11) \quad r^2 D_r X^{(\ell=1)} + Y_r = -q_r^{(\ell=1)} = O(r^{-3}),$$

$$(G.12) \quad D^b Y_b = -\frac{1}{2} \mathring{\mathfrak{g}}^{ab} h_{ab}^{(\ell=1)} = O(r^{-3}).$$

With this choice, $\widehat{h}^{(+)}$ satisfies

$$(G.13) \quad \widehat{h}_{ai}^{(+)} = 0, \quad \mathring{\mathfrak{g}}^{ab} \widehat{h}_{ab}^{(+)} = 0.$$

Note that (G.10)–(G.12) imply

$$(G.14) \quad D^b (r^2 D_b X) = \frac{1}{2} \mathring{\mathfrak{g}}^{ab} h_{ab} - D^b q_b.$$

The homogeneous version of the equation (G.14) for X has no non-trivial solutions tending to zero at infinity by the maximum principle. The operator at the left-hand side of (G.14) has indicial exponents in $\{0, -3\}$, and therefore (G.14) has a unique solution $X = O(r^{-3})$ which is a linear combination of $l = 1$ spherical harmonics.

The conditions (G.13) do not fix the gauge uniquely: an additional gauge transformation satisfying

$$(G.15) \quad r^2 D_a X + Y_a = 0, \quad D^a Y_a = 0,$$

preserves the form of $\widehat{h}^{(+)}$.

By a rotation we set $J = J^{(1)} S_{(l=1, m=1)}$. We define new variables C_a as

$$(G.16) \quad \begin{aligned} \widehat{h}_{ab}^{(\ell=1, T)} &= \frac{1}{f} \left[C_a^{(\ell=1)} D_b r + C_b^{(\ell=1)} D_a r - \tilde{g}_{ab} C_d^{(\ell=1)} D^d r \right], \\ C_a^{(\ell=1)} &= \widehat{h}_{ab}^{(\ell=1, T)} D^b r, \end{aligned}$$

and decompose them into modes

$$(G.17) \quad C_a = \sum_{m=1}^3 C_a^{(m)} S_{(l=1, m)}.$$

Eliminating C_r between the $m = 2, 3$ components of $G'_{tr}[h]$ and $G'_{t\phi}[h]$ we obtain

$$(G.18) \quad \partial_r C_t^{(2,3)} + \frac{C_t^{(2,3)}}{r - 2\mu + r^3} = 0,$$

and therefore, by integration from the axis of rotation at r_0 , the $C_t^{(2,3)}$ depend only on r . The $m = 2, 3$ components of $G'_{t\phi}[h]$ show $\partial_t C_r^{(2,3)} = -\partial_r C_t^{(2,3)}$ which implies that $C_t^{(2,3)}$ vanishes and $C_r^{(2,3)}$ can only depend on r . Eliminating second derivatives between $G'_{tt}[h]$ and $G'_{rr}[h]$ shows that $C_r^{(2,3)}$ vanishes as well.

To handle the $m = 1$ equations we define new variables $Z_r^{(1)}(t, r)$ and $Z_t^{(1)}(t, r)$ by

$$(G.19) \quad \begin{aligned} C_r^{(1)} &= 6\mu \partial_r Z_r^{(1)} + \frac{r}{2} \partial_r J^{(1)}, \\ C_t^{(1)} &= Z_t^{(1)} - \frac{3\mu - r}{2} \partial_t J^{(1)} - r f 6\mu \partial_t \partial_r Z_r^{(1)}. \end{aligned}$$

Note that this defines $Z_r^{(1)}$ only up to an irrelevant constant.

The $m = 1$ component of $G'_{tr}[h]$ directly gives $Z_t^{(1)} = 0$. Eliminating third order derivatives from the remaining equations we obtain

$$(G.20) \quad \partial_r J^{(1)} + 4r f \partial_r^2 Z_r^{(1)} + (12r^2 + 8) \partial_r Z_r^{(1)} = 0.$$

Differentiating the $m = 1$ equations by r and using (G.20) to express derivatives of J by $Z_r^{(1)}$ gives two fifth order and one fourth order equation for $Z_r^{(1)}$. Eliminating higher derivatives we finally obtain a third order equation for $Z_r^{(1)}$

$$(G.21) \quad r^2 f \partial_r D^a (r^2 D_a Z_r^{(1)}) + r^4 \partial_r (r^{-2} f) D^a (r^2 D_a Z_r^{(1)}) = 0.$$

This implies

$$(G.22) \quad D^a (r^2 D_a Z_r^{(1)}) = \frac{C r^2}{f},$$

with a constant C which has to vanish for $Z_r^{(1)}$ to be regular at r_0 .

We now consider the remaining gauge freedom. We see from (G.15) that for any X satisfying

$$(G.23) \quad D^a(r^2 D_a X) = 0$$

there exists an associated Y_a giving a gauge transformation which preserves (G.13).

Inserting the definition of our new variables into (G.9) we find that under a gauge transformation satisfying (G.15) the fields $Z_r^{(1)}$ and $J^{(1)}$ transform as

$$(G.24) \quad \partial_r Z_r^{(1)} \mapsto \partial_r(Z_r^{(1)} + X^{(1)}),$$

$$(G.25) \quad J^{(1)} \mapsto J^{(1)} - 4X^{(1)} + \frac{4f}{r} Y_r^{(1)},$$

where X and Y_a have been split into $l = 1$ modes as for C_a in (G.17).

As $Z_r^{(1)}$ satisfies (G.23) we can render it independent of r by a gauge transformation with $\partial_r X^{(1)} = -\partial_r Z_r^{(1)}$, which preserves (G.13).

With $\partial_r Z_r^{(1)} \equiv 0$ we see from (G.20) that $J^{(1)}$ can only depend on t . From the remaining equations $\partial_t^2 J^{(1)} = 0$, i.e. $J^{(1)}$ is constant.

We can exploit the remaining freedom in $X^{(1)}$ to set

$$(G.26) \quad X^{(1)} = \frac{J^{(1)}}{4}, \quad Y_a = 0,$$

obtaining $J^{(1)} \equiv 0$. This gives $Z_r^{(1)} = \text{const}$ and therefore $C_r \equiv C_t \equiv 0$.

We arrive at $h^{(+)} = \mathcal{L}_{Y^{\diamond+}} \mathring{\mathfrak{g}}$ where $Y^{\diamond+}$ is the combined gauge vector consisting of the part defined by (G.10)–(G.12), that given by (G.24) and that given by (G.26). From the asymptotics (D.2) of h and from (F.1)–(F.10) with the right-hand sides being zero now, we conclude that

$$(G.27) \quad h^{\diamond 1+} \equiv h^{(+)} = \mathcal{L}_{Y^{\diamond+}} \mathring{\mathfrak{g}}, \quad |Y^{\diamond+}|_{\mathring{\mathfrak{g}}} = O(r^{-1}).$$

An alternative, and somewhat simpler, proof can be given using [15]. Since the last reference was only available in Polish at the time of writing of this paper [29], we felt it more appropriate to provide the argument above.

Appendix H. A linearised Birkhoff-type theorem with $K = 0$

In the case $K = 0$ the gauge transformations take the form

$$\begin{aligned}
 \text{(H.1)} \quad h_{tt} &\rightarrow h_{tt} + Y^r \partial_r f + 2f \partial_t Y^t, \\
 \text{(H.2)} \quad h_{tr} &\rightarrow h_{tr} + f^{-1} \partial_t Y^r + f \partial_r Y^t, \\
 \text{(H.3)} \quad h_{rr} &\rightarrow h_{rr} + Y^r \partial_r f^{-1} + 2f^{-1} \partial_r Y^r, \\
 \text{(H.4)} \quad h_{ij} &\rightarrow h_{ij} + \delta_{ij} 2r Y^r + r^2 (\partial_i Y^j + \partial_j Y^i), \\
 \text{(H.5)} \quad h_{ti} &\rightarrow h_{ti} + f \partial_i Y^t + r^2 \partial_t Y^i, \\
 \text{(H.6)} \quad h_{ri} &\rightarrow h_{ri} + f^{-1} \partial_i Y^r + r^2 \partial_r Y^i.
 \end{aligned}$$

We consider a $k = 0 = k_V = k_T$ linearised solution of the Einstein equations, i.e. one that only depends on t and r :

$$\text{(H.7)} \quad h_{\mu\nu} = h_{\mu\nu}(t, r).$$

As there are (constant) harmonic vectors on the background with $K = 0$ we have to consider both scalar and vector perturbations (the tensor ones are always controlled by the master functions). Scalar perturbations can be treated analogously to the $K = 1$ case in Appendix F: They take the form (F.11) and by choosing a gauge vector Y as in (F.12) and (F.13), we can set, for $r > r_0 + \varepsilon$, $\widehat{h}_{ij} \equiv 0$ and $\widehat{h}_{tr} \equiv 0$ where $\widehat{h}_{\mu\nu} = h_{\mu\nu} - \mathcal{L}_Y \mathfrak{g}_{\mu\nu}$ and $\varepsilon > 0$ is an arbitrary cutoff distance.

We define \tilde{h}_{rr} , \tilde{h}_{tt} as

$$\text{(H.8)} \quad \tilde{h}_{rr} := r^{n-1} f^2 \widehat{h}_{rr}, \quad \tilde{h}_{tt} := r^{n-1} (\widehat{h}_{tt} + f^2 \widehat{h}_{rr}),$$

and insert into the linearised Einstein tensor $G'_{\mu\nu}[\widehat{h}]$ using the equations in [20, Appendix B]. For $r > r_0 + \varepsilon$ the $G'_{tt}[h]$, $G'_{tr}[h]$, $G'_{rr}[h]$ components lead to

$$\text{(H.9)} \quad \tilde{h}_{rr} = \tilde{h}_{rr}(r),$$

$$\text{(H.10)} \quad \tilde{h}_{tt} = C(t) r f$$

where C has to vanish for h to be in L^2 . Inserting this into $G'_{ij}[h]$ gives $\tilde{h}_{rr} = 2\delta\mu$, for some constant $\delta\mu$.

The only remaining perturbations, up to gauge, are variations of the mass. As these are never in L^2 for $K = 0$, the scalar part of the perturbation has to be pure gauge.

We now consider the vector part, which takes the form

$$(H.11) \quad h_{ab} = 0, \quad h_{ai} = h_{ai}(r), \quad h_{ij} = 0.$$

We choose a gauge vector, which we denote by Y^\diamond , such that $Y^{\diamond a} \equiv 0$ and $Y^{\diamond i} = Y^{\diamond i}(t, r)$. We define $(Y^\diamond)^i$ by integrating (H.6) to obtain $h_{ri} = \mathcal{L}_Y \mathring{\mathfrak{g}}_{ri}$:

$$(H.12) \quad (Y^\diamond)^i = \int r^{-2} (h_{ri} - f^{-1} \partial_i (Y^\diamond)^r) dr = \int r^{-2} h_{ri} dr = O(r^{-n-2}).$$

Equations (H.1)–(H.6) show that

$$|\mathcal{L}_{Y^\diamond} \mathring{\mathfrak{g}}|_{\mathring{\mathfrak{g}}}^2 = O(r^{-2n-2}),$$

and, as it is regular at r_0 , $\mathcal{L}_{Y^\diamond} \mathring{\mathfrak{g}} \in L^2$.

The nontrivial linearised Einstein equations turn out to be

$$(H.13) \quad r^2 \partial_r^2 h_{ti} + r(n-2) \partial_r h_{ti} - 2(n-1) h_{ti} = 0,$$

$$(H.14) \quad 2 \partial_t \widehat{h}_{ti} - r \partial_r \partial_t \widehat{h}_{ti} = 0.$$

Integrating (H.14) twice gives

$$\widehat{h}_{ti} = C_1 r^2 t + g(r),$$

where C_1 has to vanish because of the periodicity of the t coordinate. Inserting this into (H.13) leads to

$$\widehat{h}_{ti} = C_2 r^2 + C_3 r^{1-n},$$

where C_2 has to vanish to ensure \widehat{h}_{ti} is in L^2 , while C_3 has to vanish because the tensors $dt dx^i$ are not smooth at the axis of rotation $r = r_0$.

Thus $\widehat{h}_{\mu\nu} \equiv 0$ and, if we denote by h^\diamond the part of h from which all higher modes have been removed (in the notation of (4.2), and setting $\mathbb{S}^0(x^i) \equiv 1$ for simplicity,

$$(H.15) \quad h_{\mu\nu}^\diamond dx^\mu dx^\nu = f_{ab,0}^S dx^a dx^b + r f_{ai,0}^V dx^a dx^i + 2r^2 (H_{L,0}^S \gamma_{ij} + H_{T,0}^T \mathbb{T}_{ij}^0) dx^i dx^j,$$

we obtain

$$(H.16) \quad h^\diamond = \mathcal{L}_{Y^\diamond} \mathring{\mathfrak{g}}, \quad |Y^\diamond|_{\mathring{\mathfrak{g}}} = O(r^{-n-1}).$$

Appendix I. A linearised Birkhoff-type theorem with $K = -1$

In the case $K = -1$ the gauge transformations take the form

$$\begin{aligned}
 \text{(I.1)} \quad h_{tt} &\rightarrow h_{tt} + Y^r \partial_r f + 2f \partial_t Y^t, \\
 \text{(I.2)} \quad h_{tr} &\rightarrow h_{tr} + f^{-1} \partial_t Y^r + f \partial_r Y^t, \\
 \text{(I.3)} \quad h_{rr} &\rightarrow h_{rr} + Y^r \partial_r f^{-1} + 2f^{-1} \partial_r Y^r, \\
 \text{(I.4)} \quad h_{ij} &\rightarrow h_{ij} + \delta_{ij} \left(Y^r \frac{2r}{(x^{n+1})^2} - Y^{n+1} \frac{2r^2}{(x^{n+1})^3} \right) \\
 &\quad + \frac{r^2}{x^{n+1}} (\partial_i Y^j + \partial_j Y^i), \\
 \text{(I.5)} \quad h_{ti} &\rightarrow h_{ti} + f \partial_i Y^t + \frac{4r^2}{B^2} \partial_t Y^i, \\
 \text{(I.6)} \quad h_{ri} &\rightarrow h_{ri} + f^{-1} \partial_i Y^r + \frac{4r^2}{B^2} \partial_r Y^i,
 \end{aligned}$$

where we use the form $h_K = \sum_{i=2}^{n+1} (dx^i)^2 / (x^{n+1})^2$ of the hyperbolic metric.

We consider the scalar part of a $l = 0$ (i.e. $k = 0$) solution of the linearised Einstein equations in dimension $n = 2$. This will cover the full $l = 0$ case for $K = -1$, as the tensor part is controlled by the master functions and there are no vector modes with $l = 0$, as described in Section 4.

Such a perturbation takes the same form as for the $K = 1$ case, i.e.

$$\text{(I.7)} \quad h_{ab} = h_{ab}(t, r), \quad h_{ia} \equiv 0, \quad h_{ij} = \psi(t, r) \mathring{\mathfrak{g}}_{ij}.$$

We define a gauge vector Y_ε^\diamond , according to (F.12) and (F.13). This gives, for $r > r_0 + \varepsilon$ where $\varepsilon > 0$ is an arbitrary cutoff distance, $h_{tr} = \mathcal{L}_{Y_\varepsilon^\diamond} \mathring{\mathfrak{g}}_{tr}$ and $h_{ij} = \mathcal{L}_{Y_\varepsilon^\diamond} \mathring{\mathfrak{g}}_{ij}$.

As in (F.14) we define $\widehat{h}_{\mu\nu}$ as $h_{\mu\nu} - \mathcal{L}_{Y_\varepsilon^\diamond} \mathring{\mathfrak{g}}$, then \tilde{h}_{rr} , \tilde{h}_{tt} are defined as for $K = 1$,

$$\text{(I.8)} \quad \tilde{h}_{rr} := r f^2 \widehat{h}_{rr}, \quad \tilde{h}_{tt} := r (\widehat{h}_{tt} + f^2 \widehat{h}_{rr}),$$

and all this is inserted into the linearised Einstein tensor $G'_{\mu\nu}[\widehat{h}]$. We find that, for $r > r_0 + \varepsilon$

$$\text{(I.9)} \quad \tilde{h}_{rr} = 2\delta\mu, \quad \tilde{h}_{tt} = C(t)rf,$$

for some constant $\delta\mu$. As in the $K = 1$ case, $C \equiv 0$ as h is assumed to be in L^2 and, up to gauge, the only perturbations remaining are variations of the mass. For the case $K = -1$ these are never in L^2 .

We conclude that Y_0^\diamond is smooth everywhere and, if we denote again by h^\diamond the part of h from which all higher modes have been removed, we obtain

$$(I.10) \quad h^\diamond = \mathcal{L}_{Y_0^\diamond} \mathring{\mathfrak{g}}, \quad |Y^\diamond|_{\mathring{\mathfrak{g}}} = O(r^{-3}).$$

Appendix J. The Riemannian Kerr anti-de Sitter metrics

The Riemannian Schwarzschild-anti de Sitter metrics belong to the family of the Riemannian Kerr anti-de Sitter metrics, parameterised with m and an “angular momentum parameter” a . The variations of those metrics with respect to the parameter a provide non-trivial solutions of the linearised Einstein equations at the Schwarzschild-anti de Sitter metric, which we need to analyse. For this is it is convenient to start with a discussion of the family of the Riemannian Kerr anti-de Sitter metrics with small parameter a . Our presentation follows [10], where Kerr-Newman-de Sitter metrics were considered.

In Boyer-Lindquist coordinates, after the replacements $a \rightarrow ia$ and $t \rightarrow it$ the Kerr anti-de Sitter metric becomes

$$(J.1) \quad \mathfrak{g} = \frac{\Sigma}{\Delta_r} dr^2 + \frac{\Sigma}{\Delta_\theta} d\theta^2 + \frac{\sin^2(\theta)}{\Xi^2 \Sigma} \Delta_\theta (adt + (r^2 - a^2) d\varphi)^2 + \frac{1}{\Xi^2 \Sigma} \Delta_r (dt - a \sin^2(\theta) d\varphi)^2,$$

where, after setting $\lambda = \Lambda/3$, we have $\Sigma = r^2 - a^2 \cos^2(\theta)$ and

$$\Delta_r = (r^2 - a^2) (1 - \lambda r^2) - 2\mu r,$$

$\Delta_\theta = 1 - \lambda a^2 \cos^2(\theta)$, and $\Xi = 1 - \lambda a^2$.

Keeping in mind that we are interested in the metric for small a , we consider $\mu > 0$ and we assume that the largest zero of Δ_r , which we denote by r_0 , is positive. For $r \in [r_0, \infty)$ we introduce a new coordinate ρ defined as

$$(J.2) \quad \rho = \int_{r_0}^r \frac{1}{\sqrt{\Delta_r}} dr = \frac{2}{\sqrt{\kappa}} \sqrt{(r - r_0)} \mathbb{I}_1(r - r_0),$$

where

$$(J.3) \quad \kappa := |\Delta'_r|_{r=r_0} \neq 0,$$

and with a function $\mathbb{1}_1$ which is smooth near the origin and satisfies $\mathbb{1}_1(0) = 1$. Inverting, it follows that

$$(J.4) \quad r = r_0 + \frac{\kappa}{4}\rho^2\mathbb{1}_2(\rho^2), \quad \Delta_r = \frac{\kappa^2}{4}\rho^2\mathbb{1}_3(\rho^2),$$

with functions $\mathbb{1}_2, \mathbb{1}_3$ which are smooth near the origin, with $\mathbb{1}_2(0) = 1 = \mathbb{1}_3(0)$.

Smoothness at $r = r_0$ requires that t defines a $2\pi\omega$ -periodic coordinate, with

$$(J.5) \quad \omega := \frac{2\Xi(r_0^2 - a^2)}{\Delta'_r(r_0)}.$$

In order to guarantee regularity near the intersection of the axis $\{\sin\theta = 0\}$ with the axis $\{\Delta_r = 0\}$, near $\theta = 0$ we use a coordinate system $(\rho, \tau, \theta, \phi)$, with $t = \omega\tau$ and ϕ defined through the formula

$$(J.6) \quad d\varphi := \alpha d\phi + \frac{a}{a^2 - r_0^2}dt \equiv \alpha d\phi + \frac{a\omega}{a^2 - r_0^2}d\tau,$$

for some constants $\alpha, \omega \in \mathbb{R}^*$ which will be determined shortly by requiring 2π -periodicity of τ and ϕ . In this coordinate system the metric takes the form

$$(J.7) \quad g = \Sigma \left\{ d\rho^2 + \frac{1}{\Xi^2 \Sigma^2} \left[\frac{\kappa^2 \omega^2 \Sigma^2}{4(r_0^2 - a^2)^2} \mathbb{1}_4(\rho^2, \sin^2(\theta)) \rho^2 d\tau^2 + \alpha^2 (\Delta_\theta (a^2 - r^2)^2 + a^2 \Delta_r \sin^2(\theta)) \sin^2(\theta) d\phi^2 + F(\rho^2, \sin^2(\theta)) \rho^2 \sin^2(\theta) d\tau d\phi \right] + \frac{1}{\Delta_\theta} d\theta^2 \right\},$$

for some smooth functions $\mathbb{1}_4$ and F , with $\mathbb{1}_4(0, y) = 1$. As is well known, when (ρ, τ) are viewed as polar coordinates around $\rho = 0$, the one form $\rho^2 d\tau$ and the quadratic form $d\rho^2 + \rho^2 d\tau^2$ are smooth. Similarly when (θ, ϕ) are polar coordinates around $\theta = 0$, the one form $\sin^2(\theta) d\phi$ and the quadratic form $d\theta^2 + \sin(\theta)^2 d\phi^2$ are smooth. It is then easily inferred that the requirements of 2π -periodicity of τ and ϕ , together with

$$(J.8) \quad \frac{\kappa^2 \omega^2}{4\Xi^2 (r_0^2 - a^2)^2} = 1, \quad \frac{\alpha^2 \Delta_\theta^2 (a^2 - r^2)^2}{\Xi^2 (r^2 - a^2 \cos^2(\theta))^2} \Big|_{\theta=0} \equiv \alpha^2 = 1,$$

imply smoothness both of the sum of the diagonal terms of the metric g and of the off-diagonal term $g_{\tau\phi}d\tau d\phi$ on

$$\Omega := \{(r, \tau, \theta, \phi) \in [r_0, \infty) \times S^1 \times [0, \pi) \times S^1\}.$$

Here $[r_0, \infty) \times S^1$ is understood as \mathbb{R}^2 with center of rotation at r_0 , similarly $[0, \pi) \times S^1$ is understood as a disc D^2 of radius π .

The above calculations remain valid without changes near $\theta = \pi$. When $\theta \in (0, \pi]$ we will denote by $\widehat{\tau}$ and $\widehat{\phi}$ the relevant angular coordinates, and $\widehat{\omega}$, $\widehat{\alpha}$ the corresponding coefficients. Thus, for $\theta \in (0, \pi]$:

$$(J.9) \quad t = \widehat{\omega}\widehat{\tau}, \quad d\varphi = \widehat{\alpha}d\widehat{\phi} + \frac{a\widehat{\omega}}{a^2 - r_0^2}d\widehat{\tau},$$

with

$$(J.10) \quad \widehat{\omega} = \pm\omega, \quad \widehat{\alpha} = \pm 1.$$

We obtain likewise a smooth metric on the set

$$\widehat{\Omega} := \{(r, \widehat{\tau}, \theta, \widehat{\phi}) \in [r_0, \infty) \times S^1 \times (0, \pi) \times S^1\} \approx \mathbb{R}^2 \times D^2.$$

Disregarding issues of orientation, without loss of generality we can choose the plus signs above. The manifold M , obtained by patching together Ω with $\widehat{\Omega}$, using the obvious identifications resulting from the formulae

$$(J.11) \quad \omega d\tau = \widehat{\omega}d\widehat{\tau}, \quad \alpha d\phi + \frac{a\omega}{a^2 - r_0^2}d\tau = \widehat{\alpha}d\widehat{\phi} + \frac{a\widehat{\omega}}{a^2 - r_0^2}d\widehat{\tau},$$

is diffeomorphic to $\mathbb{R}^2 \times S^2$.

Note that while $d\varphi$ is a well defined one-form on M , the function φ is a well defined coordinate-modulo- 2π on M if and only if $\frac{a\omega}{a^2 - r_0^2} \in \mathbb{Z}^*$. We emphasise that it is *not* necessary to impose this last restriction to obtain a well defined smooth Riemannian metric on M , and we will not impose it.

Differentiating (J.1) with respect to a in the $(\rho, \tau, \theta, \phi)$ coordinates we obtain

$$(J.12) \quad \left. \frac{dg}{da} \right|_{a=0} = -2\alpha\omega \left(\frac{r^2 - r_0^2}{r_0^2} + f \right) \sin^2(\theta) d\tau d\phi.$$

Therefore variations with respect to the angular momentum parameter a are never in L^2 .

Appendix K. The master equation for vector perturbations

The object of this appendix is to justify the convergences of the mode-decomposition series in the vector sector. We thus consider the vector projection h^V of h , which we decompose into a complete (cf., e.g., [8]) set of vector harmonics \mathbb{V}_i^I :

$$(K.1) \quad \begin{aligned} h_{ab}^V &= 0, & h_{ai}^V &= r \sum_I f_{a,I}^V \mathbb{V}_i^I, \\ h_{ij}^V &= -r^2 \sum_{I: k_V(I) \neq 0} H_{T,I}^V \frac{1}{k_V(I)} (\widehat{D}_i \mathbb{V}_j^I + \widehat{D}_j \mathbb{V}_i^I), \end{aligned}$$

where $k_V = k_V(I)$ in the last sum is determined by the corresponding eigenvalue of the vector Laplacian $\widehat{\Delta}$ acting on \mathbb{V}_i^I :

$$(K.2) \quad \widehat{\Delta} \mathbb{V}_i^I = -k_V^2(I) \mathbb{V}_i^I.$$

For $k \in \mathbb{N}$ let $H^k(nN)$ denote the space of tensor fields on nN of Sobolev regularity with k derivatives. Standard functional analysis shows that we have

$$(K.3) \quad \|D_{i_1} \cdots D_{i_\ell} \mathbb{V}_i^I\|_{L^2(nN)} \approx (1 + k_V(I))^\ell,$$

where we use \approx to denote equivalence of norms, hence

$$(K.4) \quad \|h_{ai}^V dx^i\|_{H^k(nN)}^2 \approx r^2 \sum_I (1 + k_V(I)^2)^k |f_{a,I}^V|^2.$$

Similarly, for any j ,

$$(K.5) \quad \|D_{a_1 \dots a_j} (r^{-1} h_{ai}^V) dx^i\|_{H^k(nN)}^2 \approx \sum_I (1 + k_V(I)^2)^k |D_{a_1 \dots a_j} f_{a,I}^V|^2.$$

The Ishibashi-Kodama master functions $\Phi_{V,I}$ are defined, for

$$k_V(I) > (n - 1)K,$$

as solutions of the (integrable) system (cf., e.g., [12, Eqs. (2.13)–(2.15)]) of PDE's

$$(K.6) \quad \partial_b (r^{n/2} \Phi_{V,I}) = -r^{n-1} \varepsilon_b^a \left(f_{a,I}^V + \frac{r}{k_V(I)} D_a H_{T,I}^V \right).$$

Note that $r \geq r_0 > 0$ throughout, where r_0 is the location of the event horizon, so there is no issue of singularities arising in the equations at $r = 0$ in the current case.

If $\Phi_{V,I}$ is known we have, formally,

$$\begin{aligned}
 \text{(K.7)} \quad D_a(r^{-2}h_{ij}^V) &= - \sum_{I: k_V(I) \neq 0} \frac{1}{k_V(I)} D_a H_{T,I}^V (\widehat{D}_i \mathbb{V}_j^I + \widehat{D}_j \mathbb{V}_i^I) \\
 &= - \sum_{I: 0 < k_V(I) \leq (n-1)K} \frac{1}{k_V(I)} D_a H_{T,I}^V (\widehat{D}_i \mathbb{V}_j^I + \widehat{D}_j \mathbb{V}_i^I) \\
 &\quad + r^{-1} \sum_{I: k_V(I) > (n-1)K} (f_{a,I}^V - r^{-n+1} \varepsilon_a{}^b D_b(\Phi_{V,I})) (\widehat{D}_i \mathbb{V}_j^I + \widehat{D}_j \mathbb{V}_i^I).
 \end{aligned}$$

It is convenient to define the parts $h_{ij}^{V\star}$, respectively $h_{ai}^{V\star}$, of h_{ij}^V , respectively of h_{ai}^V , in which the low vector harmonics have been removed:

$$\text{(K.8)} \quad h_{ij}^{V\star} := h_{ij}^V + r^2 \sum_{I: 0 < k_V(I) \leq (n-1)K} \frac{1}{k_V(I)} H_{T,I}^V (\widehat{D}_i \mathbb{V}_j^I + \widehat{D}_j \mathbb{V}_i^I),$$

$$\text{(K.9)} \quad h_{ai}^{V\star} := h_{ai}^V - r \sum_{I: 0 < k_V(I) \leq (n-1)K} f_{a,I}^V \mathbb{V}_i^I.$$

From now on we assume that $\Phi_{V,I}$ vanishes. From (K.7) we then obtain

$$\text{(K.10)} \quad \|D_a(r^{-2}h_{ij}^{V\star}) dx^i dx^j\|_{H^k(nN)} \leq 2r^{-2} \|h_{ai}^{V\star} dx^i\|_{H^{k+1}(nN)},$$

which furthermore justifies the convergence and equality (K.7) in the $\Phi_{V,I} \equiv 0$ case. Set, again formally

$$\text{(K.11)} \quad Y_i^{V\star} := -r^2 \sum_{I: k_V(I) > (n-1)K} \frac{H_{T,I}^V}{k_V(I)} \mathbb{V}_i^I.$$

Then

$$\text{(K.12)} \quad \|Y_i^{V\star}\|_{H^k(nN)} = \|h_{ij}^{V\star}\|_{H^{k-1}(nN)},$$

which justifies convergence in (K.11). Next, again formally

$$\text{(K.13)} \quad D_a Y_i^{V\star} = -r^2 \sum_{I: k_V(I) > (n-1)K} \frac{D_a H_{T,I}^V}{k_V(I)} \mathbb{V}_i^I = -r \sum_{I: k_V(I) > (n-1)K} f_{a,I}^V \mathbb{V}_i^I = h_{ai}^{V\star}.$$

Hence

$$(K.14) \quad \|D_a Y^{V\star}\|_{H^k(nN)} = \|h_{ai}^{V\star} dx^i\|_{H^k(nN)}.$$

This justifies convergence in (K.13), and thus

$$(K.15) \quad h_{ai}^{V\star} = D_a Y_i^{V\star} = \mathcal{L}_{Y^{V\star}} \mathring{g}_{ai}.$$

Acknowledgements. The research of PTC was supported in part by the Austrian Research Fund (FWF), Project P29517-N27, and by the Polish National Center of Science (NCN) under grant 2016/21/B/ST1/00940. ED was supported by the grant ANR-17-CE40-0034 of the French National Research Agency ANR (project CCEM). PK was supported by a uni:docs grant of the University of Vienna. Useful discussions with Jacek Jezierski and Olivier Sarbach are acknowledged. We are grateful to Piotr Waluk and Jacek Jezierski for making their results available prior to publication.

References

- [1] L. Andersson and P. T. Chruściel, *On asymptotic behavior of solutions of the constraint equations in general relativity with “hyperboloidal boundary conditions”*, *Dissert. Math.* **355** (1996), 1–100 (English).
- [2] M. T. Anderson, P. T. Chruściel, and E. Delay, *Non-trivial, static, geodesically complete space-times with a negative cosmological constant. II. $n \geq 5$* , *AdS/CFT Correspondence: Einstein Metrics and Their Conformal Boundaries*, IRMA Lect. Math. Theor. Phys., Vol. 8, Eur. Math. Soc., Zürich, (2005), pp. 165–204. [arXiv:gr-qc/0401081](#).
- [3] M. Berger and D. Ebin, *Some decompositions of the space of symmetric tensors on a Riemannian manifold*, *Jour. Diff. Geom.* **3** (1969), 379–392.
- [4] A. L. Besse, *Einstein manifolds*, *Ergebnisse d. Math. 3. Folge*, Vol. 10, Springer, Berlin, (1987).
- [5] D. Birmingham, *Topological black holes in anti-de Sitter space*, *Class. Quantum Grav.* **16** (1999), 1197–1205. [arXiv:hep-th/9808032](#).
- [6] J. Bjoraker and Y. Hosotani, *Monopoles, dyons and black holes in the four-dimensional Einstein-Yang-Mills theory*, *Phys. Rev.* **D62** (2000), 043513. [arXiv:hep-th/0002098](#).

- [7] J. L. Blázquez-Salcedo, J. Kunz, F. Navarro-Lérida, and E. Radu, *Static Einstein-Maxwell magnetic solitons and black holes in an odd dimensional AdS spacetime*, *Entropy* **18** (2016), 438. [arXiv:1612.03747 \[gr-qc\]](#).
- [8] M. Boucetta, *Spectre des laplaciens de Lichnerowicz sur les sphères et les projectifs réels*, *Publ. Mat.* **43** (1999), 451–483.
- [9] P. T. Chruściel, E. Delay, and P. Klinger, *Non-singular spacetimes with a negative cosmological constant: IV. Stationary black hole solutions with matter fields*, [arXiv:1708.04947 \[gr-qc\]](#), (2017).
- [10] P. T. Chruściel and M. Hörzinger, *The Euclidean quantisation of Kerr-Newman-de Sitter black holes*, *Journal of High Energy Physics* **2016** (2016), no. 4, 1–37. [arXiv:1511.08496 \[hep-th\]](#).
- [11] G. Dotti, *Black hole non-modal linear stability: the Schwarzschild (A)dS cases*, *Class. Quantum Grav.* **33** (2016), 205005, 43.
- [12] A. Ishibashi and H. Kodama, *Stability of higher-dimensional Schwarzschild black holes*, *Progr. Theoret. Phys.* **110**, 901–919. [arXiv:hep-th/0305185](#).
- [13] A. Ishibashi and R. M. Wald, *Dynamics in non-globally hyperbolic static spacetimes III: Anti-de Sitter spacetime*, *Class. Quant. Grav.* **21** (2004), 2981.
- [14] J. Jezierski and P. Waluk, *Degrees of freedom of weak gravitational field on a spherically symmetric background*, *Acta Phys. Pol.* **10** (2017), 391–395.
- [15] J. Jezierski and P. Waluk, *Gauge invariant description of weak gravitational field on a spherically symmetric background with cosmological constant*, *Class. Quant. Grav.* **36** (2019), 215006. [arXiv:1806.09023 \[gr-qc\]](#).
- [16] M. Kimura, *A simple test for stability of black hole by S-deformation*, *Class. Quantum Grav.* **34** (2017), 235007. [arXiv:1706.01447 \[gr-qc\]](#).
- [17] P. Klinger, *Non-degeneracy of Riemannian Schwarzschild-anti de Sitter metrics: Birkhoff-type results in linearized gravity*, [arXiv:1806.05023 \[gr-qc\]](#), (2018).
- [18] H. Kodama and A. Ishibashi, *A master equation for gravitational perturbations of maximally symmetric black holes in higher dimensions*, *Prog. Theor. Phys.* **110** (2003), 701–722. [arXiv:hep-th/0305147](#).

- [19] H. Kodama and A. Ishibashi, *Master equations for perturbations of generalised static black holes with charge in higher dimensions*, Prog. Theor. Phys. **111** (2004), 29–73. [arXiv:hep-th/0308128](#).
- [20] H. Kodama, A. Ishibashi, and O. Seto, *Brane world cosmology: Gauge invariant formalism for perturbation*, Phys. Rev. **D62** (2000), 064022. [arXiv:hep-th/0004160](#).
- [21] H. Kodama and M. Sasaki, *Cosmological perturbation theory*, Prog. Theor. Phys. Suppl. **78** (1984), 1–166.
- [22] F. Kottler, *Über die physikalischen Grundlagen der Einsteinschen Gravitationstheorie*, Annalen der Physik **56** (1918), 401–462.
- [23] J. M. Lee, *Fredholm operators and Einstein metrics on conformally compact manifolds*, Mem. Amer. Math. Soc. **183** (2006), vi+83. [arXiv:math.DG/0105046](#).
- [24] A. Lichnerowicz, *Propagateurs et commutateurs en relativité générale*, Inst. Hautes Études Sci. Publ. Math. (1961), 56.
- [25] N. Okuyama and K. Maeda, *Five-dimensional black hole and particle solution with nonAbelian gauge field*, Phys. Rev. **D67** (2003), 104012, 18pp. [arXiv:gr-qc/0212022](#) [gr-qc].
- [26] E. Radu and D. H. Tchrakian, *Gravitating Yang-Mills fields in all dimensions*, 418th WE-Heraeus-Seminar: Models of Gravity in Higher Dimensions: From theory to Experimental search Bremen, Germany, August 25-29, 2008, (2009). [arXiv:0907.1452](#) [gr-qc].
- [27] M. A. Rubin and C. R. Ordóñez, *Eigenvalues and degeneracies for n -dimensional tensor spherical harmonics*, Journal of Mathematical Physics **25** (1984), no. 10, 2888–2894.
- [28] R. Schoen, *A lower bound for the first eigenvalue of a negatively curved manifold*, J. Differential Geom. **17** (1982), no. 2, 233–238.
- [29] P. Waluk, Master’s thesis, (2017).
- [30] E. Winstanley, *Existence of stable hairy black holes in $SU(2)$ Einstein Yang-Mills theory with a negative cosmological constant*, Class. Quant. Grav. **16** (1999), 1963–1978.

FACULTY OF PHYSICS AND ERWIN SCHRÖDINGER INSTITUTE
UNIVERSITY OF VIENNA, BOLTZMANNGASSE 5, A1090 WIEN, AUSTRIA
E-mail address: `piotr.chrusciel@univie.ac.at`

LABORATOIRE DE MATHÉMATIQUES D'AVIGNON
F-84 916 AVIGNON, ET FRUMAM - CNRS, MARSEILLE, FRANCE
AND ERWIN SCHRÖDINGER INSTITUTE, UNIVERSITY OF VIENNA
BOLTZMANNGASSE 5, A1090 WIEN, AUSTRIA
E-mail address: `Erwann.Delay@univ-avignon.fr`

FACULTY OF PHYSICS AND ERWIN SCHRÖDINGER INSTITUTE
UNIVERSITY OF VIENNA, BOLTZMANNGASSE 5, A1090 WIEN, AUSTRIA
E-mail address: `paul.klinger@univie.ac.at`

