

Interpolation Spaces and Energy Quantization for Yang-Mills Fields

TRISTAN RIVIÈRE

In [19] G. Tian proved that the defect measure of a weakly converging sequence of Yang-Mills Fields on a riemannian manifold (M, g) of dimension n ($n \geq 4$) is carried by a $n - 4$ -rectifiable subset S of M . In the present paper we complete the picture of weakly converging sequences of YM Fields by proving that, in 4 dimension, the defect measure is quantized: at any point of S it is the sum of L^2 energies of YM fields on S^4 . This result is extended to any dimension under some additional assumption on the $W^{2,1}$ norm of the curvature. In the last part we study non-linearities issued from curvatures in general and we prove that strong convergence in $W^{1,n/2}$ of Coulomb gauges of connections over a n -dimensional manifold preserves the topology of the bundle at the limit.

I. Introduction.

The first step in the program of building up a Gauge theory in higher dimension (higher than the conformal one) proposed by S.K. Donaldson and R.P. Thomas is to give a precise picture of the non-compactness in L^2 of Yang-Mills fields on a manifold M bellow some energy level.

Let E be a vector bundle over a riemannian manifold (M, g) with structure group a compact Lie Group G (denote by \mathfrak{G} its Lie Algebra). We will adopt the notations of [6] Chapter 2. A connection A on E is characterized by its corresponding covariant derivative ∇_A acting on sections of E , $\Omega^0(E)$ into sections of $E \otimes T^*M$. From ∇_A one may constuct the exterior derivative d_A from E -valued p -forms, $\Omega^p(E)$ into E -valued $p + 1$ -forms ([6] p. 35). We will denote by F_A the curvature of A which is a two form into the \mathfrak{G} -bundle deduced from the principal bundle of E for the adjoint representation ($F_A \in \Omega^2(\mathfrak{G}_E)$). A Yang-Mills Field of E is a critical point of the Yang-Mills action

$$(I.1) \quad \int_M |F_A|^2$$

where the scalar product on $\Omega^2(\mathfrak{G}_E)$ is deduced from the Killing form of G and the metric g on M (see [6]). F_A is Yang-Mills if and only if it solves the Euler-Lagrange Equation

$$(I.2) \quad d_A^* F_A = 0$$

where d_A^* is the adjoint of d_A for the above mentioned L^2 -scalar product on $\Omega^*(\mathfrak{G}_E)$.

In [19] G. Tian studied sequences of smooth solutions of (I.2) having uniformly bounded energy. One among the main results of Tian's paper can be stated in the following way.

Theorem I.1 ([19]). *Let A_i be a sequence of Yang-Mills connection with uniformly bounded energy, then one may extract a subsequence $A_{i'}$ such that, modulo gauge transformation,*

$$F_{A_{i'}} \rightharpoonup F_A \quad \text{weakly in } L^2$$

where F_A is the curvature of a smooth Yang-Mills connection outside a subset Σ of M having finite $n - 4$ -Hausdorff dimension,

$$|F_{A_{i'}}|^2 dV_g \longrightarrow |F_A|^2 dV_g + \Theta \mathcal{H}^{n-4} \llcorner S$$

as Radon measures where dV_g denotes the volume form on (M, g) , \mathcal{H}^{n-4} is the $n - 4$ Hausdorff measure, S is a $n - 4$ rectifiable subset of M containing Σ , $\mathcal{H}^{n-4} \llcorner S$ is the restriction to S of the Hausdorff measure and Θ is a measurable fonction on S admitting a positive lower bound depending only on n .

Of course the main difficulty in the above stated result is to establish the rectifiability property of the energy defect measure of the sequence $A_{i'} : \mu = \Theta \mathcal{H}^{n-4} \llcorner S$. The proof in [19] follows the approach of F.H. Lin in [9] where the rectifiability of the defect measure of stationary harmonic maps was established.

The main question we adress in the present paper concerns the multiplicity Θ of the defect measure in Theorem I.1. In the particular case when we are dealing with special YM connections which are Ω -anti-self-dual for some closed $n - 4$ form Ω (see [19]) G. Tian proved that $\Theta(x)$ is an integer, sum of the energies of anti-self-dual 4-dimensional instantons that concentrate at x , which means that there is no loss of energy between "bubbles". This phenomenon can be called the *no neck property* (see discussions about the existence or non existence of necks in bubble trees for harmonic maps and

“almost”-harmonic maps on surfaces in [5] and [16]). This no neck property was previously established in [6] in dimension 4 for sequences of anti-self-dual Yang-Mills fields and was an important step in the compactification of the moduli space of ASD-connections on a compact 4-manifolds.

Here we will extend this *no neck property* to sequences of general smooth YM Fields in dimension larger or equal to 4. We will use the following notation: $|\nabla_A \nabla_A F_A| = \sqrt{\sum_{i,j} |\nabla_i \nabla_j F_A|^2}$ and ∇_i denotes the covariant derivative induced by A along the i -th direction of \mathbb{R}^n . Our first main result is the following.

Theorem I.2. *Let $n \geq 4$ and let $\mu = \Theta S[\mathcal{H}^{n-4}$ be the defect measure of a sequence of Yang-Mills connections given in Theorem I.2, assuming in the case where $n > 4$ that $\|\nabla_A \nabla_A F_A\|_{L^1}$ is uniformly bounded, then μ is quantized:*

$$\text{for } \mathcal{H}^{n-4} \text{ a. e. } x \in S \quad \Theta(x) = \sum_{j=1}^{N_x} \|F_{B_j}\|_{L^2}^2$$

where N_x is a positive integer and B_j are Yang-Mills connections of G -bundles over S^4 issued from $A_{i'}$ and that concentrate at x as $i \rightarrow +\infty$.

Remark I.1. We believe that the assumption $\|\nabla_A \nabla_A F_A\|_{L^1}$ to be uniformly bounded in dimension strictly larger than 4 could be removed from the statement of Theorem I.2 above as it is here for $n = 4$.

Remark I.2. The result above is extended to sequences of eventually singular Yang-Mills Fields that are stationary in [13].

The meaning of *issued from $A_{i'}$ and that concentrate at x as $i \rightarrow +\infty$* has to be understood in the following way : There are conformal maps ψ_j^i from S^4 minus a fixed number of balls shrinking to zero, $\cup_k B_{i,j}^k$, into N_x disjoint subparts of the 4-dimensional plane perpendicular to $T_x S$ in $T_x M$ such that the pull-back bundle $\psi_j^{i*} \exp_x^* E$ extends to the smooth bundle over S^4 on which B_j is defined, moreover the pull back curvature $\psi_j^{i*} \exp_x^* F_{A_{i'}}$ converges strongly in $L^2(S^4 \setminus \cup_k B_{i,j}^k)$ to F_{B_j} (modulo changes of gauge). \exp_x denotes the exponential map from $(T_x M, g_x)$ into (M, g) .

The above stated quantization result we shall prove below is established using technics introduced by the author in collaboration with F.H. Lin while studying Ginzburg-Landau vortices in [10]. Right after this work we realized

that these techniques could be developed in order to prove the corresponding statement to Theorem I.2 but for harmonic maps between manifolds in [11] (see also [12] for an adaptation of these techniques to parabolic situations). As the main tool we introduced the use of Lorentz Spaces on subspaces having the conformal dimension of the problem. More specifically the originality of these works was to present a use of the $L^{2,\infty} - L^{2,1}$ duality as the right substitute of the apparently natural $L^2 - L^2$ duality in which we got stucked for quite a long time. This slight shift in duality is also the key point of the analysis we will present below. In the last part of the paper we extend the use of Lorentz Spaces to general curvatures of bundles over manifolds. One of the crucial observation is the following (for instance in 4-dimension): assume we are working in a Gauge A which is controlled in the $W^{1,2}$ -norm (the conformal invariant 1 in 4-dim) from the Peetre-Sobolev embeddings [15] we deduce that A is controlled in the Lorentz norm $L^{4,2}$ and from standard properties of products in Lorentz Spaces we obtain that $[A, A]$ is controlled in $L^{2,1}$ norm: in other words in the curvature $F_A = dA + [A, A]$ we observed that the non-linear part $[A, A]$, which makes the analysis of curvatures usually quite delicate, is in fact slightly more regular (in $L^{2,1}$) than the linear part dA itself which is a-priori only in L^2 . This kind of idea is also adapted to Yang-Mills equations (see Lemma II.3). In the last theorems of the paper we prove that strongly converging sequences of Coulomb gauges in $W^{1,n/2}$ for a given bundle over a n -manifold preserve the topology of the bundle at the limit. This was well known in $W^{1,p}$ for $p > n/2$ (see [21]) and is not true anymore in $W^{1,p}$ for $p < n/2$. These borderline situation is solved by the mean of Brezis-Wainger-Sobolev embeddings for Lorentz Spaces (see [3]).

Acknowledgments. The author is very grateful to Fanghua Lin for the exchanges of ideas which have motivated some aspects of this work. The author would like also to express his gratitude to Haïm Brezis and Luc Tartar for their comments and stimulating discussions on Lorentz Spaces.

II. Proof of Theorem I.2 for $n > 4$.

In order to simplify the presentation of the proof again, since the result is essentially local, we take the basis of E to be the flat unit ball B^n of \mathbb{R}^n .

Let S be the rectifiable subset carrying the defect measure $\nu = \Theta \mathcal{H}^{n-4} \llcorner S$ of the sequence A_i of Yang-Mills connections. For \mathcal{H}^{n-4} a.e. $x_0 \in S$, S admits a unique tangent plane P_0 at x_0 , moreover for \mathcal{H}^{n-4} a.e. $x_0 \in S$ we

have

$$(II.1) \quad \lim_{r \rightarrow 0} \frac{1}{r^{n-4}} \int_{B_r(x_0)} |F_A|^2 = 0$$

where A is the limiting admissible connection. This last fact is a consequence of a Federer-Zimmer type covering argument detailed in [19] p. 222. For any $\Lambda > 0$ we denote by G_Λ the following subset B^n which has to be a subset of S

$$G_\Lambda = \left\{ x; \liminf_{r \rightarrow 0} \liminf_{i \rightarrow +\infty} \frac{1}{r^{n-4}} \int_{B_r(x)} |\nabla_{A_i} \nabla_{A_i} F_{A_i}| \geq \Lambda \right\}.$$

For any $\delta > 0$ we may assign to each x in G_Λ a positive radius $r_x < \delta$ such that

$$\forall r < r_x \quad \liminf_{i \rightarrow +\infty} \frac{1}{r^{n-4}} \int_{B_r(x)} |\nabla_{A_i} \nabla_{A_i} F_{A_i}| \geq \frac{\Lambda}{2}.$$

From $(B_{r_x}(x))_{x \in G_\Lambda}$ we may extract a Besicovitch covering of G_Λ , $(B_{r_k}(x_k))_{k \in K}$, where K is at most countable, such that each point of B^n is covered by a number of balls $B_{r_k}(x_k)$ bounded by a universal number. Then, for any finite subfamily J of K , we have an n_J such that

$$\forall j \in J \quad \forall i \geq n_J \quad \int_{B_{r_j}(x_j)} |\nabla_{A_i} \nabla_{A_i} F_{A_i}| \geq \frac{r_j^{n-4} \Lambda}{4}$$

summing over j because of the Besicovitch property we deduce that

$$\sum_{j \in J} r_j^{n-4} \leq C \frac{\|\nabla_{A_i} \nabla_{A_i} F_{A_i}\|_{L^1(B^n)}}{\Lambda}$$

where C is a universal number independent on J . Thus we deduce $\sum_{k \in K} r_k^{n-4} \leq \frac{C}{\Lambda}$ and since δ can be chosen as small as we want, we have $\mathcal{H}^{n-4}(G_\Lambda) \leq C/\Lambda$. From this we deduce

$$\mathcal{H}^{n-4}(\cap_\Lambda G_\Lambda) = 0.$$

In other words we have For \mathcal{H}^{n-4} a.e. $x_0 \in \Sigma$

$$(II.2) \quad \liminf_{r \rightarrow 0} \liminf_{i \rightarrow +\infty} \frac{1}{r^{n-4}} \int_{B_r(x_0)} |\nabla_{A_i} \nabla_{A_i} F_{A_i}| < +\infty.$$

Choose now x_0 in Σ such that Σ admits a tangent plane P_0 at x_0 and both (II.1) and (II.2) hold. Let $\phi_r(x) = rx + x_0$, the previous statements imply

$$\begin{aligned} \lim_{r \rightarrow 0} \lim_{i \rightarrow +\infty} \frac{1}{r^{n-4}} \phi_r^* \mu_i &= \lim_{r \rightarrow 0} \frac{1}{r^{n-4}} \phi_r^* |F_A|^2 dx + \frac{1}{r^{n-4}} \phi_r^* \Theta \mathcal{H}^{n-4} \lfloor S \\ &= \Theta(x_0) \mathcal{H}^{n-4} \lfloor P_0 \end{aligned}$$

where μ_i denotes the measure $\mu_i = |F_{A_i}|^2 dx$ (dx is the Lebesgue measure). From (II.2) there exists $r_k \rightarrow 0$ as $k \rightarrow +\infty$ such that for all k

$$\liminf_{i \rightarrow +\infty} \frac{1}{r_k^{n-4}} \int_{B_{r_k}(x_0)} |\nabla_{A_i} \nabla_{A_i} F_{A_i}| \leq C$$

where C is fixed. Then for any k we may find i_k large enough such that

$$\lim_{k \rightarrow +\infty} \frac{1}{r_k^{n-4}} \phi_{r_k}^* \mu_{i_k} = \Theta(x_0) \mathcal{H}^{n-4} \lfloor P_0$$

and

$$\forall k \quad \frac{1}{r_k^{n-4}} \int_{B_{r_k}(x_0)} |\nabla_{A_{i_k}} \nabla_{A_{i_k}} F_{A_{i_k}}| \leq 2C.$$

Let $\tilde{A}_k(x) = r_k A_{i_k}(r_k x + x_0)$ for x in $B_1^n(0)$, we deduce that

$$\int_{B_1^n} |F_{\tilde{A}_k}|^2(x) dx = \frac{1}{r_k^{n-4}} \phi_{r_k}^* \mu_{i_k} \rightarrow \Theta(x_0) \mathcal{H}^{n-4} \lfloor P_0$$

and

$$\int_{B_1^n} |\nabla_{\tilde{A}_k} \nabla_{\tilde{A}_k} F_{\tilde{A}_k}| \leq 2C.$$

Thus modulo an adapted rescaling of A_i at such x_0 , our tangent defect measure $\Theta^{n-4}(\mu, x_0) = \Theta(x_0) \mathcal{H}^{n-4} \lfloor P_0$ is the defect measure of a sequence of Yang-Mills connections converging weakly to a flat connection A ($F_A = 0$):

$$(II.3) \quad |F_{A_i}|^2 dx \rightharpoonup \Theta(x_0) \mathcal{H}^{n-4} \lfloor P_0 \quad \text{in } B_1.$$

To simplify notations we can take $x_0 = 0$ and P_0 to be the $n - 4$ -dimensional plane $\{x_i = 0\}$ for $i = n - 3 \dots n$. We will simply denote $\Theta(x_0)$ by Θ that we assume to be non zero. Observe that we can also have chosen x_0

and the rescaled rate such that our rescaled connection A_i verifies for any $k = 1 \cdots n - 4$

$$(II.4) \quad \lim_{i \rightarrow +\infty} \int_{B_1} \left| \frac{\partial}{\partial x_k} \lfloor F_{A_i} \right|^2 dx = 0$$

(cf. Lemma 3.3.2 of [19]) where $\frac{\partial}{\partial x_k} \lfloor F_{A_i}$ denotes the contraction of $\frac{\partial}{\partial x_k}$ and F_{A_i} that lies in $\Omega^1(\mathfrak{G}_E)$.

We use the following notations $X_1 = (x_1 \cdots x_{n-4})$ and $X_2 = (x_{n-3} \cdots x_n)$. Let

$$f_i(X_1) = \int_{\{X_1\} \times B_1^4(0)} \sum_{k=1}^{n-4} \left| \frac{\partial}{\partial x_k} \lfloor F_{A_i} \right|^2.$$

Since $f_i \rightarrow 0$ in $L^1(B_1^{n-4}(0))$, its maximal function

$$M f_i(X_1) = \sup_{r>0} \frac{1}{r^{n-4}} \int_{B_r^{n-4}(X_1)} f_i$$

tends to zero in L^1_{weak} :

$$\lim_{i \rightarrow +\infty} \|\lambda \{X_1 ; M f_i(X_1) \geq \lambda\}\|_{L^\infty(\mathbb{R})} = 0.$$

Thus there exists a set $E_i, |E_i| \geq 0,99|B_1^{n-4}(0)|$ such that

$$(II.5) \quad \|M f_i\|_{L^\infty(E_i)} \rightarrow 0.$$

Using the fact that $\|\nabla_{A_i} \nabla_{A_i} F_{A_i}\|_{L^1(B_1^n)}$ is uniformly bounded and Kato inequality $|d|\nabla_{A_i} F_i| \leq |\nabla_{A_i} \nabla_{A_i} F_i|$, (we keep denoting the curvature of A_i by F_i) we deduce that

$$(II.6) \quad \int_{B_1(0)} |\nabla_{A_i} F_i|^{\frac{4}{3}} + |d|\nabla_{A_i} F_i| \leq C$$

where C is independant on i . By Fubini there exists a measurable subset G_i of $B_1^{n-4}(0)$, verifying $|G_i| \geq 0,99|B_1^{n-4}|$ and such that

$$(II.7) \quad \forall X_1 \in G_i \quad \int_{\{X_1\} \times B_1^4(0)} |\nabla_{A_i} F_i|^{\frac{4}{3}} + |d|\nabla_{A_i} F_i| \leq C$$

where C is independant on i and X_1 in G_i . Using now the embedding of $W^{1,1}(B^4)$ into the Lorentz Space $L^{\frac{4}{3},1}(B^4)$ (this embedding was first noticed

by J. Peetre [15], see also [18] and a short proof in [8] Theorem 3.3.9) we get that

$$(II.8) \quad \forall X_1 \in G_i \quad \|\nabla_{A_i} F_i\|_{L^{\frac{4}{3},1}(\{X_1\} \times B_1^4(0))} \leq C$$

which yields, by Kato inequality,

$$(II.9) \quad \forall X_1 \in G_i \quad \|d|F_i|\|_{L^{\frac{4}{3},1}(\{X_1\} \times B_1^4(0))} \leq C.$$

C again is independant of i . We may also have chosen G_i such that, for any X_1 in G_i , $\int_{\{X_1\} \times B_1^4(0)} |F_i|^2$ is bounded independantly of i preserving the size of $G_i : |G_i| \geq 0,99|B_1^{n-4}|$. Using now theorem 8 of [18] we get that the $L^{2,1}$ norm of F_i on $\{X_1\} \times B_1^4(0)$ is uniformly bounded

$$(II.10) \quad \forall X_1 \in G_i \quad \|F_i\|_{L^{2,1}(\{X_1\} \times B_1^4(0))} \leq C.$$

We will call a “good slice” a point $X_1 \in G_i \cap E_i$ (this set has a measure bounded from bellow independantly of i).

We claim that for any $X_1 \in B_1^{n-4}(0)$ and any fixed δ , for i large enough,

$$(II.11) \quad \sup_{X_2 \in B_1^4(0)} \delta^{4-n} \int_{B_\delta(X_1, X_2)} |F_{A_i}|^2 \geq \varepsilon(n)$$

where $\varepsilon(n)$ is the Uhlenbeck constant given in Theorem 2.2.1 of [19]. That yields to the control of the L^∞ norm of the curvature in the ball of half radius. If (II.11) would not hold we would contradicts the concentration phenomenon and the fact that $\Theta \neq 0$.

Pick now an arbitrary sequence of “good slices” $\{X_1^i\}$ as $i \rightarrow +\infty$. Because of (II.11), since each A_i is smooth, we may find a sequence $\delta_i \rightarrow 0$ such that

$$(II.12) \quad \max_{X_2} \delta_i^{4-n} \int_{B_{\delta_i}(X_1^i, X_2)} |F_{A_i}|^2 = \delta_i^{4-n} \int_{B_{\delta_i}(X_1^i, X_2^i)} |F_{A_i}|^2 = \frac{\varepsilon(n)}{2}$$

($1/2$ can be replaced by any positive number between 0 and 1). In other words X_2^i is located at a bubble of characteristic size δ_i . More precisely, if one introduces $B_i = \tau_i^* A_i$ where $\tau_i : X \rightarrow \delta_i X + (X_1^i, X_2^i)$, we have, following [19] p. 237 that, modulo extraction of a subsequence and eventually changes of gauge,

$$(II.13) \quad F_{B_i} \rightarrow F_B \quad \text{in } C_{\text{loc}}^k(B_1^{n-4}(0) \times \mathbb{R}^4)$$

moreover B is gauge equivalent to the pull-back by the orthogonal projection map of a non flat Yang-Mills connection on $\{0\} \times \mathbb{R}^4$. This is the first bubble we detect.

At this stage of the proof it is tempted to compare Θ and $\int_{\{0\} \times \mathbb{R}^4} |F_B|^2$ and to deduce from the above convergence that $\Theta \geq \int_{\{0\} \times \mathbb{R}^4} |F_B|^2$. This will be a consequence of the following lemma

Lemma II.1. *Let A_i be a sequence of Yang-Mills connections on B_1^n such that*

$$(II.14) \quad |F_{A_i}|^2 dx \longrightarrow \Theta_0 \mathcal{H}^{n-4} \llcorner P_0$$

as Radon measures, where Θ_0 is a positive constant and P_0 is the $n-4$ plane $\{x_k = 0; k = n - 3 \cdots n\}$, assume also that

$$(II.15) \quad \forall k = 1 \cdots n - 4 \quad \frac{\partial}{\partial x_k} \llcorner F_{A_i} \longrightarrow 0 \quad \text{in } L^2(B_1^n)$$

then for a.e. $X_1 \in P_0$

$$(II.16) \quad \lim_{i \rightarrow +\infty} \int_{\{X_1\} \times B_1^4(0)} |F_{A_i}|^2 = \Theta_0.$$

Proof of Lemma II.1. The idea of the proof comes from Lin's work [9] in the context of stationary harmonic maps. Computations which are required to prove the lemma can be found in [19] part 3.

Let $\phi \in C_0^\infty(B_1^4(0))$ such that $\phi(X_2) \equiv 1$ in $B_{1/2}^4(0)$, introduce

$$m_i(X_1) = \int_{B_1^4(0)} \phi^2 |F_{A_i}|^2(X_1, X_2) dX_2$$

following (3.3.19) of [19] we obtain that m_i solves the following equation:

$$(II.17) \quad \text{grad } m_i = f_i + \text{div } u_i$$

where $f_i : B_1^{n-4} \rightarrow \mathbb{R}^{n-4}$ and $u_i : B_1^{n-4} \rightarrow \mathbb{R}^{n-4} \times \mathbb{R}^{n-4}$ verify

$$(II.18) \quad \|f_i\|_{L^1(B_1^{n-4})} + \|u_i\|_{L^1(B_1^{n-4})} \longrightarrow 0.$$

(II.15) is of course used to establish (II.18.) From Allard strong constancy Lemma [1] we deduce the existence of a constant C_i such that

$$(II.19) \quad \|m_i - C_i\|_{L^1_{\text{loc}}(B_1^{n-4})} \longrightarrow 0.$$

Since $\int_{B_r^{n-4}} m_i \rightarrow |B_r^{n-4}| \Theta_0$, we deduce that $C_i \rightarrow \Theta_0$ as $i \rightarrow +\infty$ and Lemma II.3 is proved since $|F_{A_i}|$ converges to 0 in $B_1^n \setminus P_0$. \square

Because of (II.16) we may of course choose X_1^i in $E_i \cap G_i$ such that

$$(II.20) \quad \lim_{i \rightarrow \infty} \int_{\{X_1^i\} \times B_1^4(0)} |F_{A_i}|^2 = \Theta.$$

Thus at this stage of the proof we have shown that

$$(II.21) \quad \Theta \geq \int_{\mathbb{R}^4} |F_B|^2$$

where B is a non flat YM-connection of \mathbb{R}^4 .

We can restate the goal to achieve in order to prove Theorem I.2: the aim is to show that

$$(II.22) \quad \lim_{i \rightarrow +\infty} \int_{\{X_1^i\} \times B_1^4(0)} |F_{A_i}|^2 = \sum_{j=1}^m \int_{\mathbb{R}^4} |F_{B_j}|^2$$

where F_{B_j} are non flat Yang-Mills connections on \mathbb{R}^4 . This will be shown by induction on the total number of bubbles m . Indeed we know that each concentration generating a bubble has a cost of energy bounded from below by a positive number $\varepsilon(4)$ (recall that $\varepsilon(4)$ is the Uhlenbeck constant in 4 dimension see [20]). Thus the number of possible bubbles is uniformly bounded.

Assume first we have only one bubble which is the one already detected of characteristic size δ_i . Clearly we have

$$(II.23) \quad \int_{\mathbb{R}^4} |F_B|^2 = \lim_{R \rightarrow +\infty} \lim_{i \rightarrow +\infty} \int_{\{X_1^i\} \times B_{R\delta_i}^4(X_2^i)} |F_{A_i}|^2.$$

In order to prove that

$$(II.24) \quad \Theta = \lim_{i \rightarrow +\infty} \int_{\{X_1^i\} \times B_1^4(0)} |F_{A_i}|^2 = \int_{\mathbb{R}^4} |F_B|^2$$

we have to prove that

$$(II.25) \quad \lim_{R \rightarrow +\infty} \lim_{i \rightarrow +\infty} \int_{\{X_1^i\} \times B_1^4 \setminus B_{R\delta_i}^4(X_2^i)} |F_{A_i}|^2 = 0$$

in other words that there is no “neck” formation and no loss of energy which is not quantized (see [7] and [16] for “neck” phenomenon in bubble trees).

To simplify notations we take (X_1^i, X_2^i) to be the origin $(0, 0)$.

We claim that for any ε there are sufficiently large $R > 0$ and $i_0 \in \mathbb{N}$ such that

$$(II.26) \quad \forall i \geq i_0 \quad \forall R\delta_i \leq r \leq \frac{1}{2}$$

$$\frac{1}{r^{n-4}} \int_{B_r^{n-4}(0)} \int_{B_{2r}^4(0) \setminus B_r^4(0)} |F_{A_i}|^2 \leq \varepsilon.$$

Assume this is not the case, then we may find an $\varepsilon_0 > 0$, a subsequence $i' \rightarrow +\infty$ (still denoted i) and a sequence r_i such that

$$(II.27) \quad \frac{1}{r_i^{n-4}} \int_{B_{r_i}^{n-4}(0)} \int_{B_{2r_i}^4(0) \setminus B_{r_i}^4(0)} |F_{A_i}|^2 \geq \varepsilon_0$$

$$\frac{r_i}{\delta_i} \rightarrow +\infty.$$

(We may always assume that $r_i \rightarrow 0$ since $F_{A_i} \rightarrow 0$ in $B^n \setminus P_0$.) Let $\lambda_i \rightarrow 0$ such that $\frac{r_i}{\lambda_i} = o(1)$ and $\lambda_i \frac{r_i}{\delta_i} \rightarrow +\infty$. Consider the dilation map $\tau_i(X) = r_i X$ in \mathbb{R}^n and let $B_i = \tau_i^* A_i$, thus B_i is a Yang-Mills connection on $B_1^{n-4}(0) \times B_{2/\lambda_i}^4(0) \setminus B_{\lambda_i}^4$ and satisfies

$$(II.28) \quad \int_{B_1^{n-4}(0)} \int_{B_2^4(0) \setminus B_1^4(0)} |F_{B_i}|^2 \geq \varepsilon_0.$$

Taking into account the translation by (X_1^i, X_2^i) , (II.5) implies

$$(II.29) \quad \limsup_{i \rightarrow +\infty} \frac{1}{r} \frac{1}{r^{n-4}} \int_{B_r^{n-4}} \int_{B_{2/\lambda_i}^4(0) \setminus B_{\lambda_i}^4(0)} \left| \frac{\partial}{\partial x_k} [F_{B_i}] \right|^2 = 0.$$

Thus, from Proposition 3.1.2 of [19], modulo extraction of a subsequence and eventually changes of gauge, we have

$$F_{B_i} \rightharpoonup F_{B_\infty} \quad \text{in } L^2_{\text{loc}}(B_1^{n-4}(0) \times \mathbb{R}^4 \setminus \{0\})$$

where

$$(II.30) \quad \forall k = 1 \cdots n - 4 \quad \frac{\partial}{\partial x_k} [F_{B_\infty}] = 0$$

$$\int_{B_1^{n-4}(0) \times \mathbb{R}^4} |F_{B_\infty}|^2 < +\infty$$

B_∞ is an admissible YM connection

arguing like in [19] p. 237 we deduce that F_{B_∞} is gauge equivalent to the pull-back of the projection on $\{0\} \times \mathbb{R}^4$ of a Yang-Mills connection on $\{0\} \times \mathbb{R}^4 \setminus \{0\}$ which has to be smooth at $\{0\}$ because of the removability singularity result of Uhlenbeck [20]. Thus B_∞ is gauge equivalent to the pull-back of an \mathbb{R}^4 YM-connection. If B_∞ is not flat we already get a contradiction because we just exhibited a new bubble. If F_{B_∞} is flat, that means that the convergence of F_{B_i} to F_{B_∞} admits a non zero defect measure

$$|F_{B_i}|^2 \rightarrow \nu = f(x) \mathcal{H}^{n-4} \llcorner \Sigma$$

and because of (II.29) we easily deduce, using arguments like in the proof of Lemma II.1, that Σ is contained in $n - 4$ -planes P_i parallel to $\{x_k = 0; k \geq n - 3\}$ and that f is constant on each of the P_i s:

$$(II.31) \quad \nu = \sum_j C_j \mathcal{H}^{n-4} \llcorner P_j$$

where the C_j are non zero constants. Following again [19] page 235 we get a new bubble and this contradict our assumption that there is only one bubble.

Thus in any cases (II.27) yields a contradiction and (II.26) holds. For ε small enough in (II.26) we are in the setting of Uhlenbeck ε -regularity theorem ([19] Theorem 2.2.1 or [14]) and we deduce the existence of ε_0 such that for any $\varepsilon < \varepsilon_0$, there exists $R > 0$ and $i_0 \in \mathbb{N}$ such that

$$(II.32) \quad \forall i \geq i_0 \quad \forall |X| \geq R\delta_i \quad |X|^4 |F_{A_i}|^2 \leq \varepsilon$$

restricting F_{A_i} to the plane $\{0\} \times \mathbb{R}^4$ we deduce from (II.32)

$$(II.33) \quad \|F_{A_i}\|_{L^{2,\infty}(\{0\} \times B_{1/2}^4 \setminus B_{R\delta_i}^4)}^2 \leq \varepsilon$$

where $L^{2,\infty}$ is the Lorentz Space defined for instance in [17]. Combining (II.33) with (II.10) and using the $L^{2,\infty} - L^{2,1}$ duality (see [17]) we deduce that for every $\varepsilon < \varepsilon_0$ there exists an R such that for any $i \geq i_0$ we have

$$(II.34) \quad \|F_{A_i}\|_{L^2(B_{1/2}^4 \setminus B_{R\delta_i}^4)}^2 \leq C\varepsilon.$$

This implies (II.25) and Theorem I.2 is proved in the case of 1 bubble.

The case of more than 1 bubble can be handled in a very similar way and we just give few details for $m = 2$.

The proof starts the same untill (II.26) which cannot hold anymore otherwise we would have had 1 bubble only as it is explained above. Thus we

may extract r_i such that (II.27) holds. We have two possibilities the first: F_{B_i} has no defect measure and the convergence is strong and it remains to show that

$$(II.35) \quad \lim_{R \rightarrow +\infty} \lim_{i \rightarrow +\infty} \int_{\{0\} \times B_{r_i/R}^4 \setminus B_{R\delta_i}^4} |F_{B_i}|^2 = 0$$

and

$$(II.36) \quad \lim_{R \rightarrow +\infty} \lim_{i \rightarrow +\infty} \int_{\{0\} \times B_{1/2}^4 \setminus B_{Rr_i}^4} |F_{B_i}|^2 = 0.$$

This can be done exactly like in the proof of (II.27) since we already got the second bubble given by F_{B_∞} . The second situation is when $|F_{B_i}|^2$ has some non-zero defect measure and it is given by (II.31). We need to have $F_{B_\infty} = 0$ otherwise there would be too many bubbles. Thus since $F_{B_\infty} = 0$ we are back to the case $m = 1$. In this case also one has to establish (II.35) and (II.36) and this can be done again following word by word the proof of (II.27). The case $m = 2$ is proved. For larger m one follows exactly the same strategy. □

III. The conformal dimension $n = 4$.

In this part we explain, in the case $n = 4$ how we can omit the hypothesis on the $W^{2,1}$ bound of the curvature to prove Theorem I.2. Theorem I.2 was not known at the conformal dimension $n = 4$ in general but only established for ASD connections in this dimension (see the analysis in [20], [21] and [6]). We follow the same strategy as the one developed in part II and we keep denoting δ_i the characteristic size of the first bubble given by (II.12) that we assume to be centered at the origin. In order to clarify the presentation we assume that there is only one bubble so that (II.26) holds:

$$(III.1) \quad \forall \varepsilon \quad \exists R \equiv R_\varepsilon \quad \text{s.t.} \quad \forall i \geq i_0 \quad \forall r \text{ s.t. } R_\varepsilon \delta_i \leq r \leq 4$$

$$\int_{B_{16r}^4(0) \setminus B_r^4(0)} |F_{A_i}|^2 \leq \varepsilon.$$

We claim the following lemma holds:

Lemma III.1. *There exists $\varepsilon_0 > 0$ such that for any $\varepsilon < \varepsilon_0$ There exists a trivialization of the bundle on $B^4 \setminus B_{R_\varepsilon \delta_i}(0)$ such that, in this trivialization*

$$\|A_i\|_{W^{1,2}(B^4 \setminus B_{R_\varepsilon \delta_i}(0))}$$

is uniformly bounded, independently on i and ε .

Proof of Lemma III.1. We first have a fixed trivialisation over B^4 (coming from the bundle we fixed) in which we have no control of the sequence of connections A_i which are 1-forms defined on all of B^4 into the Lie algebra \mathfrak{G} of the structure group G . Let R given by (III.1) for an ε that will be fixed later on in the proof. We introduce the following family of annuli centered at $0 : T^k = B_{2^{-k+2}}(0) \setminus B_{2^{-k-2}}(0)$ for the integers going from 0 to $N_i = [\log_2(1/R\delta_i) + 2]$ ($[s]$ denotes the integer part of the real s). Applying Uhlenbeck's Theorem 1.3 in [21], for ε chosen small enough, independent on i or k , for any $k = 0 \dots N_i$, there exists a change of gauge σ_i^k on T^k such that if we denote A_i^k the connection expressed in this gauge ($A_i^k = (\sigma_i^k)^{-1}A_i\sigma_i^k + (\sigma_i^k)^{-1}d\sigma_i^k$) we have

$$(III.2) \quad \int_{T_i^k} |\nabla A_i^k|^2 + \frac{|A_i^k|^2}{|x|^2} \leq c \int_{T_i^k} |F_{A_i}|^2 \leq c\varepsilon$$

(observe that we have no control on the σ_i^k but only on the transition functions $(\sigma_i^k)^{-1}\sigma_i^{k-1}$ as we will see bellow). Using again Uhlenbeck regularity theorem in ([20] p. 18), we have the estimate in the smaller annuli $\hat{T}_i^k = B_{2^{-k+1}}(0) \setminus B_{2^{-k-1}}(0)$

$$(III.3) \quad |\nabla^l A_i^k| \leq C_l \frac{\varepsilon^{1/2}}{|x|^{l+1}}.$$

On $\hat{T}^k \cap \hat{T}^{k-1} = B_{2^{-k+1}}(0) \setminus B_{2^{-k}}(0)$ let $g_i^{k,k-1}$ be the transition function given by

$$g_i^{k,k-1} = (\sigma_i^k)^{-1}\sigma_i^{k-1}.$$

We have

$$(III.4) \quad dg_i^{k,k-1} = g_i^{k,k-1}A_i^{k-1} - A_i^k g_i^{k,k-1}$$

and from (III.3) we deduce

$$(III.5) \quad |\nabla g_i^{k,k-1}| \leq C\varepsilon/|x| \quad \text{and} \quad |\nabla^2 g_i^{k,k-1}| \leq C\varepsilon/|x|^2.$$

Then there exists $\bar{g}_i^{k,k-1}$ in G such that

$$(III.6) \quad \|g_i^{k,k-1} - \bar{g}_i^{k,k-1}\|_{L^\infty(\hat{T}^k \cap \hat{T}^{k-1})} \leq C\varepsilon.$$

Let $h^k = \prod_{l \leq k-1} \bar{g}_i^{l,l-1}$ (ordered product $\bar{g}_i^{k-1,k-2} \bar{g}_i^{k-2,k-3} \dots$), replacing $g_i^{k,k-1}$ by $\tilde{g}_i^{k,k-1} = (h^k)^{-1} g_i^{k,k-1} h^{k-1}$ we have

$$(III.7) \quad \|\tilde{g}_i^{k,k-1} - \text{id}\|_{L^\infty(\hat{T}^k \cap \hat{T}^{k-1})} = \|\bar{g}_i^{k,k-1} g_i^{k,k-1} - \text{id}\|_{L^\infty(\hat{T}^k \cap \hat{T}^{k-1})} \leq \varepsilon.$$

So changing σ_i^k into $\sigma_i^k h^k$, we may assume $\bar{g}_i^{k,k-1} = \text{id}$. We assume now that ε is chosen small enough so that Uhlenbeck's Theorem (III.2) applies and so that, in view of (III.6), $g_i^{k,k-1}$ is in the neighborhood of the identity where \exp is invertible. The goal is to express $g_i^{k,k-1}$ as transition functions (eventually on smaller intersecting set than the $\hat{T}^k \cap \hat{T}^{k-1}$) of a sequence of trivialisations ρ_i^k on which we have some control. Let ϕ be a smooth map on $B_2(0)$ such that $\phi \equiv 1$ on $B_2 \setminus B_{2^{1/2}}$ and $\phi \equiv 0$ on B_1 . Denote by $\phi^k(x) = \phi(2^k x)$. ρ^k is now defined by induction on k . Let $\rho_i^0 \equiv 1$. assume now ρ_i^{k-1} is constructed, we denote

$$Y_i^k = \exp^{-1}(\rho_i^{k-1} g_i^{k,k-1})$$

and we choose on $\tilde{T}^k = B_{2^{-k+1}}(0) \setminus B_{2^{-k-1/2}}(0)$, $\rho^k := \exp(-\phi^k Y_i^k)$. Since $\rho^{k-1} \equiv \text{id}$ on $B_{2^{-k+1}}$ we have

$$|\nabla^l \rho_i^k| \leq C \sum_{j=0}^l |\nabla^j \phi^k| |\nabla^{l-j} Y^k| \leq C \sum_{j=0}^l 2^{jk} |\nabla^{l-j} g_i^{k,k-1}|.$$

Thus from (III.6) and (III.7), we have for $l = 0, 1, 2$

$$(III.8) \quad |\nabla^l \rho_i^k| \leq \varepsilon / |x|^l$$

and from (III.2) and (III.4)

$$(III.9) \quad \int_{\tilde{T}^k} |\nabla^2 \rho_i^k|^2 + \frac{|\nabla \rho_i^k|^2}{|x|^2} \leq C \int_{T^k} |F_{A_i}|^2.$$

Moreover on $\tilde{T}^k \cap \tilde{T}^{k-1} = B_{2^{-k+1}} \setminus B_{2^{-k+1/2}}$, by construction $g_i^{k,k-1} = (\rho_i^k)^{-1} \rho_i^{k-1}$. Let \tilde{F}_i be the two form on $B_2 \setminus B_{R\delta_i}$ into the Lie algebra of G that coincide with $(\rho_i^{k-1})^{-1} F_i^{k-1} \rho_i^{k-1} = (\rho_i^k)^{-1} F_i^k \rho_i^k$ in tiT^k , it is our original sequence of curvatures F_{A_i} expressed in the trivialisations that coincide with $\rho_i^k \sigma_i^k$ on the tiT^k 's. the corresponding 1-form expressing the connection in this gauge is $\tilde{A}_i = (\rho_i^k)^{-1} A_i^k \rho_i^k + (\rho_i^k)^{-1} d\rho_i^k$ on \tilde{T}^k . Combining

(III.2) and (III.8) and (III.9), we have

$$(III.10) \quad \int_{B_2 \setminus B_{2R\delta_i}} \frac{|\tilde{A}_i|^2}{|x|^2} + |\nabla \tilde{A}_i|^2 \leq C \sum_{k=0}^{N_i} \int_{T^k} \frac{|A_i^k|^2}{|x|^2} + |\nabla A_i^k|^2 \leq C \int_{B_4 \setminus B_{R\delta_i}} |F_{A_i}|^2.$$

Finally the $L^{2,1}$ estimate of the curvature and the quantization property (Theorem I.1 for $n = 4$) is a consequence of inequality (III.10) and the following observation.

Lemma III.2. *Let A be the 1-form of a Yang-Mills connection on a given trivialization of a principal bundle over B^4 , the following estimate holds*

$$(III.11) \quad \|F_A\|_{L^{2,1}(B^4_{1/2})} \leq C \left[\|A\|_{W^{1,2}(B^4)} + \|A\|_{W^{1,2}(B^4)}^3 \right]$$

where F_A is the curvature $F_A = dA + [A, A]$ and $\|\cdot\|_{L^{2,1}}$ denotes the Lorentz norm of the space dual to the weak L^2 space.

Proof of Lemma III.2. Using whitney extension theorem we may find an extension \tilde{A} of A on all of \mathbb{R}^4 such that $\|\tilde{A}\|_{W^{1,2}(\mathbb{R}^4)} \leq \|A\|_{W^{1,2}(B^4)}$. Then Using the Lorentz-Sobolev embedding of J. Peetre [15], we have

$$(III.12) \quad \|A\|_{L^{4,2}(B^4)} \leq \|\tilde{A}\|_{L^{4,2}(\mathbb{R}^4)} \leq C \|\tilde{A}\|_{W^{1,2}(\mathbb{R}^4)}$$

where $L^{4,2}$ denotes the Lorentz Space of function f such that $\int_{\mathbb{R}} f_*^2(t) t^{-1/2} < +\infty$ (f_* is the decreasing rearrangement of f). Now using the following property on multiplication of functions in Lorentz Spaces

$$(III.13) \quad \forall a, b \text{ funct. on } B^4 \quad \|ab\|_{L^{4/3,1}(B^4)} \leq \|a\|_{L^{4,2}(B^4)} \|b\|_{L^{2,2}(B^4)}$$

($L^{2,2}$ is the usual L^2 Space) see [18]. Since $F = dA + [A, A]$ solves

$$dF = -[A, F] \quad \text{and} \quad d^*F = -*[A, *F].$$

Combining (III.12) and (III.13) we get

$$(III.14) \quad \|dF\|_{L^{4/3,1}(B^4)} + \|d^*F\|_{L^{4/3,1}(B^4)} \leq \|A\|_{W^{1,2}(B^4)} \|F\|_{L^2(B^4)}.$$

let G on B^4 such that $dG = dF$, $d^*G = d^*F$ and $\iota_{\partial B^4}^* G = 0$ (where $\iota_{\partial B^4}$ is the embedding of ∂B^4 in B^4). The operator that assign to the pair (dG, d^*G)

the tensor $\nabla G = \sum_{k=1}^4 \frac{\partial G}{\partial x_k} \otimes dx_k$ is a weak (r, r) operator for r between 1 and ∞ , then using Theorem 3.15 Chapter V of [17] we obtain

(III.15)

$$\begin{aligned} \|G\|_{L^2(B^4)} + \|\nabla G\|_{L^{4/3,1}(B^4)} &\leq C \|dG\|_{L^{4/3,1}(B^4)} + \|d^*G\|_{L^{4/3,1}(B^4)} \\ &\leq \|A\|_{W^{1,2}(B^4)} \|F\|_{L^2(B^4)}. \end{aligned}$$

Since $F - G$ is an harmonic form on B^4 with an L^2 control on F and an L^2 control on G we get

$$(III.16) \quad \|\nabla(F - G)\|_{L^{4/3,1}(B_{1/2}^4)} \leq C (\|A\|_{W^{1,2}(B^4)} + 1) \|F\|_{L^2(B^4)}.$$

Combining (III.15) and (III.16) we obtain

$$(III.17) \quad \|\nabla F\|_{L^{4/3,1}(B_{1/2}^4)} \leq C (\|A\|_{W^{1,2}(B^4)} + 1) \|F\|_{L^2(B^4)}.$$

Using now the Lorentz-Sobolev embedding of J. Peetre (see [18] Theorem 8) we get the desired estimate on the $L^{2,1}$ norm of the curvature and Lemma III.2 is proved. \square

We show how to adapt the arguments above to the present situation: we have only a gauge in $B_1^4 \setminus B_{R\delta_i}^4$ in which we control the $W^{1,2}$ norm of the Gauge independently of i and ε . Let χ be a cut-off function equal to 1 on $B_1 \setminus B_{2R\delta_i}$ and 0 on $B_{R\delta_i}$, such that $\|\nabla\chi\|_\infty \leq C/R\delta_i$. Let's take F the curvature in the Gauge given by Lemma III.1 and consider χF . It solves

$$d(\chi F) = -\chi[A, F] + d\chi \wedge F \quad \text{and} \quad d(*(\chi F)) = -\chi[A, *F] + d\chi \wedge *F.$$

We use the arguments above to deduce that $\|\chi[A, F]\|_{L^{\frac{4}{3},1}(B^4)} \leq C$ and similarly $\|\chi[A, *F]\|_{L^{\frac{4}{3},1}(B^4)} \leq C$. From the L^∞ estimates established for the curvature in the previous part we deduce that

$$|d\chi \wedge *F| \leq \frac{C}{(R\delta_i)^3} \quad \text{and} \quad |d\chi \wedge F| \leq \frac{C}{(R\delta_i)^3}$$

and this terms are supported in the annulus $B_{2R\delta_i} \setminus B_{R\delta_i}$ and using the definition of the $L^{\frac{4}{3},1}$ norm we easily get that

$$\|d\chi \wedge *F\|_{L^{\frac{4}{3},1}(B^4)} \leq C \quad \text{and} \quad \|d\chi \wedge F\|_{L^{\frac{4}{3},1}(B^4)} \leq C$$

where C is independent on i and ε . So arguing as in the previous lemma we have established that

$$(III.18) \quad \|F\|_{L^{2,1}(B_1^4 \setminus B_{R\delta_i}^4)} \leq C$$

where C is independent on i and ε and the proof of Theorem I.1 follows from the arguments of section 2. \square

IV. Some more applications of Lorentz Spaces to the analysis of non-linearities issued from curvatures.

In this part, which is independent from the rest of the paper, we aim to present how Lorentz spaces are helpful to establish properties relative to strong converging sequences of connections of a given bundle over a riemannian manifold. From now on the Lie group G is a subgroup of $SO(k)$ seen as a submanifold of the vector space of k by k matrices.

We first establish the following result:

Theorem IV.1. *Let Q be a principal G -bundle over a compact n -dimensional riemannian manifold M . Assume there exists on Q a sequence of connections D_i , a fixed covering of M (\mathcal{U}^α) and a trivialization σ_i^α over \mathcal{U}^α such that if $A_i^\alpha = \sigma_i^\alpha D_i (\sigma_i^\alpha)^{-1} - d$ is a Coulomb Gauge for D_i ($d^* A_i^\alpha = 0$) that converges strongly to some A_∞^α in $W^{1,n/2}$, then A_∞^α defines a connection of the same bundle Q : there exists a connection D_∞ of Q and a system of trivialization σ_∞^α such that*

$$A_\infty^\alpha = \sigma_\infty^\alpha D_\infty (\sigma_\infty^\alpha)^{-1} - d$$

where d denotes the exterior differentiation.

Remark IV.1. In other words we prove that the strong convergence of coulomb gauges of connections in $W^{1,n/2}$ preserves the topology of the bundle. It was well known (see [6]) that weak convergence of connections in $W^{1,n/2}$ does not preserve the topology of the bundle. It is straightforward to check that strong convergences of connections in $W^{1,p}$ for $p > n/2$ preserves the topology. It is well known that strong convergence in $W^{1,p}$ for $p < n/2$ does not preserve the topology of the bundle (Ex: the bubbling phenomenon for ASD connection in 4-dimension). So the theorem above solves a borderline case.

Proof of Theorem IV.1. Denote $g_i^{\alpha,\beta}$ the sequences of transition function from $U_\alpha \cap U_\beta$ into G . The transition functions solve

$$(IV.1) \quad dg_i^{\alpha,\beta} = g_i^{\alpha,\beta} A_i^\alpha - A_i^\beta g_i^{\alpha,\beta}.$$

From Sobolev embedding we deduce that

$$(IV.2) \quad \int_{U^{\alpha,\beta}} |dg_i^{\alpha,\beta}|^n \leq C$$

where $U^{\alpha,\beta} = U^\alpha \cap U^\beta$. Let $p < n$, for any $i, j \in \mathbb{N}$ we have

$$(IV.3) \quad \int_{U^{\alpha,\beta}} |dg_i^{\alpha,\beta} - dg_j^{\alpha,\beta}|^p \leq \int_{U^{\alpha,\beta}} |A_i^\alpha - A_j^\alpha|^p + \int_{U^{\alpha,\beta}} |A_i^\beta - A_j^\beta|^p + \left(\int_{U^{\alpha,\beta}} |A_i^\alpha|^n + |A_i^\beta|^n + |A_j^\alpha|^n + |A_j^\beta|^n \right)^{\frac{p}{n}} \left(\int_{U^{\alpha,\beta}} |g_i^{\alpha,\beta} - g_j^{\alpha,\beta}|^{\frac{np}{n-p}} \right)^{\frac{n-p}{n}}.$$

From (IV.2) there exists a subsequence (still denoted $g_i^{\alpha,\beta}$) such that $g_i^{\alpha,\beta} \rightarrow g_\infty^{\alpha,\beta}$ in $L^q(U^{\alpha,\beta})$ for any $q < +\infty$, thus from (IV.3) we deduce that $g_i^{\alpha,\beta} \rightarrow g_\infty^{\alpha,\beta}$ in $W^{1,p}(U^{\alpha,\beta})$ for any $p < n$. Consider a strict subcover of U^α denoted \mathcal{V}^α with $\overline{\mathcal{V}^\alpha} \cap \overline{\mathcal{V}^\beta} \subset U^{\alpha,\beta}$. Let $\delta > 0$, there exists ρ_δ such that for any $\rho < \rho_\delta$, $\max_{z,\alpha} \|A^\alpha\|_{W^{1,n/2}(B_\rho(z))} \leq (\frac{\delta}{2})^{1/2}$. We can cover $\mathcal{V}^{\alpha,\beta}$ by a fixed union of balls $B_\rho(z)$ included in $U^{\alpha,\beta}$ such that $\rho < \rho_\delta$ and such that, applying Fubini Theorem, modulo extraction of a subsequence

$$(IV.4) \quad \int_{\partial B_\rho(z)} |\nabla(g_i^{\alpha,\beta} - g_\infty^{\alpha,\beta})|^{n-\frac{1}{2}} \rightarrow 0.$$

On any such a $B_{2\rho}(z)$ from (IV.1) and the Coulomb gauge condition we have

$$(IV.5) \quad -\Delta g_i^{\alpha,\beta} = * [dg_i^{\alpha,\beta} \wedge * A_i^\alpha] + * [* A_i^\beta \wedge dg_i^{\alpha,\beta}].$$

Since $g_i^{\alpha,\beta} \in W^{2,n/2}(U^{\alpha,\beta})$ (from (IV.1)) using the Peetre-Lorentz-Sobolev embedding presented in [18] Theorem 8, we have $g_i^{\alpha,\beta} \in L^{n,n/2}$. Using now the multiplication property of functions in Lorentz Spaces given in [18]: the product of 2 functions in $L^{n,n/2}$ is in $L^{n/2,n/4}$. Thus, from (IV.5) again, we deduce that $\Delta g_i^{\alpha,\beta}$ is bounded in $L^{n/2,n/4}(B_{2\rho}(z))$, reducing a bit the size of the ball (from 2ρ to $3\rho/2$), using the weak (r, r) property of Riesz operator and Theorem 3.15 of Chap. V of [17], we have that $\|\nabla^2 g_i^{\alpha,\beta}\|_{L^{n/2,n/4}}$ is

bounded independently on i . Bootstrapping this information again in (IV.5) we get that $\Delta g_i^{\alpha,\beta}$ is bounded in $L^{n/2, n/6}(B_{2\rho}(z))\dots$ and so on until the second exponent reaches $\frac{n}{n-2}$. So we deduce that

$$(IV.6) \quad \|\nabla g_i^{\alpha,\beta}\|_{L^{n, \frac{n}{n-2}}(B_\rho(z))} \leq C$$

where C is independent on i . We have (all norms above being taken over $B_\rho(z)$)

$$(IV.7) \quad \begin{aligned} & \|\Delta(g_i^{\alpha,\beta} - g_\infty^{\alpha,\beta})\|_{L^{\frac{n}{2}, 1}} \\ & \leq \|\nabla(g_i^{\alpha,\beta} - g_\infty^{\alpha,\beta})\|_{L^{n, n/n-2}} \left[\|A_i^\alpha\|_{L^{n, \frac{n}{2}}(B_\rho(z))} + \|A_i^\beta\|_{L^{n, \frac{n}{2}}} \right] \\ & + \left[\|A_i^\alpha - A_\infty^\alpha\|_{L^{n, \frac{n}{2}}} + \|A_i^\beta - A_\infty^\beta\|_{L^{n, \frac{n}{2}}} \right] \left[\|\nabla g_i^{\alpha,\beta}\|_{L^{n, \frac{n}{n-2}}} + \|\nabla g_\infty^{\alpha,\beta}\|_{L^{n, \frac{n}{n-2}}} \right]. \end{aligned}$$

Since we have $\int_{B_\rho(z)} |A_i^\alpha|^{\frac{n}{2}} + |\nabla A_i^\alpha|^{\frac{n}{2}} \leq \delta$, we deduce that $\int_{B_\rho(z)} |A_i^\alpha|^n \leq (\delta^{2/n} + \rho^{n-2}\delta^{2/n})^n \leq C\delta^2$ where C is a universal constant (we may always assume that $\rho < 1$). Using the embedding of $W^{1, n/2}(B_\rho)$ into $L^{n, n/2}(B_\rho)$, taking into account the two previous estimates and the common scaling behavior of the norms L^n , $L^{n, n/2}$ and the pseudo-norm: gradient in $L^{n/2}$ in n dimension, we have

$$(IV.8) \quad \|A_i^\alpha\|_{L^{n, n/2}(B_\rho)} \leq C\delta^2.$$

Let h solving

$$(IV.9) \quad \begin{cases} \Delta h = \Delta(g_i^{\alpha,\beta} - g_\infty^{\alpha,\beta}) & \text{in } B_\rho(z) \\ h = 0 & \text{on } \partial B_\rho(z). \end{cases}$$

Standard result on Riesz Kernel gives that the operator that assign $\nabla^2 h$ to $\Delta(g_i^{\alpha,\beta} - g_\infty^{\alpha,\beta})$ is a weak (r, r) operator for r between 0 and ∞ . Thus we have

$$(IV.10) \quad \begin{aligned} \|\nabla h\|_{L^{n, \frac{n}{n-2}}(B_\rho)} & \leq \|\nabla h\|_{L^{n, 1}(B_\rho)} + \|\nabla^2 h\|_{L^{n/2, 1}(B_\rho)} \\ & \leq C\|\Delta(g_i^{\alpha,\beta} - g_\infty^{\alpha,\beta})\|_{L^{n/2, 1}(B_\rho)} \end{aligned}$$

where C is independent on ρ . Let now k solving

$$(IV.11) \quad \begin{cases} \Delta k = 0 & \text{in } B_\rho(z) \\ k = g_i^{\alpha,\beta} - g_\infty^{\alpha,\beta} & \text{on } \partial B_\rho(z). \end{cases}$$

From standard elliptic estimates we have

$$(IV.12) \quad \|k\|_{W^{1-\frac{1}{s},s}(\partial B_\rho(z))} \leq C \|g_i^{\alpha,\beta} - g_\infty^{\alpha,\beta}\|_{W^{1,n-\frac{1}{2}}(\partial B_\rho(z))}$$

for $s = n \left(\frac{2n-1}{2(n-1)} \right) > n$. Thus we deduce

$$(IV.13) \quad \|\nabla k\|_{L^{n,\frac{n}{n-2}}(B_\rho(z))} \leq C_\rho \|g_i^{\alpha,\beta} - g_\infty^{\alpha,\beta}\|_{W^{1,n-\frac{1}{2}}(\partial B_\rho(z))}.$$

Combining (IV.7)...(IV.13)

$$(IV.14) \quad \begin{aligned} & \|\nabla(g_i^{\alpha,\beta} - g_\infty^{\alpha,\beta})\|_{L^{n,\frac{n}{n-2}}(B_\rho(z))} \\ & \leq C\delta^2 \|\nabla(g_i^{\alpha,\beta} - g_\infty^{\alpha,\beta})\|_{L^{n,\frac{n}{n-2}}(B_\rho(z))} \\ & \quad + C \left[\|A_i^\alpha - A_\infty^\alpha\|_{L^{n,n/2}(B_\rho(z))} + \|A_i^\beta - A_\infty^\beta\|_{L^{n,n/2}(B_\rho(z))} \right] \\ & \quad + C_\rho \|g_i^{\alpha,\beta} - g_\infty^{\alpha,\beta}\|_{W^{1,n-\frac{1}{2}}(\partial B_\rho(z))} \end{aligned}$$

and from (IV.4) and the embedding of $W^{1,n/2}$ into $L^{n,n/2}$ we deduce that $A_i^\alpha \rightarrow A_\infty^\alpha$ in $L^{n,\frac{n}{n-2}}$, thus

$$(IV.15) \quad \|\nabla(g_i^{\alpha,\beta} - g_\infty^{\alpha,\beta})\|_{L^{n,\frac{n}{n-2}}(B_\rho(z))} \rightarrow 0.$$

Bootstrapping this information into (IV.7) we get

$$(IV.16) \quad \|\Delta(g_i^{\alpha,\beta} - g_\infty^{\alpha,\beta})\|_{L^{n/2,1}(B_\rho(z))} \rightarrow 0$$

thus for any $\lambda < 1$ we have, using again the weak (r, r) property of the Riesz operator,

$$\|\nabla^2(g_i^{\alpha,\beta} - g_\infty^{\alpha,\beta})\|_{L^{n/2,1}(B_{\lambda\rho}(z))} \rightarrow 0.$$

Using now a Lorentz-Sobolev embedding (Theorem 8 [18]) we deduce

$$(IV.17) \quad \|\nabla(g_i^{\alpha,\beta} - g_\infty^{\alpha,\beta})\|_{L^{n,1}(B_{\lambda\rho}(z))} \rightarrow 0.$$

At this stage of the proof we are going to use the key embedding established by H. Brezis and S. Waigner [3] which says that a function on \mathbb{R}^n whose derivatives are in $L^{n,1}$ is in L^∞ . This can be shown in the following way: assume $\nabla f \in L^{n,1}(\mathbb{R}^n)$, we have $f(x) = \operatorname{div} \int_{\mathbb{R}^n} G(x-y) f(y)$ where G is the Kernel of the Laplacian on \mathbb{R}^n . Then since $|\nabla G(x)| \leq C/|x|^{n-1}$ we have $|\{x ; |\nabla G|(x) \geq \lambda\}| \leq C/\lambda^{-n/n-1}$ which means that the weak $L^{n/n-1}$ -norm of the derivatives of G is bounded: $\|\nabla G\|_{L^{\frac{n}{n-1},\infty}(\mathbb{R}^n)} \leq C$. The claim follows then from the duality $L^{n,1} - L^{\frac{n}{n-1},\infty}$. Thus, granting this embedding and inequality (IV.17), we deduce, since the balls B_ρ cover $\mathcal{V}^\alpha \cap \mathcal{V}^\beta$,

$$\|g_i^{\alpha,\beta} - g_\infty^{\alpha,\beta}\|_{L^\infty(\mathcal{V}^\alpha \cap \mathcal{V}^\beta)} \rightarrow 0.$$

Using now Proposition 3.2 of [21] we deduce that, for i large enough, $g_i^{\alpha,\beta}$ and $g_\infty^{\alpha,\beta}$ are equivalent cocycle for the Čech cohomology (see [4]), in other words there exist trivialisations ρ^α over sets (eventually slightly smaller than the \mathcal{V}^α) such that $(g_i^{\alpha,\beta})^{-1}g_\infty^{\alpha,\beta} = \rho^\alpha (\rho^\beta)^{-1}$. The equivalence of the cocycles imply that they define topologically the same bundles and Theorem IV.1 is proved. □

If we do not assume anymore that we are working with Coulomb gauges we have the following result that generalises the previous theorem.

Theorem IV.2. *Let Q be a principal G -bundle over a compact n dimensional riemannian manifold M . Assume there exists on Q a sequence of connections D_i , a fixed covering of M (\mathcal{U}^α) and a trivialization σ_i^α over \mathcal{U}^α such that if $A_i^\alpha = \sigma_i^\alpha D_i (\sigma_i^\alpha)^{-1} - d$ converges strongly to some A_∞^α in $W^{1,n/2}$ and if $d^* A_i^\alpha$ strongly converges to $d^* A_\infty^\alpha$ in the Lorentz space $L^{n/2,1}$, then A_∞^α defines a connection of the same bundle Q : there exists a connection D_∞ of Q and a system of trivialization σ_∞^α such that*

$$A_\infty^\alpha = \sigma_\infty^\alpha D_i (\sigma_\infty^\alpha)^{-1} - d$$

where d denotes the exterior differentiation.

Remark IV.2. In the particular cases where the topology of the bundle is given by the second Chern class only, for instance for $G = SU(2)$ over an orientable 4-manifold, the result above does not require anymore $d^* A_i$ to converge in $L^{2,1}$ but only in $L^{2,2}$ (we only assume that A_i converges in $W^{1,2}$). This is a consequence of the integral formula for the Chern class (see [22]). To know whether the strong $L^{n/2,1}$ convergence of $d^* A_i^\alpha$ is really needed in the statement of Theorem IV.2, remains an open question.

Proof of Theorem IV.2. The goal is to find a change of trivialisation in which we fulfill the assumption of Theorem IV.1: strong $W^{1,n/2}$ convergence of the new A_i^α that will be Coulomb $d^*A_i^\alpha = 0$.

We can assume that the U^α are diffeomorphic to B^n and that the $W^{1,n/2}$ norm of A_∞^α on U^α is below the range κ where Uhlenbeck's theorems in [21] apply. First of all we show that we can assume that $d^*A_i^\alpha \rightarrow 0$ in $L^{n/2,1}(U^\alpha)$. Indeed let σ^α be the change of gauge for A_∞^α given by Corollary 2.2 in [21]: we have that $\hat{A}_\infty^\alpha = (\sigma^\alpha)^{-1}A_\infty^\alpha\sigma^\alpha + (\sigma^\alpha)^{-1}d\sigma^\alpha$ verifies

$$(IV.18) \quad d^*\hat{A}_\infty^\alpha = 0 \quad \text{and} \quad \|\hat{A}_\infty^\alpha\|_{W^{1,n/2}(U^\alpha)} \leq C\kappa$$

since $d^*\hat{A}_\infty^\alpha = 0$, σ^α solves

$$(IV.19) \quad \Delta\sigma^\alpha = *[d\sigma^\alpha \wedge *\hat{A}_\infty^\alpha] - d^*A_\infty^\alpha\sigma^\alpha + *[*A_\infty^\alpha \wedge d\sigma^\alpha].$$

Argueing as in the proof of the previous theorem, we have $\|A_\infty^\alpha\|_{L^{n,n/2}} \leq C$, $\|\hat{A}_\infty^\alpha\|_{L^{n,n/2}} \leq C$ and thus $\|d\sigma^\alpha\|_{L^{n,n/2}} \leq C$, using the hypothesis of the theorem and multiplication properties in the Lorentz spaces mentioned above we have

$$(IV.20) \quad \|\Delta\sigma^\alpha\|_{L^{n/2,n/4}(U^\alpha)} \leq C.$$

Using the weak (r,r) property of the Riesz operator and Theorem 3.15 Chapter V of [17], we deduce that for any $\omega \subset\subset U^\alpha$

$$(IV.21) \quad \|\nabla^2\sigma^\alpha\|_{L^{n/2,n/4}(\omega)} \leq C.$$

Having taken a slightly larger ball than U^α from the begining, we may always assume that the inequality above holds for $\omega = U^\alpha$. Then we deduce that $\|nabla\sigma^\alpha\|_{L^{n,n/4}(U^\alpha)} \leq C$ and as in the proof of Theorem IV.1 we keep bootstrapping this information in (IV.19) untill reaching

$$(IV.22) \quad \|\nabla^2\sigma^\alpha\|_{L^{n/2,1}(U^\alpha)} \leq C.$$

We then proceed to the change of gauge given by σ^α for the complete sequence A_i^α and we denote $\hat{A}_i^\alpha = (\sigma^\alpha)^{-1}A_i^\alpha\sigma^\alpha + (\sigma^\alpha)^{-1}d\sigma^\alpha$ clearly we have that the hypothesis of the theorem are preserved and that $d^*\hat{A}_i^\alpha \rightarrow 0$ in $L^{n,1}$. At this stage of the proof we will need the corresponding lemma to Lemma 2.7 of [21] but with Lorentz norms: Denote

$$\tilde{W}_1^{1,n/2}(\Lambda^1 B^n, \mathfrak{G}) = \left\{ a \in W^{1,n/2}(\Lambda^1 B^n, \mathfrak{G}); d^*a \in L^{n/2,1} \right\},$$

$$W_{1,0}^{2,n/2}(B^n, G) = \left\{ s \in W^{2,n/2}(B^n, G); \nabla^2 s \in L^{n/2,1} \text{ and } s = \text{id on } \partial B^n \right\}.$$

Using the above notations we have

Lemma IV.1. *There exists $\delta > 0$ such that if A in $W^{1,n/2}(\wedge^1 B^n; \mathfrak{G})$ verifying $d^*A = 0$ and $\|A\|_{W^{1,n/2}} \leq \delta$, then there exists $\varepsilon > 0$ such that for any $a \in \tilde{W}_{1,\nu}^{1,n/2}(\wedge^1 B^n, \mathfrak{G})$ verifying $\|a\|_{\tilde{W}_1^{1,2}} \leq \varepsilon$, the non linear equation*

$$d^*(s^{-1}ds + s^{-1}(A + a)s) = 0$$

has a solution s in $W_{1,0}^{2,n/2}(B^n, G)$ that depends smoothly on $a \in \tilde{W}_1^{1,n/2}$.

Proof of Lemma IV.1. Consider the spaces

$$\hat{W}_{1,0}^{2,n/2}(B^n, \mathfrak{G}) = \left\{ U \in W^{2,n/2}(B^n, G); \nabla^2 U \in L^{n/2,1}, U = 0 \text{ on } \partial B^n \right\}.$$

Using Brezis-Wainger-Sobolev embedding we have

$$(IV.23) \quad \|U\|_{L^\infty(B^n)} \leq C \|U\|_{\hat{W}_{1,0}^{2,n/2}(B^n)}.$$

This is the key point that permits to adapt Uhlenbeck arguments to the present situation and the $L^{n/2,1}$ assumption on the second derivatives is crucial at this stage. From (IV.23) we deduce that the maps $U \rightarrow \exp(U)$ or $\exp(-U)$ from $\hat{W}_{1,0}^{2,n/2}$ into L^∞ are smooth. Thus the map

$$\hat{W}_{1,0}^{2,n/2} \otimes \tilde{W}_1^{1,n/2} \rightarrow L^{n/2,1}$$

$$(U, a) \longrightarrow d^* [\exp(-U)d\exp(U) + \exp(-U)(A + a)\exp(U)]$$

is smooth (we also use that $d^*A = 0$, and again that for U in $\hat{W}_{1,0}^{2,n/2}$ and a in $\tilde{W}_1^{1,n/2}$, $dU \in L^{n, \frac{n}{n-2}}$ and $a \in L^{n,n/2}$ too - we even have $dU \in L^{n,1}$). At $(U, \lambda) = (0, 0)$ we consider the linearized operator

$$H(\psi) = d^*d\psi + *([*A, d\psi]).$$

Because of the boundary condition $U = 0$ on ∂B^n , $d^*d\psi = \Delta\psi$ is invertible from $\hat{W}_{1,0}^{2,n/2}$ into $L^{n/2,1}$, where we use again that Riesz operator is weak (r, r) and Theorem 3.15 Chapter V of [17]. Thus we have

$$\|H(\psi)\|_{L^{n/2,1}} \geq C_1 \|\psi\|_{\hat{W}_{1,0}^{2,2}} - C_2 \|A\|_{L^{n,n/2}} \|\psi\|_{\hat{W}_{1,0}^{2,2}}.$$

Thus choosing $\|A\|_{W^{1,n/2}}$ small enough makes $\|A\|_{L^{n,n/2}}$ small, which makes H invertible and applying the local inversion theorem we establish Lemma IV.1. □

End of the proof of Theorem IV.2. We can apply Lemma IV.18 above to $A = \hat{A}_\infty^\alpha$ and $a = \hat{A}_i^\alpha - \hat{A}_\infty^\alpha$ for i large enough such that $\|\hat{A}_i^\alpha - \hat{A}_\infty^\alpha\|_{\dot{W}_1^{1,n/2}} \leq \varepsilon$ and we generate a last change of gauge that put us in the setting of Theorem IV.1 above that we can apply and Theorem IV.2 is proved. \square

References.

- [1] W. Allard, *An integrality theorem and a regularity theorem for surfaces whose first variation with respect to a parametric elliptic integrand is controlled*, Proc. Symp. Pure Math., **44** (1986), 1-28.
- [2] H. Brezis, *Laser beams and limiting cases of Sobolev inequalities*, Non linear P.D.E., College de France Seminar, **II**, H. Brezis & J.L. Lions, eds., Pitman, 1982.
- [3] H. Brezis and S. Wainger, *A note on limiting cases of Sobolev embeddings and convolution inequalities*, Comm. P.D.E., **5** (1980), 773-789.
- [4] R. Bott and L. Tu, *Differential forms in Algebraic Topology*, Springer, 1982.
- [5] W.Y. Ding and G. Tian, *Energy identity for a class of approximate harmonic maps from surfaces*, Comm. Anal. Geom., **3** (1995), 543-554.
- [6] S.K. Donaldson and P.B. Kronheimer, *The geometry of Four-Manifolds*, Oxford, 1990.
- [7] S.K. Donaldson and R.P. Thomas, *Gauge Theory in higher dimensions*, in 'The geometric Universe' (Oxford, 1996), Oxford Univ. Press, (1998), 31-47.
- [8] F. Hélein, *Harmonic maps, Conservation laws and moving frames*, Diderot, 1996.
- [9] F.H. Lin, *Gradient estimates and blow-up analysis for stationary harmonic maps*, Ann. Math., **149** (1999), 785-829.
- [10] F.H. Lin and T. Rivière, *A Quantization Property for Static Ginzburg-Landau Vortices*, Comm. Pure App. Math., **54** (2001), 206-208.
- [11] F.H. Lin and T. Rivière, *Energy Quantization for Harmonic Maps*, Duke Math. Journal, 2002.

- [12] F.H. Lin and T. Rivière, *A quantization property for moving line vortices*, Comm. Pure App. Math., **54** (2001), 826-850.
- [13] Y. Meyer and T. Rivière, *A partial regularity result for a class of stationary Yang-Mills Fields in high dimensions*, to appear in Rev. Mat. Iberoamericana, 20002.
- [14] H. Nakajima, *Compactness of the moduli space of Yang-Mills connections in higher dimensions*, J. Math. Soc. Japan, **40** (1988), 383-392.
- [15] J. Peetre, *Espaces d'interpolation et théorème de Soboleff*, Ann. Inst. Fourier, Grenoble, **16** (1966), 279-317.
- [16] T. Parker, *Bubble tree convergence for harmonic maps*, J. Differential Geom., **44** (1996), 545-633.
- [17] E. Stein and G. Weiss, *Introduction to Fourier analysis on Euclidian spaces*, Princeton Univ. Press, 1971.
- [18] L. Tartar, *Imbedding Theorems of Sobolev Spaces into Lorentz Spaces*, Boll. U.M.I., **1B** (1998) 479-500.
- [19] G. Tian, *Gauge theory and calibrated geometry, I*, Ann. Math., **151** (2000), 193-268.
- [20] K. Uhlenbeck, *Removable singularities in Yang-Mills fields*, Comm. Math. Phys., **83** (1982), 11-29.
- [21] K.Uhlenbeck, *Connections with L^p bounds on curvature*, Comm. Math. Phys., **83** (1982), 31-42.
- [22] K.Uhlenbeck, *The Chern classes of Sobolev connections*, Comm. Math. Phys., **101** (1985), 449-457.

D-MATH
ETH-ZENTRUM
8092 ZÜRICH
SWITZERLAND

RECEIVED DECEMBER 12, 2000.