

Left-orderable, non- L -space surgeries on knots

KIMIHIKO MOTEKI AND MASAKAZU TERAGAITO

Let K be a knot in the 3-sphere S^3 . An r -surgery on K is *left-orderable* if the resulting 3-manifold $K(r)$ of the surgery has left-orderable fundamental group, and an r -surgery on K is called an *L -space surgery* if $K(r)$ is an L -space. A conjecture of Boyer, Gordon and Watson says that non-reducing surgeries on K can be classified into left-orderable surgeries or L -space surgeries. We introduce a way to provide knots with left-orderable, non- L -space surgeries. As an application we present infinitely many hyperbolic knots on each of which every non-trivial surgery is a hyperbolic, left-orderable, non- L -space surgery.

In honor of Dale Rolfsen

1. Introduction

A non-trivial group G is said to be *left-orderable* if there exists a strict total ordering $<$ on its elements such that $g < h$ implies $fg < fh$ for all elements $f, g, h \in G$. The left-orderability of fundamental groups of 3-manifolds has been studied by Boyer, Rolfsen and Wiest [5]. In particular, they prove that the fundamental group of a P^2 -irreducible 3-manifold is left-orderable if and only if it has an epimorphism to a left-orderable group [5, Theorem 1.1(1)]. Since the infinite cyclic group \mathbb{Z} is left-orderable, a P^2 -irreducible 3-manifold with first Betti number $b_1 \geq 1$ has left-orderable fundamental group. One obstruction for G being left-orderable is an existence of torsion elements in G . Thus, for instance, lens spaces cannot have left-orderable fundamental groups. It is interesting to characterize rational homology 3-spheres whose fundamental groups are left-orderable. Examples suggest that there exists a correspondence between rational homology 3-spheres whose fundamental groups cannot be left-ordered and L -spaces, which appear in the Heegaard Floer homology theory [45, 46]. For a rational homology 3-sphere M , we have $\widehat{\text{rkHF}}(M) \geq |H_1(M; \mathbb{Z})|$. If the equality holds, then M is called an *L -space*.

Following [4, 1.1], for homogeneity, we use \mathbb{Z}_2 -coefficients for Heegaard Floer homology.

The present paper is motivated by the following conjecture formulated by Boyer, Gordon and Watson [4].

Conjecture 1.1 [4]. *An irreducible rational homology 3-sphere is an L -space if and only if its fundamental group is not left-orderable.*

In [4] the conjecture is verified for geometric, non-hyperbolic 3-manifolds and the 2-fold branched covers of non-splitting alternating links. See also [2, 8, 19, 27, 49] for related results.

A useful way to construct rational homology 3-spheres is Dehn surgery on knots in the 3-sphere S^3 . Henceforth, we will focus on Conjecture 1.1 for rational homology 3-spheres obtained by Dehn surgery on knots in S^3 . For any knot K in S^3 the exterior $E(K) = S^3 - \text{int}N(K)$ has left-orderable fundamental group [5, Corollary 3.5]. However, the result $K(r)$ of r -Dehn surgery may not have such a fundamental group; see Examples 1.6, [11] and [30].

A Dehn surgery is said to be *left-orderable* if the resulting 3-manifold of the surgery has left-orderable fundamental group. Define the set of left-orderable surgeries on K as

$$\mathcal{S}_{\text{LO}}(K) = \{r \in \mathbb{Q} \mid \pi_1(K(r)) \text{ is left-orderable}\}.$$

Similarly a Dehn surgery is called an *L -space surgery* if the resulting 3-manifold of the surgery is an L -space, and the set of L -space surgeries on K is defined as

$$\mathcal{S}_L(K) = \{r \in \mathbb{Q} \mid K(r) \text{ is an } L\text{-space}\}.$$

Remark 1.2. (1) Note that 0-surgery does not yield a rational homology 3-sphere, and hence $K(0)$ is not an L -space and $0 \notin \mathcal{S}_L(K)$. On the other hand, if K is a trivial knot, then $K(0) \cong S^2 \times S^1$ which has left orderable fundamental group. If K is a non-trivial knot, then $K(0)$ is irreducible [15, Corollary 8.3] and $H_1(K(0)) \cong \mathbb{Z}$, hence $0 \in \mathcal{S}_{\text{LO}}(K)$ [5, Theorem 1.1].

(2) Let K^* be the mirror image of a knot K , and put $-\mathcal{S} = \{-r \mid r \in \mathcal{S}\}$ for $\mathcal{S} \subset \mathbb{Q}$. Since $K^*(-r)$ is orientation reversingly diffeomorphic to

$K(r)$ and the conditions of a 3-manifold M having left-orderable fundamental group and being an L -space are independent of the orientation of M [47, p. 1288], we have $\mathcal{S}_{\text{LO}}(K^*) = -\mathcal{S}_{\text{LO}}(K)$ and $\mathcal{S}_L(K^*) = -\mathcal{S}_L(K)$.

If $K(r)$ is a reducible 3-manifold for a non-trivial knot K , it has a lens space summand [18, Theorem 3], hence $r \notin \mathcal{S}_{\text{LO}}(K)$, but r may or may not be in $\mathcal{S}_L(K)$; see Remark 1.4 and Example 1.6.

If $K(r)$ is irreducible, Conjecture 1.1 asserts that r belongs to exactly one of $\mathcal{S}_{\text{LO}}(K)$ and $\mathcal{S}_L(K)$. Taking the cabling conjecture [16] into consideration, Conjecture 1.1 suggests:

Conjecture 1.3. *Let K be a knot in S^3 which is not a cable of a non-trivial knot. Then $\mathcal{S}_{\text{LO}}(K) \cup \mathcal{S}_L(K) = \mathbb{Q}$ and $\mathcal{S}_{\text{LO}}(K) \cap \mathcal{S}_L(K) = \emptyset$.*

Remark 1.4. The cabling conjecture [16] asserts that if $K(r)$ is reducible for a non-trivial knot K , then K is cabled and r is a cabling slope. There exists a cable knot K for which $\mathcal{S}_{\text{LO}}(K) \cup \mathcal{S}_L(K) \neq \mathbb{Q}$. For instance, let K be a (p, q) cable of a non-fibered knot k ($q > 0$). Then $K(pq) = k(\frac{p}{q})\sharp L(q, p)$ [17, Corollary 7.3]. Since $\pi_1(K(pq))$ has a torsion, $pq \notin \mathcal{S}_{\text{LO}}(K)$. To see that $pq \notin \mathcal{S}_L(K)$, we note that $\widehat{HF}(K(pq)) \cong \widehat{HF}(k(\frac{p}{q})) \otimes \widehat{HF}(L(q, p))$; see [53, 8.1(5)] ([46]). Since k is a non-fibered knot, $k(\frac{p}{q})$ is not an L -space [43, 44]. Hence the rank of $\widehat{HF}(K(pq))$ is strictly bigger than $|p|q$, and $K(pq)$ is not an L -space. It follows that $pq \notin \mathcal{S}_{\text{LO}}(K) \cup \mathcal{S}_L(K)$.

For the trivial knot and non-trivial torus knots, Examples 1.5 and 1.6 describe $\mathcal{S}_{\text{LO}}(K)$ and $\mathcal{S}_L(K)$ explicitly. Note that these knots satisfy Conjecture 1.3.

Example 1.5 trivial knot. Let K be the trivial knot in S^3 . Then $\mathcal{S}_{\text{LO}}(K) = \{0\}$ and $\mathcal{S}_L(K) = \mathbb{Q} - \{0\}$.

Example 1.6 torus knots. For a non-trivial torus knot $T_{p,q}$ ($p > q \geq 2$), the argument in the proof of [10, Theorem 1.4] shows that $\mathcal{S}_{\text{LO}}(T_{p,q}) = (-\infty, pq - p - q) \cap \mathbb{Q}$ and $\mathcal{S}_L(T_{p,q}) = [pq - p - q, \infty) \cap \mathbb{Q}$.

Example 1.7 figure-eight knot. Let K be the figure-eight knot. Following [47, 48], $\mathcal{S}_L(K) = \emptyset$. Thus, it is expected that $\mathcal{S}_{\text{LO}}(K) = \mathbb{Q}$. Boyer, Gordon and Watson [4] show that $\mathcal{S}_{\text{LO}}(K) \supset (-4, 4) \cap \mathbb{Q}$, and Clay, Lidman and Watson [8] improve that $\mathcal{S}_{\text{LO}}(K) \supset [-4, 4] \cap \mathbb{Q}$. Furthermore, Fenley [14] implies that $\mathcal{S}_{\text{LO}}(K) \supset \mathbb{Z}$.

For related results, see [9, 11, 20, 33, 54, 56].

It is known that there exist some constraints for knots which admit L -space surgeries. For instance, such knots have specific Alexander polynomials [47], and must be fibered [43, 44]. Thus, generically we have $\mathcal{S}_L(K) = \emptyset$. Hence, Conjecture 1.3 suggests that $\mathcal{S}_{LO}(K) = \mathbb{Q}$ for most knots. Despite being expected, there is no literature giving explicitly knots with $\mathcal{S}_{LO}(K) = \mathbb{Q}$ and $\mathcal{S}_L(K) = \emptyset$. In the present note we give infinitely many satellite knots and hyperbolic knots with this property.

Theorem 1.8. *Given a non-trivial knot K' , there are infinitely many prime satellite knots K each of which has K' as a companion knot and enjoys the following properties:*

- (1) $K(r)$ is a toroidal 3-manifold which is not a graph manifold for all but finitely many $r \in \mathbb{Q}$.
- (2) $\mathcal{S}_{LO}(K) = \mathbb{Q}$.
- (3) $\mathcal{S}_L(K) = \emptyset$.

This is an application of Proposition 7.1 due to Clay and Watson [10, Proposition 4.1]. In Theorem 1.8, K is satellite knot and the resulting 3-manifold $K(r)$ has a non-trivial Jaco–Shalen–Johannson (JSJ) decomposition [28, 29]. Since Proposition 7.1 does not work for creation of hyperbolic knots, we will introduce an effective way to provide infinitely many hyperbolic knots having left-orderable, non- L -space surgeries from a given knot with left-orderable surgeries; see Section 4. Then we will apply the construction to prove the following:

Theorem 1.9. *There exist infinitely many hyperbolic knots K each of which enjoys the following properties.*

- (1) $K(r)$ is a hyperbolic 3-manifold for all $r \in \mathbb{Q}$.
- (2) $\mathcal{S}_{LO}(K) = \mathbb{Q}$.
- (3) $\mathcal{S}_L(K) = \emptyset$.

2. Left-orderable surgeries on periodic knots

A knot K in S^3 is called a *periodic knot* with period p if there is an orientation preserving diffeomorphism $f : S^3 \rightarrow S^3$ such that $f(K) = K$, $f^p = id$ ($p > 1$), $\text{Fix}(f) \neq \emptyset$, and $\text{Fix}(f) \cap K = \emptyset$, where $\text{Fix}(f)$ is the set




Figure 1: A periodic knot K with an axis C and its factor knot \overline{K} ; T is a tangle.

of fixed points of f . By the positive answer to the Smith conjecture [40], f is a rotation of S^3 about the unknotted circle $C = \text{Fix}(f)$. So by taking the quotient $S^3/\langle f \rangle$, we obtain the *factor knot* $\overline{K} = K/\langle f \rangle$ and the unknotted circle $\overline{C} = C/\langle f \rangle$ in $S^3 = S^3/\langle f \rangle$. We often call C the *axis* and \overline{C} the *branch circle*. See Figure 1. Since K is connected, the linking number $lk(\overline{K}, \overline{C})$ and the period p are relatively prime. Note that if the periodic knot K is unknotted, then the equivariant loop theorem [36] implies that $K \cup C$ is the Hopf link and $\overline{K} \cup \overline{C}$ is also the Hopf link. To exclude such a trivial case, in the following we consider non-trivial periodic knots.

The next theorem asserts that the fundamental groups of 3-manifolds obtained by Dehn surgeries on the periodic knot K inherit left-orderability from those of 3-manifolds obtained by Dehn surgeries on the factor knot \overline{K} . For a subset $\mathcal{S} \subset \mathbb{Q}$ and a positive integer p , we denote by $p\mathcal{S}$ the subset $\{pr \mid r \in \mathcal{S}\} \subset \mathbb{Q}$. Note that if $\mathcal{S} = \mathbb{Q}$, then $p\mathcal{S} = \mathbb{Q}$.

Theorem 2.1. *Let K be a non-trivial knot in S^3 with cyclic period p , and let \overline{K} be its factor knot. Then $\mathcal{S}_{\text{LO}}(K) \supset p\mathcal{S}_{\text{LO}}(\overline{K})$.*

Proof of Theorem 2.1. Let $f : S^3 \rightarrow S^3$ be an orientation preserving diffeomorphism giving the cyclic period p of K with axis $C = \text{Fix}(f)$ and factor knot $\overline{K} = K/\langle f \rangle$. Take an $\langle f \rangle$ -invariant tubular neighborhood $N(K)$ of K . Let $N(\overline{K})$ be the quotient $N(K)/\langle f \rangle$. In the following $E(K) = S^3 - \text{int}N(K)$ and $E(\overline{K}) = E(K)/\langle f \rangle = S^3 - \text{int}N(\overline{K})$. Denote by (μ, λ) (resp. $(\overline{\mu}, \overline{\lambda})$) a preferred meridian-longitude pair of $\pi_1(\partial N(K))$ (resp. $\pi_1(\partial N(\overline{K}))$). We can choose a simple closed curve representing the preferred longitude λ which is invariant under $\langle f \rangle$; see [12].

Let $\pi : E(K) \rightarrow E(\overline{K})$ be the cyclic branched covering branched along $\overline{C} = C/\langle f \rangle$. □

Lemma 2.2. *The branched cover $\pi : E(K) \rightarrow E(\overline{K})$ can be extended to a branched cover $\pi' : K(\frac{m}{n}) \rightarrow \overline{K}(\frac{m}{pn})$.*

Proof of Lemma 2.2. Let $\pi_* : \pi_1(E(K)) \rightarrow \pi_1(E(\overline{K}))$ be the homomorphism induced by π . Then $\pi_*|_{\pi_1(\partial E(K))} : \pi_1(\partial E(K)) \rightarrow \pi_1(\partial E(\overline{K}))$ sends μ to $\overline{\mu}$, and λ to $p\overline{\lambda}$. Hence, $\pi_*|_{\pi_1(\partial E(K))}(m\mu + n\lambda) = m\overline{\mu} + pn\overline{\lambda} = (m, p)\{\frac{m}{(m, p)}\overline{\mu} + \frac{pn}{(m, p)}\overline{\lambda}\}$. Then we can extend $\pi : E(K) \rightarrow E(\overline{K})$ to $\pi' : K(\frac{m}{n}) = E(K) \cup V \rightarrow \overline{K}(\frac{m}{pn}) = E(\overline{K}) \cup \overline{V}$, where V, \overline{V} are filled solid tori. If $(m, p) \geq 2$, then π' branches along the core of the filled solid torus \overline{V} as well as \overline{C} . □

Thus we have a commutative diagram:

$$\begin{array}{ccc} E(K) & \xrightarrow{\pi} & E(\overline{K}) \\ \text{Dehn filling} \downarrow & & \downarrow \text{Dehn filling} \\ K(\frac{m}{n}) & \xrightarrow{\pi'} & \overline{K}(\frac{m}{pn}) \end{array}$$

Assume that $\frac{m}{pn} \in \mathcal{S}_{\text{LO}}(\overline{K})$, i.e., $\overline{K}(\frac{m}{pn})$ has left-orderable fundamental group. Let us prove that $K(\frac{m}{n})$ has also left-orderable fundamental group, i.e., $p \times \frac{m}{pn} = \frac{m}{n} \in \mathcal{S}_{\text{LO}}(K)$.

Lemma 2.3. *$K(\frac{m}{n})$ is irreducible.*

Proof of Lemma 2.3. Suppose for a contradiction that $K(\frac{m}{n})$ is reducible. Since K is a non-trivial periodic knot, K is cabled and $\frac{m}{n}$ is the cabling slope [21, 22, 34]. First, we assume that K is a torus knot. Then $E(K)$ has a unique Seifert fibration (up to isotopy). Following [35, Theorem 2.2], we choose a Seifert fibration of $E(K)$ which is preserved by f . If C is not a fiber, we take a regular fiber t intersecting C . Since f fixes a point in $t \cap C$, $f(t) = t$ and f reverses the orientation of t . This then implies that f reverses the orientation of K , and hence C intersects K in exactly two points, a contradiction. Thus C is a fiber in the $\langle f \rangle$ -invariant Seifert fibration of $E(K)$. Since a regular fiber is knotted in S^3 , C is one of two exceptional fibers in $E(K)$. Then the quotient $E(\overline{K}) = E(K)/\langle f \rangle$ has also a Seifert fibration induced from that of $E(K)$ and thus \overline{K} is a torus knot; the surgery slope $\frac{m}{pn}$ on $\partial E(\overline{K})$ is the fiber slope. Since $\frac{m}{pn}$ is the fiber (i.e., cabling)

slope, $\pi_1(\overline{K}(\frac{m}{pn}))$ has a non-trivial torsion, contradicting the left-orderability of $\pi_1(\overline{K}(\frac{m}{pn}))$.

Next assume that K is an (x, y) -cable in a knotted solid torus W , where $y \geq 2$. By the $\langle f \rangle$ -invariant version ([35, Theorem 8.6]) of the torus decomposition theorem [28, 29], we may assume that f leaves a companion solid torus W invariant. First, we note that $W \cap C = \emptyset$. For otherwise, $f|_{\partial W}$ has fixed points and hence it is an involution, and f reverses the orientation of an $\langle f \rangle$ -invariant core of W . Hence, it also reverses the orientation of K (which has winding number $y \geq 2$ in W). This then implies that C intersects K in exactly two points, a contradiction. Thus $W \subset S^3 - C$. We denote the quotient $W/\langle f \rangle$ by \overline{W} . We may assume that the cable space $W - \text{int}N(K)$ has a Seifert fibration preserved by f [35, Theorem 2.2]. Then $\overline{W} - \text{int}N(\overline{K}) = (W - \text{int}N(K))/\langle f \rangle$ has an induced Seifert fibration in which a regular fiber on $\partial N(\overline{K})$ represents the surgery slope $\frac{m}{pn}$. This implies that the result of $\frac{m}{pn}$ -surgery of \overline{W} along \overline{K} , and hence $\overline{K}(\frac{m}{pn})$, has a non-trivial lens space summand whose fundamental group has order $y \geq 2$. Since $\pi_1(\overline{K}(\frac{m}{pn}))$ has a non-trivial torsion, it cannot be left-orderable, contradicting the assumption. \square

The above diagram induces the commutative diagram of fundamental groups below.

$$\begin{array}{ccc} \pi_1(E(K)) & \xrightarrow{\pi_*} & \pi_1(E(\overline{K})) \\ \downarrow & & \downarrow \\ \pi_1(K(\frac{m}{n})) & \xrightarrow{\pi'_*} & \pi_1(\overline{K}(\frac{m}{pn})). \end{array}$$

Lemma 2.4. $\pi'_* : \pi_1(K(\frac{m}{n})) \rightarrow \pi_1(\overline{K}(\frac{m}{pn}))$ is surjective.

Proof of Lemma 2.4. Choose a point $x \in C = \text{Fix}(f)$ (resp. $\pi(x) \in \overline{C}$) as a base point of $\pi_1(E(K))$ (resp. $\pi_1(E(\overline{K}))$). We take obvious meridians $\overline{\mu}_i$ of \overline{K} which are generators of $\pi_1(E(\overline{K}), \pi(x))$ (with respect to the Wirtinger presentation of $\pi_1(E(\overline{K}), \pi(x))$). Then their lifts $\mu_i \in \pi_1(E(K))$ satisfy $\pi_*(\mu_i) = \overline{\mu}_i$, and hence $\pi_* : \pi_1(E(K)) \rightarrow \pi_1(E(\overline{K}))$ is an epimorphism. Since vertical homomorphisms are also epimorphisms, $\pi'_* : \pi_1(K(\frac{m}{n})) \rightarrow \pi_1(\overline{K}(\frac{m}{pn}))$ is also an epimorphism. \square

By Lemma 2.3 $K(\frac{m}{n})$ is irreducible, and by Lemma 2.4 we have an epimorphism from $\pi_1(K(\frac{m}{n}))$ to the left-orderable group $\pi_1(\overline{K}(\frac{m}{pn}))$. Then

it follows from [5, Theorem 1.1(1)] that $\pi_1(K(\frac{m}{n}))$ is also left-orderable. Thus if $r = \frac{m}{pn} \in \mathcal{S}_{LO}(\overline{K})$, then $pr = \frac{m}{n} \in \mathcal{S}_{LO}(K)$. \square (Theorem 2.1)

3. *L*-space surgeries on periodic knots

In [43, 44] Ni proves that if a knot K in S^3 has an *L*-space surgery, then K is a fibered knot, i.e., $E(K)$ has a fibering over the circle. For a periodic knot K , the next theorem gives a necessary condition on the factor knot for K having an *L*-space surgery.

Theorem 3.1. *Let K be a periodic knot in S^3 with axis C , and let \overline{K} be its factor knot with branch circle \overline{C} . Suppose that K has an *L*-space surgery. Then $E(\overline{K})$ has a fibering over the circle with a fiber surface \overline{S} such that $|\overline{S} \cap \overline{C}|$ equals the algebraic intersection number between \overline{S} and \overline{C} , i.e., the linking number $lk(\overline{K}, \overline{C})$.*

In particular, we have:

Corollary 3.2. *Let K be a periodic knot with factor knot \overline{K} . If \overline{K} is not fibered, then $\mathcal{S}_L(K) = \emptyset$.*

Proof of Theorem 3.1. Let $f : S^3 \rightarrow S^3$ be an orientation preserving diffeomorphism of finite order satisfying $f(K) = K$. Note that $C = \text{Fix}(f)$, $\overline{K} = K/\langle f \rangle$ and $\overline{C} = C/\langle f \rangle$. Let $N(K)$ be an $\langle f \rangle$ -invariant tubular neighborhood of K .

Assume that K has an *L*-space surgery. Then Ni [43, Corollary 1.3] ([44]) proves that $E(K) = S^3 - \text{int}N(K)$ has a fibering over the circle. Following Proposition 6.1 in [13], we can isotope the fibering to a fibering preserved by the action of $\langle f \rangle$ so that the axis C is transverse to the fibers. Thus $E(\overline{K})$ inherits a fibering over the circle such that all the fibers are transverse to the branch circle $\overline{C} = C/\langle f \rangle$. Let \overline{S} be a fiber surface of $E(\overline{K})$. Since \overline{C} intersects each fiber surface of the fibering of $E(\overline{K})$ transversely, $|\overline{S} \cap \overline{C}|$ coincides with the algebraic intersection number between \overline{S} and \overline{C} , i.e., the linking number $lk(\partial\overline{S}, \overline{C})$, which equals the linking number $lk(\overline{K}, \overline{C})$. \square

As Ni [43, 44] proves, the fiberedness of K is necessary for K to have an *L*-space surgery. On the other hand, the periodicity of K itself also puts strong restrictions on 3-manifolds obtained by Dehn surgeries on K . For instance, if a periodic knot K with period $p > 2$ has a finite surgery, which is also an *L*-space surgery, then K is a torus knot or a cable of a torus knot [38, Proposition 5.6]. So we would like to ask:

Question 3.3. *Let K be a knot in S^3 with cyclic period $p > 2$ other than a torus knot or a cable of a torus knot. Then does K admit an L -space surgery?*

4. Periodic constructions

Given a periodic knot, taking the quotient by the periodic automorphism, we obtain its factor knot; see Section 2. Reversing this procedure, we have:

Definition 4.1 Periodic construction. Let $(\overline{K}, \overline{C})$ be a pair of a knot \overline{K} and an unknotted circle \overline{C} which is disjoint from \overline{K} . Then for an integer $p \geq 2$ with $(p, lk(\overline{K}, \overline{C})) = 1$, take the p -fold cyclic branched cover of S^3 branched along \overline{C} to obtain a knot $K_{\overline{C}}^p$ which is the preimage of \overline{K} . We call $K_{\overline{C}}^p$ the knot obtained from the pair $(\overline{K}, \overline{C})$ by p -periodic construction.

Note that $K_{\overline{C}}^p$ is a knot with cyclic period p whose factor knot is \overline{K} . Hence, Theorems 2.1 and 3.1 immediately imply the following result.

Theorem 4.2. *Let $(\overline{K}, \overline{C})$ be a pair as in Definition 4.1. If \overline{K} is a fibered knot, \overline{C} is chosen so that any fiber surface (i.e., minimal genus Seifert surface) \overline{S} satisfies the inequality $|\overline{S} \cap \overline{C}| > lk(\overline{K}, \overline{C})$. Then a knot $K_{\overline{C}}^p$ obtained from the pair $(\overline{K}, \overline{C})$ by p -periodic construction enjoys the following properties:*

- (1) $\mathcal{S}_{\text{LO}}(K_{\overline{C}}^p) \supset p\mathcal{S}_{\text{LO}}(\overline{K})$.
- (2) $\mathcal{S}_L(K_{\overline{C}}^p) = \emptyset$.

If \overline{K} is a trivial knot, then $\mathcal{S}_{\text{LO}}(\overline{K}) = \{0\}$ and hence $p\mathcal{S}_{\text{LO}}(\overline{K}) = \{0\}$. So we will apply Theorem 4.2 to non-trivial knots.

Remark 4.3. We denote the genus of a knot k in S^3 by $g(k)$. For \overline{K} and $K_{\overline{C}}^p$, we have $g(K_{\overline{C}}^p) \geq pg(\overline{K})$ [42, Theorem 3.2]. In particular, for a non-trivial knot \overline{K} , $g(K_{\overline{C}}^p) \rightarrow \infty$ as $p \rightarrow \infty$.

Theorem 4.2 is accompanied by the following theorems.

Theorem 4.4. *Given a non-trivial knot \overline{K} in S^3 , we can take an unknotted circle \overline{C} so that $\overline{K} \cup \overline{C}$ is a hyperbolic link with arbitrary linking number.*

Proof of Theorem 4.4. The following argument is based on the proofs of Theorems 2.1 and 2.2 in [1]. Arrange \overline{K} as a closed n -braid for some integer n .




Figure 2: Insertion of $(n - 1)$ -strands; $n = 2$.




Figure 3: Arrangement of C_1, \dots, C_{n-1} ; $n = 4$.

If necessary, stabilizing the braid, we may assume that the braid contains both a positive crossing and a negative crossing (Figure 2). Then introduce $(n - 1)$ -strands C_i ($i = 1, \dots, n - 1$) between the n -strands of the original braid so that the crossings introduced, together with the original crossings, are alternately positive and negative. See Figure 2.

Then we arrange C_i as in Figure 3 so that the closed braid is a 2-component link consisting of \overline{K} and an unknotted circle $\overline{C} = C_1 \cup \dots \cup C_{n-1}$ and $\overline{K} \cup \overline{C}$ is a non-split prime alternating link [37, Theorem 1].

Since our braid contains both a positive crossing and a negative crossing, we can add some negative twists or positive twists as in Figure 4 to make \overline{C} so that $lk(\overline{K}, \overline{C}) = l$ for a given integer l .

Note that the resulting link $\overline{K} \cup \overline{C}$ is also a non-split prime alternating link. It follows from [37, Corollary 2] that $\overline{K} \cup \overline{C}$ is either a torus link or a hyperbolic link. Since \overline{K} is non-trivial, but \overline{C} is trivial, the former cannot occur, and thus $\overline{K} \cup \overline{C}$ is a hyperbolic link. \square




Figure 4: Adding negative or positive twistings.

Theorem 4.5. (1) If $\overline{K} \cup \overline{C}$ is a hyperbolic link and $p > 2$, then $K_{\overline{C}}^p$ is a hyperbolic knot, and $K_{\overline{C}}^p(r)$ is a hyperbolic 3-manifold for all $r \in \mathbb{Q}$.

(2) Assume that $p > 2$ and \overline{C}_i ($i = 1, 2$) is an unknotted circle such that $lk(\overline{K}, \overline{C}_i)$ and p are relatively prime, and $\overline{K} \cup \overline{C}_i$ is a hyperbolic link. If $K_{\overline{C}_1}^p$ and $K_{\overline{C}_2}^p$ are isotopic in S^3 , then $\overline{K} \cup \overline{C}_1$ and $\overline{K} \cup \overline{C}_2$ are isotopic.

Proof of Theorem 4.5. (1) Assume for a contradiction that $K_{\overline{C}}^p$ is not hyperbolic. Then it is either a torus knot or a satellite knot. Let $f : S^3 \rightarrow S^3$ be the deck transformation of the p -fold cyclic branched cover given in Theorem 4.2, which is an orientation preserving diffeomorphism giving the cyclic period p of $K_{\overline{C}}^p$. In the following, we take an $\langle f \rangle$ -invariant tubular neighborhood $N(K_{\overline{C}}^p)$ and denote $S^3 - \text{int}N(K_{\overline{C}}^p)$ by $E(K_{\overline{C}}^p)$. The preimage of the branch circle \overline{C} is an unknotted circle $C = \text{Fix}(f)$, which is contained in the interior of $E(K_{\overline{C}}^p)$. Note also that $K_{\overline{C}}^p$ is a non-trivial knot. For otherwise, the equivariant loop theorem [36] implies that $K_{\overline{C}}^p \cup C$ is the Hopf link and $\overline{K} \cup \overline{C}$ is also the Hopf link, contradicting the hyperbolicity of $\overline{K} \cup \overline{C}$.

Claim 4.6. $K_{\overline{C}}^p$ is not a torus knot.

Proof of Claim 4.6. Assume for a contradiction that $K_{\overline{C}}^p$ is a torus knot. Then $E(K_{\overline{C}}^p)$ has a unique Seifert fibration up to isotopy. We choose a Seifert fibration of $E(K_{\overline{C}}^p)$ which is preserved by f [35, Theorem 2.2]. Then the argument in the proof of Lemma 2.3 shows that C is one of two exceptional fibers in $E(K_{\overline{C}}^p)$. Then the quotient $E(\overline{K}) - \text{int}N(\overline{C}) = (E(K_{\overline{C}}^p) - \text{int}N(C))/\langle f \rangle$

has also a Seifert fibration. Thus $S^3 - \text{int}N(\overline{K} \cup \overline{C}) = E(\overline{K}) - \text{int}N(\overline{C})$ is a Seifert fiber space, contradicting its hyperbolicity. \square

Claim 4.7. $K_{\overline{C}}^p$ is not a satellite knot.

Proof of Claim 4.7. Suppose for a contradiction that $K_{\overline{C}}^p$ is a satellite knot. Then we have an $\langle f \rangle$ -invariant torus decomposition of $E(K_{\overline{C}}^p)$ [35, Theorem 8.6]. Let Σ be the invariant family of essential tori in $E(K_{\overline{C}}^p)$.

Case (i). There is an essential torus $T \in \Sigma$ such that $f(T) = T$. Then T bounds an $\langle f \rangle$ -invariant companion solid torus W containing $K_{\overline{C}}^p$. Note that $K_{\overline{C}}^p$ is not a core of W . We see that $W \cap C = \emptyset$, for otherwise $f|_{\partial W}$ has a fixed point and it is an involution, i.e., $(f|_{\partial W})^2$ is the identity map. By the classical Smith theory [52] f itself is an involution, contradicting the assumption. Thus W lies in $S^3 - C$. We may assume that $W \subset S^3 - \text{int}N(C)$ for a small tubular neighborhood $N(C)$ of C . Since the core of W is not a core of $S^3 - \text{int}N(C)$, $S^3 - \text{int}N(K_{\overline{C}}^p \cup C)$ contains the $\langle f \rangle$ -invariant essential torus $T = \partial W$. This then implies that $S^3 - \text{int}N(\overline{K} \cup \overline{C})$ contains an essential torus $\partial W / \langle f \rangle$. This contradicts the hyperbolicity of $S^3 - \text{int}N(\overline{K} \cup \overline{C})$.

Case (ii). For each $T \in \Sigma$, $f(T) \neq T$ (hence, $f(T) \cap T = \emptyset$). Let us pick an essential torus $T \in \Sigma$. Note that T is essential in $S^3 - \text{int}N(K_{\overline{C}}^p \cup C)$. Then the image $\overline{T} \subset E(\overline{K} \cup \overline{C})$ of T by the covering projection is also essential. This contradicts the hyperbolicity of $S^3 - \text{int}N(\overline{K} \cup \overline{C})$. \square

It follows that $K_{\overline{C}}^p$ is a hyperbolic knot in S^3 .

Since $K_{\overline{C}}^p$ is a hyperbolic knot with period $p > 2$, it follows from [39, Corollary 1.4] that $K_{\overline{C}}^p(r)$ is a hyperbolic 3-manifold for all $r \in \mathbb{Q}$, or $p = 3$, $r = 0$ and $g(K_{\overline{C}}^p) = 1$. Since $g(K_{\overline{C}}^p) \geq pg(\overline{K}) \geq p > 2$, the latter cannot occur. Hence $K_{\overline{C}}^p(r)$ is a hyperbolic 3-manifold for all $r \in \mathbb{Q}$ as desired.

(2) In the following, for notational simplicity, we write $K_i = K_{C_i}^p$.

The assumption, together with (1), implies that K_i ($i = 1, 2$) is a hyperbolic knot. Recall that K_i has an orientation preserving diffeomorphism f_i such that $f_i(K_i) = K_i$, $f_i^p = id$ and $\text{Fix}(f_i) = C_i$. Note that $\overline{K} = K_i / \langle f_i \rangle$ and $\overline{C}_i = C_i / \langle f_i \rangle$. Suppose that K_1 and K_2 are isotopic in S^3 . Then we have an orientation preserving diffeomorphism φ of S^3 such that $\varphi(K_1) = K_2$. Note that $f'_2 = \varphi^{-1} \circ f_2 \circ \varphi$ is an orientation preserving diffeomorphism of S^3 , which satisfies $f'_2(K_1) = K_1$ and gives also a cyclic period p for K_1 . Let us put $C'_2 = \varphi^{-1}(C_2)$. Then we see that $\text{Fix}(f'_2) = C'_2$. Since $\varphi \circ f'_2 = f_2 \circ \varphi$, φ induces an orientation preserving diffeomorphism

$\bar{\varphi} : S^3 = S^3/\langle f'_2 \rangle \rightarrow S^3 = S^3/\langle f_2 \rangle$ sending $K_1/\langle f'_2 \rangle$ to $\bar{K} = K_2/\langle f_2 \rangle$ and $C'_2/\langle f'_2 \rangle$ to $\bar{C}_2 = C_2/\langle f_2 \rangle$.

Now the hyperbolic knot K_1 has two orientation preserving, periodic diffeomorphisms f_1 and f'_2 of period $p > 2$. Then [3, 2.1 Theorem (a)] shows that the pairwise isotopy classes $[f_1]$ and $[f'_2]$ in the symmetry group $\text{Sym}(S^3, K_1)$ have order p . Furthermore, since K_1 is hyperbolic, $\text{Sym}(S^3, K_1)$ is isomorphic to a finite cyclic group or a dihedral group [31, Theorems 10.5.3 and 10.6.2(2)]. This implies that subgroups $\langle [f_1] \rangle$ and $\langle [f'_2] \rangle$ of order p coincide, since $p > 2$. Then it follows from [3, 2.1 Theorem (c)] that $\langle f_1 \rangle$ and $\langle f'_2 \rangle$ are conjugate by a diffeomorphism g in $\text{Diff}(S^3, K_1)$, which is isotopic to the identity. Hence, $(f'_2)^k = g \circ f_1 \circ g^{-1}$ for some integer k ($1 \leq k \leq p - 1$), which has also period p . Note that $(f'_2)^k$ leaves K_1 invariant and $K_1/\langle (f'_2)^k \rangle = K_1/\langle f'_2 \rangle$, and that $\text{Fix}((f'_2)^k) = C'_2$ and $C'_2/\langle (f'_2)^k \rangle = C'_2/\langle f'_2 \rangle$. For any $x \in C_1 = \text{Fix}(f_1)$, we have $(f'_2)^k(g(x)) = g(f_1(x)) = g(x)$, thus $g(x) \in \text{Fix}((f'_2)^k) = C'_2$, and hence $g(C_1) \subset C'_2$. Conversely if $x' \in C'_2 = \text{Fix}((f'_2)^k)$, then we see that $g^{-1}(x') \in C_1$ and $x' \in g(C_1)$, hence $C'_2 \subset g(C_1)$. Thus, we have $g(C_1) = C'_2$. Therefore we have an orientation preserving diffeomorphism $\bar{g} : S^3 = S^3/\langle f_1 \rangle \rightarrow S^3 = S^3/\langle (f'_2)^k \rangle = S^3/\langle f'_2 \rangle$ sending $\bar{K} = K_1/\langle f_1 \rangle$ to $K_1/\langle (f'_2)^k \rangle = K_1/\langle f'_2 \rangle$ and $\bar{C}_1 = C_1/\langle f_1 \rangle$ to $C'_2/\langle (f'_2)^k \rangle = C'_2/\langle f'_2 \rangle$.

Now the orientation preserving diffeomorphism $\bar{\varphi} \circ \bar{g}$ of S^3 satisfies $\bar{\varphi} \circ \bar{g}(\bar{K}) = \bar{K}$ and $\bar{\varphi} \circ \bar{g}(\bar{C}_1) = \bar{C}_2$. Thus, $\bar{K} \cup \bar{C}_1$ and $\bar{K} \cup \bar{C}_2$ are isotopic. \square

5. Examples

In this section, we present two examples illustrating how the periodic construction works according to whether the initial knot \bar{K} is fibered or not fibered.

First we apply Theorem 4.2 in the case where \bar{K} is not fibered. In such a case we can choose \bar{C} arbitrarily with $lk(\bar{K}, \bar{C}) \neq 0$ to obtain a knot $K_{\bar{C}}^p$ having properties (1) and (2) in Theorem 4.2.

Let T_n ($n \neq 0, \pm 1$) be the twist knot illustrated in Figure 5.

Then T_n is a hyperbolic knot, and since the Alexander polynomial of T_n is not monic, it is not fibered [6, 8.16 Proposition]. Suppose that $n > 1$. Then it follows from [20, 56] that $\pi_1(T_n(r))$ is left-orderable for $r \in (-4n, 4)$. Furthermore, it is known by [54] that $\pi_1(T_n(4))$ is left-orderable. Hence, $\mathcal{S}_{\text{LO}}(T_n) \supset (-4n, 4] \cap \mathbb{Q}$.

Example 5.1. Let us take a 2-component link $T_2 \cup \bar{C}$ as in Figure 6; $lk(T_2, \bar{C}) = 1$. Let p be any integer with $p > 2$ and $K_{2, \bar{C}}^p$ a knot obtained




Figure 5: The twist knot T_n .




Figure 6: The twist knot T_2 and an axis \overline{C} .

from (T_2, \overline{C}) by p -periodic construction. Then $K_{2, \overline{C}}^p$ enjoys the following properties:

- (1) $K_{2, \overline{C}}^p$ is a hyperbolic knot in S^3 .
- (2) $K_{2, \overline{C}}^p(r)$ is a hyperbolic 3-manifold for all $r \in \mathbb{Q}$.
- (3) $\mathcal{S}_{\text{LO}}(K_{2, \overline{C}}^p) \supset (-8p, 4p] \cap \mathbb{Q}$.
- (4) $\mathcal{S}_L(K_{2, \overline{C}}^p) = \emptyset$.

Proof. Assertions (1) and (2) follow from Theorem 4.5(1) once we show that $T_2 \cup \overline{C}$ is a hyperbolic link. Since $T_2 \cup \overline{C}$ is a non-split prime alternating link [37, Theorem 1], it is either a torus link or a hyperbolic link [37, Corollary 2]. The former cannot happen, because T_2 is non-trivial, but \overline{C} is trivial. Hence $T_2 \cup \overline{C}$ is a hyperbolic link as desired. Since T_2 is not fibered and




Figure 7: The trefoil knot $T_{-3,2}$ and the unknotted circle \bar{C} .

$\pi_1(T_2(r))$ is left-orderable for $r \in (-8, 4]$, assertions (3) and (4) follow from Theorem 4.2. \square

Next we apply Theorem 4.2 in the case where \bar{K} is a fibered knot. In the next example, we take a trefoil knot $T_{-3,2}$ as \bar{K} .

Example 5.2. Let us take the 2-component link $T_{-3,2} \cup \bar{C}$ shown in Figure 7; $lk(T_{-3,2}, \bar{C}) = 1$. Let p be any integer with $p > 2$ and $K_{-3,2,\bar{C}}^p$ a knot obtained from $(T_{-3,2}, \bar{C})$ by p -periodic construction. Then $K_{-3,2,\bar{C}}^p$ enjoys the following properties:

- (1) $K_{-3,2,\bar{C}}^p$ is a hyperbolic knot in S^3 .
- (2) $K_{-3,2,\bar{C}}^p(r)$ is a hyperbolic 3-manifold for all $r \in \mathbb{Q}$.
- (3) $\mathcal{S}_{\text{LO}}(K_{-3,2,\bar{C}}^p) \supset (-p, \infty) \cap \mathbb{Q}$.
- (4) $\mathcal{S}_L(K_{-3,2,\bar{C}}^p) = \emptyset$.

Proof of Example 5.2. Recall that $\mathcal{S}_{\text{LO}}(T_{-3,2}) = (-1, \infty) \cap \mathbb{Q}$; see Remark 1.2(2) and Example 1.6.

Since as illustrated in Figure 7(i) $T_{-3,2} \cup \bar{C}$ is a non-split prime alternating link [37, Theorem 1], it is either a torus link or a hyperbolic link [37, Corollary 2]. If we have the former case, then $T_{-3,2}$ is isotopic to \bar{C} which is

a trivial knot, a contradiction. Hence, $T_{-3,2} \cup \overline{C}$ is a hyperbolic link. Then (1) and (2) follow from Theorem 4.5(1).

Let us prove (3) and (4) using Theorem 4.2. Since $T_{-3,2}$ is fibered, we need to check the condition of Theorem 4.2: for any fiber surface \overline{S} of $E(T_{-3,2})$, $|\overline{S} \cap \overline{C}|$ is strictly bigger than the algebraic intersection number between \overline{S} and \overline{C} , i.e., $lk(T_{-3,2}, \overline{C})$.

In Figure 7(ii), we give a minimal genus Seifert surface F of $T_{-3,2}$, which is a once-punctured torus with $\partial F = T_{-3,2}$. Put $\overline{S} = F \cap E(T_{-3,2})$. Then by [13, Lemma 5.1] \overline{S} is a fiber surface of $E(T_{-3,2})$. We see that $|\overline{S} \cap \overline{C}| = 5$ and the algebraic intersection number between \overline{S} and \overline{C} is one. Assume for a contradiction that we have another fiber surface \overline{S}' of $E(T_{-3,2})$ such that $|\overline{S}' \cap \overline{C}| < |\overline{S} \cap \overline{C}|$. Since \overline{S} and \overline{S}' are fiber surfaces of $E(T_{-3,2})$, they are isotopic; see [13, Lemma 5.1], [55]. This then implies that we can isotope \overline{C} to \overline{C}' in $E(T_{-3,2})$ so that $|\overline{S} \cap \overline{C}'| < |\overline{S} \cap \overline{C}|$.

Claim 5.3. *There exists a smooth map φ from a semi-disk D into $E(T_{-3,2})$ such that $\varphi^{-1}(\overline{C})$ is an arc $c \subset \partial D$ and $\varphi^{-1}(\overline{S})$ is the arc $\alpha = \overline{\partial D} - c$.*

Proof of Claim 5.3. Let $\Phi : S^1 \times [0, 1] \rightarrow E(T_{-3,2})$ be a smooth map giving an isotopy between $\overline{C}(= \Phi(S^1 \times \{0\}))$ to $\overline{C}'(= \Phi(S^1 \times \{1\}))$. We may assume Φ is transverse to \overline{S} . Furthermore, the essentiality of \overline{S} in $E(T_{-3,2})$ enables us to modify Φ to eliminate the circle components as usual. Since $|\overline{S} \cap \overline{C}'| < |\overline{S} \cap \overline{C}| = 5$ and the algebraic intersection number between \overline{S} and \overline{C}' coincides with the algebraic intersection number between \overline{S} and \overline{C} , we have $|\overline{S} \cap \overline{C}'| = 1$ or 3 . Thus $\Phi^{-1}(\overline{S})$ consists of three properly embedded arcs α, α' and β , where $\partial\alpha \subset S^1 \times \{0\}$, $\partial\alpha' \subset S^1 \times \{0\}$, and β connects $S^1 \times \{0\}$ and $S^1 \times \{1\}$ (Figure 8(i), (ii)), consists of four properly embedded arcs α, β, β' and β'' , where $\partial\alpha \subset S^1 \times \{0\}$, and each of β, β', β'' connects $S^1 \times \{0\}$ and $S^1 \times \{1\}$ (Figure 8(iii)), or consists of four properly embedded arcs α, α', β and γ , where $\partial\alpha \subset S^1 \times \{0\}$, $\partial\alpha' \subset S^1 \times \{0\}$, β connects $S^1 \times \{0\}$ and $S^1 \times \{1\}$, and $\partial\gamma \subset S^1 \times \{1\}$ (Figure 8(iv), (v)). In either case there is a semi-disk D cobounded by α and an arc $c \subset S^1 \times \{0\}$.

Putting $\varphi = \Phi|_D : D \rightarrow E(T_{-3,2})$, we obtain a desired smooth map. \square

Cut open $E(T_{-3,2})$ along \overline{S} to obtain a product 3-manifold $\overline{S} \times [0, 1]$. The circle \overline{C} is cut into five arcs c_1, c_2, c_3, c_4 and c_5 as in Figure 7(ii). Note that $\partial c_1 \subset \overline{S} \times \{0\}$, $\partial c_3 \subset \overline{S} \times \{1\}$, and each of c_2, c_4, c_5 connects $\overline{S} \times \{0\}$ and $\overline{S} \times \{1\}$. Moreover, we see that c_1 and c_3 are linking once relative their boundaries.




Figure 8: $\Phi^{-1}(\bar{S})$ in $S^1 \times [0, 1]$.

On the other hand, since c is either c_1 or c_3 , Claim 5.3 shows that c_1 and c_3 are unlinked relative their boundaries. This contradiction shows that for any fiber surface \bar{S} , $|\bar{S} \cap \bar{C}| = 5$ and $|\bar{S} \cap \bar{C}| > lk(T_{-3,2}, \bar{C})$. Since $\pi_1(T_{-3,2}(r))$ is left-orderable if $r \in (-1, \infty)$, the conclusions (3) and (4) follow from Theorem 4.2. This completes the proof of Example 5.2. \square

6. Surgeries on alternating knots

Theorem 1.5 in [47], together with [48, Proposition 9.6] ([43, Proof of Corollary 1.3], [26, Claim 2]), shows that for an alternating knot K which is not a $(p, 2)$ -torus knot, $K(r)$ is not an L -space for all $r \in \mathbb{Q}$.

We say that an alternating knot is *positive* (resp. *negative*) if it has a reduced alternating diagram such that each of the crossings is positive (resp. negative). An alternating knot is *special* if it is either positive or negative.

In [4] Boyer, Gordon and Watson prove:

Proposition 6.1 [4]. *Let K be a prime alternating knot in S^3 .*

- (1) *If K is not a special alternating knot, then $\pi_1(K(\frac{1}{n}))$ is left-orderable for all non-zero integers n .*
- (2) *If K is a positive (resp. negative) alternating knot, then $\pi_1(K(\frac{1}{n}))$ is left-orderable for all positive (resp. negative) integers n .*




Figure 9: An alternating knot \overline{K} and a perpendicular circle \overline{C} .

Let \overline{K} be an alternating knot. For convenience, we position $\overline{K} \subset \mathbb{R}^3 = S^3 - \{\infty\}$ so that \overline{K} lies in the xy -plane except near crossings of \overline{K} , where \overline{K} lies on a “bubble” as in [37]. Then we say an unknotted circle $\overline{C} \subset S^3 - \overline{K}$ is *perpendicular* if it passes ∞ and intersects the xy -plane exactly once. Note that $\overline{C} \cap \mathbb{R}^3$ is perpendicular to the xy -plane. See Figure 9, in which the dot indicates a perpendicular circle \overline{C} .

Proposition 6.2. *Let \overline{K} be a prime alternating knot and \overline{C} a perpendicular circle. Let p be an integer such that $p \geq 2$ and $(p, \text{lk}(\overline{K}, \overline{C})) = 1$, and let $K_{\overline{C}}^p$ be the knot obtained from $(\overline{K}, \overline{C})$ by p -periodic construction. Then we have:*

- (1) $K_{\overline{C}}^p$ is an alternating knot.
- (2) If \overline{K} is not a special alternating knot, then $\pi_1(K_{\overline{C}}^p(\frac{p}{n}))$ is left-orderable for all non-zero integers n .
- (3) If \overline{K} is a positive (resp. negative) alternating knot, then $\pi_1(K_{\overline{C}}^p(\frac{p}{n}))$ is left-orderable for all positive (resp. negative) integers n .

Proof of Proposition 6.2. The first assertion follows immediately from diagrammatic consideration. The conclusions (2) and (3) follow from Proposition 6.1 and Theorem 2.1. \square

Remark 6.3. In Proposition 6.2, if \overline{K} is not a $(p, 2)$ -torus knot, then $K_{\overline{C}}^p$ is not a $(p', 2)$ -torus knot. For otherwise, the argument in the proof of Lemma 2.3 implies that \overline{K} is a torus knot. Since it is alternating, it is a $(p, 2)$ -torus knot for some odd integer p [41, Theorem 3.2], a contradiction. Therefore, as mentioned in the beginning of this section, $K_{\overline{C}}^p(r)$ is not an L -space for all $r \in \mathbb{Q}$.

Applying Proposition 6.2 and Remark 6.3, we have:

Example 6.4. Take an alternating knot \overline{K} and a perpendicular circle \overline{C} as illustrated in Figure 9; $lk(\overline{K}, \overline{C}) = 1$. Note that \overline{K} is not a special alternating knot. Hence for any integer $p \geq 2$, $K_{\overline{C}}^p$ is an alternating knot, $K_{\overline{C}}^p(r)$ is not an L -spaces for all $r \in \mathbb{Q}$, and $\pi_1(K_{\overline{C}}^p(\frac{p}{n}))$ is left-orderable for all non-zero integers n .

7. Knots with $\mathcal{S}_{LO}(K) = \mathbb{Q}$ and $\mathcal{S}_L(K) = \emptyset$

The goal of this section is to prove Theorems 1.8 and 1.9. We start with Proposition 7.1 below, which was shown by Clay and Watson [10, Proposition 4.1].

Let k be a knot in S^3 , which is contained in a standardly embedded solid torus $V \subset S^3$. Assume that k is not contained in a 3-ball in V . We call k a *pattern knot* in S^3 and the pair (V, k) a *pattern*. Let f be an orientation preserving embedding from V into S^3 which sends a preferred longitude of V to that of $f(V) \subset S^3$. Then we obtain a knot $K = f(k)$ in S^3 , which is called a *satellite knot* with a pattern knot k and a *companion knot* $K' = f(c)$, where c is a core of V .

Proposition 7.1 [10]. *Let K be a satellite knot with a pattern knot k . If $K(r)$ is irreducible and $r \in \mathcal{S}_{LO}(k)$, then $r \in \mathcal{S}_{LO}(K)$.*

7.1. Composite knots K with $\mathcal{S}_{LO}(K) = \mathbb{Q}$ and $\mathcal{S}_L(K) = \emptyset$

In this subsection, we prove that the connected sum of two torus knots $T_{-p,q}$ and $T_{r,s}$ where $p > q \geq 2$ and $r > s \geq 2$, satisfies $\mathcal{S}_{LO}(T_{-p,q} \# T_{r,s}) = \mathbb{Q}$ and $\mathcal{S}_L(T_{-p,q} \# T_{r,s}) = \emptyset$ (Proposition 7.3). Thus $T_{-p,q} \# T_{r,s}$ satisfies Conjecture 1.3.

Proposition 7.1 and Theorem 2.1 immediately imply:

Proposition 7.2. *Let K and K' be nontrivial knots. Then we have:*

- (1) $\mathcal{S}_{\text{LO}}(K \# K') \supset \mathcal{S}_{\text{LO}}(K) \cup \mathcal{S}_{\text{LO}}(K')$.
- (2) $\mathcal{S}_{\text{LO}}(pK) \supset p\mathcal{S}_{\text{LO}}(K)$, where pK denotes the connected sum of p copies of K .

Proof of Proposition 7.2. (1) Following [17, Lemma 7.1], we see that $(K \# K')$ (r) is irreducible for all $r \in \mathbb{Q}$.

Let us regard $K \# K'$ as a satellite knots with a pattern knot K and a companion knot K' . Then Proposition 7.1 shows that $\mathcal{S}_{\text{LO}}(K \# K') \supset \mathcal{S}_{\text{LO}}(K)$. Exchanging the roles of K and K' , we have $\mathcal{S}_{\text{LO}}(K \# K') \supset \mathcal{S}_{\text{LO}}(K')$ as well. Thus $\mathcal{S}_{\text{LO}}(K \# K') \supset \mathcal{S}_{\text{LO}}(K) \cup \mathcal{S}_{\text{LO}}(K')$.

(2) Since pK is a knot with cyclic period p whose factor knot is K , the result follows from Theorem 2.1. \square

As a step toward proofs of Theorems 1.8 and 1.9, we prove:

Proposition 7.3. *For torus knots $T_{-p,q}$ and $T_{r,s}$, where $p > q \geq 2$ and $r > s \geq 2$, we have $\mathcal{S}_{\text{LO}}(T_{-p,q} \# T_{r,s}) = \mathbb{Q}$ and $\mathcal{S}_L(T_{-p,q} \# T_{r,s}) = \emptyset$.*

Proof of Proposition 7.3. Recall that $\mathcal{S}_{\text{LO}}(T_{-p,q}) = (-pq + p + q, \infty) \cap \mathbb{Q}$ and $\mathcal{S}_{\text{LO}}(T_{r,s}) = (-\infty, rs - r - s) \cap \mathbb{Q}$. Note that $-pq + p + q < 0 < rs - r - s$. Now apply Proposition 7.2 to $T_{-p,q} \# T_{r,s}$ to conclude that $\mathcal{S}_{\text{LO}}(T_{-p,q} \# T_{r,s}) \supset ((-pq + p + q, \infty) \cup (-\infty, rs - r - s)) \cap \mathbb{Q} = \mathbb{Q}$. Hence, $\mathcal{S}_{\text{LO}}(T_{-p,q} \# T_{r,s}) = \mathbb{Q}$.

Next we show that $T_{-p,q} \# T_{r,s}$ has no L -space surgeries.

Claim 7.4. *The coefficient of t in the Alexander polynomial of $T_{-p,q} \# T_{r,s}$ is -2 .*

Proof of Claim 7.4. Recall that $T_{-p,q}$ has the Alexander polynomial $\Delta_{T_{-p,q}}(t) = \Delta_{T_{p,q}}(t) = \frac{(t^{pq} - 1)(t - 1)}{(t^p - 1)(t^q - 1)}$, and $T_{r,s}$ has the Alexander polynomial $\Delta_{T_{r,s}}(t) = \frac{(t^{rs} - 1)(t - 1)}{(t^r - 1)(t^s - 1)}$. Since $\Delta_{T_{-p,q} \# T_{r,s}}(t) = \Delta_{T_{-p,q}}(t)\Delta_{T_{r,s}}(t) = \Delta_{T_{p,q}}(t)\Delta_{T_{r,s}}(t)$, $\Delta'_{T_{-p,q} \# T_{r,s}}(0) = \Delta'_{T_{p,q}}(0)\Delta_{T_{r,s}}(0) + \Delta_{T_{p,q}}(0)\Delta'_{T_{r,s}}(0)$. Note that $\Delta_{T_{p,q}}(0) = \Delta_{T_{r,s}}(0) = 1$ and a simple computation shows that $\Delta'_{T_{p,q}}(0) = \Delta'_{T_{r,s}}(0) = -1$. Thus $\Delta'_{T_{-p,q} \# T_{r,s}}(0) = (-1) + (-1) = -2$. This then implies that the coefficient of t in the Alexander polynomial of $T_{-p,q} \# T_{r,s}$ is -2 . \square




Figure 10: $k \cup C$.

Apply [47, Corollary 1.3], together with [48, Proposition 9.6] ([43, Proof of Corollary 1.3], [26, Claim 2]), to conclude that $T_{-p,q} \# T_{r,s}$ has no L -space surgeries. \square

Let us consider the connected sum $T_{p,q} \# T_{r,s}$ instead of $T_{-p,q} \# T_{r,s}$, where $p > q \geq 2$ and $r > s \geq 2$. The argument in the proof of Claim 7.4 shows that $\mathcal{S}_L(T_{p,q} \# T_{r,s}) = \emptyset$. On the other hand, putting $m_0 = \max\{pq - p - q, rs - r - s\}$, Example 1.6 and Proposition 7.2 merely imply $\mathcal{S}_{LO}(T_{p,q} \# T_{r,s}) \supset (-\infty, m_0)$. So we would like to ask:

Question 7.5. *Does $\mathcal{S}_{LO}(T_{p,q} \# T_{r,s}) = \mathbb{Q}$ hold for integers $p > q \geq 2$ and $r > s \geq 2$?*

7.2. Proof of Theorem 1.8

Let us consider $k = T_{-3,2} \# T_{3,2}$ and take an unknotted circle C as in Figure 10. Following Proposition 7.3, $\mathcal{S}_{LO}(k) = \mathbb{Q}$.

Note that the link $k \cup C$ is an alternating link. Since $k \cup C$ is a non-split prime alternating link [37, Theorem 1], it is either a torus link or a hyperbolic link [37, Corollary 2]. The former is not the case, because k is non-trivial, but C is trivial. Thus, $k \cup C$ is hyperbolic, hence letting $V = S^3 - \text{int}N(C)$, k is a hyperbolic knot in V . Apply the satellite construction with the pattern (V, k) and the companion knot K' to obtain a satellite knot K with a pattern knot $k = T_{-3,2} \# T_{3,2}$. Since k is hyperbolic in V , the satellite knot K is prime, and the 3-manifold obtained from V by r -surgery on k is again hyperbolic for all but finitely many $r \in \mathbb{Q}$. This then implies that $K(r)$ is a toroidal 3-manifold with a hyperbolic piece, in particular, $K(r)$ is not a graph manifold, for all but finitely many $r \in \mathbb{Q}$. This establishes (1).

If $K(r)$ were reducible for some $r \in \mathbb{Q}$, then K is cabled [51, 4.5 Corollary]. However, this is impossible, because $V - k$ is hyperbolic. Hence $K(r)$

is irreducible for any $r \in \mathbb{Q}$. Now Proposition 7.1 shows that $\mathcal{S}_{\text{LO}}(K) \supset \mathcal{S}_{\text{LO}}(k) = \mathbb{Q}$.

Let us see that $\mathcal{S}_L(K) = \emptyset$. Since $lk(k, C) = 0$, i.e., the winding number of K in V is zero, (V, k) is not fibered, and hence neither is the satellite knot K ; see [25, Theorem 1]. Hence, [43, Corollary 1.3] shows that $\mathcal{S}_L(K) = \emptyset$.

Finally, we show that there are infinitely many satellite knots K with a companion knot K' and enjoy the required properties in Theorem 1.8. For instance, let us take $k_p = T_{-3,2} \# T_{p,2}$ ($p \geq 3$). As shown in Proposition 7.3, $\mathcal{S}_{\text{LO}}(k_p) = \mathbb{Q}$ for all $p \geq 3$. It follows from Theorem 4.4, there is an unknotted circle C_p so that $k_p \cup C_p$ is hyperbolic and $lk(k_p, C_p) = 0$. Each C_p gives a pattern (V, k_p) . Let K_p be a satellite knot with a companion knot K' and pattern (V, k_p) . Then the same argument as above shows that K_p satisfies the properties of Theorem 1.8. If $p \neq p' \geq 3$, then $k_p \cup C_p$ are not isotopic to $k_{p'} \cup C_{p'}$. Hence, there is no orientation preserving diffeomorphism of V which leaves the preferred longitude of V invariant and maps k_p to $k_{p'}$. Thus we see that the resulting satellite knots K_p and $K_{p'}$ are never isotopic.

7.3. Proof of Theorems 1.9

As in the proof of Theorem 1.8, we consider the connected sum $T_{-3,2} \# T_{3,2}$, which has the property: $\mathcal{S}_{\text{LO}}(T_{-3,2} \# T_{3,2}) = \mathbb{Q}$ (Proposition 7.3).

Although we can apply the periodic construction and Theorem 4.2 to the fibered knot $T_{-3,2} \# T_{3,2}$, for ease of handling, we take the connected sum $(T_{-3,2} \# T_{3,2}) \# T_2$, where T_2 is the twist knot shown in Figure 5. The Alexander polynomial of $(T_{-3,2} \# T_{3,2}) \# T_2$ is $(t^2 - t + 1)^2(2t^2 - 5t + 2)$, which is not monic, and hence $(T_{-3,2} \# T_{3,2}) \# T_2$ is not fibered. Proposition 7.2 shows that $\mathcal{S}_{\text{LO}}((T_{-3,2} \# T_{3,2}) \# T_2) \supset \mathcal{S}_{\text{LO}}(T_{-3,2} \# T_{3,2}) = \mathbb{Q}$, and hence $\mathcal{S}_{\text{LO}}((T_{-3,2} \# T_{3,2}) \# T_2) = \mathbb{Q}$.

Let us put $\bar{K} = T_{-3,2} \# T_{3,2} \# T_2$ and take an unknotted circle \bar{C} as in Figure 11; $lk(\bar{K}, \bar{C}) = 1$.

Since $\bar{K} \cup \bar{C}$ is a non-split prime alternating link [37, Theorem 1], it is either a torus link or a hyperbolic link [37, Corollary 2]. The former cannot happen, because \bar{K} is non-trivial, but \bar{C} is trivial. Hence, $\bar{K} \cup \bar{C}$ is a hyperbolic link. Let $p > 2$ be any integer, and apply the p -periodic construction to the pair (\bar{K}, \bar{C}) to obtain a knot $K_{\bar{C}}^p$. It follows from Theorem 4.2 and Theorem 4.5(1) that $K_{\bar{C}}^p$ is a hyperbolic knot and enjoys the properties (1)–(3) in Theorem 1.9. By changing p , we obtain infinitely many such knots. For instance, see Remark 4.3.




Figure 11: $\overline{K} \cup \overline{C}$.

- Remark 7.6.**
- 1) By Theorem 4.4 there are infinitely many unknotted circles for $\overline{K} = T_{-3,2} \# T_{3,2} \# T_2$, and for each unknotted circle \overline{C} we obtain infinitely many hyperbolic knots $K_{\overline{C}}^p$, where p and $lk(\overline{K}, \overline{C})$ are relatively prime. See also Theorem 4.5(2).
 - 2) Recall that any knot K obtained by the periodic construction (Section 4), for instance a knot obtained in the proof of Theorem 1.9, is not fibered and every non-trivial surgery on K is a left-orderable, non- L -space surgery. So we can apply Theorem 4.2 again to the knot K and an arbitrarily chosen unknotted circle to obtain yet further infinitely many non-fibered knots K' each of which has the (same) factor knot K . Then r -surgery on K' is also a left-orderable, non- L -space surgery for all $r \in \mathbb{Q}$. We can apply this procedure repeatedly arbitrarily many times.
 - 3) Let K be the knot 10_{99} in Rolfsen's knot table [50]. Recently, Clay [7] used an epimorphism from $E(K)$ to $E(T_{3,2})$ which preserves the peripheral subgroup [32] to show that every non-trivial surgery on K is left-orderable surgery. Since K has no cyclic period [31, Appendix F], this example cannot be explained by the periodic construction.

8. Shapes of $\mathcal{S}_{LO}(K)$ — questions and conjectures

As we mentioned in Remark 1.2(1), $0 \in \mathcal{S}_{LO}(K)$ for any knot K . If K is the trivial knot then $\mathcal{S}_{LO}(K) = \{0\}$, which has the smallest size. On the other hand, Theorems 1.8, 1.9 and Proposition 7.3 demonstrate that there are infinitely many knots K with $\mathcal{S}_{LO}(K) = \mathbb{Q}$, which has largest size.

It seems interesting to determine the shape of $\mathcal{S}_{LO}(K)$ when it is neither $\{0\}$ nor \mathbb{Q} .

Question 8.1. *If K is a non-trivial knot in S^3 , then does $\mathcal{S}_{\text{LO}}(K)$ contain $(-1, 1) \cap \mathbb{Q}$?*

Recently, Li and Roberts [33, Corollary 1.2] prove that for any hyperbolic knot K , there exists a constant N_K such that $\{\frac{1}{n} \mid |n| > N_K\} \subset \mathcal{S}_{\text{LO}}(K)$.

More strongly, we would like to ask:

Question 8.2. *If K is a non-trivial knot in S^3 , then does $\mathcal{S}_{\text{LO}}(K)$ contain $(-\infty, 1) \cap \mathbb{Q}$ or $(-1, \infty) \cap \mathbb{Q}$?*

For the simplest non-trivial knot $T_{3,2}$ (resp. $T_{-3,2}$), we have $\mathcal{S}_{\text{LO}}(T_{3,2}) = (-\infty, 1) \cap \mathbb{Q}$ (resp. $\mathcal{S}_{\text{LO}}(T_{-3,2}) = (-1, \infty) \cap \mathbb{Q}$); see Remark 1.2(2) and Example 1.6.

Question 8.3. *If $\mathcal{S}_{\text{LO}}(K) = (-\infty, 1) \cap \mathbb{Q}$ or $\mathcal{S}_{\text{LO}}(K) = (-1, \infty) \cap \mathbb{Q}$, then is K a trefoil knot $T_{3,2}$ or $T_{-3,2}$, respectively?*

Question 8.4. *Let K be a non-trivial knot in S^3 . Then does $\mathcal{S}_{\text{LO}}(K)$ have a maximum or minimum?*

Conjecture 1.3 says that $\mathcal{S}_L(K)$ and $\mathcal{S}_{\text{LO}}(K)$ are complementary to each other in \mathbb{Q} if K is not a cable of a non-trivial knot. So let us look at the shape of $\mathcal{S}_L(K)$, which is described by Proposition 9.6 in [48] ([23, Lemma 2.13]).

Theorem 8.5 [23, 48]. *Suppose that K is a non-trivial knot and $\mathcal{S}_L(K) \neq \emptyset$. Then $\mathcal{S}_L(K) = [2g(K) - 1, \infty) \cap \mathbb{Q}$ or $\mathcal{S}_L(K) = (-\infty, -2g(K) + 1] \cap \mathbb{Q}$.*

Theorem 8.5 makes us expect the following explicit form of $\mathcal{S}_{\text{LO}}(K)$.

Conjecture 8.6. *Let K be a non-trivial knot in S^3 which is not a cable of a non-trivial knot. Then $\mathcal{S}_{\text{LO}}(K)$ coincides with one of \mathbb{Q} , $(-\infty, 2g(K) - 1) \cap \mathbb{Q}$ or $(-2g(K) + 1, \infty) \cap \mathbb{Q}$.*

Finally, we give a comment on Question 8.3 in case of $\mathcal{S}_{\text{LO}}(K) = (-\infty, 1) \cap \mathbb{Q}$; the other case follows by taking the mirror image. By the assumption $1 \notin \mathcal{S}_{\text{LO}}(K)$. If Conjecture 1.1 is true, then $1 \in \mathcal{S}_L(K)$ or $K(1)$ is reducible. The latter possibility is eliminated by [18, Corollary 3.1], and hence $K(1)$ is an L -space. Then Proposition 8.7 [24, Proposition 6] below shows that K is a trefoil knot $T_{3,2}$.

Proposition 8.7 [24]. *Suppose K is a non-trivial knot and $K(\frac{1}{n})$ is an L -space. Then $n = 1$ (resp. -1) and K is a trefoil knot $T_{3,2}$ (resp. $T_{-3,2}$).*

Acknowledgments

We would like to thank Cameron Gordon, Hiroshi Matsuda, Yi Ni and Motoo Tange for private communications concerning L -spaces, and thank Joshua Greene and Adam Levine for suggesting the use of Alexander polynomials in the proof of Proposition 7.3. We would also like to thank Adam Clay for informing us his curious example mentioned in Remark 7.6(3), and Kazuhiro Ichihara for letting us know about the paper [30]. Finally, we would like to thank the referees for their careful reading and useful comments.

The first author has been partially supported by JSPS Grants-in-Aid for Scientific Research (C), 21540098, The Ministry of Education, Culture, Sports, Science and Technology, Japan and Joint Research Grant of Institute of Natural Sciences at Nihon University for 2013.

The second author has been partially supported by JSPS Grant-in-Aid for Scientific Research (C), 25400093, The Ministry of Education, Culture, Sports, Science and Technology, Japan.

References

- [1] I. Aitchison, *Isotoping and twisting knots and ribbons*, Thesis—University of California, Berkeley, May 1984.
- [2] M. Boileau and S. Boyer, *Graph manifolds \mathbb{Z} -homology 3-spheres and Taut foliations*, preprint.
- [3] M. Boileau and B. Zimmermann, *The π -orbifold group of a link*, Math. Z., **200** (1989), 187–208.
- [4] S. Boyer, C.McA. Gordon and L. Watson, *On L -spaces and left-orderable fundamental groups*, Math. Ann. **356** (2013), 1213–1245.
- [5] S. Boyer, D. Rolfsen and B. Wiest, *Orderable 3-manifold groups*, Ann. Inst. Fourier, **55** (2005), 243–288.
- [6] G. Burde and H. Zieschang, *Knots, de Gruyter Studies in Mathematics*, 5. Walter de Gruyter & Co., Berlin, 2003.
- [7] A. Clay, private communication.
- [8] A. Clay, T. Lidman and L. Watson, *Graph manifolds, left-orderability and amalgamation*, Algebr. Geom. Topol. **13** (2013), 2347–2368.

- [9] A. Clay and M. Teragaito, *Left-orderability and exceptional Dehn surgery on two-bridge knots*, Contemp. Math. Amer. Math. Soc. **597** (2013), 225–233.
- [10] A. Clay and L. Watson, *On cabled knots, Dehn surgery, and left-orderable fundamental groups*, Math. Res. Lett. **18** (2011), 1085–1095.
- [11] A. Clay and L. Watson, *Left-orderable fundamental groups and Dehn surgery*, Int. Math. Res. Not., IMRN 2013, no. 12, 2862–2890.
- [12] A. Edmonds, *Least area Seifert surfaces and periodic knots*, Topology Appl. **18** (1984), 109–113.
- [13] A. Edmonds and C. Livingston, *Group actions on fibered three-manifolds*, Comment. Math. Helv. **58** (1983), 529–542.
- [14] S. Fenley, *Anosov flows in 3-manifolds*, Ann. Math. **139** (1994), 79–115.
- [15] D. Gabai, *Foliations and the topology of 3-manifolds. III*, J. Differ. Geom. **26** (1987), 479–536.
- [16] F. González-Acuña and H. Short, *Knot surgery and primeness*, Math. Proc. Camb. Phil. Soc. **99** (1986), 89–102.
- [17] C.McA. Gordon, *Dehn surgery and satellite knots*, Trans. Amer. Math. Soc. **275** (1983), 687–708.
- [18] C.McA. Gordon and J. Luecke, *Knots are determined by their complements*, J. Amer. Math. Soc. **2** (1989), 371–415.
- [19] J. Greene, *Alternating links and left-orderability*, preprint.
- [20] R. Hakamata and M. Teragaito, *Left-orderable fundamental groups and Dehn surgery on genus one two-bridge knots*, to appear in Algebr. Geom. Topol.
- [21] C. Hayashi and K. Motegi, *Dehn surgery on knots in solid tori creating essential annuli*, Trans. Amer. Math. Soc. **349** (1997), 4897–4930.
- [22] C. Hayashi and K. Shimokawa, *Symmetric knots satisfy the cabling conjecture*, Math. Proc. Camb. Phil. Soc. **123** (1998), 501–529.
- [23] M. Hedden, *On knot Floer homology and cabling. II*, Int. Math. Res. Not. IMRN, (12) (2009), 2248–2274.
- [24] M. Hedden and L. Watson, *Does Khovanov homology detect the unknot?*, Amer. J. Math. **132** (2010), 1339–1345.

- [25] M. Hirasawa, K. Murasugi and D. Silver, *When does a satellite knot fiber?*, Hiroshima Math. J. **38** (2008), 411–423.
- [26] K. Ichihara and I.D. Jong, *Cyclic and finite surgeries on Montesinos knots*, Algebr. Geom. Topol. **9** (2009), 731–742.
- [27] T. Ito, *Non-left-orderable double branched coverings*, Algebr. Geom. Topol. **13** (2013), 1937–1965.
- [28] W. Jaco and P. Shalen, *Seifert fibered spaces in 3-manifolds*, Mem. Amer. Math. Soc. **21**(220) (1979), viii+192.
- [29] K. Johannson, *Homotopy equivalences of 3-manifolds with boundaries*, Lect. Notes in Math. **761**, Springer-Verlag, Berlin 1979.
- [30] J. Jun, *$(-2, 3, 7)$ -pretzel knot and Reebless foliation*, Topology Appl. **145** (2004), 209–232.
- [31] A. Kawachi, *A survey of knot theory*, Birkhäuser-Verlag, Basel, Boston, and Berlin, 1996.
- [32] T. Kitano and M. Suzuki, *A partial order in the knot table*, Exp. Math. **14** (2005), 385–390.
- [33] T. Li and R. Roberts, *Taut foliations in knot complements*, to appear in Pacific J. Math.
- [34] E. Luft and X. Zhang, *Cabling conjecture for symmetric knots*, Math. Ann. **298** (1994), 489–496.
- [35] W. Meeks and P. Scott, *Finite group actions on 3-manifolds*, Invent. Math. **86** (1986), 287–346.
- [36] W. Meeks and S.-T. Yau, *The equivariant Dehn’s lemma and loop theorem*, Comment. Math. Helv. **56** (1981), 225–239.
- [37] W. Menasco, *Closed incompressible surfaces in alternating knots and link complements*, Topology **23** (1984), 37–44.
- [38] K. Miyazaki and K. Motegi, *Seifert fibered manifolds and Dehn surgery III*, Commun. Anal. Geom. **7** (1999), 551–582.
- [39] K. Miyazaki and K. Motegi, *Toroidal surgery on periodic knots*, Pacific J. Math. **193** (2000), 381–396.
- [40] J. Morgan and H. Bass (eds.), *The Smith conjecture*, Academic Press, Orlando, FL, 1984.

- [41] K. Murasugi, *On the Alexander polynomial of the alternating knot*, Osaka Math. J. **10** (1958), 181–189.
- [42] S. Naik, *Periodicity, genera and Alexander polynomials of knots*, Pacific J. Math. **166** (1994), 357–371.
- [43] Y. Ni, *Knot Floer homology detects fibred knots*, Invent. Math. **170** (2007), 577–608.
- [44] Y. Ni, *Erratum: Knot Floer homology detects fibred knots*, Invent. Math. **177** (2009), 235–238.
- [45] P. Ozsváth and Z. Szabó, *Holomorphic disks and topological invariants for closed three-manifolds*, Ann. Math. **159** (2004), 1027–1158.
- [46] P. Ozsváth and Z. Szabó, *Holomorphic disks and three-manifold invariants: properties and applications*, Ann. Math. **159** (2004), 1159–1245.
- [47] P. Ozsváth and Z. Szabó, *On knot Floer homology and lens space surgeries*, Topology **44** (2005), 1281–1300.
- [48] P. Ozsváth and Z. Szabó, *Knot Floer homology and rational surgeries*, Algebr. Geom. Topol. **11** (2011), 1–68.
- [49] T. Peters, *On L -spaces and non left-orderable 3-manifold groups*, preprint.
- [50] D. Rolfsen, *Knots and links*, Publish or Perish, Berkeley, 1976.
- [51] M. Scharlemann, *Producing reducible 3-manifolds by surgery on a knot*, Topology **29** (1990), 481–500.
- [52] P.A. Smith, *Transformations of finite period*, Ann. Math. **39** (1938), 127–164.
- [53] Z. Szabó, *Lecture Notes on Heegaard Floer Homology*, LAS/Park City Mathematics Series, **15**, 2006, 199–228.
- [54] M. Teragaito, *Left-orderability and exceptional Dehn surgery on twist knots*, Canad. Math. Bull. **56** (2013), 850–859.
- [55] W. Thurston, *A norm for the homology of 3-manifolds*, Mem. Amer. Math. Soc. **59**(339) (1986), i–vi, 99–130.
- [56] A. Tran, *On left-orderable fundamental groups and Dehn surgeries on double twist knots*, to appear in J. Math. Soc. Japan.

DEPARTMENT OF MATHEMATICS
NIHON UNIVERSITY
TOKYO 156-8550
JAPAN

E-mail address: `motegi@math.chs.nihon-u.ac.jp`

DEPARTMENT OF MATHEMATICS AND MATHEMATICS EDUCATION
HIROSHIMA UNIVERSITY
1-1-1 KAGAMIYAMA, HIGASHI-HIROSHIMA 739-8524
JAPAN

E-mail address: `teragai@hiroshima-u.ac.jp`

RECEIVED JULY 22, 2013

