

Second-order equations and local isometric immersions of pseudo-spherical surfaces

NABIL KAHOUADJI, NIKY KAMRAN, AND KETI TENENBLAT

We consider the class of differential equations that describe pseudo-spherical surfaces of the form $u_t = F(u, u_x, u_{xx})$ and $u_{xt} = F(u, u_x)$. We answer the following question: Given a pseudo-spherical surface determined by a solution u of such an equation, do the coefficients of the second fundamental form of the local isometric immersion in \mathbb{R}^3 depend on a jet of finite order of u ? We show that, except for the sine-Gordon equation, where the coefficients depend on a jet of order zero, for all other differential equations, whenever such an immersion exists, the coefficients are universal functions of x and t , independent of u .

1. Introduction

The class of partial differential equations describing pseudo-spherical surfaces, which has been defined and studied in depth in a foundational paper by Chern and Tenenblat [3], contains a large subclass of equations enjoying remarkable integrability properties, such as the existence of infinite hierarchies of conservation laws, Bäcklund transformations and associated linear problems. Recall that a partial differential equation

$$(1) \quad \Delta \left(t, x, u, \frac{\partial u}{\partial x}, \frac{\partial u}{\partial t}, \dots, \frac{\partial^k u}{\partial t^l \partial x^{k-l}} \right) = 0,$$

is said to describe pseudo-spherical surfaces if there exist 1-forms

$$(2) \quad \omega^i = f_{i1}dx + f_{i2}dt, \quad 1 \leq i \leq 3,$$

2010 Mathematics Subject Classification: 35L60, 37K25, 47J35, 53B10, 53B25.

Key words and phrases: evolution equations, nonlinear hyperbolic equations, pseudo-spherical surfaces, isometric immersions.

Research partially supported by a CRM-ISM post-doctoral Fellowship, by NSERC Grant RGPIN 105490-2011 and by the Ministério de Ciência e Tecnologia, Brazil, CNPq Proc. No. 303774/2009-6.

where the coefficients f_{ij} , $1 \leq i \leq 3$, $1 \leq j \leq 2$, are smooth functions of t, x, u and finitely many derivatives of u with respect to t and x , such that the structure equations

$$(3) \quad d\omega^1 = \omega^3 \wedge \omega^2, \quad d\omega^2 = \omega^1 \wedge \omega^3, \quad d\omega^3 = \omega^1 \wedge \omega^2$$

hold if, and only if, u is a solution of (1) for which $\omega^1 \wedge \omega^2 \neq 0$. In other words, every smooth solution of an equation (1) describing pseudo-spherical surfaces defines on its domain $U \subset \mathbb{R}^2$ a Riemannian metric

$$(4) \quad ds^2 = (\omega^1)^2 + (\omega^2)^2,$$

of constant Gaussian curvature equal to -1 , with ω^3 being the Levi-Civita connection 1-form of the metric (4).

One of the most important examples of a partial differential equation describing pseudo-spherical surfaces is the sine-Gordon equation

$$(5) \quad \frac{\partial^2 u}{\partial t \partial x} = \sin u,$$

for which a choice of 1-forms (2) satisfying the structure equations (3) is given by

$$(6) \quad \omega^1 = \cos \frac{u}{2} (dx + dt),$$

$$(7) \quad \omega^2 = \sin \frac{u}{2} (dx - dt),$$

$$(8) \quad \omega^3 = \frac{u_x}{2} dx - \frac{u_t}{2} dt.$$

It should be noted that this choice of 1-forms is by no means unique. In particular, we could also have used

$$(9) \quad \omega^1 = \frac{1}{\eta} \sin u dt,$$

$$(10) \quad \omega^2 = \eta dx + \frac{1}{\eta} \cos u dt,$$

$$(11) \quad \omega^3 = u_x dx,$$

where η is a continuous non-vanishing real parameter. This continuous parameter is closely related to the parameter appearing in the classical Bäcklund transformation for the sine-Gordon equation and accounts for the existence of infinitely many conservation laws for the sine-Gordon equation. More

generally, partial differential equations (1) which describe pseudo-spherical surfaces and for which one of the components f_{ij} (say f_{21}) can be chosen to be a continuous parameter will be said to describe η pseudo-spherical surfaces.

In [3], Chern and Tenenblat provided a complete classification of the evolution equations of the form

$$(12) \quad u_t = F(u, u_x, \dots, \partial^k u / \partial x^k),$$

which describe pseudo-spherical surfaces under the assumption that $f_{21} = \eta$, where η is a real parameter, providing an extensive class of non-linear partial differential equations, in two independent variables, describing pseudo-spherical surfaces. Rabelo in [10], [11] characterized equations of the form $u_{xt} = F(u, u_x, \dots, \partial u / \partial x^k)$, with $f_{21} = \eta$. The complete classification for equations of type $u_{xt} = F(u, u_x)$ and $u_t = u_{xxx} + G(u, u_x, u_{xx})$ was given in [12] and [13], respectively.

In general, the importance of the class of differential equations that describe pseudo-spherical surfaces is due to the fact that such a differential equation is always the integrability condition of a linear system of differential equations, which may be used in the inverse scattering method to solve the differential equation (see for example [1], where the method was applied to a subclass of equations obtained in [11]). While the assumption of $f_{21} = \eta$ is natural in the context of the inverse scattering method, the problem of classifying the differential equations describing pseudo-spherical surfaces, without any other assumption, is important in its own right and was considered by Kamran and Tenenblat in [8], where one can find a complete classification of evolution equations of the form (12) which describe pseudo-spherical surfaces, as opposed to η pseudo-spherical surfaces. These results provide a systematic way of verifying if a given differential equation of this type describes pseudo-spherical surfaces. The results obtained in [8] were extended by Reyes in [14] to differential equations of the form $u_t = F(x, t, u, u_x, \dots, \partial^k u / \partial x^k)$. The concept of a differential equation that describes pseudo-spherical surfaces was extended by Ding and Tenenblat in [4] to a system of differential equations that describes constant curvature surfaces (pseudo-spherical and also spherical), where classification results for such systems were obtained. More recently, in order to determine new classes of differential equations that describe pseudo-spherical surfaces, as a consequence of [8], assuming that f_{21} and f_{31} are linear combinations of f_{11} , Gomes [6] classified and obtained large new classes of such equations by considering fifth order equations of type (12).

We should point out that the classification results mentioned above, contain not only general statements, but also examples of interesting new and well known non linear differential equations. Other aspects of the theory of differential equations which describe pseudo-spherical surfaces and its applications thereof can be found in [2], [7], [9], [5], [14]-[18].

A classical theorem in the theory of surfaces states that any pseudo-spherical surface can be locally isometrically immersed into three-dimensional Euclidean space \mathbb{E}^3 . This result can thus be applied to the metrics arising from the solutions u of any partial differential equation (1) describing pseudo-spherical surfaces, thereby associating to any solution u a local isometric immersion of a metric with constant Gaussian curvature equal to -1 . This theorem is however largely an existence result, which does not give an explicit expression for the second fundamental form of the local isometric immersion. It is therefore a most remarkable property of the sine-Gordon equation that the second fundamental form of any such immersion can be expressed in closed form as a function of u and finitely many derivatives. Indeed, let us first recall that the components a, b, c of the second fundamental form of any local isometric immersion of a metric of constant curvature equal to -1 into \mathbb{E}^3 are defined by the 1-forms ω_1^3, ω_2^3 according to

$$(13) \quad \omega_1^3 = a\omega^1 + b\omega^2, \quad \omega_2^3 = b\omega^1 + c\omega^2,$$

where these forms satisfy the structure equations

$$(14) \quad d\omega_1^3 = -\omega_2^3 \wedge \omega^3, \quad d\omega_2^3 = \omega_1^3 \wedge \omega^3,$$

and the Gauss equation

$$ac - b^2 = -1.$$

For the sine-Gordon equation, with the choice of 1-forms ω^1, ω^2 and ω^3 given by (6), (7) and (8), it is easily verified that the 1-forms ω_1^3, ω_2^3 are given by

$$\begin{aligned} \omega_1^3 &= \sin \frac{u}{2} (dx + dt) = \tan \frac{u}{2} \omega^1, \\ \omega_2^3 &= -\cos \frac{u}{2} (dx - dt) = -\cot \frac{u}{2} \omega^2. \end{aligned}$$

In general, given a partial differential equation (1) describing pseudo-spherical surfaces, it is straightforward to derive a set of necessary and sufficient conditions, in terms of the coefficients f_{ij} of the 1-forms (2), for a, b and c to be the components of the second fundamental form of a local isometric

immersion corresponding to a solution of (1). We write

$$(15) \quad d\omega_1^3 = (db(e_1) - da(e_2))\omega^1 \wedge \omega^2 - a\omega^2 \wedge \omega^3 + b\omega^1 \wedge \omega^3,$$

$$(16) \quad d\omega_2^3 = (dc(e_1) - db(e_2))\omega^1 \wedge \omega^2 - b\omega^2 \wedge \omega^3 + c\omega^1 \wedge \omega^3,$$

where (e_1, e_2) is the pair of vector fields dual to the coframe (ω^1, ω^2) , given by

$$\begin{vmatrix} f_{11} & f_{21} \\ f_{12} & f_{22} \end{vmatrix} e_1 = f_{22}\partial_x - f_{21}\partial_t, \quad \begin{vmatrix} f_{11} & f_{21} \\ f_{12} & f_{22} \end{vmatrix} e_2 = -f_{12}\partial_x + f_{11}\partial_t.$$

Thus, using the notation D_t and D_x for the total derivative operators, we obtain

$$(17) \quad f_{11}D_t a + f_{21}D_t b - f_{12}D_x a - f_{22}D_x b - 2b \begin{vmatrix} f_{11} & f_{31} \\ f_{12} & f_{32} \end{vmatrix} + (a - c) \begin{vmatrix} f_{21} & f_{31} \\ f_{22} & f_{32} \end{vmatrix} = 0,$$

$$(18) \quad f_{11}D_t b + f_{21}D_t c - f_{12}D_x b - f_{22}D_x c + (a - c) \begin{vmatrix} f_{11} & f_{31} \\ f_{12} & f_{32} \end{vmatrix} + 2b \begin{vmatrix} f_{21} & f_{31} \\ f_{22} & f_{32} \end{vmatrix} = 0,$$

where a, b and c , which are assumed to depend on t, x, u and finitely many derivatives of u with respect to t and x , satisfy the Gauss equation

$$(19) \quad ac - b^2 = -1.$$

In view of the above discussion, it is natural to ask the following question: *Do there exist equations other than the sine-Gordon equation within the class of partial differential equations describing pseudo-spherical (or η pseudo-spherical) surfaces, for which the components a, b, c of the second fundamental form of the local isometric immersion depend on a jet of finite order of u , that is on x, t, u and finitely many derivatives of u ?*

If such equations were to exist, they would have an important geometric property in common with the sine-Gordon equation. In this paper, we give a complete answer to the above question in the case of second-order hyperbolic equations of the form

$$(20) \quad u_{xt} = F(u, u_x),$$

and evolution equations of the form

$$(21) \quad u_t = F(u, u_x, u_{xx}),$$

which describe η pseudo-spherical surfaces as in [12] and [3].

We begin with the case of evolution equations (21), for which our main result is the following:

Theorem 1. *Except for second-order evolution equations of the form*

$$(22) \quad u_t = \frac{f_{12,u_x}}{f_{11,u}} u_{xx} + \frac{f_{12,u}}{f_{11,u}} u_x \mp \frac{\lambda f_{11} - \eta f_{12}}{f_{11,u}},$$

where $f_{11,u} \neq 0$ and $f_{12,u_x} \neq 0$, there exists no second-order evolution equation describing η pseudo-spherical surfaces, given as in [3], with the property that the coefficients of the second fundamental forms of the local isometric immersions of the surfaces associated to the solutions u of the equation depend on a jet of finite order of u . Moreover, the coefficients of the second fundamental forms of the local isometric immersions of the surfaces determined by the solutions u of (22) are universal, i.e., they are universal functions of x and t , independent of u .

Theorem 1 suggests that there is no real analogue of the sine-Gordon equation within the class of second-order evolution equations describing η pseudo-spherical surfaces, from the perspective of the local isometric immersions of pseudo-spherical surfaces associated to their solutions. Indeed, even though the special class of evolution equations (22) has the property that the components of the second fundamental forms of the immersions associated to its solutions depend on jets of finite order of u , this dependence is given in terms of functions of x and t for all choices of solutions u (see Proposition 2).

The results for second-order hyperbolic equations (20) are similar, with the notable exception that they single out the sine-Gordon equation as the only equation, up to constants, for which the second fundamental form of the local isometric immersion is not universal. In order to state these results, we begin by recalling the classification theorem proved by Rabelo and Tenenblat [12] for equations (20) describing pseudo-spherical surfaces:

Theorem 2 (Rabelo & Tenenblat [12]). *Let F be a differentiable function defined on an open connected subset $U \subset \mathbb{R}^2$. An equation*

$$u_{xt} = F(u, u_x)$$

describes an η pseudo-spherical surface for $\eta \in \mathcal{P} \subset \mathbb{R}$, where \mathcal{P} is a dense subset of \mathbb{R} and F independent of η if, and only if, F satisfies one of the following:

- i) F is independent of u_x and $F''(u) + \alpha F(u) = 0$, $U = \mathbb{R}^2$, $\mathcal{P} = \mathbb{R} \setminus \{0\}$, and α is a non-zero real constant.
- ii) $F = \nu e^{\delta u} \sqrt{\beta + \gamma u_x^2}$, where $U = \{(u, z) \in \mathbb{R}^2; \beta + \gamma z^2 > 0\}$, $\mathcal{P} = \mathbb{R}$, $\delta, \gamma, \beta, \nu$ are real constants, with δ, γ, ν nonzero, and $\beta = 0$ when $\gamma = 1$; or
- iii) $F = \lambda u + \zeta u_x + \tau$, where $U = \mathbb{R}^2$, $\mathcal{P} = \mathbb{R} \setminus \{0\}$, and λ, ζ, τ are real constants.

The expressions of functions f_{ij} of the 1-forms ω^i for each equation of Theorem 2 are recalled in Section 4 (Lemmas 6–8). We are now ready to state our main result for the case of second-order hyperbolic equations (20).

Theorem 3. *Let F be an equation of the form $u_{xt} = F(u, u_x)$ that describes η pseudo-spherical surfaces as in Theorem 2, with the 1-forms as in [12].*

- 1) *If F is independent of u_x and satisfies $F''(u) + \alpha F(u) = 0$, where α is a positive real constant, then there exists a local isometric immersion in \mathbb{R}^3 of the pseudo-spherical surface determined by a solution u , for which the coefficients of the second fundamental form depend on a jet of finite order of u if, and only if, they depend on the jet of order zero.*
- 2) *If $F = \lambda u + \zeta u_x + \tau$, then there exists a local isometric immersion in \mathbb{R}^3 of the pseudo-spherical surface determined by a solution u , for which the coefficients of the second fundamental form depend on a jet of finite order of u if, and only if, λ, ξ and τ do not vanish simultaneously, and the coefficients are independent of u , that is they are universal functions of x and t .*
- 3) *For the remaining equations, that is, if F is independent of u_x and satisfies $F''(u) + \alpha F(u) = 0$, where α is a negative real constant, $F = \nu e^{\delta u} \sqrt{\beta + \gamma u_x^2}$ and $F = 0$, there is no local isometric immersion of the pseudo-spherical surface determined by a solution u , for which the coefficients of the second fundamental form depend on a jet of finite order of u .*

The coefficients of the second fundamental form of the local isometric immersions stated in Theorem 3 are given explicitly in Section 4 (Propositions 3 and 5). Theorem 3 shows likewise that when viewed through the perspective of the local isometric immersions associated to its solutions, the sine-Gordon equation occupies a special position within the class of hyperbolic equations (20) as the unique equation, up to normalization constants,

for which the coefficients of the second fundamental form of the local isometric immersion of the surface determined by a solution u , depends on a jet of finite order of u , without being universal, *i.e.* independent of u .

While Theorems 1 and 3 give a complete answer to the general question we have raised in this paper in the case of second-order evolution equations (21) and second-order hyperbolic equations (20), the question still remains open for all the other classes of equations describing pseudo-spherical surfaces. We believe that it should be possible to extend the proof of Theorem 1 to the case of k -th order evolution equations with $k \geq 3$ in order to obtain a similar result to the effect that all the second-fundamental forms that depend only on jets of finite order of the solutions of evolution equation should be universal.

Our paper is organized as follows. In Section 2, we recall the results of Chern and Tenenblat [3] on the classification of evolution equations describing pseudo-spherical surfaces and use these to give an analogue of the normal forms of Theorem 2 for the case of second-order evolution equations (21). These normal forms are then used as the starting point in Section 3 of the proof of Theorem 1. Section 4 is devoted to the proof of Theorem 3. The proofs involve a careful analysis of the possible dependence on higher-order jets of u of the solutions of the system of differential constraints (17) and (18) that must be satisfied by components a, b, c of the second fundamental form, together with the algebraic constraint given by the Gauss equation (19).

2. The classification of second-order evolution equations describing η pseudo-spherical surfaces

In [3], Chern and Tenenblat obtained necessary and sufficient conditions in the form of differential equations on the functions f_{ij} for the existence of an evolution equation of the form

$$(23) \quad \frac{\partial u}{\partial t} = F \left(u, u_x, \dots, \frac{\partial^k u}{\partial x^k} \right),$$

which describes η pseudo-spherical surfaces, *i.e.*, with $f_{21} = \eta$, where η is a nonzero parameter. They also performed a complete classification of the evolution equations of the form (23) which describe η pseudo-spherical surfaces. They obtained four classes of evolution equations (Theorems 2.2 to 2.5 in [3]). These four classes of equations are determined algebraically by f_{11}, f_{31}, f_{22} and their derivatives, up to some differential constraints. In what follows, we consider only second-order evolution equations of the form (23) and solve

the differential constraints that f_{11}, f_{31} and f_{22} must satisfy in order for (23) to describe η pseudo-spherical surfaces. We shall deal with two of the four classes (Theorems 2.2 and 2.4 in [3]) since the two remaining classes of evolution equations (Theorems 2.3 and 2.5 in [3]) lead to evolution equations of the first order, when $k = 2$.

It will be convenient to introduce the following notation for the spatial derivatives of u (used in [3] and also in [8]),

$$z_i = \frac{\partial^i u}{\partial x^i}, \quad 0 \leq i \leq k,$$

and to view $(x, t, z_0, z_1, \dots, z_k)$ as local coordinates in an open subset U of a manifold.

Lemma 1 (Chern & Tenenblat [3]). *Consider a second-order evolution equation of the form $z_{0,t} = F(z_0, z_1, z_2)$ which describes an η pseudo-spherical surface with associated forms $\omega^i = f_{i1}dx + f_{i2}dt$. If f_{ij} are differentiable functions of z_0, z_1, z_2 , then*

$$(24) \quad f_{ij,z_2} = 0, \quad f_{11,z_1} = f_{31,z_1} = f_{22,z_1} = 0,$$

$$(25) \quad f_{11,z_0}^2 + f_{31,z_0}^2 \neq 0.$$

In order to state the results, we introduce the following notation

$$(26) \quad \begin{aligned} H &= f_{11}f_{11,z_0} - f_{31}f_{31,z_0}, & L &= f_{11}f_{31,z_0} - f_{31}f_{11,z_0}, \\ P &= f_{11,z_0}f_{31,z_0z_0} - f_{31,z_0}f_{11,z_0z_0}, & M &= f_{31,z_0}^2 - f_{11,z_0}^2. \end{aligned}$$

Lemma 2. *Let f_{ij} , $1 \leq i \leq 3$, $1 \leq j \leq 2$, be differentiable functions of z_0, z_1, z_2 such that (24) and (25) hold and $f_{21} = \eta$ a nonzero parameter. Suppose $HL \neq 0$. Then $z_{0,t} = F(z_0, z_1, z_2)$ describes an η pseudo-spherical surface with associated 1-forms $\omega^i = f_{i1}dx + f_{i2}dt$, if and only if*

$$(27) \quad \begin{aligned} F &= \frac{\mp f_{22,z_0}}{\eta\sqrt{1-\alpha^2}f_{11,z_0}}z_2 \mp \frac{f_{22,z_0z_0}}{\eta\sqrt{1-\alpha^2}f_{11,z_0}}z_1^2 \\ &+ \left(\frac{(\eta^2 + f_{11}^2 - f_{31}^2)f_{22,z_0}}{\eta[(1-\alpha^2)f_{11} \mp \alpha\eta\sqrt{1-\alpha^2}]f_{11,z_0}} + \frac{f_{22}}{\eta} \right) z_1, \end{aligned}$$

and

$$(28) \quad f_{31} = \alpha f_{11} \pm \eta \sqrt{1 - \alpha^2},$$

$$(29) \quad f_{12} = \frac{f_{11} f_{22}}{\eta} \mp \frac{f_{22, z_0}}{\eta \sqrt{1 - \alpha^2}} z_1,$$

$$(30) \quad f_{32} = \frac{(\alpha f_{11} \pm \eta \sqrt{1 - \alpha^2}) f_{22}}{\eta} \mp \frac{\alpha f_{22, z_0}}{\eta \sqrt{1 - \alpha^2}} z_1,$$

where $f_{22, z_0} \neq 0$, $f_{11, z_0} \neq 0$, and $\alpha^2 < 1$.

Proof. If $k = 2$, Theorem 2.2 in [3] gives the general expression of second-order evolution equations $z_{0,t} = F$ which describe η pseudo-spherical surfaces, namely

$$(31) \quad F = \frac{1}{L} \sum_{i=0}^1 z_{i+1} B_{z_i} + \frac{1}{HL} \left(-z_1 \frac{L}{\eta} + f_{31}^2 - f_{11}^2 \right) z_1 A^0 \\ + \frac{B}{HL} (z_1 M + \eta L) + z_1 \frac{f_{22}}{\eta},$$

where

$$B = f_{22, z_0} z_1, \quad A^0 = \frac{1}{L} (-z_1 P + \eta M) B_{z_1} + f_{22, z_1} H,$$

where the functions f_{12} and f_{32} are given by

$$(32) \quad f_{12} = \frac{f_{11} f_{22}}{\eta} + \frac{1}{H} \left(-\frac{f_{11} A^0}{\eta} z_1 + f_{31, z_0} B \right), \\ f_{32} = \frac{f_{31} f_{22}}{\eta} + \frac{1}{H} \left(-\frac{f_{31} A^0}{\eta} z_1 + f_{11, z_0} B \right),$$

and where (2.12) in [3] gives two differential equations that the functions f_{11} , f_{31} and f_{22} must satisfy. When $k = 2$, these equations reduce to

$$(33) \quad \frac{L}{\eta} f_{22, z_0} - \frac{L}{\eta} \left(z_1 \frac{A^0}{H} \right)_{z_0} + A^0 + \frac{M}{H} B_{z_0} + \frac{B}{H^2} (LP + M^2) = 0,$$

$$(34) \quad \frac{L}{\eta} f_{22, z_1} - \frac{L}{\eta} \left(z_1 \frac{A^0}{H} \right)_{z_1} + \frac{M}{H} B_{z_1} = 0.$$

If $L \neq 0$, then the differential equation (34) leads to $P f_{22, z_0} = 0$. The vanishing of f_{22, z_0} contradicts the fact that F is a second order evolution

equation. We conclude then that

$$(35) \quad f_{22,z_0} \neq 0, \quad P = 0.$$

Differentiating (33) with respect to z_1 leads to $-L(M/HL)_{z_0} + M^2/L^2 = 0$ and hence, the differential equation (33) leads to

$$(36) \quad M = -\frac{L^2}{\eta^2}.$$

The vanishing of P implies that

$$(37) \quad f_{31} = \alpha f_{11} + \beta, \quad \text{where } \alpha, \beta \in \mathbb{R}.$$

We have then

$$(38) \quad L = -\beta f_{11,z_0}, \quad H = [(1 - \alpha^2)f_{11} - \alpha\beta]f_{11,z_0}.$$

The non-vanishing of L implies that $\beta \neq 0$ and $f_{11,z_0} \neq 0$. Substituting (37) in (36) and in the expression of M as in (26) leads to

$$(39) \quad \beta^2 = \eta^2(1 - \alpha^2).$$

The non-vanishing of β and η , and the latter equation imply that $\alpha \in (-1, 1)$. Finally, substituting $\beta = \pm\eta\sqrt{1 - \alpha^2}$, $P = 0$ and (37) in the expressions (31) and (32) leads to expressions (27), (28), (29), and (30). \square

If $HL = 0$, then there are three classes of evolution equations to consider, which are given in Theorems 2.3-2.5 in [3]. However, F is of second order only when $H = L = 0$, as in Theorem 2.4 in [3].

Lemma 3. *Let f_{ij} , $1 \leq i \leq 3$, $1 \leq j \leq 2$, be differentiable functions of z_0, z_1, z_2 such that (24) and (25) hold and $f_{21} = \eta$ a nonzero parameter. Suppose $f_{31} = \pm f_{11} \neq 0$. Then $z_{0,t} = F(z_0, z_1, z_2)$ describes an η pseudo-spherical surface with associated 1-forms $\omega^i = f_{i1}dx + f_{i2}dt$, if and only if $f_{22} = \lambda$, where λ is constant, $f_{32} = \pm f_{12}$, and*

$$(40) \quad F = \frac{f_{12,z_1}}{f_{11,z_0}} z_2 + \frac{f_{12,z_0}}{f_{11,z_0}} z_1 \mp \frac{\lambda f_{11} - \eta f_{12}}{f_{11,z_0}}.$$

Proof. Immediate when $k = 2$ in Theorem 2.4 in [3]. \square

3. Proof of Theorem 1

Lemma 4. *Let $u_t = F(u, u_x, u_{xx})$ be a second-order evolution equation describing η pseudo-spherical surfaces as in Lemma 2 or in Lemma 3. If there exists a local isometric immersion of a surface determined by a solution u for which the coefficients of the second fundamental form (13) depend on a jet of finite order of u , i.e., a, b and c depend on $x, t, u, \dots, \partial^\ell u / \partial x^\ell$, where ℓ is finite, then a, b and c are universal, i.e., a, b and c depend only on x and t .*

Proof. Assume a, b and c depend on a jet of finite order, i.e., they depend on x, t, z_0, \dots and z_ℓ , where ℓ is fixed. Then (17) becomes

$$(41) \quad \begin{aligned} & f_{11}a_t + \eta b_t - f_{12}a_x - f_{22}b_x \\ & - 2b(f_{11}f_{32} - f_{31}f_{12}) + (a - c)(\eta f_{32} - f_{22}f_{31}) \\ & - \sum_{i=0}^{\ell} (f_{12}a_{z_i} + f_{22}b_{z_i})z_{i+1} + \sum_{i=0}^{\ell} (f_{11}a_{z_i} + \eta b_{z_i})z_{i,t} = 0, \end{aligned}$$

and (18) becomes

$$(42) \quad \begin{aligned} & f_{11}b_t + \eta c_t - f_{12}b_x - f_{22}c_x \\ & + (a - c)(f_{11}f_{32} - f_{12}f_{31}) + 2b(\eta f_{32} - f_{22}f_{31}) \\ & - \sum_{i=0}^{\ell} (f_{12}b_{z_i} + f_{22}c_{z_i})z_{i+1} + \sum_{i=0}^{\ell} (f_{11}b_{z_i} + \eta c_{z_i})z_{i,t} = 0. \end{aligned}$$

Since $f_{22,z_0} \neq 0$ and $f_{11,z_0} \neq 0$ for evolution equations (27), and $f_{11,z_0} \neq 0$ and $f_{12,z_1} \neq 0$ for evolution equations (40), differentiating (41) and (42) with respect to $z_{\ell+2}$ leads to $f_{11}a_{z_\ell} + \eta b_{z_\ell} = f_{11}b_{z_\ell} + \eta c_{z_\ell} = 0$, and hence

$$(43) \quad b_{z_\ell} = -\frac{f_{11}}{\eta} a_{z_\ell}, \quad \text{and} \quad c_{z_\ell} = \frac{f_{11}^2}{\eta^2} a_{z_\ell}.$$

Differentiating the Gauss equation (19) with respect to z_ℓ leads to $ca_{z_\ell} + ac_{z_\ell} - 2bb_{z_\ell} = 0$, and substituting (43) in the latter leads to

$$(44) \quad \left[c + \left(\frac{f_{11}}{\eta} \right)^2 a + 2\frac{f_{11}}{\eta} b \right] a_{z_\ell} = 0.$$

If $c + \left(\frac{f_{11}}{\eta} \right)^2 a + 2\frac{f_{11}}{\eta} b = 0$ on an open set, then substituting the expression of c in the Gauss equation $-ac + b^2 = 1$ leads to $(f_{11}a/\eta + b)^2 = 1$, so

that

$$b = \pm 1 - \frac{f_{11}}{\eta} a, \quad \text{and} \quad c = \left(\frac{f_{11}}{\eta} \right)^2 a \mp 2 \frac{f_{11}}{\eta}.$$

We have then

$$\begin{aligned} D_t b &= -\frac{f_{11}}{\eta} D_t a - \frac{a}{\eta} f_{11, z_0} F, \\ D_t c &= \left(\frac{f_{11}}{\eta} \right)^2 D_t a + \frac{2}{\eta} \left(\frac{f_{11}}{\eta} a \mp 1 \right) f_{11, z_0} F, \\ D_x b &= -\frac{f_{11}}{\eta} D_x a - \frac{a}{\eta} f_{11, z_0} z_1, \\ D_x c &= \left(\frac{f_{11}}{\eta} \right)^2 D_x a + \frac{2}{\eta} \left(\frac{f_{11}}{\eta} a \mp 1 \right) f_{11, z_0} z_1, \end{aligned}$$

and hence

$$(45) \quad f_{11} D_t a + \eta D_t b = -a f_{11, z_0} F,$$

$$(46) \quad f_{11} D_t b + \eta D_t c = \left(\frac{f_{11}}{\eta} a \mp 2 \right) f_{11, z_0} F,$$

$$(47) \quad f_{12} D_x a + f_{22} D_x b = -\frac{\Delta_{12}}{\eta} D_x a - \frac{a f_{22}}{\eta} f_{11, z_0} z_1,$$

$$(48) \quad \begin{aligned} f_{12} D_x b + f_{22} D_x c &= \frac{f_{11}}{\eta} \frac{\Delta_{12}}{\eta} D_x a + \frac{\Delta_{12}}{\eta^2} a f_{11, z_0} z_1 \\ &\quad + \frac{f_{22}}{\eta} \left(\frac{f_{11}}{\eta} a \mp 2 \right) f_{11, z_0} z_1. \end{aligned}$$

where $\Delta_{12} = f_{11} f_{22} - \eta f_{12}$. Substituting the latter four equalities in (17) lead to

$$\begin{aligned} &-a f_{11, z_0} F + \frac{\Delta_{12}}{\eta} D_x a + \frac{a f_{22}}{\eta} f_{11, z_0} z_1 \\ &-2b(f_{11} f_{32} - f_{31} f_{12}) + (a - c)(\eta f_{32} - f_{31} f_{22}) = 0, \end{aligned}$$

which is equivalent to

$$(49) \quad \begin{aligned} &-a f_{11, z_0} F + \frac{\Delta_{12}}{\eta} \sum_{i=0}^{\ell} a_{z_i} z_{i+1} + \frac{a f_{22}}{\eta} f_{11, z_0} z_1 \\ &-2b(f_{11} f_{32} - f_{31} f_{12}) + (a - c)(\eta f_{32} - f_{31} f_{22}) = 0. \end{aligned}$$

Substituting the four equalities (45)–(48) into (18) lead to

$$\left(\frac{f_{11}}{\eta}a \mp 2\right)f_{11,z_0}F - \frac{f_{11}}{\eta} \frac{\Delta_{12}}{\eta}D_x a - \frac{\Delta_{12}}{\eta^2}af_{11,z_0}z_1 - \frac{f_{22}}{\eta}\left(\frac{f_{11}}{\eta}a \mp 2\right)f_{11,z_0}z_1 + (a - c)(f_{11}f_{32} - f_{31}f_{12}) + 2b(\eta f_{32} - f_{31}f_{22}) = 0,$$

which is equivalent to

$$(50) \quad \left(\frac{f_{11}}{\eta}a \mp 2\right)f_{11,z_0}F - \frac{f_{11}}{\eta} \frac{\Delta_{12}}{\eta} \sum_{i=0}^{\ell} a_{z_i}z_{i+1} - \frac{\Delta_{12}}{\eta^2}af_{11,z_0}z_1 - \frac{f_{22}}{\eta}\left(\frac{f_{11}}{\eta}a \mp 2\right)f_{11,z_0}z_1 + (a - c)(f_{11}f_{32} - f_{31}f_{12}) + 2b(\eta f_{32} - f_{31}f_{22}) = 0.$$

- If $\ell \geq 2$, then differentiating (49) with respect to $z_{\ell+1}$ leads to $\Delta_{12}a_{z_\ell} = 0$. Thus $a_{z_\ell} = 0$ and also $b_{z_\ell} = c_{z_\ell} = 0$.
- If $\ell = 1$, then differentiating (49) and (50) with respect to z_2 lead to

$$-af_{11,z_0}F_{z_2} + \frac{\Delta_{12}}{\eta}a_{z_1} = 0,$$

$$\left(\frac{f_{11}}{\eta}a \mp 2\right)f_{11,z_0}F_{z_2} - \frac{f_{11}}{\eta} \frac{\Delta_{12}}{\eta}a_{z_1} = 0.$$

The latter system leads to $f_{11,z_0}F_{z_2} = 0$, which runs into a contradiction.

- If $\ell = 0$, then differentiating (49) and (50) with respect to z_2 lead to

$$-af_{11,z_0}F_{z_2} = 0,$$

$$\left(\frac{f_{11}}{\eta}a \mp 2\right)f_{11,z_0}F_{z_2} = 0.$$

The latter system leads to $f_{11,z_0}F_{z_2} = 0$, which runs into a contradiction.

Therefore, for all ℓ , (17), (18) and the Gauss equation is an inconsistent system.

If $c + \left(\frac{f_{11}}{\eta}\right)^2 a + 2\frac{f_{11}}{\eta}b \neq 0$, then $a_{z_\ell} = 0$, and hence $b_{z_\ell} = c_{z_\ell} = 0$, and successive differentiation leads to $a_{z_i} = b_{z_i} = c_{z_i} = 0$ for all $i = 0, \dots, \ell$.

Finally, if the functions a, b and c depend on a jet of finite order, then they are universal, i.e., they are functions of x and t only. □

Proposition 1. *For the second-order evolution equations which describe η pseudo-spherical surfaces as in Lemma 2, there is no local isometric immersion in \mathbb{R}^3 of a pseudo-spherical surface determined by a solution u , for which the coefficients a, b, c of the second fundamental form depend on a jet of finite order of u .*

Proof. Let a, b , and c be coefficients of the second fundamental form satisfying the Gauss equation $ac - b^2 = -1$. By Lemma 4, if a, b and c depend on a jet of finite order, then a, b and c depend only on x and t . From (28)–(30), we have

$$(51) \quad f_{11}f_{32} - f_{12}f_{31} = f_{22,z_0}z_1,$$

$$(52) \quad \eta f_{32} - f_{22}f_{31} = \mp \frac{\alpha f_{22,z_0}}{\sqrt{1 - \alpha^2}} z_1.$$

Taking into account the expressions (51), (52) and (29), equations (17) and (18) become

$$(53) \quad f_{11}a_t + \eta b_t - \left(\frac{f_{11}f_{22}}{\eta} \mp \frac{f_{22,z_0}}{\eta\sqrt{1 - \alpha^2}} z_1 \right) a_x - f_{22}b_x - 2bf_{22,z_0}z_1 \mp (a - c) \frac{\alpha f_{22,z_0}}{\sqrt{1 - \alpha^2}} z_1 = 0,$$

$$(54) \quad f_{11}b_t + \eta c_t - \left(\frac{f_{11}f_{22}}{\eta} \mp \frac{f_{22,z_0}}{\eta\sqrt{1 - \alpha^2}} z_1 \right) b_x - f_{22}c_x + (a - c)f_{22,z_0}z_1 \mp 2b \frac{\alpha f_{22,z_0}}{\sqrt{1 - \alpha^2}} z_1 = 0.$$

Differentiating (53) and (54) with respect to z_1 and the fact that $f_{22,z_0} \neq 0$ lead to

$$\begin{pmatrix} a_x \\ b_x \end{pmatrix} = \begin{pmatrix} \pm\eta\sqrt{1 - \alpha^2} & \alpha\eta \\ \alpha\eta & \mp\eta\sqrt{1 - \alpha^2} \end{pmatrix} \begin{pmatrix} 2b \\ a - c \end{pmatrix}.$$

The determinant of the 2×2 matrix appearing in the above equation is non-zero, therefore, a_x and b_x can not vanish simultaneously. Otherwise, $a - c = b = 0$, and this contradicts the Gauss equation. (53) and (54) become then

$$(55) \quad \eta f_{11}a_t + \eta^2 b_t - f_{11}f_{22}a_x - \eta f_{22}b_x = 0,$$

$$(56) \quad \eta f_{11}b_t + \eta^2 c_t - f_{11}f_{22}b_x - \eta f_{22}c_x = 0.$$

Differentiating (55) and (56) with respect to z_0 , and dividing by $\eta f_{11,z_0}$ lead to

$$(57) \quad a_t = \frac{(f_{11}f_{22})_{z_0}}{\eta f_{11,z_0}} a_x + \frac{f_{22,z_0}}{f_{11,z_0}} b_x,$$

$$(58) \quad b_t = \frac{(f_{11}f_{22})_{z_0}}{\eta f_{11,z_0}} b_x + \frac{f_{22,z_0}}{f_{11,z_0}} c_x.$$

Observe that $\frac{(f_{11}f_{22})_{z_0}}{f_{11,z_0}}$ and $\frac{f_{22,z_0}}{f_{11,z_0}}$ cannot both be constant. Otherwise, $f_{22,z_0} = 0$ which is a contradiction. Differentiating (57) and (58) with respect to z_0 leads to

$$\begin{aligned} \left(\frac{(f_{11}f_{22})_{z_0}}{\eta f_{11,z_0}} \right)_{z_0} a_x + \left(\frac{f_{22,z_0}}{f_{11,z_0}} \right)_{z_0} b_x &= 0, \\ \left(\frac{(f_{11}f_{22})_{z_0}}{\eta f_{11,z_0}} \right)_{z_0} b_x + \left(\frac{f_{22,z_0}}{f_{11,z_0}} \right)_{z_0} c_x &= 0. \end{aligned}$$

We conclude that

$$(59) \quad a_x c_x - b_x^2 = 0.$$

Subtracting (58) multiplied by a_x from (57) multiplied by b_x , it follows from (59) that

$$(60) \quad a_x b_t - a_t b_x = 0.$$

From (55), we have $f_{11}(\eta a_t - f_{22}a_x) + \eta^2 b_t - \eta f_{22}b_x = 0$. Note that $(\eta a_t - f_{22}a_x) \neq 0$. Otherwise, since $f_{22,z_0} \neq 0$, we have $a_x = a_t = 0$ and hence it follows from (59) that $b_x = 0$, which runs into a contradiction. Therefore,

$$(61) \quad f_{11} = \frac{\eta(\eta b_t - f_{22}b_x)}{\eta a_t - f_{22}a_x}.$$

Differentiating (61) with respect to z_0 and taking into account (60) lead to $f_{11,z_0} = 0$, which is a contradiction. \square

Proposition 2. *Let $u_t = F(u, u_x, u_{xx})$ be a second-order evolution equation which describes η pseudo-spherical surfaces, as in Lemma 3. There exists a local isometric immersion in \mathbb{R}^3 of a pseudo-spherical surface, determined by a solution u , for which the coefficients of the second fundamental form*

(13) depend on a jet of finite order of u if, and only if, the coefficients are universal and are given by

$$(62) \quad a = \sqrt{le^{\pm 2(\eta x + \lambda t)} - \gamma^2 e^{\pm 4(\eta x + \lambda t)} - 1},$$

$$(63) \quad b = \gamma e^{\pm 2(\eta x + \lambda t)},$$

$$(64) \quad c = \frac{\gamma^2 e^{\pm 4(\eta x + \lambda t)} - 1}{\sqrt{le^{\pm 2(\eta x + \lambda t)} - \gamma^2 e^{\pm 4(\eta x + \lambda t)} - 1}},$$

$l, \gamma \in \mathbb{R}, l > 0$ and $l^2 > 4\gamma^2$. The 1-forms are defined on a strip of \mathbb{R} where

$$(65) \quad \log \sqrt{\frac{l - \sqrt{l^2 - 4\gamma^2}}{2\gamma^2}} < \pm(\eta x + \lambda t) < \log \sqrt{\frac{l + \sqrt{l^2 - 4\gamma^2}}{2\gamma^2}}.$$

Moreover, the constants l and γ have to be chosen so that the strip intersects the domain of the solution of the evolution equation.

Proof. As for the previous proposition, if a, b and c depend on a jet of finite order, it follows from Lemma 4 that a, b and c depend only on x and t . We assume also that $f_{12, z_1} \neq 0$, otherwise, the evolution equation is not of second-order. Equations (17) and (18) become

$$(66) \quad f_{11}a_t + \eta b_t - f_{12}a_x - \lambda b_x \pm (\eta f_{12} - \lambda f_{11})(a - c) = 0,$$

$$(67) \quad f_{11}b_t + \eta c_t - f_{12}b_x - \lambda c_x \pm (\eta f_{12} - \lambda f_{11})2b = 0.$$

Differentiating (66) and (67) with respect to z_1 , and the fact that $f_{12, z_1} \neq 0$ lead to

$$(68) \quad a_x \mp \eta(a - c) = 0,$$

$$(69) \quad b_x \mp 2\eta b = 0.$$

Taking into account (68) and (69) and differentiating (66) and (67) with respect to z_0 leads to

$$(70) \quad a_t \mp \lambda(a - c) = 0,$$

$$(71) \quad b_t \mp 2\lambda b = 0,$$

and hence, (66) and (67) become

$$(72) \quad \eta b_t - \lambda b_x = 0,$$

$$(73) \quad \eta c_t - \lambda c_x = 0.$$

Note that (69) and (71) imply (72), and (68) and (70) imply

$$(74) \quad \eta a_t - \lambda a_x = 0,$$

and hence imply (73). From (69) and (71), we conclude that

$$(75) \quad b = \gamma e^{\pm 2(\eta x + \lambda t)}, \quad \gamma \in \mathbb{R}.$$

Note that $a \neq 0$. Otherwise, if $a = 0$, then (68) implies that $c = 0$ and the Gauss equation leads to $b = \pm 1$ which contradicts (69). Therefore, from the Gauss equation we have $c = (b^2 - 1)a^{-1}$. Then, in view of (75), equations (68) and (70) reduce to

$$\begin{aligned} aa_x \mp \eta(a^2 - \gamma^2 e^{\pm 4(\eta x + \lambda t)} + 1) &= 0, \\ aa_t \mp \lambda(a^2 - \gamma^2 e^{\pm 4(\eta x + \lambda t)} + 1) &= 0. \end{aligned}$$

The latter system leads then to

$$a = \sqrt{le^{\pm 2(\eta x + \lambda t)} - \gamma^2 e^{\pm 4(\eta x + \lambda t)} - 1}, \quad l \in \mathbb{R},$$

which is defined wherever $le^{\pm 2(\eta x + \lambda t)} - \gamma^2 e^{\pm 4(\eta x + \lambda t)} - 1 > 0$. Hence $l > 0$ and

$$\frac{l - \sqrt{l^2 - 4\gamma^2}}{2\gamma^2} < e^{\pm 2(\eta x + \lambda t)} < \frac{l + \sqrt{l^2 - 4\gamma^2}}{2\gamma^2},$$

i.e., a is defined on the strip described by (65). Now, from either (68) or (70), we obtain

$$c = \frac{\gamma^2 e^{\pm 4(\eta x + \lambda t)} - 1}{\sqrt{le^{\pm 2(\eta x + \lambda t)} - \gamma^2 e^{\pm 4(\eta x + \lambda t)} - 1}}.$$

A straightforward computation shows that the converse holds. Finally, we observe that given a solution of the evolution equation, in order to have an immersion, one has to choose the constants l and γ , such that the strip (65) intersects the domain of the solution in \mathbb{R}^2 . \square

4. Proof of Theorem 3

We begin by introducing some notations. Given a differentiable function $u(x, t)$, we denote its partial derivatives by

$$(76) \quad z_i = \frac{\partial^i u}{\partial x^i}, \quad w_i = \frac{\partial^i u}{\partial t^i}, \quad \text{where} \quad z_0 = w_0 = u.$$

We have therefore

$$z_{i,x} = z_{i+1}, \quad z_{i,t} = \frac{\partial^{i-1}u_{xt}}{\partial x^{i-1}}, \quad w_{i,x} = \frac{\partial^{i-1}u_{xt}}{\partial t^{i-1}}, \quad w_{i,t} = w_{i+1},$$

and the total derivatives of a differentiable function $\varphi = \varphi(x, t, z_0, z_1, w_1, \dots, z_\ell, w_\ell)$ are given by

$$(77) \quad D_x\varphi = \varphi_x + \sum_{i=0}^{\ell} \varphi_{z_i} z_{i+1} + \sum_{i=1}^{\ell} \varphi_{w_i} w_{i,x},$$

$$(78) \quad D_t\varphi = \varphi_t + \sum_{i=1}^{\ell} \varphi_{z_i} z_{i,t} + \sum_{i=0}^{\ell} \varphi_{w_i} w_{i+1}.$$

We also introduce the notation

$$(79) \quad \Delta_{ij} = f_{i1}f_{j2} - f_{j1}f_{i2}.$$

Observe that

$$(80) \quad \Delta_{12} \neq 0, \quad \Delta_{13}^2 + \Delta_{23}^2 \neq 0.$$

In fact, $\Delta_{12} \neq 0$ is equivalent to $\omega^1 \wedge \omega^2 \neq 0$. Moreover, $\omega^1 \wedge \omega^3 = \Delta_{13}dx \wedge dt$ and $\omega^2 \wedge \omega^3 = \Delta_{23}dx \wedge dt$. If $\Delta_{13} = \Delta_{23} = 0$, then it follows from (3) that $d\omega^1 = d\omega^2 = 0$. Therefore, $\omega^3(e_1) = \omega^3(e_2) = 0$ and hence $\omega^3 = 0$ that is in contradiction with $d\omega^3 = \omega^1 \wedge \omega^2$.

The classification theorem of Rabelo and Tenenblat (see Theorem 2) for hyperbolic equations describing η pseudo-spherical surfaces makes use of a number of lemmas. Its proof also provides the coefficients f_{ij} of the 1-forms (2) for each equation of Theorem 2. We will need the lemmas and these coefficients for the proof of Theorem 3. We therefore recall them from [12] without proof. However, the reader can easily check, in each case stated in Lemmas 6-8, that the structure equations (3) hold if, and only if, the corresponding hyperbolic equation holds.

Lemma 5. [12] *Let $u_{xt} = F(u, u_x)$ be a differential equation describing η pseudo-spherical surfaces, with associated one-forms $\omega_i = f_{i1}dx + f_{i2}dt$, where f_{ij} and F are real differentiable (C^∞) functions on a open connected*

set $U \subset \mathbb{R}^2$. Then

$$\begin{aligned} f_{11,u} &\equiv f_{31,u} \equiv 0, \\ f_{12,u_x} &\equiv f_{22,u_x} \equiv f_{32,u_x} \equiv 0, \\ f_{11,u_x}^2 + f_{31,u_x}^2 &\neq 0, \quad \text{in } U. \end{aligned}$$

Lemma 6. [12] *The coefficients f_{ij} of the 1-forms (2) for the equation*

$$(81) \quad u_{xt} = F(u), \quad \text{where} \quad F''(u) + \alpha F(u) = 0, \quad \alpha \in \mathbb{R} \setminus \{0\}$$

are given by

$$(82) \quad \begin{pmatrix} f_{11} & f_{12} \\ f_{21} & f_{22} \\ f_{31} & f_{32} \end{pmatrix} = \begin{pmatrix} -\alpha(Bz_1 - AQ) & \alpha A(QF' - \eta F)/(Q^2\alpha + \eta^2) \\ \eta & (\eta F' + \alpha QF)/(Q^2\alpha + \eta^2) \\ -\alpha(Az_1 - BQ) & B\alpha(QF' - \eta F)/(Q^2\alpha + \eta^2) \end{pmatrix},$$

where $z_1 = u_x$, $A, B, Q \in \mathbb{R}$ are such that $\alpha = 1/(A^2 - B^2)$, $A^2 - B^2 \neq 0$ and $Q^2\alpha + \eta^2 \neq 0$ and $\eta \in \mathbb{R} \setminus \{0\}$. In particular, if $B = 0$ and hence $A \neq 0$, one has $\alpha = 1/A^2 > 0$ and

$$(83) \quad \begin{pmatrix} f_{11} & f_{12} \\ f_{21} & f_{22} \\ f_{31} & f_{32} \end{pmatrix} = \begin{pmatrix} \alpha A Q & \alpha A(QF' - \eta F)/(Q^2\alpha + \eta^2) \\ \eta & (\eta F' + Q\alpha F)/(Q^2\alpha + \eta^2) \\ -\alpha A z_1 & 0 \end{pmatrix}.$$

Lemma 7. [12] *The coefficients f_{ij} of the 1-forms (2) for the equation*

$$(84) \quad u_{xt} = \nu e^{\delta u} \sqrt{\beta + \gamma u_x^2}, \quad \text{where} \quad \delta, \gamma, \nu \in \mathbb{R} \setminus \{0\} \quad \text{and} \quad \beta = 0, \quad \text{when } \gamma = 1,$$

are given as follows:

a) If $\gamma \neq 1$, then

$$(85) \quad \begin{pmatrix} f_{11} & f_{12} \\ f_{21} & f_{22} \\ f_{31} & f_{32} \end{pmatrix} = \begin{pmatrix} \eta A \delta - (Bz_1 \mp A\sqrt{\Delta})\delta^2/(\gamma - 1) & \pm A\delta\nu e^{\delta z_0} \\ \eta & \pm \nu e^{\delta z_0} \\ \eta B \delta - (Az_1 \mp B\sqrt{\Delta})\delta^2/(\gamma - 1) & \pm B\delta\nu e^{\delta z_0} \end{pmatrix},$$

where $z_0 = u$, $z_1 = u_x$, $\Delta = \beta + \gamma z_1^2 > 0$, $A^2 - B^2 = (\gamma - 1)/\delta^2$ and $\eta \in \mathbb{R} \setminus \{0\}$.

b) If $\gamma = 1$,

$$(86) \quad \begin{pmatrix} f_{11} & f_{12} \\ f_{21} & f_{22} \\ f_{31} & f_{32} \end{pmatrix} = \begin{pmatrix} \frac{1}{2}(\frac{1}{A} + \delta^2 A)z_1 + \eta\delta A & \pm A\delta\nu e^{\delta z_0} \\ \eta & \pm \nu e^{\delta z_0} \\ \frac{1}{2}(-\frac{1}{A} + \delta^2 A)z_1 \pm \eta\delta A & A\delta\nu e^{\delta z_0} \end{pmatrix},$$

where $A, \eta \in \mathbb{R} \setminus \{0\}$.

Lemma 8. [12] *The coefficients f_{ij} of the 1-forms (2) for the equation*

$$(87) \quad u_{xt} = \lambda u + \xi u_x + \tau, \quad \lambda, \xi, \tau \in \mathbb{R}$$

are given as follows:

a) If $\lambda = \xi = \tau = 0$, then

$$(88) \quad \begin{pmatrix} f_{11} & f_{12} \\ f_{21} & f_{22} \\ f_{31} & f_{32} \end{pmatrix} = \begin{pmatrix} z_1 & 0 \\ \eta & e^{z_0} \\ \eta & e^{z_0} \end{pmatrix},$$

where $z_0 = u$, $z_1 = u_x$ and $\eta \neq 0$.

b) If $\lambda \neq 0$, then

$$(89) \quad \begin{pmatrix} f_{11} & f_{12} \\ f_{21} & f_{22} \\ f_{31} & f_{32} \end{pmatrix} = \begin{pmatrix} \pm\eta T z_1/\lambda & T z_0 + \tau T/\lambda \\ \eta & \lambda/\eta \mp \xi \\ \eta T z_1/\lambda & \pm T z_0 \pm \tau T/\lambda \end{pmatrix}$$

where $T, \eta \in \mathbb{R} \setminus \{0\}$.

c) If $\lambda = 0$ and $\xi^2 + \tau^2 \neq 0$, then

$$(90) \quad \begin{pmatrix} f_{11} & f_{12} \\ f_{21} & f_{22} \\ f_{31} & f_{32} \end{pmatrix} = \begin{pmatrix} \int dz_1/F(z_1) & 1/\eta \\ \eta & 0 \\ \int dz_1/F(z_1) & 1/\eta \end{pmatrix}$$

where $\eta \in \mathbb{R} \setminus \{0\}$.

Having recalled these results from [12], we are now ready to proceed with the proof of Theorem 3. The proof consists of a number of technical lemmas and propositions, in which we analyze the existence of solutions for the system of equations (17), (18) and (19) that depend on u and finitely many derivatives, for each of the classes of hyperbolic equations obtained by Rabelo and Tenenblat in Theorem 2.

With the notation introduced in (79), equations (17) and (18) are written as

$$(91) \quad f_{11}D_t a + \eta D_t b - f_{12}D_x a - f_{22}D_x b - 2b\Delta_{13} + (a - c)\Delta_{23} = 0,$$

$$(92) \quad f_{11}D_t b + \eta D_t c - f_{12}D_x b - f_{22}D_x c + (a - c)\Delta_{13} + 2b\Delta_{23} = 0.$$

Lemma 9. *Consider an equation $u_{xt} = F(u, u_x)$ describing η pseudo-spherical surfaces, with 1-forms ω^i as in (2) where the functions f_{ij} are given by (82)–(90). Assume there is a local isometric immersion of any pseudo-spherical surface, determined by a solution $u(x, t)$, for which the coefficients a, b, c of the forms ω_1^3 and ω_2^3 depend on a jet of finite order of u . Then*

i) $a \neq 0$ on any open set.

ii) $c = 0$ on an open set U if, and only if, $f_{11} = 0$ on U , i.e., F satisfies (81) and f_{ij} are given by (83) with $Q = 0$. In this case, $\alpha = 1/A^2 > 0$,

$$(93) \quad a = \pm \frac{2}{A\alpha} \frac{F'}{F}, \quad b = \pm 1, \quad \text{and} \quad c = 0.$$

Proof. If there is a local isometric immersion of the pseudo-spherical surface, then (91), (92) and (19) must be satisfied by a, b and c .

i) Assume $a = 0$ on an open set, then it follows from (19) that $b \pm 1$. Substituting into (91) and (92) leads to

$$(94) \quad \mp 2\Delta_{13} - c\Delta_{23} = 0,$$

$$(95) \quad \eta D_t c - f_{22}D_x c - c\Delta_{13} \pm 2\Delta_{23} = 0.$$

It follows from (94) and (80) that $\Delta_{23} \neq 0$ and $c = \mp 2\Delta_{13}/\Delta_{23}$. Since Δ_{13} and Δ_{23} depend only on z_0 and z_1 , we conclude that c depends only on z_0 and z_1 and (95) reduces to

$$(96) \quad \eta(c_{z_1}F + c_{z_0}w_1) - f_{22}(c_{z_0}z_1 + c_{z_1}z_2) - c\Delta_{13} \pm 2\Delta_{23} = 0.$$

Taking the derivative of this equation with respect to z_2 and w_1 implies that $f_{22}c_{z_1} = 0$ and $c_{z_0} = 0$. If $f_{22} \neq 0$ then c is constant and (96) reduces to $-c\Delta_{13} \pm 2\Delta_{23} = 0$ i.e., we have

$$\begin{pmatrix} -c & \pm 2 \\ \mp 2 & -c \end{pmatrix} \begin{pmatrix} \Delta_{13} \\ \Delta_{23} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

Since the determinant is nonzero, it implies that $\Delta_{13} = \Delta_{23} = 0$ which contradicts (80). If $f_{22} = 0$ on an open set, then the functions f_{ij} are given

by (90) and hence $\Delta_{13} = 0$ and $\Delta_{23} = 1$. Then (94) implies that $c = 0$ and (95) gives a contradiction. This concludes the proof of i).

ii) Observe that except for the functions f_{ij} given by (83) with $Q = 0$, f_{11} does not vanish on an open set. We will first show that if $f_{11} = 0$ on an open set i.e, F satisfies (81) and f_{ij} are given by (83) and $Q = 0$, then $c = 0$. In fact, for such f_{ij} s we have $\Delta_{13} = -A^2\alpha^2F(u)z_1/\eta$, $\Delta_{23} = A\alpha F'(u)z_1/\eta$, $\alpha = 1/A^2 > 0$ and $A \neq 0$. Hence (91) and (92) reduce to

$$\begin{aligned} \eta D_t b - f_{12} D_x a - f_{22} D_x b - 2b\Delta_{13} + (a - c)\Delta_{23} &= 0, \\ \eta D_t c - f_{12} D_x b - f_{22} D_x c + (a - c)\Delta_{13} + 2b\Delta_{23} &= 0, \end{aligned}$$

where $f_{12} = -\alpha AF/\eta$ and $f_{22} = F'/\eta$. Assume $c \neq 0$, then it follows from (19) that $a = (b^2 - 1)/c$. Assume that a, b and c depend on a jet of order ℓ of u . For $\ell \geq 1$, taking derivatives of both equations with respect to $w_{\ell+1}$ implies that $b_{w_\ell} = c_{w_\ell} = 0$ and hence $a_{w_k} = 0$. Successive differentiation with respect to w_k, \dots, w_1 imply that a, b and c do not depend on w_ℓ, \dots, w_0 . Successive differentiation with respect to $z_{\ell+1}, \dots, z_2$ imply that a, b and c do not depend on z_ℓ, \dots, z_1 . Hence, a, b and c depend only on x and t . Therefore, the above system of equations reduce to

$$\begin{aligned} \eta b_t + \frac{\alpha AF}{\eta} a_x - \frac{F'}{\eta} b_x + 2b \frac{\alpha^2 A^2 F}{\eta} z_1 + (a - c) \frac{\alpha AF'}{\eta} z_1 &= 0, \\ \eta c_t + \frac{\alpha AF}{\eta} b_x - \frac{F'}{\eta} c_x + (a - c) \frac{\alpha^2 A^2 F}{\eta} z_1 + 2b \frac{\alpha AF'}{\eta} z_1 &= 0. \end{aligned}$$

Taking the derivative with respect to z_1 we get

$$(97) \quad \begin{pmatrix} 2b & a - c \\ a - c & 2b \end{pmatrix} \begin{pmatrix} \alpha AF \\ F' \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

Since αAF and F' are not zero we get $a - c = \pm 2b$ and the derivative with respect to z_0 of any equation of (97) reduces to $b(AF' \mp F) = 0$ as a consequence of (81). If $b = 0$ then Gauss equation (19) reduces to $a^2 = -1$. If $F = \pm AF'$ then the derivative with respect to z_0 implies that $\alpha A^2 = -1$. In both cases we get a contradiction. Therefore, $c = 0$.

Conversely, assume $c = 0$ on an open set, then (19) implies $b = \pm 1$ and (91) and (92) reduce to

$$(98) \quad f_{11} D_t a - f_{12} D_x a \mp 2\Delta_{13} + a\Delta_{23} = 0,$$

$$(99) \quad a\Delta_{13} \pm 2\Delta_{23} = 0.$$

It follows from (99) and (80) that $\Delta_{13} \neq 0$ and $a = \mp 2\Delta_{23}/\Delta_{13}$. Since Δ_{13} and Δ_{23} depend only on z_0 and z_1 , we conclude that a depends only on z_0 and z_1 and (98) reduces to

$$(100) \quad f_{11}(a_{z_1}F + a_{z_0}w_1) - f_{12}(a_{z_0}z_1 + a_{z_1}z_2) \mp 2\Delta_{13} + a\Delta_{23} = 0.$$

Differentiation with respect to w_1 and z_2 implies

$$(101) \quad f_{11}a_{z_0} = f_{12}a_{z_1} = 0.$$

Since $\Delta_{12} \neq 0$, we observe that f_{11} and f_{12} cannot vanish simultaneously.

If both $f_{11} \neq 0$ and $f_{12} \neq 0$ then from (101) we conclude that a is constant and (100) reduces to $\mp 2\Delta_{13} + a\Delta_{23} = 0$. This equation with (99) implies that $\Delta_{13} = \Delta_{23} = 0$ which contradicts (80).

If $f_{12} = 0$ on an open set, then f_{ij} are given by (82) with $A = 0, B \neq 0$ or (85) with $A = 0, B \neq 0$ or (88). Since $f_{11} \neq 0$, it follows from (101) that $a_{z_0} = 0$ and (100) reduces to

$$(102) \quad f_{11}a_{z_1}F \mp 2\Delta_{13} + a\Delta_{23} = 0.$$

If f_{ij} are given by (82) with $A = 0, B \neq 0$, then

$$\Delta_{13} = \frac{\alpha(QF' - \eta F)}{Q^2\alpha + \eta^2}z_1, \quad \Delta_{23} = -B\alpha F, \quad a_{z_1}z_1 = -a.$$

Substituting into (102) and differentiating twice with respect to z_1 runs into a contradiction. If f_{ij} are given by (85) with $A = 0, B \neq 0$, then

$$\Delta_{13} = \mp B\nu\delta^2 z_1 e^{\delta z_0}, \quad \Delta_{23} = \pm \nu\delta z_1 e^{\delta z_0}, \quad a = \pm \frac{2}{B\delta}.$$

Therefore, (102) reduces to $\mp 2\Delta_{13} + a\Delta_{23} = 0$ which is in contradiction with (99). Finally if f_{ij} are given by (88), then $\Delta_{23} = 0$, hence it follows from (80) and (99) that $a = 0$ which is a contradiction.

We conclude that if $c = 0$ on an open set, then $f_{11} = 0$ i.e., f_{ij} are given by (83) with $Q = 0$. Therefore $\Delta_{13} = -A^2\alpha^2 F(u)z_1/\eta$, $\Delta_{23} = A\alpha F'(u)z_1/\eta$ and hence (99) implies that $a = \pm 2F'/(A\alpha F)$. Moreover, (100) is an identity since $A^2\alpha = 1$. This concludes the proof of Lemma 9. □

Consider an equation $u_{xt} = F(u, u_x)$ describing η pseudo-spherical surfaces given by Lemmas 6-8. The existence of a local isometric immersion in \mathbb{R}^3 of any pseudo-spherical surface, determined by a solution u , for which

the coefficients a , b and c depend on $x, t, z_0, z_1, w_1, \dots, z_\ell, w_\ell$, is equivalent to requiring that (91), (92) and (19) must be satisfied. Substituting the expressions of the total derivatives with respect to x and t given by (77) and (78), we rewrite (91) and (92) as

$$\begin{aligned}
 (103) \quad & f_{11}a_t + \eta b_t + \sum_{i=0}^{\ell} (f_{11}a_{w_i} + \eta b_{w_i})w_{i+1} \\
 & + \sum_{i=1}^{\ell} (f_{11}a_{z_i} + \eta b_{z_i}) \frac{\partial^{i-1}F}{\partial x^{i-1}} - (f_{12}a_x + f_{22}b_x) \\
 & - \sum_{i=0}^{\ell} (f_{12}a_{z_i} + f_{22}b_{z_i})z_{i+1} - \sum_{i=1}^{\ell} (f_{12}a_{w_i} + f_{22}b_{w_i}) \frac{\partial^{i-1}F}{\partial t^{i-1}} \\
 & - 2b\Delta_{13} + (a - c)\Delta_{23} = 0,
 \end{aligned}$$

and

$$\begin{aligned}
 (104) \quad & f_{11}b_t + \eta c_t + \sum_{i=0}^{\ell} (f_{11}b_{w_i} + \eta c_{w_i})w_{i+1} \\
 & + \sum_{i=1}^{\ell} (f_{11}b_{z_i} + \eta c_{z_i}) \frac{\partial^{i-1}F}{\partial x^{i-1}} - (f_{12}b_x + f_{22}c_x) \\
 & - \sum_{i=0}^{\ell} (f_{12}b_{z_i} + f_{22}c_{z_i})z_{i+1} - \sum_{i=1}^{\ell} (f_{12}b_{w_i} + f_{22}c_{w_i}) \frac{\partial^{i-1}F}{\partial t^{i-1}} \\
 & + (a - c)\Delta_{13} + 2b\Delta_{23} = 0.
 \end{aligned}$$

Differentiating (103) and (104) with respect to $w_{\ell+1}$ leads to

$$(105) \quad f_{11}a_{w_\ell} + \eta b_{w_\ell} = 0 \quad f_{11}b_{w_\ell} + \eta c_{w_\ell} = 0.$$

Differentiation of the Gauss equation (19) with respect to w_ℓ gives $ca_{w_\ell} + ac_{w_\ell} - 2bb_{w_\ell} = 0$. Taking into account (105) in the latter, we obtain

$$(106) \quad \left[c + \left(\frac{f_{11}}{\eta} \right)^2 a + 2 \frac{f_{11}}{\eta} b \right] a_{w_\ell} = 0.$$

The following two lemmas will consider the cases in which the expression between brackets in (106) vanishes or not on an open set.

Lemma 10. *Consider an equation $u_{xt} = F(u, u_x)$ describing η pseudo-spherical surfaces, with 1-forms ω^i as in (2) where the functions f_{ij} are given by (82)–(90). Assume there is a local isometric immersion of a pseudo-spherical surface determined by a solution $u(x, t)$, for which the coefficients a, b, c of the second fundamental form depend on a jet of finite order of u . If*

$$(107) \quad c + \left(\frac{f_{11}}{\eta}\right)^2 a + 2\frac{f_{11}}{\eta}b = 0$$

on a non empty open set, then

i) For equation (81) with f_{ij} as in (83) a, b and c are given by

$$(108) \quad \begin{aligned} a &= \pm \frac{2\eta}{A(Q^2\alpha + \eta^2)} \left(\frac{\eta F'}{\alpha F} + Q \right), \\ b &= \mp \frac{1}{Q^2\alpha + \eta^2} \left(2\eta Q \frac{F'}{F} + Q^2\alpha - \eta^2 \right), \\ c &= \pm \frac{2QA\alpha}{Q^2\alpha + \eta^2} \left(Q \frac{F'}{F} - \eta \right), \end{aligned}$$

where $\alpha = 1/A^2$. In particular when $Q = 0$, a, b, c are given by (93).

ii) For all equations, except those considered in i), equations (91), (92) and (19) form an inconsistent system.

Proof. If (107) holds then substituting c into the Gauss equation (19) leads to $(f_{11}a/\eta + b)^2 = 1$, and hence

$$(109) \quad b = \pm 1 - \frac{f_{11}}{\eta}a \quad \text{and} \quad c = \left(\frac{f_{11}}{\eta}\right)^2 a \mp 2\frac{f_{11}}{\eta}.$$

Therefore,

$$\begin{aligned} f_{11}D_t a + \eta D_t b &= -af_{11,z_1}F, \\ f_{12}D_x a + f_{22}D_x b &= -\frac{\Delta_{12}}{\eta}D_x a - \frac{af_{22}f_{11,z_1}}{\eta}z_2, \\ f_{11}D_t b + \eta D_t c &= \frac{af_{11}f_{11,z_1}}{\eta}F \mp 2f_{11,z_1}F, \\ f_{12}D_x b + f_{22}D_x c &= \frac{f_{11}\Delta_{12}}{\eta^2}D_x a + \frac{\Delta_{12}af_{11,z_1}}{\eta^2}z_2 \\ &\quad + \frac{af_{22}f_{11}f_{11,z_1}}{\eta^2}z_2 \mp 2\frac{f_{22}f_{11,z_1}}{\eta}z_2. \end{aligned}$$

Equation (91) becomes

$$(110) \quad -af_{11,z_1}F + \frac{\Delta_{12}}{\eta}D_xa + \frac{af_{22}f_{11,z_1}}{\eta}z_2 \mp 2\Delta_{13} \\ + 2\frac{f_{11}}{\eta}a\Delta_{13} + \left[1 - \left(\frac{f_{11}}{\eta}\right)^2\right]a\Delta_{23} \pm 2\frac{f_{11}}{\eta}\Delta_{23} = 0$$

and (92) becomes

$$(111) \quad \frac{af_{11}f_{11,z_1}}{\eta}F \mp 2f_{11,z_1}F - \frac{f_{11}\Delta_{12}}{\eta^2}D_xa \\ - \frac{\Delta_{12}af_{11,z_1}}{\eta^2}z_2 - \frac{af_{22}f_{11}f_{11,z_1}}{\eta^2}z_2 \pm 2\frac{f_{22}f_{11,z_1}}{\eta}z_2 \\ + \left[1 - \left(\frac{f_{11}}{\eta}\right)^2\right]a\Delta_{13} \pm 2\frac{f_{11}}{\eta}\Delta_{13} \pm 2\Delta_{23} - 2\frac{f_{11}}{\eta}a\Delta_{23} = 0.$$

If $\ell \geq 2$, then differentiating (110) with respect to $z_{\ell+1}$ leads to $a_{z_\ell} = 0$. Successive differentiation with respect to z_ℓ, \dots, z_3 leads to $a_{z_\ell} = a_{z_{\ell-1}} = \dots = a_{z_2} = 0$. If $\ell \geq 1$, then differentiating (110) and (111) with respect to z_2 leads to

$$\Delta_{12}a_{z_1} + af_{22}f_{11,z_1} = 0, \\ -f_{11}\Delta_{12}a_{z_1} - \Delta_{12}af_{11,z_1} - af_{22}f_{11}f_{11,z_1} \pm 2\eta f_{22}f_{11,z_1} = 0,$$

which is equivalent to

$$(112) \quad \Delta_{12}a_{z_1} + af_{22}f_{11,z_1} = 0,$$

$$(113) \quad (\Delta_{12}a \mp 2\eta f_{22})f_{11,z_1} = 0.$$

i) For equation (81) with f_{ij} given by (83) we have $f_{11,z_1} = 0$. Hence (113) is trivially satisfied and (112) implies that $a_{z_1} = 0$. Moreover, (110) and (111) reduce to

$$(114) \quad \frac{\Delta_{12}}{\eta}D_xa + 2\left(\frac{f_{11}}{\eta}a \mp 1\right)\Delta_{13} + \left[\left(1 - \frac{f_{11}^2}{\eta^2}\right)a \pm 2\frac{f_{11}}{\eta}\right]\Delta_{23},$$

$$(115) \quad -f_{11}\frac{\Delta_{12}}{\eta^2}D_xa + \left[\left(1 - \frac{f_{11}^2}{\eta^2}\right)a \pm 2\frac{f_{11}}{\eta}\right]\Delta_{13} - 2\left(\frac{f_{11}}{\eta}a \mp 1\right)\Delta_{23}.$$

Adding equation (114) multiplied by f_{11}/η with (115) and cancelling a nonzero factor, we get

$$a\Delta_{13} - \left(\frac{f_{11}a}{\eta} \mp 2\right)\Delta_{23} = 0.$$

Since $\Delta_{13} - f_{11}\Delta_{23}/\eta = f_{31}\Delta_{12}/\eta$, we conclude that $a = \mp 2\Delta_{23}\eta/(f_{31}\Delta_{12})$. For the functions f_{ij} as in (83) we have $f_{31} = -\alpha Az_1 \neq 0$ and

$$\Delta_{12} = \alpha AF, \quad \Delta_{13} = \frac{\alpha(QF' - \eta F)}{Q^2\alpha + \eta^2}z_1, \quad \Delta_{23} = \frac{\alpha A(\eta F' + \alpha QF)}{Q^2\alpha + \eta^2}z_1.$$

Therefore, we conclude that a is given by

$$a = \pm \frac{2\eta}{\alpha Q^2 + \eta^2} \left(\eta A \frac{F'}{F} + \frac{Q}{A} \right).$$

A straightforward computation shows that substituting the expressions of a , $D_x a = a_{z_0} z_1$, $f_{11} = \alpha A Q$ and using the fact that $\alpha A^2 = 1$ equation (114) is trivially satisfied. It follows from (109) that b and c are given as in (108). Observe that when $Q = 0$ then (108) reduces to (93).

ii) For all equations except those considered in i) we have $f_{11,z_1} \neq 0$.

If $\ell = 0$, then differentiating (110) and (111) with respect to z_2 leads to $a f_{22} f_{11,z_1} = 0$ and to $\Delta_{12} a f_{11,z_1} + a f_{22} f_{11} f_{11,z_1} \mp 2\eta f_{22} f_{11,z_1} = 0$. From Lemma 9 $a \neq 0$, hence $f_{22} f_{11,z_1} = 0$ and $\Delta_{12} f_{11,z_1} = 0$. This implies that $f_{11,z_1} = 0$ which is a contradiction. Therefore, $\ell \geq 1$.

If $f_{22} = 0$, which is the case for equation (87) with f_{ij} given by (90), then (112) and (113) leads to $a = 0$ which contradicts Lemma 9. Thus, (91), (92), and the Gauss equation form an inconsistent system.

If $f_{22} \neq 0$, (which is the case for all equations except (87) with $\lambda = 0$, $\xi^2 + \tau^2 \neq 0$) then dividing (113) by f_{11,z_1} leads to $\Delta_{12} a \mp 2\eta f_{22} = 0$, and differentiating the latter with respect to z_1 gives $\Delta_{12} a_{z_1} + a \Delta_{12,z_1} = 0$, where from (79) we have $\Delta_{12,z_1} = f_{22} f_{11,z_1}$. Therefore, (112) is a consequence of (113). From (113), we have

$$(116) \quad a = \pm 2\eta \frac{f_{22}}{\Delta_{12}},$$

which means that $a_{w_1} = a_x = a_t = 0$, i.e., a is a function of z_0 and z_1 only. Equations (110) and (111) become

$$\begin{aligned}
 & -af_{11,z_1}F + \frac{\Delta_{12}}{\eta}a_{z_0}z_1 \mp 2\Delta_{13} \\
 & \quad + 2\frac{f_{11}}{\eta}a\Delta_{13} + \left[1 - \left(\frac{f_{11}}{\eta}\right)^2\right]a\Delta_{23} \pm 2\frac{f_{11}}{\eta}\Delta_{23} = 0, \\
 \frac{af_{11}f_{11,z_1}F}{\eta} \mp 2f_{11,z_1}F - f_{11}\frac{\Delta_{12}}{\eta^2}a_{z_0}z_1 + \left[1 - \left(\frac{f_{11}}{\eta}\right)^2\right]a\Delta_{13} \\
 & \quad \pm 2\frac{f_{11}}{\eta}\Delta_{13} \pm 2\Delta_{23} - 2\frac{f_{11}}{\eta}a\Delta_{23} = 0,
 \end{aligned}$$

which are equivalent to

$$\begin{aligned}
 (117) \quad & af_{11,z_1}F - \frac{\Delta_{12}}{\eta}a_{z_0}z_1 - \frac{f_{11}}{\eta}a\Delta_{13} \pm 2\Delta_{13} - a\Delta_{23} \\
 & = \frac{f_{11}}{\eta} \left(af_{31}\frac{\Delta_{12}}{\eta} \pm 2\Delta_{23} \right)
 \end{aligned}$$

and

$$\begin{aligned}
 (118) \quad \pm 2f_{11,z_1}F = & \frac{f_{11}}{\eta} \left[af_{11,z_1}F - \frac{\Delta_{12}}{\eta}a_{z_0}z_1 - \frac{f_{11}}{\eta}a\Delta_{13} \pm 2\Delta_{13} - a\Delta_{23} \right] \\
 & + af_{31}\frac{\Delta_{12}}{\eta} \pm 2\Delta_{23}.
 \end{aligned}$$

Substituting (117) in (118), we obtain

$$F = \pm \frac{1}{f_{11,z_1}} \left(1 + \frac{f_{11}^2}{\eta^2} \right) \left(af_{31}\frac{\Delta_{12}}{2\eta} \mp \Delta_{23} \right),$$

which simplifies to

$$(119) \quad F = \frac{(f_{11}^2 + \eta^2)f_{32}}{\eta f_{11,z_1}}.$$

Observe that we are considering $f_{22} \neq 0$ and $f_{11,z_1} \neq 0$. A straightforward computation shows that (119) leads to a contradiction for equation (81) with f_{ij} as in (82) with $B \neq 0$ and equations (84), (87) with f_{ij} given as in (85)–(90). This concludes the proof of Lemma 10. \square

Lemma 11. *Consider an equation $u_{xt} = F(u, u_x)$ describing η pseudo-spherical surfaces, with 1-forms ω^i as in (2) where the functions f_{ij} are given by (82)-(90). Assume there is a local isometric immersion of a pseudo-spherical surface, determined by a solution $u(x, t)$, for which the coefficients a, b, c of the second fundamental form depend on a jet of finite order of u . If*

$$(120) \quad c + \left(\frac{f_{11}}{\eta}\right)^2 a + 2\frac{f_{11}}{\eta}b \neq 0,$$

holds then a, b and c are functions of x and t , and thus universal.

Proof. If (120) holds then, it follows from Lemma 9 that $c \neq 0$ and $f_{11} \neq 0$. Moreover, from (106) we get $a_{w_\ell} = 0$ and hence (105) implies that $b_{w_\ell} = c_{w_\ell} = 0$.

If $\ell = 0$, then a, b , and c are functions of x and t , and thus universal. If $\ell \geq 1$, then consecutive differentiation of (103), (104) and (19) with respect to w_ℓ, \dots, w_1 lead to $a_{w_i} = b_{w_i} = c_{w_i} = 0$ for $i = 0, \dots, \ell$. In particular, a, b and c do not depend on z_0 . Therefore, a, b , and c are functions of x, t, z_1, \dots, z_ℓ . Differentiating (103) and (104) with respect to $z_{\ell+1}$ leads to

$$(121) \quad f_{12}a_{z_\ell} + f_{22}b_{z_\ell} = 0 \quad \text{and} \quad f_{12}b_{z_\ell} + f_{22}c_{z_\ell} = 0.$$

Differentiation of the Gauss equation (19) with respect to z_ℓ gives

$$(122) \quad ca_{z_\ell} + ac_{z_\ell} - 2bb_{z_\ell} = 0.$$

If $f_{22} = 0$, which is the case for equation (87) with f_{ij} as in (90), since $f_{12} \neq 0$, (121) implies that $a_{z_\ell} = b_{z_\ell} = 0$, and (122) leads to $ac_{z_\ell} = 0$. From Lemma (9) we have $a \neq 0$, hence $c_{z_\ell} = 0$. Successive differentiation of (103), (104) and (19) with respect to z_ℓ, \dots, z_2 leads to $a_{z_i} = b_{z_i} = 0$, and hence $c_{z_i} = 0$ for $i = 1, \dots, \ell$. Therefore, a, b , and c are functions of x and t .

If $f_{22} \neq 0$, then (121) leads to

$$(123) \quad b_{z_\ell} = -\frac{f_{12}}{f_{22}}a_{z_\ell} \quad \text{and} \quad c_{z_\ell} = \frac{f_{12}^2}{f_{22}^2}a_{z_\ell}.$$

Substituting these expressions into (122) we get

$$(124) \quad \left[c + \left(\frac{f_{12}}{f_{22}}\right)^2 a + 2\frac{f_{12}}{f_{22}}b \right] a_{z_\ell} = 0.$$

If

$$(125) \quad c + \left(\frac{f_{12}}{f_{22}}\right)^2 a + 2\frac{f_{12}}{f_{22}}b \neq 0,$$

then $a_{z_\ell} = 0$ and (123) implies that $b_{z_\ell} = c_{z_\ell} = 0$. Consecutive differentiations of (103) and (104) with respect to z_ℓ, \dots, z_2 lead to $a_{z_i} = b_{z_i} = c_{z_i} = 0$ for $i = 1, \dots, \ell$, and hence, a, b and c are functions of x and t only.

If

$$(126) \quad c + \left(\frac{f_{12}}{f_{22}}\right)^2 a + 2\frac{f_{12}}{f_{22}}b = 0$$

on a non empty open set, then Lemma 9 and (120) imply that $c \neq 0$ and hence $f_{12} \neq 0$. It follows from (126) and (19) that

$$(127) \quad b = \pm 1 - \frac{f_{12}}{f_{22}}a \quad \text{and} \quad c = \left(\frac{f_{12}}{f_{22}}\right)^2 a \mp 2\frac{f_{12}}{f_{22}}.$$

Therefore,

$$\begin{aligned} f_{11}D_t a + \eta D_t b &= \frac{\Delta_{12}}{f_{22}}D_t a - \eta a \left(\frac{f_{12}}{f_{22}}\right)_{z_0} w_1, \\ f_{12}D_x a + f_{22}D_x b &= -a f_{22} \left(\frac{f_{12}}{f_{22}}\right)_{z_0} z_1, \\ f_{11}D_t b + n D_t c &= \left(\frac{2\eta f_{12}}{f_{22}}a - f_{11}a \mp 2\eta\right) \left(\frac{f_{12}}{f_{22}}\right)_{z_0} w_1 - \frac{f_{12}}{f_{22}^2} \Delta_{12} D_t a, \\ f_{12}D_x b + f_{22}D_x c &= (a f_{12} \mp 2f_{22}) \left(\frac{f_{12}}{f_{22}}\right)_{z_0} z_1. \end{aligned}$$

Therefore, equation (91) becomes

$$\frac{\Delta_{12}}{f_{22}}D_t a - \eta a \left(\frac{f_{12}}{f_{22}}\right)_{z_0} w_1 + a f_{22} \left(\frac{f_{12}}{f_{22}}\right)_{z_0} z_1 - 2b\Delta_{13} + (a - c)\Delta_{23} = 0$$

and (92) becomes

$$\begin{aligned} -\frac{f_{12}}{f_{22}} \frac{\Delta_{12}}{f_{22}} D_t a + \left\{ \left(\frac{2\eta f_{12}}{f_{22}}a - f_{11}a \mp 2\eta\right) w_1 - (a f_{12} \mp 2f_{22})z_1 \right\} \left(\frac{f_{12}}{f_{22}}\right)_{z_0} \\ + (a - c)\Delta_{13} + 2b\Delta_{23} = 0. \end{aligned}$$

Differentiating the first equation with respect to w_1 leads to $\eta a \left(\frac{f_{12}}{f_{22}}\right)_{z_0} = 0$. Since $\eta a \neq 0$ we have $\left(\frac{f_{12}}{f_{22}}\right)_{z_0} = 0$ and the equations reduce to

$$(128) \quad \frac{\Delta_{12}}{f_{22}} D_t a - 2b\Delta_{13} + (a - c)\Delta_{23} = 0,$$

$$(129) \quad -\frac{f_{12}}{f_{22}} \frac{\Delta_{12}}{f_{22}} D_t a + (a - c)\Delta_{13} + 2b\Delta_{23} = 0,$$

Adding (128) multiplied by f_{12}/f_{22} with (129) we get

$$a\Delta_{13} + \left(\pm 2 - \frac{f_{12}}{f_{22}} a\right) \Delta_{23} = 0,$$

which reduces to

$$(130) \quad \frac{f_{32}}{f_{22}} \Delta_{12} a \pm 2\Delta_{23} = 0.$$

Observe that we have $f_{22} \neq 0$, $f_{12} \neq 0$ and $(f_{12}/f_{22})_{z_0} = 0$. Therefore, the only equation that satisfies these conditions is (84) with f_{ij} as in Lemma 7.

If $\gamma = 1$, it follows from (86) that $f_{32} \neq 0$ and (130) implies that a is constant hence, $D_t a = 0$. Therefore, (128) and (129) reduce to

$$\begin{pmatrix} -2b & a - c \\ a - c & 2b \end{pmatrix} \begin{pmatrix} \Delta_{13} \\ \Delta_{23} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

It follows from (80) that $b = 0$ and $a = c$, which contradicts the Gauss equation.

If $\gamma \neq 1$, the functions f_{ij} are given by (85). If $f_{32} = 0$ then $B = 0$ and (130) implies that $\Delta_{23} = 0$. Then it follows from the expression of Δ_{23} that $A = 0$, which contradicts the fact that $A^2 - B^2 \neq 0$. If $f_{32} \neq 0$ i.e., $B \neq 0$, then (130) implies that

$$(131) \quad a = \mp 2 \frac{\Delta_{23} f_{22}}{\Delta_{12} f_{32}}.$$

Substituting the expressions of b and c as in (127) into (128) we get

$$(132) \quad \frac{\Delta_{12}}{f_{22}} D_t a + 2 \left(\frac{f_{12}}{f_{22}} a \mp 1 \right) \Delta_{13} + \left[a - \left(\frac{f_{12}}{f_{22}} \right)^2 a \pm 2 \frac{f_{12}}{f_{22}} \right] \Delta_{23} = 0.$$

Computing the total derivative of a with respect to t , using the expression of a as in (131), equation (132) leads to

$$F(\Delta_{23,z_1}\Delta_{12} - \Delta_{12,z_1}\Delta_{23}) = -f_{22}(\Delta_{13}^2 + \Delta_{23}^2),$$

which in view of (84) and (85) reduces to $(B^2 - A^2\gamma)z_1^2 - A^2\beta = 0$, which is also a contradiction. Therefore, we conclude that the system (91), (92) and the Gauss equation is an inconsistent system. This concludes the proof of Lemma 11. \square

Lemma 12. *Consider the equation $u_{xt} = F(u, u_x)$ which describes η pseudo-spherical surfaces where F is given by (81) and f_{ij} as in (82). If the coefficients of the second fundamental form of the isometric immersion in \mathbb{R}^3 of the pseudo-spherical surface, determined by a solution u , are universal, then the system of equations (91), (92) and the Gauss equation (19) is inconsistent.*

Proof. If the coefficients of the second fundamental form of the isometric immersion of the η pseudo-spherical surfaces described by the differential equation are universal, then equations (91) and (92) reduce to:

$$\begin{aligned} f_{11}a_t + \eta b_t - f_{12}a_x - f_{22}b_x - 2b\Delta_{13} + (a - c)\Delta_{23} &= 0 \\ f_{11}b_t + \eta c_t - f_{12}b_x - f_{22}c_x + (a - c)\Delta_{13} + 2b\Delta_{23} &= 0, \end{aligned}$$

where f_{ij} are given by (82). Differentiating both equations with respect to z_1 leads to

$$(133) \quad -\alpha B a_t - 2b \frac{\alpha(QF' - \eta F)}{Q^2\alpha + \eta^2} + (a - c) \frac{\alpha A(\eta F' + \alpha QF)}{Q^2\alpha + \eta^2} = 0$$

$$(134) \quad -\alpha B b_t + (a - c) \frac{\alpha(QF' - \eta F)}{Q^2\alpha + \eta^2} + 2b \frac{\alpha A(\eta F' + \alpha QF)}{Q^2\alpha + \eta^2} = 0$$

Multiplying (133) and (134) by $Q^2\alpha + \eta^2/\alpha$, and differentiating with respect to z_0 , and taking into account that $F'' = -\alpha F$, we obtain

$$\begin{pmatrix} 2b & \alpha A(a - c) \\ -(a - c) & 2\alpha Ab \end{pmatrix} \begin{pmatrix} \alpha QF + \eta F' \\ QF' - \eta F \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

Since $\alpha QF + \eta F'$ and $QF' - \eta F$ are not zero, we conclude that $\alpha A[4b^2 + (a - c)^2] = 0$. If $b = 0$ and $a = c$ then Gauss equation leads to a contradiction.

If $A = 0$ then equations (133) and (134) reduce to

$$\begin{aligned} -\alpha B a_t - 2b \frac{\alpha(QF' - \eta F)}{Q^2\alpha + \eta^2} &= 0, \\ -\alpha B b_t + (a - c) \frac{\alpha(QF' - \eta F)}{Q^2\alpha + \eta^2} &= 0. \end{aligned}$$

taking derivative with respect to z_0 of both equations, we conclude that $b = a - c = 0$ which is again a contradiction. Therefore, the system (91), (92) and the Gauss equation is inconsistent. \square

Proposition 3. *Consider an equation*

$$u_{xt} = F(u), \text{ with } F'' + \alpha F = 0, \quad \alpha \neq 0,$$

describing η pseudo-spherical surfaces with f_{ij} given by (82). There exists a local isometric immersion in \mathbb{R}^3 of a pseudo-spherical surface, defined by a solution u , for which the coefficients of the second fundamental form depend on a jet of finite order of u , that is, a, b and c depend on $x, t, u, w_1 \dots, \partial^\ell u / \partial x^\ell, w_\ell$, where ℓ is finite if, and only if, $\alpha > 0$ and f_{ij} are given by (83), a, b, c depend on the jet of order zero of u and are given by (108).

Proof. Assume the local isometric immersion exists. If $c + (f_{11}/\eta)^2 a + 2f_{11}b/\eta = 0$ on a non empty open set, then it follows from Lemma 10 that $B = 0$, i.e. $\alpha > 0$ and f_{ij} are given by (83). Moreover, a, b, c depend on the jet of order zero of u and are given by (108). If $c + (f_{11}/\eta)^2 a + 2f_{11}b/\eta \neq 0$, then Lemma 11 implies that a, b, c are universal. However, it follows from Lemma 12 that such an immersion does not exist.

Conversely, a straightforward computation shows that if f_{ij} are given as in (83) and a, b, c as in (108), then the connection forms ω_1^3 and ω_2^3 given by (13) satisfy the structure equations (14) of an immersion in \mathbb{R}^3 and the Gauss equation (19). \square

Proposition 4. *Consider an equation of type $u_{xt} = \nu e^{\delta u} \sqrt{\beta + \gamma u_x^2}$ describing η pseudo-spherical surfaces, with f_{ij} given by Lemma 7. There is no local isometric immersion in \mathbb{R}^3 of a pseudo-spherical surface determined by a solution u of the equation, for which the coefficients of the second fundamental form depend on a jet of finite order of u .*

Proof. If the immersion exists, then Lemma 10 ii) implies that $c + (f_{11}/\eta)^2 a + 2f_{11}b/\eta \neq 0$, and it follows from Lemma 11 that a, b, c are universal. Therefore, equations (91) and (92) reduce to

$$\begin{aligned} f_{11}a_t + \eta b_t - f_{12}a_x - f_{22}b_x - 2b\Delta_{13} + (a - c)\Delta_{23} &= 0, \\ f_{11}b_t + \eta c_t - f_{12}b_x - f_{22}c_x + (a - c)\Delta_{13} + 2b\Delta_{23} &= 0, \end{aligned}$$

where f_{ij} are given by (85) if $\gamma \neq 1$ and (86) if $\gamma = 1$. Differentiating these equations with respect to z_1 and then with respect to z_0 leads to

$$\begin{pmatrix} -2b & a - c \\ a - c & 2b \end{pmatrix} \begin{pmatrix} \Delta_{13, z_1 z_0} \\ \Delta_{23, z_1 z_0} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

In both cases, i.e., $\gamma = 1$ or $\gamma \neq 1$, since $\Delta_{13, z_1 z_0} \Delta_{23, z_1 z_0} \neq 0$, these equations imply that $b = 0$ and $a = c$ which is inconsistent with the Gauss equation. \square

Proposition 5. *Consider an equation $u_{xt} = \lambda u + \xi u_x + \tau$ describing η -pseudospherical surfaces with f_{ij} given by (88)-(90). There exists a local isometric immersion in \mathbb{R}^3 of a pseudo-spherical surface, defined by a solution u , for which the coefficients of the second fundamental form a, b, c depend of a jet of finite order of u if, and only if, λ, ξ and τ do not vanish simultaneously and a, b, c are universal and given by:*

i) When $\lambda \neq 0$,

$$(135) \quad a = \sqrt{lL(x, t) - \gamma^2 L^2(x, t) - 1}, \quad b = \gamma L(x, t), \quad c = \frac{b^2 - 1}{a},$$

where $L(x, t) = e^{\pm 2[\eta x + (\lambda/\eta \mp \zeta)t]}$ $l, \gamma \in \mathbb{R}$ and $l^2 > 4\gamma^2$ and the 1-forms are defined on a strip of \mathbb{R} where

$$(136) \quad \log \sqrt{\frac{l - \sqrt{l^2 - 4\gamma^2}}{2\gamma^2}} < \pm[\eta x + (\lambda/\eta \mp \zeta)t] < \log \sqrt{\frac{l + \sqrt{l^2 - 4\gamma^2}}{2\gamma^2}}.$$

ii) When $\lambda = 0$ and $\xi^2 + \tau^2 \neq 0$,

$$(137) \quad a = \sqrt{le^{2\eta x} - \gamma^2 e^{4\eta x} - 1}, \quad b = \gamma e^{2\eta x}, \quad c = \frac{b^2 - 1}{a},$$

$l, \gamma \in \mathbb{R}$ and $l^2 > 4\gamma^2$ and the 1-forms are defined on a strip of \mathbb{R}^2 where

$$(138) \quad \log \sqrt{\frac{l - \sqrt{l^2 - 4\gamma^2}}{2\gamma^2}} < \eta x < \log \sqrt{\frac{l + \sqrt{l^2 - 4\gamma^2}}{2\gamma^2}}.$$

Moreover, the constants l and γ have to be chosen so that the strip intersects the domain of the solution of the evolution equation.

Proof. If the coefficients of the second fundamental form of the local isometric immersion of η pseudo-spherical surfaces described by the equation of type iii) depend of a jet of finite order of u , then they are universal by Lemmas 10 and 11, and hence (91) and (92) becomes

$$(139) \quad f_{11}a_t + \eta b_t - f_{12}a_x - f_{22}b_x - 2b\Delta_{13} + (a - c)\Delta_{23} = 0,$$

$$(140) \quad f_{11}b_t + \eta c_t - f_{12}b_x - f_{22}c_x + (a - c)\Delta_{13} + 2b\Delta_{23} = 0.$$

If $\lambda = \xi = \tau = 0$ and f_{ij} are given by (88) then taking the derivative of both equations with respect to z_0 , and using the fact that $\Delta_{13} = e^{z_0} z_1$ and $\Delta_{23} = 0$ we get

$$\begin{aligned} b_x + 2bz_1 &= 0, \\ c_x - (a - c)z_1 &= 0. \end{aligned}$$

Since a, b, c are universal we conclude that $b = 0$ and $a = c$ which contradicts Gauss equation. Therefore the immersion does not exit.

i) If $\lambda \neq 0$ and the functions f_{ij} are as in (89) then $\Delta_{13} = 0$. Differentiating (139) and (140) with respect to z_1 leads to (after dividing by f_{11,z_1})

$$(141) \quad a_t = \pm f_{22}(a - c),$$

$$(142) \quad b_t = \pm 2bf_{22}.$$

Differentiating (139) and (140) with respect to z_0 leads to (after dividing by f_{12,z_0})

$$(143) \quad a_x = \pm \eta(a - c),$$

$$(144) \quad b_x = \pm 2\eta b.$$

and hence, (139) and (140) reduce to

$$(145) \quad \eta b_t - f_{22} b_x = 0,$$

$$(146) \quad \eta c_t - f_{22} c_x = 0.$$

The equations (141), (142), (143), (144), (145), and (146) are the same as (70), (71), (68), (69), (72), and (73) respectively, since f_{22} is constant. Therefore, a is as in (62), b is as in (63), and c is as in (64) and are subject to (65), where λ is replaced by $f_{22} = \lambda/\eta \mp \zeta$. Therefore, we obtain a, b, c given as in (135) defined on the strip (136).

ii) If $\lambda = 0$, $\xi^2 + \tau^2 \neq 0$ and the functions f_{ij} are as in (90), then $\Delta_{13} = 0$ and $\Delta_{23} = 1$, hence (139) and (140) reduce to

$$(147) \quad f_{11} a_t + \eta b_t - f_{12} a_x + (a - c) = 0,$$

$$(148) \quad f_{11} b_t + \eta c_t - f_{12} b_x + 2b = 0.$$

Differentiating with respect to z_1 leads to $a_t = b_t = 0$. Since from Lemma 9 we have $a \neq 0$, Gauss equation implies that $c_t = 0$ and thus (147) and (148) become

$$a_x = \eta(a - c),$$

$$b_x = 2\eta b,$$

where $c = (b^2 - 1)/a$. The arguments used in the proof of Proposition 2, with $\lambda = 0$ and \pm replaced by $+$, imply that a, b, c are given by (137), that are defined on the strip given by (138).

The converse follows from a straightforward computation. \square

Finally, the proof of Theorem 3 follows from Propositions 3, 4 and 5. \square

References

- [1] R. Beals, M. Rabelo, and K. Tenenblat, *Bäcklund transformations and inverse scattering solutions for some pseudospherical surface equations*, Stud. Appl. Math., **81** (1989), no. 2, 125–151.
- [2] J. A. Cavalcante and K. Tenenblat, *Conservation laws for nonlinear evolution equations*, J. Math. Phys., **29** (1988), no. 4, 1044–1049.
- [3] S. S. Chern and K. Tenenblat, *Pseudospherical surfaces and evolution equations*, Stud. Appl. Math., **74** (1986), 55–83.

- [4] Q. Ding and K. Tenenblat, *On differential systems describing surfaces of constant curvature*, J. of Differential Equations, **184** (2002), 185–214.
- [5] V. M. Foursov, P. J. Olver, and E.G. Reyes, *On formal integrability of evolution equations and local geometry of surfaces*, Differential Geom. Appl., **15** (2001), 183–199.
- [6] V. P. Gomes Neto, *Fifth-order evolution equations describing pseudo-spherical surfaces*, J. of Differential Equations, **249** (2010), 2822–2865.
- [7] P. Górká and E. G. Reyes, *The modified Hunter-Saxton equation*, J. Geom. Phys., **62** (2012), 1793–1809.
- [8] N. Kamran and K. Tenenblat, *On differential equations describing pseudo-spherical surfaces*, J. of Differential Equations, **115** (1995), no. 1, 75–98.
- [9] L. Jorge and K. Tenenblat, *Linear problems associated to evolution equations of type $u_{tt} = F(u, u_x, u_{xx}, u_t)$* , Stud. Appl. Math., **77** (1987), 103–117.
- [10] M. Rabelo *A characterization of differential equations of type $u_{xt} = F(u, \partial u/\partial x, \dots, \partial^k/\partial x^k)$ which describe pseudospherical surfaces*, An. Acad. Brasil. Ciênc., **60** (1988), no. 2, 119–126.
- [11] M. Rabelo *On equations which describe pseudospherical surfaces*, Stud. Appl. Math., **81** (1989), 221–248.
- [12] M. Rabelo and K. Tenenblat, *On equations of the type $u_{xt} = F(u, u_x)$ which describe pseudospherical surfaces*, J. Math. Phys., **6** (1990), 1400–1407.
- [13] M. Rabelo and K. Tenenblat, *A classification of pseudospherical surfaces equations of the type $u_t = u_{xxx} + G(u, u_x, u_{xx})$ which describe pseudo-spherical surfaces*, J. Math. Phys., **33** (1992), 537–549.
- [14] E. G. Reyes, *Pseudo-spherical surfaces and integrability of evolution equations*, J. of Differential Equations, **147** (1998), no. 1, 195–230.
- [15] E. G. Reyes, *Conservation laws and Calapso-Guichard deformations of equations describing pseudo-spherical surfaces*, J. Math. Phys., **41** (2000), 2968–2989.
- [16] E. G. Reyes, *Geometric integrability of the Camassa-Holm equation*, Lett. Math. Phys., **59** (2002), 117–131.

- [17] E. G. Reyes, *Pseudo-potentials, nonlocal symmetries and integrability of some shallow water equations*, *Selecta Math. (N.S.)*, **12** (2006), 241–270.
- [18] E. G. Reyes, *Correspondence theorems for hierarchies of equations of pseudo-spherical type*, *J. Differential Equations*, **225** (2006), 26–56.

DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY
EVANSTON, IL 60208-2370, USA

E-mail address: `nabil@math.northwestern.edu`

Current address:

DEPARTMENT OF MATHEMATICS
NORTHEASTERN ILLINOIS UNIVERSITY
CHICAGO, IL 60625-4699, USA

DEPARTMENT OF MATHEMATICS AND STATISTICS, MCGILL UNIVERSITY
MONTREAL, QUEBEC, H3A 0B9, CANADA

E-mail address: `nkamran@math.mcgill.ca`

DEPARTMENT OF MATHEMATICS, UNIVERSIDADE DE BRASÍLIA
BRASÍLIA – DF, 70910-900, BRAZIL

E-mail address: `K.Tenenblat@mat.unb.br`

RECEIVED SEPTEMBER 20, 2013

