

A Riemann-Hilbert Correspondence in Positive Characteristic

BHARGAV BHATT AND JACOB LURIE

We explain a version of the Riemann-Hilbert correspondence for p -torsion étale sheaves on an arbitrary \mathbf{F}_p -scheme.

1	Introduction	74
1.1	Outline	75
1.2	The Work of Böckle-Pink	78
1.3	The Work of Emerton-Kisin	79
1.4	Other Related Works	82
1.5	Acknowledgements	83
2	Overview	83
2.1	Frobenius Modules	83
2.2	Étale Sheaves	86
2.3	The Solution Functor	87
2.4	The Riemann-Hilbert Correspondence	88
3	The Category of Frobenius Modules	89
3.1	Comparison of $R[F]$ -Modules with R -Modules	90
3.2	Perfect Frobenius Modules	92
3.3	Restriction and Extension of Scalars	95
3.4	Perfect Rings	97
3.5	Exactness Properties of f^\diamond	98
4	Holonomic Frobenius Modules	101
4.1	Holonomicity	102

4.2	Comparison with Algebraic Frobenius Modules	103
4.3	The Noetherian Case	107
4.4	The Support of a Holonomic Frobenius Module	109
5	Compactly Supported Direct Images	111
5.1	Definitions	112
5.2	Extension by Zero	114
5.3	Kashiwara's Theorem	117
5.4	Existence of Compactly Supported Direct Images	119
6	The Riemann-Hilbert Functor	123
6.1	Existence of the Riemann-Hilbert Functor	124
6.2	Functoriality	126
6.3	Constructible Sheaves	128
6.4	Exactness of the Riemann-Hilbert Functor	130
6.5	Comparison of Finite Direct Images	131
7	The Riemann-Hilbert Correspondence	133
7.1	Derived Solution Functors	133
7.2	Full Faithfulness of the Riemann-Hilbert Functor	135
7.3	The Case of a Field	137
7.4	Proof of the Main Theorem	139
8	Tensor Products	141
8.1	Tensor Products of Frobenius Modules	142
8.2	Derived Tensor Products	144
8.3	Tensor Products of Holonomic Modules	147
8.4	Compatibility with the Riemann-Hilbert Correspondence	148
9	The p^n -Torsion Case	149
9.1	Frobenius Modules over the Witt Vectors	150

9.2	Functoriality	152
9.3	The Solution Functor	154
9.4	Derived Solution Functors	157
9.5	Algebraic Frobenius Modules over $W_n(R)$	160
9.6	The Riemann-Hilbert Correspondence for $\mathbf{Z}/p^n\mathbf{Z}$ -Sheaves	163
10	Globalization	166
10.1	Frobenius Sheaves on a Scheme	167
10.2	The Riemann-Hilbert Correspondence	170
10.3	Functoriality	172
10.4	Holonomic Frobenius Sheaves	176
10.5	Proper Direct Images	178
10.6	Application: The Proper Base Change Theorem	181
11	The Contravariant Riemann-Hilbert Correspondence	185
11.1	Finitely Generated Unit Frobenius Modules	186
11.2	Existence of Kernels	188
11.3	Finitely Generated Unit Complexes	192
11.4	The Emerton-Kisin Correspondence	197
12	Duality for Frobenius Modules	198
12.1	The Derived Riemann-Hilbert Correspondence	199
12.2	Duality for R -Projective Frobenius Modules	202
12.3	Weak Duality in $D(R[F])$	204
12.4	Presentations of Holonomic Complexes	207
12.5	The Duality Functor	210
12.6	Comparison of Solution Functors	214
	References	216

1. Introduction

Let p be a prime number, which we regard as fixed throughout this paper. Our starting point is the following theorem of Katz (see [10, Proposition 4.1.1]):

Theorem 1.0.1 (Katz). *Let k be a perfect field of characteristic p and let \bar{k} be an algebraic closure of k . Then the construction $V \mapsto (V \otimes_{\mathbf{F}_p} \bar{k})^{\text{Gal}(\bar{k}/k)}$ induces an equivalence from the category of finite-dimensional \mathbf{F}_p -vector spaces V with a continuous action of $\text{Gal}(\bar{k}/k)$ to the category of finite-dimensional k -vector spaces M equipped with a Frobenius-semilinear automorphism φ_M .*

The equivalence of Theorem 1.0.1 can be extended to infinite-dimensional vector spaces; in this case, we must add the requirement that M is *locally finite* in the sense that each element $x \in M$ belongs to a finite-dimensional φ_M -stable subspace. Our primary goal in this paper is to prove the following more general result; see Theorem 7.4.1 for a more precise formulation that also describes constructible sheaves.

Theorem 1.0.2. *Let R be a commutative \mathbf{F}_p -algebra. Then there is a fully faithful embedding of abelian categories*

$$\left\{ \begin{array}{l} p\text{-torsion étale sheaves} \\ \text{on } \text{Spec}(R) \end{array} \right\} \xrightarrow{\text{RH}} \left\{ \begin{array}{l} R\text{-modules } M \text{ equipped with a} \\ \text{Frobenius-semilinear automorphism } \varphi_M \end{array} \right\}.$$

Moreover, the essential image of RH consists of those pairs (M, φ_M) that satisfy the following condition: every element $x \in M$ satisfies an equation of the form

$$\varphi_M^n x + a_1 \varphi_M^{n-1} x + \cdots + a_n x = 0$$

for some coefficients $a_1, \dots, a_n \in R$.

We also establish various extensions of Theorem 1.0.2. For instance, the theorem readily extends from the affine scheme $\text{Spec}(R)$ to arbitrary \mathbf{F}_p -schemes, as we summarize next; see Theorem 10.2.7 for a more precise formulation that also identifies constructible sheaves.

Theorem 1.0.3. *Let X be an \mathbf{F}_p -scheme. Then there is a fully faithful embedding of abelian categories*

$$\left\{ \begin{array}{l} p\text{-torsion étale} \\ \text{sheaves on } X \end{array} \right\} \xrightarrow{\text{RH}} \left\{ \begin{array}{l} \text{quasi-coherent } \mathcal{O}_X\text{-modules } \mathcal{F} \text{ equipped with} \\ \text{a Frobenius-semilinear automorphism } \varphi_{\mathcal{F}} \end{array} \right\}.$$

The essential image consists of those pairs $(\mathcal{F}, \varphi_{\mathcal{F}})$ that Zariski locally on X satisfy the condition appearing in Theorem 1.0.2.

Let Mod_R^{Fr} denote the category whose objects are pairs (M, φ_M) , where M is an R -module and φ_M is a Frobenius semilinear endomorphism of M ; we will refer to such pairs as *Frobenius modules over R* . The functor RH appearing in Theorem 1.0.2 may be viewed as an exact and fully faithful embedding of the abelian category of p -torsion étale sheaves on $\text{Spec}(R)$ into the abelian category Mod_R^{Fr} . This functor also induces a bijection on Ext-groups, and thus passes to the derived category to give the following result; we refer to Theorem 12.1.5 for a more precise formulation that also identifies the constructible derived category.

Theorem 1.0.4. *Let R be an \mathbf{F}_p -algebra, and let $D_{\text{ét}}^b(\text{Spec}(R), \mathbf{F}_p)$ be the bounded derived category of p -torsion étale sheaves on $\text{Spec}(R)$. Then the functor in Theorem 1.0.2 lifts to a t -exact and fully faithful embedding of triangulated categories*

$$\text{RH} : D_{\text{ét}}^b(\text{Spec}(R), \mathbf{F}_p) \rightarrow D^b(\text{Mod}_R^{\text{Fr}})$$

whose essential image comprises those complexes $K \in D^b(\text{Mod}_R^{\text{Fr}})$ with the property that each Frobenius module $H^i(K)$ satisfies the conditions in Theorem 1.0.2.

Theorem 1.0.2 also admits an analog describing p^n -torsion étale sheaves on $\text{Spec}(R)$ in terms of suitably defined Frobenius modules over the ring $W_n(R)$ of length n Witt vectors of R ; see Theorem 9.6.1.

1.1. Outline

The first half of this paper is devoted to the proof of Theorem 1.0.2. Note that Theorem 1.0.2 supplies a description of the category of (p -torsion) étale sheaves on $\text{Spec}(R)$ as quasi-coherent sheaves on $\text{Spec}(R)$ with additional structure, and can therefore be viewed as a positive-characteristic analogue of the Riemann-Hilbert correspondence. We will emphasize this perspective by referring to the functor RH appearing in Theorem 1.0.2 as the *Riemann-Hilbert functor*. It is not so easy to describe this functor directly. Instead, we begin in §2 by constructing a functor in the opposite direction. Let Mod_R^{Fr} denote the category of *Frobenius modules over R* introduced above. If M is an R -module and \widetilde{M} denotes the associated quasi-coherent sheaf on the étale site of $\text{Spec}(R)$, then every Frobenius-semilinear automorphism φ_M of M determines an automorphism of \widetilde{M} , which we will also denote by φ_M . We let $\text{Sol}(M)$ denote subsheaf of φ_M -fixed points in \widetilde{M} : that is, the kernel of the map $\text{id} - \varphi_M : \widetilde{M} \rightarrow \widetilde{M}$, formed in the category $\text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$

of p -torsion étale sheaves on $\mathrm{Spec}(R)$. The construction $(M, \varphi_M) \mapsto \mathrm{Sol}(M)$ determines a functor

$$\mathrm{Sol} : \mathrm{Mod}_R^{\mathrm{Fr}} \rightarrow \mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(R), \mathbf{F}_p),$$

which we will refer to as the *solution functor* (Construction 2.3.1).

We will say that a Frobenius module (M, φ_M) is *perfect* if the map φ_M is invertible. In §6, we will show that, when restricted to category $\mathrm{Mod}_R^{\mathrm{perf}}$ of perfect Frobenius modules, the solution functor Sol has a left adjoint (Theorem 6.1.1). This left adjoint is the Riemann-Hilbert functor $\mathrm{RH} : \mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(R), \mathbf{F}_p) \rightarrow \mathrm{Mod}_R^{\mathrm{perf}}$ appearing in the statement Theorem 1.0.2. The existence of the functor RH is not evident: to construct it, we will need to develop a theory of *compactly supported direct images* in the setting of (perfect) Frobenius modules; this is the subject of §5. We also prove in §6 that the functor RH is exact (Proposition 6.4.1). This is easy to see in the case where R is an algebraically closed field: in this case, the category $\mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(R), \mathbf{F}_p)$ is equivalent to the category of vector spaces over \mathbf{F}_p , where every exact sequence is split. We handle the general case by reducing to the case of an algebraically closed field, using a theory of base change for perfect Frobenius modules (which we study in §3) and its compatibility with the Riemann-Hilbert correspondence (which we prove as Proposition 6.2.2).

We will complete the proof of Theorem 1.0.2 in §7 by showing that the Riemann-Hilbert functor RH is fully faithful and characterizing its essential image. The full-faithfulness is actually fairly easy, once we know that the functor RH is exact: it essentially follows from the exactness of the Artin-Schreier sequence $0 \rightarrow \underline{\mathbf{F}}_p \rightarrow \tilde{R} \rightarrow \tilde{R} \rightarrow 0$ in the category of étale sheaves $\mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(R), \mathbf{F}_p)$ (see Proposition 7.2.1). To understand the essential image of the Riemann-Hilbert functor, it will be convenient to consider first the functor RH^c obtained by restricting RH to the subcategory $\mathrm{Shv}_{\mathrm{ét}}^c(\mathrm{Spec}(R), \mathbf{F}_p) \subseteq \mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(R), \mathbf{F}_p)$ of *constructible* étale sheaves on $\mathrm{Spec}(R)$. The functor RH^c takes values in the subcategory $\mathrm{Mod}_R^{\mathrm{hol}} \subseteq \mathrm{Mod}_R^{\mathrm{perf}}$ of *holonomic* Frobenius modules (Definition 4.1.1), which we study in §4. Theorem 1.0.2 will then follow by combining the following two assertions:

- The functor $\mathrm{RH}^c : \mathrm{Shv}_{\mathrm{ét}}^c(\mathrm{Spec}(R), \mathbf{F}_p) \rightarrow \mathrm{Mod}_R^{\mathrm{hol}}$ is an equivalence of categories (Theorem 7.4.1): that is, every holonomic Frobenius module M has the form $\mathrm{RH}(\mathcal{F})$, for some constructible étale sheaf \mathcal{F} on $\mathrm{Spec}(R)$. We will prove this using formal arguments to reduce to the case where R is a field, in which case the desired result follows from Theorem 1.0.1 (which we reprove here as Proposition 7.3.1).

- A perfect Frobenius module M can be written as a filtered colimit of holonomic Frobenius modules if and only if every element $x \in M$ satisfies an equation $\varphi_M^n x + a_1 \varphi_M^{n-1} x + \cdots + a_n x = 0$ for some coefficients $a_1, \dots, a_n \in R$ (Theorem 4.2.9). We will refer to such Frobenius modules as *algebraic*.

In the second half of this paper, we consider several refinements of Theorem 1.0.2:

- Let (M, φ_M) and (N, φ_N) be Frobenius modules over R . If M and N are perfect, then they can also be regarded as Frobenius modules over the perfection R^{1/p^∞} (Proposition 3.4.3). In this case, we can regard the tensor product $M \otimes_{R^{1/p^\infty}} N$ as a (perfect) Frobenius module over R , with Frobenius endomorphism given by $x \otimes y \mapsto \varphi_M(x) \otimes \varphi_N(y)$. In §8, we show that the Riemann-Hilbert functor RH of Theorem 1.0.2 is compatible with tensor products, in the sense that there are canonical isomorphisms

$$\text{RH}(\mathcal{F} \otimes_{\mathbf{F}_p} \mathcal{G}) \simeq \text{RH}(\mathcal{F}) \otimes_{R^{1/p^\infty}} \text{RH}(\mathcal{G})$$

(Corollary 8.4.2). Our proof relies on the vanishing of the Tor-groups $\text{Tor}_n^{R^{1/p^\infty}}(M, N)$ for $n > 0$ when M and N are algebraic Frobenius modules, which we establish as Theorem 8.3.1.

- In §9, we prove a generalization of Theorem 1.0.2 where the category $\text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ of p -torsion étale sheaves is replaced by the larger category $\text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{Z}/p^n \mathbf{Z})$ of p^n -torsion étale sheaves, for some integer $n \geq 0$. In this case, we must also replace the category Mod_R^{Fr} of Frobenius modules over R by the larger category $\text{Mod}_{W_n(R)}^{\text{Fr}}$ of *Frobenius modules over $W_n(R)$* ; here $W_n(R)$ denotes the ring of n -truncated Witt vectors of R (see Theorem 9.6.1).
- In §10, we prove a generalization of Theorem 1.0.2 where the affine scheme $\text{Spec}(R)$ is replaced by an arbitrary \mathbf{F}_p -scheme X (Theorem 10.2.7). We also show that the Riemann-Hilbert correspondence is compatible with the formation of (higher) direct images along proper morphisms $f : X \rightarrow Y$ of finite presentation (Theorem 10.5.5). As an application, we reprove a special case of the proper base change theorem in étale cohomology (namely, the case of p -torsion sheaves on \mathbf{F}_p -schemes; see Corollary 10.6.2).
- In §12, we study the derived category $D(R[F])$ of Frobenius modules over R . The equivalence of abelian categories $\text{RH}^c : \text{Shv}_{\text{ét}}^c(\text{Spec}(R),$

$\mathbf{F}_p) \simeq \text{Mod}_R^{\text{hol}}$ extends to an equivalence of triangulated categories

$$D_c^b(\text{Spec}(R), \mathbf{F}_p) \simeq D_{\text{hol}}^b(R[F]),$$

where $D_{\text{hol}}^b(R[F]) \subseteq D(R[F])$ denotes the full subcategory spanned by the cohomologically bounded chain complexes with holonomic cohomology and $D_c^b(\text{Spec}(R), \mathbf{F}_p)$ is the constructible derived category of $\text{Spec}(R)$; see Corollary 12.1.7. We also construct a duality functor $\mathbb{D} : D_{\text{hol}}^b(R[F]) \rightarrow D(R[F])^{\text{op}}$, and show that it is a fully faithful embedding (Theorem 12.5.4). Combining this duality functor with our Riemann-Hilbert correspondence, we obtain a second embedding from the constructible derived category $D_c^b(\text{Spec}(R), \mathbf{F}_p)$ to the derived category $D(R[F])$. Using this construction, we recover the contravariant Riemann-Hilbert correspondence of [6] (in a strong form, which does not require the F_p -algebra R to be regular or even Noetherian; see Theorem 11.4.4), whose statement we review in §11 (see also §1.3).

These sections are more or less independent of one another, and can be read in any order (except that §12 depends on §11). One can also develop a theory which incorporates several of these refinements simultaneously (for example, one can compare derived categories of $\mathbf{Z}/p^n\mathbf{Z}$ -sheaves on an arbitrary \mathbf{F}_p -scheme X with derived categories of quasi-coherent Frobenius modules over the Witt sheaf $W_n(\mathcal{O}_X)$); we leave such extensions to the reader.

1.2. The Work of Böckle-Pink

In the case where the R is Noetherian, Theorem 1.0.2 is essentially due to Böckle and Pink. Let us briefly summarize some of their work. Assume that R is Noetherian, and let $\text{Mod}_{R,\text{fg}}^{\text{Fr}}$ denote the full category of Frobenius modules (M, φ_M) which are *finitely generated* as modules over R . In [4], Böckle and Pink construct an equivalence of abelian categories

$$\text{Shv}_{\text{ét}}^c(\text{Spec}(R), \mathbf{F}_p) \simeq \text{Mod}_{R,\text{fg}}^{\text{Fr}}/\text{Nil},$$

where Nil is the full subcategory of $\text{Mod}_{R,\text{fg}}^{\text{Fr}}$ spanned by those Frobenius modules (M, φ_M) where φ_M is nilpotent, and $\text{Mod}_{R,\text{fg}}^{\text{Fr}}/\text{Nil}$ denotes the Serre quotient (Böckle and Pink denote this Serre quotient by $\text{Crys}(R)$ and refer to it as the category of *crystals* on $\text{Spec}(R)$). From the perspective of [4], the main innovation of this paper is to realize the category $\text{Crys}(R)$ concretely as a full subcategory Mod_R^{Fr} , rather than abstractly as a Serre quotient. To achieve this, we note that every Frobenius module (M, φ_M) admits a

perfection M^{1/p^∞} , given as an abelian group by the direct limit

$$\varinjlim (M \xrightarrow{\varphi_M} M \xrightarrow{\varphi_M} M \xrightarrow{\varphi_M} M \rightarrow \cdots);$$

we will study this construction in detail in §3.2. This construction annihilates every Frobenius module (M, φ_M) for which φ_M is nilpotent, and therefore determines a functor

$$\mathrm{Crys}(R) = \mathrm{Mod}_{R, \mathrm{fg}}^{\mathrm{Fr}} / \mathrm{Nil} \rightarrow \mathrm{Mod}_R^{\mathrm{Fr}}.$$

This functor is fully faithful, and its essential image is the subcategory $\mathrm{Mod}_R^{\mathrm{hol}} \subseteq \mathrm{Mod}_R^{\mathrm{Fr}}$ of *holonomic* Frobenius modules that we study in §4. The resulting identification of $\mathrm{Crys}(R)$ with $\mathrm{Mod}_R^{\mathrm{hol}}$ carries the equivalence $\mathrm{Shv}_{\mathrm{ét}}^c(\mathrm{Spec}(R), \mathbf{F}_p) \simeq \mathrm{Crys}(R)$ to the Riemann-Hilbert equivalence $\mathrm{RH}^c : \mathrm{Shv}_{\mathrm{ét}}^c(\mathrm{Spec}(R), \mathbf{F}_p) \simeq \mathrm{Mod}_R^{\mathrm{hol}}$ of Theorem 7.4.1.

One advantage of our approach is that it does not require the ring R to be Noetherian. Beware that if R is not Noetherian, then the subcategory $\mathrm{Mod}_{R, \mathrm{fg}}^{\mathrm{Fr}} \subseteq \mathrm{Mod}_R^{\mathrm{Fr}}$ is not abelian, and the formalism of Serre quotients is not available. Nevertheless, we will see that the category $\mathrm{Mod}_R^{\mathrm{hol}}$ of holonomic Frobenius modules is still a well-behaved abelian subcategory of $\mathrm{Mod}_R^{\mathrm{Fr}}$ (Corollary 4.3.3). Note that the extra generality afforded by allowing non-Noetherian rings can quite useful in practice: one of the main themes of the present paper is that the theory is often clarified by replacing an \mathbf{F}_p -algebra R by its perfection R^{1/p^∞} , which is almost never Noetherian.

Our realization of crystals as holonomic Frobenius modules also has the advantage of essentially trivializing the passage to derived categories in §12.1 (see Theorem 12.1.5). The corresponding statement in [4] requires more categorical preliminaries (largely to deal with the derived category of the Serre quotient category $\mathrm{Crys}(R)$ in a useful fashion), and does not describe the constructible derived category $D_c^b(\mathrm{Spec}(R), \mathbf{F}_p)$ as explicitly as Corollary 12.1.7.

Remark 1.2.1. When R is not Noetherian, we cannot realize the category $\mathrm{Mod}_R^{\mathrm{hol}}$ as a Serre quotient of the category $\mathrm{Mod}_{R, \mathrm{fg}}^{\mathrm{Fr}}$. Nevertheless, it is still possible to realize the holonomic derived category $D_{\mathrm{hol}}^b(R[F])$ as a *Verdier* quotient of the triangulated category of complexes of Frobenius modules which are finitely generated and projective over R ; see Remark 12.4.5.

1.3. The Work of Emerton-Kisin

The Riemann-Hilbert correspondence of Theorem 1.0.2 is also closely related to the work of Emerton and Kisin (see [6]). In the case where R is a smooth

algebra over a field k of characteristic p , Emerton and Kisin construct an equivalence of triangulated categories

$$\mathrm{RSol}_{\mathrm{EK}} : D_{\mathrm{fgu}}^b(R[F])^{\mathrm{op}} \simeq D_c^b(\mathrm{Spec}(R), \mathbf{F}_p),$$

where $D_{\mathrm{fgu}}^b(R[F])$ denotes the full subcategory of $D(R[F])$ spanned by the cohomologically bounded chain complexes whose cohomology groups *finitely generated unit* Frobenius modules (see Definition 11.1.3). This differs from our Riemann-Hilbert equivalence $\mathrm{Sol} : \mathrm{Mod}_R^{\mathrm{hol}} \simeq \mathrm{Shv}_{\mathrm{\acute{e}t}}^c(\mathrm{Spec}(R), \mathbf{F}_p)$ in two important respects:

- The functor $\mathrm{Sol} : \mathrm{Mod}_R^{\mathrm{hol}} \simeq \mathrm{Shv}_{\mathrm{\acute{e}t}}^c(\mathrm{Spec}(R), \mathbf{F}_p)$ is an equivalence of abelian categories (though it can be extended to an equivalence of suitable derived categories, see Corollary 12.1.7). However, the functor $\mathrm{RSol}_{\mathrm{EK}}$ is well-defined only at the level of derived categories: in other words, it is not t-exact. Gabber has identified the image of the abelian category of finitely generated unit Frobenius modules under the equivalence $\mathrm{RSol}_{\mathrm{EK}}$ with a certain category of *perverse* \mathbf{F}_p -sheaves inside $D_c^b(\mathrm{Spec}(R), \mathbf{F}_p)$ (see [8]).
- The equivalence $\mathrm{Sol} : \mathrm{Mod}_R^{\mathrm{hol}} \simeq \mathrm{Shv}_{\mathrm{\acute{e}t}}^c(\mathrm{Spec}(R), \mathbf{F}_p)$ is a covariant functor, while $\mathrm{RSol}_{\mathrm{EK}}$ is contravariant.

Example 1.3.1. To illustrate the contrast between the Riemann-Hilbert correspondence of Theorem 1.0.2 and the Riemann-Hilbert correspondence of [6] in more concrete terms, let us consider an arbitrary \mathbf{F}_p -algebra R . A choice of non-zero divisor $t \in R$ determines closed and open immersions

$$i : \mathrm{Spec}(R/tR) \hookrightarrow \mathrm{Spec}(R) \quad j : \mathrm{Spec}(R[t^{-1}]) \hookrightarrow \mathrm{Spec}(R),$$

so that the constant sheaf $\underline{\mathbf{F}}_p$ on $\mathrm{Spec}(R)$ fits into an exact sequence

$$(1) \quad 0 \rightarrow j_! \underline{\mathbf{F}}_p \rightarrow \underline{\mathbf{F}}_p \rightarrow i_* \underline{\mathbf{F}}_p \rightarrow 0.$$

Then the functor RH of Theorem 1.0.2 carries the étale sheaves $\underline{\mathbf{F}}_p$, $j_! \underline{\mathbf{F}}_p$, and $i_* \underline{\mathbf{F}}_p$ to the Frobenius modules R^{1/p^∞} , $(tR)^{1/p^\infty}$, and $(R/tR)^{1/p^\infty}$, respectively, and the exact sequence (1) to the short exact sequence

$$0 \rightarrow (tR)^{1/p^\infty} \rightarrow R^{1/p^\infty} \rightarrow (R/tR)^{1/p^\infty} \rightarrow 0.$$

of Frobenius modules.

On the other hand, if R is a smooth algebra over a field k of characteristic p , then the contravariant functor $\mathrm{RSol}_{\mathrm{EK}}$ from [6] carries the étale sheaves $\underline{\mathbf{F}}_p$, $j_! \underline{\mathbf{F}}_p$, and $i_* \underline{\mathbf{F}}_p$ to the chain complexes of (finitely generated

unit) Frobenius modules R , $R[t^{-1}]$, and $(R[t^{-1}]/R)[-1]$, respectively (up to a cohomological shift of $\dim(R)$: see Remark 11.4.5). In other words, RSol_{EK} carries exact sequence (1) into the distinguished triangle $(R[t^{-1}]/R)[-1] \rightarrow R \rightarrow R[t^{-1}]$ in the derived category $D_{\text{fgu}}^b(\text{Mod}_R^{\text{Fr}})$, obtained by “rotating” the exact sequence

$$0 \rightarrow R \rightarrow R[t^{-1}] \rightarrow R[t^{-1}]/R \rightarrow 0$$

in the category of Frobenius modules.

In §12, we will show that the functor RSol_{EK} fits into a commutative diagram of triangulated categories

$$\begin{array}{ccc} & D_{\text{hol}}^b(R[F]) & \\ \mathbb{D} \swarrow & & \searrow \text{RSol} \\ D_{\text{fgu}}^b(R[F])^{\text{op}} & \xrightarrow{\text{RSol}_{\text{EK}}} & D_c^b(\text{Spec}(R); \mathbf{F}_p), \end{array}$$

where RSol is a derived version of our solution functor Sol , and \mathbb{D} denotes a certain duality functor on the derived category of Frobenius modules. Using the fact that RSol is an equivalence of categories (which follows easily from Theorem 1.0.2 and its proof) and that \mathbb{D} is an equivalence of categories (which we prove as Theorem 12.5.4), we give a new proof of the assertion that RSol_{EK} is an equivalence of categories. Moreover, our argument does not require the assumption that R is a smooth algebra over a field: we allow R to be an arbitrary \mathbf{F}_p -algebra, with the caveat that the triangulated category $D_{\text{fgu}}^b(R[F])$ must be suitably defined (if R is not a regular Noetherian ring, then the criteria for membership in the subcategory $D_{\text{fgu}}^b(R[F]) \subseteq D(R[F])$ must be imposed at the derived level, rather than at the level of individual cohomology groups; see Definition 11.3.4).

Remark 1.3.2. Let R be a commutative \mathbf{F}_p -algebra, let $X = \text{Spec}(R)$ denote the associated affine scheme, and let $\varphi_X : X \rightarrow X$ denote the absolute Frobenius map. A Frobenius module over R can be identified with a quasi-coherent sheaf \mathcal{E} on X equipped with a map $\varphi_{\mathcal{E}} : \mathcal{E} \rightarrow \varphi_{X*} \mathcal{E}$, or equivalently with a map $\psi_{\mathcal{E}} : \varphi_X^* \mathcal{E} \rightarrow \mathcal{E}$. In this paper, we will be primarily concerned with the class of *perfect* Frobenius modules, characterized by the requirement that the map $\varphi_{\mathcal{E}}$ is an isomorphism. By contrast, the book [6] is mainly concerned with the class of *unit* Frobenius modules, characterized by the requirement that the map $\psi_{\mathcal{E}}$ is an isomorphism. Note that the direct image functor φ_{X*} is always exact (since φ_X is affine morphism), but the exactness

of the pullback functor φ_X^* requires some strong hypotheses on R (for example, that R is a regular Noetherian ring). Consequently, the category $\text{Mod}_R^{\text{perf}}$ of perfect Frobenius modules is always abelian, but the category of finitely generated unit Frobenius modules is well-behaved only in special cases.

1.4. Other Related Works

Extensions of the contravariant Riemann-Hilbert correspondence of [6] to singular schemes have also been explored in the papers of Blickle-Böckle [2, 3], Schedlmeier [14], and Ohkawa [13]. Under mild finiteness conditions on R , the papers [2, 3] develop a theory of “Cartier” modules: these are coherent sheaves \mathcal{E} on $X = \text{Spec}(R)$ equipped with a map $C_{\mathcal{E}} : \varphi_{X*} \mathcal{E} \rightarrow \mathcal{E}$. Passing to a quotient by a naturally defined subcategory of nilpotent objects yields the category $\text{Crys}_{\text{Cart}}(R)$ of *Cartier crystals* on R . For X smooth over a perfect field, the category $\text{Crys}_{\text{Cart}}(R)$ is identified in [2] with the category of finitely generated unit Frobenius modules (and thus with the category of perverse \mathbf{F}_p -sheaves on $\text{Spec}(R)_{\text{ét}}$ via [6] and [8]). Roughly speaking, the smoothness of R ensures that $\varphi^! \mathcal{E} \simeq \varphi^* \mathcal{E} \otimes \omega_X^{1-p}$, so a Cartier module $C_{\mathcal{E}} : \varphi_{X*} \mathcal{E} \rightarrow \mathcal{E}$ gives by adjunction a map $\alpha : \mathcal{E} \otimes \omega_X^{-1} \rightarrow \varphi_X^*(\mathcal{E} \otimes \omega_X^{-1})$ whose unitalization (see Construction 11.2.2) yields the desired finitely generated unit Frobenius module. For R not necessarily regular, even though finitely generated unit Frobenius modules may be badly behaved, the abelian category $\text{Crys}_{\text{Cart}}(R)$ is shown to have good behaviour in [2, 3]; see also [8]. Using this, the paper [14] shows that, given an embedding $X \hookrightarrow \text{Spec}(P)$ with P smooth over a perfect field, a suitably defined derived category of $\text{Crys}_{\text{Cart}}(R)$ is equivalent to the full subcategory of $D_{\text{fgu}}^b(P[F])$ spanned by complexes supported on X . Combining this with the Riemann-Hilbert correspondence from [6] for P and a suitable analogue of Kashiwara’s theorem, one obtains a description of the constructible derived category $D_c^b(\text{Spec}(R), \mathbf{F}_p)$ in terms of Cartier crystals on R ([13, 14]).

Remark 1.4.1. The most important example of a Cartier module is the coherent dualizing sheaf ω_X for $X = \text{Spec}(R)$, where we take $C_{\omega_X} : \varphi_{X*} \omega_X \rightarrow \omega_X$ to be the Grothendieck dual of the unit map $\mathcal{O}_X \rightarrow \varphi_{X*} \mathcal{O}_X$. In the case where X is a smooth scheme over a perfect field k , this map can also be described in terms of the Cartier operator on the de Rham complex of X , which motivates the terminology.

In summary, the papers [2, 3, 13, 14] extend the Riemann-Hilbert correspondence of [6] to algebras of finite type over a field by developing the theory of Cartier modules and reducing to the smooth case. In contrast, the discussion in §11 gives an intrinsic extension of the Riemann-Hilbert corre-

spondence of [6] to all \mathbf{F}_p -algebras R , which is described using Frobenius modules (that is, quasi-coherent sheaves with a map $\mathcal{E} \rightarrow \varphi_{X*} \mathcal{E}$). The presentation via Cartier crystals gives a module-theoretic description of perverse \mathbf{F}_p -sheaves on $\mathrm{Spec}(R)_{\acute{e}t}$ (which has an important precursor in [8]), while the presentation in §11 works entirely in the derived category and thus avoids discussion of the abelian category of perverse sheaves.

1.5. Acknowledgements

During the period in which this work was carried out, the first author was supported by National Science Foundation under grant number 1501461 and a Packard fellowship, and the second author was supported by the National Science Foundation under grant number 1510417. We thank Manuel Blickle and Johan de Jong for useful conversations. We also thank Mohsen Asgharzadeh, Micah Darrell, Adeel Khan, and the anonymous referee for helpful remarks on earlier versions of this manuscript.

2. Overview

Our goal in this section is to give a precise formulation of Theorem 1.0.2. We begin by introducing two of the principal objects of interest in this paper: the category $\mathrm{Mod}_R^{\mathrm{Fr}}$ of *Frobenius modules* over a commutative \mathbf{F}_p -algebra R (Definition 2.1.1) and the category $\mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(R), \mathbf{F}_p)$ of *p -torsion étale sheaves* on $\mathrm{Spec}(R)$ (Definition 2.2.2). The Riemann-Hilbert correspondence of Theorem 1.0.2 is a fully faithful embedding of categories

$$\mathrm{RH} : \mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(R), \mathbf{F}_p) \rightarrow \mathrm{Mod}_R^{\mathrm{Fr}}.$$

It will take a bit of work to construct this functor (this is the main objective of §6). In this section, we consider instead the *solution sheaf* functor $\mathrm{Sol} : \mathrm{Mod}_R^{\mathrm{Fr}} \rightarrow \mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(R), \mathbf{F}_p)$, which is left inverse to the Riemann-Hilbert functor RH and admits a very simple description (Definition 2.3.3). We then formulate a variant of Theorem 1.0.2, which asserts that the functor Sol becomes an equivalence when restricted to a certain full subcategory $\mathrm{Mod}_R^{\mathrm{alg}} \subseteq \mathrm{Mod}_R^{\mathrm{Fr}}$ (Theorem 2.4.3).

2.1. Frobenius Modules

We begin by introducing some definitions.

Definition 2.1.1. Let R be a commutative \mathbf{F}_p -algebra. A *Frobenius module over R* is an R -module M equipped with an additive map $\varphi_M : M \rightarrow M$ satisfying the identity $\varphi_M(\lambda x) = \lambda^p \varphi_M(x)$ for $x \in M$, $\lambda \in R$.

Let (M, φ_M) and (N, φ_N) be Frobenius modules over R . A *morphism of Frobenius modules* from (M, φ_M) to (N, φ_N) is an R -module homomorphism $\rho : M \rightarrow N$ for which the diagram

$$\begin{array}{ccc} M & \xrightarrow{\rho} & N \\ \downarrow \varphi_M & & \downarrow \varphi_N \\ M & \xrightarrow{\rho} & N \end{array}$$

commutes. We let Mod_R^{Fr} denote the category whose objects are Frobenius modules (M, φ_M) and whose morphisms are morphisms of Frobenius modules. We will refer to Mod_R^{Fr} as the *category of Frobenius modules over R* .

Notation 2.1.2. Let M and N be Frobenius modules over a commutative \mathbf{F}_p -algebra R . We let $\text{Hom}_R^{\text{Fr}}(M, N)$ denote the set of Frobenius module morphisms from (M, φ_M) to (N, φ_N) .

Remark 2.1.3. Let (M, φ_M) be a Frobenius module over a commutative \mathbf{F}_p -algebra R . We will often abuse terminology by simply referring to M as a Frobenius module over R : in this case, we are implicitly asserting that M is equipped with a Frobenius-semilinear map $\varphi_M : M \rightarrow M$.

Example 2.1.4. Let R be a commutative \mathbf{F}_p -algebra. Then we can regard R as a Frobenius module over itself, via the Frobenius map

$$\varphi_R : R \rightarrow R \quad \varphi_R(\lambda) = \lambda^p.$$

More generally, the same comment applies to an ideal $I \subseteq R$.

It will sometimes be helpful to identify Frobenius modules over a commutative \mathbf{F}_p -algebra R with modules over a certain (noncommutative) enlargement of R .

Notation 2.1.5. Let R be an \mathbf{F}_p -algebra. We let $R[F]$ denote the noncommutative ring whose elements are finite sums $\sum_{n \geq 0} c_n F^n$, with multiplication given by

$$\left(\sum_{m \geq 0} c_m F^m \right) \left(\sum_{n \geq 0} c'_n F^n \right) = \sum_{k \geq 0} \left(\sum_{i+j=k} c_i c'_j \right) F^k.$$

We will identify R with the subring of $R[F]$ consisting of those sums $\sum_{n \geq 0} c_n F^n$ for which the coefficients c_i vanish for $i > 0$. Unwinding the definitions, we see that the category Mod_R^{Fr} is equivalent to the category

of left $R[F]$ -modules. In particular, Mod_R^{Fr} is an abelian category which has enough projective objects and enough injective objects. Given objects $M, N \in \text{Mod}_R^{\text{Fr}}$, we let $\text{Ext}_{R[F]}^n(M, N)$ denote the n th Ext-group of M and N , computed in the abelian category Mod_R^{Fr} .

We now consider the behavior of the category Mod_R^{Fr} as the commutative \mathbf{F}_p -algebra R varies.

Construction 2.1.6 (Extension of Scalars). Let $f : A \rightarrow B$ be a homomorphism of commutative \mathbf{F}_p -algebras. If M is a Frobenius module over A , then we can regard the tensor product $B \otimes_A M$ as a Frobenius module over B , with Frobenius map given by

$$\varphi_{B \otimes_A M}(b \otimes x) = b^p \otimes \varphi_M(x).$$

The construction $M \mapsto B \otimes_A M$ determines a functor from Mod_A^{Fr} to Mod_B^{Fr} , which we will denote by f_{Fr}^* and refer to as *extension of scalars along f* .

Remark 2.1.7 (Restriction of Scalars). In the situation of Construction 2.1.6, the extension of scalars $f_{\text{Fr}}^*(M) = B \otimes_A M$ is characterized by the following universal property: for every Frobenius module N over B , composition with the map $M \rightarrow B \otimes_A M$ induces a bijection

$$\text{Hom}_{B[F]}(B \otimes_A M, N) \rightarrow \text{Hom}_{A[F]}(M, N).$$

In other words, we can regard the functor f_{Fr}^* as a left adjoint to the forgetful functor $\text{Mod}_B^{\text{Fr}} \rightarrow \text{Mod}_A^{\text{Fr}}$. We will denote this forgetful functor by f_* and refer to it as *restriction of scalars along f* .

Warning 2.1.8. Let $f : A \rightarrow B$ be a homomorphism of commutative \mathbf{F}_p -algebras. Then f extends to a homomorphism of noncommutative rings $f^+ : A[F] \rightarrow B[F]$, where $A[F]$ and $B[F]$ are defined as in Notation 2.1.5. The content of Construction 2.1.6 and Remark 2.1.7 is that, for every Frobenius module M over A , the canonical map

$$B \otimes_A M \rightarrow B[F] \otimes_{A[F]} M$$

is an isomorphism. This relies on the fact that $A[F]$ is freely generated as a *left* A -module by the elements $\{F^n\}_{n \geq 0}$. Beware that $A[F]$ is usually not free when regarded as a *right* A -module (so the analogous compatibility would fail if we were to study right modules over $A[F]$, rather than left modules). In fact, following Notation 3.1.1, the ring $A[F]$ identifies with $\bigoplus_{n \geq 0} A \cdot F^n$ as a left A -module, and with $\bigoplus_{n \geq 0} F^n \cdot A^{1/p^n}$ as a right A -module. In particular, the latter is free over A only under strong conditions (such as regularity).

Remark 2.1.9. Let $f : A \rightarrow B$ be a homomorphism of commutative \mathbf{F}_p -algebras, and suppose that the multiplication map $m : B \otimes_A B \rightarrow B$ is an isomorphism (this condition is satisfied, for example, if f is surjective, or if f exhibits B as a localization of A). Then, for any Frobenius module M over B , the counit map $v : f_{\mathbf{F}_r}^* f_*(M) \rightarrow M$ is also an isomorphism: note that the domain of v can be identified with the tensor product $B \otimes_A M \simeq (B \otimes_A B) \otimes_B M$. It follows that the restriction of scalars functor $f_* : \text{Mod}_B^{\text{Fr}} \rightarrow \text{Mod}_A^{\text{Fr}}$ is fully faithful.

2.2. Étale Sheaves

For the reader's convenience, we briefly review the theory of étale sheaves. We consider here only the case of *affine* schemes (we will discuss sheaves on more general schemes in §10).

Notation 2.2.1. Let R be a commutative ring. We let $\text{CAlg}_R^{\text{ét}}$ denote the category whose objects are étale R -algebras, and whose morphisms are R -algebra homomorphisms.

Definition 2.2.2. Let R and Λ be commutative rings, and let Mod_Λ denote the category of Λ -modules. An *étale sheaf of Λ -modules on $\text{Spec}(R)$* is a functor

$$\mathcal{F} : \text{CAlg}_R^{\text{ét}} \rightarrow \text{Mod}_\Lambda$$

which satisfies the following pair of conditions:

- For every faithfully flat map $u : A \rightarrow B$ in $\text{CAlg}_R^{\text{ét}}$, the sequence

$$0 \rightarrow \mathcal{F}(A) \xrightarrow{\mathcal{F}(u)} \mathcal{F}(B) \xrightarrow{\mathcal{F}(u \otimes \text{id}) - \mathcal{F}(\text{id} \otimes u)} \mathcal{F}(B \otimes_A B)$$

is exact.

- For every finite collection of étale R -algebras $\{A_i\}_{i \in I}$, the map

$$\mathcal{F}\left(\prod_{i \in I} A_i\right) \rightarrow \prod_{i \in I} \mathcal{F}(A_i)$$

is an isomorphism.

We let $\text{Shv}_{\text{ét}}(\text{Spec}(R), \Lambda)$ denote the category whose objects are étale sheaves of Λ -modules on $\text{Spec}(R)$ (where morphisms are given by natural transformations).

Remark 2.2.3. In this paper, we will be concerned almost exclusively with the case where Λ is the finite field \mathbf{F}_p . The only exception is in §9, where we take $\Lambda = \mathbf{Z}/p^n\mathbf{Z}$ for some integer $n \geq 0$.

Example 2.2.4 (Constant Sheaves). Let R be a commutative ring and let M be a module over a commutative ring Λ . We let $\underline{M} \in \text{Shv}_{\text{ét}}(\text{Spec}(R), \Lambda)$ denote the functor which associates to each étale R -algebra A the set $\underline{M}(A)$ of locally constant M -valued functions on $\text{Spec}(A)$. We will refer to \underline{M} as the *constant sheaf with value M* .

Example 2.2.5 (Quasi-Coherent Sheaves). Let R be an \mathbf{F}_p -algebra. For every R -module M , the construction $(A \in \text{CAlg}_R^{\text{ét}}) \mapsto A \otimes_R M$ determines an étale sheaf of \mathbf{F}_p -modules on M , which we denote by \widetilde{M} (see [15, Tag 03DX]). Note that if M is a Frobenius module over R , then φ_M determines a map of étale sheaves $\widetilde{\varphi}_M : \widetilde{M} \rightarrow \widetilde{M}$; this map is \mathbf{F}_p -linear, but not R -linear in general.

Notation 2.2.6. Let R and Λ be commutative rings. If \mathcal{F} and \mathcal{G} are étale sheaves of Λ -modules on $\text{Spec}(R)$, we let $\text{Hom}_{\underline{\Lambda}}(\mathcal{F}, \mathcal{G})$ denote the abelian group of morphisms from \mathcal{F} to \mathcal{G} in the category $\text{Shv}_{\text{ét}}(\text{Spec}(R), \Lambda)$ (emphasizing the idea that \mathcal{F} and \mathcal{G} can be regarded as modules over the constant sheaf $\underline{\Lambda}$).

Note that $\text{Shv}_{\text{ét}}(\text{Spec}(R), \Lambda)$ is an abelian category with enough injective objects, so that we can consider Ext-groups in $\text{Shv}_{\text{ét}}(\text{Spec}(R), \Lambda)$. We denote these Ext-groups by $\text{Ext}_{\underline{\Lambda}}^n(\mathcal{F}, \mathcal{G})$ for $n \geq 0$.

Remark 2.2.7 (Functoriality). Let $f : A \rightarrow B$ be a homomorphism of commutative rings. Then f induces a base change functor

$$\text{CAlg}_A^{\text{ét}} \rightarrow \text{CAlg}_B^{\text{ét}} \quad A' \mapsto A' \otimes_A B.$$

Precomposition with this functor determines a pushforward functor

$$f_* : \text{Shv}_{\text{ét}}(\text{Spec}(B), \Lambda) \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(A), \Lambda),$$

given concretely by the formula $(f_* \mathcal{F})(A') = \mathcal{F}(A' \times_A B)$. The functor f_* admits a left adjoint $f^* : \text{Shv}_{\text{ét}}(\text{Spec}(A), \Lambda) \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(B), \Lambda)$, which we refer to as *pullback along f* .

2.3. The Solution Functor

Construction 2.3.1. Let R be a commutative \mathbf{F}_p -algebra and let M be a Frobenius module over R . For every étale R -algebra A , we regard the tensor

product $A \otimes_R M$ as a Frobenius module over A (see Construction 2.1.6). We define

$$\mathrm{Sol}(M)(A) = \{x \in (A \otimes_R M) : \varphi_{A \otimes_R M}(x) = x\}.$$

The construction $A \mapsto \mathrm{Sol}(M)(A)$ determines a functor $\mathrm{CAlg}_R^{\text{ét}} \rightarrow \mathrm{Mod}_{\mathbf{F}_p}$, which we will refer to as *the solution sheaf* of M .

Proposition 2.3.2. *Let R be a commutative \mathbf{F}_p -algebra and let M be a Frobenius module over R . Then $\mathrm{Sol}(M)$ is an étale sheaf of \mathbf{F}_p -modules on $\mathrm{Spec}(R)$.*

Proof. Let \widetilde{M} denote the quasi-coherent sheaf associated to M (Example 2.2.5). It follows immediately from the definition that $\mathrm{Sol}(M)$ can be described as the kernel of the map

$$(\mathrm{id} - \widetilde{\varphi}_M) : \widetilde{M} \rightarrow \widetilde{M}.$$

Since \widetilde{M} is an étale sheaf of \mathbf{F}_p -modules on $\mathrm{Spec}(R)$, the functor $\mathrm{Sol}(M)$ has the same property. \square

Definition 2.3.3. Let R be a commutative \mathbf{F}_p -algebra. We will regard the construction

$$(M \in \mathrm{Mod}_R^{\mathrm{Fr}}) \mapsto (\mathrm{Sol}(M) \in \mathrm{Shv}_{\text{ét}}(\mathrm{Spec}(R), \mathbf{F}_p))$$

as a functor $\mathrm{Sol} : \mathrm{Mod}_R^{\mathrm{Fr}} \rightarrow \mathrm{Shv}_{\text{ét}}(\mathrm{Spec}(R), \mathbf{F}_p)$. We will refer to Sol as *the solution functor*.

Remark 2.3.4. The solution functor $\mathrm{Sol} : \mathrm{Mod}_R^{\mathrm{Fr}} \rightarrow \mathrm{Shv}_{\text{ét}}(\mathrm{Spec}(R), \mathbf{F}_p)$ is left exact. However, it is usually not right exact.

2.4. The Riemann-Hilbert Correspondence

We now introduce a class of Frobenius modules on which the solution functor of Definition 2.3.3 is particularly well-behaved.

Definition 2.4.1. Let R be a commutative \mathbf{F}_p -algebra and let M be a Frobenius module over R . We will say that M is *algebraic* if it satisfies the following conditions:

- (a) The map $\varphi_M : M \rightarrow M$ is an isomorphism of abelian groups.
- (b) Every element $x \in M$ satisfies an equation of the form

$$\varphi_M^n(x) + a_1 \varphi_M^{n-1}(x) + \cdots + a_n x = 0$$

for some coefficients $a_i \in R$.

We let $\text{Mod}_R^{\text{alg}}$ denote the full subcategory of Mod_R^{Fr} spanned by the algebraic Frobenius modules over R .

Remark 2.4.2. Let R be a commutative \mathbf{F}_p -algebra and let M be a Frobenius module over R . Then condition (b) of Definition 2.4.1 is equivalent to the following:

- (b') For every finitely generated R -submodule $M_0 \subseteq M$, the Frobenius submodule of M generated by M_0 is also finitely generated as an R -module.

To see that (b) implies (b'), suppose that $M_0 \subseteq M$ is the R -submodule generated by finitely many elements $\{x_i\}_{i \in I}$. Then condition (b) guarantees that the R -submodule generated by $\{\varphi_M^k(x_i)\}_{i \in I, k \leq n}$ is stable under the action of φ_M for some $n \gg 0$; this is clearly the smallest φ_M -stable submodule of M which contains M_0 .

Conversely, suppose that (b') is satisfied and let x be an element of M . Applying condition (b') to the submodule $M_0 = Rx \subseteq M$, we see that the sum $\sum_{k \geq 0} R\varphi_M^k(x) \subseteq M$ is generated by finitely many elements, and is therefore contained in $\sum_{0 \leq k < n} R\varphi_M^k(x)$ for some integer n . It follows that $\varphi_M^n(x)$ can be written as a linear combination $a_1\varphi_M^{n-1}(x) + \dots + a_nx$ for some coefficients $a_1, a_2, \dots, a_n \in R$.

We can now formulate the main result of this paper:

Theorem 2.4.3. *Let R be a commutative \mathbf{F}_p -algebra. Then the solution sheaf functor $\text{Sol} : \text{Mod}_R^{\text{alg}} \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ is an equivalence of categories.*

Note that Theorem 2.4.3 immediately implies Theorem 1.0.2: assuming Theorem 2.4.3, the Riemann-Hilbert functor RH can be defined as the composition

$$\text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p) \xrightarrow{\text{Sol}^{-1}} \text{Mod}_R^{\text{alg}} \hookrightarrow \text{Mod}_R^{\text{Fr}}.$$

We will give a different (but ultimately equivalent) definition of the functor RH in §6: the construction of this functor is one of the key ingredients in our proof of Theorem 2.4.3, which we present in §7.

3. The Category of Frobenius Modules

Let R be a commutative \mathbf{F}_p -algebra. Our goal in this section is to establish some elementary properties of the abelian category Mod_R^{Fr} of Frobenius

modules over R . We begin in §3.1 by studying the forgetful functor from Mod_R^{Fr} to the category of R -modules. The main observation is that the ring R admits a very simple resolution

$$0 \rightarrow R[F] \xrightarrow{F-1} R[F] \rightarrow R \rightarrow 0$$

in the category of left modules over the noncommutative ring $R[F]$ appearing in Notation 2.1.5. This allows us to reduce calculations of Ext-groups in the category Mod_R^{Fr} to calculations of Ext-groups in the category Mod_R : see Construction 3.1.7.

In §3.2, we restrict our attention to the class of *perfect* Frobenius modules: that is, Frobenius modules M for which the map $\varphi_M : M \rightarrow M$ is bijective (Definition 3.2.1). The collection of Frobenius modules with this property form a category which we denote by $\text{Mod}_R^{\text{perf}}$. Our main result is that the inclusion functor $\text{Mod}_R^{\text{perf}} \hookrightarrow \text{Mod}_R^{\text{Fr}}$ admits an exact left adjoint $M \rightarrow M^{1/p^\infty}$, given informally by “inverting the Frobenius” (Proposition 3.2.7).

Let $f : A \rightarrow B$ be a homomorphism of commutative \mathbf{F}_p -algebras. In §2.1, we observed that extension of scalars along f determines a functor $f_{\text{Fr}}^* : \text{Mod}_A^{\text{Fr}} \rightarrow \text{Mod}_B^{\text{Fr}}$. Beware that this construction does *not* carry perfect Frobenius modules to perfect Frobenius modules. To remedy this, we introduce in §3.3 another functor $f^\diamond : \text{Mod}_A^{\text{perf}} \rightarrow \text{Mod}_B^{\text{perf}}$, given concretely by the formula $f^\diamond(M) = (f_{\text{Fr}}^* M)^{1/p^\infty}$. The functors f^\diamond and f_{Fr}^* are generally different, but they agree when the ring homomorphism f is étale (Corollary 3.4.7). The proof of this fact will require some elementary facts about perfect rings of characteristic p , which we review in §3.4.

3.1. Comparison of $R[F]$ -Modules with R -Modules

Throughout this section, we fix a commutative \mathbf{F}_p -algebra R .

Notation 3.1.1. Let Mod_R denote the abelian category of R -modules. For each $n \geq 0$, restriction of scalars along the n th power of the Frobenius map $\varphi_R : R \rightarrow R$ determines a forgetful functor from Mod_R to itself, which we will denote by $M \mapsto M^{1/p^n}$.

Let M be an R -module. Then there is a canonical isomorphism of abelian groups $M \simeq M^{1/p^n}$. For each element $x \in M$, we will denote the image of x under this isomorphism by x^{1/p^n} . The action of R on M^{1/p^n} can then be described by the formula $\lambda(x^{1/p^n}) = (\lambda^{p^n} x)^{1/p^n}$, for $\lambda \in R$ and $x \in M$.

Construction 3.1.2. Let N be an R -module. We let N^\dagger denote the R -module given by the product $\prod_{n \geq 0} N^{1/p^n}$. We identify elements of N^\dagger with the collection of all sequences (x_0, x_1, x_2, \dots) in N , where the action of R is given by

$$\lambda(x_0, x_1, x_2, \dots) = (\lambda x_0, \lambda^p x_1, \lambda^{p^2} x_2, \dots).$$

We regard N^\dagger as an Frobenius module over R , with endomorphism $\varphi_{N^\dagger} : N^\dagger \rightarrow N^\dagger$ given by

$$\varphi_{N^\dagger}(x_0, x_1, x_2, \dots) = (x_1, x_2, x_3, \dots).$$

Lemma 3.1.3. *Let M be a Frobenius module over R , let N be an arbitrary R -module, and let $v : N^\dagger \rightarrow N$ be the R -module homomorphism given by*

$$v(x_0, x_1, x_2, \dots) = x_0.$$

Then composition with v induces a bijection $\text{Hom}_R^{\text{Fr}}(M, N^\dagger) \rightarrow \text{Hom}_R(M, N)$.

Proof. For every R -module homomorphism $f : M \rightarrow N$, define $f^+ : M \rightarrow N^\dagger$ by the formula $f^+(x) = (f(x), f(\varphi_M(x)), f(\varphi_M^2(x)), \dots)$. An elementary calculation shows that the construction $f \mapsto f^+$ determines an inverse to the map $\text{Hom}_R^{\text{Fr}}(M, N^\dagger) \rightarrow \text{Hom}_R(M, N)$ given by composition with v . \square

Remark 3.1.4. It follows from Lemma 3.1.3 that we can regard the construction $N \mapsto N^\dagger$ as a right adjoint to the forgetful functor $\text{Mod}_R^{\text{Fr}} \rightarrow \text{Mod}_R$.

Remark 3.1.5. Using the equivalence of Notation 2.1.5, we can identify the forgetful map $\text{Mod}_R^{\text{Fr}} \rightarrow \text{Mod}_R$ with the functor given by restriction of scalars along the ring homomorphism $R \rightarrow R[F]$. The right adjoint to this restriction of scalars functor is given by $M \mapsto \text{Hom}_R(R[F], M)$, which we can identify with M^\dagger using the canonical left R -module basis of $R[F]$ given by $\{F^n\}_{n \geq 0}$.

Remark 3.1.6. Let M be a Frobenius module over R and let N be an arbitrary R -module. Then the identification $\text{Hom}_{R[F]}(M, N^\dagger) \simeq \text{Hom}_R(M, N)$ of Lemma 3.1.3 extends to an isomorphism of graded abelian groups

$$\text{Ext}_{R[F]}^*(M, N^\dagger) \simeq \text{Ext}_R^*(M, N).$$

This isomorphism can be described explicitly by choosing a projective resolution

$$\dots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow M$$

in the abelian category Mod_R^{Fr} . Note that each P_n is also projective when regarded as an R -module (this follows from the observation that the algebra $R[F]$ of Notation 2.1.5 is free as a left R -module), so that both $\text{Ext}_{R[F]}^*(M, N^\dagger)$ and $\text{Ext}_R^*(M, N)$ can be computed as the cohomology of the cochain complex of abelian groups $\text{Hom}_{R[F]}(P_*, N^\dagger) \simeq \text{Hom}_R(P_*, N)$.

Construction 3.1.7. Let N be a Frobenius module over R . Then the construction $(x \in N) \mapsto (x, \varphi_N(x), \varphi_N^2(x), \dots)$ determines a morphism of Frobenius modules $u : N \rightarrow N^\dagger$ (which is a unit map for the adjunction of Remark 3.1.4). Note that u is a monomorphism which fits into a short exact sequence of Frobenius modules

$$0 \rightarrow N \xrightarrow{u} N^\dagger \xrightarrow{\alpha} (N^{1/p})^\dagger \rightarrow 0,$$

where α is given by the formula

$$\alpha(x_0, x_1, x_2, \dots) = (\varphi_N(x_0) - x_1, \varphi_N(x_1) - x_2, \varphi_N(x_2) - x_3, \dots).$$

It follows that for any other Frobenius module M over R , we have a short exact sequence of abelian groups

$$0 \rightarrow \text{Hom}_R^{\text{Fr}}(M, N) \rightarrow \text{Hom}_R(M, N) \xrightarrow{\beta} \text{Hom}_R(M, N^{1/p}),$$

where β is given by the formula $\beta(f)(x) = \varphi_N(f(x))^{1/p} - f(\varphi_M(x))^{1/p}$. Moreover, if M is a projective object of Mod_R^{Fr} , then β is surjective. More generally, Remark 3.1.6 supplies a long exact sequence of abelian groups

$$\begin{aligned} \dots \rightarrow \text{Ext}_R^{*-1}(M, N^{1/p}) &\rightarrow \text{Ext}_{R[F]}^*(M, N) \rightarrow \text{Ext}_R^*(M, N) \\ &\rightarrow \text{Ext}_R^*(M, N^{1/p}) \rightarrow \dots \end{aligned}$$

Remark 3.1.8. Let R be a commutative \mathbf{F}_p -algebra and let M be a Frobenius module over R . It follows from Construction 3.1.7 that if M has projective (injective) dimension $\leq n$ as a module over R , then it has projective (injective) dimension $\leq n + 1$ as a module over $R[F]$.

3.2. Perfect Frobenius Modules

Let R be a commutative \mathbf{F}_p -algebra, which we regard as fixed throughout this section.

Definition 3.2.1. Let M be a Frobenius module over R . We will say that M is *perfect* if the map $\varphi_M : M \rightarrow M$ is an isomorphism of abelian groups. We let $\text{Mod}_R^{\text{perf}}$ denote the full subcategory of Mod_R^{Fr} spanned by the perfect Frobenius modules over R .

Remark 3.2.2. The full subcategory $\text{Mod}_R^{\text{perf}} \subseteq \text{Mod}_R^{\text{Fr}}$ is closed under limits, colimits, and extensions. In particular, $\text{Mod}_R^{\text{perf}}$ is an abelian category, and the inclusion functor $\text{Mod}_R^{\text{perf}} \hookrightarrow \text{Mod}_R^{\text{Fr}}$ is exact.

Notation 3.2.3. Let M be a Frobenius module over R . We let M^{1/p^∞} denote the colimit of the sequence

$$M \xrightarrow{\varphi_M} M^{1/p} \xrightarrow{\varphi_M} M^{1/p^2} \rightarrow \dots$$

We will refer to M^{1/p^∞} as the *perfection* of M .

Example 3.2.4. Let us regard R as a Frobenius module over itself as in Example 2.1.4. Then R^{1/p^∞} is the perfection of R in the sense of commutative algebra: that is, it is an initial object in the category of R -algebras in which every element admits a unique p th root.

Example 3.2.5. Let $R[F]$ be as in Notation 2.1.5, which we regard as a Frobenius module over R . We will denote the perfection of $R[F]$ by $R^{1/p^\infty}[F^{\pm 1}]$. Unwinding the definitions, we can identify elements of $R^{1/p^\infty}[F^{\pm 1}]$ with expressions of the form $\sum_{n \in \mathbf{Z}} c_n F^n$ where the coefficients $c_n \in R^{1/p^\infty}$ vanish for all but finitely many integers n .

Remark 3.2.6. The set $R^{1/p^\infty}[F^{\pm 1}]$ has the structure of an associative ring, with multiplication given by the formula

$$\left(\sum_{m \in \mathbf{Z}} c_m F^m\right) \left(\sum_{n \in \mathbf{Z}} c'_n F^n\right) = \sum_{k \in \mathbf{Z}} \left(\sum_{i+j=k} c_i c'_j\right) F^k.$$

This ring can be obtained from the associative ring $R[F]$ by formally adjoining an inverse of the element F . It follows that the equivalence of Mod_R^{Fr} with the category of left $R[F]$ -modules restricts to an equivalence of $\text{Mod}_R^{\text{perf}}$ with the category of left $R^{1/p^\infty}[F^{\pm 1}]$ -modules. In particular, $\text{Mod}_R^{\text{perf}}$ is an abelian category which has enough projective objects and enough injective objects.

In the situation of Notation 3.2.3, the perfection M^{1/p^∞} inherits the structure of a Frobenius module. Moreover, it enjoys the following universal property:

Proposition 3.2.7. *The inclusion functor $\iota : \text{Mod}_R^{\text{perf}} \hookrightarrow \text{Mod}_R^{\text{Fr}}$ admits a left adjoint, which carries a Frobenius module M to its perfection M^{1/p^∞} .*

Proof. Under the equivalence of Remark 3.2.6, a left adjoint to ι corresponds to the functor of extension of scalars along the map $R[F] \rightarrow R^{1/p^\infty}[F^{\pm 1}]$, which is given by $M \mapsto M^{1/p^\infty}$. \square

Remark 3.2.8. The perfection functor $M \mapsto M^{1/p^\infty}$ is exact. It follows that the inclusion functor $\text{Mod}_R^{\text{perf}} \hookrightarrow \text{Mod}_R^{\text{Fr}}$ carries injective objects to injective objects. In particular, if M and N are perfect Frobenius module over R , then the canonical map $\text{Ext}_{\text{Mod}_R^{\text{perf}}}^*(M, N) \rightarrow \text{Ext}_{\text{Mod}_R^{\text{Fr}}}^*(M, N)$ is an isomorphism. We will denote either of these Ext-groups by $\text{Ext}_{R[F]}^*(M, N)$.

For the purpose of comparing Frobenius modules with étale sheaves, there is no harm in restricting our considerations to perfect Frobenius modules:

Proposition 3.2.9. *Let $f : M \rightarrow N$ be a morphism of Frobenius modules over R . If the induced map $M^{1/p^\infty} \rightarrow N^{1/p^\infty}$ is an isomorphism of perfect Frobenius modules, then the induced map $\text{Sol}(M) \rightarrow \text{Sol}(N)$ is an isomorphism of étale sheaves.*

Proof. Factoring f as a composition $M \rightarrow \text{im}(f) \rightarrow N$, we can reduce to proving Proposition 3.2.9 in the special case where f is assumed to be either surjective or injective. Suppose first that f is injective. Our hypothesis that f induces an isomorphism $M^{1/p^\infty} \rightarrow N^{1/p^\infty}$ guarantees that the perfection $(N/M)^{1/p^\infty}$ vanishes: that is, the Frobenius map $\varphi_{N/M}$ is locally nilpotent. It follows that for any étale R -algebra A , the Frobenius map $\varphi_{A \otimes_R (N/M)}$ is also locally nilpotent, and therefore has no nonzero fixed points. It follows that the étale sheaf $\text{Sol}(N/M)$ vanishes. Since the solution functor is left exact (Remark 2.3.4), we have an exact sequence of étale sheaves $0 \rightarrow \text{Sol}(M) \rightarrow \text{Sol}(N) \rightarrow \text{Sol}(N/M)$, which proves that $\text{Sol}(f)$ is an isomorphism.

We now treat the case where f is surjective. We wish to prove that f induces an isomorphism $\text{Sol}(M)(A) \rightarrow \text{Sol}(N)(A)$ for every étale R -algebra A . Replacing M and N by $A \otimes_R M$ and $A \otimes_R N$ (which does not injure our assumption that f induces an equivalence of perfection: see Proposition 3.3.2), we can reduce to the case $A = R$. We have a commutative diagram of short exact sequence

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \ker(f) & \longrightarrow & M & \xrightarrow{f} & N \longrightarrow 0 \\
 & & \downarrow \text{id} - \varphi_{\ker(f)} & & \downarrow \text{id} - \varphi_M & & \downarrow \text{id} - \varphi_N \\
 0 & \longrightarrow & \ker(f) & \longrightarrow & M & \xrightarrow{f} & N \longrightarrow 0
 \end{array}$$

which gives a short exact sequence

$$\ker(\text{id} - \varphi_{\ker(f)}) \rightarrow \text{Sol}(M)(R) \rightarrow \text{Sol}(N)(R) \rightarrow \text{coker}(\text{id} - \varphi_{\ker(f)}).$$

It will therefore suffice to show that the map $\text{id} - \varphi_{\ker(f)}$ is an isomorphism. This is clear: our assumption that f induces an equivalence of perfections guarantees that $\varphi_{\ker(f)}$ is locally nilpotent, so that $\text{id} - \varphi_{\ker(f)}$ has an inverse given by the infinite sum $\sum_{n \geq 0} \varphi_{\ker(f)}^n$. \square

3.3. Restriction and Extension of Scalars

Let $f : A \rightarrow B$ be a homomorphism of commutative \mathbf{F}_p -algebras and let M be a Frobenius module over B . Then M is perfect as a Frobenius module over B if and only if it is perfect when regarded as a Frobenius module over A . In particular, the restriction of scalars functor $f_* : \text{Mod}_B^{\text{Fr}} \rightarrow \text{Mod}_A^{\text{Fr}}$ carries $\text{Mod}_B^{\text{perf}}$ into $\text{Mod}_A^{\text{perf}}$. We will abuse notation by denoting the induced map $\text{Mod}_B^{\text{perf}} \rightarrow \text{Mod}_A^{\text{perf}}$ also by f_* , so that we have a commutative diagram σ :

$$\begin{array}{ccc} \text{Mod}_B^{\text{perf}} & \longrightarrow & \text{Mod}_B^{\text{Fr}} \\ \downarrow f_* & & \downarrow f_* \\ \text{Mod}_A^{\text{perf}} & \longrightarrow & \text{Mod}_A^{\text{Fr}}. \end{array}$$

Remark 3.3.1. Let $f : A \rightarrow B$ be a homomorphism of commutative \mathbf{F}_p -algebras and let M be a Frobenius module over B . Then the canonical map $u : M \rightarrow M^{1/p^\infty}$ induces a map $f_*(u) : f_*(M) \rightarrow f_*(M^{1/p^\infty})$ of Frobenius modules over A whose target is perfect. It follows that $f_*(u)$ extends uniquely to a map $v : f_*(M)^{1/p^\infty} \rightarrow f_*(M^{1/p^\infty})$. Moreover, the map v is an isomorphism: this follows by inspecting the construction of the perfection given in §3.2. Put more informally, the formation of the perfection M^{1/p^∞} does not depend on whether we regard M as a Frobenius module over B or over A (or over \mathbf{F}_p).

In the situation of Remark 3.3.1, the extension of scalars functor $f_{\text{Fr}}^* : \text{Mod}_A^{\text{Fr}} \rightarrow \text{Mod}_B^{\text{Fr}}$ usually does not carry perfect Frobenius modules to perfect Frobenius modules. However, we do have the following:

Proposition 3.3.2. *Let $f : A \rightarrow B$ be a homomorphism of commutative \mathbf{F}_p -algebras. Then the forgetful functor $\text{Mod}_B^{\text{perf}} \rightarrow \text{Mod}_A^{\text{perf}}$ admits a left*

adjoint f^\diamond . Moreover, the diagram of categories

$$\begin{array}{ccc} \mathrm{Mod}_A^{\mathrm{Fr}} & \xrightarrow{(-)^{1/p^\infty}} & \mathrm{Mod}_A^{\mathrm{perf}} \\ \downarrow f_{\mathrm{Fr}}^* & & \downarrow f^\diamond \\ \mathrm{Mod}_B^{\mathrm{Fr}} & \xrightarrow{(-)^{1/p^\infty}} & \mathrm{Mod}_B^{\mathrm{perf}} \end{array}$$

commutes up to canonical isomorphism. More precisely, for every object $M \in \mathrm{Mod}_A^{\mathrm{Fr}}$, the canonical map $f^\diamond(M^{1/p^\infty}) \rightarrow (f_{\mathrm{Fr}}^* M)^{1/p^\infty}$ is an equivalence.

Proof. Defining f^\diamond by the formula $f^\diamond(M) = (f_{\mathrm{Fr}}^* M)^{1/p^\infty}$, it follows immediately from the definitions that f^\diamond is left adjoint to the forgetful functor $f_* : \mathrm{Mod}_B^{\mathrm{perf}} \rightarrow \mathrm{Mod}_A^{\mathrm{perf}}$. The desired commutativity follows from the commutativity of the diagram σ above (by passing to left adjoints). \square

In the situation of Proposition 3.3.2, the functors

$$f_{\mathrm{Fr}}^* : \mathrm{Mod}_A^{\mathrm{Fr}} \rightarrow \mathrm{Mod}_B^{\mathrm{Fr}} \quad f^\diamond : \mathrm{Mod}_A^{\mathrm{perf}} \rightarrow \mathrm{Mod}_B^{\mathrm{perf}}$$

are right exact, but generally not left exact (unless B is flat over A). We can therefore consider their left derived functors.

Construction 3.3.3. Let $f : A \rightarrow B$ be a homomorphism of \mathbf{F}_p -algebras. The abelian category $\mathrm{Mod}_A^{\mathrm{Fr}}$ has enough projective objects, so that the extension of scalars functor $f_{\mathrm{Fr}}^* : \mathrm{Mod}_A^{\mathrm{Fr}} \rightarrow \mathrm{Mod}_B^{\mathrm{Fr}}$ has left derived functors $L_n f_{\mathrm{Fr}}^* : \mathrm{Mod}_A^{\mathrm{Fr}} \rightarrow \mathrm{Mod}_B^{\mathrm{Fr}}$ for $n \geq 0$. More concretely, for $M \in \mathrm{Mod}_A^{\mathrm{Fr}}$, we can describe $L_n f_{\mathrm{Fr}}^* M$ as the n th homology of the chain complex $f_{\mathrm{Fr}}^*(P_*)$, where P_* is a projective resolution of M in the abelian category $\mathrm{Mod}_A^{\mathrm{Fr}}$. Note that P_* is then also a projective resolution of M in the category Mod_A of A -modules, and that the chain complex $f_{\mathrm{Fr}}^*(P_*)$ can be identified with $B \otimes_A P_*$. It follows that for $n \geq 0$, we have canonical B -module isomorphisms $L_n f_{\mathrm{Fr}}^* M = \mathrm{Tor}_n^A(M, B)$. We can summarize the situation as follows:

- (*) If $f : A \rightarrow B$ is a homomorphism of \mathbf{F}_p -algebras and M is a Frobenius module over A , then the Tor-groups $\mathrm{Tor}_n^A(M, B)$ can be regarded as Frobenius modules over B . Moreover, the construction

$$\mathrm{Mod}_A^{\mathrm{Fr}} \rightarrow \mathrm{Mod}_B^{\mathrm{Fr}} \quad M \mapsto \mathrm{Tor}_n^A(M, B)$$

can be identified with the n th left derived functor of the construction $M \mapsto f_{\mathrm{Fr}}^* M$.

Variante 3.3.4. In the situation of Construction, the abelian category $\text{Mod}_A^{\text{perf}}$ also has enough projective objects, so we can consider the left derived functors $L_n f^\diamond$ of the functor $f^\diamond : \text{Mod}_A^{\text{perf}} \rightarrow \text{Mod}_B^{\text{perf}}$. Note that if M is a perfect Frobenius module over A and P_* is a projective resolution of M in the category Mod_A^{Fr} of all Frobenius modules over A , then P_*^{1/p^∞} is a projective resolution of $M^{1/p^\infty} \simeq M$ in the category $\text{Mod}_A^{\text{perf}}$ of perfect Frobenius modules over A . We can therefore identify $(L_n f^\diamond)(M)$ with the n th homology group of the chain complex $f^\diamond P_*^{1/p^\infty} \simeq (f_{\text{Fr}}^* P_*)^{1/p^\infty}$. Using the exactness of the functor $N \mapsto N^{1/p^\infty}$, we obtain isomorphisms

$$(L_n f^\diamond)(M) \simeq (L_n f_{\text{Fr}}^*)(M)^{1/p^\infty} \simeq \text{Tor}_n^A(M, B)^{1/p^\infty}.$$

3.4. Perfect Rings

Let R be a commutative \mathbf{F}_p -algebra. Recall that R is said to be *perfect* if the Frobenius homomorphism $\varphi_R : R \rightarrow R$ is an isomorphism.

Remark 3.4.1. A commutative \mathbf{F}_p -algebra R is perfect if and only if it is perfect when regarded as a Frobenius module over itself, in the sense of Definition 3.2.1.

Example 3.4.2. Let R be any commutative \mathbf{F}_p -algebra. Then the perfection R^{1/p^∞} of Example 3.2.4 is a perfect \mathbf{F}_p -algebra.

Let R be an \mathbf{F}_p -algebra. If M is a perfect Frobenius module over R , then M admits the structure of a module over R^{1/p^∞} . More precisely, we have the following result, whose proof is left to the reader:

Proposition 3.4.3. *Let R be an algebra over \mathbf{F}_p . Then the restriction of scalars functor $\text{Mod}_{R^{1/p^\infty}}^{\text{perf}} \rightarrow \text{Mod}_R^{\text{perf}}$ is an equivalence of categories.*

Remark 3.4.4. Let R be a Noetherian \mathbf{F}_p -algebra of finite Krull dimension d , and suppose that the Frobenius map $\varphi_R : R \rightarrow R^{1/p}$ exhibits $R^{1/p}$ as a finite module over R . Then the abelian category of all R^{1/p^∞} -modules has global dimension $\leq 2d+1$ (see [1, Remark 11.33]). It follows from Proposition 3.4.3 and Construction 3.1.7 that the category $\text{Mod}_R^{\text{perf}}$ has global dimension $\leq 2d+2$.

Proposition 3.4.5. *Let $f : A \rightarrow B$ be a homomorphism of perfect \mathbf{F}_p -algebras. Then the extension of scalars functor $f_{\text{Fr}}^* : \text{Mod}_A^{\text{Fr}} \rightarrow \text{Mod}_B^{\text{Fr}}$ carries $\text{Mod}_A^{\text{perf}}$ into $\text{Mod}_B^{\text{perf}}$.*

Proof. Let M be a perfect Frobenius module over A . Then the maps

$$\varphi_B : B \rightarrow B \quad \varphi_A : A \rightarrow A \quad \varphi_M : M \rightarrow M$$

are isomorphisms, so the induced map $\varphi_{B \otimes_A M} : B \otimes_A M \rightarrow B \otimes_A M$ is also an isomorphism. \square

Warning 3.4.6. In the proof of Proposition 3.4.5, it is not enough to assume that M and B are perfect. For example, the tensor product $\mathbf{F}_p[x]^{1/p^\infty} \otimes_{\mathbf{F}_p[x]} \mathbf{F}_p[x]^{1/p^\infty}$ is not a perfect ring.

Corollary 3.4.7. *Let $f : A \rightarrow B$ be an étale morphism of \mathbf{F}_p -algebras. Then the extension of scalars functor $f_{\text{Fr}}^* : \text{Mod}_A^{\text{Fr}} \rightarrow \text{Mod}_B^{\text{Fr}}$ carries $\text{Mod}_A^{\text{perf}}$ into $\text{Mod}_B^{\text{perf}}$.*

Proof. Let M be a perfect Frobenius module over A . Then we can also regard M as a Frobenius module over A^{1/p^∞} . Since f is étale, the diagram of commutative rings

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow \\ A^{1/p^\infty} & \longrightarrow & B^{1/p^\infty} \end{array}$$

is a pushout square (see, for example, [15, Tag 0EBS]). It follows that we can identify $f_{\text{Fr}}^* M$ with the tensor product $B^{1/p^\infty} \otimes_{A^{1/p^\infty}} M$, which is perfect by Proposition 3.4.5. \square

3.5. Exactness Properties of f^\diamond

Our final goal in this section is to establish the following fundamental exactness property for pullbacks of algebraic Frobenius modules:

Theorem 3.5.1. *Let $f : A \rightarrow B$ be a homomorphism of commutative \mathbf{F}_p -algebras and let M be an algebraic Frobenius module over A . Then the abelian groups $\text{Tor}_n^A(M, B)^{1/p^\infty}$ vanish for $n > 0$.*

Before giving the proof of Theorem 3.5.1, let us collect some consequences:

Corollary 3.5.2. *Let $f : A \rightarrow B$ be a homomorphism of commutative \mathbf{F}_p -algebras and suppose we are given an exact sequence $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ in $\text{Mod}_A^{\text{perf}}$. If M'' is algebraic, then the sequence $0 \rightarrow f^\diamond M' \rightarrow f^\diamond M \rightarrow f^\diamond M'' \rightarrow 0$ is also exact.*

Proof. Variant 3.3.4 supplies an exact sequence

$$\mathrm{Tor}_1^A(M'', B)^{1/p^\infty} \rightarrow f^\diamond M' \rightarrow f^\diamond M \rightarrow f^\diamond M'' \rightarrow 0,$$

where the first term vanishes by virtue of Theorem 3.5.1. □

Corollary 3.5.3. *Let $f : A \rightarrow B$ be a morphism of \mathbf{F}_p -algebras and suppose we are given objects $M \in \mathrm{Mod}_A^{\mathrm{Fr}}$, $N \in \mathrm{Mod}_B^{\mathrm{Fr}}$. If M is algebraic and N is perfect, then the canonical map $\mathrm{Ext}_{A[F]}^*(M, N) \rightarrow \mathrm{Ext}_{B[F]}^*(f^\diamond M, N)$ is an isomorphism.*

Proof. Let P_* be a projective resolution of M in the category $\mathrm{Mod}_A^{\mathrm{perf}}$. Then Theorem 3.5.1 guarantees that $f^\diamond P_*$ is a projective resolution of $f^\diamond M$ in the category $\mathrm{Mod}_B^{\mathrm{perf}}$, so that both $\mathrm{Ext}_{A[F]}^*(M, N)$ and $\mathrm{Ext}_{B[F]}^*(f^\diamond M, N)$ can be identified with the cohomology of the cochain complex $\mathrm{Hom}_{A[F]}(P_*, N) \simeq \mathrm{Hom}_{B[F]}(f^\diamond P_*, N)$. □

We now turn to the proof of Theorem 3.5.1. The main ingredient is the following observation from [1]:

Lemma 3.5.4. *Let A be a perfect \mathbf{F}_p -algebra containing an element a , and let $I = (a, a^{1/p}, a^{1/p^2}, \dots)$ denote the kernel of the map $A \rightarrow (A/(a))^{1/p^\infty}$. Then the elements $a^{1/p^n} \in I$ determine an A -module isomorphism of I with the direct limit of the diagram*

$$A \xrightarrow{a^{1-1/p}} A \xrightarrow{a^{1/p-1/p^2}} A \xrightarrow{a^{1/p^2-1/p^3}} A \xrightarrow{a^{1/p^3-1/p^4}} \dots$$

Proof. Unwinding the definitions, we must show that if an element $x \in A$ satisfies the equation $xa^{1/p^m} = 0$ for some $m \geq 0$, then we have $xa^{1/p^m-1/p^n} = 0$ for some $n \gg m$. We now compute

$$\begin{aligned} xa^{1/p^m-1/p^{m+1}} &= x^{1/p} x^{(p-1)/p} a^{1/p^{m+1}} a^{(p-2)/p^{m+1}} \\ &= (xa^{1/p^m})^{1/p} x^{(p-1)/p} a^{(p-2)/p^{m+1}} \\ &= 0. \end{aligned}$$

□

Proof of Theorem 3.5.1. Let $f : A \rightarrow B$ be a homomorphism of commutative \mathbf{F}_p -algebras and let M be an algebraic Frobenius module over A ; we wish to show that the groups $\mathrm{Tor}_n^A(M, B)^{1/p^\infty}$ vanish for $n > 0$. Writing B as a filtered direct limit of finitely generated A -algebras, we can assume that B is finitely presented over A : that is, we can write $B \simeq A[x_1, \dots, x_k]/I$ for some

finitely generated ideal $I \subseteq A[x_1, \dots, x_k]$. Set $N = (A[x_1, \dots, x_k] \otimes_A M)^{1/p^\infty}$. Using the flatness of $A[x_1, \dots, x_k]$ over A , we obtain isomorphisms

$$\mathrm{Tor}_*^A(M, B)^{1/p^\infty} \simeq \mathrm{Tor}_*^{A[x_1, \dots, x_k]}(N, B)^{1/p^\infty}.$$

Moreover, N is an algebraic Frobenius module over $A[x_1, \dots, x_k]$ (see Corollary 4.2.8). We may therefore replace A by $A[x_1, \dots, x_k]$ (and M by N) and thereby reduce to the case where f is surjective.

Proceeding by induction on the number of generators of I , we can reduce to the case where $I = (a)$ is a principal ideal. Since M is perfect, we can regard M as a module over the perfection A^{1/p^∞} , so that we have canonical isomorphisms $\mathrm{Tor}_*^A(M, B)^{1/p^\infty} \simeq \mathrm{Tor}_*^{A^{1/p^\infty}}(M, B^{1/p^\infty})$. We have an exact sequence

$$0 \rightarrow I^{1/p^\infty} \rightarrow A^{1/p^\infty} \rightarrow B^{1/p^\infty} \rightarrow 0$$

in the category of modules over A^{1/p^∞} , where I^{1/p^∞} is flat over A^{1/p^∞} by virtue of Lemma 3.5.4. It follows that the groups $\mathrm{Tor}_n^A(M, B)^{1/p^\infty}$ vanish for $n \geq 2$, and $\mathrm{Tor}_1^A(M, B)^{1/p^\infty}$ can be identified with the kernel of the map

$$\rho : M \otimes_{A^{1/p^\infty}} I^{1/p^\infty} \rightarrow M \otimes_{A^{1/p^\infty}} A^{1/p^\infty} \simeq M.$$

We will complete the proof by showing that ρ is injective. Using the description of I^{1/p^∞} supplied by Lemma 3.5.4, we see that the injectivity of ρ can be reformulated as follows:

- (*) Let x be an element of M which satisfies the equation $a^{1/p^m} x = 0$, for some integer m . Then $a^{1/p^m - 1/p^{m'}} x = 0$ for some $m' \gg m$.

To prove (*), we use our assumption that M is algebraic to write

$$\varphi_M^n(x) = c_1 \varphi_M^{n-1}(x) + \dots + c_n x$$

for some coefficients $c_1, \dots, c_n \in A$. We then compute

$$\begin{aligned} \varphi_M^n(a^{1/p^{m+1}} x) &= a^{p^n/p^{m+1}} \varphi_M^n(x) \\ &= \sum_{i=1}^n a^{p^n/p^{m+1}} c_i \varphi_M^{n-i}(x) \\ &= \sum_{i=1}^n c_i \varphi_M^{n-i}(a^{p^{i-1}/p^m} x) \\ &= 0. \end{aligned}$$

Using the bijectivity of φ_M , it follows that $a^{1/p^{m+1}}x = 0$, which immediately implies (*). \square

Remark 3.5.5. The reasoning used to prove Theorem 3.5.1 can also be used to show the following result (see [1]): if $B \leftarrow A \rightarrow C$ is a diagram of perfect rings, then $\mathrm{Tor}_A^i(B, C) = 0$ for $i > 0$. Indeed, as in the proof of Theorem 3.5.1, one reduces to the case $B = A/I$, where $I = \bigcup_n f^{1/p^n}A$ is the radical of an ideal generated by a single element $f \in A$. In this case, the presentation given in Lemma 3.5.4 and the perfectness of C imply that I is a flat A -module, and that $I \otimes_A C \simeq J$, where $J = \bigcup_n f^{1/p^n}C \subseteq C$ is also an ideal. The desired claim follows immediately.

4. Holonomic Frobenius Modules

Let R be a commutative \mathbf{F}_p -algebra and let $\mathrm{Mod}_R^{\mathrm{Fr}}$ denote the category of Frobenius modules over R . In this section, we consider a full subcategory $\mathrm{Mod}_R^{\mathrm{hol}} \subseteq \mathrm{Mod}_R^{\mathrm{Fr}}$ whose objects we refer to as *holonomic Frobenius modules*. Roughly speaking, the class of holonomic Frobenius modules can be regarded as a characteristic p analogue of the class of (regular) holonomic \mathcal{D} -modules on complex analytic varieties. We will later show that the category $\mathrm{Mod}_R^{\mathrm{hol}}$ can be characterized as the essential image of the category $\mathrm{Shv}_{\mathrm{ét}}^c(\mathrm{Spec}(R), \mathbf{F}_p)$ of *constructible étale sheaves* under the Riemann-Hilbert equivalence

$$\mathrm{RH} : \mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(R), \mathbf{F}_p) \simeq \mathrm{Mod}_R^{\mathrm{alg}} \subseteq \mathrm{Mod}_R^{\mathrm{Fr}}$$

of Theorem 1.0.2 (see Theorem 7.4.1). Our goal in this section is to lay the groundwork by establishing the basic formal properties of $\mathrm{Mod}_R^{\mathrm{hol}}$.

We begin in §4.1 by defining the class of holonomic Frobenius modules (Definition 4.1.1) and verifying some elementary closure properties. In §4.2, we show that every holonomic Frobenius module is algebraic (Proposition 4.2.1) and that, conversely, every algebraic Frobenius module can be realized as a filtered colimit of holonomic Frobenius modules (Theorem 4.2.9). This result will allow us to reduce certain questions about algebraic modules to the case of holonomic modules, which enjoy good finiteness properties. For example, we prove in §4.3 that if R is Noetherian, then the category $\mathrm{Mod}_R^{\mathrm{hol}}$ is also Noetherian (Proposition 4.3.1). In §4.4 we associate to each holonomic Frobenius module M a constructible subset $\mathrm{supp}(M) \subseteq \mathrm{Spec}(R)$ which we refer to as *the support of M* . We will see later that the support $\mathrm{supp}(M)$ exerts strong control over the behavior of M : for example, it is empty if and only if $M \simeq 0$ (Proposition 5.3.3).

4.1. Holonomicity

Definition 4.1.1. Let R be a commutative \mathbf{F}_p -algebra and let M be a Frobenius module over R . We will say that M is *holonomic* if there exists an isomorphism $M \simeq M_0^{1/p^\infty}$, where $M_0 \in \text{Mod}_R^{\text{Fr}}$ is finitely presented as an R -module. We let $\text{Mod}_R^{\text{hol}}$ denote the full subcategory of $\text{Mod}_R^{\text{perf}}$ spanned by the holonomic Frobenius modules over R .

Proposition 4.1.2. *Let $f : A \rightarrow B$ be a homomorphism of \mathbf{F}_p -algebras. If $M \in \text{Mod}_A^{\text{perf}}$ is holonomic, then $f^\diamond M \in \text{Mod}_B^{\text{perf}}$ is holonomic.*

Proof. Without loss of generality we may assume that $M = M_0^{1/p^\infty}$ for some $M_0 \in \text{Mod}_A^{\text{Fr}}$ which is finitely presented as an A -module. Then $f^\diamond M \simeq (B \otimes_A M_0)^{1/p^\infty}$, and $B \otimes_A M_0$ is finitely presented as a B -module. \square

We also have the following converse of Proposition 4.1.2, whose proof we leave to the reader:

Proposition 4.1.3. *Let R be an \mathbf{F}_p -algebra and let M be a holonomic Frobenius module over R . Then there exists an inclusion $\iota : R' \hookrightarrow R$ where R' is finitely generated over \mathbf{F}_p and an isomorphism $M \simeq \iota^\diamond M'$, where M' is a holonomic Frobenius module over R' .*

Remark 4.1.4. Let R be a commutative \mathbf{F}_p -algebra and suppose we are given objects $M, N \in \text{Mod}_R^{\text{perf}}$, where M is holonomic. Then we can choose an isomorphism $M \simeq M_0^{1/p^\infty}$ for some $M_0 \in \text{Mod}_R^{\text{Fr}}$ which is finitely presented as an R -module. Using Construction 3.1.7 (and the observation that $\varphi_N : N \rightarrow N^{1/p}$ is an isomorphism), we obtain a long exact sequence

$$\text{Ext}_{R[F]}^*(M, N) \rightarrow \text{Ext}_R^*(M_0, N) \xrightarrow{\gamma} \text{Ext}_R^*(M_0, N) \rightarrow \text{Ext}_{R[F]}^{*+1}(M, N),$$

where γ is the map given by $\gamma(f) = f - \varphi_N^{-1} \circ f \circ \varphi_{M_0}$.

Proposition 4.1.5. *Let R be an \mathbf{F}_p -algebra and let M be a Frobenius module over R which is holonomic. Then M is a compact object of the category $\text{Mod}_R^{\text{perf}}$: that is, the functor $N \mapsto \text{Hom}_R^{\text{Fr}}(M, N)$ commutes with filtered colimits.*

Proof. Choose an isomorphism $M \simeq M_0^{1/p^\infty}$ for some $M_0 \in \text{Mod}_R^{\text{Fr}}$ which is finitely presented as an R -module, and observe that the exact sequence of Remark 4.1.4 depends functorially on N . \square

4.2. Comparison with Algebraic Frobenius Modules

Our next goal is to compare the theory of holonomic Frobenius modules (introduced in Definition 4.1.1) with the theory of algebraic Frobenius (introduced in Definition 2.4.1). Our starting point is the following:

Proposition 4.2.1. *Let R be a commutative \mathbf{F}_p -algebra and let M be a holonomic Frobenius module over R . Then M is algebraic.*

Proof. Let x be an element of M . Using Proposition 4.1.3, we can choose a finitely generated subalgebra $R' \subseteq R$ and an isomorphism $M \simeq (R \otimes_{R'} M')^{1/p^\infty}$ for some $M' \in \text{Mod}_{R'}^{\text{hol}}$. Enlarging R' if necessary, we may assume that x is the image of some element $x' \in M'$. Since M' is holonomic, we can write $M' = M'_0{}^{1/p^\infty}$ for some $M'_0 \in \text{Mod}_{R'}^{\text{Fr}}$ which is finitely presented as an R' -module. We can then write $x' = \varphi_{M'}^{-k}(y)$ for some $y \in M'$ which lifts to an element $y_0 \in M'_0$. Since M'_0 is a Noetherian R' -module, the submodule generated by the elements $\{\varphi_{M'_0}^n(y_0)\}_{n \geq 0}$ is finitely generated. It follows that y_0 satisfies an equation $\varphi_{M'_0}^n(y) + a_1 \varphi_{M'_0}^{n-1}(y) + \dots + a_n y = 0$ for some elements $a_i \in R'$, so that x satisfies the equation $\varphi_M^n(x) + a_1^{p^k} \varphi_M^{n-1}(x) + \dots + a_n^{p^k} x = 0$. \square

Corollary 4.2.2. *Let R be a commutative \mathbf{F}_p -algebra. Then the collection of holonomic Frobenius modules over R is closed under finite direct sums and cokernels.*

Remark 4.2.3. We will see later that the collection of holonomic Frobenius modules is also closed under the formation of kernels and extensions; in particular, it is an abelian subcategory of Mod_R^{Fr} (Corollary 4.3.3).

Proof of Corollary 4.2.2. Closure under finite direct sums is obvious. Let $u : M \rightarrow N$ be a morphism in $\text{Mod}_R^{\text{hol}}$. Then we can choose isomorphisms $M \simeq M_0^{1/p^\infty}$ and $N \simeq N_0^{1/p^\infty}$ for some objects $M_0, N_0 \in \text{Mod}_R^{\text{Fr}}$ which are finitely presented as R -modules. Let x_1, \dots, x_k be a set of generators for M_0 as an R -module. Let us abuse notation by writing $u(x_i)$ for the image of x_i under the composite map $M_0 \rightarrow M \xrightarrow{u} N$. Then we can choose some integer $n \gg 0$ for which each $u(x_i)$ has the form $\varphi_N^{-n}(y_i)$ for some y_i which lifts to an element $\bar{y}_i \in N_0$. Since N is holonomic, it is algebraic (Proposition 4.2.1). It follows that each y_i satisfies some equation $\varphi_N^{d_i}(y_i) + c_{1,i} \varphi_N^{d_i-1}(y_i) + \dots + c_{d_i,i} y_i = 0$ in $N \simeq N_0^{1/p^\infty}$, so that $\varphi_{N_0}^{d_i+e}(\bar{y}_i) + c_{1,i}^{p^e} \varphi_{N_0}^{d_i-1+e}(\bar{y}_i) + \dots + c_{d_i,i}^{p^e} \varphi_{N_0}(\bar{y}_i) = 0$ in N_0 for $e \gg 0$. In particular, the elements $\{\varphi_{N_0}^a(\bar{y}_i)\}$ generate a Frobenius submodule $N'_0 \subseteq N_0$ which is finitely generated as an R -module. Then N_0/N'_0 is finitely presented as

an R -module. Using evident isomorphism $\text{coker}(u) \simeq (N_0/N'_0)^{1/p^\infty}$, we conclude that $\text{coker}(u)$ is holonomic. \square

Our next goal is to establish a converse to Proposition 4.2.1, which asserts that every algebraic Frobenius module can be “built from” holonomic Frobenius modules (Theorem 4.2.9). First, we need some general facts about algebraicity.

Proposition 4.2.4. *Let R be an \mathbf{F}_p -algebra. Then $\text{Mod}_R^{\text{alg}}$ is a localizing subcategory of $\text{Mod}_R^{\text{perf}}$. That is:*

- (a) *Given a short exact sequence $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ of perfect Frobenius modules over R , M is algebraic if and only if M' and M'' are algebraic.*
- (b) *The collection of algebraic Frobenius modules is closed under (possibly infinite) direct sums.*

Proof. The “only if” direction of (a) follows immediately from the definitions. To prove the reverse direction, suppose we are given an exact sequence $0 \rightarrow M' \xrightarrow{\alpha} M \rightarrow M'' \rightarrow 0$ where M' and M'' are algebraic. Let x be an element of M . Since M'' is algebraic, we deduce that there is an equation of the form $\varphi_M^m(x) + a_1\varphi_M^{m-1}(x) + \cdots + a_mx = \alpha(y)$ for some $a_i \in R$ and some $y \in M'$. Since M' is algebraic, we obtain an equation of the form $\varphi_{M'}^n(y) + b_1\varphi_{M'}^{n-1}(y) + \cdots + b_ny = 0$ for some $b_j \in R$. It follows that

$$\sum_{0 \leq i \leq m, 0 \leq j \leq n} a_i b_j^i \varphi_M^{i+j}(x) = 0$$

with the convention that $a_0 = b_0 = 1$. Allowing x to vary, we deduce that M is algebraic.

To prove (b), we observe that the general case immediately reduces to the case of a finite direct sum, which follows from (a). \square

Proposition 4.2.5. *Let R be a commutative \mathbf{F}_p -algebra, let M be a perfect Frobenius module over R , and let $x \in M$ be an element. The following conditions are equivalent:*

- (1) *There exists a map of Frobenius modules $f : M' \rightarrow M$, where M' is holonomic, and an element $x' \in M'$ satisfying $f(x') = x$.*
- (2) *There exists an algebraic Frobenius submodule $M_0 \subseteq M$ which contains x .*

(3) *The element x satisfies an equation*

$$\varphi_M^n(x) + a_1\varphi_M^{n-1}(x) + \cdots + a_nx$$

for some coefficients $a_1, \dots, a_n \in R$.

Proof. We first show that (1) implies (2). Let $f : M' \rightarrow M$ be a morphism of Frobenius modules. If M' is holonomic, then it is also algebraic (Proposition 4.2.1). Consequently, if there exists an element $x' \in M'$ satisfying $f(x') = x$, then x belongs to the submodule $\text{im}(f) \subseteq M$, which is algebraic by virtue of Proposition 4.2.4.

The implication (2) \Rightarrow (3) is immediate from the definitions. We will complete the proof by showing that (3) \Rightarrow (1). Assume that x satisfies an equation $\varphi_M^n(x) + a_1\varphi_M^{n-1}(x) + \cdots + a_nx = 0$. Let N denote the free R -module on a basis $\{y_0, \dots, y_{n-1}\}$, which we regard as a Frobenius module over R by setting

$$\varphi_N(y_i) = \begin{cases} y_{i+1} & \text{if } i < n-1 \\ -a_1y_{n-1} - a_2y_{n-2} - \cdots - a_ny_0 & \text{if } i = n-1. \end{cases}$$

Then the construction $y_i \mapsto \varphi_M^i(x)$ determines a map of Frobenius modules $f_0 : N \rightarrow M$. Since M is perfect, we can extend f_0 to a map $f : N^{1/p^\infty} \rightarrow M$. It follows immediately from the construction that N^{1/p^∞} is holonomic and that x belongs to the image of f . \square

Corollary 4.2.6. *Let R be a commutative \mathbf{F}_p -algebra and let M be an algebraic Frobenius module over R . Then there exists an epimorphism of Frobenius modules $\bigoplus M_\alpha \rightarrow M$, where each M_α is holonomic.*

Corollary 4.2.7. *Let R be a commutative \mathbf{F}_p -algebra and let M be a perfect Frobenius module over R . Then there exists a largest Frobenius submodule $M_0 \subseteq M$ which is algebraic. Moreover, an element $x \in M$ belongs to M_0 if and only if it satisfies an equation*

$$\varphi_M^n(x) + a_1\varphi_M^{n-1}(x) + \cdots + a_nx$$

for some coefficients $a_1, \dots, a_n \in R$.

Proof. Let M_0 be the sum of all algebraic Frobenius submodules of M . It follows from Proposition 4.2.4 that M_0 is also algebraic, so that each element $x \in M_0$ satisfies an equation $\varphi_M^n(x) + a_1\varphi_M^{n-1}(x) + \cdots + a_nx$ for some coefficients $a_1, \dots, a_n \in R$. The reverse implication follows from Proposition 4.2.5. \square

Corollary 4.2.8. *Let $f : A \rightarrow B$ be a homomorphism of commutative \mathbf{F}_p -algebras and let M be an algebraic Frobenius module over A . Then $f^\circ(M)$ is an algebraic Frobenius module over B .*

Proof. Applying Corollary 4.2.7, we deduce that there is a largest algebraic submodule $N \subseteq f^\circ(M)$, and that N contains the image of the map

$$M \rightarrow f^\circ(M) \simeq (B \otimes_A M)^{1/p^\infty}.$$

Since N is a B -submodule of $f^\circ(M)$ which is stable under the automorphism $\varphi_{f^\circ(M)}^{-1}$, it follows that $N = f^\circ(M)$. \square

Theorem 4.2.9. *Let R be a commutative \mathbf{F}_p -algebra. Then the inclusion functor $\mathrm{Mod}_R^{\mathrm{hol}} \hookrightarrow \mathrm{Mod}_R^{\mathrm{alg}}$ extends to an equivalence of categories $\mathrm{Ind}(\mathrm{Mod}_R^{\mathrm{hol}}) \simeq \mathrm{Mod}_R^{\mathrm{alg}}$.*

Proof. It follows from Proposition 4.1.5 that the inclusion $\mathrm{Mod}_R^{\mathrm{hol}} \hookrightarrow \mathrm{Mod}_R^{\mathrm{perf}}$ extends to a fully faithful embedding $\iota : \mathrm{Ind}(\mathrm{Mod}_R^{\mathrm{hol}}) \rightarrow \mathrm{Mod}_R^{\mathrm{perf}}$. Since every holonomic Frobenius module is algebraic (Proposition 4.2.1) and the collection of algebraic Frobenius modules is closed under filtered colimits, the essential image of ι is contained in the full subcategory $\mathrm{Mod}_R^{\mathrm{alg}} \subseteq \mathrm{Mod}_R^{\mathrm{perf}}$. To complete the proof, it will suffice to verify the reverse inclusion. Let M be an algebraic Frobenius module; we wish to show that M can be written as a filtered colimit $\varinjlim M_\alpha$, where each M_α is holonomic. Using Corollary 4.2.6, we can choose an epimorphism $\rho : \bigoplus_{\alpha \in I} M_\alpha \rightarrow M$ for some set I , where each M_α is holonomic. The kernel $\ker(\rho)$ is then algebraic (Proposition 4.2.4), so we can apply Corollary 4.2.6 again to choose an epimorphism $\rho' : \bigoplus_{\beta \in J} M'_\beta \rightarrow \ker(\rho)$, where each M'_β is holonomic. We can identify ρ' with a system of maps $\{\rho'_{\alpha,\beta} : M'_\beta \rightarrow M_\alpha\}_{\alpha \in I, \beta \in J}$. Using Proposition 4.1.5, we see that for each $\beta \in J$ there are only finitely many $\alpha \in I$ for which $\rho'_{\alpha,\beta}$ is nonzero. It follows that we can write M as a filtered colimit of Frobenius modules of the form $\mathrm{coker}(\bigoplus_{\beta \in J_0} M'_\beta \rightarrow \bigoplus_{\alpha \in I_0} M_\alpha)$ where $I_0 \subseteq I$ and $J_0 \subseteq J$ are finite. Each of these Frobenius modules is holonomic by virtue of Corollary 4.2.2. \square

Let $f : A \rightarrow B$ be a morphism of commutative \mathbf{F}_p -algebras. In general, the restriction of scalars functor $f_* : \mathrm{Mod}_B^{\mathrm{Fr}} \rightarrow \mathrm{Mod}_A^{\mathrm{Fr}}$ does not preserve holonomicity. However, we do have the following:

Proposition 4.2.10. *Let $f : A \rightarrow B$ be an \mathbf{F}_p -algebra homomorphism which is finite and of finite presentation. Then the restriction of scalars functor $\mathrm{Mod}_B^{\mathrm{Fr}} \rightarrow \mathrm{Mod}_A^{\mathrm{Fr}}$ carries $\mathrm{Mod}_B^{\mathrm{hol}}$ into $\mathrm{Mod}_A^{\mathrm{hol}}$ and $\mathrm{Mod}_B^{\mathrm{alg}}$ into $\mathrm{Mod}_A^{\mathrm{alg}}$.*

Proof. Because restriction of scalars commutes with filtered colimits, it will suffice to show that if $M \in \text{Mod}_B^{\text{Fr}}$ is a holonomic Frobenius module over B , then it is also a holonomic Frobenius module over A (Theorem 4.2.9). Write $M = M_0^{1/p^\infty}$ for some $M_0 \in \text{Mod}_B^{\text{Fr}}$ which is finitely presented as a B -module. We now complete the proof by observing that our assumption on f guarantees that M_0 is also finitely presented as an A -module. \square

Warning 4.2.11. The finite presentation hypothesis in Proposition 4.2.10 cannot be relaxed to finite generation. For example, take $A = \mathbf{F}_p[x_1, x_2, x_3, \dots]$ to be a polynomial ring on countably many generators, and let $B = A/(x_1, x_2, x_3, \dots) = \mathbf{F}_p$ be its residue field at the origin. Then B is holonomic when regarded as a Frobenius module over itself, but not when regarded as a Frobenius module over A ; one can see this directly, but a quick proof is provided by Theorem 4.4.4 (note that $\text{Spec}(B)$ is not a constructible subset of $\text{Spec}(A)$).

4.3. The Noetherian Case

Recall that an object X of an abelian category \mathcal{A} is said to be *Noetherian* if the collection of subobjects of X satisfies the ascending chain condition. A Grothendieck abelian category \mathcal{A} is said to be *locally Noetherian* if every object of \mathcal{A} can be written as a union of Noetherian subobjects.

Proposition 4.3.1. *Let R be a Noetherian \mathbf{F}_p -algebra. Then the abelian category $\text{Mod}_R^{\text{alg}}$ is locally Noetherian. Moreover, an object of $\text{Mod}_R^{\text{alg}}$ is Noetherian if and only if it is holonomic.*

Proof. We first show that every holonomic R -module M is a Noetherian object of the abelian category $\text{Mod}_R^{\text{perf}}$. Write $M = M_0^{1/p^\infty}$ for some $M_0 \in \text{Mod}_R^{\text{Fr}}$ which is finitely generated as an R -module. Replacing M_0 by its image in M , we can assume without loss of generality that M_0 is a submodule of M . For any subobject $M' \subseteq M$ in the abelian category $\text{Mod}_R^{\text{alg}}$, let $M'_0 = M' \cap M_0$. Note that for any $x \in M'$, we have $\varphi_M^n(x) \in M_0$ for $n \gg 0$. It follows that $M' = \{x \in M : (\exists n)[\varphi_M^n(x) \in M'_0]\} \simeq M'_0^{1/p^\infty}$. Consequently, the construction $M' \mapsto M'_0$ determines an monomorphism from the partially ordered set of subobjects of M (in the abelian category $\text{Mod}_R^{\text{perf}}$) to the partially ordered set of subobjects of M_0 (in the abelian category Mod_R). Since M_0 is a Noetherian R -module, the latter partially ordered set satisfies the ascending chain condition, so the former does as well.

Now suppose that M is an arbitrary algebraic Frobenius module over R . Using Corollary 4.2.6, we can choose an epimorphism of Frobenius modules $\bigoplus_{\alpha \in I} M_\alpha \rightarrow M$, where each M_α is holonomic. For every finite subset $I_0 \subseteq I$,

let M_{I_0} denote the image of the composite map

$$\bigoplus_{\alpha \in I_0} M_\alpha \hookrightarrow \bigoplus_{\alpha \in I} M_\alpha \rightarrow M.$$

Then $M = \bigcup_{I_0} M_{I_0}$, and the first part of the proof shows that each M_{I_0} is Noetherian. This proves that the category $\text{Mod}_R^{\text{alg}}$ is locally Noetherian. It follows that every Noetherian object of $\text{Mod}_R^{\text{alg}}$ is compact, and therefore arises as a direct summand of a holonomic Frobenius module by virtue of Theorem 4.2.9. Since the collection of holonomic Frobenius modules is closed under passage to direct summands, it follows that $\text{Mod}_R^{\text{hol}}$ is precisely the collection of Noetherian objects of $\text{Mod}_R^{\text{alg}}$. \square

Remark 4.3.2. Let R be a Noetherian \mathbf{F}_p -algebra and let M be a holonomic Frobenius module over R . Then, for every integer k , the construction $N \mapsto \text{Ext}_{R[F]}^k(M, N)$ commutes with filtered direct limits when restricted to *perfect* Frobenius modules over R . This follows from the exact sequence of Remark 4.1.4 (together with the fact that the construction $N \mapsto \text{Ext}_R^k(M_0, N)$ commutes with filtered colimits, whenever M_0 is a finitely generated R -module).

Corollary 4.3.3. *Let R be a commutative \mathbf{F}_p -algebra. Then $\text{Mod}_R^{\text{hol}}$ is an abelian subcategory of $\text{Mod}_R^{\text{perf}}$ which is closed under the formation of kernels, cokernels, and extensions.*

Proof. Closure under the formation of cokernels was established in Corollary 4.2.2. We next show that it is closed under kernels. Let $u : M \rightarrow N$ be a morphism of holonomic Frobenius modules over R ; we wish to show that the kernel $\ker(u)$ is holonomic. Using a direct limit argument, we can write $u = f^\circ(u_0)$ for some map $f : R_0 \rightarrow R$ where R_0 is a finitely generated \mathbf{F}_p -algebra and some $u_0 : M_0 \rightarrow N_0$ in $\text{Mod}_{R_0}^{\text{hol}}$. Since R_0 is Noetherian, Proposition 4.3.1 guarantees that M_0 is a Noetherian object of $\text{Mod}_{R_0}^{\text{hol}}$. It follows that $\ker(u_0) \subseteq M_0$ is also a Noetherian object of $\text{Mod}_{R_0}^{\text{hol}}$. Corollary 3.5.2 supplies an isomorphism $\ker(u) \simeq f^\circ \ker(u_0)$, so that $\ker(u)$ is a holonomic Frobenius module over R by virtue of Proposition 4.1.2.

We now prove closure under extensions. Suppose that we are given a short exact sequence of Frobenius modules over R ,

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0,$$

where M' and M'' are holonomic; we wish to show that M is also holonomic. Using Proposition 4.1.3, we can write $M' = f^\circ M'_0$ and $M'' = f^\circ M''_0$, where

$f : R_0 \hookrightarrow R$ is the inclusion of a finitely generated subring and the Frobenius modules M'_0 and M''_0 are holonomic over R_0 . The preceding exact sequence is then classified by an element

$$\eta \in \text{Ext}_{R[F]}^1(M'', M') \simeq \text{Ext}_{R_0[F]}^1(M''_0, R \otimes_{R_0} M'_0).$$

Applying Remark 4.3.2, we can arrange (after enlarging R_0 if necessary) that η can be lifted to an element $\eta_0 \in \text{Ext}_{R_0[F]}^1(M''_0, M'_0)$, which classifies a short exact sequence

$$0 \rightarrow M'_0 \rightarrow M_0 \rightarrow M''_0 \rightarrow 0,$$

of Frobenius modules over R_0 . Since R_0 is Noetherian, we can regard M'_0 and M''_0 as Noetherian objects of the category $\text{Mod}_{R_0}^{\text{perf}}$ (Proposition 4.3.1). It follows that M_0 is also a Noetherian object of the abelian category $\text{Mod}_{R_0}^{\text{perf}}$, and is therefore a holonomic Frobenius module over R_0 (Proposition 4.3.1). Applying Proposition 4.1.2, we deduce that $M \simeq f^\diamond M_0$ is a holonomic Frobenius module over R . \square

4.4. The Support of a Holonomic Frobenius Module

Definition 4.4.1. Let R be a commutative \mathbf{F}_p -algebra and let M be a perfect Frobenius module over R . For each point $x \in \text{Spec}(R)$, we let $\kappa(x)$ denote the residue field of R at x , and we let $f_x : R \rightarrow \kappa(x)$ denote the canonical map. We let $\text{supp}(M)$ denote the set $\{x \in \text{Spec}(R) : f_x^\diamond(M) \neq 0\}$. We will refer to $\text{supp}(M)$ as the *support* of M .

Remark 4.4.2. Let R be an \mathbf{F}_p -algebra, let $M \in \text{Mod}_R^{\text{perf}}$, and let x be a point of $\text{Spec}(R)$, corresponding to a prime ideal $\mathfrak{p} \subseteq R$. The following conditions are equivalent:

- (1) The point x belongs to the support of M .
- (2) There exists a field κ and an R -algebra homomorphism $f : R \rightarrow \kappa$ such that $\ker(f) = \mathfrak{p}$ and $f^\diamond M \neq 0$.
- (3) For every field κ and every map $f : R \rightarrow \kappa$ with $\ker(f) = \mathfrak{p}$, we have $f^\diamond M \neq 0$.

To see this, we note that any map $f : R \rightarrow \kappa$ with $\ker(f) = \mathfrak{p}$ factors uniquely as a composition $R \xrightarrow{f_x} \kappa(x) \xrightarrow{\iota} \kappa$, so that we have a canonical isomorphism

$$f^\diamond M \simeq \kappa^{1/p^\infty} \otimes_{\kappa(x)^{1/p^\infty}} f_x^\diamond(M).$$

Remark 4.4.3. Let $f : A \rightarrow B$ be an \mathbf{F}_p -algebra homomorphism and let $M \in \text{Mod}_A^{\text{perf}}$. Then $\text{supp}(f^\diamond M)$ is the inverse image of $\text{supp}(M)$ under the map $\text{Spec}(B) \rightarrow \text{Spec}(A)$ determined by f (this follows immediately from Remark 4.4.2).

The support $\text{supp}(M)$ of Definition 4.4.1 is well-behaved when M is holonomic:

Theorem 4.4.4. *Let R be a commutative \mathbf{F}_p -algebra and let M be a holonomic Frobenius module over R . Then $\text{supp}(M)$ is a constructible subset of $\text{Spec}(R)$.*

Proof. Using a direct limit argument, we can choose a finitely generated subring $R' \subseteq R$ and an isomorphism $M \simeq (R \otimes_{R'} M')^{1/p^\infty}$ for some $M' \in \text{Mod}_{R'}^{\text{hol}}$. Replacing R by R' and M by M' , we can reduce to the case where R is Noetherian. By general nonsense, it will suffice to prove the following:

- (a) If $x \in \text{Spec}(R)$ belongs to the support of M , then there exists an open subset $U \subseteq \overline{\{x\}}$ which is contained in $\text{supp}(M)$.
- (b) If $x \in \text{Spec}(R)$ does not belong to the support M , then there exists an open subset $U \subseteq \overline{\{x\}}$ which is disjoint from $\text{supp}(M)$.

Let $\mathfrak{p} \subseteq R$ be the prime ideal determined by the point x . Using Remark 4.4.3, we can replace R by R/\mathfrak{p} and thereby reduce to the case where R is an integral domain and x is the generic point of $\text{Spec}(R)$.

Write $M = M_0^{1/p^\infty}$, where M_0 is finitely generated as an R -module. Let K denote the fraction field of R . Then $V = K^{1/p^\infty} \otimes_R M_0$ is a finite-dimensional vector space over K^{1/p^∞} , equipped with a Frobenius-semilinear endomorphism $\varphi_V : V \rightarrow V$. Then $\bigcup_n \ker(\varphi_V^n)$ is a K^{1/p^∞} -subspace of V , which admits a basis $\{v_i\}_{1 \leq i \leq k}$. Replacing R by a localization $R[t^{-1}]$ for some nonzero element $t \in R$, we can assume that each v_i can be lifted to an element of $R^{1/p^\infty} \otimes_R M_0$, and therefore also to an element $\bar{v}_i \in R' \otimes_R M_0$ for some subalgebra $R' \subseteq R^{1/p^\infty}$ which is finitely generated over R . Note that the inclusion $R \hookrightarrow R'$ induces a homeomorphism $\text{Spec}(R') \rightarrow \text{Spec}(R)$. We may therefore replace R by R' (and M_0 by $R' \otimes_R M_0$) and thereby reduce to the case where each \bar{v}_i belongs to M_0 . Replacing R by a localization if necessary, we may further assume that each \bar{v}_i is annihilated by some power of φ_{M_0} . Then the set $\{\bar{v}_i\}_{1 \leq i \leq k}$ generates a Frobenius submodule $M'_0 \subseteq M_0$ whose perfection vanishes. We may therefore replace M_0 by M_0/M'_0 and thereby reduce to the case where the map φ_V is injective.

Let us identify φ_{M_0} with an R -linear map $\beta : M_0^{(1)} \rightarrow M_0$, where $M_0^{(1)}$ is obtained from M_0 by extension of scalars along the Frobenius map $\varphi_R :$

$R \rightarrow R$. Note that the induced map $\beta_{K^{1/p^\infty}} : K^{1/p^\infty} \otimes_R M_0^{(1)} \rightarrow K^{1/p^\infty} \otimes_R M_0$ can be identified with φ_V and is therefore a monomorphism. Since the domain and codomain of $\beta_{K^{1/p^\infty}}$ are vector spaces of the same dimension over K^{1/p^∞} , it follows that $\beta_{K^{1/p^\infty}}$ is an isomorphism. Replacing R by a localization $R[t^{-1}]$ if necessary, we can assume that M_0 is a free module of finite rank r and that β is an isomorphism. In this case, it is easy to see that $\text{supp}(M) = \begin{cases} \text{Spec}(R) & \text{if } r > 0 \\ \emptyset & \text{if } r = 0, \end{cases}$ from which assertions (a) and (b) follow immediately. \square

5. Compactly Supported Direct Images

Let $f : A \rightarrow B$ be a morphism of commutative rings. Then the direct image functor $f_* : \text{Shv}_{\text{ét}}(\text{Spec}(B), \mathbf{F}_p) \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(A), \mathbf{F}_p)$ admits a left adjoint, which we denote by $f^* : \text{Shv}_{\text{ét}}(\text{Spec}(A), \mathbf{F}_p) \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(B), \mathbf{F}_p)$ and refer to as *pullback along f* . In the special case where f is étale, the pullback functor can be described concretely by the formula $(f^* \mathcal{F})(B') = \mathcal{F}(B')$. In particular, f^* preserves inverse limits. It follows (either by the adjoint functor theorem, or by direct construction) that the functor f^* admits a further left adjoint, which we denote by $f_! : \text{Shv}_{\text{ét}}(\text{Spec}(B), \Lambda) \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(A), \Lambda)$ and refer to as the *compactly supported direct image functor*.

Now suppose that $f : A \rightarrow B$ is a homomorphism of commutative \mathbf{F}_p -algebras. Under the Riemann-Hilbert correspondence of Theorem 1.0.2, the pullback functor $f^* : \text{Shv}_{\text{ét}}(\text{Spec}(A), \mathbf{F}_p) \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(B), \mathbf{F}_p)$ on étale sheaves corresponds to the extension of scalars functor $f^\diamond : \text{Mod}_A^{\text{alg}} \rightarrow \text{Mod}_B^{\text{alg}}$ on algebraic Frobenius modules (see Proposition 6.2.2). One consequence of this is that, if the morphism f is étale, the functor $f^\diamond : \text{Mod}_A^{\text{alg}} \rightarrow \text{Mod}_B^{\text{alg}}$ must also admit a left adjoint. Our goal in this section is to give a direct proof of this statement, which does not appeal to the Riemann-Hilbert correspondence (in fact, the work of this section will be needed in §6 to *construct* the Riemann-Hilbert functor).

We begin in §5.1 by introducing a notion of *compactly supported direct image* in the setting of Frobenius modules (Definition 5.1.2). From the definition, it will be immediately clear that if $f : A \rightarrow B$ is an étale morphism of commutative \mathbf{F}_p -algebras, then the formation of compactly supported direct images supplies a *partially defined* functor $f_! : \text{Mod}_A^{\text{alg}} \rightarrow \text{Mod}_B^{\text{alg}}$. Our main result, which we prove in §5.4, is that this functor is actually total: that is, compactly supported direct images of algebraic modules always exist (Theorem 5.4.1). The strategy of proof is to use the structure theory of étale

morphisms to reduce to the case where B is a localization $A[t^{-1}]$, which we handle in §5.2. In this case, compactly supported direct images of holonomic Frobenius modules admit a very simple characterization (Proposition 5.2.2) which makes them easy to construct explicitly. In §5.3, we apply this characterization to prove an analogue of Kashiwara’s theorem for Frobenius modules: the datum of a holonomic Frobenius module over a quotient ring $R/(t)$ is equivalent to the datum of a holonomic Frobenius module over R whose support is contained in the vanishing locus of t (Theorem 5.3.1).

5.1. Definitions

We begin by introducing some terminology.

Definition 5.1.1. Let $f : A \rightarrow B$ be an étale morphism of \mathbf{F}_p -algebras. Let M be a perfect Frobenius module over B and let \overline{M} be a perfect Frobenius module over A . We will say that a morphism $u : M \rightarrow f^\diamond \overline{M}$ exhibits \overline{M} as a weak compactly supported direct image of M if, for every perfect Frobenius module N over A , the composite map

$$\mathrm{Hom}_{A[F]}(\overline{M}, N) \rightarrow \mathrm{Hom}_{B[F]}(f^\diamond \overline{M}, f^\diamond N) \xrightarrow{\circ u} \mathrm{Hom}_{B[F]}(M, f^\diamond N)$$

is a bijection.

Let $f : A \rightarrow B$ be an étale morphism of \mathbf{F}_p -algebras and let M be a perfect Frobenius module over B . It follows immediately from the definition that if there exists a morphism $u : M \rightarrow f^\diamond \overline{M}$ which exhibits \overline{M} as a weak compactly supported direct image of M , then the Frobenius module \overline{M} (and the morphism u) are determined up to canonical isomorphism. In general, such a module need not exist. The main result of this section (Theorem 5.4.1) asserts that every algebraic Frobenius module M over B admits a weak compactly supported direct image \overline{M} . Moreover, we will have the following additional properties:

- (a) The module \overline{M} is also algebraic (as a Frobenius module over A).
- (b) For every perfect Frobenius module N over A , the canonical map

$$\mathrm{Ext}_{A[F]}^n(\overline{M}, N) \rightarrow \mathrm{Ext}_{B[F]}^n(M, f^\diamond N)$$

is an isomorphism for all integers n , rather than merely for $n = 0$.

- (c) The Frobenius module \overline{M} remains a weak compactly supported direct image of M after any extension of scalars. More precisely, for any

pushout diagram of commutative rings

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow g' & & \downarrow g \\ A' & \xrightarrow{f'} & B', \end{array}$$

the induced map $g'^{\diamond}(u) : g'^{\diamond}(M) \rightarrow g'^{\diamond}(f^{\diamond}(\overline{M})) \simeq f'^{\diamond}(g^{\diamond}(\overline{M}))$ exhibits $g^{\diamond}(\overline{M})$ as a weak compactly supported direct image of $g'^{\diamond}(M)$.

Our proof for the existence of weak compactly supported direct images will proceed by a somewhat complicated induction on the structure of the étale morphism $f : A \rightarrow B$. In order to carry out the details, it will be important to strengthen our inductive hypothesis: that is, we need to show not only that weak compactly supported direct images exist, but also that they have the properties listed above. For this reason, it will be convenient to introduce a more complicated version of Definition 5.1.1 which incorporates properties (a), (b), and (c) automatically.

Definition 5.1.2. Let $f : A \rightarrow B$ be an étale morphism of \mathbf{F}_p -algebras. Suppose we are given algebraic Frobenius modules $M \in \text{Mod}_B^{\text{alg}}$ and $\overline{M} \in \text{Mod}_A^{\text{alg}}$. We will say that a morphism $u : M \rightarrow f^{\diamond}\overline{M}$ exhibits \overline{M} as a compactly supported direct image of M if the following condition is satisfied:

(*) For every pushout diagram of commutative rings

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow g' & & \downarrow g \\ A' & \xrightarrow{f'} & B' \end{array}$$

and every object $N \in \text{Mod}_{A'}^{\text{perf}}$, the composite map

$$\begin{aligned} \text{Ext}_{A'[F]}^*(g'^{\diamond}\overline{M}, N) &\rightarrow \text{Ext}_{B'[F]}^*(f'^{\diamond}g'^{\diamond}\overline{M}, f'^{\diamond}N) \\ &\simeq \text{Ext}_{B'[F]}^*(g^{\diamond}f^{\diamond}\overline{M}, f'^{\diamond}N) \\ &\rightarrow \text{Ext}_{B'[F]}^*(g^{\diamond}M, f'^{\diamond}N). \end{aligned}$$

is an isomorphism.

Remark 5.1.3. Let $f : A \rightarrow B$ and $u : M \rightarrow f^{\diamond}\overline{M}$ be as in Definition 5.1.2. If f exhibits \overline{M} as a compactly supported direct image of M , then it also

exhibits \overline{M} as a weak compactly supported direct image of M . In fact, the converse holds as well (assuming that M is algebraic): this follows from the uniqueness of weak compactly supported direct images, once we have shown that compactly supported direct images exist (Theorem 5.4.1).

Notation 5.1.4. Let $f : A \rightarrow B$ be an étale morphism of \mathbf{F}_p -algebras and let $M \in \text{Mod}_B^{\text{alg}}$. If there exists an object $\overline{M} \in \text{Mod}_A^{\text{alg}}$ and a morphism $u : M \rightarrow f^\circ \overline{M}$ which exhibits \overline{M} as a compactly supported direct image of M , then we will denote \overline{M} by $f_! M$. In this case, we will say that $f_! M$ exists. Note that, in this event, the Frobenius module $f_! M$ depends functorially on M .

5.2. Extension by Zero

Our next goal is to prove the existence of compactly supported direct images in the case of an elementary open immersion

$$\text{Spec}(A[t^{-1}]) \hookrightarrow \text{Spec}(A)$$

(Proposition 5.2.4). In this case, Definition 5.1.2 can be formulated more simply, at least for holonomic Frobenius modules.

Definition 5.2.1. Let A be a commutative \mathbf{F}_p -algebra containing an element t and let M be a holonomic Frobenius module over $A[t^{-1}]$. An *extension by zero of M* is a holonomic Frobenius module \overline{M} over A such that $\overline{M}[t^{-1}]$ is isomorphic to M and $(\overline{M}/t\overline{M})^{1/p^\infty} \simeq 0$.

Proposition 5.2.2. *Let A be an \mathbf{F}_p -algebra containing an element t , and let $f : A \rightarrow A[t^{-1}]$ be the canonical map. Suppose we are given holonomic Frobenius modules $M \in \text{Mod}_{A[t^{-1}]}^{\text{hol}}$ and $\overline{M} \in \text{Mod}_A^{\text{hol}}$ together with a map $u : M \rightarrow f^\circ \overline{M} \simeq \overline{M}[t^{-1}]$. The following conditions are equivalent:*

- (a) *The morphism u exhibits \overline{M} as a compactly supported direct image of M , in the sense of Definition 5.1.2.*
- (b) *The morphism u exhibits \overline{M} as an extension by zero of M : that is, u is an isomorphism and the Frobenius module $(\overline{M}/t\overline{M})^{1/p^\infty}$ vanishes.*

The proof of Proposition 5.2.2 will require an elementary fact from commutative algebra:

Lemma 5.2.3. *Let M and N be modules over a commutative ring A , and let γ be an element of $\text{Ext}_A^n(M, N)$ for some $n \geq 0$. Suppose that M is Noetherian. If γ is annihilated by some power of an element $t \in A$, then there exists $d \geq 0$ such that the image of γ vanishes in $\text{Ext}_A^n(t^d M, N)$.*

Proof. Let $M_0 \subseteq M$ be the submodule consisting of those elements which are annihilated by some power of t . Since M is Noetherian, we can choose an integer $k \gg 0$ such that M_0 is annihilated by t^k . For each $d \geq k$, the kernel of the surjection $M \xrightarrow{t^d} t^d M$ is annihilated by t^k , so there exists a dotted arrow as indicated in the diagram

$$\begin{array}{ccc} M & \xrightarrow{t^k} & M \\ \downarrow t^d & \nearrow & \downarrow t^{d-k} \\ t^d M & \longrightarrow & M. \end{array}$$

It follows that the restriction map $\text{Ext}_A^n(M, N) \rightarrow \text{Ext}_A^n(t^d M, N)$ factors through the map $t^{d-k} : \text{Ext}_A^n(M, N) \rightarrow \text{Ext}_A^n(M, N)$. It therefore suffices to choose d large enough that $t^{d-k}\gamma = 0$. \square

Proof of Proposition 5.2.2. Assume first that (a) is satisfied. Applying condition (*) of Definition 5.1.2 in the case $A' = A[t^{-1}]$, we deduce that u is an isomorphism. Applying condition (*) of Definition 5.1.2 in the case $A' = A/(t)$, we deduce that $(\overline{M}/t\overline{M})^{1/p^\infty} \simeq 0$.

We now prove the converse. Assume that u is an isomorphism and that $(\overline{M}/t\overline{M})^{1/p^\infty} \simeq 0$, and suppose we are given a pushout diagram of \mathbf{F}_p -algebras

$$\begin{array}{ccc} A & \xrightarrow{f} & A[t^{-1}] \\ \downarrow g' & & \downarrow g \\ A' & \xrightarrow{f'} & A'[t^{-1}] \end{array}$$

and an object $N \in \text{Mod}_{A'}^{\text{perf}}$. To verify condition (*) of Definition 5.1.2, it will suffice to show that the canonical map

$$\theta : \text{Ext}_{A'[F]}^*(g'^\diamond \overline{M}, N) \rightarrow \text{Ext}_{A'[t^{-1}][F]}^*(f'^\diamond g'^\diamond \overline{M}, f'^\diamond N)$$

is an isomorphism. Replacing A by A' and \overline{M} by $g'^\diamond \overline{M}$, we can reduce to the case $A = A'$. Let $Q \in \text{Mod}_A^{\text{perf}}$ denote the image of the unit map $N \rightarrow f^\diamond N \simeq N[t^{-1}]$, so that we have short exact sequences

$$0 \rightarrow K \rightarrow N \rightarrow Q \rightarrow 0 \quad 0 \rightarrow Q \rightarrow N[t^{-1}] \rightarrow K' \rightarrow 0.$$

In order to show that the composite map

$$\text{Ext}_{A[F]}^*(\overline{M}, N) \rightarrow \text{Ext}_{A[F]}^*(\overline{M}, Q) \rightarrow \text{Ext}_{A[F]}^*(\overline{M}, N[t^{-1}])$$

is an isomorphism, it will suffice to show that the groups $\text{Ext}_{A[F]}^*(\overline{M}, K)$ and $\text{Ext}_{A[F]}^*(\overline{M}, K')$ vanish. This is a special case of the following:

- (*) If $\overline{M} \in \text{Mod}_A^{\text{hol}}$ satisfies $(\overline{M}/t\overline{M})^{1/p^\infty} \simeq 0$ and $N \in \text{Mod}_A^{\text{perf}}$ satisfies $N[t^{-1}] \simeq 0$, then $\text{Ext}_{A[F]}^*(\overline{M}, N)$ vanishes.

Write $\overline{M} = T^{1/p^\infty}$ for some Frobenius module $T \in \text{Mod}_A^{\text{Fr}}$ in A which is finitely presented as an A -module. Using our assumption that $(\overline{M}/t\overline{M})^{1/p^\infty} \simeq 0$, we deduce that $\varphi_T^m T \subseteq tT$ for $m \gg 0$. By a direct limit argument, we can assume that $T \simeq A \otimes_{A_0} T_0$ for some finitely generated subalgebra $A_0 \subseteq A$ which contains t and some $T_0 \in \text{Mod}_{A_0}^{\text{Fr}}$ which is finitely presented as an A_0 -module, and that T_0 satisfies $\varphi_{T_0}^m T_0 \subseteq tT_0$. Using Corollary 3.5.3, we can replace A by A_0 and T by T_0 , and thereby reduce to proving (*) in the special case where A is Noetherian.

Using Remark 4.1.4, we obtain a long exact sequence

$$\text{Ext}_{A[F]}^*(\overline{M}, N) \rightarrow \text{Ext}_A^*(T, N) \xrightarrow{\text{id}-U} \text{Ext}_A^*(T, N) \rightarrow \text{Ext}_{A[F]}^{*+1}(\overline{M}, N),$$

where U is the endomorphism of $\text{Ext}_A^*(T, N)$ given by $U(\gamma) = \varphi_N^{-1} \circ \gamma \circ \varphi_T$. To prove (*), it will suffice to show that the map $1 - U$ is an isomorphism of $\text{Ext}_A^*(T, N)$ with itself. In fact, we will show that U is locally nilpotent (so that $1 - U$ has an inverse given by the formal infinite sum $1 + U + U^2 + \dots$).

Fix an element $\gamma \in \text{Ext}_A^k(T, N)$; we wish to show that $U^m(\gamma) \simeq 0$ for $m \gg 0$. Since A is Noetherian and T is a finitely generated A -module, the construction $S \mapsto \text{Ext}_A^k(T, S)$ commutes with filtered colimits. In particular, there exists a finitely generated A -submodule $N_0 \subseteq N$ such that γ can be lifted to an element $\gamma_0 \in \text{Ext}_A^k(T, N_0)$. Using our assumption that $N[t^{-1}] \simeq 0$, we deduce that N_0 is annihilated by t^c for $c \gg 0$. It follows that the image of γ_0 in $\text{Ext}_A^k(t^c T, N_0)$ vanishes for $c' \gg c$ (Lemma 5.2.3). We now observe that for $m \gg 0$, the map φ_T^m factors through $t^c T$, so that $\gamma \circ \varphi_T^m = 0$ and therefore $U^m(\gamma) = 0$ as desired. \square

Proposition 5.2.4. *Let A be an \mathbf{F}_p -algebra containing an element t , and let $f : A \rightarrow A[t^{-1}]$ be the canonical map. Then every algebraic Frobenius module M over $A[t^{-1}]$ admits a compactly supported direct image along f .*

Proof. Using Theorem 4.2.9, we can reduce to the case where M is holonomic. Write $M = M_0^{1/p^\infty}$, where $M_0 \in \text{Mod}_{A[t^{-1}]}^{\text{Fr}}$ is finitely presented as an $A[t^{-1}]$ -module. Then we can choose an isomorphism $\alpha : M_0 \simeq \overline{M}_0[t^{-1}]$ for some finitely presented object $\overline{M}_0 \in \text{Mod}_A$. Then φ_{M_0} determines an A -module homomorphism $\rho : \overline{M}_0 \rightarrow \overline{M}_0[t^{-1}]^{1/p}$. Since \overline{M}_0 is finitely presented

as an A -module, we can assume that ρ factors as a composition

$$\overline{M}_0 \rightarrow \overline{M}_0^{1/p} \xrightarrow{t^n} \overline{M}_0[t^{-1}]^{1/p}$$

for some integer n . Multiplying the isomorphism α by a suitable power of t , we can arrange that $n > 0$. Set $\overline{M} = \overline{M}_0^{1/p^\infty}$. Then α induces an isomorphism $M \simeq \overline{M}[t^{-1}]$ and $\varphi_{\overline{M}}$ is locally nilpotent on $\overline{M}/t\overline{M}$, so that α exhibits \overline{M} as an extension by zero of M , which is also a compactly supported direct image of M by virtue of Proposition 5.2.2. \square

5.3. Kashiwara’s Theorem

Let X be a smooth algebraic variety over the field \mathbf{C} of complex numbers, and let $Y \subseteq X$ be a smooth subvariety of X . A theorem of Kashiwara (see [9, §1.6]) asserts that the category of algebraic \mathcal{D} -modules on Y is equivalent to the category of algebraic \mathcal{D} -modules on X which vanish over the open set $X - Y$. In this section, we prove the following analogue for (holonomic) Frobenius modules:

Theorem 5.3.1. *Let M be a holonomic Frobenius module over a commutative \mathbf{F}_p -algebra A , and let $I \subseteq A$ be an ideal. The following conditions are equivalent:*

- *The support $\text{supp}(M)$ is contained in the vanishing locus $\text{Spec}(A/I) \subseteq \text{Spec}(A)$.*
- *The submodule $IM \subseteq M$ vanishes: that is, M has the structure of a Frobenius module over A/I .*

Remark 5.3.2. If the equivalent conditions of Theorem 5.3.1 are satisfied, then M is also holonomic when regarded as a Frobenius module over A/I . Conversely, if the ideal I is finitely generated, then any holonomic Frobenius module over A/I is a holonomic Frobenius module over A which satisfies the conditions of Theorem 5.3.1 (see Proposition 4.2.10). In other words, the category $\text{Mod}_{A/I}^{\text{hol}}$ can be identified with the full subcategory of $\text{Mod}_A^{\text{hol}}$ spanned by objects set-theoretically supported on $\text{Spec}(A/I) \subseteq \text{Spec}(A)$. Beware that this is generally not true if I is not finitely generated (Warning 4.2.11).

We begin by treating the following special case of Theorem 5.3.1 (which is the only case we will actually need):

Proposition 5.3.3. *Let M be a holonomic Frobenius module over a commutative \mathbf{F}_p -algebra A . Then $M \simeq 0$ if and only if the support $\text{supp}(M)$ is empty.*

Proof. The “only if” direction is obvious. To prove the converse, let us assume that $\text{supp}(M) = \emptyset$; we wish to prove that $M \simeq 0$. Using Proposition 4.1.3, we can choose a finitely generated subring $A' \subseteq A$ and an equivalence $M \simeq (A \otimes_{A'} M')^{1/p^\infty}$ for some $M' \in \text{Mod}_{A'}^{\text{hol}}$. Set $Z = \text{supp}(M') \subseteq \text{Spec}(A')$. Then Z is constructible (Theorem 4.4.4). Using Remark 4.4.3, we deduce that the image of the map $\text{Spec}(A) \rightarrow \text{Spec}(A')$ is disjoint from Z . Enlarging A' if necessary, we can arrange that $Z = \emptyset$. We may therefore replace A by A' and M by M' , and thereby reduce to the case where A is Noetherian.

Proceeding by Noetherian induction, we may assume that for every nonzero ideal $I \subseteq A$, we have $(M/IM)^{1/p^\infty} \simeq 0$. We may assume that $A \neq 0$ (otherwise, there is nothing to prove). If A is not reduced, then taking I to be the nilradical of A we deduce that $M = M^{1/p^\infty} \simeq (M/IM)^{1/p^\infty} \simeq 0$. We may therefore assume that A is reduced. Using Proposition 5.2.2, we deduce that M is the compactly supported direct image of $M[x^{-1}]$ for every nonzero element $x \in A$. It will therefore suffice to show that we can choose a nonzero element $x \in A$ such that $M[x^{-1}] \simeq 0$. Since A is reduced and Noetherian, we can choose a non-zero divisor $t \in A$ such that $A[t^{-1}]$ is an integral domain. Replacing A by $A[t^{-1}]$, we can assume that A is an integral domain. Write $M = M_0^{1/p^\infty}$ for some $M_0 \in \text{Mod}_A^{\text{Fr}}$ which is finitely presented as an A -module. Let K be the fraction field of A . Since the support $\text{supp}(M)$ does not contain the generic point of $\text{Spec}(A)$, the Frobenius module $(K \otimes_A M_0)^{1/p^\infty}$ vanishes. Using the finite generation of M_0 , we conclude that the Frobenius endomorphism of $K \otimes_A M_0$ is nilpotent. It follows that there exists a nonzero element $x \in A$ for which the Frobenius map $\varphi_{M_0[x^{-1}]}$ is nilpotent, so that $M[x^{-1}] \simeq M_0[x^{-1}]^{1/p^\infty} \simeq 0$ as desired. \square

Proof of Theorem 5.3.1. Let M be a holonomic Frobenius module over a commutative \mathbf{F}_p -algebra A and let $I \subseteq A$ be an ideal. It follows immediately from the definitions that if M is annihilated by I , then the support $\text{supp}(M)$ is contained in the vanishing locus of I . Conversely, suppose that $\text{supp}(M) \subseteq \text{Spec}(A/I)$; we wish to show that M is annihilated by each element $x \in I$. Note that the inclusion $\text{supp}(M) \subseteq \text{Spec}(A/I)$ guarantees that the support of $M[x^{-1}]$ is empty, where we regard $M[x^{-1}]$ as a holonomic Frobenius module over $A[x^{-1}]$. Applying Proposition 5.3.3, we conclude that $M[x^{-1}] \simeq 0$.

Choose an isomorphism $M \simeq M_0^{1/p^\infty}$, where M_0 is a Frobenius module over A which is finitely presented as an A -module. Let N denote the image of the map $M_0 \rightarrow M$, so that $N \subseteq M$ is a Frobenius submodule which is finitely generated over A . The vanishing of $M[x^{-1}]$ guarantees that $x^k N = 0$ for some $k \gg 0$. Applying φ_M^{-n} , we conclude that $x^{\frac{k}{p^n}} N^{1/p^n} = 0$ for all $n \geq 0$. As $M = \varinjlim_n N^{1/p^n}$, it follows that $x^{\frac{k}{p^n}} M = 0$ for all $n \geq 0$ (here we regard

M as a Frobenius module over the perfection A^{1/p^∞}). For $n \gg 0$, this implies $xM = 0$, as desired. \square

5.4. Existence of Compactly Supported Direct Images

Our goal in this section is to prove the following:

Theorem 5.4.1. *Let $f : A \rightarrow B$ be an étale morphism of \mathbf{F}_p -algebras. Then, for every object $M \in \text{Mod}_B^{\text{alg}}$, there exists a compactly supported direct image $f_!M \in \text{Mod}_A^{\text{alg}}$ (see Notation 5.1.4). Moreover, the functor $f_! : \text{Mod}_B^{\text{alg}} \rightarrow \text{Mod}_A^{\text{alg}}$ is exact.*

Remark 5.4.2. In the situation of Theorem 5.4.1, the right exactness of the functor $f_! : \text{Mod}_B^{\text{alg}} \rightarrow \text{Mod}_A^{\text{alg}}$ is automatic (since $f_!$ is left adjoint to the functor $f^\diamond : \text{Mod}_A^{\text{alg}} \rightarrow \text{Mod}_B^{\text{alg}}$, which is exact by virtue of Corollary 3.5.2). Moreover, since the functor $f^\diamond : \text{Mod}_A^{\text{alg}} \rightarrow \text{Mod}_B^{\text{alg}}$ preserves filtered colimits, the functor $f_!$ preserves compact objects: that is, it carries $\text{Mod}_B^{\text{hol}}$ into $\text{Mod}_A^{\text{hol}}$ (see Theorem 4.2.9).

The proof of Theorem 5.4.1 will require some preliminaries. We begin with some elementary remarks, whose proofs follow immediately from our definitions.

Lemma 5.4.3. *Suppose we are given a pushout diagram of \mathbf{F}_p -algebras*

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow g' & & \downarrow g \\ A' & \xrightarrow{f'} & B' \end{array}$$

where f is étale. If $\overline{M} \in \text{Mod}_B^{\text{alg}}$ and $u : M \rightarrow f^\diamond \overline{M}$ is a morphism in $\text{Mod}_A^{\text{alg}}$ which exhibits \overline{M} as a compactly supported direct image of M , then the induced map $g^\diamond M \rightarrow g^\diamond f^\diamond \overline{M} \simeq f'^\diamond g'^\diamond \overline{M}$ exhibits $g'^\diamond \overline{M}$ as a compactly supported direct image of $g^\diamond M$.

In particular, if $f_!M$ exists, then $f'_!(g^\diamond M)$ exists (and is canonically isomorphic to $g'^\diamond(f_!M)$).

Lemma 5.4.4. *Let $f : A \rightarrow B$ and $g : B \rightarrow C$ be étale morphisms of \mathbf{F}_p -algebras. Suppose we are given an objects $M_C \in \text{Mod}_C^{\text{alg}}$, a morphism $u : M_B \rightarrow g^\diamond M_C$ in $\text{Mod}_B^{\text{alg}}$, and a morphism $v : M_A \rightarrow f^\diamond M_B$ in $\text{Mod}_A^{\text{alg}}$. Assume that u exhibits M_B as a compactly supported direct image of M_C .*

Then v exhibits M_A as a compactly supported direct image of M_B if and only if the composite map

$$M_A \xrightarrow{v} f^\diamond M_B \xrightarrow{g^\diamond(u)} g^\diamond f^\diamond M_C$$

exhibits M_A as a compactly supported direct image of M_C .

In particular, if $g_!M$ exists, then $f_!(g_!M)$ exists if and only if $(g \circ f)_!M$ exists (and, in this case, they are canonically isomorphic).

Lemma 5.4.5. *Let $f : A \rightarrow B$ be an étale morphism of \mathbf{F}_p -algebras and suppose we are given an exact sequence $0 \rightarrow M' \xrightarrow{u} M \rightarrow M'' \rightarrow 0$ in the abelian category $\text{Mod}_B^{\text{alg}}$. Suppose that $f_!M'$ and $f_!M$ exist, and that the canonical map $f_!(u) : f_!M' \rightarrow f_!M$ is a monomorphism. Then $f_!M''$ exists, and is given by $\text{coker}(f_!(u))$.*

Lemma 5.4.6. *Let $f : A \rightarrow B$ be a faithfully flat étale morphism of \mathbf{F}_p -algebras and let $M \in \text{Mod}_A^{\text{perf}}$. Then M is algebraic if and only if $f^\diamond M$ is algebraic.*

Proof. The “only if” direction follows from Corollary 4.2.8. Conversely, suppose that $f^\diamond M$ is algebraic. Choose an element $x \in M$. For each $n \geq 0$, let $M(n)$ denote the A -submodule of M generated by the elements $\{\varphi_M^k(x)\}_{k < n}$, so we have inclusions of A -submodules

$$\{0\} = M(0) \subseteq M(1) \subseteq M(2) \subseteq \cdots \subseteq M.$$

Using Corollary 3.4.7, we can identify $f^\diamond M$ with $B \otimes_A M$, so that each $B \otimes_A M(n)$ can be identified with the B -submodule $M'(n) \subseteq f^\diamond M$ generated by $\{\varphi_{B \otimes_A M}^k(1 \otimes x)\}_{k < n}$. Since $f^\diamond M$ is algebraic, there exists an integer n such that $M'(n) = M'(n+1)$. The faithful flatness of B over A then guarantees that $M(n) = M(n+1)$, so that x satisfies an equation of the form $\varphi_M^n(x) + a_1 \varphi_M^{n-1}(x) + \cdots + a_n x = 0$. \square

Lemma 5.4.7. *Suppose we are given a pushout square of étale morphisms between \mathbf{F}_p -algebras*

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow & & \downarrow g \\ A' & \xrightarrow{f'} & B' \end{array},$$

where the vertical maps are faithfully flat. Let $M \in \text{Mod}_B^{\text{alg}}$. If $f'_!(g^\diamond M)$ exists, then $f_!M$ exists.

Proof. Use faithfully flat descent together with Lemma 5.4.6. \square

Lemma 5.4.8. *Let A be an \mathbf{F}_p -algebra containing an element t , let $f : A \rightarrow A[t^{-1}]$ be the canonical map, and suppose we are given objects $M \in \text{Mod}_{A[t^{-1}]}^{\text{alg}}$ and a morphism $u : f_!M \rightarrow N$ in $\text{Mod}_A^{\text{alg}}$. If $f^\diamond(u)$ is a monomorphism, then u is a monomorphism.*

Proof. Set $K = \ker(u)$. Then $f_!M/K$ is algebraic (Proposition 4.2.4), so Corollary 3.5.2 implies that the map $(K/tK)^{1/p^\infty} \rightarrow ((f_!M)/t(f_!M))^{1/p^\infty}$ is a monomorphism. Invoking Proposition 5.2.2, we deduce that the natural map $f_!f^\diamond K \rightarrow K$ is an equivalence. Since $f^\diamond K \simeq \ker(f^\diamond u) \simeq 0$, we conclude that $K \simeq 0$ so that u is a monomorphism as desired. \square

Proof of Theorem 5.4.1. Let us say that an étale ring homomorphism $f : A \rightarrow B$ is *good* if the functor $f_! : \text{Mod}_B^{\text{alg}} \rightarrow \text{Mod}_A^{\text{alg}}$ is well-defined and exact. Our proof now proceeds in several steps:

- (i) Every localization $f : A \rightarrow A[t^{-1}]$ is good. The existence of $f_!$ follows from Proposition 5.2.4, and the exactness of $f_!$ follows from Remark 5.4.2 and Lemma 5.4.8.
- (ii) Let $f : A \rightarrow B$ and $g : B \rightarrow C$ be étale ring homomorphisms. If f and g are good, then $(g \circ f) : A \rightarrow C$ is good. This follows immediately from Lemma 5.4.4.
- (iii) Let $f : A \rightarrow B$ be an étale \mathbf{F}_p -algebra homomorphism and suppose we are given elements $t_0, t_1 \in B$ which generate the unit ideal. Set $B_0 = B[t_0^{-1}]$, $B_1 = B[t_1^{-1}]$, and $B_{01} = B[t_0^{-1}, t_1^{-1}]$. If the induced maps $f_0 : A \rightarrow B_0$ and $f_1 : A \rightarrow B_1$ are good, then $f_!$ exists. To prove this, choose any object $M \in \text{Mod}_B^{\text{alg}}$, and define

$$M_0 = M[t_0^{-1}] \in \text{Mod}_{B_0}^{\text{alg}} \quad M_1 = M[t_1^{-1}] \in \text{Mod}_{B_1}^{\text{alg}}$$

$$M_{01} = M[t_0^{-1}, t_1^{-1}] \in \text{Mod}_{B_{01}}^{\text{alg}}.$$

We have a commutative diagram

$$\begin{array}{ccc} B & \xrightarrow{g_0} & B_0 \\ \downarrow g_1 & \searrow g_{01} & \downarrow h \\ B_1 & \longrightarrow & B_{01} \end{array}$$

which yields a short exact sequence

$$0 \rightarrow g_{01}!M_{01} \xrightarrow{u} g_0!M_0 \oplus g_1!M_1 \rightarrow M \rightarrow 0$$

in $\text{Mod}_B^{\text{alg}}$. Using our assumptions that f_0 and f_1 are good (which also implies that the induced map $f_{01} : A \rightarrow B_{01}$ is good, using (ii)

and (iii) together with Lemma 5.4.4, we deduce that $f_!(g_{01!}M_{01})$ and $f_!(g_{0!}M_0 \oplus g_{1!}M_1)$ exist. By virtue of Lemma 5.4.5, to prove the existence of $f_!M$, it will suffice to show that $f_!u$ is a monomorphism. In fact, we claim that the composite map

$$f_!g_{01!}M_{01} \xrightarrow{f_!u} f_!(g_{0!}M_0 \oplus g_{1!}M_1) \rightarrow f_!g_{0!}M_0$$

is a monomorphism. Using Lemma 5.4.4 and our assumption that $f_{0!}$ is exact, we are reduced to showing that the map $h_!M_{01} \rightarrow M_0$ is a monomorphism in $\text{Mod}_{B_0}^{\text{alg}}$, which is a special case of Lemma 5.4.8.

- (iv) Let $f : A \rightarrow B$ be as in (iii). Then f is good. To prove this, we must show that for every short exact sequence $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ in $\text{Mod}_B^{\text{alg}}$, the induced map $f_!M' \rightarrow f_!M$ is a monomorphism. Define M_0 , M_1 , and M_{01} as above, and define M'_0 , M'_1 , M'_{01} , M''_0 , M''_1 , and M''_{01} similarly. We then have a diagram of short exact sequences

$$\begin{array}{ccccccccc} 0 & \longrightarrow & f_{01!}M'_{01} & \longrightarrow & f_{0!}M'_0 \oplus f_{1!}M'_1 & \longrightarrow & f_!M' & \longrightarrow & 0 \\ & & \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma & & \\ 0 & \longrightarrow & f_{01!}M_{01} & \longrightarrow & f_{0!}M_0 \oplus f_{1!}M_1 & \longrightarrow & f_!M & \longrightarrow & 0 \end{array}$$

Using the exactness of the functors $f_{0!}$, $f_{1!}$, and $f_{01!}$, the snake lemma yields an exact sequence

$$0 \rightarrow \ker(\gamma) \rightarrow f_{01!}M''_{01} \xrightarrow{\rho} f_{0!}M''_0 \oplus f_{1!}M''_1.$$

It will therefore suffice to show that ρ is a monomorphism, which was established in the proof of (iii).

- (v) Let $f : A \rightarrow B$ be an étale \mathbf{F}_p -algebra homomorphism, and suppose that the unit ideal of B is generated by elements $\{t_i \in B\}_{1 \leq i \leq n}$ for which each of the induced maps $A \rightarrow B[t_i^{-1}]$ is good. Then f is good. This follows from (iii) and (iv), using induction on n .
- (vi) Suppose we are given a pushout square of étale maps

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow & & \downarrow g \\ A' & \xrightarrow{f'} & B' \end{array},$$

where the vertical maps are faithfully flat. If f' is good, then f is good. This follows from Lemma 5.4.7.

We now wish to prove that every étale morphism $f : A \rightarrow B$ is good. For each point $x \in \text{Spec}(A)$, let $\kappa(x)$ denote the residue field of A at x and let $d(x)$ denote the dimension $\dim_{\kappa(x)}(\kappa(x) \otimes_A B)$. Set $d(B) = \sup_{x \in \text{Spec}(A)} d(x)$. We proceed by induction on $d(B)$. If $d = 0$, then $B \simeq 0$ and there is nothing to prove. To carry out the inductive step, we note that since f is étale, the induced map $\text{Spec}(B) \rightarrow \text{Spec}(A)$ has open image. The complement of this image can be written as the vanishing locus of an ideal $I = (a_1, \dots, a_n) \subseteq A$. Then I generates the unit ideal of B . By virtue of (v), to prove that f is good, it will suffice to show that each of the composite maps $A \rightarrow A[a_i^{-1}] \xrightarrow{f_i} B[a_i^{-1}]$ is good. Using (i) and (ii), we are reduced to showing that the maps $f_i : A[a_i^{-1}] \rightarrow B[a_i^{-1}]$ are good. Replacing f by f_i , we may reduce to the case where f is faithfully flat. Form a pushout square

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow f & & \downarrow \\ B & \xrightarrow{f'} & B \otimes_A B. \end{array}$$

By virtue of (vi), we can replace f by f' and thereby reduce to the case where B splits as a direct product $A \times B'$. We then have $d(B') = d(B) - 1 < d(B)$, so our inductive hypothesis implies that the map $A \rightarrow B'$ is good. From this, we immediately deduce that f is also good. \square

6. The Riemann-Hilbert Functor

Let R be a commutative \mathbf{F}_p -algebra. In §2, we defined the solution functor

$$\text{Sol} : \text{Mod}_R^{\text{Fr}} \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$$

and asserted that it becomes an equivalence of categories when restricted to the category $\text{Mod}_R^{\text{alg}} \subseteq \text{Mod}_R^{\text{Fr}}$ of algebraic Frobenius modules (Theorem 2.4.3). We will prove this by defining a functor $\text{RH} : \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p) \rightarrow \text{Mod}_R^{\text{alg}}$, which we will refer to as the *Riemann-Hilbert functor*, and then showing that it is an inverse to the solution functor. Our goal in this section is to construct the Riemann-Hilbert functor and to establish its basic properties. Our principal results can be summarized as follows:

- (a) When restricted to perfect Frobenius modules, the solution functor $\text{Sol} : \text{Mod}_R^{\text{perf}} \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ admits a left adjoint (Theorem 6.1.1). We will take this left adjoint as a definition of the Riemann-Hilbert functor.

- (b) The Riemann-Hilbert functor $\mathrm{RH} : \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(R), \mathbf{F}_p) \rightarrow \mathrm{Mod}_R^{\mathrm{perf}}$ depends functorially on R , in the sense that it is compatible with pullback (Proposition 6.2.2). We also show that it is compatible with compactly supported direct images along étale morphisms (Proposition 6.2.3), and direct images along morphisms which are finite and of finite presentation (Theorem 6.5.1).
- (c) The Riemann-Hilbert functor $\mathrm{RH} : \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(R), \mathbf{F}_p) \rightarrow \mathrm{Mod}_R^{\mathrm{perf}}$ carries constructible étale sheaves on $\mathrm{Spec}(R)$ to holonomic Frobenius modules over R (Theorem 6.3.1).
- (d) The Riemann-Hilbert functor $\mathrm{RH} : \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(R), \mathbf{F}_p) \rightarrow \mathrm{Mod}_R^{\mathrm{perf}}$ is exact (Proposition 6.4.1).

In §7, we will apply these results to show that RH is an inverse of the solution functor (once we restrict our attention to algebraic Frobenius modules), and thereby obtain a proof of Theorem 1.0.2.

6.1. Existence of the Riemann-Hilbert Functor

Our starting point is the following:

Theorem 6.1.1. *Let R be a commutative \mathbf{F}_p -algebra. Then the solution functor $\mathrm{Sol} : \mathrm{Mod}_R^{\mathrm{perf}} \rightarrow \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(R), \mathbf{F}_p)$ admits a left adjoint*

$$\mathrm{RH} : \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(R), \mathbf{F}_p) \rightarrow \mathrm{Mod}_R^{\mathrm{perf}}.$$

Moreover, for every p -torsion étale sheaf $\mathcal{F} \in \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(R), \mathbf{F}_p)$, the Frobenius module $\mathrm{RH}(\mathcal{F})$ is algebraic.

Warning 6.1.2. In the statement of Theorem 6.1.1, it is important to restrict the solution functor Sol to the category of *perfect* Frobenius modules. The defining property of the Riemann-Hilbert functor RH is that we have bijections

$$\mathrm{Hom}_{\mathbf{F}_p}(\mathcal{F}, \mathrm{Sol}(M)) \simeq \mathrm{Hom}_{R[F]}(\mathrm{RH}(\mathcal{F}), M)$$

for $\mathcal{F} \in \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(R), \mathbf{F}_p)$ and M a perfect Frobenius module over R . One does not generally have such a bijection when M is not perfect.

To prove Theorem 6.1.1, it will be convenient to introduce a temporary bit of terminology. Let R be a commutative \mathbf{F}_p -algebra, and suppose we are given a p -torsion étale sheaf \mathcal{F} . A *Riemann-Hilbert associate* of \mathcal{F} is an

object of $\text{Mod}_R^{\text{perf}}$ which corepresents the functor

$$\text{Mod}_R^{\text{perf}} \rightarrow \text{Set} \quad M \mapsto \text{Hom}_{\underline{\mathbf{F}}_p}(\mathcal{F}, \text{Sol}(M)).$$

If \mathcal{F} is a perfect Frobenius module over R which admits a Riemann-Hilbert associate, we will denote that associate by $\text{RH}(\mathcal{F})$; note that it is well-defined up to unique isomorphism and depends functorially on \mathcal{F} . Theorem 6.1.1 can then be reformulated as the statement that every étale sheaf $\mathcal{F} \in \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ admits an algebraic Riemann-Hilbert associate. The proof of this assertion is based on three simple observations:

Proposition 6.1.3. *Let R be a commutative \mathbf{F}_p -algebra. Then the perfection R^{perf} is a Riemann-Hilbert associate of the constant sheaf $\underline{\mathbf{F}}_p$.*

Proof. For every perfect Frobenius module M over R , we have canonical bijections

$$\begin{aligned} \text{Hom}_{R[F]}(R^{\text{perf}}, M) &\simeq \text{Hom}_{R[F]}(R, M) \\ &\simeq \{x \in M : \varphi_M(x) = x\} \\ &\simeq \text{Sol}(M)(R) \\ &\simeq \text{Hom}_{\underline{\mathbf{F}}_p}(\underline{\mathbf{F}}_p, \text{Sol}(M)). \end{aligned}$$

□

Proposition 6.1.4. *Let R be a commutative \mathbf{F}_p -algebra, and suppose we are given some diagram of étale sheaves $\{\mathcal{F}_\alpha\}$ having a colimit $\mathcal{F} = \varinjlim \mathcal{F}_\alpha$ in the category $\text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$. Suppose that each \mathcal{F}_α admits a Riemann-Hilbert associate $\text{RH}(\mathcal{F}_\alpha)$. Then \mathcal{F} admits a Riemann-Hilbert associate, given by $\varinjlim \text{RH}(\mathcal{F}_\alpha)$ (where the colimit is formed in the category $\text{Mod}_R^{\text{perf}}$).*

Proof. For any perfect Frobenius module M , we have canonical bijections

$$\begin{aligned} \text{Hom}_{R[F]}(\varinjlim_{\alpha} \text{RH}(\mathcal{F}_\alpha), M) &\simeq \varprojlim_{\alpha} \text{Hom}_{R[F]}(\text{RH}(\mathcal{F}_\alpha), M) \\ &\simeq \varprojlim_{\alpha} \text{Hom}_{\underline{\mathbf{F}}_p}(\mathcal{F}_\alpha, \text{Sol}(M)) \\ &\simeq \text{Hom}_{\underline{\mathbf{F}}_p}(\mathcal{F}, \text{Sol}(M)). \end{aligned}$$

□

Proposition 6.1.5. *Let $f : A \rightarrow B$ be an étale morphism of commutative \mathbf{F}_p -algebras and let \mathcal{F} be a p -torsion étale sheaf on $\text{Spec}(B)$. Suppose that \mathcal{F} admits a Riemann-Hilbert associate $\text{RH}(\mathcal{F})$ which is algebraic. Then the*

compactly supported direct image $f_! \text{RH}(\mathcal{F})$ is a Riemann-Hilbert associate of $f_! \mathcal{F} \in \text{Shv}_{\text{ét}}(\text{Spec}(A), \mathbf{F}_p)$.

Proof. Let M be a perfect Frobenius module over A . It follows immediately from the definitions that the solution sheaf $\text{Sol}(f^\diamond M) \in \text{Shv}_{\text{ét}}(\text{Spec}(B), \mathbf{F}_p)$ can be identified with the pullback $f^* \text{Sol}(M)$. We therefore obtain canonical bijections

$$\begin{aligned} \text{Hom}_{A[F]}(f_! \text{RH}(\mathcal{F}), M) &\simeq \text{Hom}_{B[F]}(\text{RH}(\mathcal{F}), f^\diamond M) \\ &\simeq \text{Hom}_{\mathbf{F}_p}(\mathcal{F}, \text{Sol}(f^\diamond M)) \\ &\simeq \text{Hom}_{\mathbf{F}_p}(\mathcal{F}, f^* \text{Sol}(M)) \\ &\simeq \text{Hom}_{\mathbf{F}_p}(f_! \mathcal{F}, \text{Sol}(M)). \end{aligned}$$

□

Proof of Theorem 6.1.1. Let R be a commutative \mathbf{F}_p -algebra and let \mathcal{F} be a p -torsion étale sheaf on $\text{Spec}(R)$; we wish to show that \mathcal{F} admits an algebraic Riemann-Hilbert associate. For every étale ring homomorphism $j : R \rightarrow R'$ and element $\eta \in \mathcal{F}(R')$, we can identify η with a map of étale sheaves $u_\eta : j_! \mathbf{F}_p \rightarrow \mathcal{F}$. Amalgamating these, we obtain an epimorphism $u : \mathcal{F}' \rightarrow \mathcal{F}$ in the category $\text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$, where \mathcal{F}' is a direct sum of étale sheaves of the form $j_! \mathbf{F}_p$ (where j varies over étale morphisms $R \rightarrow R'$). Repeating this argument for $\ker(u)$, we can construct an exact sequence

$$\mathcal{F}'' \xrightarrow{v} \mathcal{F}' \xrightarrow{u} \mathcal{F} \rightarrow 0,$$

where \mathcal{F}'' is also a direct sum of sheaves of the form $j_! \mathbf{F}_p$. By virtue of Proposition 6.1.4, it will suffice to show that each of the sheaves $j_! \mathbf{F}_p$ admits an algebraic Riemann-Hilbert associate. Using Proposition 6.1.5, we are reduced to showing that if R' is an étale R -algebra, then the constant sheaf $\mathbf{F}_p \in \text{Shv}_{\text{ét}}(\text{Spec}(R'), \mathbf{F}_p)$ admits an algebraic Riemann-Hilbert associate. This follows from Proposition 6.1.3. □

6.2. Functoriality

We now study the behavior of the Riemann-Hilbert functor

$$\text{RH} : \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p) \rightarrow \text{Mod}_R^{\text{perf}}$$

as the commutative \mathbf{F}_p -algebra R varies. We begin with a simple observation:

Proposition 6.2.1. *Let $f : A \rightarrow B$ be a homomorphism of commutative \mathbf{F}_p -algebras. Then the diagram of categories*

$$\begin{array}{ccc} \mathrm{Mod}_B^{\mathrm{perf}} & \xrightarrow{f_*} & \mathrm{Mod}_A^{\mathrm{perf}} \\ \downarrow \mathrm{Sol} & & \downarrow \mathrm{Sol} \\ \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(B), \mathbf{F}_p) & \xrightarrow{f_*} & \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(A), \mathbf{F}_p) \end{array}$$

commutes up to canonical isomorphism.

Proof. Let M be a perfect Frobenius module over B , let A' be an étale A -algebra, and set $B' = A' \otimes_A B$. We then have canonical bijections

$$\begin{aligned} (f_* \mathrm{Sol}(M))(A') &\simeq \mathrm{Sol}(M)(B') \\ &\simeq \{x \in B' \otimes_B M : \varphi_{B' \otimes_B M}(x) = x\} \\ &\simeq \{x \in A' \otimes_A M : \varphi_{A' \otimes_A M}(x) = x\} \\ &\simeq \mathrm{Sol}(f_* M)(A'). \end{aligned}$$

□

Proposition 6.2.2. *Let $f : A \rightarrow B$ be an \mathbf{F}_p -algebra homomorphism. Then the diagram of categories*

$$\begin{array}{ccc} \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(A), \mathbf{F}_p) & \xrightarrow{f^*} & \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(B), \mathbf{F}_p) \\ \downarrow \mathrm{RH} & & \downarrow \mathrm{RH} \\ \mathrm{Mod}_A^{\mathrm{perf}} & \xrightarrow{f^\diamond} & \mathrm{Mod}_B^{\mathrm{perf}} \end{array}$$

commutes up to canonical isomorphism.

Proof. This follows immediately from Proposition 6.2.1 by passing to left adjoints. □

In the situation of Proposition 6.2.2, the vertical maps carry étale sheaves to algebraic Frobenius modules, so we also have a commutative diagram

$$\begin{array}{ccc} \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(A), \mathbf{F}_p) & \xrightarrow{f^*} & \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(B), \mathbf{F}_p) \\ \downarrow \mathrm{RH} & & \downarrow \mathrm{RH} \\ \mathrm{Mod}_A^{\mathrm{alg}} & \xrightarrow{f^\diamond} & \mathrm{Mod}_B^{\mathrm{alg}}. \end{array}$$

In the case where $f : A \rightarrow B$ is étale, the horizontal maps in this diagram admit left adjoints (Theorem 5.4.1). We therefore obtain a natural transformation $f_! \circ \mathrm{RH} \rightarrow \mathrm{RH} \circ f_!$ in the category of functors from $\mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(B), \mathbf{F}_p)$ to $\mathrm{Mod}_A^{\mathrm{alg}}$.

Proposition 6.2.3. *Let $f : A \rightarrow B$ be an étale morphism of \mathbf{F}_p -algebras. Then the Beck-Chevalley transformation $f_! \circ \mathrm{RH} \rightarrow \mathrm{RH} \circ f_!$ described above is an isomorphism. In other words, the diagram of categories*

$$\begin{array}{ccc} \mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(A), \mathbf{F}_p) & \xleftarrow{f_!} & \mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(B), \mathbf{F}_p) \\ \downarrow \mathrm{RH} & & \downarrow \mathrm{RH} \\ \mathrm{Mod}_A^{\mathrm{alg}} & \xleftarrow{f_!} & \mathrm{Mod}_B^{\mathrm{alg}} \end{array}$$

commutes up to canonical isomorphism.

Proof. This is a translation of Proposition 6.1.5 (or, more precisely, of its proof). \square

Remark 6.2.4. We can also formulate Proposition 6.2.3 in terms of solution sheaves: it follows from the commutativity of the diagram

$$\begin{array}{ccc} \mathrm{Mod}_A^{\mathrm{perf}} & \xrightarrow{f^\circ} & \mathrm{Mod}_B^{\mathrm{perf}} \\ \downarrow \mathrm{Sol} & & \downarrow \mathrm{Sol} \\ \mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(A), \mathbf{F}_p) & \xrightarrow{f^*} & \mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(B), \mathbf{F}_p) \end{array}$$

when $f : A \rightarrow B$ is an étale morphism of \mathbf{F}_p -algebras, which in turn follows immediately from the definitions (and was invoked in the proof of Proposition 6.1.5).

6.3. Constructible Sheaves

Let R be a commutative ring. Recall (see [15, Tag 05BE]) that a sheaf $\mathcal{F} \in \mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(R), \mathbf{F}_p)$ is said to be *constructible* if there is finite stratification

$$\emptyset = X_0 \subseteq X_1 \subseteq \cdots \subseteq X_n = \mathrm{Spec}(R),$$

where each open stratum $X_m - X_{m-1}$ is a constructible subset of $\mathrm{Spec}(R)$ and admits an étale surjection $U_m \rightarrow (X_m - X_{m-1})$ such that the restriction $\mathcal{F}|_{U_m}$ is isomorphic to a constant sheaf \underline{V} , for some finite-dimensional

vector space V over \mathbf{F}_p . We let $\mathrm{Shv}_{\acute{e}t}^c(\mathrm{Spec}(R), \mathbf{F}_p)$ denote the full subcategory of $\mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(R), \mathbf{F}_p)$ spanned by the constructible sheaves. Note that $\mathrm{Shv}_{\acute{e}t}^c(\mathrm{Spec}(R), \mathbf{F}_p)$ is closed under the formation of kernels and cokernels in $\mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(R), \mathbf{F}_p)$; in particular, it is an abelian category.

Theorem 6.1.1 admits the following refinement:

Theorem 6.3.1. *Let R be a commutative \mathbf{F}_p -algebra. Then the Riemann-Hilbert functor $\mathrm{RH} : \mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(R), \mathbf{F}_p) \rightarrow \mathrm{Mod}_R^{\mathrm{perf}}$ carries constructible étale sheaves to holonomic Frobenius modules over R .*

Notation 6.3.2. For every \mathbf{F}_p -algebra R , we let $\mathrm{RH}^c : \mathrm{Shv}_{\acute{e}t}^c(\mathrm{Spec}(R), \mathbf{F}_p) \rightarrow \mathrm{Mod}_R^{\mathrm{hol}}$ denote the restriction of the Riemann-Hilbert functor RH to constructible sheaves.

For the proof of Theorem 6.3.1, we will need a few standard facts about constructible sheaves, which we assert here without proof:

Proposition 6.3.3. *Let $f : A \rightarrow B$ be an étale ring homomorphism. Then the functor $f_! : \mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(B), \mathbf{F}_p) \rightarrow \mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(A), \mathbf{F}_p)$ carries constructible sheaves to constructible sheaves.*

Proof. This is contained in [15, Tag 03S8]. □

Proposition 6.3.4. *Let A be a commutative ring and let $\mathcal{F} \in \mathrm{Shv}_{\acute{e}t}^c(\mathrm{Spec}(A), \mathbf{F}_p)$. Then there exists an étale morphism $f : A \rightarrow B$ and an epimorphism $f_! \underline{\mathbf{F}}_p \rightarrow \mathcal{F}$ in the abelian category $\mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(A), \mathbf{F}_p)$.*

Proof. This follows from [15, Tag 09YT]. □

Proof of Theorem 6.3.1. Let R be a commutative \mathbf{F}_p -algebra and let \mathcal{F} be a constructible p -torsion étale sheaf on $\mathrm{Spec}(R)$. We wish to show that the Frobenius module $\mathrm{RH}(\mathcal{F})$ is holonomic. We first apply Proposition 6.3.4 to choose an epimorphism $u : \mathcal{F}' \rightarrow \mathcal{F}$, where \mathcal{F}' has the form $f_! \underline{\mathbf{F}}_p$ for some étale morphism $f : A \rightarrow B$. Then \mathcal{F}' is constructible, so $\ker(u)$ is also constructible. Applying Proposition 6.3.4 again, we can choose an epimorphism $v : \mathcal{F}'' \rightarrow \ker(u)$, where \mathcal{F}'' has the form $g_! \underline{\mathbf{F}}_p$ for some étale morphism $g : A \rightarrow C$. We then have an exact sequence

$$\mathcal{F}'' \xrightarrow{v} \mathcal{F}' \xrightarrow{u} \mathcal{F} \rightarrow 0.$$

The Riemann-Hilbert functor RH is right exact (since it is a left adjoint), so we obtain an exact sequence of Frobenius modules

$$\mathrm{RH}(\mathcal{F}'') \rightarrow \mathrm{RH}(\mathcal{F}') \rightarrow \mathrm{RH}(\mathcal{F}) \rightarrow 0.$$

Since the collection of holonomic Frobenius modules over R is closed under the formation of cokernels (Corollary 4.2.2), it will suffice to show that $\mathrm{RH}(\mathcal{F}'')$ and $\mathrm{RH}(\mathcal{F}')$ are holonomic. Using Propositions 6.2.3 and 6.1.3, we obtain isomorphisms $\mathrm{RH}(\mathcal{F}') \simeq f_! B^{\mathrm{perf}}$ and $\mathrm{RH}(\mathcal{F}'') \simeq g_! C^{\mathrm{perf}}$. The desired holonomicity now follows from Remark 5.4.2. \square

We close this section by recording (without proof) a few more elementary facts about constructible sheaves which will be needed in the proof of Theorem 1.0.2. First, we have the following duals to Propositions 6.3.3 and 6.3.4:

Proposition 6.3.5. *Let $f : A \rightarrow B$ be a ring homomorphism which is finite and of finite presentation. Then the direct image functor $f_* : \mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(B), \mathbf{F}_p) \rightarrow \mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(A), \mathbf{F}_p)$ carries constructible sheaves to constructible sheaves.*

Proof. See [7, §1, Lemma 4.11]. \square

Proposition 6.3.6. *Let A be a commutative ring and let $\mathcal{F} \in \mathrm{Shv}_{\acute{e}t}^c(\mathrm{Spec}(A), \mathbf{F}_p)$. Then there exists a ring homomorphism $f : A \rightarrow B$ which is finite and of finite presentation and a monomorphism $\mathcal{F} \rightarrow f_* \underline{\mathbf{F}}_p$ in the category $\mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(A), \mathbf{F}_p)$.*

Proof. See [7, §1, Proposition 4.12]. \square

Proposition 6.3.7. *Let R be a commutative ring. Then the inclusion functor $\mathrm{Shv}_{\acute{e}t}^c(\mathrm{Spec}(R), \mathbf{F}_p) \hookrightarrow \mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(R), \mathbf{F}_p)$ extends to an equivalence of categories $\mathrm{Ind}(\mathrm{Shv}_{\acute{e}t}^c(\mathrm{Spec}(R), \mathbf{F}_p)) \simeq \mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(R), \mathbf{F}_p)$.*

Proof. See [15, Tag 03SA]. \square

6.4. Exactness of the Riemann-Hilbert Functor

Proposition 6.4.1. *Let R be a commutative \mathbf{F}_p -algebra. Then the Riemann-Hilbert functor $\mathrm{RH} : \mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(R), \mathbf{F}_p) \rightarrow \mathrm{Mod}_R^{\mathrm{perf}}$ is exact.*

Proof. Since the Riemann-Hilbert functor RH is defined as the left adjoint to the solution functor, it is automatically right exact. It will therefore suffice to show that if $u : \mathcal{F} \rightarrow \mathcal{G}$ is a monomorphism of p -torsion étale sheaves on $\mathrm{Spec}(R)$, then the induced map $\mathrm{RH}(u) : \mathrm{RH}(\mathcal{F}) \rightarrow \mathrm{RH}(\mathcal{G})$ is also a monomorphism. Using Proposition 6.3.7, we can reduce to the case where \mathcal{F} and \mathcal{G} are constructible, so that the Frobenius modules $\mathrm{RH}(\mathcal{F})$ and $\mathrm{RH}(\mathcal{G})$ are holonomic (Theorem 6.3.1). It follows that the kernel of the map $\mathrm{RH}(u)$ is also holonomic (Corollary 4.3.3). By virtue of Proposition 5.3.3,

to show that $\text{RH}(u)$ is a monomorphism, it will suffice to show that the support $\text{supp}(\ker(\text{RH}(u)))$ is empty. Fix a point $x \in \text{Spec}(R)$; we will show that $x \notin \text{supp}(\ker(\text{RH}(u))) \subseteq \text{Spec}(R)$. Let κ be an algebraic closure of the residue field of R at the point x and let $f : R \rightarrow \kappa$ be the canonical map; we wish to show that $f^\diamond(\ker(\text{RH}(u)))$ vanishes. Since the functor f^\diamond is exact on algebraic Frobenius modules (Corollary 3.5.2) and compatible with the Riemann-Hilbert functor (Proposition 6.2.2), we have

$$f^\diamond(\ker(\text{RH}(u))) \simeq \ker(f^\diamond(\text{RH}(u))) \simeq \ker(\text{RH}(f^*(u))).$$

We can therefore replace R by κ and thereby reduce to the case where R is an algebraically closed field. In this case, the category $\text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ is equivalent to the category of vector spaces over \mathbf{F}_p . It follows that every exact sequence in the category $\text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ is split, so the exactness of the Riemann-Hilbert functor RH is automatic. \square

Corollary 6.4.2. *Let R be a commutative \mathbf{F}_p -algebra. Then the solution functor $\text{Sol} : \text{Mod}_R^{\text{perf}} \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ carries injective objects of the abelian category $\text{Mod}_R^{\text{perf}}$ to injective objects of the abelian category $\text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$.*

6.5. Comparison of Finite Direct Images

Let $f : A \rightarrow B$ be a homomorphism of commutative \mathbf{F}_p -algebras, so that Proposition 6.2.2 supplies a commutative diagram of categories

$$\begin{array}{ccc} \text{Shv}_{\text{ét}}(\text{Spec}(A), \mathbf{F}_p) & \xrightarrow{f^*} & \text{Shv}_{\text{ét}}(\text{Spec}(B), \mathbf{F}_p) \\ \downarrow \text{RH} & & \downarrow \text{RH} \\ \text{Mod}_A^{\text{perf}} & \xrightarrow{f^\diamond} & \text{Mod}_B^{\text{perf}} \end{array} .$$

Note that the horizontal maps in this diagram admit right adjoints

$$f_* : \text{Mod}_B^{\text{perf}} \rightarrow \text{Mod}_A^{\text{perf}} \quad f_* : \text{Shv}_{\text{ét}}(\text{Spec}(B), \mathbf{F}_p) \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(A), \mathbf{F}_p).$$

By general nonsense, we obtain a Beck-Chevalley transformation $\text{RH} \circ f_* \rightarrow f_* \circ \text{RH}$ in the category of functors from $\text{Shv}_{\text{ét}}(\text{Spec}(B), \mathbf{F}_p)$ to $\text{Mod}_A^{\text{perf}}$. In general, this map need not be an isomorphism: for example, if $B^{\text{perf}} = (f_* \circ \text{RH})(\mathbf{F}_p)$ is not algebraic when regarded as a Frobenius module over A , then it cannot belong to the essential image of the Riemann-Hilbert functor

$\mathrm{RH} : \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(A), \mathbf{F}_p) \rightarrow \mathrm{Mod}_A^{\mathrm{perf}}$. However, under some mild finiteness hypotheses, this phenomenon does not arise:

Theorem 6.5.1. *Let $f : A \rightarrow B$ be a morphism of commutative \mathbf{F}_p -algebras which is finite and of finite presentation. Then, for every p -torsion étale sheaf \mathcal{F} on $\mathrm{Spec}(B)$, the canonical map $\epsilon_{\mathcal{F}} : \mathrm{RH}(f_* \mathcal{F}) \rightarrow f_*(\mathrm{RH}(\mathcal{F}))$ is an isomorphism of Frobenius modules over A . Consequently, the diagram of categories*

$$\begin{array}{ccc} \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(A), \mathbf{F}_p) & \xleftarrow{f_*} & \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(B), \mathbf{F}_p) \\ \downarrow \mathrm{RH} & & \downarrow \mathrm{RH} \\ \mathrm{Mod}_A^{\mathrm{perf}} & \xleftarrow{f_*} & \mathrm{Mod}_B^{\mathrm{perf}} \end{array}$$

commutes (up to canonical isomorphism).

Remark 6.5.2. In §10, we will prove a more general version of Theorem 6.5.1, which applies in the situation of a morphism $f : X \rightarrow Y$ of \mathbf{F}_p -schemes which is proper and of finite presentation (Corollary 10.5.6).

Proof of Theorem 6.5.1. The functors

$$\mathrm{RH} : \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(A), \mathbf{F}_p) \rightarrow \mathrm{Mod}_A^{\mathrm{perf}} \quad \mathrm{RH} : \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(B), \mathbf{F}_p) \rightarrow \mathrm{Mod}_B^{\mathrm{perf}}$$

$$f_* : \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(B), \mathbf{F}_p) \rightarrow \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(A), \mathbf{F}_p) \quad f_* : \mathrm{Mod}_B^{\mathrm{perf}} \rightarrow \mathrm{Mod}_A^{\mathrm{perf}}$$

all commute with filtered colimits. Consequently, to show that the map $\epsilon_{\mathcal{F}}$ is an equivalence for every object $\mathcal{F} \in \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(B), \mathbf{F}_p)$, it will suffice to show that $\epsilon_{\mathcal{F}}$ is an equivalence when the sheaf \mathcal{F} is constructible (Proposition 6.3.7). In this case, the direct image $f_* \mathcal{F} \in \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(A), \mathbf{F}_p)$ is constructible (Proposition 6.3.5). Applying Theorem 6.3.1, we deduce that $\mathrm{RH}(\mathcal{F})$ and $\mathrm{RH}(f_* \mathcal{F})$ are holonomic. It follows from Proposition 4.2.10 that $f_* \mathrm{RH}(\mathcal{F})$ is also holonomic. Applying Corollary 4.3.3, we deduce that $\ker(\epsilon_{\mathcal{F}})$ and $\mathrm{coker}(\epsilon_{\mathcal{F}})$ are holonomic. By virtue of Propositions 5.3.3 and 4.2.4, to show that $\epsilon_{\mathcal{F}}$ is an isomorphism, it will suffice to show that $g^\circ(\epsilon_{\mathcal{F}})$ is an isomorphism for every map $g : A \rightarrow \kappa$ where κ is an algebraically closed field. Using Proposition 6.2.2 (and the fact that pushforward of étale sheaves along finite morphisms commutes with base change), we can replace A by κ and thereby reduce to the case where A is an algebraically closed field. In this case, B is a finite-dimensional algebra over κ . Writing B as a product of local rings, we can assume that B is local with residue field κ . Then the constructible sheaf $\mathcal{F} \in \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(B), \mathbf{F}_p)$ has the form \underline{V} for some finite-dimensional vector space V over \mathbf{F}_p . Choosing a basis for V , we

can reduce to the case where $V = \mathbf{F}_p$. Using Proposition 6.1.3, we see that $\epsilon_{\mathcal{F}}$ can be identified with the canonical map $A^{1/p^\infty} \rightarrow B^{1/p^\infty}$, which is an isomorphism since the radical of B is nilpotent. \square

7. The Riemann-Hilbert Correspondence

Let R be a commutative \mathbf{F}_p -algebra. Our goal in this section is to prove Theorem 1.0.2 by showing that the Riemann-Hilbert functor

$$\text{RH} : \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p) \rightarrow \text{Mod}_R^{\text{alg}}$$

is an equivalence of categories. Let us outline the strategy we will use. Our first objective (which is achieved in §7.2) is to show that the Riemann-Hilbert functor is fully faithful: that is, that the unit map $\mathcal{F} \rightarrow \text{Sol}(\text{RH}(\mathcal{F}))$ is an isomorphism for any p -torsion étale sheaf \mathcal{F} on $\text{Spec}(R)$ (see Proposition 7.2.1). One obstacle to proving this is that the solution functor $\text{Sol} : \text{Mod}_R^{\text{perf}} \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ is not exact. However, we show in §7.1 that it is *almost* exact: more precisely, it has only one derived functor, which can be explicitly described (Proposition 7.1.1).

The rest of this section is devoted to showing that every algebraic Frobenius module M over R belongs to the essential image of the Riemann-Hilbert functor. To prove this, we may assume without loss of generality that M is holonomic. In this case, we prove something stronger: the Frobenius module M can be realized as $\text{RH}(\mathcal{F})$, where \mathcal{F} is a *constructible* p -torsion étale sheaf on $\text{Spec}(R)$ (Theorem 7.4.1). In the case where R is a field, this assertion is classical; we give a proof in §7.3 for the reader's convenience (Proposition 7.3.1). The general case is treated in §7.4, using a dévissage which reduces to the case where R is a field.

7.1. Derived Solution Functors

Let R be a commutative \mathbf{F}_p -algebra. The solution functor

$$\text{Sol} : \text{Mod}_R^{\text{perf}} \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$$

of Construction 2.3.1 is left exact (Remark 2.3.4), but is usually not exact. Since the category $\text{Mod}_R^{\text{perf}}$ has enough injective objects (Remark 3.2.6), we can consider its right derived functors. For each $n \geq 0$, we let $\text{Sol}^n : \text{Mod}_R^{\text{perf}} \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ denote the n th right derived functor of Sol . These derived functors admit a simple explicit description:

Proposition 7.1.1. *Let R be a commutative \mathbf{F}_p -algebra and let M be a perfect Frobenius module over R . Then we have a canonical short exact sequence*

$$0 \rightarrow \mathrm{Sol}(M) \rightarrow \widetilde{M} \xrightarrow{\mathrm{id} - \widetilde{\varphi}_M} \widetilde{M} \rightarrow \mathrm{Sol}^1(M) \rightarrow 0$$

and the sheaves $\mathrm{Sol}^n(M)$ vanish for $n \geq 2$.

Here \widetilde{M} denotes the quasi-coherent sheaf associated to the R -module M (see Example 2.2.5). We will deduce Proposition 7.1.1 from the following:

Lemma 7.1.2. *Let R be a commutative \mathbf{F}_p -algebra. If M is an injective object of $\mathrm{Mod}_R^{\mathrm{perf}}$, then the sequence $0 \rightarrow \mathrm{Sol}(M) \rightarrow \widetilde{M} \xrightarrow{\mathrm{id} - \widetilde{\varphi}_M} \widetilde{M} \rightarrow 0$ is exact in the category of abelian presheaves on the category $\mathrm{CAlg}_R^{\acute{e}t}$.*

Proof. Choose an étale morphism $f : R \rightarrow A$. We then have a diagram of exact sequences

$$\begin{array}{ccc} 0 & & 0 \\ \downarrow & & \downarrow \\ \mathrm{Sol}(M)(A) & \xrightarrow{\sim} & \mathrm{Ext}_{A[F]}^0(A, A \otimes_R M) \\ \downarrow & & \downarrow \\ \widetilde{M} & \xrightarrow{\sim} & \mathrm{Ext}_{A[F]}^0(A[F], A \otimes_R M) \\ \downarrow \mathrm{id} - \varphi & & \downarrow \\ \widetilde{M} & \xrightarrow{\sim} & \mathrm{Ext}_{A[F]}^0(A[F], A \otimes_R M) \\ & & \downarrow \\ & & \mathrm{Ext}_{A[F]}^1(A, A \otimes_R M). \end{array}$$

To complete the proof, it will suffice to show that the group $\mathrm{Ext}_{A[F]}^1(A, A \otimes_R M) \simeq \mathrm{Ext}_{A[F]}^1(A^{1/p^\infty}, A \otimes_R M)$ vanishes. Using Theorem 5.4.1, we obtain an isomorphism

$$\mathrm{Ext}_{A[F]}^1(A^{1/p^\infty}, A \otimes_R M) \simeq \mathrm{Ext}_{R[F]}^1(f_! A^{1/p^\infty}, M),$$

where the right hand side vanishes by virtue of our assumption that M is injective. \square

Proof of Proposition 7.1.1. Let M be a perfect Frobenius module over a commutative \mathbf{F}_p -algebra R , and choose an injective resolution $0 \rightarrow M \rightarrow Q^0 \rightarrow Q^1 \rightarrow \cdots$ in the abelian category $\text{Mod}_R^{\text{perf}}$. Using Lemma 7.1.2, we obtain a short exact sequence of cochain complexes

$$0 \rightarrow \text{Sol}(Q^*) \rightarrow \tilde{Q}^* \xrightarrow{\text{id}-\varphi} \tilde{Q}^* \rightarrow 0.$$

Since the construction $N \mapsto \tilde{N}$ is exact, the chain complex \tilde{Q}^* is an acyclic resolution of \tilde{M} . The associated long exact sequence now supplies the desired isomorphisms. \square

Corollary 7.1.3. *Let R be a commutative \mathbf{F}_p -algebra. Then the functor $\text{Sol}^n : \text{Mod}_R^{\text{perf}} \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ commutes with filtered colimits for each $n \geq 0$.*

Corollary 7.1.4. *Let R be a commutative \mathbf{F}_p -algebra, let \mathcal{F} be a p -torsion étale sheaf on $\text{Spec}(R)$, and let M be a perfect Frobenius module over R . Then we have canonical short exact sequences*

$$0 \rightarrow \text{Ext}_{\underline{\mathbf{F}}_p}^n(\mathcal{F}, \text{Sol}(M)) \rightarrow \text{Ext}_{R[F]}^n(\text{RH}(\mathcal{F}), M) \rightarrow \text{Ext}_{\underline{\mathbf{F}}_p}^{n-1}(\mathcal{F}, \text{Sol}^1(M)) \rightarrow 0.$$

Proof. Since the solution functor $\text{Sol} : \text{Mod}_R^{\text{perf}} \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ carries injective objects of $\text{Mod}_R^{\text{perf}}$ to injective objects of $\text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ (Corollary 6.4.2), we have a Grothendieck spectral sequence

$$\text{Ext}_{\underline{\mathbf{F}}_p}^s(\mathcal{F}, \text{Sol}^t(M)) \Rightarrow \text{Ext}_{R[F]}^{s+t}(\text{RH}(\mathcal{F}), M).$$

The existence of the desired short exact sequences now follows from the vanishing of the groups $\text{Sol}^t(M)$ for $t \geq 2$ (Proposition 7.1.1). \square

7.2. Full Faithfulness of the Riemann-Hilbert Functor

We are now ready to prove a weak version of Theorem 1.0.2.

Proposition 7.2.1. *Let R be a commutative \mathbf{F}_p -algebra and let \mathcal{F} be a p -torsion étale sheaf on $\text{Spec}(R)$. Then the unit map $u_{\mathcal{F}} : \mathcal{F} \rightarrow \text{Sol}(\text{RH}(\mathcal{F}))$ is an isomorphism and the sheaf $\text{Sol}^1(\text{RH}(\mathcal{F}))$ vanishes.*

We first treat a special case of Proposition 7.2.1:

Lemma 7.2.2. *Let R be a commutative \mathbf{F}_p -algebra. Then the unit map $u : \underline{\mathbf{F}}_p \rightarrow \text{Sol}(\text{RH}(\underline{\mathbf{F}}_p))$ is an isomorphism and the sheaf $\text{Sol}^1(\text{RH}(\underline{\mathbf{F}}_p))$ vanishes.*

Proof. Without loss of generality, we may assume that R is perfect. Using Propositions 6.1.3 and 7.1.1, we see that Lemma 7.2.2 is equivalent to the exactness of the Artin-Schreier sequence

$$0 \rightarrow \underline{\mathbf{F}}_p \rightarrow \tilde{R} \xrightarrow{\text{id}-\varphi} \tilde{R} \rightarrow 0$$

in the category $\text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$. \square

Proof of Proposition 7.2.1. Using Corollary 7.1.3 and Proposition 6.3.7, we can reduce to the case where the sheaf \mathcal{F} is constructible. Using Proposition 6.3.6, we can choose an exact sequence of constructible sheaves

$$0 \rightarrow \mathcal{F} \rightarrow \mathcal{G} \rightarrow \mathcal{H} \rightarrow 0,$$

where $\mathcal{G} = f_* \underline{\mathbf{F}}_p$ for some \mathbf{F}_p -algebra homomorphism $f : R \rightarrow A$ which is finite and of finite presentation. Using Proposition 6.4.1 and Proposition 7.1.1, we obtain a commutative diagram

$$\begin{array}{ccc}
 0 & & 0 \\
 \downarrow & & \downarrow \\
 \mathcal{F} & \xrightarrow{u_{\mathcal{F}}} & \text{Sol}(\text{RH}(\mathcal{F})) \\
 \downarrow & & \downarrow \\
 \mathcal{G} & \xrightarrow{u_{\mathcal{G}}} & \text{Sol}(\text{RH}(\mathcal{G})) \\
 \downarrow & & \downarrow \\
 \mathcal{H} & \xrightarrow{u_{\mathcal{H}}} & \text{Sol}(\text{RH}(\mathcal{H})) \\
 \downarrow & & \downarrow \delta \\
 0 & \longrightarrow & \text{Sol}^1(\text{RH}(\mathcal{F})) \\
 & & \downarrow \\
 & & \text{Sol}^1(\text{RH}(\mathcal{G}))
 \end{array}$$

whose columns are exact. It follows from Lemma 7.2.2, Theorem 6.5.1, and Proposition 6.2.1 that $u_{\mathcal{G}}$ is an isomorphism and $\text{Sol}^1(\text{RH}(\mathcal{G}))$ vanishes. Inspecting the diagram, we deduce that $u_{\mathcal{F}}$ is a monomorphism. Applying the same argument to \mathcal{H} , we see that $u_{\mathcal{H}}$ is also monomorphism, so a diagram chase shows that $u_{\mathcal{F}}$ is an epimorphism. Applying the same argument to \mathcal{H} ,

we conclude that $u_{\mathcal{H}}$ is also an epimorphism. The commutativity of the diagram shows that $\delta \circ u_{\mathcal{H}}$ vanishes, so that $\delta = 0$. Since δ is an epimorphism, we conclude that $\text{Sol}^1(\text{RH}(\mathcal{F})) \simeq 0$. \square

It follows from Proposition 7.2.1 that the Riemann-Hilbert functor is fully faithful, even at the “derived” level:

Corollary 7.2.3. *Let R be a commutative \mathbf{F}_p -algebra. Then the Riemann-Hilbert functor $\text{RH} : \text{Shv}_{\acute{e}t}(\text{Spec}(R), \mathbf{F}_p) \rightarrow \text{Mod}_R^{\text{perf}}$ is fully faithful. Moreover, for every pair of objects $\mathcal{F}, \mathcal{G} \in \text{Shv}_{\acute{e}t}(\text{Spec}(R), \mathbf{F}_p)$, the induced map $\text{Ext}_{\mathbf{F}_p}^*(\mathcal{F}, \mathcal{G}) \rightarrow \text{Ext}_{R[F]}^*(\text{RH}(\mathcal{F}), \text{RH}(\mathcal{G}))$ is an isomorphism.*

Proof. Combine Proposition 7.2.1 with Corollary 7.1.4. \square

7.3. The Case of a Field

It follows from Corollary 7.2.3 that, for any commutative \mathbf{F}_p -algebra R , the functor

$$\text{RH}^c : \text{Shv}_{\acute{e}t}^c(\text{Spec}(R), \mathbf{F}_p) \rightarrow \text{Mod}_R^{\text{hol}}$$

of Notation 6.3.2 is fully faithful. We now show that it is an equivalence in the special case where R is a field, which is essentially a restatement of Theorem 1.0.1:

Proposition 7.3.1. *[10, Proposition 4.1.1] Let κ be a field of characteristic p . Then the functor $\text{RH}^c : \text{Shv}_{\acute{e}t}^c(\text{Spec}(\kappa), \mathbf{F}_p) \rightarrow \text{Mod}_{\kappa}^{\text{hol}}$ is an equivalence of categories.*

We begin by treating the case where κ is algebraically closed (compare [5, §III, Lemma 3.3]):

Lemma 7.3.2. *Let κ be an algebraically closed field of characteristic p . Then the functor $\text{RH}^c : \text{Shv}_{\acute{e}t}^c(\text{Spec}(\kappa), \mathbf{F}_p) \rightarrow \text{Mod}_{\kappa}^{\text{hol}}$ is an equivalence of categories.*

Proof. Using Corollary 7.2.3 and Proposition 6.4.1, we see that RH^c is a fully faithful embedding whose essential image $\mathcal{C} \subseteq \text{Mod}_{\kappa}^{\text{hol}}$ is an abelian subcategory which is closed under extensions. We wish to show that \mathcal{C} contains every object $M \in \text{Mod}_{\kappa}^{\text{hol}}$. Applying Proposition 4.3.1, we see that M is a Noetherian object of the abelian category $\text{Mod}_{\kappa}^{\text{alg}}$. Consequently, there exists a subobject $M' \subseteq M$ (in the abelian category $\text{Mod}_{\kappa}^{\text{alg}}$) which is maximal among those subobjects which belong to \mathcal{C} . It follows from the maximality of M' (and the stability of \mathcal{C} under extensions) that the quotient M/M' does not contain any nonzero subobjects which belong to \mathcal{C} . Replacing M

by M/M' , we can reduce to the case where M does not have any nonzero subobjects which belong to \mathcal{C} .

Suppose that M is nonzero. Choose a nonzero element $x \in M$. Since M is algebraic, the element x satisfies an equation

$$\varphi_M^n(x) + \lambda_1 \varphi_M^{n-1}(x) + \cdots + \lambda_n x = 0$$

for some coefficients $\lambda_1, \lambda_2, \dots, \lambda_n \in \kappa$. We may assume that x has been chosen so that n is as small as possible; this guarantees that the set $\{x, \varphi_M(x), \dots, \varphi_M^{n-1}(x)\}$ is linearly independent over κ , and therefore $\lambda_n \neq 0$. Since $x \neq 0$, we must have $n > 0$.

Note that

$$f(t) = t^{p^n} + \lambda_1^{p^{n-1}} t^{p^{n-1}} + \lambda_2^{p^{n-2}} t^{p^{n-2}} + \cdots + \lambda_n x$$

is a separable polynomial of degree $p^n > 1$, and therefore has p^n distinct roots in the field κ . Consequently, there exists a nonzero element $a \in \kappa$ such that $f(a) = 0$. Let

$$y = ax + (a^p + a\lambda_1)\varphi_M(x) + \cdots + (a^{p^{n-1}} + a^{p^{n-2}}\lambda_1^{p^{n-2}} + \cdots + a\lambda_{n-1})\varphi_M^{n-1}(x).$$

Since the elements $\{\varphi_M^i(x)\}_{0 \leq i < n}$ are linearly independent and $a \neq 0$, y is a nonzero element of M . An explicit calculation gives

$$\begin{aligned} y - \varphi_M(y) &= ax + \sum_{0 < i < n} a\lambda_i \varphi_M^i(x) + (a\lambda_n - f(a))\varphi_M^n(x) \\ &= a(x + \lambda_1 \varphi_M(x) + \cdots + \lambda_n \varphi_M^n(x)) \\ &= 0. \end{aligned}$$

It follows that y generates a nonzero Frobenius submodule of M which is isomorphic to $\kappa \simeq \mathrm{RH}(\mathbf{F}_p)$, contradicting our assumption that M does not contain any nonzero subobjects which belong to \mathcal{C} . \square

Proof of Proposition 7.3.1. Let κ be an arbitrary field of characteristic p . As in the proof of Lemma 7.3.2, we see that the functor $\mathrm{RH}^c : \mathrm{Shv}_{\mathrm{\acute{e}t}}^c(\mathrm{Spec}(\kappa), \mathbf{F}_p) \rightarrow \mathrm{Mod}_{\kappa}^{\mathrm{hol}}$ is a fully faithful embedding whose essential image $\mathcal{C} \subseteq \mathrm{Mod}_{\kappa}^{\mathrm{hol}}$ is an abelian category which is closed under extensions. We wish to show that \mathcal{C} contains every object $M \in \mathrm{Mod}_{\kappa}^{\mathrm{hol}}$. Let $\bar{\kappa}$ be an algebraic closure of κ . Lemma 7.3.2 shows that $(\bar{\kappa} \otimes_{\kappa} M)^{1/p^\infty} \in \mathrm{Mod}_{\bar{\kappa}}^{\mathrm{hol}}$ belongs to the essential image of the functor $\mathrm{RH}^c : \mathrm{Shv}_{\bar{\kappa}}^c \rightarrow \mathrm{Mod}_{\bar{\kappa}}^{\mathrm{hol}}$. Using a direct limit argument, we see that there exists a finite algebraic extension κ' of κ

such that $M' = (\kappa' \otimes_{\kappa} M)^{1/p^\infty}$ belongs to the essential image of the functor $\mathrm{RH}^c : \mathrm{Shv}_{\kappa'}^c \rightarrow \mathrm{Mod}_{\kappa'}^{\mathrm{hol}}$. By restriction of scalars, we can regard M' as an object of $\mathrm{Mod}_{\kappa}^{\mathrm{hol}}$ (Proposition 4.2.10), and the resulting object belongs to the subcategory \mathcal{C} (Theorem 6.5.1). We have an evident monomorphism $M \rightarrow M'$ in the abelian category $\mathrm{Mod}_{\kappa}^{\mathrm{hol}}$. Applying the same argument to the quotient M/M' , we can choose a monomorphism $M/M' \hookrightarrow M''$ for some $M'' \in \mathcal{C}$. It follows that M can be identified with the kernel of the composite map $M' \rightarrow M/M' \hookrightarrow M''$, and therefore belongs to \mathcal{C} (since \mathcal{C} is an abelian subcategory of $\mathrm{Mod}_{\kappa}^{\mathrm{hol}}$). \square

7.4. Proof of the Main Theorem

We now generalize Proposition 7.3.1 to the case of an arbitrary \mathbf{F}_p -algebra:

Theorem 7.4.1. *Let R be a commutative \mathbf{F}_p -algebra. Then the Riemann-Hilbert functor $\mathrm{RH}^c : \mathrm{Shv}_{\mathrm{ét}}^c(\mathrm{Spec}(R), \mathbf{F}_p) \rightarrow \mathrm{Mod}_R^{\mathrm{hol}}$ (see Notation 6.3.2) is an equivalence of categories.*

Before giving the proof of Theorem 7.4.1, let us collect some of its consequences. First, we note that it immediately implies the results of this paper:

Proof of Theorem 1.0.2 from Theorem 7.4.1. Let R be an \mathbf{F}_p -algebra. It follows from Theorem 7.4.1 that the functor $\mathrm{RH}^c = \mathrm{RH}|_{\mathrm{Shv}_{\mathrm{ét}}^c(\mathrm{Spec}(R), \mathbf{F}_p)}$ is a fully faithful embedding, whose essential image consists of compact objects of $\mathrm{Mod}_R^{\mathrm{perf}}$ (see Proposition 4.1.5). Moreover, the functor RH preserves filtered colimits (by virtue of the fact that it is left adjoint to the solution functor). Using Proposition 6.3.7, we deduce that RH is a fully faithful embedding whose essential image consists of those perfect Frobenius modules which can be realized as filtered colimits of holonomic Frobenius modules. By virtue of Theorem 4.2.9, this essential image is exactly $\mathrm{Mod}_R^{\mathrm{alg}}$. \square

Proof of Theorem 2.4.3 from Theorem 7.4.1. Let R be an \mathbf{F}_p -algebra. Then $\mathrm{Sol} : \mathrm{Mod}_R^{\mathrm{alg}} \rightarrow \mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(R), \mathbf{F}_p)$ is right adjoint to the Riemann-Hilbert functor $\mathrm{RH} : \mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(R), \mathbf{F}_p) \rightarrow \mathrm{Mod}_R^{\mathrm{alg}}$. Since the latter is an equivalence of categories, the former must also be an equivalence of categories. \square

Corollary 7.4.2. *Let $f : A \rightarrow B$ be a homomorphism of \mathbf{F}_p -algebras and let M be an algebraic A -module. Then the comparison map $f^*(\mathrm{Sol}(M)) \rightarrow \mathrm{Sol}(f^\diamond M)$ is an isomorphism in $\mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(B), \mathbf{F}_p)$.*

Proof. Combine Theorem 2.4.3 with Proposition 6.2.2. \square

Corollary 7.4.3. *Let $A \rightarrow B$ be a homomorphism of commutative \mathbf{F}_p -algebras which is étale and faithfully flat, and let M be a perfect Frobenius module over A . If $f^\diamond(M) = B \otimes_A M$ is a holonomic Frobenius module over B , then M is a holonomic Frobenius module over A .*

Proof. It follows from Lemma 5.4.6 that M is algebraic. Consequently, to show that M is holonomic, it will suffice (by virtue of Theorems 1.0.2 and 7.4.1) to show that $\mathrm{Sol}(M)$ is a constructible sheaf. This follows from Remark 6.2.4, since constructibility of étale sheaves can be tested locally with respect to the étale topology. \square

Corollary 7.4.4. *Let R be a commutative \mathbf{F}_p -algebra and let M be an algebraic Frobenius module over R . Then $\mathrm{Sol}^i(M) \simeq 0$ for $i > 0$.*

Proof. By virtue of Theorem 1.0.2 we can write $M = \mathrm{RH}(\mathcal{F})$ for some $\mathcal{F} \in \mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(R), \mathbf{F}_p)$. In this case, the desired result follows from Proposition 7.2.1. \square

Proof of Theorem 7.4.1. Let R be a commutative \mathbf{F}_p -algebra. As in the proof of Lemma 7.3.2, we see that the functor RH^c is a fully faithful embedding whose essential image $\mathcal{C} \subseteq \mathrm{Mod}_R^{\mathrm{hol}}$ is an abelian subcategory which is closed under extensions. We wish to show that \mathcal{C} contains every object $M \in \mathrm{Mod}_R^{\mathrm{hol}}$. Using a direct limit argument, we can choose an \mathbf{F}_p -algebra homomorphism $\iota : R_0 \rightarrow R$ and an equivalence $M \simeq \iota^\diamond M_0$ for some $M_0 \in \mathrm{Mod}_{R_0}^{\mathrm{hol}}$, where R_0 is finitely generated over \mathbf{F}_p . By virtue of Proposition 6.2.2, it will suffice to show that M_0 belongs to the essential image of the functor $\mathrm{RH}^c : \mathrm{Shv}_{R_0}^c \rightarrow \mathrm{Mod}_{R_0}^{\mathrm{hol}}$. We may therefore replace R by R_0 (and M by M_0) and thereby reduce to the case where R is Noetherian.

Applying Proposition 4.3.1, we see that M is a Noetherian object of the abelian category $\mathrm{Mod}_R^{\mathrm{alg}}$. Consequently, there exists a subobject $M' \subseteq M$ (in the abelian category $\mathrm{Mod}_R^{\mathrm{alg}}$) which is maximal among those subobjects which belong to \mathcal{C} . It follows from the maximality of M' (and the stability of \mathcal{C} under extensions) that the quotient M/M' does not contain any nonzero subobjects which belong to \mathcal{C} . Replacing M by M/M' , we can reduce to the case where M does not have any nonzero subobjects which belong to \mathcal{C} .

Let $K \subseteq \mathrm{Spec}(R)$ be the closure of the support $\mathrm{supp}(M)$. Then K is the vanishing locus of a radical ideal $I \subseteq \mathrm{Spec}(R)$. Using Theorem 5.3.1, we see that M can be regarded as a holonomic Frobenius module over the quotient ring R/I . Using Theorem 6.5.1, we can replace A by A/I and thereby reduce to the case where R is reduced and $K = \mathrm{Spec}(R)$.

If $R \simeq 0$, then $M \simeq 0$ and there is nothing to prove. Otherwise, R contains a minimal prime ideal \mathfrak{p} . Since R is reduced, the localization

$R_{\mathfrak{p}}$ is a field. Applying Proposition 7.3.1, we deduce that $M_{\mathfrak{p}}$ belongs to the essential image of the functor $\mathrm{RH}^c : \mathrm{Shv}_{R_{\mathfrak{p}}}^c \rightarrow \mathrm{Mod}_{R_{\mathfrak{p}}}^{\mathrm{hol}}$. It follows by a direct limit argument that there exists some element $t \in R - \mathfrak{p}$ for which the localization $M[t^{-1}]$ belongs to the essential image of the functor $\mathrm{RH}^c : \mathrm{Shv}_{R[t^{-1}]}^c \rightarrow \mathrm{Mod}_{R[t^{-1}]}^{\mathrm{hol}}$. Let $f : R \rightarrow R[t^{-1}]$ be the localization map, and set $M' = f_! M[t^{-1}]$; using Proposition 6.2.3, we deduce that M' belongs to the essential image of the Riemann-Hilbert functor $\mathrm{RH}^c : \mathrm{Shv}_R^c \rightarrow \mathrm{Mod}_R^{\mathrm{hol}}$. Note that Lemma 5.4.8 guarantees that the counit map $M' = f_! f^* M \rightarrow M$ is a monomorphism, so we must have $M' \simeq 0$. It follows that the localization $M[t^{-1}]$ vanishes, so that the prime ideal \mathfrak{p} cannot belong to the support of M . Using the constructibility of $\mathrm{supp}(M)$ (Theorem 4.4.4), we deduce that there exists an open neighborhood of \mathfrak{p} which does not intersect $\mathrm{supp}(M)$, contradicting the equality $K = \mathrm{Spec}(R)$. \square

8. Tensor Products

Let A be a commutative ring and let $\mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(A), \mathbf{F}_p)$ denote the category of p -torsion étale sheaves on $\mathrm{Spec}(A)$. This category is equipped with a tensor product functor

$$\otimes_{\mathbf{F}_p} : \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(A), \mathbf{F}_p) \times \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(A), \mathbf{F}_p) \rightarrow \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(A), \mathbf{F}_p)$$

which carries a pair of étale sheaves $(\mathcal{F}, \mathcal{G})$ to the sheafification of the presheaf

$$(B \in \mathrm{CAlg}_A^{\acute{\mathrm{e}}\mathrm{t}}) \mapsto \mathcal{F}(B) \otimes_{\mathbf{F}_p} \mathcal{G}(B).$$

In the case where A is an \mathbf{F}_p -algebra, Theorem 2.4.3 supplies an equivalence of categories

$$\mathrm{Sol} : \mathrm{Mod}_A^{\mathrm{alg}} \rightarrow \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(A), \mathbf{F}_p)$$

Our goal this section is to promote the solution functor Sol to an equivalence of *symmetric monoidal* categories: that is, to show that it is compatible with tensor products.

We begin in §8.1 by studying an analogous tensor product operation on the category $\mathrm{Mod}_A^{\mathrm{Fr}}$ of Frobenius modules over A . In fact, there are two such operations (which are closely related):

- If M and N are Frobenius modules over A , then the tensor product $M \otimes_A N$ inherits the structure of a Frobenius module over A (Construction 8.1.1).

- If M and N are perfect Frobenius modules over A , then they can also be regarded as modules of the perfection A^{1/p^∞} ; in this case, the tensor product $M \otimes_{A^{1/p^\infty}} N$ inherits the structure of a perfect Frobenius module over A (Remark 8.1.5).

Like the usual tensor product on the category of A -modules, the tensor product on Frobenius modules is right exact but generally not left exact. One can partially remedy this failure of exactness by studying left derived functors of the tensor product: in §8.2, we show that these agree with the usual Tor-functors of commutative algebra (Proposition 8.2.1). The central result of this section asserts that if we restrict our attention to *algebraic* Frobenius modules, then these Tor-groups automatically vanish (when computed relative to the perfection A^{1/p^∞} ; see Theorem 8.3.1). We prove this statement in §8.3, and apply it in §8.4 to show that the Riemann-Hilbert correspondence is compatible with tensor products (Theorem 8.4.1).

8.1. Tensor Products of Frobenius Modules

We begin with some general remarks.

Construction 8.1.1. Let A be a commutative \mathbf{F}_p -algebra. If M and N are Frobenius modules over A , then we regard the tensor product $M \otimes_A N$ as a Frobenius module over A , with Frobenius map

$$\varphi_{M \otimes_A N} : M \otimes_A N \rightarrow M \otimes_A N$$

given by the formula $\varphi_{M \otimes_A N}(x \otimes y) = \varphi_M(x) \otimes \varphi_N(y)$. Note that the commutativity and associativity isomorphisms

$$M \otimes_A N \simeq N \otimes_A M \quad (M \otimes_A N) \otimes_A P \simeq M \otimes_A (N \otimes_A P)$$

are isomorphisms of Frobenius modules, and therefore endow Mod_A^{Fr} with the structure of a symmetric monoidal category.

Example 8.1.2 (Tensor Products of Free Modules). Let A be a commutative \mathbf{F}_p -algebra and let M and N be Frobenius modules over A which are freely generated (as left $A[F]$ -modules) by elements $x \in M$ and $y \in N$. Then the tensor product $M \otimes_A N$ is freely generated by the elements $F^n x \otimes y$ and $x \otimes F^n y$ (which coincide when $n = 0$).

Remark 8.1.3 (Compatibility with Extension of Scalars). Let $f : A \rightarrow B$ be a homomorphism of commutative \mathbf{F}_p -algebras, and let $f_{\text{Fr}}^* : \text{Mod}_A^{\text{Fr}} \rightarrow \text{Mod}_B^{\text{Fr}}$ be the functor of extension of scalars along f (given by $M \mapsto B \otimes_A M$).

Then f_{Fr}^* is a symmetric monoidal functor: in particular, we have canonical isomorphisms $f_{\text{Fr}}^*(M \otimes_A N) \simeq (f_{\text{Fr}}^*M) \otimes_B (f_{\text{Fr}}^*N)$.

Remark 8.1.4. Let A be an \mathbf{F}_p -algebra. If M and N are Frobenius modules over A , then we have a canonical isomorphism

$$(M \otimes_A N)^{1/p^\infty} \simeq M^{1/p^\infty} \otimes_{A^{1/p^\infty}} N^{1/p^\infty}.$$

In particular, if A , M , and N are perfect, then the tensor product $M \otimes_A N$ is also perfect.

Remark 8.1.5. Let A be a perfect \mathbf{F}_p -algebra. It follows from Remark 8.1.4 that the full subcategory $\text{Mod}_A^{\text{perf}} \subseteq \text{Mod}_A^{\text{Fr}}$ is closed under tensor products, and therefore inherits the structure of a symmetric monoidal category (with tensor product \otimes_A).

More generally, if A is an arbitrary \mathbf{F}_p -algebra, then the restriction-of-scalars functor $\theta : \text{Mod}_{A^{1/p^\infty}}^{\text{perf}} \rightarrow \text{Mod}_A^{\text{perf}}$ is an equivalence of categories (Proposition 3.4.3). It follows that there is an essentially unique symmetric monoidal structure on the category $\text{Mod}_A^{\text{perf}}$ for which the functor θ is symmetric monoidal. We will denote the underlying tensor product by $(M, N) \mapsto M \otimes_{A^{1/p^\infty}} N$ (note that if M and N are perfect Frobenius modules over A , then they can be regarded as modules over A^{1/p^∞} in an essentially unique way).

Warning 8.1.6. Let A be a commutative \mathbf{F}_p -algebra. Then the inclusion functor $\text{Mod}_A^{\text{perf}} \hookrightarrow \text{Mod}_A^{\text{Fr}}$ is usually not a symmetric monoidal functor, if we regard Mod_A^{Fr} as equipped with the symmetric monoidal structure of Construction 8.1.1 (given by tensor product over A) and $\text{Mod}_A^{\text{perf}}$ with the symmetric monoidal structure of Remark 8.1.5 (given by tensor product over A^{1/p^∞}). However, it has a symmetric monoidal left adjoint, given by the perfection construction $M \mapsto M^{1/p^\infty}$ (note that Remark 8.1.4 supplies an isomorphism $(M \otimes_A N)^{1/p^\infty} \simeq M \otimes_{A^{1/p^\infty}} N$ in the case where M and N are perfect).

Remark 8.1.7 (Compatibility with Extension of Scalars). Let $f : A \rightarrow B$ be a homomorphism of commutative \mathbf{F}_p -algebras, and let $f^\diamond : \text{Mod}_A^{\text{perf}} \rightarrow \text{Mod}_B^{\text{perf}}$ be the functor of Proposition 3.3.2. Then f^\diamond is symmetric monoidal with respect to the tensor products described in Remark 8.1.5: in particular, if M and N are perfect Frobenius modules over A , then we have a canonical isomorphism $f^\diamond(M \otimes_{A^{1/p^\infty}} N) \simeq (f^\diamond M) \otimes_{B^{1/p^\infty}} (f^\diamond N)$. This follows from Remark 8.1.3, applied to the map $f^{1/p^\infty} : A^{1/p^\infty} \rightarrow B^{1/p^\infty}$.

8.2. Derived Tensor Products

Let A be an \mathbf{F}_p -algebra and let M be a Frobenius module over A . Then the construction $N \mapsto M \otimes_A N$ determines a right exact functor from the abelian category Mod_A^{Fr} to itself. Since the abelian category Mod_A^{Fr} has enough projective objects (it is equivalent to the category of left modules over the noncommutative ring $A[F]$ of Notation 2.1.5), the construction $N \mapsto M \otimes_A N$ admits left derived functors, which we will temporarily denote by $N \mapsto T_*(M, N)$. More concretely, we define $T_k(M, N)$ to be the k th homology group of the chain complex

$$\cdots \rightarrow M \otimes_A P_2 \rightarrow M \otimes_A P_1 \rightarrow M \otimes_A P_0 \rightarrow 0,$$

where $\cdots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow N \rightarrow 0$ is a projective resolution of N in the category Mod_A^{Fr} (it follows from elementary homological algebra that the Frobenius modules $T_*(M, N)$ are independent of the choice of resolution, up to canonical isomorphism).

Proposition 8.2.1. *Let A be an \mathbf{F}_p -algebra. For every pair of Frobenius modules M and N over A , we have canonical A -module isomorphisms $\text{Tor}_*^A(M, N) \xrightarrow{\sim} T_*(M, N)$.*

Proof. Since $A[F]$ is free as a left A -module, every projective left $A[F]$ -module is also projective when viewed as a left A -module. Consequently, if P_* is a resolution of N by projective left $A[F]$ -modules, then it is also a resolution of N by projective A -modules, so the homology groups of the chain complex $M \otimes_A P_*$ can be identified with $\text{Tor}_*^A(M, N)$. \square

We can formulate Proposition 8.2.1 more informally as follows: if M and N are Frobenius modules over A , then the Tor-groups $\text{Tor}_*^A(M, N)$ inherit the structure of Frobenius modules over A .

Remark 8.2.2. Our description of the Frobenius structure on the Tor-groups $\text{Tor}_*^A(M, N)$ is *a priori* asymmetric in M and N , since it depends on taking the left derived functors of the construction $M \otimes_A \bullet$. However, one can give a more symmetric description as follows. Let \mathcal{C} denote the category whose objects are triples (A, M, N) , where A is an associative ring, M is a right module over A , and N is a left module over A . For each integer k , the construction $(A, M, N) \mapsto \text{Tor}_k^A(M, N)$ can be regarded as a functor from \mathcal{C} to the category of abelian groups. In the special case where A is a commutative \mathbf{F}_p -algebra and $M, N \in \text{Mod}_A^{\text{Fr}}$, we can regard the triple $(\varphi_A, \varphi_M, \varphi_N)$ as a morphism from (A, M, N) to itself in the category \mathcal{C} , and

therefore induces a map of abelian groups $\varphi : \mathrm{Tor}_k^A(M, N) \rightarrow \mathrm{Tor}_k^A(M, N)$. It is easy to check that φ corresponds to the Frobenius map on $T_k(M, N)$ under the isomorphism of Proposition 8.2.1.

Proposition 8.2.3. *Let A be a perfect \mathbf{F}_p -algebra. If M and N are perfect Frobenius modules over A , then the Tor-groups $\mathrm{Tor}_*^A(M, N)$ are also perfect Frobenius modules over A .*

Proof. This follows immediately from the description of the Frobenius structure on $\mathrm{Tor}_*^A(M, N)$ given in Remark 8.2.2. Alternatively, we can show that each $\mathrm{Tor}_k^A(M, N)$ is perfect using induction on k . When $k = 0$, the desired result follows from Remark 8.1.4. For $k > 0$, we can choose a short exact sequence $0 \rightarrow N' \rightarrow P \rightarrow N \rightarrow 0$, where P is a free module over $A^{1/p^\infty}[F^{\pm 1}]$ (see Example 3.2.5). Then N' is also a perfect Frobenius module over A . Moreover, since A is perfect, the ring $A^{1/p^\infty}[F^{\pm 1}]$ is free as a left A -module, so the groups $\mathrm{Tor}_*^A(M, P)$ vanish for $* > 0$. We therefore have a short exact sequence

$$0 \rightarrow \mathrm{Tor}_k^A(M, N) \rightarrow \mathrm{Tor}_{k-1}^A(M, N') \rightarrow \mathrm{Tor}_{k-1}^A(M, P)$$

which exhibits $\mathrm{Tor}_k^A(M, N)$ as the kernel of a map between perfect Frobenius modules, so that $\mathrm{Tor}_k^A(M, N)$ is itself perfect. \square

Variante 8.2.4. Let A be an arbitrary \mathbf{F}_p -algebra, and let M and N be perfect Frobenius modules over A . Then we can regard M and N as Frobenius modules over A^{1/p^∞} in an essentially unique way (Proposition 3.4.3). Using Proposition 8.2.3, we can regard the Tor-groups $\mathrm{Tor}_*^{A^{1/p^\infty}}(M, N)$ as perfect Frobenius modules over A^{1/p^∞} , and therefore also (by restriction of scalars) as perfect Frobenius modules over A .

Proposition 8.2.5. *Let A be an \mathbf{F}_p -algebra and let M and N be Frobenius modules over A . Then the canonical map $\mathrm{Tor}_*^A(M, N) \rightarrow \mathrm{Tor}_*^{A^{1/p^\infty}}(M^{1/p^\infty}, N^{1/p^\infty})$ induces an isomorphism of Frobenius modules*

$$\rho_* : \mathrm{Tor}_*^A(M, N)^{1/p^\infty} \simeq \mathrm{Tor}_*^{A^{1/p^\infty}}(M^{1/p^\infty}, N^{1/p^\infty}).$$

Proof. Let us regard M as fixed. We will show that for every Frobenius module N and every nonnegative integer k , the map

$$\rho_k : \mathrm{Tor}_k^A(M, N)^{1/p^\infty} \rightarrow \mathrm{Tor}_k^{A^{1/p^\infty}}(M^{1/p^\infty}, N^{1/p^\infty})$$

is an isomorphism. The proof proceeds by induction on k . When $k = 0$, the desired result is the content of Remark 8.1.4. Assume that $k > 0$, and

choose a short exact sequence of Frobenius modules $0 \rightarrow N' \rightarrow P \rightarrow N \rightarrow 0$ where P is a free left module over $A[F]$. Then P^{1/p^∞} is a free left module over $A^{1/p^\infty}[F^{\pm 1}]$, and is therefore also free as an A^{1/p^∞} -module. It follows that the groups $\mathrm{Tor}_k^A(M, P)$ and $\mathrm{Tor}_k^{A^{1/p^\infty}}(M^{1/p^\infty}, P^{1/p^\infty})$ both vanish. Consequently, the map ρ_k fits into a commutative diagram of exact sequences

$$\begin{array}{ccc}
 0 & & 0 \\
 \downarrow & & \downarrow \\
 \mathrm{Tor}_k^A(M, N)^{1/p^\infty} & \xrightarrow{\rho_k} & \mathrm{Tor}_k^{A^{1/p^\infty}}(M^{1/p^\infty}, N^{1/p^\infty}) \\
 \downarrow & & \downarrow \\
 \mathrm{Tor}_{k-1}^A(M, N')^{1/p^\infty} & \xrightarrow{\rho'} & \mathrm{Tor}_{k-1}^{A^{1/p^\infty}}(M^{1/p^\infty}, N'^{1/p^\infty}) \\
 \downarrow & & \downarrow \\
 \mathrm{Tor}_{k-1}^A(M, P)^{1/p^\infty} & \xrightarrow{\rho''} & \mathrm{Tor}_{k-1}^{A^{1/p^\infty}}(M^{1/p^\infty}, P^{1/p^\infty}).
 \end{array}$$

The maps ρ' and ρ'' are isomorphisms by our inductive hypothesis, so that ρ_k is an isomorphism as well. □

Remark 8.2.6 (Compatibility with Extension of Scalars). Let $f : A \rightarrow B$ be a homomorphism of \mathbf{F}_p -algebras. Then the extension of scalars functor

$$f_{\mathrm{Fr}}^* : \mathrm{Mod}_A^{\mathrm{Fr}} \rightarrow \mathrm{Mod}_B^{\mathrm{Fr}}$$

is right exact, having left derived functors $N \mapsto \mathrm{Tor}_*^A(B, N)$. Let M be a Frobenius module over B . Then we can regard the functors $\{\mathrm{Tor}_k^A(M, \bullet)\}_{k \geq 0}$ the left derived functor of the construction $N \mapsto M \otimes_B (f_{\mathrm{Fr}}^* N)$. Since the functor $f_{\mathrm{Fr}}^* : \mathrm{Mod}_A^{\mathrm{Fr}} \rightarrow \mathrm{Mod}_B^{\mathrm{Fr}}$ carries projective objects to projective objects, we have a Grothendieck spectral sequence (in the abelian category $\mathrm{Mod}_A^{\mathrm{Fr}}$)

$$\mathrm{Tor}_s^B(M, \mathrm{Tor}_t^A(B, N)) \Rightarrow \mathrm{Tor}_{s+t}^A(M, N).$$

If M and N are perfect, then we can apply the same reasoning to the induced map $A^{1/p^\infty} \rightarrow B^{1/p^\infty}$ to obtain a Grothendieck spectral sequence

$$\mathrm{Tor}_s^{B^{1/p^\infty}}(M, \mathrm{Tor}_t^{A^{1/p^\infty}}(B^{1/p^\infty}, N)) \Rightarrow \mathrm{Tor}_{s+t}^{A^{1/p^\infty}}(M, N).$$

8.3. Tensor Products of Holonomic Modules

Our next goal is to prove the following variant of Theorem 3.5.1:

Theorem 8.3.1. *Let A be an \mathbf{F}_p -algebra and let M and N be algebraic Frobenius modules over A . Then:*

- (1) *The tensor product $M \otimes_{A^{1/p^\infty}} N$ is algebraic.*
- (2) *The Tor-groups $\mathrm{Tor}_*^{A^{1/p^\infty}}(M, N)$ vanish for $* > 0$.*
- (3) *If M and N are holonomic, then $M \otimes_{A^{1/p^\infty}} N$ is holonomic.*

The proof of Theorem 8.3.1 will require some preliminaries.

Lemma 8.3.2. *Let A be a Noetherian \mathbf{F}_p -algebra, and let M and N be holonomic Frobenius modules over A . Then:*

- (1) *The Tor-groups $\mathrm{Tor}_*^{A^{1/p^\infty}}(M, N)$ are also holonomic Frobenius modules over A .*
- (2) *Let $f : A \rightarrow B$ be any homomorphism of commutative rings. Then the canonical map*

$$f^\diamond \mathrm{Tor}_*^{A^{1/p^\infty}}(M, N) \rightarrow \mathrm{Tor}_*^{B^{1/p^\infty}}(f^\diamond M, f^\diamond N)$$

is an isomorphism.

Proof. Since M and N are holonomic, we can write $M = M_0^{1/p^\infty}$ and $N = N_0^{1/p^\infty}$, where $M_0, N_0 \in \mathrm{Mod}_A^{\mathrm{Fr}}$ are finitely generated as A -modules. The assumption that A is Noetherian guarantees that the Tor-groups $\mathrm{Tor}_k^A(M_0, N_0)$ is finitely generated as an A -module. Using the isomorphisms $\mathrm{Tor}_k^{A^{1/p^\infty}}(M_0^{1/p^\infty}, N_0^{1/p^\infty}) \simeq \mathrm{Tor}_k^A(M_0, N_0)^{1/p^\infty}$ of Proposition 8.2.5, we conclude that each $\mathrm{Tor}_k^{A^{1/p^\infty}}(M, N)$ is holonomic. This proves (1).

We now prove (2). Let $f : A \rightarrow B$ be a homomorphism of commutative rings. Let P_* and Q_* be resolutions of M and N by projective objects of $\mathrm{Mod}_A^{\mathrm{perf}}$. Then P_* and Q_* are also resolutions of M and N by projective A^{1/p^∞} -modules. It follows that the homology groups of the complexes $f^\diamond P_*$ and $f^\diamond Q_*$ can be identified with the groups $\mathrm{Tor}_*^{A^{1/p^\infty}}(B^{1/p^\infty}, M)$ and $\mathrm{Tor}_*^{A^{1/p^\infty}}(B^{1/p^\infty}, N)$, which vanish for $* > 0$ by virtue of Theorem 3.5.1. In other words, we can regard $f^\diamond P_*$ and $f^\diamond Q_*$ as projective resolutions of $f^\diamond M$ and $f^\diamond N$, respectively. It follows that $\mathrm{Tor}_*^{B^{1/p^\infty}}(f^\diamond M, f^\diamond N)$ can be identified with the homology of the tensor product complex

$$(f^\diamond P_*) \otimes_{B^{1/p^\infty}} (f^\diamond Q_*) \simeq B^{1/p^\infty} \otimes_{A^{1/p^\infty}} (P_* \otimes_{A^{1/p^\infty}} Q_*).$$

We therefore have a convergent spectral sequence

$$E_{s,t}^2 = \mathrm{Tor}_s^{A^{1/p^\infty}}(B^{1/p^\infty}, \mathrm{Tor}_t^{A^{1/p^\infty}}(M, N)) \Rightarrow \mathrm{Tor}_{s+t}^{B^{1/p^\infty}}(f^\diamond M, f^\diamond N).$$

To prove assertion (2), it will suffice to show that the groups $E_{s,t}^2$ vanish for $s > 0$, which follows from assertion (1) and Theorem 3.5.1. \square

Proof of Theorem 8.3.1. Let M and N be algebraic Frobenius modules over an \mathbf{F}_p -algebra A ; we wish to prove that the tensor product $M \otimes_{A^{1/p^\infty}} N$ is algebraic and that the Tor-groups $\mathrm{Tor}_*^{A^{1/p^\infty}}(M, N)$ vanish for $* > 0$. Using Theorem 4.2.9, we can write M as a filtered colimit of holonomic Frobenius modules and thereby reduce to the case where M is holonomic. Similarly, we can assume that N is holonomic. Applying Proposition 4.1.3, we can assume that $M = \iota^\diamond M'$ and $N = \iota^\diamond N'$, where $\iota : A' \hookrightarrow A$ is the inclusion of a finitely generated subalgebra and $M', N' \in \mathrm{Mod}_A^{\mathrm{hol}}$. In this case, Lemma 8.3.2 supplies isomorphisms $\mathrm{Tor}_*^{A^{1/p^\infty}}(M, N) \simeq \iota^\diamond \mathrm{Tor}_*^{A'^{1/p^\infty}}(M', N')$. We may therefore replace A by A' , and thereby reduce to the case where A is Noetherian. It now follows from Lemma 8.3.2 that the Tor-groups $\mathrm{Tor}_s^{A^{1/p^\infty}}(M, N)$ are holonomic for each $s \geq 0$; we wish to show that they vanish for $s > 0$. By virtue of Proposition 5.3.3, it will suffice to show that $f^\diamond \mathrm{Tor}_s^{A^{1/p^\infty}}(M, N) \simeq 0$ for every homomorphism $f : A \rightarrow \kappa$, where κ is a field. Applying Lemma 8.3.2 again, we can reduce to the case where $A = \kappa$, in which case the vanishing is automatic. \square

8.4. Compatibility with the Riemann-Hilbert Correspondence

Let A be a commutative \mathbf{F}_p -algebra and let $\mathrm{Sol} : \mathrm{Mod}_A^{\mathrm{perf}} \rightarrow \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(A), \mathbf{F}_p)$ be the solution sheaf functor (Construction 9.3.1), given by the formula

$$\mathrm{Sol}(M)(B) = \{x \in (B \otimes_A M) : \varphi_{B \otimes_A M}(x) = x\}.$$

Note that if $x \in \mathrm{Sol}(M)(B)$ and $y \in \mathrm{Sol}(N)(B)$, then the tensor $x \otimes y$ can be regarded as an element of $\mathrm{Sol}(M \otimes_{A^{1/p^\infty}} N)(B)$. This observation determines a bilinear map

$$\mathrm{Sol}(M)(B) \times \mathrm{Sol}(N)(B) \rightarrow \mathrm{Sol}(M \otimes_{A^{1/p^\infty}} N)(B)$$

which depends functorially on B , and therefore gives rise to a map of sheaves $\mathrm{Sol}(M) \otimes_{\mathbf{F}_p} \mathrm{Sol}(N) \rightarrow \mathrm{Sol}(M \otimes_{A^{1/p^\infty}} N)$.

Theorem 8.4.1. *Let A be a commutative \mathbf{F}_p -algebra and suppose that M and N are algebraic A -modules. Then the comparison map*

$$\theta : \mathrm{Sol}(M) \otimes_{\mathbf{F}_p} \mathrm{Sol}(N) \rightarrow \mathrm{Sol}(M \otimes_{A^{1/p^\infty}} N)$$

is an isomorphism in the category $\mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(A), \mathbf{F}_p)$.

Proof. It will suffice to show that for every algebraically closed field κ and every homomorphism $f : A \rightarrow \kappa$, the pullback $f^*(\theta)$ is an isomorphism in $\mathrm{Shv}_{\kappa}^{\acute{e}t}$. Since M , N , and $M \otimes_{A^{1/p^\infty}} N$ are algebraic (Theorem 8.3.1), we can identify $f^*(\theta)$ with the tautological map

$$\mathrm{Sol}(f^\circ M) \otimes_{\mathbf{F}_p} \mathrm{Sol}(f^\circ N) \rightarrow \mathrm{Sol}(f^\circ M \otimes_{\kappa} f^\circ N).$$

We may therefore replace A by κ and thereby reduce to the case where A is an algebraically closed field. In this case, Theorem 2.4.3 implies that $\mathrm{Mod}_A^{\mathrm{alg}}$ is equivalent to the category of vector spaces over \mathbf{F}_p . Consequently, the Frobenius modules M and N can be decomposed as a direct sum of copies of $A^{1/p^\infty} = \kappa$, and the desired result is obvious. \square

Corollary 8.4.2. *Let A be a commutative \mathbf{F}_p -algebra. Then the Riemann-Hilbert functor $\mathrm{RH} : \mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(A), \mathbf{F}_p) \rightarrow \mathrm{Mod}_A^{\mathrm{perf}}$ admits the structure of a symmetric monoidal functor (where the symmetric monoidal structure on $\mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(A), \mathbf{F}_p)$ is given by the usual tensor product of sheaves, and the symmetric monoidal structure on $\mathrm{Mod}_A^{\mathrm{perf}}$ is given by the tensor product $\otimes_{A^{1/p^\infty}}$ of Remark 8.1.5).*

Proof. It follows from Theorem 8.4.1 that the lax symmetric monoidal functor $\mathrm{Sol} : \mathrm{Mod}_A^{\mathrm{perf}} \rightarrow \mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(A), \mathbf{F}_p)$ is symmetric monoidal when restricted to $\mathrm{Mod}_A^{\mathrm{alg}}$. Combining this observation with Theorem 2.4.3, we see that the functor $\mathrm{Sol}|_{\mathrm{Mod}_A^{\mathrm{alg}}}$ is an equivalence of symmetric monoidal categories. We conclude by observing that the functor RH can be obtained by composing an inverse of $\mathrm{Sol}|_{\mathrm{Mod}_A^{\mathrm{alg}}}$ with the inclusion functor $\mathrm{Mod}_A^{\mathrm{alg}} \hookrightarrow \mathrm{Mod}_A^{\mathrm{perf}}$. \square

9. The p^n -Torsion Case

Let R be a commutative \mathbf{F}_p -algebra. Theorem 1.0.2 supplies a fully faithful embedding from the category $\mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(R), \mathbf{F}_p)$ p -torsion étale sheaves on $\mathrm{Spec}(R)$ to the category of Frobenius modules over R . Our goal in this section is to prove a generalization of Theorem 1.0.2, which gives an analogous

realization for the category $\mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(R), \mathbf{Z}/p^n\mathbf{Z})$ of $\mathbf{Z}/p^n\mathbf{Z}$ -torsion étale sheaves, for any nonnegative integer n . Our first step will be to introduce an analogous enlargement of the category $\mathrm{Mod}_R^{\mathrm{Fr}}$ of Frobenius module over R . In §9.1, we define a notion of Frobenius module over $W_n(R)$, where $W_n(R)$ is the ring of length n Witt vectors over R (Definition 9.1.1). The collection of such Frobenius modules can be organized into a category $\mathrm{Mod}_{W_n(R)}^{\mathrm{Fr}}$. In §9.2, we study the dependence of this category on the \mathbf{F}_p -algebra R (emphasizing in particular the effect of replacing R by its perfection R^{1/p^∞} , which makes Witt vectors much more pleasant to work with). In §9.3, we introduce a *solution functor*

$$\mathrm{Sol} : \mathrm{Mod}_{W_n(R)}^{\mathrm{Fr}} \rightarrow \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(R), \mathbf{Z}/p^n\mathbf{Z})$$

connecting Frobenius modules over $W_n(R)$ to p^n -torsion étale sheaves (which reduces to Construction 2.3.1 in the case $n = 1$). Like its p -torsion counterpart, this solution functor is not exact. However, we show in §9.4 that it is *almost* exact when restricted to perfect Frobenius modules, in the sense that it has only one nonvanishing derived functor (Proposition 9.4.1). We apply this result in §9.6 to show that the functor Sol restricts to an equivalence of categories $\mathrm{Mod}_{W_n(R)}^{\mathrm{alg}} \simeq \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(R), \mathbf{Z}/p^n\mathbf{Z})$ (Theorem 9.6.1). Here $\mathrm{Mod}_{W_n(R)}^{\mathrm{alg}}$ denotes the full subcategory of $\mathrm{Mod}_{W_n(R)}^{\mathrm{Fr}}$ spanned by the *algebraic* Frobenius modules over $W_n(R)$, which we introduce in §9.5 (Definition 9.5.2).

9.1. Frobenius Modules over the Witt Vectors

We begin by extending some of the notions introduced in §3. Let R be a commutative \mathbf{F}_p -algebra. For every nonnegative integer n , we let $W_n(R)$ denote the ring of length n Witt vectors of R . The Frobenius map $\varphi_R : R \rightarrow R$ induces a ring homomorphism $F : W_n(R) \rightarrow W_n(R)$, which we will refer to as the *Witt vector Frobenius*.

Definition 9.1.1. Let R be a commutative \mathbf{F}_p -algebra and let $n \geq 0$ be an integer. A *Frobenius module over $W_n(R)$* is an $W_n(R)$ -module M equipped with an additive map $\varphi_M : M \rightarrow M$ satisfying the identity $\varphi_M(\lambda x) = F(\lambda)\varphi_M(x)$ for $x \in M$, $\lambda \in W_n(R)$. We will say that a Frobenius module M is *perfect* if the map $\varphi_M : M \rightarrow M$ is an isomorphism of abelian groups.

Let (M, φ_M) and (N, φ_N) be Frobenius modules over $W_n(R)$. A *morphism of Frobenius modules* from (M, φ_M) to (N, φ_N) is an $W_n(R)$ -module

homomorphism $\rho : M \rightarrow N$ for which the diagram

$$\begin{array}{ccc} M & \xrightarrow{\rho} & N \\ \downarrow \varphi_M & & \downarrow \varphi_N \\ M & \xrightarrow{\rho} & M' \end{array}$$

commutes. We let $\text{Mod}_{W_n(R)}^{\text{Fr}}$ denote the category whose objects are Frobenius modules (M, φ_M) over $W_n(R)$, and whose morphisms are morphisms of Frobenius modules. We let $\text{Mod}_{W_n(R)}^{\text{perf}}$ denote the full subcategory of $\text{Mod}_{W_n(R)}^{\text{Fr}}$ spanned by the perfect Frobenius modules over $W_n(R)$.

Remark 9.1.2. In the special case $n = 1$, Definition 9.1.1 reduces to Definitions 2.1.1 and 3.2.1. In particular, we have an equivalence of categories $\text{Mod}_{W_1(R)}^{\text{Fr}} \simeq \text{Mod}_R^{\text{Fr}}$, which restricts to an equivalence $\text{Mod}_{W_1(R)}^{\text{perf}} \simeq \text{Mod}_R^{\text{perf}}$.

Remark 9.1.3. Let R be a commutative ring in which $p = 0$ and let $n \geq 0$. Then $\text{Mod}_{W_n(R)}^{\text{Fr}}$ can be identified with the category of left modules over the noncommutative ring $W_n(R)[F]$ whose elements are finite sums $\sum_{i \geq 0} c_i F^i$, where each coefficient c_i belongs to $W_n(R)$, with multiplication given by

$$\left(\sum_{i \geq 0} c_i F^i\right) \left(\sum_{j \geq 0} c'_j F^j\right) = \sum_{k \geq 0} \left(\sum_{i+j=k} c_i F^i(c'_j)\right) F^k.$$

In particular, $\text{Mod}_{W_n(R)}^{\text{Fr}}$ is an abelian category with enough projective objects and enough injective objects.

Remark 9.1.4. In the situation of Definition 9.1.1, the full subcategory

$$\text{Mod}_{W_n(R)}^{\text{perf}} \subseteq \text{Mod}_{W_n(R)}^{\text{Fr}}$$

is closed under limits, colimits, and extensions. In particular, $\text{Mod}_{W_n(R)}^{\text{perf}}$ is an abelian category, and the inclusion functor $\text{Mod}_{W_n(R)}^{\text{perf}} \hookrightarrow \text{Mod}_{W_n(R)}^{\text{Fr}}$ is exact.

Remark 9.1.5. For each $n > 0$, we can identify $\text{Mod}_{W_{n-1}(R)}^{\text{Fr}}$ with the full subcategory of $\text{Mod}_{W_n(R)}^{\text{Fr}}$ spanned by those objects (M, φ_M) where M is annihilated by the kernel of the restriction map $W_n(R) \rightarrow W_{n-1}(R)$. We therefore obtain (exact) fully faithful embeddings

$$\text{Mod}_R^{\text{Fr}} \simeq \text{Mod}_{W_1(R)}^{\text{Fr}} \hookrightarrow \text{Mod}_{W_2(R)}^{\text{Fr}} \hookrightarrow \text{Mod}_{W_3(R)}^{\text{Fr}} \hookrightarrow \dots$$

Similarly, we have fully faithful embeddings

$$\mathrm{Mod}_R^{\mathrm{perf}} \simeq \mathrm{Mod}_{W_1(R)}^{\mathrm{perf}} \hookrightarrow \mathrm{Mod}_{W_2(R)}^{\mathrm{perf}} \hookrightarrow \mathrm{Mod}_{W_3(R)}^{\mathrm{perf}} \hookrightarrow \cdots$$

Remark 9.1.6. In the situation of Definition 9.1.1, the inclusion $\mathrm{Mod}_{W_n(R)}^{\mathrm{perf}} \hookrightarrow \mathrm{Mod}_{W_n(R)}^{\mathrm{Fr}}$ admits a left adjoint. Concretely, this left adjoint carries a Frobenius module M to the direct limit of the sequence

$$M \xrightarrow{\varphi_M} M^{F^{-1}} \xrightarrow{\varphi_M} M^{F^{-2}} \rightarrow \cdots ;$$

here $M^{F^{-k}}$ denotes the $W_n(R)$ -module obtained from M by restriction of scalars along the ring homomorphism $F^k : W_n(R) \rightarrow W_n(R)$. We will denote this direct limit by M^{1/p^∞} and refer to it as the *perfection* of M . Note that when $n = 1$, this agrees with the construction of Notation 3.2.3.

Example 9.1.7. Let R be a commutative \mathbf{F}_p -algebra. For each $n \geq 0$, we can regard $M = W_n(R)$ as a Frobenius module over itself by taking φ_M to be the Witt vector Frobenius map $F : W_n(R) \rightarrow W_n(R)$. Then the perfection M^{1/p^∞} can be identified with $W_n(R^{1/p^\infty})$.

9.2. Functoriality

If $f : A \rightarrow B$ is a homomorphism of \mathbf{F}_p -algebras, then there is an evident forgetful functor $\mathrm{Mod}_{W_n(B)}^{\mathrm{Fr}} \rightarrow \mathrm{Mod}_{W_n(A)}^{\mathrm{Fr}}$. This functor admits a left adjoint $f_{\mathrm{Fr}}^* : \mathrm{Mod}_{W_n(A)}^{\mathrm{Fr}} \rightarrow \mathrm{Mod}_{W_n(B)}^{\mathrm{Fr}}$, given by extension of scalars along the evident ring homomorphism $W_n(A)[F] \rightarrow W_n(B)[F]$. Since the natural map $W_n(B) \otimes_{W_n(A)} W_n(A)[F] \rightarrow W_n(B)[F]$ is an isomorphism, we have canonical isomorphisms $f_{\mathrm{Fr}}^* M \simeq W_n(B) \otimes_{W_n(A)} M$ in the category of $W_n(B)$ -modules.

Remark 9.2.1. Let $f : A \rightarrow B$ be a homomorphism of commutative \mathbf{F}_p -algebras and let M be a Frobenius module over $W_n(B)$. Then M is perfect as a Frobenius module over $W_n(B)$ if and only if it is perfect when regarded as a Frobenius module over $W_n(A)$. Moreover, the perfection M^{1/p^∞} does not depend on whether we regard M as a Frobenius module over $W_n(B)$ or over $W_n(A)$. It follows that the diagram of forgetful functors

$$\begin{array}{ccc} \mathrm{Mod}_{W_n(B)}^{\mathrm{perf}} & \longrightarrow & \mathrm{Mod}_{W_n(B)}^{\mathrm{Fr}} \\ \downarrow & & \downarrow \\ \mathrm{Mod}_{W_n(A)}^{\mathrm{perf}} & \longrightarrow & \mathrm{Mod}_{W_n(A)}^{\mathrm{Fr}} \end{array}$$

commutes (up to canonical isomorphism).

The following result is a formal consequence of Remark 9.2.1:

Proposition 9.2.2. *Let $f : A \rightarrow B$ be a homomorphism of commutative \mathbf{F}_p -algebras. Then the forgetful functor $\text{Mod}_{W_n(B)}^{\text{perf}} \rightarrow \text{Mod}_{W_n(A)}^{\text{perf}}$ admits a left adjoint f^\diamond . Moreover, the diagram of categories*

$$\begin{array}{ccc} \text{Mod}_{W_n(A)}^{\text{Fr}} & \xrightarrow{(-)^{1/p^\infty}} & \text{Mod}_{W_n(A)}^{\text{perf}} \\ \downarrow f_{\text{Fr}}^* & & \downarrow f^\diamond \\ \text{Mod}_{W_n(B)}^{\text{Fr}} & \xrightarrow{(-)^{1/p^\infty}} & \text{Mod}_{W_n(B)}^{\text{perf}} \end{array}$$

commutes up to canonical isomorphism. More precisely, for every object $M \in \text{Mod}_A^{\text{Fr}}$, the canonical map $f^\diamond(M^{1/p^\infty}) \rightarrow (f_{\text{Fr}}^ M)^{1/p^\infty}$ is an equivalence.*

Proposition 9.2.3. *Let R be a commutative \mathbf{F}_p -algebra. For each $n \geq 0$, the restriction of scalars functor $\text{Mod}_{W_n(R^{1/p^\infty})}^{\text{perf}} \rightarrow \text{Mod}_{W_n(R)}^{\text{perf}}$ is an equivalence of categories.*

Proof. Let $f : R \rightarrow R^{1/p^\infty}$ be the tautological map. Since the restriction of scalars functor is evidently conservative, it suffices to observe that for each object $M \in \text{Mod}_{W_n(R)}^{\text{perf}}$, the unit map

$$M \rightarrow f^\diamond(M) = (W_n(R^{1/p^\infty}) \otimes_{W_n(R)} M)^{1/p^\infty}$$

is an isomorphism of (perfect) Frobenius modules over $W_n(R)$. □

Corollary 9.2.4. *Let R be an \mathbf{F}_p -algebra and let $0 \leq m \leq n$. Then the essential image of the tautological map $\text{Mod}_{W_m(R)}^{\text{perf}} \hookrightarrow \text{Mod}_{W_n(R)}^{\text{perf}}$ consists of those perfect Frobenius modules over $W_n(R)$ which are annihilated by p^m .*

Proof. By virtue of Proposition 9.2.3, we can assume without loss of generality that R is perfect. In this case, the desired result follows from Remark 9.1.5, since the kernel of the restriction map $W_n(R) \rightarrow W_m(R)$ is the principal ideal (p^m) . □

Proposition 9.2.5. *Let $f : A \rightarrow B$ be a homomorphism of perfect \mathbf{F}_p -algebras. Then the extension of scalars functor $f_{\text{Fr}}^* : \text{Mod}_{W_n(A)}^{\text{Fr}} \rightarrow \text{Mod}_{W_n(B)}^{\text{Fr}}$ carries perfect Frobenius modules over $W_n(A)$ to perfect Frobenius modules over $W_n(B)$.*

Proof. Let M be a perfect Frobenius module over A . Then the maps

$$F_B : W_n(B) \rightarrow W_n(B) \quad F_A : W_n(A) \rightarrow W_n(A) \quad \varphi_M : M \rightarrow M$$

are isomorphisms, so the induced map

$$\varphi_{f_{\text{Fr}}^* M} : W_n(B) \otimes_{W_n(A)} M \rightarrow W_n(B) \otimes_{W_n(A)} M$$

is also an isomorphism. \square

Corollary 9.2.6. *Let $f : A \rightarrow B$ be an étale morphism of \mathbf{F}_p -algebras. Then the extension of scalars functor $f_{\text{Fr}}^* : \text{Mod}_{W_n(A)}^{\text{Fr}} \rightarrow \text{Mod}_{W_n(B)}^{\text{Fr}}$ carries $\text{Mod}_{W_n(A)}^{\text{perf}}$ into $\text{Mod}_{W_n(B)}^{\text{perf}}$.*

Proof. Let M be a perfect Frobenius module over A . Then we can also regard M as a Frobenius module over A^{1/p^∞} . Since f is étale, the diagram of commutative rings

$$\begin{array}{ccc} W_n(A) & \longrightarrow & W_n(B) \\ \downarrow & & \downarrow \\ W_n(A^{1/p^\infty}) & \longrightarrow & W_n(B^{1/p^\infty}) \end{array}$$

is a pushout square by the result [16, 2.4] of van der Kallen. It follows that we can identify $f_{\text{Fr}}^* M$ with the tensor product $W_n(B^{1/p^\infty}) \otimes_{W_n(A^{1/p^\infty})} M$, which is perfect by Proposition 9.2.5. \square

9.3. The Solution Functor

We now adapt Construction 2.3.1 to the setting of Frobenius modules over the Witt vectors.

Construction 9.3.1. Let R be a commutative \mathbf{F}_p -algebra and let M be a Frobenius module over $W_n(R)$. We let $\text{Sol}(M)$ denote the functor $\text{CAlg}_R^{\text{ét}} \rightarrow \text{Mod}_{\mathbf{Z}/p^n\mathbf{Z}}$ given by the formula

$$\text{Sol}(M)(R') = \{x \in (W_n(R') \otimes_{W_n(R)} M) : \varphi_{W_n(R') \otimes_{W_n(R)} M}(x) = x\}.$$

We will refer to $\text{Sol}(M)$ as the *solution sheaf* of M .

Remark 9.3.2. In the situation of Construction 9.3.1, suppose that the action of $W_n(R)$ on M factors through the restriction map $W_n(R) \rightarrow R$,

so that M can be regarded as a Frobenius module over R (if M is perfect, this is equivalent to the requirement that $pM = 0$, by virtue of Corollary 9.2.4). Then the functor $\text{Sol}(M)$ of Construction 9.3.1 agrees with the functor $\text{Sol}(M)$ of Construction 2.3.1: this follows from the fact that the diagram of commutative rings

$$\begin{array}{ccc} W_n(R) & \longrightarrow & W_n(R') \\ \downarrow & & \downarrow \\ R & \longrightarrow & R' \end{array}$$

is a pushout square, for any étale R -algebra R' .

Our first goal is to show that the functor $\text{Sol}(M)$ of Construction 9.3.1 is actually a sheaf with respect to the étale topology on $\text{Spec}(R)$. To prove this, it will be convenient to consider the following variant of Example 2.2.5:

Notation 9.3.3. Let R be a commutative \mathbf{F}_p -algebra and let M be a module over $W_n(R)$. We let $\widetilde{M} \in \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{Z}/p^n\mathbf{Z})$ denote the sheaf given by the formula $\widetilde{M}(R') = W_n(R') \otimes_{W_n(R)} M$. Note that, when M is annihilated by the kernel of the restriction map $W_n(R) \rightarrow R$ (so that M can be regarded as an R -module), this agrees with the sheaf of $\mathbf{Z}/p^n\mathbf{Z}$ -modules introduced in Example 2.2.5.

Remark 9.3.4. Let R be a commutative \mathbf{F}_p -algebra. Then the construction $R' \mapsto W_n(R')$ induces an equivalence from the category of étale R -algebras to the category of étale $W_n(R)$ -algebras. In particular, the category of étale sheaves on $\text{Spec}(R)$ is equivalent to the category of étale sheaves on $\text{Spec}(W_n(R))$. If M is a module over $W_n(R)$, then it determines a quasi-coherent sheaf on $\text{Spec}(W_n(R))$, which corresponds (under the preceding equivalence) to the functor $\widetilde{M} : \text{CAlg}_R^{\text{ét}} \rightarrow \text{Mod}_{\mathbf{Z}/p^n\mathbf{Z}}$ of Notation 9.3.3. In particular, the functor \widetilde{M} is always a sheaf with respect to the étale topology on $\text{CAlg}_R^{\text{ét}}$.

Remark 9.3.5. In the situation of Notation 9.3.3, suppose that R is perfect and that M is flat as a module over $\mathbf{Z}/p^n\mathbf{Z}$. Then, for each $R' \in \text{CAlg}_R^{\text{ét}}$, the abelian group $\widetilde{M}(R')$ is also flat as a $\mathbf{Z}/p^n\mathbf{Z}$ -module. In particular, the sheaf \widetilde{M} is flat over $\mathbf{Z}/p^n\mathbf{Z}$.

If M is a Frobenius module over $W_n(R)$, then the Frobenius map φ_M determines an endomorphism of the associated étale sheaf \widetilde{M} . By construction, we have an exact sequence of presheaves

$$0 \rightarrow \text{Sol}(M) \rightarrow \widetilde{M} \xrightarrow{\text{id} - \varphi_M} \widetilde{M}.$$

It follows that $\text{Sol}(M)$ is always a sheaf with respect to the étale topology. We may therefore regard the construction $M \mapsto \text{Sol}(M)$ as a functor from the category of Frobenius modules over $W_n(R)$ to the category of p^n -torsion sheaves on $\text{Spec}(R)$. We will denote this functor by

$$\text{Sol} : \text{Mod}_{W_n(R)}^{\text{Fr}} \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{Z}/p^n\mathbf{Z})$$

and refer to it as the *solution sheaf functor*.

Proposition 9.3.6. *Let R be a commutative \mathbf{F}_p -algebra and let M be an injective object of the abelian category $\text{Mod}_{W_n(R)}^{\text{perf}}$. Then we have a short exact sequence $0 \rightarrow \text{Sol}(M) \rightarrow \widetilde{M} \xrightarrow{\text{id} - \varphi_M} \widetilde{M} \rightarrow 0$ in the category of abelian presheaves on $\text{CAlg}_R^{\text{ét}}$.*

The proof of Proposition 9.3.6 is based on the following:

Lemma 9.3.7. *Let R be an \mathbf{F}_p -algebra and let M be an injective object of the abelian category $\text{Mod}_{W_n(R)}^{\text{perf}}$. Then M is free when regarded as a module over $\mathbf{Z}/p^n\mathbf{Z}$.*

Proof. Choose a collection of elements $\{x_i\}_{i \in I}$ of M , whose images form a basis for M/pM as a vector space over \mathbf{F}_p . Then the elements x_i determine a map of $\mathbf{Z}/p^n\mathbf{Z}$ -modules $f : \bigoplus_{i \in I} \mathbf{Z}/p^n\mathbf{Z} \rightarrow M$. The map f is surjective by virtue of Nakayama’s lemma; we will complete the proof by showing that it is injective. Assume otherwise: then there exists some nonzero element $\vec{c} \in \ker(f)$, which we can identify with a collection of elements $\{c_i\}_{i \in I}$ of $\mathbf{Z}/p^n\mathbf{Z}$ (almost all of which vanish). Let us assume that \vec{c} has been chosen so that the ideal $(c_i)_{i \in I} \subseteq \mathbf{Z}/p^n\mathbf{Z}$ is as large as possible. Since the elements x_i have images in M/pM which are linearly independent over \mathbf{F}_p , we must have $(c_i)_{i \in I} \neq \mathbf{Z}/p^n\mathbf{Z}$. It follows that we can write $\vec{c} = p\vec{b}$ for some element $\vec{b} \in \bigoplus_{i \in I} \mathbf{Z}/p^n\mathbf{Z}$. Then $pf(\vec{b}) = f(\vec{c}) = 0$, so there is a unique map of (perfect) Frobenius modules $g : R^{1/p^\infty}[F^{\pm 1}] \rightarrow M$ satisfying $g(1) = f(\vec{b})$.

Let $W_n(R^{1/p^\infty})[F^{\pm 1}]$ denote the perfection of the Frobenius module $W_n(R)[F]$. Note that multiplication by p^{n-1} induces a monomorphism

$$R^{1/p^\infty}[F^{\pm 1}] \rightarrow W_n(R^{1/p^\infty})[F^{\pm 1}].$$

Invoking our assumption that M is injective, we conclude that g factors as a composition $R^{1/p^\infty}[F^{\pm 1}] \xrightarrow{p^{n-1}} W_n(R^{1/p^\infty})[F^{\pm 1}] \xrightarrow{h} M$. Since f is surjective, we can write $h(1) = f(\vec{a})$ for some element $\vec{a} \in \bigoplus_{i \in I} \mathbf{Z}/p^n\mathbf{Z}$. We then have

$$f(\vec{b} - p^{n-1}\vec{a}) = f(\vec{b}) - p^{n-1}f(\vec{a}) = g(1) - p^{n-1}h(1) = 0,$$

so that $\vec{b} - p^{n-1}\vec{a}$ belongs to $\ker(f)$. However, the ideal generated by the coefficients of $\vec{b} - p^{n-1}\vec{a}$ is strictly larger than the ideal $(c_i)_{i \in I}$, which contradicts our choice of \vec{c} . \square

Proof of Proposition 9.3.6. Let M be an injective object of the abelian category $\text{Mod}_{W_n(R)}^{\text{perf}}$; we wish to show that the map

$$\text{id} - \varphi_M : \widetilde{M} \rightarrow \widetilde{M}$$

is a epimorphism of $\mathbf{Z}/p^n\mathbf{Z}$ -valued presheaves. We proceed by induction on n . The case $n = 0$ is vacuous and the case $n = 1$ follows from Lemma 7.1.2, so we may assume that $n \geq 2$. For each $k \geq 0$, let $M[p^k]$ denote the kernel of the map $p^k : M \rightarrow M$. Write $n = i + j$, for some positive integers i and j . Since M is injective, Lemma 9.3.7 implies that we have a short exact sequence of (perfect) Frobenius modules

$$0 \rightarrow M[p^i] \rightarrow M \xrightarrow{p^i} M[p^j] \rightarrow 0.$$

Applying the construction $N \mapsto \widetilde{N}$, we obtain a commutative diagram of short exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & \widetilde{M[p^i]} & \longrightarrow & \widetilde{M} & \xrightarrow{p^i} & \widetilde{M[p^j]} \longrightarrow 0 \\ & & \downarrow \text{id} - \varphi_{M[p^i]} & & \downarrow \text{id} - \varphi_M & & \downarrow \text{id} - \varphi_{M[p^j]} \\ 0 & \longrightarrow & \widetilde{M[p^i]} & \longrightarrow & \widetilde{M} & \xrightarrow{p^i} & \widetilde{M[p^j]} \longrightarrow 0 \end{array}$$

in the category of presheaves of abelian groups on $\text{CALg}_R^{\text{ét}}$. Since M is an injective object of $\text{Mod}_{W_n(R)}^{\text{perf}}$, the submodules $M[p^i]$ and $M[p^j]$ are injective objects of $\text{Mod}_{W_i(R)}^{\text{perf}}$ and $\text{Mod}_{W_j(R)}^{\text{perf}}$, respectively (this follows from Corollary 9.2.4, since the inclusion functors $\text{Mod}_{W_i(R)}^{\text{perf}} \hookrightarrow \text{Mod}_{W_n(R)}^{\text{perf}} \hookleftarrow \text{Mod}_{W_j(R)}^{\text{perf}}$ are exact). Applying our inductive hypothesis, we deduce that the outer vertical maps in the preceding diagram are epimorphisms, so that the middle vertical map is also an epimorphism (by the snake lemma). \square

9.4. Derived Solution Functors

For every commutative \mathbf{F}_p -algebra R and every integer $n \geq 0$, the solution functor $\text{Sol} : \text{Mod}_{W_n(R)}^{\text{perf}} \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{Z}/p^n\mathbf{Z})$ is left exact, and therefore

admits right derived functors

$$\mathrm{Sol}^m : \mathrm{Mod}_{W_n(R)}^{\mathrm{perf}} \rightarrow \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(R), \mathbf{Z}/p^n\mathbf{Z})$$

for $m \geq 0$. These functors are described by the following generalization of Proposition 7.1.1:

Proposition 9.4.1. *Let R be a commutative \mathbf{F}_p -algebra and let M be a perfect Frobenius module over $W_n(R)$. Then we have a canonical short exact sequence*

$$0 \rightarrow \mathrm{Sol}^0(M) \rightarrow \widetilde{M} \xrightarrow{\mathrm{id} - \varphi_M} \widetilde{M} \rightarrow \mathrm{Sol}^1(M) \rightarrow 0,$$

and the sheaves $\mathrm{Sol}^m(M)$ vanish for $m \geq 2$.

Proof. Choose an injective resolution $0 \rightarrow M \rightarrow Q^0 \rightarrow Q^1 \rightarrow \cdots$ in the abelian category $\mathrm{Mod}_{W_n(R)}^{\mathrm{perf}}$. Using Proposition 9.3.6, we obtain a short exact sequence of cochain complexes

$$0 \rightarrow \mathrm{Sol}(Q^*) \rightarrow \widetilde{Q}^* \xrightarrow{\mathrm{id} - \varphi} \widetilde{Q}^* \rightarrow 0.$$

Since the construction $N \mapsto \widetilde{N}$ is exact, the chain complex \widetilde{Q}^* is an acyclic resolution of \widetilde{M} . The associated long exact sequence now supplies the desired isomorphisms. \square

Remark 9.4.2. Let R be a commutative \mathbf{F}_p -algebra and let M be a perfect Frobenius module over $W_n(R)$. Then M can also be regarded as a perfect Frobenius module over $W_m(R)$ for $m \geq n$. The étale sheaf $\mathrm{Sol}^i(M)$ depends *a priori* on whether we choose to regard M as an object of the abelian category $\mathrm{Mod}_{W_n(R)}^{\mathrm{perf}}$ (in which case $\mathrm{Sol}^i(M)$ is defined as sheaf of $\mathbf{Z}/p^n\mathbf{Z}$ -modules on $\mathrm{Spec}(R)$), or as an object of the larger abelian category $\mathrm{Mod}_{W_m(R)}^{\mathrm{perf}}$ (in which case $\mathrm{Sol}^i(M)$ is defined as a sheaf of $\mathbf{Z}/p^m\mathbf{Z}$ -modules on $\mathrm{Spec}(R)$). However, Proposition 9.4.1 shows that the resulting étale sheaves are canonically isomorphic.

We will also need a generalization of Corollary 6.4.2:

Proposition 9.4.3. *Let R be a commutative \mathbf{F}_p -algebra and let Q be an injective object of $\mathrm{Mod}_{W_n(R)}^{\mathrm{perf}}$. Then $\mathrm{Sol}(Q)$ is an injective object of $\mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(R), \mathbf{Z}/p^n\mathbf{Z})$.*

The proof of Proposition 9.4.3 will require the following:

Lemma 9.4.4. *Let R be a commutative \mathbf{F}_p -algebra, let M be a $W_n(R)$ -module which is flat over $\mathbf{Z}/p^n\mathbf{Z}$, and let $\mathcal{F} \in \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$. Then the canonical map*

$$\text{Ext}_{\mathbf{F}_p}^1(\mathcal{F}, \widetilde{M}[p]) \rightarrow \text{Ext}_{\mathbf{Z}/p^n\mathbf{Z}}^1(\mathcal{F}, \widetilde{M})$$

is bijective.

Proof. Suppose we are given an extension $0 \rightarrow \widetilde{M} \rightarrow \mathcal{G} \rightarrow \mathcal{F} \rightarrow 0$ in the abelian category $\text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{Z}/p^n\mathbf{Z})$. We wish to show that there exists a commutative diagram of short exact sequences

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \widetilde{M}[p] & \longrightarrow & \mathcal{G}' & \longrightarrow & \mathcal{F} & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow \text{id} & & \\ 0 & \longrightarrow & \widetilde{M} & \longrightarrow & \mathcal{G} & \longrightarrow & \mathcal{F} & \longrightarrow & 0, \end{array}$$

where \mathcal{G}' is annihilated by p , and that the extension class of the upper exact sequence is uniquely determined. The uniqueness is clear: note that if such a diagram exists, then it induces an isomorphism $\mathcal{G}' \simeq \mathcal{G}[p] = \ker(p : \mathcal{G} \rightarrow \mathcal{G})$. To prove existence, it will suffice to show that the composite map $\mathcal{G}[p] \rightarrow \mathcal{G} \rightarrow \mathcal{F}$ is an epimorphism. To prove this, we note that the commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{G}[p] & \longrightarrow & \mathcal{G} & \longrightarrow & p\mathcal{G} & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathcal{F} & \xrightarrow{\text{id}} & \mathcal{F} & \longrightarrow & 0 & \longrightarrow & 0 \end{array}$$

yields a long exact sequence

$$\widetilde{M} \rightarrow \mathcal{G} / \mathcal{G}[p] \rightarrow \text{coker}(\mathcal{G}[p] \rightarrow \mathcal{F}) \rightarrow \text{coker}(\mathcal{G} \rightarrow \mathcal{F}),$$

where the last term vanishes (since the map $\mathcal{G} \rightarrow \mathcal{F}$ is an epimorphism). We are therefore reduced to showing that the canonical map $\widetilde{M} \rightarrow \mathcal{G} / \mathcal{G}[p]$ is an epimorphism. Since \mathcal{F} is annihilated by p , the map $p : \mathcal{G} \rightarrow \mathcal{G}$ induces a monomorphism $v : \mathcal{G} / \mathcal{G}[p] \rightarrow \widetilde{M}$. It will therefore suffice to show that the image of v is contained in the image of the map $p : \widetilde{M} \rightarrow \widetilde{M}$. This follows from Remark 9.3.5, since $\text{im}(v)$ is annihilated by p^{n-1} . \square

Proof of Proposition 9.4.3. Let Q be an injective object of $\text{Mod}_{W_n(R)}^{\text{perf}}$. We wish to show that $\text{Sol}(Q)$ is an injective object of $\text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{Z}/p^n\mathbf{Z})$: that is, that the group $\text{Ext}_{\mathbf{Z}/p^n\mathbf{Z}}^1(\mathcal{F}, \text{Sol}(Q))$ vanishes for every sheaf $\mathcal{F} \in$

$\mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(R), \mathbf{Z}/p^n\mathbf{Z})$. Since the collection of those objects \mathcal{F} for which the group $\mathrm{Ext}_{\mathbf{Z}/p^n\mathbf{Z}}^1(\mathcal{F}, \mathrm{Sol}(Q))$ vanishes is closed under extensions, we may assume without loss of generality that \mathcal{F} is annihilated by p .

By virtue of Proposition 9.3.6, we have short exact sequences of étale sheaves

$$\begin{aligned} 0 \rightarrow \mathrm{Sol}(Q[p]) \rightarrow \widetilde{Q}[p] \xrightarrow{\mathrm{id} - \varphi_{Q[p]}} \widetilde{Q}[p] \rightarrow 0 \\ 0 \rightarrow \mathrm{Sol}(Q) \rightarrow \widetilde{Q} \xrightarrow{\mathrm{id} - \varphi_Q} \widetilde{Q} \rightarrow 0 \end{aligned}$$

which supply a commutative diagram of long exact sequences

$$\begin{array}{ccc} \mathrm{Ext}_{\mathbf{F}_p}^0(\mathcal{F}, \widetilde{Q}[p]) & \xrightarrow{\alpha} & \mathrm{Ext}_{\mathbf{Z}/p^n\mathbf{Z}}^0(\mathcal{F}, \widetilde{Q}) \\ \downarrow \mathrm{id} - \varphi_{Q[p]} & & \downarrow \mathrm{id} - \varphi_Q \\ \mathrm{Ext}_{\mathbf{F}_p}^0(\mathcal{F}, \widetilde{Q}[p]) & \xrightarrow{\alpha} & \mathrm{Ext}_{\mathbf{Z}/p^n\mathbf{Z}}^0(\mathcal{F}, \widetilde{Q}) \\ \downarrow & & \downarrow \\ \mathrm{Ext}_{\mathbf{F}_p}^1(\mathcal{F}, \mathrm{Sol}(Q[p])) & \xrightarrow{\beta} & \mathrm{Ext}_{\mathbf{Z}/p^n\mathbf{Z}}^1(\mathcal{F}, \mathrm{Sol}(Q)) \\ \downarrow & & \downarrow \\ \mathrm{Ext}_{\mathbf{F}_p}^1(\mathcal{F}, \widetilde{Q}[p]) & \xrightarrow{\gamma} & \mathrm{Ext}_{\mathbf{Z}/p^n\mathbf{Z}}^1(\mathcal{F}, \widetilde{Q}) \\ \downarrow \mathrm{id} - \varphi_{Q[p]} & & \downarrow \mathrm{id} - \varphi_Q \\ \mathrm{Ext}_{\mathbf{F}_p}^1(\mathcal{F}, \widetilde{Q}[p]) & \xrightarrow{\gamma} & \mathrm{Ext}_{\mathbf{Z}/p^n\mathbf{Z}}^1(\mathcal{F}, \widetilde{Q}). \end{array}$$

The map α is obviously an isomorphism, and γ is an isomorphism by virtue of Lemma 9.4.4. It follows that β is also an isomorphism. We are therefore reduced to proving that the group $\mathrm{Ext}_{\mathbf{F}_p}^1(\mathcal{F}, \mathrm{Sol}(Q[p]))$ vanishes. In fact, we claim that $\mathrm{Sol}(Q[p])$ is an injective object of the abelian category $\mathrm{Shv}_{\acute{e}t}(\mathrm{Spec}(R), \mathbf{F}_p)$: this is a special case of Corollary 6.4.2, since $Q[p]$ is an injective object of the abelian category $\mathrm{Mod}_R^{\mathrm{perf}}$. \square

9.5. Algebraic Frobenius Modules over $W_n(R)$

Let R be a commutative \mathbf{F}_p -algebra and let M be a perfect Frobenius module over $W_n(R)$. We let $M[p]$ and M/pM denote the kernel and cokernel of the map $p : M \rightarrow M$. Then $M[p]$ and M/pM are perfect Frobenius modules

over $W_n(R)$ which are annihilated by p , and can therefore be identified with perfect Frobenius modules over R (Corollary 9.2.4).

Proposition 9.5.1. *Let R be a commutative \mathbf{F}_p -algebra and let M be a perfect Frobenius module over $W_n(R)$. The following conditions are equivalent:*

- (1) *The quotient $M/pM \in \text{Mod}_R^{\text{perf}}$ is algebraic, in the sense of Definition 2.4.1.*
- (2) *The submodule $M[p] \in \text{Mod}_R^{\text{perf}}$ is algebraic, in the sense of Definition 2.4.1.*
- (3) *Every element $x \in M$ satisfies an equation of the form*

$$\varphi_M^k(x) + a_1\varphi_M^{k-1}(x) + \cdots + a_kx = 0$$

for some coefficients $a_i \in W_n(R)$.

Definition 9.5.2. Let R be a commutative \mathbf{F}_p -algebra and let M be a perfect Frobenius module over $W_n(R)$. We will say that M is *algebraic* if it satisfies the equivalent conditions of Proposition 9.5.1. We let $\text{Mod}_{W_n(R)}^{\text{alg}}$ denote the full subcategory of $\text{Mod}_{W_n(R)}^{\text{perf}}$ spanned by the algebraic Frobenius modules over $W_n(R)$.

Remark 9.5.3. In the situation of Definition 9.5.2, an object $M \in \text{Mod}_{W_n(R)}^{\text{Fr}}$ is algebraic if and only if it is algebraic when viewed as a Frobenius module over $W_m(R)$, for any $m \geq n$.

Remark 9.5.4. In the situation of Definition 9.5.2, let M be a perfect Frobenius module over $W_n(R)$ which is annihilated by p . Then M can be regarded as a perfect Frobenius module over R (Corollary 9.2.4). Moreover, M is algebraic in the sense of Definition 9.5.2 if and only if it is algebraic in the sense of Definition 2.4.1.

Proof of Proposition 9.5.1. The implication (3) \Rightarrow (2) is obvious. We now show that (2) \Rightarrow (1). Assume that M is a perfect Frobenius module over $W_n(R)$ and that the p -torsion submodule $M[p]$ is algebraic (as a perfect Frobenius module over R). For each integer $i \geq 0$, we have a short exact sequence

$$0 \rightarrow (M[p] \cap p^i M)/(M[p] \cap p^{i+1} M) \rightarrow p^i M/p^{i+1} M \xrightarrow{p} p^{i+1} M/p^{i+2} M \rightarrow 0$$

Since the collection of algebraic objects of $\text{Mod}_R^{\text{perf}}$ is closed under the formation of subobjects and quotient objects (Proposition 4.2.4), condition (2) guarantees that each $(M[p] \cap p^i M)/(M[p] \cap p^{i+1} M)$ is algebraic. Since the

collection of algebraic objects of $\text{Mod}_R^{\text{perf}}$ is closed under extensions (Proposition 4.2.4), it follows by descending induction on i that each $p^i M/p^{i+1} M$ is algebraic. Taking $i = 0$, we deduce that (1) is satisfied.

We now complete the proof by showing that (1) implies (3). We proceed by induction on n , the case $n = 0$ being trivial. Assume that $n > 0$ and let x be an element of M having image $\bar{x} \in M/pM$. Condition (1) guarantees that we can find an element $\bar{\mu} = F^m + \bar{a}_1 F^{m-1} + \cdots + \bar{a}_{m-1} F + \bar{a}_m \in R[F]$ such that $\bar{\mu}(\bar{x}) = 0$. Lift $\bar{\mu}$ to an element $\mu = F^m + a_1 F^{m-1} + \cdots + a_m \in W_n(R)[F]$, so that $\mu(x) \in pM$. Note that pM/p^2M is a quotient of M/pM , and is therefore algebraic by virtue of Proposition 4.2.4. The Frobenius module pM is annihilated by p^{n-1} , and can therefore be regarded as a perfect Frobenius module over $W_{n-1}(R)$ by virtue of Corollary 9.2.4. Applying our inductive hypothesis, we deduce that there exists an expression $\nu = F^{m'} + b_1 F^{m'-1} + \cdots + b_{m'-1} F + b_{m'} \in W_n(R)[F]$ such that $\nu(\mu(x)) = 0$, so that x is annihilated by $\nu\mu \in W_n(R)[F]$. \square

We have the following generalization of Proposition 4.2.4:

Proposition 9.5.5. *Let R be a commutative \mathbf{F}_p -algebra and let $n \geq 0$. Then $\text{Mod}_{W_n(R)}^{\text{alg}}$ is a localizing subcategory of $\text{Mod}_{W_n(R)}^{\text{perf}}$. That is:*

- (a) *Given a short exact sequence $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ of perfect Frobenius modules over R , M is algebraic if and only if M' and M'' are algebraic.*
- (b) *The collection of algebraic Frobenius modules over $W_n(R)$ is closed under (possibly infinite) direct sums.*

Proof. We will prove (a); assertion (b) is immediate from the definitions. Suppose we are given an exact sequence $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ of perfect Frobenius modules over $W_n(R)$. Then we also have an exact sequence $M'/pM' \rightarrow M/pM \rightarrow M''/pM'' \rightarrow 0$. If M'/pM' is algebraic, then Proposition 4.2.4 implies that M/pM is algebraic if and only if M''/pM'' is algebraic. Using characterization (1) of Proposition 9.5.1, we conclude that if M' is algebraic, then M is algebraic if and only if M'' is algebraic. Applying the same argument to the exact sequence $0 \rightarrow M'[p] \rightarrow M[p] \rightarrow M''[p]$ (and using characterization (2) of Proposition 9.5.1), we deduce that if M'' is algebraic, then M is algebraic if and only if M' is algebraic. \square

Proposition 9.5.6. *Let R be a commutative \mathbf{F}_p -algebra, let $n \geq 0$, and let M be an algebraic Frobenius module over $W_n(R)$. Then the étale sheaves $\text{Sol}^i(M) \in \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{Z}/p^n\mathbf{Z})$ vanish for $i \neq 0$.*

Proof. We prove the following assertion by induction on m :

($*_m$) Let M be an algebraic Frobenius module over $W_n(R)$ which is annihilated by p^m . Then $\text{Sol}^i(M) = 0$.

Note that assertion ($*_0$) is trivial, and assertion ($*_n$) implies Proposition 9.5.6. It will therefore suffice to show that ($*_m$) implies ($*_{m+1}$). Note that if M is an algebraic Frobenius module which is annihilated by p^{m+1} , then the short exact sequence $0 \rightarrow M[p] \rightarrow M \rightarrow pM \rightarrow 0$ yields an exact sequence of sheaves $\text{Sol}^i(M[p]) \rightarrow \text{Sol}^i(M) \rightarrow \text{Sol}^i(pM)$. Here pM and $M[p]$ are also algebraic (Proposition 9.5.5), and pM is annihilated by p^m . Our inductive hypothesis then guarantees that $\text{Sol}^i(pM) = 0$. To complete the proof, it will suffice to show that $\text{Sol}^i(M[p]) = 0$. To prove this, we can replace $W_n(R)$ by R (Remark 9.4.2), in which case the desired result follows from Corollary 7.4.4. \square

9.6. The Riemann-Hilbert Correspondence for $\mathbf{Z}/p^n\mathbf{Z}$ -Sheaves

We can now formulate the main result of this section:

Theorem 9.6.1 (Riemann-Hilbert Correspondence). *Let R be a commutative \mathbf{F}_p -algebra and let $n \geq 0$. Then the functor $\text{Sol} : \text{Mod}_{W_n(R)}^{\text{perf}} \rightarrow \text{Shv}_{\acute{e}t}(\text{Spec}(R), \mathbf{Z}/p^n\mathbf{Z})$ induces an equivalence of categories $\text{Mod}_{W_n(R)}^{\text{alg}} \rightarrow \text{Shv}_{\acute{e}t}(\text{Spec}(R), \mathbf{Z}/p^n\mathbf{Z})$.*

We will deduce Theorem 9.6.1 from the following comparison result:

Proposition 9.6.2. *Let R be a commutative \mathbf{F}_p -algebra and let M and N be perfect Frobenius modules over $W_n(R)$. Assume that M is algebraic and that $\text{Sol}^1(N) \simeq 0$. Then the canonical map*

$$\text{Ext}_{W_n(R)[F]}^i(M, N) \rightarrow \text{Ext}_{\mathbf{Z}/p^n\mathbf{Z}}^i(\text{Sol}(M), \text{Sol}(N))$$

is an isomorphism for $i \geq 0$.

Proof of Theorem 9.6.1 from Proposition 9.6.2. We first claim that the composite functor

$$\text{Mod}_{W_n(R)}^{\text{alg}} \hookrightarrow \text{Mod}_{W_n(R)}^{\text{perf}} \xrightarrow{\text{Sol}} \text{Shv}_{\acute{e}t}(\text{Spec}(R), \mathbf{Z}/p^n\mathbf{Z})$$

is fully faithful. Let M and N be algebraic Frobenius modules over $W_n(R)$; we wish to show that the canonical map

$$\text{Hom}_{W_n(R)[F]}(M, N) \rightarrow \text{Hom}_{\mathbf{F}_p}(\text{Sol}(M), \text{Sol}(N))$$

is an isomorphism. This is a special case of Proposition 9.6.2, since $\mathrm{Sol}^1(N) \simeq 0$ by virtue of Proposition 9.5.6.

Let $\mathcal{C} \subseteq \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(R), \mathbf{Z}/p^n\mathbf{Z})$ denote the full subcategory spanned by those sheaves of the form $\mathrm{Sol}(M)$, where M is an algebraic Frobenius module over $W_n(R)$. To complete the proof of Theorem 9.6.1, it will suffice to show that every object of $\mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spec}(R), \mathbf{Z}/p^n\mathbf{Z})$ belongs to \mathcal{C} . Note that Theorem 2.4.3 guarantees \mathcal{C} contains every sheaf of $\mathbf{Z}/p\mathbf{Z}$ -modules on $\mathrm{Spec}(R)$. We will complete the proof by showing that \mathcal{C} is closed under the formation of extensions. Suppose we are given a short exact sequence of étale sheaves

$$0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0,$$

where \mathcal{F}' and \mathcal{F}'' belong to \mathcal{C} ; we wish to show that \mathcal{F} also belongs to \mathcal{C} . Without loss of generality, we may assume that $\mathcal{F}' = \mathrm{Sol}(M')$ and $\mathcal{F}'' = \mathrm{Sol}(M'')$ for some algebraic Frobenius modules M' and M'' over $W_n(R)$. In this case, the preceding exact sequence is classified by an element $\eta \in \mathrm{Ext}_{\mathbf{Z}/p^n\mathbf{Z}}^1(\mathrm{Sol}(M''), \mathrm{Sol}(M'))$. Invoking Proposition 9.6.2 again, we deduce that η can be lifted (uniquely) to an element $\bar{\eta} \in \mathrm{Ext}_{W_n(R)[F]}^1(M'', M')$, which classifies a short exact sequence of Frobenius modules $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$. Proposition 9.5.5 guarantees that M is algebraic, so that $\mathcal{F} \simeq \mathrm{Sol}(M)$ also belongs to the category \mathcal{C} . \square

We now turn to the proof of Proposition 9.6.2. We begin with some special cases.

Lemma 9.6.3. *Let R be an \mathbf{F}_p -algebra and let M be an algebraic Frobenius module over R , and let N be any object of $\mathrm{Mod}_{W_n(R)}^{\mathrm{perf}}$. Then the canonical map*

$$\theta : \mathrm{Hom}_{W_n(R)[F]}(M, N) \rightarrow \mathrm{Hom}_{\mathbf{Z}/p^n\mathbf{Z}}(\mathrm{Sol}(M), \mathrm{Sol}(N))$$

is an isomorphism.

Proof. Since the functor Sol is left exact, we have an isomorphism $\mathrm{Sol}(N)[p] \simeq \mathrm{Sol}(N[p])$. Since M and $\mathrm{Sol}(M)$ are annihilated by p , we can identify θ with the canonical map

$$\mathrm{Hom}_{R[F]}(M, N[p]) \rightarrow \mathrm{Hom}_{\mathbf{F}_p}(\mathrm{Sol}(M), \mathrm{Sol}(N[p])).$$

We may therefore replace N by $N[p]$ and thereby reduce to the case $n = 1$. Using Theorem 2.4.3, we can choose an isomorphism $M \simeq \mathrm{RH}(\mathcal{F})$ for some

object $\mathcal{F} \in \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$. In this case, we θ has a left inverse, given by the map

$$\text{Hom}_{\mathbf{F}_p}(\text{Sol}(\text{RH}(\mathcal{F})), \text{Sol}(N)) \rightarrow \text{Hom}_{\mathbf{F}_p}(\mathcal{F}, \text{Sol}(N))$$

given by precomposition with the unit map $u : \mathcal{F} \rightarrow \text{Sol}(\text{RH}(\mathcal{F}))$. This map is an isomorphism by virtue of Proposition 7.2.1. \square

Lemma 9.6.4. *Let R be an \mathbf{F}_p -algebra and let M be an algebraic Frobenius module over R . Let N be any perfect Frobenius module over $W_n(R)$. If $\text{Sol}^1(N) \simeq 0$, then the canonical map*

$$\text{Ext}_{W_n(R)[F]}^i(M, N) \rightarrow \text{Ext}_{\mathbf{Z}/p^n\mathbf{Z}}^i(\text{Sol}(M), \text{Sol}(N))$$

is an isomorphism for $i \geq 0$.

Proof. Choose an injective resolution $0 \rightarrow N \rightarrow Q^0 \rightarrow Q^1 \rightarrow \dots$ in the abelian category $\text{Mod}_{W_n(R)}^{\text{perf}}$. Our hypothesis that $\text{Sol}^1(N)$ vanishes guarantees that the complex $0 \rightarrow \text{Sol}(N) \rightarrow \text{Sol}(Q^0) \rightarrow \text{Sol}(Q^1) \rightarrow \dots$ is exact in the abelian category $\text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{Z}/p^n\mathbf{Z})$ (Proposition 9.4.1). Moreover, each $\text{Sol}(Q^i)$ is an injective object of $\text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{Z}/p^n\mathbf{Z})$ (Proposition 9.4.3). It will therefore suffice to show that the canonical map

$$\text{Hom}_{W_n(R)[F]}(M, Q^*) \rightarrow \text{Hom}_{\mathbf{Z}/p^n\mathbf{Z}}(\text{Sol}(M), \text{Sol}(Q^*))$$

is a quasi-isomorphism of chain complexes. In fact, this map is an isomorphism of chain complexes: this is a special case of Lemma 9.6.3. \square

Proof of Proposition 9.6.2. Let N be a perfect Frobenius module over $W_n(R)$, and suppose that $\text{Sol}^1(N) \simeq 0$. Let us say that an object $M \in \text{Mod}_{W_n(R)}^{\text{alg}}$ is *good* if the canonical map

$$\rho_i : \text{Ext}_{W_n(R)[F]}^i(M, N) \rightarrow \text{Ext}_{\mathbf{Z}/p^n\mathbf{Z}}^i(\text{Sol}(M), \text{Sol}(N))$$

is an isomorphism for $i \geq 0$. It follows from Lemma 9.6.4 that if $M \in \text{Mod}_{W_n(R)}^{\text{alg}}$ is annihilated by p , then M is good. We wish to show that every object of $\text{Mod}_{W_n(R)}^{\text{alg}}$ is good. For this, it will suffice to establish the following:

- (*) Let $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ be a short exact sequence of algebraic Frobenius modules over $W_n(R)$. If M' and M'' are good, then M is also good.

To prove (*), we note that the vanishing of $\mathrm{Sol}^1(M')$ (Proposition 9.5.6) guarantees the exactness of the sequence $0 \rightarrow \mathrm{Sol}(M') \rightarrow \mathrm{Sol}(M) \rightarrow \mathrm{Sol}(M'') \rightarrow 0$. It follows that each ρ_i fits into a commutative diagram of exact sequences

$$\begin{array}{ccc}
 \mathrm{Ext}_{W_n(R)[F]}^{i-1}(M', N) & \xrightarrow{\rho'_{i-1}} & \mathrm{Ext}_{\underline{\mathbf{Z}/p^n\mathbf{Z}}}^{i-1}(\mathrm{Sol}(M'), \mathrm{Sol}(N)) \\
 \downarrow & & \downarrow \\
 \mathrm{Ext}_{W_n(R)[F]}^i(M'', N) & \xrightarrow{\rho''_i} & \mathrm{Ext}_{\underline{\mathbf{Z}/p^n\mathbf{Z}}}^i(\mathrm{Sol}(M''), \mathrm{Sol}(N)) \\
 \downarrow & & \downarrow \\
 \mathrm{Ext}_{W_n(R)[F]}^i(M, N) & \xrightarrow{\rho_i} & \mathrm{Ext}_{\underline{\mathbf{Z}/p^n\mathbf{Z}}}^i(\mathrm{Sol}(M), \mathrm{Sol}(N)) \\
 \downarrow & & \downarrow \\
 \mathrm{Ext}_{W_n(R)[F]}^i(M', N) & \xrightarrow{\rho'_i} & \mathrm{Ext}_{\underline{\mathbf{Z}/p^n\mathbf{Z}}}^i(\mathrm{Sol}(M'), \mathrm{Sol}(N)) \\
 \downarrow & & \downarrow \\
 \mathrm{Ext}_{W_n(R)[F]}^{i+1}(M'', N) & \xrightarrow{\rho''_{i+1}} & \mathrm{Ext}_{\underline{\mathbf{Z}/p^n\mathbf{Z}}}^{i+1}(\mathrm{Sol}(M''), \mathrm{Sol}(N)).
 \end{array}$$

Our hypothesis that M' and M'' are good guarantees that the maps ρ'_{i-1} , ρ''_i , ρ'_i , and ρ''_{i+1} are isomorphisms, so that ρ_i is also an isomorphism. \square

10. Globalization

For any commutative \mathbf{F}_p -algebra R , the Riemann-Hilbert correspondence of Theorem 1.0.2 supplies a description of the category of p -torsion étale sheaves on the affine \mathbf{F}_p -scheme $X = \mathrm{Spec}(R)$ in terms of Frobenius modules over R . Our goal in this section is to extend the Riemann-Hilbert correspondence to the case of an arbitrary \mathbf{F}_p -scheme X . We begin in §10.1 by introducing the notion of a *Frobenius sheaf* on X : that is, a quasi-coherent sheaf \mathcal{E} on X equipped with a Frobenius-semilinear endomorphism $\varphi_{\mathcal{E}}$ (Definition 10.1.2). The collection of Frobenius sheaves on X forms a category, which we will denote by $\mathrm{QCoh}_X^{\mathrm{Fr}}$. In §10.2 we construct an equivalence RH from the category $\mathrm{Shv}_{\mathrm{ét}}(X; \mathbf{F}_p)$ of p -torsion étale sheaves on X to a full subcategory $\mathrm{QCoh}_X^{\mathrm{alg}} \subseteq \mathrm{QCoh}_X^{\mathrm{Fr}}$ (Theorem 10.2.7 and Notation 10.2.10). This is essentially a formal exercise (given the earlier results of this paper): roughly speaking, the Riemann-Hilbert functor RH is constructed by amalgamating

the equivalences $\mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(U; \mathbf{F}_p) \simeq \mathrm{QCoh}_U^{\mathrm{alg}}$ where U ranges over affine open subsets of X . Consequently, any *local* question about the functor RH can be reduced to the affine case: we use this observation in §10.3 to argue that the Riemann-Hilbert correspondence is compatible with the formation of pullbacks along an arbitrary morphism of \mathbf{F}_p -schemes $f : X \rightarrow Y$ (Variant 10.3.12). However, we do encounter a genuinely new *global* phenomenon: the Riemann-Hilbert correspondence is also compatible with direct images (and higher direct images) along a morphism $f : X \rightarrow Y$ which is proper and of finite presentation (Theorem 10.5.5). We prove this in §10.5 using a global characterization for holonomic Frobenius sheaves (Theorem 10.4.1), which we establish in §10.4. In §10.6, we apply these ideas to give a proof of the proper base change theorem in étale cohomology (in the special case of p -torsion sheaves on \mathbf{F}_p -schemes; see Corollary 10.6.2).

Remark 10.0.1. Throughout this section, we confine our study of Frobenius sheaves on X to the case where X is an \mathbf{F}_p -scheme. However, the results of this section can be extended to more general geometric objects, such as algebraic spaces over \mathbf{F}_p . Similarly, the results can also be extended to have “coefficients in \mathbf{Z}/p^n ” in the sense of §9. We leave such extensions to the reader.

10.1. Frobenius Sheaves on a Scheme

We begin by introducing some terminology.

Notation 10.1.1. For any scheme X , we let QCoh_X denote the category of quasi-coherent sheaves on X . If X is an \mathbf{F}_p -scheme, we let $\varphi_X : X \rightarrow X$ denote the absolute Frobenius morphism from X to itself.

Definition 10.1.2. Let X be an \mathbf{F}_p -scheme. A *Frobenius sheaf* on X is a pair $(\mathcal{E}, \varphi_{\mathcal{E}})$, where \mathcal{E} is a quasi-coherent sheaf on X and $\varphi_{\mathcal{E}} : \mathcal{E} \rightarrow \varphi_{X*} \mathcal{E}$ is a morphism of quasi-coherent sheaves. If $(\mathcal{E}, \varphi_{\mathcal{E}})$ and $(\mathcal{F}, \varphi_{\mathcal{F}})$ are Frobenius sheaves on X , then we will say that a \mathcal{O}_X -module map $f : \mathcal{E} \rightarrow \mathcal{F}$ is a *morphism of Frobenius sheaves* if the diagram

$$\begin{array}{ccc}
 \mathcal{E} & \xrightarrow{f} & \mathcal{F} \\
 \downarrow \varphi_{\mathcal{E}} & & \downarrow \varphi_{\mathcal{F}} \\
 \varphi_{X*} \mathcal{E} & \xrightarrow{\varphi_{X*}(f)} & \varphi_{X*} \mathcal{F}
 \end{array}$$

commutes. We let $\mathrm{QCoh}_X^{\mathrm{Fr}}$ denote the category whose objects are Frobenius sheaves on X and whose morphisms are morphisms of Frobenius sheaves.

We will generally abuse terminology by identifying a Frobenius sheaf $(\mathcal{E}, \varphi_{\mathcal{E}})$ with its underlying quasi-coherent sheaf \mathcal{E} , and simply referring to \mathcal{E} as a *Frobenius sheaf on X* .

Example 10.1.3. Let $X = \text{Spec}(R)$ be an affine \mathbf{F}_p -scheme. Then the global sections functor $\mathcal{E} \mapsto \Gamma(X, \mathcal{E})$ induces an equivalence of categories $\text{QCoh}_X^{\text{Fr}} \rightarrow \text{Mod}_R^{\text{Fr}}$.

Remark 10.1.4. Let X be an \mathbf{F}_p -scheme. Then the category $\text{QCoh}_X^{\text{Fr}}$ is abelian. Moreover, the forgetful functor $\text{QCoh}_X^{\text{Fr}} \rightarrow \text{QCoh}_X$ is exact.

Variante 10.1.5. Let X be an \mathbf{F}_p -scheme. Using the adjointness of the functors φ_{X*} and φ_X^* , we can obtain a slightly different description of the category $\text{QCoh}_X^{\text{Fr}}$ of Frobenius sheaves:

- The objects of $\text{QCoh}_X^{\text{Fr}}$ can be identified with pairs $(\mathcal{E}, \psi_{\mathcal{E}})$, where \mathcal{E} is a quasi-coherent sheaf on X and $\psi_{\mathcal{E}} : \varphi_X^* \mathcal{E} \rightarrow \mathcal{E}$ is a morphism of quasi-coherent sheaves.
- A morphism from $(\mathcal{E}, \psi_{\mathcal{E}})$ to $(\mathcal{F}, \psi_{\mathcal{F}})$ in the category $\text{QCoh}_X^{\text{Fr}}$ is a morphism of quasi-coherent sheaves $f : \mathcal{E} \rightarrow \mathcal{F}$ for which the diagram

$$\begin{array}{ccc} \varphi_X^* \mathcal{E} & \xrightarrow{\varphi_X^*(f)} & \varphi_X^* \mathcal{F} \\ \downarrow \psi_{\mathcal{E}} & & \downarrow \psi_{\mathcal{F}} \\ \mathcal{E} & \xrightarrow{f} & \mathcal{F} \end{array}$$

commutes.

In what follows, we will regard quasi-coherent sheaves on a scheme X as sheaves on the étale site of X (see Example 2.2.5). Given a quasi-coherent sheaf $\mathcal{E} \in \text{QCoh}_X$ and an étale morphism $f : U \rightarrow X$, we let $\mathcal{E}(U)$ denote the abelian group of global sections $\Gamma(U, f^* \mathcal{E})$. Note that if $U = \text{Spec}(R)$ is affine, then $\mathcal{E}(U)$ has the structure of an R -module; if X is an \mathbf{F}_p -scheme and \mathcal{E} is a Frobenius sheaf, then $\mathcal{E}(U)$ inherits the structure of a Frobenius module over R .

Proposition 10.1.6. *Let X be an \mathbf{F}_p -scheme and let \mathcal{E} be a Frobenius sheaf on X . The following conditions are equivalent:*

- (1) *For every étale morphism $f : U \rightarrow X$ where $U \simeq \text{Spec}(R)$ is affine, the group of sections $\mathcal{E}(U)$ is perfect (respectively algebraic, holonomic) when regarded as a Frobenius module over R .*
- (2) *For every open subset $U \subseteq X$ where $U \simeq \text{Spec}(R)$ is affine, the group of sections \mathcal{E} is perfect (respectively algebraic, holonomic) when regarded as a Frobenius module over R .*

- (3) *There exists an étale covering $\{U_\alpha \rightarrow X\}$ where each $U_\alpha \simeq \text{Spec}(R_\alpha)$ is affine, and each $\mathcal{E}(U_\alpha)$ is perfect (respectively algebraic, holonomic) when regarded as a Frobenius module over R_α .*

Proof. The implications (1) \Rightarrow (2) \Rightarrow (3) are obvious. The implication (3) \Rightarrow (1) follows from Corollary 3.4.7 (respectively Lemma 5.4.6, Corollary 7.4.3). \square

Definition 10.1.7. Let X be an \mathbf{F}_p -scheme and let \mathcal{E} be a Frobenius sheaf on X . We will say that \mathcal{E} is *perfect* (respectively *algebraic*, *holonomic*) if it satisfies the equivalent conditions of Proposition 10.1.6. We let $\text{QCoh}_X^{\text{perf}}$ (respectively $\text{QCoh}_X^{\text{alg}}$, $\text{QCoh}_X^{\text{hol}}$) denote the full subcategory of $\text{QCoh}_X^{\text{Fr}}$ spanned by those Frobenius sheaves which are perfect (respectively algebraic, holonomic), so that we have inclusions

$$\text{QCoh}_X^{\text{hol}} \subseteq \text{QCoh}_X^{\text{alg}} \subseteq \text{QCoh}_X^{\text{perf}} \subseteq \text{QCoh}_X^{\text{Fr}}.$$

Example 10.1.8. Let $X = \text{Spec}(R)$ be an affine \mathbf{F}_p -scheme. Then a Frobenius sheaf $\mathcal{E} \in \text{QCoh}_X^{\text{Fr}}$ is perfect (respectively algebraic, holonomic) if and only if $\Gamma(X, \mathcal{E})$ is perfect (respectively algebraic, holonomic) when regarded as a Frobenius module over R .

Remark 10.1.9. Let X be an \mathbf{F}_p -scheme. Then the subcategories

$$\text{QCoh}_X^{\text{hol}} \subseteq \text{QCoh}_X^{\text{alg}} \subseteq \text{QCoh}_X^{\text{perf}} \subseteq \text{QCoh}_X^{\text{Fr}}$$

are closed under the formation of kernels, cokernels, and extensions. In particular, they are abelian subcategories of $\text{QCoh}_X^{\text{Fr}}$. Moreover, the subcategories $\text{QCoh}_X^{\text{alg}} \subseteq \text{QCoh}_X^{\text{perf}} \subseteq \text{QCoh}_X$ are closed under (possibly infinite) direct sums (and therefore under all colimits). To prove these assertions, we can work locally and thereby reduce to the case where X is affine: in this case, the desired results follow from Remark 3.2.2, Proposition 4.2.4, and Corollary 4.3.3.

Remark 10.1.10 (Descent). Let X be an \mathbf{F}_p -scheme. Then the theory of Frobenius sheaves satisfies effective descent with respect to the étale topology on X , and is therefore determined (in some sense) by its behavior when X is affine. In other words, the construction $(U \rightarrow X) \mapsto \text{QCoh}_U^{\text{Fr}}$ determines a stack on the étale site of X . The same remark applies to the subcategories $\text{QCoh}_U^{\text{hol}}$, $\text{QCoh}_U^{\text{alg}}$, and $\text{QCoh}_U^{\text{perf}}$ (by virtue of Proposition 10.1.6).

Remark 10.1.11 (Perfection). Let X be an \mathbf{F}_p -scheme and let \mathcal{E} be a Frobenius sheaf on X . We let \mathcal{E}^{1/p^∞} denote the direct limit of the diagram

$$\mathcal{E} \xrightarrow{\varphi_{\mathcal{E}}} \varphi_{X*} \mathcal{E} \xrightarrow{\varphi_{X*}(\varphi_{\mathcal{E}})} \varphi_{X*}^2 \mathcal{E} \rightarrow \dots$$

Then we have a canonical isomorphism $\mathcal{E}^{1/p^\infty} \simeq \varphi_{X*} \mathcal{E}^{1/p^\infty}$ which endows \mathcal{E}^{1/p^∞} with the structure of a perfect Frobenius sheaf on X . Moreover, the canonical map $u : \mathcal{E} \rightarrow \mathcal{E}^{1/p^\infty}$ is a morphism of Frobenius sheaves with the following universal property: for any perfect Frobenius sheaf \mathcal{F} on X , composition with u induces a bijection

$$\mathrm{Hom}_{\mathrm{QCoh}_X^{\mathrm{perf}}}(\mathcal{E}^{1/p^\infty}, \mathcal{F}) \rightarrow \mathrm{Hom}_{\mathrm{QCoh}_X^{\mathrm{Fr}}}(\mathcal{E}, \mathcal{F}).$$

In other words, we can regard the construction $\mathcal{E} \mapsto \mathcal{E}^{1/p^\infty}$ as a left adjoint to the inclusion functor $\mathrm{QCoh}_X^{\mathrm{perf}} \subseteq \mathrm{QCoh}_X^{\mathrm{Fr}}$. Note that the perfection functor $\mathcal{E} \mapsto \mathcal{E}^{1/p^\infty}$ is exact (since filtered direct limits in QCoh_X are exact; see [15, Tag 077K]).

10.2. The Riemann-Hilbert Correspondence

We now extend the Riemann-Hilbert correspondence of Theorem 1.0.2 to the case of a general \mathbf{F}_p -scheme.

Notation 10.2.1. For any scheme X , we let $\mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(X, \mathbf{F}_p)$ denote the abelian category of p -torsion sheaves on the étale site of X . If X is an \mathbf{F}_p -scheme, then we have a forgetful functor $\mathrm{QCoh}_X \rightarrow \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(X, \mathbf{F}_p)$ which carries a sheaf of \mathcal{O}_X -modules to its underlying sheaf of \mathbf{F}_p -modules. We will generally abuse notation by not distinguishing between a quasi-coherent sheaf \mathcal{E} and its image under this functor. Moreover, we will also abuse notation by identify \mathcal{E} with its direct image $\varphi_{X*} \mathcal{E}$ under the absolute Frobenius map $\varphi_X : X \rightarrow X$: note that there is a canonical isomorphism $\mathcal{E} \simeq \varphi_{X*} \mathcal{E}$ in the category $\mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(X, \mathbf{F}_p)$, though this isomorphism is not \mathcal{O}_X -linear.

Construction 10.2.2 (The Solution Functor). Let X be an \mathbf{F}_p -scheme and let $(\mathcal{E}, \varphi_{\mathcal{E}})$ be a Frobenius sheaf on X . We let $\mathrm{Sol}(\mathcal{E})$ denote the kernel of the map $(\mathrm{id} - \varphi_{\mathcal{E}}) : \mathcal{E} \rightarrow \mathcal{E}$, formed in the abelian category $\mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(X, \mathbf{F}_p)$. The construction $(\mathcal{E}, \varphi_{\mathcal{E}}) \mapsto \mathrm{Sol}(\mathcal{E})$ determines a functor $\mathrm{Sol} : \mathrm{QCoh}_X^{\mathrm{Fr}} \rightarrow \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(X, \mathbf{F}_p)$, which we will refer to as *the solution functor*.

Remark 10.2.3. In the special case where $X = \mathrm{Spec}(R)$ is affine, the solution functor of Construction 10.2.2 agrees with the solution functor of Construction 2.3.1. More precisely, for any Frobenius sheaf \mathcal{E} on X , we have a canonical isomorphism $\mathrm{Sol}(\mathcal{E}) \simeq \mathrm{Sol}(\Gamma(X, \mathcal{E}))$ in the category $\mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(X, \mathbf{F}_p)$.

Remark 10.2.4. Construction 10.2.2 is local with respect to the étale topology. More precisely, if $f : U \rightarrow X$ is an étale morphism of \mathbf{F}_p -schemes, then we have a canonical isomorphism $f^* \text{Sol}(\mathcal{E}) \simeq \text{Sol}(f^* \mathcal{E})$ for every Frobenius sheaf \mathcal{E} on X .

Remark 10.2.5. Let X be an \mathbf{F}_p -scheme and let \mathcal{E} be a Frobenius sheaf on X . Then the canonical map $\mathcal{E} \rightarrow \mathcal{E}^{1/p^\infty}$ induces an isomorphism of étale sheaves $\text{Sol}(\mathcal{E}) \rightarrow \text{Sol}(\mathcal{E}^{1/p^\infty})$. To prove this, we can reduce to the case where X is affine, in which case the desired result follows from Proposition 3.2.9.

Remark 10.2.6. Let X be an \mathbf{F}_p -scheme and let \mathcal{E} be an algebraic Frobenius sheaf on X . Then the sequence

$$0 \rightarrow \text{Sol}(\mathcal{E}) \rightarrow \mathcal{E} \xrightarrow{\text{id} - \varphi_\mathcal{E}} \mathcal{E} \rightarrow 0$$

is exact (in the abelian category $\text{Shv}_{\text{ét}}(X, \mathbf{F}_p)$). To prove this, we can work locally on X and thereby reduce to the case where X is affine, in which case the desired result follows from Propositions 7.1.1 and 7.2.1.

Theorem 10.2.7. *Let X be an \mathbf{F}_p -scheme. Then the solution functor Sol induces equivalences of abelian categories*

$$\text{QCoh}_X^{\text{alg}} \simeq \text{Shv}_{\text{ét}}(X, \mathbf{F}_p) \quad \text{QCoh}_X^{\text{hol}} \rightarrow \text{Shv}_{\text{ét}}^c(X, \mathbf{F}_p).$$

Here $\text{Shv}_{\text{ét}}^c(X, \mathbf{F}_p)$ denotes the full subcategory of $\text{Shv}_{\text{ét}}(X, \mathbf{F}_p)$ spanned by those p -torsion étale sheaves \mathcal{F} which are locally constructible (that is, for which the restriction $\mathcal{F}|_U \in \text{Shv}_{\text{ét}}(U, \mathbf{F}_p)$ is constructible for each affine open subset $U \subseteq X$).

Remark 10.2.8. If the scheme X is quasi-compact and quasi-separated, then a sheaf $\mathcal{F} \in \text{Shv}_{\text{ét}}(X, \mathbf{F}_p)$ belongs to the subcategory $\text{Shv}_{\text{ét}}^c(X, \mathbf{F}_p)$ if and only if it is constructible: that is, if and only if it becomes locally constant along some constructible stratification of X .

Proof of Theorem 10.2.7. Since the constructions

$$(U \subseteq X) \mapsto \text{QCoh}_U^{\text{hol}}, \text{QCoh}_U^{\text{alg}}, \text{Shv}_{\text{ét}}(U, \mathbf{F}_p), \text{Shv}_{\text{ét}}^c(U, \mathbf{F}_p)$$

satisfy effective descent with respect to the Zariski topology (or even the étale topology), we can reduce to the case where $X = \text{Spec}(R)$ is affine. In this case, the desired equivalences follow from Theorems 1.0.2 and 7.4.1. \square

Corollary 10.2.9. *Let X be an \mathbf{F}_p -scheme which is quasi-compact and quasi-separated. Then the inclusion functor $\text{QCoh}_X^{\text{hol}} \hookrightarrow \text{QCoh}_X^{\text{alg}}$ extends to an equivalence of categories $\text{Ind}(\text{QCoh}_X^{\text{hol}}) \simeq \text{QCoh}_X^{\text{alg}}$.*

Proof. By virtue of Theorem 10.2.7, it will suffice to show that the inclusion functor $\mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}^c(X, \mathbf{F}_p) \hookrightarrow \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(X, \mathbf{F}_p)$ extends to an equivalence $\mathrm{Ind}(\mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}^c(X, \mathbf{F}_p)) \simeq \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(X, \mathbf{F}_p)$, which follows from [15, Tag 03SA]. \square

Notation 10.2.10. Let X be an \mathbf{F}_p -scheme. We let $\mathrm{RH} : \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(X, \mathbf{F}_p) \rightarrow \mathrm{QCoh}_X^{\mathrm{alg}}$ denote an inverse of the solution functor. We will refer to RH as the *Riemann-Hilbert functor*.

Remark 10.2.11. Let X be an \mathbf{F}_p -scheme and let $\mathcal{F} \in \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(X, \mathbf{F}_p)$ be a p -torsion étale sheaf on X . Then the Frobenius sheaf $\mathrm{RH}(\mathcal{F})$ is characterized by the following universal property: for every perfect Frobenius sheaf \mathcal{E} on X , the canonical map

$$\begin{aligned} \mathrm{Hom}_{\mathrm{QCoh}_X^{\mathrm{perf}}}(\mathrm{RH}(\mathcal{F}), \mathcal{E}) &\rightarrow \mathrm{Hom}_{\mathbf{F}_p}(\mathrm{Sol}(\mathrm{RH}(\mathcal{F})), \mathrm{Sol}(\mathcal{E})) \\ &\simeq \mathrm{Hom}_{\mathbf{F}_p}(\mathcal{F}, \mathrm{Sol}(\mathcal{E})) \end{aligned}$$

is a bijection. To prove this, we can reduce to the case where X is affine, in which case the desired result follows from the properties of the Riemann-Hilbert functor given in Theorem 6.1.1.

We can summarize the situation as follows: when regarded as a functor from $\mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(X, \mathbf{F}_p)$ to $\mathrm{QCoh}_X^{\mathrm{perf}}$, the Riemann-Hilbert functor of Notation 10.2.10 is left adjoint to the solution functor $\mathrm{Sol} : \mathrm{QCoh}_X^{\mathrm{perf}} \rightarrow \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(X, \mathbf{F}_p)$.

10.3. Functoriality

We now consider the behavior of Frobenius sheaves as the \mathbf{F}_p -scheme X varies.

Construction 10.3.1 (Pullback of Frobenius Sheaves). Let $f : X \rightarrow Y$ be a morphism of \mathbf{F}_p -schemes, so that we have a commutative diagram of schemes

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow \varphi_X & & \downarrow \varphi_Y \\ X & \xrightarrow{f} & Y \end{array}$$

and therefore a canonical isomorphism $f^* \circ \varphi_Y^* \simeq \varphi_X^* \circ f^*$ in the category of functors from QCoh_Y to QCoh_X .

Let \mathcal{E} be a Frobenius sheaf on Y , and let $\psi_{\mathcal{E}} : \varphi_Y^* \mathcal{E} \rightarrow \mathcal{E}$ be as in Variant 10.1.5. We let $\psi_{f^* \mathcal{E}}$ denote the composite map

$$\varphi_X^* f^* \mathcal{E} \simeq f^* \varphi_Y^* \mathcal{E} \xrightarrow{f^* \psi_{\mathcal{E}}} f^* \mathcal{E}.$$

The construction $(\mathcal{E}, \psi_{\mathcal{E}}) \mapsto (f^* \mathcal{E}, \psi_{f^* \mathcal{E}})$ determines a functor $\mathrm{QCoh}_Y^{\mathrm{Fr}} \rightarrow \mathrm{QCoh}_X^{\mathrm{Fr}}$. We will denote this functor also by f^* , and refer to it as the functor of *pullback along f* .

Remark 10.3.2. In the special case where X and Y are affine, the pullback functor of Construction 10.3.1 agrees with the extension of scalars functor of Construction 2.1.6.

Under some mild assumptions, the pullback functor f^* of Construction 10.3.1 admits a right adjoint:

Proposition 10.3.3. *Let $f : X \rightarrow Y$ be a morphism of schemes which is quasi-compact and quasi-separated. Then the pullback functor $f^* : \mathrm{QCoh}_Y^{\mathrm{Fr}} \rightarrow \mathrm{QCoh}_X^{\mathrm{Fr}}$ admits a right adjoint $f_* : \mathrm{QCoh}_X^{\mathrm{Fr}} \rightarrow \mathrm{QCoh}_Y^{\mathrm{Fr}}$. Moreover, the functor f_* is compatible with the usual direct image functor on quasi-coherent sheaves: that is, the diagram*

$$\begin{array}{ccc} \mathrm{QCoh}_X^{\mathrm{Fr}} & \xrightarrow{f_*} & \mathrm{QCoh}_Y^{\mathrm{Fr}} \\ \downarrow & & \downarrow \\ \mathrm{QCoh}_X & \xrightarrow{f_*} & \mathrm{QCoh}_Y \end{array}$$

commutes up to canonical isomorphism.

Proof. The assumption that f is quasi-compact and quasi-separated guarantees that the pullback functor $f^* : \mathrm{QCoh}_Y \rightarrow \mathrm{QCoh}_X$ admits a right adjoint $f_* : \mathrm{QCoh}_X \rightarrow \mathrm{QCoh}_Y$. If \mathcal{E} is a Frobenius sheaf on X , we can equip the direct image $f_* \mathcal{E}$ with the structure of a Frobenius sheaf on Y by defining $\varphi_{f_* \mathcal{E}}$ to be the composition

$$f_* \mathcal{E} \xrightarrow{f_* \varphi_{\mathcal{E}}} f_* \varphi_{X^*} \mathcal{E} \simeq \varphi_{Y^*} f_* \mathcal{E}.$$

We leave it to the reader to verify that the construction $(\mathcal{E}, \varphi_{\mathcal{E}}) \mapsto (f_* \mathcal{E}, \varphi_{f_* \mathcal{E}})$ determines a functor from $\mathrm{QCoh}_X^{\mathrm{Fr}}$ to $\mathrm{QCoh}_Y^{\mathrm{Fr}}$ which is right adjoint to the pullback functor of Construction 10.3.1. \square

Remark 10.3.4. In the situation of Proposition 10.3.3, if $\mathcal{E} \in \mathrm{QCoh}_X^{\mathrm{Fr}}$ has the property that $\varphi_{\mathcal{E}}$ is an isomorphism, then $\varphi_{f_* \mathcal{E}}$ is also an isomorphism. In other words, the direct image functor f_* carries $\mathrm{QCoh}_X^{\mathrm{perf}}$ into $\mathrm{QCoh}_Y^{\mathrm{perf}}$.

The pullback functor of Construction 10.3.1 generally does not carry perfect Frobenius sheaves to perfect Frobenius sheaves. To remedy this, we consider the following variant:

Construction 10.3.5. Let $f : X \rightarrow Y$ be a morphism of \mathbf{F}_p -schemes. We define a functor $f^\diamond : \mathrm{QCoh}_Y^{\mathrm{perf}} \rightarrow \mathrm{QCoh}_X^{\mathrm{perf}}$ by the formula $f^\diamond(\mathcal{E}) = (f^* \mathcal{E})^{1/p^\infty}$.

Remark 10.3.6. If $f : X \rightarrow Y$ is a quasi-compact, quasi-separated morphism of \mathbf{F}_p -schemes, then the functor $f^\diamond : \mathrm{QCoh}_Y^{\mathrm{perf}} \rightarrow \mathrm{QCoh}_X^{\mathrm{perf}}$ is left adjoint to the direct image functor $f_* : \mathrm{QCoh}_X^{\mathrm{perf}} \rightarrow \mathrm{QCoh}_Y^{\mathrm{perf}}$.

Remark 10.3.7. In the situation where X and Y are affine, the functor $f^\diamond : \mathrm{QCoh}_Y^{\mathrm{perf}} \rightarrow \mathrm{QCoh}_X^{\mathrm{perf}}$ agrees (using the identification of Example 10.1.3) with the functor described in Proposition 3.3.2.

In some cases, there is no difference between the functors f^* and f^\diamond :

Proposition 10.3.8. *Let $f : X \rightarrow Y$ be a morphism of \mathbf{F}_p -schemes. Assume either that f is étale, or that both X and Y are perfect (that is, the Frobenius maps $\varphi_X : X \rightarrow X$ and $\varphi_Y : Y \rightarrow Y$ are isomorphisms). Then the pullback functor $f^* : \mathrm{QCoh}_Y^{\mathrm{Fr}} \rightarrow \mathrm{QCoh}_X^{\mathrm{Fr}}$ carries $\mathrm{QCoh}_Y^{\mathrm{perf}}$ into $\mathrm{QCoh}_X^{\mathrm{perf}}$. Consequently, the functors f^* and f^\diamond coincide on $\mathrm{QCoh}_Y^{\mathrm{perf}}$.*

Proof. The assertion is local on both X and Y , and therefore follows from Corollary 3.4.7 (in the case where f is étale) and Proposition 3.4.5 (in the case where X and Y are perfect). \square

Proposition 10.3.9. *Let $f : X \rightarrow Y$ be a morphism of \mathbf{F}_p -schemes and let \mathcal{E} be an algebraic Frobenius sheaf on Y . Then $f^\diamond \mathcal{E}$ is an algebraic Frobenius sheaf on X . If \mathcal{E} is holonomic, then $f^\diamond \mathcal{E}$ is also holonomic.*

Proof. Both assertions are local on X and Y . We may therefore assume that X and Y are affine, in which case the desired results follow from Corollary 4.2.8 and Proposition 4.1.2. \square

Proposition 10.3.10. *Let X be an \mathbf{F}_p -scheme and let X^{perf} denote the perfection of X (so that $\mathcal{O}_{X^{\mathrm{perf}}} = \mathcal{O}_X^{1/p^\infty}$). Then the canonical map $f : X^{\mathrm{perf}} \rightarrow X$ induces an equivalence of categories $f_* : \mathrm{QCoh}_{X^{\mathrm{perf}}}^{\mathrm{perf}} \rightarrow \mathrm{QCoh}_X^{\mathrm{perf}}$.*

Proof. The assertion is local on X and we may therefore assume that X is affine, in which case the desired conclusion follows from Proposition 3.4.3. \square

We now consider behavior of direct and inverse image functors under the Riemann-Hilbert correspondence. We first observe that any morphism of schemes $f : X \rightarrow Y$ induces a left exact functor $f_* : \mathrm{Shv}_{\mathrm{ét}}(X, \mathbf{F}_p) \rightarrow \mathrm{Shv}_{\mathrm{ét}}(Y, \mathbf{F}_p)$, which is compatible with the direct image functor on quasi-coherent sheaves when f is quasi-compact and quasi-separated. We therefore obtain the following:

Proposition 10.3.11. *Let $f : X \rightarrow Y$ be a morphism of \mathbf{F}_p -schemes which is quasi-compact and quasi-separated. Then the diagram of functors*

$$\begin{array}{ccc} \mathrm{QCoh}_X^{\mathrm{Fr}} & \xrightarrow{\mathrm{Sol}} & \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(X, \mathbf{F}_p) \\ \downarrow f_* & & \downarrow f_* \\ \mathrm{QCoh}_Y^{\mathrm{Fr}} & \xrightarrow{\mathrm{Sol}} & \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(Y, \mathbf{F}_p) \end{array}$$

commutes (up to canonical isomorphism).

Variante 10.3.12. Let $f : X \rightarrow Y$ be any morphism of \mathbf{F}_p -schemes. Then the diagram of functors

$$\begin{array}{ccc} \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(Y, \mathbf{F}_p) & \xrightarrow{f^*} & \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(X, \mathbf{F}_p) \\ \downarrow \mathrm{RH} & & \downarrow \mathrm{RH} \\ \mathrm{QCoh}_Y^{\mathrm{perf}} & \xrightarrow{f^\circ} & \mathrm{QCoh}_X^{\mathrm{perf}} \end{array}$$

commutes (up to canonical isomorphism). In the case where f is quasi-compact and quasi-separated, this follows formally from Proposition 10.3.11 (by passing to left adjoints; see Remark 10.2.11). The general case can be handled by working locally on X and Y (which reduces us to the situation of Proposition 6.2.2).

Construction 10.3.13 (Étale Compactly Supported Direct Images). Let $f : X \rightarrow Y$ be an étale morphism of \mathbf{F}_p -schemes. Then the functor $f^* : \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(Y, \mathbf{F}_p) \rightarrow \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(X, \mathbf{F}_p)$ admits a left adjoint $f_! : \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(X, \mathbf{F}_p) \rightarrow \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(Y, \mathbf{F}_p)$. Using Theorem 10.2.7, we deduce that there is an essentially unique functor $f_! : \mathrm{QCoh}_X^{\mathrm{alg}} \rightarrow \mathrm{QCoh}_Y^{\mathrm{alg}}$ for which the diagram

$$\begin{array}{ccc} \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(X, \mathbf{F}_p) & \xrightarrow{\mathrm{RH}} & \mathrm{QCoh}_X^{\mathrm{alg}} \\ \downarrow f_! & & \downarrow f_! \\ \mathrm{Shv}_{\acute{\mathrm{e}}\mathrm{t}}(Y, \mathbf{F}_p) & \xrightarrow{\mathrm{RH}} & \mathrm{QCoh}_Y^{\mathrm{alg}} \end{array}$$

commutes up to isomorphism. We will refer to $f_!$ as the functor of *compactly supported direct image along f* .

Example 10.3.14. In the situation of Construction 10.3.13, if X and Y are étale, then $f_! : \mathrm{QCoh}_X^{\mathrm{alg}} \rightarrow \mathrm{QCoh}_Y^{\mathrm{alg}}$ agrees with the functor constructed in Section 5.

In the situation of Construction 10.3.13, the functor $f_! : \mathrm{QCoh}_X^{\mathrm{alg}} \rightarrow \mathrm{QCoh}_Y^{\mathrm{alg}}$ can be characterized a left adjoint to the pullback functor $f^* \simeq f^\diamond : \mathrm{QCoh}_Y^{\mathrm{alg}} \rightarrow \mathrm{QCoh}_X^{\mathrm{alg}}$. However, it has a slightly stronger property:

Proposition 10.3.15. *Let $f : X \rightarrow Y$ be an étale morphism of \mathbf{F}_p -schemes and let \mathcal{E} be an algebraic Frobenius sheaf on X . Then, for any perfect Frobenius sheaf \mathcal{F} on Y , the canonical map*

$$\begin{aligned} \theta : \mathrm{Hom}_{\mathrm{QCoh}_Y^{\mathrm{perf}}}(f_! \mathcal{E}, \mathcal{F}) &\rightarrow \mathrm{Hom}_{\mathrm{QCoh}_X^{\mathrm{perf}}}(f^* f_! \mathcal{E}, f^\diamond \mathcal{F}) \\ &\rightarrow \mathrm{Hom}_{\mathrm{QCoh}_X^{\mathrm{perf}}}(\mathcal{E}, f^* \mathcal{F}) \end{aligned}$$

is a bijection.

Proof. By virtue of Theorem 10.2.7, we can assume that $\mathcal{E} = \mathrm{RH}(\mathcal{E})$ for some p -torsion étale sheaf \mathcal{E} on X . In this case, the map θ fits into a commutative diagram

$$\begin{array}{ccc} \mathrm{Hom}_{\mathrm{QCoh}_Y^{\mathrm{perf}}}(f_! \mathrm{RH}(\mathcal{E}), \mathcal{F}) & \xrightarrow{\theta} & \mathrm{Hom}_{\mathrm{QCoh}_X^{\mathrm{perf}}}(\mathrm{RH}(\mathcal{E}), f^* \mathcal{F}) \\ \downarrow & & \downarrow \\ \mathrm{Hom}_{\underline{\mathbf{F}}_p}(f_! \mathcal{E}, \mathrm{Sol}(\mathcal{F})) & \xrightarrow{\quad} & \mathrm{Hom}_{\underline{\mathbf{F}}_p}(\mathcal{E}, \mathrm{Sol}(f^* \mathcal{F})) \end{array}$$

where the bottom horizontal map is an isomorphism because the formation of solution sheaves is local for the étale topology, and the vertical maps are isomorphisms by virtue of Remark 10.2.11. \square

Example 10.3.16. Let $j : U \rightarrow X$ be a quasi-compact open immersion of \mathbf{F}_p -schemes. Then the functor $j_! : \mathrm{QCoh}_U^{\mathrm{alg}} \rightarrow \mathrm{QCoh}_X^{\mathrm{alg}}$ can be described explicitly as follows: if $\mathcal{I} \subseteq \mathcal{O}_{X^{\mathrm{perf}}}$ denotes the (necessarily quasi-coherent) radical ideal sheaf defining $(X - U)^{\mathrm{perf}}$, then the Frobenius automorphism of $\mathcal{O}_{X^{\mathrm{perf}}}$ endows \mathcal{I} with the structure of a holonomic Frobenius module on X , and we have $j_!^{\mathrm{alg}}(\mathcal{O}_{U^{\mathrm{perf}}}) \simeq \mathcal{I}$. More generally, for any $E \in \mathrm{QCoh}_X^{\mathrm{alg}}$, we have $j_!^{\mathrm{alg}}(j^\diamond E) \simeq \mathcal{I} \otimes E$.

10.4. Holonomic Frobenius Sheaves

Recall that a Frobenius module M over a commutative \mathbf{F}_p -algebra R is holonomic if and only if there exists an isomorphism $M \simeq M_0^{1/p^\infty}$, where $M_0 \in \mathrm{Mod}_R^{\mathrm{Fr}}$ is finitely presented as an R -module. We now show that holonomic Frobenius sheaves admit an analogous characterization:

Theorem 10.4.1. *Let X be a Noetherian \mathbf{F}_p -scheme and let \mathcal{E} be a Frobenius sheaf on X . The following conditions are equivalent:*

- (1) *There exists a Frobenius subsheaf $\mathcal{E}_0 \subseteq \mathcal{E}$ such that \mathcal{E}_0 is coherent as an \mathcal{O}_X -module and the inclusion $\mathcal{E}_0 \hookrightarrow \mathcal{E}$ induces an isomorphism $\mathcal{E}_0^{1/p^\infty} \simeq \mathcal{E}$.*
- (2) *There exists an isomorphism $\mathcal{E} \simeq \mathcal{E}_0^{1/p^\infty}$ for some $\mathcal{E}_0 \in \text{QCoh}_X^{\text{perf}}$ which is coherent as a \mathcal{O}_X -module.*
- (3) *The Frobenius sheaf \mathcal{E} is holonomic.*

The proof of Theorem 10.4.1 will require some preliminaries.

Remark 10.4.2. Let X and \mathcal{E} be as in Theorem 10.4.1, and suppose that we are given Frobenius subsheaves $\mathcal{E}_0 \subseteq \mathcal{E}_1 \subseteq \mathcal{E}$. If the inclusion $\mathcal{E}_0 \hookrightarrow \mathcal{E}$ induces an isomorphism $\mathcal{E}_0^{1/p^\infty} \simeq \mathcal{E}$, then the inclusion $\mathcal{E}_1 \hookrightarrow \mathcal{E}$ has the same property. This follows immediately from the exactness of the perfection construction $\mathcal{F} \mapsto \mathcal{F}^{1/p^\infty}$.

Lemma 10.4.3. *Let X be an \mathbf{F}_p -scheme, let \mathcal{E} be a Frobenius sheaf on X , and let $\mathcal{E}_0 \subseteq \mathcal{E}$ be a quasi-coherent \mathcal{O}_X -submodule of \mathcal{E} . Then there exists a smallest Frobenius subsheaf $\mathcal{E}' \subseteq \mathcal{E}$ which contains \mathcal{E}_0 .*

Proof. Take \mathcal{E}' to be the image of the composite map

$$\bigoplus_{n \geq 0} (\varphi_X^n)^* \mathcal{E}_0 \rightarrow \bigoplus_{n \geq 0} (\varphi_X^n)^* \mathcal{E} \xrightarrow{\psi_{\mathcal{E}}^n} \mathcal{E}.$$

□

Remark 10.4.4. In the situation of Lemma 10.4.3, the construction $\mathcal{E}_0 \mapsto \mathcal{E}'$ is compatible with pullbacks along flat morphisms; in particular, it is compatible with restrictions to open sets.

Lemma 10.4.5. *Let X be a Noetherian \mathbf{F}_p -scheme, let \mathcal{E} be an algebraic Frobenius sheaf on X , and let $\mathcal{E}_0 \subseteq \mathcal{E}$ be a coherent \mathcal{O}_X -submodule of \mathcal{E} . Then the Frobenius subsheaf $\mathcal{E}' \subseteq \mathcal{E}$ of Lemma 10.4.3 is also coherent as a \mathcal{O}_X -module.*

Proof. By virtue of Remark 10.4.4, we can assume without loss of generality that $X = \text{Spec}(R)$ is affine. In this case, the desired result follows from Remark 2.4.2. □

Proof of Theorem 10.4.1. The implications (1) \Rightarrow (2) \Rightarrow (3) are obvious. We will prove that (3) implies (1). Let \mathcal{E} be a holonomic Frobenius sheaf on X . Choose a finite cover $\{U_i\}$ of X by affine open sets $U_i \simeq \text{Spec}(R_i)$, and

set $M_i = \mathcal{E}(U_i)$. Then each M_i is a holonomic Frobenius module over R_i . We can therefore choose isomorphisms $M_i \simeq N_i^{1/p^\infty}$, where each $N_i \in \text{Mod}_{R_i}^{\text{Fr}}$ is finitely generated as a module over R_i . Replacing each N_i with its image in M_i , we can assume that N_i corresponds to a Frobenius-stable subsheaf $\overline{\mathcal{F}}_i \subseteq \mathcal{E}|_{U_i}$. Applying [15, Tag 01PF], we can find choose a coherent subsheaf $\mathcal{F}_i \subseteq \mathcal{E}$ satisfying $\mathcal{F}_i|_{U_i} = \overline{\mathcal{F}}_i$. Let \mathcal{F} denote the smallest Frobenius subsheaf of \mathcal{E} which contains each \mathcal{F}_i (Lemma 10.4.3). It follows from Lemma 10.4.5 that \mathcal{F} is coherent as a \mathcal{O}_X -module. We claim that the inclusion $\mathcal{F} \hookrightarrow \mathcal{E}$ induces an isomorphism $\mathcal{F}^{1/p^\infty} \simeq \mathcal{E}$. To prove this, it suffices to show that each restriction $\mathcal{E}|_{U_i}$ is the perfection of $\mathcal{F}|_{U_i}$. This follows from Remark 10.4.2, since $\mathcal{F}|_{U_i}$ contains $\overline{\mathcal{F}}_i$ by construction. \square

Remark 10.4.6. Theorem 10.4.1 can be generalized to the non-Noetherian case. If X is an \mathbf{F}_p -scheme which is quasi-compact and quasi-separated and \mathcal{E} is a holonomic Frobenius sheaf on X , then there exists an isomorphism $\mathcal{E} \simeq \mathcal{E}_0^{1/p^\infty}$, where $\mathcal{E}_0 \in \text{QCoh}_X^{\text{Fr}}$ is locally finitely presented as a \mathcal{O}_X -module. To prove this, we first apply Theorem 10.2.7 to choose an isomorphism $\mathcal{E} = \text{RH}(\mathcal{F})$ for some constructible p -torsion étale sheaf \mathcal{F} on X . Using a Noetherian approximation argument, we can choose a map $f : X \rightarrow Y$ and an isomorphism $\mathcal{F} \simeq f^* \mathcal{F}'$, where Y is a Noetherian \mathbf{F}_p -scheme and \mathcal{F}' is a constructible p -torsion étale sheaf on Y . Applying Lemma 10.4.1, we can choose an isomorphism $\text{RH}(\mathcal{F}') \simeq \mathcal{E}'_0^{1/p^\infty}$ for some $\mathcal{E}'_0 \in \text{QCoh}_Y^{\text{Fr}}$ which is coherent as an \mathcal{O}_Y -module. Then

$$\mathcal{E} \simeq \text{RH}(\mathcal{F}) \simeq \text{RH}(f^* \mathcal{F}') \simeq f^\diamond(\text{RH}(\mathcal{F}')) \simeq f^\diamond(\mathcal{E}'_0^{1/p^\infty}) \simeq (f^* \mathcal{E}'_0)^{1/p^\infty},$$

where $f^* \mathcal{E}'_0$ is locally finitely presented as a \mathcal{O}_X -module.

10.5. Proper Direct Images

In §6.5, we proved that the Riemann-Hilbert equivalence $\text{Sol} : \text{Mod}_R^{\text{alg}} \simeq \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ is compatible with direct images along ring homomorphisms which are finite and of finite presentation (Theorem 6.5.1). In this section, we prove a generalization of this result: the global Riemann-Hilbert correspondence of Theorem 10.2.7 is compatible with direct images along morphisms of \mathbf{F}_p -schemes $f : X \rightarrow Y$ which are proper and of finite presentation. In the global setting, there is more to the story, since neither of the direct image functors

$$f_* : \text{QCoh}_X^{\text{Fr}} \rightarrow \text{QCoh}_Y^{\text{Fr}} \quad f_* : \text{Shv}_{\text{ét}}(X, \mathbf{F}_p) \rightarrow \text{Shv}_{\text{ét}}(Y, \mathbf{F}_p)$$

is necessarily exact. In this case, we also have a comparison result for higher direct images (see Theorem 10.5.5 below).

We begin with some general remarks. Let $f : X \rightarrow Y$ be a quasi-compact and quasi-separated morphism of schemes. Then we have higher direct image functors $R^i f_* : \mathrm{QCoh}_X \rightarrow \mathrm{QCoh}_Y$ (see [15, Tag 01XJ]). These functors are equipped with canonical isomorphisms $\varphi_{Y,*} \circ R^i f_* \simeq R^i f_* \circ \varphi_{X,*}$, and therefore carry (perfect) Frobenius sheaves on X to (perfect) Frobenius sheaves on Y . The central observation is the following:

Theorem 10.5.1. *Let $f : X \rightarrow Y$ be a morphism of \mathbf{F}_p -schemes which is proper and of finite presentation. If \mathcal{E} is an algebraic Frobenius sheaf on X , then the higher direct images $R^i f_* \mathcal{E}$ are algebraic Frobenius sheaves on Y .*

We begin by studying the Noetherian case.

Lemma 10.5.2. *Let $f : X \rightarrow Y$ be a proper morphism of Noetherian \mathbf{F}_p -schemes. If \mathcal{E} is a holonomic Frobenius sheaf on X , then the higher direct images $R^i f_* \mathcal{E}$ are holonomic Frobenius sheaves on Y .*

Proof. Invoking Theorem 10.4.1, we can write $\mathcal{E} = \mathcal{E}_0^{1/p^\infty}$, where \mathcal{E}_0 is a Frobenius sheaf on X which is coherent as an \mathcal{O}_X -module. It follows from the direct image theorem [15, Tag 02O3] that the higher direct images $R^i f_* \mathcal{E}_0$ are coherent \mathcal{O}_Y -modules. Since the functors $R^i f_*$ commute with filtered direct limits, we have canonical isomorphisms

$$R^i f_* \mathcal{E} \simeq R^i f_* (\mathcal{E}_0^{1/p^\infty}) \simeq (R^i f_* \mathcal{E}_0)^{1/p^\infty}.$$

Applying Theorem 10.4.1 again, we see that each $R^i f_* \mathcal{E}$ is holonomic. \square

Lemma 10.5.3. *Let R be a commutative \mathbf{F}_p -algebra, let $f : X \rightarrow \mathrm{Spec}(R)$ be a morphism of schemes which is proper and of finite presentation, and let \mathcal{E} be an holonomic Frobenius sheaf on X . Then, for every integer i , the cohomology group $H^i(X, \mathcal{E})$ is an algebraic Frobenius module over R .*

Remark 10.5.4. In the situation of Lemma 10.5.3, one can say more: the cohomology groups $H^i(X, \mathcal{E})$ are actually holonomic Frobenius modules over R (see Corollary 10.6.3 below).

Proof of Lemma 10.5.3. Applying Theorem 10.2.7, we can choose an isomorphism $\mathcal{E} \simeq \mathrm{RH}(\mathcal{F})$ for some constructible sheaf $\mathcal{F} \in \mathrm{Shv}_{\mathrm{et}}^c(X, \mathbf{F}_p)$. Us-

ing Noetherian approximation [15, Tags 01ZM and 081F], we can choose a finitely generated \mathbf{F}_p -subalgebra $R_0 \subseteq R$ and a pullback diagram of schemes

$$\begin{array}{ccc} X & \xrightarrow{\pi} & X_0 \\ \downarrow f & & \downarrow f_0 \\ \mathrm{Spec}(R) & \longrightarrow & \mathrm{Spec}(R_0), \end{array}$$

where f_0 is proper. Enlarging R_0 if necessary, we can further arrange that $\mathcal{F} = \pi^* \mathcal{F}_0$, where \mathcal{F}_0 is a constructible p -torsion étale sheaf on X_0 (see [7, §1, Proposition 4.17]). Set $\mathcal{E}_0 = \mathrm{RH}(\mathcal{F}_0)$, so that $\mathcal{E} \simeq \pi^\diamond \mathcal{E}_0$ (see Variant 10.3.12).

Write R as a filtered direct limit of finitely generated subrings $R_\alpha \subseteq R$ containing R_0 . For each index α , set $X_\alpha = \mathrm{Spec}(R_\alpha) \times_{\mathrm{Spec}(R_0)} X_0$, let $\pi_\alpha : X_\alpha \rightarrow X_0$ be the projection onto the second factor, and set $\mathcal{E}_\alpha = \pi_\alpha^\diamond \mathcal{E}_0$. Then each \mathcal{E}_α is a holonomic Frobenius sheaf on X_α (Proposition 10.3.9). Invoking Lemma 10.5.2, we see that the cohomology group $H^i(X_\alpha, \mathcal{E}_\alpha)$ is a holonomic Frobenius module over R_α , so that the tensor product $R^{1/p^\infty} \otimes_{R_\alpha^{1/p^\infty}} \mathcal{E}_\alpha$ is a holonomic Frobenius module over R (Proposition 4.1.2). We now compute

$$\begin{aligned} H^i(X, \mathcal{E}) &\simeq \varinjlim H^i(X_\alpha, \mathcal{E}_\alpha) \\ &\simeq R^{1/p^\infty} \otimes_{R^{1/p^\infty}} \varinjlim H^i(X_\alpha, \mathcal{E}_\alpha) \\ &\simeq \varinjlim R^{1/p^\infty} \otimes_{R_\alpha^{1/p^\infty}} H^i(X_\alpha, \mathcal{E}_\alpha). \end{aligned}$$

Since the collection of holonomic Frobenius modules over R is closed under direct limits (Proposition 4.2.4), it follows that $H^i(X, \mathcal{E})$ is algebraic. \square

Proof of Theorem 10.5.1. Let $f : X \rightarrow Y$ be a morphism of \mathbf{F}_p -schemes which is proper and of finite presentation, and let \mathcal{E} be an algebraic Frobenius sheaf on X . We wish to show that each higher direct image $R^i f_* \mathcal{E}$ is an algebraic Frobenius sheaf on Y . This assertion is local on Y , so we may assume without loss of generality that $Y = \mathrm{Spec}(R)$ is affine. In this case, X is quasi-compact and quasi-separated, so Corollary 10.2.9 guarantees that we can write \mathcal{E} as a filtered direct limit $\varinjlim \mathcal{E}_\alpha$, where each \mathcal{E}_α is holonomic. Since the functor $\mathcal{E} \mapsto H^i(X, \mathcal{E})$ commutes with filtered direct limits and the collection of algebraic R -modules is closed under direct limits (Proposition 4.2.4), we may replace \mathcal{E} by \mathcal{E}_α and thereby reduce to the case where \mathcal{E} is holonomic. In this case, the desired result follows from Lemma 10.5.3. \square

We now apply Theorem 10.5.1 to the study of our Riemann-Hilbert correspondence.

Theorem 10.5.5. *Let $f : X \rightarrow Y$ be a morphism of \mathbf{F}_p -schemes which is proper and of finite presentation. For every algebraic Frobenius sheaf \mathcal{E} on X , we have canonical isomorphisms $\mathrm{Sol}(R^i f_* \mathcal{E}) \simeq R^i f_* \mathrm{Sol}(\mathcal{E})$ in the category $\mathrm{Shv}_{\acute{e}t}(Y, \mathbf{F}_p)$.*

Proof. Since \mathcal{E} is algebraic, we have an exact sequence

$$0 \rightarrow \mathrm{Sol}(\mathcal{E}) \rightarrow \mathcal{E} \xrightarrow{\mathrm{id} - \varphi_{\mathcal{E}}} \mathcal{E} \rightarrow 0$$

in the category of étale sheaves on X (Remark 10.2.6). This gives rise to a long exact sequence of higher direct images

$$R^{i-1} f_* \mathcal{E} \xrightarrow{\mathrm{id} - \varphi} R^{i-1} f_* \mathcal{E} \rightarrow R^i f_* \mathrm{Sol}(\mathcal{E}) \rightarrow R^i f_* \mathcal{E} \xrightarrow{\mathrm{id} - \varphi} R^i f_* \mathcal{E}$$

which gives rise to short exact sequence of étale sheaves

$$0 \rightarrow \mathcal{F} \rightarrow R^i f_* \mathrm{Sol}(\mathcal{E}) \rightarrow \mathrm{Sol}(R^i f_* \mathcal{E}) \rightarrow 0,$$

where \mathcal{F} denotes the cokernel of the map $(\mathrm{id} - \varphi) : R^{i-1} f_* \mathcal{E} \rightarrow R^{i-1} f_* \mathcal{E}$. It will therefore suffice to show that the map $(\mathrm{id} - \varphi) : R^{i-1} f_* \mathcal{E} \rightarrow R^{i-1} f_* \mathcal{E}$ is an epimorphism of étale sheaves on Y . This follows from Remark 10.2.6, since the Frobenius sheaf $R^{i-1} f_* \mathcal{E}$ is algebraic by virtue of Theorem 10.5.1. \square

Corollary 10.5.6. *Let $f : X \rightarrow Y$ be a morphism of \mathbf{F}_p -schemes which is proper and of finite presentation. Then, for any p -torsion étale sheaf \mathcal{F} on X , we have canonical isomorphisms $\mathrm{RH}(R^i f_* \mathcal{F}) \simeq R^i f_* \mathrm{RH}(\mathcal{F})$.*

Proof. Using Theorem 10.5.5, we have a canonical isomorphism

$$R^i f_* \mathcal{F} \simeq R^i f_* \mathrm{Sol}(\mathrm{RH}(\mathcal{F})) \simeq \mathrm{Sol}(R^i f_* \mathrm{RH}(\mathcal{F})),$$

which is adjoint to a comparison map $\gamma : \mathrm{RH}(R^i f_* \mathcal{F}) \rightarrow R^i f_* \mathrm{RH}(\mathcal{F})$ (Remark 10.2.11). Since the Frobenius sheaf $R^i f_* \mathrm{RH}(\mathcal{F})$ is algebraic (Theorem 10.5.1), the map γ is an isomorphism. \square

10.6. Application: The Proper Base Change Theorem

Suppose we are given a pullback diagram of schemes σ :

$$\begin{array}{ccc} X' & \xrightarrow{g'} & X \\ \downarrow f' & & \downarrow f \\ Y' & \xrightarrow{g} & Y. \end{array}$$

For every étale sheaf \mathcal{F} on X and every integer $n \geq 0$, we have a natural comparison map $\alpha : g^* R^n f_* \mathcal{F} \rightarrow R^n f'_* g'^* \mathcal{F}$ in the category of étale sheaves on Y' . The *proper base change theorem* in étale cohomology asserts that, if the morphism f is proper and \mathcal{F} is a torsion sheaf, then the map α is an isomorphism [15, Tag 095S]. Our goal in this section is to show that, in special case where σ is a diagram of \mathbf{F}_p -schemes and \mathcal{F} is a p -torsion sheaf, the proper base change theorem can be deduced from the results of this paper in an essentially formal way.

We begin with some general remarks. Let σ be as above, and suppose that the morphisms f and f' are quasi-compact and quasi-separated. In this case, we can associate to every quasi-coherent sheaf \mathcal{E} on X a comparison map

$$\beta : g^* R^n f_* \mathcal{E} \rightarrow R^n f'_* g'^* \mathcal{E}$$

in the category QCoh_Y of quasi-coherent sheaves on Y' . Moreover, if σ is a diagram of \mathbf{F}_p -schemes and \mathcal{E} is a Frobenius sheaf on X , then β is a morphism of Frobenius sheaves. If, in addition, the Frobenius sheaf \mathcal{E} is perfect, then the perfection of β supplies a comparison map $\gamma : g^\diamond R^n f_* \mathcal{E} \rightarrow R^n f'_* g'^\diamond \mathcal{E}$ in the category $\mathrm{QCoh}_{Y'}^{\mathrm{perf}}$.

Proposition 10.6.1. *Suppose we are given a pullback diagram of \mathbf{F}_p -schemes*

$$\begin{array}{ccc} X' & \xrightarrow{g'} & X \\ \downarrow f' & & \downarrow f \\ Y' & \xrightarrow{g} & Y, \end{array}$$

where f is proper and of finite presentation. Then, for any algebraic Frobenius sheaf \mathcal{E} on X , the comparison maps $\gamma : g^\diamond R^n f_* \mathcal{E} \rightarrow R^n f'_* g'^\diamond \mathcal{E}$ are isomorphisms.

Proof. The assertion is local on Y and Y' ; we may therefore assume without loss of generality that $Y = \mathrm{Spec}(R)$ and $Y' = \mathrm{Spec}(S)$ are affine. In this case, we wish to show that the canonical map $S^{1/p^\infty} \otimes_{R^{1/p^\infty}} H^*(X, \mathcal{E}) \rightarrow H^*(X', g'^\diamond \mathcal{E})$ is an isomorphism. Choose a finite covering $\{U_i\}$ of X by affine open subsets and let $\{U'_i\}$ denote the open covering of X' given by $U'_i = g'^{-1}U_i$. Let C^* denote the Čech complex of $\{U_i\}$ with coefficients in the sheaf \mathcal{E} , so that we can identify $H^*(X, \mathcal{E})$ with the cohomology of the cochain complex C^* . Note that for any affine open subset $U \subseteq X$ having inverse image $U' \subseteq X'$, we have a canonical isomorphism $(g'^\diamond \mathcal{E})(U') = S^{1/p^\infty} \otimes_{R^{1/p^\infty}} \mathcal{E}(U)$, so that $S^{1/p^\infty} \otimes_{R^{1/p^\infty}} C^*$ is the Čech complex of the open covering

$\{U'_i\}$ with coefficients in the sheaf $g^\circledast \mathcal{E}$. We can therefore identify γ with the canonical map

$$S^{1/p^\infty} \otimes_{R^{1/p^\infty}} H^n(C^*) \rightarrow H^n(S^{1/p^\infty} \otimes_{R^{1/p^\infty}} C^*).$$

For every affine open subset $U \subseteq X$ with inverse image $U' \subseteq X'$, Remark 3.5.5 supplies isomorphisms

$$\mathrm{Tor}_m^{R^{1/p^\infty}}(S^{1/p^\infty}, \mathcal{O}_{X'}^{1/p^\infty}(U)) = \begin{cases} \mathcal{O}_{X'}^{1/p^\infty}(U') & \text{if } m = 0 \\ 0 & \text{otherwise.} \end{cases}$$

It follows that the canonical maps

$$\mathrm{Tor}_k^{R^{1/p^\infty}}(S^{1/p^\infty}, \mathcal{E}(U)) \rightarrow \mathrm{Tor}_k^{\mathcal{O}_{X'}^{1/p^\infty}(U)}(\mathcal{O}_{X'}^{1/p^\infty}(U'), \mathcal{E}(U))$$

are isomorphisms. Our assumption that \mathcal{E} is algebraic guarantees that $\mathcal{E}(U)$ is an algebraic Frobenius module over $\mathcal{O}_X(U)$, so that the groups

$$\mathrm{Tor}_k^{\mathcal{O}_{X'}^{1/p^\infty}(U)}(\mathcal{O}_{X'}^{1/p^\infty}(U'), \mathcal{E}(U))$$

vanish for $k > 0$. We therefore also have $\mathrm{Tor}_k^{R^{1/p^\infty}}(S^{1/p^\infty}, \mathcal{E}(U)) \simeq 0$ for $k > 0$. Allowing U to vary, we conclude that the tensor product $S^{1/p^\infty} \otimes_{R^{1/p^\infty}} C^*$ is equivalent to the left derived tensor product $S^{1/p^\infty} \otimes_{R^{1/p^\infty}}^L C^*$. We therefore have a convergent spectral sequence

$$E_2^{s,t} : \mathrm{Tor}_s^{R^{1/p^\infty}}(S^{1/p^\infty}, H^t(C^*)) \Rightarrow H^{t-s}(R^{1/p^\infty} \otimes_{S^{1/p^\infty}} C^*),$$

in which γ appears as an edge map. To show that γ is an isomorphism, it suffices to show that the groups $E_2^{s,t}$ vanish for $s > 0$. This follows from Theorem 3.5.1, since each $H^t(C^*) \simeq H^t(X, \mathcal{E})$ is an algebraic Frobenius module over R by virtue of Proposition 10.5.1. \square

Corollary 10.6.2 (Proper Base Change). *Suppose we are given a pullback diagram of \mathbf{F}_p -schemes*

$$\begin{array}{ccc} X' & \xrightarrow{g'} & X \\ \downarrow f' & & \downarrow f \\ Y' & \xrightarrow{g} & Y, \end{array}$$

where f is proper and of finite presentation. Then, for every p -torsion étale sheaf \mathcal{F} on X , the comparison map $\alpha : g^* R^n f_* \mathcal{F} \rightarrow R^n f'_* g'^* \mathcal{F}$ is an isomorphism

Proof. Using Corollary 10.5.6, we can identify the image of α under the Riemann-Hilbert correspondence $\mathrm{RH} : \mathrm{Shv}_{\acute{e}t}(Y', \mathbf{F}_p) \rightarrow \mathrm{QCoh}_Y^{\mathrm{perf}}$ with the comparison map $\gamma : g^\diamond R^n f_* \mathrm{RH}(\mathcal{F}) \rightarrow R^n f'_* g'^\diamond \mathrm{RH}(\mathcal{F})$ of Proposition 10.6.1. Since $\mathrm{RH}(\mathcal{F})$ is algebraic, the map γ is an isomorphism, so that α is also an isomorphism. \square

We can use Proposition 10.6.1 to show that Lemma 10.5.2 holds in the non-Noetherian case:

Corollary 10.6.3. *Let $f : X \rightarrow Y$ be a morphism of \mathbf{F}_p -schemes which is proper and of finite presentation. If \mathcal{E} is a holonomic Frobenius sheaf on X , then the higher direct images $R^n f_* \mathcal{E}$ are holonomic Frobenius sheaves on Y .*

Proof of Corollary 10.6.3. The assertion is local on Y , so we may assume without loss of generality that $Y = \mathrm{Spec}(R)$ is affine. Proceeding as in the proof of Lemma 10.5.3, we can choose a pullback diagram

$$\begin{array}{ccc} X & \xrightarrow{\pi} & X_0 \\ \downarrow f & & \downarrow f_0 \\ \mathrm{Spec}(R) & \xrightarrow{\pi'} & \mathrm{Spec}(R_0) \end{array}$$

where f_0 is proper, $R_0 \subseteq R$ is a finitely generated \mathbf{F}_p -subalgebra, and $\mathcal{E} \simeq \pi^\diamond \mathcal{E}_0$ for some holonomic Frobenius module \mathcal{E}_0 on X_0 . Lemma 10.5.2 guarantees that $R^n f_{0*} \mathcal{E}_0$ is holonomic, so that $\pi'^\diamond R^n f_{0*} \mathcal{E}_0$ is also holonomic (Proposition 4.1.2). Proposition 10.6.1 supplies an isomorphism

$$\gamma : \pi'^\diamond R^n f_{0*} \mathcal{E}_0 \rightarrow R^n f_* \pi^\diamond \mathcal{E}_0 \simeq R^n f_* \mathcal{E},$$

so that $R^n f_* \mathcal{E}$ is holonomic as well. \square

Corollary 10.6.4. *Let $f : X \rightarrow Y$ be a morphism of \mathbf{F}_p -schemes which is proper and of finite presentation. Then the higher direct image functors $R^n f_* : \mathrm{Shv}_{\acute{e}t}(X, \mathbf{F}_p) \rightarrow \mathrm{Shv}_{\acute{e}t}(Y, \mathbf{F}_p)$ carry constructible sheaves to constructible sheaves.*

Proof. Combine Corollary 10.6.3 with Theorem 10.2.7 and Corollary 10.5.6. \square

Remark 10.6.5. Let $f : R \rightarrow S$ be a morphism of \mathbf{F}_p -algebras, let X be an R -scheme which is proper and of finite presentation, and set $X_S = X \times_{\mathrm{Spec}(R)} \mathrm{Spec}(S)$. In this situation, we have a comparison map

$$\beta : \mathrm{H}^*(X, \mathcal{O}_X) \otimes_R S \rightarrow \mathrm{H}^*(X_S, \mathcal{O}_{X_S}).$$

In general, this map need not be an isomorphism, even if X is assumed to be smooth and projective over R (see Example 10.6.6). However, the domain and codomain of β can be regarded as Frobenius modules over S , and Proposition 10.6.1 implies that β^{1/p^∞} is an isomorphism: in other words, every element of $\ker(\beta)$ or $\operatorname{coker}(\beta)$ is annihilated by some power of the Frobenius. In other words, the proper base change theorem holds in the setting of coherent cohomology, provided that we work “up to perfection.”

Example 10.6.6. Let k be a field of characteristic p and set $R = k[[t]]$. Let $G \rightarrow \operatorname{Spec}(R)$ be a finite flat group scheme with generic fibre μ_p and special fibre α_p . For each $k \geq 2$, we can approximate the stack $BG \rightarrow \operatorname{Spec}(R)$ by a smooth projective R -scheme X with geometrically connected fibres, i.e., the \mathcal{O} -cohomology of the generic fibre X_η agrees with that of $B(\mu_p)$ in degrees $\leq k$, while that for the special fibre X_s agrees with that of $B(\alpha_p)$ in degrees $\leq k$; an explicit example is provided when $p = k = 2$ by degenerating a “classical” Enriques surface X_η to a “supersingular” Enriques surface X_s . Assume now that $k = 2$ for simplicity. As μ_p is linearly reductive, it follows that $H^i(X_\eta, \mathcal{O}_{X_\eta}) = 0$ for $i \in \{1, 2\}$. On the other hand, $H^i(X_s, \mathcal{O}_{X_s}) \neq 0$ for $i = 1, 2$. Now consider the R -module $H^1(X, \mathcal{O}_X)$. Since $H^0(X_s, \mathcal{O}_{X_s}) = 0$, it is easy to see that $H^1(X, \mathcal{O}_X)$ is t -torsionfree. But $H^1(X, \mathcal{O}_X)[\frac{1}{t}] = H^1(X_\eta, \mathcal{O}_{X_\eta}) = 0$, so it follows that $H^1(X, \mathcal{O}_X) = 0$. On the other hand, $H^1(X_s, \mathcal{O}_{X_s}) \neq 0$, so we have constructed an example where the base change map

$$H^1(X, \mathcal{O}_X) \otimes_R k \rightarrow H^1(X_s, \mathcal{O}_{X_s})$$

is not an isomorphism.

11. The Contravariant Riemann-Hilbert Correspondence

Let R be a smooth algebra over a field k of characteristic p . In [6], Emerton and Kisin construct an equivalence of triangulated categories

$$\operatorname{RSol}_{\text{EK}} : D_{\text{fgu}}^b(R[F])^{\text{op}} \simeq D_c^b(\operatorname{Spec}(R), \mathbf{F}_p),$$

where $D_{\text{fgu}}^b(R[F])$ denotes the full subcategory of $D(R[F])$ spanned by the cohomologically bounded chain complexes whose cohomology groups *finitely generated unit* Frobenius modules and $D_c^b(\operatorname{Spec}(R), \mathbf{F}_p)$ the constructible derived category of $\operatorname{Spec}(R)$: that is, the full subcategory of the derived category of $\operatorname{Shv}_{\text{ét}}(\operatorname{Spec}(R), \mathbf{F}_p)$ spanned by those chain complexes which are cohomologically bounded with constructible cohomology.

Our goal in this section is to review (and generalize) the construction of the functor $\mathrm{RSol}_{\mathrm{EK}}$. We begin in §11.1 by reviewing the notion of a *finitely generated unit Frobenius module* over a commutative \mathbf{F}_p -algebra R (Definition 11.1.3), following [12] and [6]. The collection of finitely generated unit modules is always closed under the formation of cokernels and extensions (Propositions 11.1.4 and 11.1.5). In §11.2 we show that, when R is a regular Noetherian \mathbf{F}_p -algebra, it is also closed under the formation of kernels (Proposition 11.2.1). In this case, we let $D_{\mathrm{fgu}}^b(R[F])$ denote the full subcategory of the derived category $D(R[F])$ spanned by those cochain complexes $M = M^*$ whose cohomology groups $H^n(M)$ are locally finitely generated unit Frobenius modules which vanish for all but finitely many values of n . In §11.3, we show that there is a sensible way to define the definition of the subcategory $D_{\mathrm{fgu}}^b(R[F]) \subseteq D(R[F])$ for an arbitrary \mathbf{F}_p -algebra R , by restricting our attention to cochain complexes which satisfy suitable “derived” versions of the requirements defining finitely generated unit modules (see Definition 11.3.4 and Proposition 11.3.9). In §11.4 we define a solution functor $\mathrm{RSol}_{\mathrm{EK}} : D_{\mathrm{fgu}}^b(\mathrm{Mod}_R^{\mathrm{Fr}})^{\mathrm{op}} \rightarrow D(\mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(R), \mathbf{F}_p))$ and assert that it restricts to an equivalence of categories $D_{\mathrm{fgu}}^b(\mathrm{Mod}_R^{\mathrm{Fr}})^{\mathrm{op}} \simeq D_c^b(\mathrm{Spec}(R), \mathbf{F}_p)$ (Theorem 11.4.4). Taking R to be a smooth algebra of finite type over a field k , this recovers the main result of [6] in the case of the affine scheme $X = \mathrm{Spec}(R)$. However, our equivalence is a bit more general, since we allow R to be an arbitrary \mathbf{F}_p -algebra. The proof of Theorem 11.4.4 will be given in §12 by comparing the functor $\mathrm{RSol}_{\mathrm{EK}}$ with the solution functor Sol of Construction 2.3.1 (and its derived functors).

11.1. Finitely Generated Unit Frobenius Modules

We now introduce the class of finitely generated unit Frobenius modules, following [6].

Notation 11.1.1. Let R be a commutative \mathbf{F}_p -algebra and let M be an R -module. We let $\varphi_R^* M$ denote the R -module obtained from M by extending scalars along the Frobenius homomorphism $\varphi_R : R \rightarrow R$. If M is a Frobenius module over R , then the Frobenius map $\varphi_M : M \rightarrow M^{1/p}$ determines an R -module homomorphism $\varphi_R^* M \rightarrow M$, which we will denote by ψ_M .

Remark 11.1.2. In the situation of Notation 11.1.1, we can regard $\varphi_R^* M$ as a Frobenius module over R (Construction 2.1.6), and ψ_M is a morphism of Frobenius modules over R . Moreover, the morphism ψ_M induces an isomorphism of perfections $(\varphi_R^* M)^{1/p^\infty} \rightarrow M^{1/p^\infty}$. To prove this, we can extend scalars to the perfection R^{1/p^∞} and thereby reduce to the case where R is

perfect. In this case, the morphism ψ_M coincides with (the Frobenius pull-back of) the map $\varphi_M : M \rightarrow M^{1/p}$, which is evidently an isomorphism of perfections.

Definition 11.1.3. Let R be a commutative \mathbf{F}_p -algebra and let M be a Frobenius module over R . We will say that M is *finitely generated unit* if it satisfies the following pair of conditions:

- (a) The module M is finitely generated as a left module over the noncommutative ring $R[F]$ of Notation 2.1.5.
- (b) The map $\psi_M : \varphi_R^* M \rightarrow M$ of Notation 11.1.1 is an isomorphism.

We now record some easy closure properties of the class finitely generated unit Frobenius modules.

Proposition 11.1.4. *Let R be a commutative \mathbf{F}_p -algebra and let $f : M \rightarrow N$ be a morphism of Frobenius modules over R . If M and N are finitely generated unit, then the cokernel $\text{coker}(f)$ is finitely generated unit.*

Proof. Since N is finitely generated as a left module over $R[F]$, the quotient $\text{coker}(f)$ is also finitely generated as a left module over $R[F]$. We have a commutative diagram of exact sequences

$$\begin{array}{ccccccc}
 \varphi_R^* M & \xrightarrow{\varphi_R^*(f)} & \varphi_R^* N & \longrightarrow & \varphi_R^*(\text{coker}(f)) & \longrightarrow & 0 \\
 \downarrow \psi_M & & \downarrow \psi_N & & \downarrow \psi_{\text{coker}(f)} & & \\
 M & \xrightarrow{f} & N & \longrightarrow & \text{coker}(f) & \longrightarrow & 0.
 \end{array}$$

Since ψ_M and ψ_N are isomorphisms, it follows that $\psi_{\text{coker}(f)}$ is also an isomorphism. □

Proposition 11.1.5. *Let R be a commutative \mathbf{F}_p -algebra and suppose we are given an exact sequence of Frobenius modules $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$. If M' and M'' are finitely generated unit, then M is finitely generated unit.*

Proof. Since the collection of finitely generated left $R[F]$ -modules is closed under extensions, the module M is finitely generated over $R[F]$. It will therefore suffice to show that the map $\psi_M : \varphi_R^* M \rightarrow M$ is an isomorphism. Let K denote the kernel of the map $\varphi_R^* M \rightarrow \varphi_R^* M''$. The morphism ψ_M fits into a commutative diagram of exact sequences

$$\begin{array}{ccccccc}
 0 & \longrightarrow & K & \longrightarrow & \varphi_R^* M & \longrightarrow & \varphi_R^* M'' \longrightarrow 0 \\
 & & \downarrow f & & \downarrow \psi_M & & \downarrow \psi_{M''} \\
 0 & \longrightarrow & M' & \longrightarrow & M & \longrightarrow & M'' \longrightarrow 0.
 \end{array}$$

Note that the map $\psi_{M'}$ factors as a composition $\varphi_R^* M' \xrightarrow{g} K \xrightarrow{f} M'$, where g is surjective. Since M' is finitely generated unit, the map $\psi_{M'}$ is an isomorphism. It follows that f is also an isomorphism. Applying the five lemma to the preceding diagram, we conclude that ψ_M is also an isomorphism. \square

11.2. Existence of Kernels

Our next goal is to prove a counterpart of Proposition 11.1.4 for *kernels* of morphisms between finitely generated unit Frobenius modules. This will require a stronger assumption on R :

Proposition 11.2.1. *Let R be a regular Noetherian \mathbf{F}_p -algebra and let $f : M \rightarrow N$ be a morphism of Frobenius modules over R . If M and N are finitely generated unit, then $K = \ker(f)$ is also finitely generated unit.*

The proof of Proposition 11.2.1 is essentially contained in [12] (see also [6]). We include a proof here for the convenience of the reader, and because the proof uses an auxiliary construction which will play a central role in §12.

Construction 11.2.2 (Unitalization). Let R be a commutative \mathbf{F}_p -algebra and let M be an R -module equipped with an R -linear map $\alpha_M : M \rightarrow \varphi_R^* M$. We let M^u denote the direct limit of the diagram

$$M \xrightarrow{\alpha_M} \varphi_R^* M \xrightarrow{\varphi_R^* \alpha_M} \varphi_R^{2*} M \xrightarrow{\varphi_R^{2*} \alpha_M} \varphi_R^{3*} M \rightarrow \dots$$

We will refer to M^u as the *unitalization* of the pair (M, α_M) . Note that there is a canonical isomorphism $M^u \simeq \varphi_R^* M^u$, whose inverse endows M^u with the structure of a Frobenius module over R .

Example 11.2.3. Let R be a commutative \mathbf{F}_p -algebra and let M be a Frobenius module over R for which the map $\psi_M : \varphi_R^* M \rightarrow M$ of Notation 11.1.1 is an isomorphism. Then the unitalization of the pair (M, ψ_M^{-1}) can be identified with M .

Remark 11.2.4 (Functoriality). Let R be a commutative \mathbf{F}_p -algebra and suppose we are given a commutative diagram of R -modules

$$\begin{array}{ccc} M & \xrightarrow{f} & N \\ \downarrow \alpha_M & & \downarrow \alpha_N \\ \varphi_R^* M & \xrightarrow{\varphi_R^*(f)} & \varphi_R^* N. \end{array}$$

Then f induces a map of unitalizations $f^u : M^u \rightarrow N^u$. Moreover:

- The cokernel of f^u can be identified with the unitalization of $\text{coker}(f)$ (with respect to the induced map $\alpha_{\text{coker}(f)} : \text{coker}(f) \rightarrow \text{coker}(\varphi_R^* f) \simeq \varphi_R^* \text{coker}(f)$).
- If the Frobenius map $\varphi_R : R \rightarrow R$ is flat (for example, if R is regular and Noetherian), then the kernel of f^u can be identified with the unitalization of $\text{ker}(f)$ (with respect to the map $\text{ker}(f) \rightarrow \text{ker}(\varphi_R^* f) \simeq \varphi_R^* \text{ker}(f)$).

Let R be a commutative \mathbf{F}_p -algebra and let $R[F]$ denote the noncommutative ring of Notation 2.1.5. For any R -module M , we have a canonical isomorphism

$$R[F] \otimes_R M \simeq M \oplus \varphi_R^* M \oplus \varphi_R^{2*} M \oplus \dots .$$

Suppose that M is equipped with a map $\alpha_M : M \rightarrow \varphi_R^* M$. Then the construction $x \mapsto (x, -\alpha_M(x))$ determines an R -linear map $M \rightarrow M \oplus \varphi_R^* M \subseteq R[F] \otimes_R M$, which extends to an $R[F]$ -linear map $\alpha' : R[F] \otimes_R M \rightarrow R[F] \otimes_R M$. A simple calculation shows that the map α' is a monomorphism with cokernel M^u . We therefore obtain the following:

Proposition 11.2.5. *Let R be a commutative \mathbf{F}_p -algebra and let M be an R -module equipped with an R -linear map $\psi_M : M \rightarrow \varphi_R^* M$. Then the preceding construction determines an exact sequence of Frobenius modules $0 \rightarrow R[F] \otimes_R M \rightarrow R[F] \otimes_R M \rightarrow M^u \rightarrow 0$.*

Corollary 11.2.6. *Let R be a commutative \mathbf{F}_p -algebra and let M be a finitely generated R -module equipped with an R -linear map $\alpha_M : M \rightarrow \varphi_R^* M$. Then the unitalization M^u is a finitely generated unit Frobenius module.*

Proof. Condition (a) of Definition 11.1.3 follows from Proposition 11.2.5, and condition (b) is immediate from the construction. □

We will be primarily interested in the following special case of Construction 11.2.2:

Construction 11.2.7. Let R be a commutative \mathbf{F}_p -algebra and let M be a Frobenius module over R , which we regard as an R -module equipped with an R -linear map $\psi_M : \varphi_R^* M \rightarrow M$ (Notation 11.1.1). Suppose that M is finitely generated and projective as an R -module, with R -linear dual $M^\vee = \text{Hom}_R(M, R)$. Then the dual of ψ_M is an R -linear map $\psi_M^\vee : M^\vee \rightarrow \varphi_R^* M^\vee$. We let $\mathbb{D}(M)$ denote the unitalization of the pair (M^\vee, ψ_M^\vee) .

Example 11.2.8. Let R be a commutative \mathbf{F}_p -algebra and let M be a Frobenius module over R . Suppose that M is a projective R -module of finite rank and that the map $\psi_M : \varphi_R^* M \rightarrow M$ is an isomorphism. In this case,

the Frobenius module $\mathbb{D}(M)$ of Construction 11.2.7 can be identified with the R -linear dual M^\vee , endowed with the Frobenius structure characterized by the formula $\psi_{M^\vee} = (\psi_M^\vee)^{-1}$ (Example 11.2.3).

Proposition 11.2.9. *Let R be a commutative \mathbf{F}_p -algebra and let M be a Frobenius module over R which is finitely generated and projective as an R -module. Then $\mathbb{D}(M)$ has projective dimension ≤ 1 as a left $R[F]$ -module.*

Proof. Use the exact sequence $0 \rightarrow R[F] \otimes_R M^\vee \rightarrow R[F] \otimes_R M^\vee \rightarrow \mathbb{D}(M) \rightarrow 0$ supplied by Proposition 11.2.5. \square

Remark 11.2.10. In the situation of Construction 11.2.7, the R -module $\mathbb{D}(M)$ is presented as a filtered direct limit of projective R -modules of finite rank, and is therefore flat over R .

In the case where R is a smooth algebra over a field k , Emerton and Kisin prove a converse to Corollary 11.2.6: every finitely generated unit Frobenius module arises as the unitalization of a finitely generated R -module M , equipped with some map $\alpha_M : M \rightarrow \varphi_R^* M$. The proof given in [6] applies more generally whenever R is a regular Noetherian \mathbf{F}_p -algebra (Corollary 11.2.12). We begin with an observation which is valid for *any* \mathbf{F}_p -algebra R :

Proposition 11.2.11. *Let R be a commutative \mathbf{F}_p -algebra and let M be a finitely generated unit Frobenius module over R . Then there exists a Frobenius module N over R which is finitely generated and free as an R -module and a surjective map of Frobenius modules $f : \mathbb{D}(N) \rightarrow M$ (here $\mathbb{D}(N)$ is the Frobenius module given by Construction 11.2.7).*

Proof. Choose a finite collection of elements $\{x_i\}_{i \in I}$ of M which generate M as a left $R[F]$ -module. Invoking the assumption that the map $\psi_M : \varphi_R^* M \rightarrow M$ is an isomorphism, we conclude that M is generated as an R -module by the elements $F^k x_j$ for $k > 0$ and $j \in I$. We may therefore choose some integer $n > 0$ such that each x_i belongs to the R -submodule of M generated by the elements $\{F^k x_j\}_{j \in I, 1 \leq k \leq n}$. Replacing the set $\{x_i\}_{i \in I}$ by the finite set $\{F^k x_i\}_{i \in I, k < n}$, we can reduce to the case $n = 1$: that is, we can arrange that there are relations $x_i = \sum_{j \in I} a_{i,j} \varphi_M(x_j)$ for some coefficients $a_{i,j} \in R$. Let $N = R^I$ be the free R -module on generators y_i^\vee for $i \in I$, and equip N with the structure of a Frobenius module by setting $\varphi_N(y_i^\vee) = \sum_{j \in I} a_{j,i} y_j^\vee$. Using Proposition 11.2.5 (or by inspection), we see that $\mathbb{D}(N)$ can be identified with the left $R[F]$ -module generated by symbols $\{y_i\}_{i \in I}$, subject to the relations $y_i = \sum_{j \in I} a_{i,j} F x_j$. It follows that there is a unique morphism of Frobenius modules $f : \mathbb{D}(N) \rightarrow M$ satisfying $f(y_i) = x_i$. Since the elements x_i generate M as an $R[F]$ -module, the morphism f is surjective. \square

Proof of Proposition 11.2.1. Let R be a regular Noetherian \mathbf{F}_p -algebra and let $f : M \rightarrow N$ be a morphism of finitely generated unit Frobenius modules over R . We wish to show that the kernel $K = \ker(f)$ is also finitely generated unit. The regularity of R guarantees that the Frobenius morphism $\varphi_R : R \rightarrow R$ is flat. It follows that we can identify the pullback φ_R^*K with the kernel of the induced map $\varphi_R^*(f) : \varphi_R^*M \rightarrow \varphi_R^*N$. We therefore have a commutative diagram of short exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & \varphi_R^*K & \longrightarrow & \varphi_R^*M & \xrightarrow{\varphi_R^*(f)} & \varphi_R^*N \\ & & \downarrow \psi_K & & \downarrow \psi_M & & \downarrow \psi_N \\ 0 & \longrightarrow & K & \longrightarrow & M & \xrightarrow{f} & N. \end{array}$$

Since ψ_M and ψ_N are isomorphisms, it follows that ψ_K is also an isomorphism.

We now complete the proof by showing that K is finitely generated as a left $R[F]$ -module. Using Proposition 11.2.11, we can choose a finitely generated projective R -module M_0 , a map $\alpha_{M_0} : M_0 \rightarrow \varphi_R^*M_0$, and a surjection of Frobenius modules $g : M_0^u \rightarrow M$. It follows that the induced map $\ker(g \circ f) \rightarrow K$ is also surjective. We may therefore replace f by $g \circ f$ and thereby reduce to the case $M = M_0^u$. Let f_0 denote the composition of f with the tautological map $M_0 \rightarrow M_0^u \simeq M$. Applying Remark 11.2.4 to the commutative diagram

$$\begin{array}{ccc} M_0 & \xrightarrow{f_0} & N \\ \downarrow \alpha_{M_0} & & \downarrow \psi_N^{-1} \\ \varphi_R^*M_0 & \longrightarrow & \varphi_R^*N, \end{array}$$

we deduce that the kernel of f can be identified with the unitalization of $\ker(f_0)$. Since R is Noetherian, the kernel $\ker(f_0)$ is finitely generated as an R -module, so that $\ker(f)$ is finitely generated as an $R[F]$ -module by virtue of Proposition 11.2.5. □

Corollary 11.2.12 ([6]). *Let R be a regular Noetherian \mathbf{F}_p -algebra and let M be a finitely generated unit Frobenius module over R . Then there exists an isomorphism $M \simeq M_0^u$, where M_0 is a finitely generated R -module equipped with a map $\alpha : M_0 \rightarrow \varphi_R^*M_0$.*

Proof. Using Proposition 11.2.11, we can choose a surjection of Frobenius modules $f : N^u \rightarrow M$, where N is a free R -module of finite rank equipped with a map $\alpha_N : N \rightarrow \varphi_R^*N$. As in the proof of Proposition 11.2.1, we

can write $\ker(f) \simeq K^u$, where K denotes the kernel of the composite map $N \rightarrow N^u \xrightarrow{f} M$ and $\alpha_K : K \rightarrow \varphi_R^* K$ is the restriction of α_N . Applying Remark 11.2.4 to the diagram

$$\begin{array}{ccc} K & \longrightarrow & N \\ \downarrow \alpha_K & & \downarrow \alpha_N \\ \varphi_R^* K & \longrightarrow & \varphi_R^* N, \end{array}$$

we deduce that $M \simeq \operatorname{coker}(K^u \rightarrow N^u)$ can be identified with the unitalization of the quotient N/K (with respect to the map $\alpha_{N/K} : N/K \rightarrow \varphi_R^*(N/K)$ induced by α_N). \square

11.3. Finitely Generated Unit Complexes

Our next goal is to introduce an analogue of Definition 11.1.3 for *cochain complexes* $M = M^*$ of Frobenius modules over a commutative \mathbf{F}_p -algebra R . When R is a regular Noetherian \mathbf{F}_p -algebra, the collection of finitely generated unit Frobenius modules span an abelian subcategory of $\operatorname{Mod}_R^{\operatorname{Fr}}$ which is closed under extensions (Propositions 11.1.4, 11.1.5, and 11.2.1), so we obtain a sensible finiteness condition on cochain complexes by requiring that the cohomology groups $H^n(M)$ are finitely generated unit. However, to get a theory which works well for *arbitrary* \mathbf{F}_p -algebras, we must abandon the idea of having a finiteness condition that can be tested at the level of individual cohomology groups: instead, we will require that the entire cochain complex M^* satisfies suitable analogues of conditions (a) and (b) of Definition 11.1.3, when regarded as an object of a suitable derived category.

Notation 11.3.1. For every associative ring A , we let $D(A)$ denote the derived category of the abelian category of left A -modules. We will be particularly interested in the case where $A = R[F]$ for some commutative \mathbf{F}_p -algebra R ; in this case, we refer to $D(R[F])$ as the *derived category of Frobenius modules over R* . We will generally abuse notation by identifying $\operatorname{Mod}_R^{\operatorname{Fr}}$ with its essential image in $D(R[F])$ (by regarding every Frobenius module over R as a chain complex concentrated in degree zero).

Remark 11.3.2. Let R be a commutative \mathbf{F}_p -algebra and let M be an object of $D(R[F])$. We will generally abuse notation by identifying M with its image under the forgetful functor $D(R[F]) \rightarrow D(R)$. Note that we have a canonical map $M \rightarrow M^{1/p}$ in $D(R)$ (where $M^{1/p}$ denotes the image of M under the functor $D(R) \rightarrow D(R)$ given by restriction of scalars along the

Frobenius). We will denote this map by φ_M and refer to it as *the Frobenius morphism* of M .

Remark 11.3.3 (Comparison with $D(R)$). Let R be a commutative \mathbf{F}_p -algebra. Recall that the forgetful functor $\text{Mod}_R^{\text{Fr}} \rightarrow \text{Mod}_R$ has an exact right adjoint, given by the functor $M \mapsto M^\dagger$ of Construction 3.1.2. Passing to derived categories, we see that the forgetful functor $D(R[F]) \rightarrow D(R)$ also has a right adjoint, given by applying the functor $M \mapsto M^\dagger$ levelwise. For any cochain complex $N = N^\bullet$ of Frobenius modules over R , Construction 3.1.7 produces a short exact sequence of cochain complexes $0 \rightarrow N^\bullet \rightarrow N^{\bullet\dagger} \rightarrow (N^\bullet)^{1/p\dagger} \rightarrow 0$, which we can regard as a distinguished triangle in the derived category $D(R[F])$. It follows that for any object $M \in D(R[F])$, we have a long exact sequence

$$\begin{aligned} \cdots \rightarrow \text{Hom}_{D(R)}(M, N^{1/p}[-1]) &\rightarrow \text{Hom}_{D(R[F])}(M, N) \\ &\rightarrow \text{Hom}_{D(R)}(M, N) \rightarrow \cdots, \end{aligned}$$

which specializes to the exact sequence of Construction 3.1.7 in the special case where M and N are concentrated in a single degree.

We now introduce a “derived” analogue of Definition 11.1.3:

Definition 11.3.4. Let R be a commutative \mathbf{F}_p -algebra and let M be an object of $D(R[F])$. We will say that M is *derived finitely generated unit* if it satisfies the following pair of conditions:

- (a) The module M is a compact object of the triangulated category $D(R[F])$: that is, it is quasi-isomorphic to a bounded chain complex of finitely generated projective left $R[F]$ -modules.
- (b) The Frobenius map $\varphi_M : M \rightarrow M^{1/p}$ induces an isomorphism $R^{1/p} \otimes_R^L M \rightarrow M$ in the derived category $D(R)$.

We let $D_{\text{fgu}}^b(R[F])$ denote the full subcategory of $D(R[F])$ spanned by the derived finitely generated unit objects.

Remark 11.3.5. Let R be a commutative \mathbf{F}_p -algebra. Then $D_{\text{fgu}}^b(R[F])$ is a triangulated subcategory of $D(R[F])$. In other words, for any distinguished triangle $M' \rightarrow M \rightarrow M'' \rightarrow M'[1]$ in $D(R[F])$, if any two of the objects M , M' , and M'' are derived finitely generated unit, then so is the third.

Example 11.3.6. Let R be a commutative \mathbf{F}_p -algebra and let M be a Frobenius module over R which is finitely generated and projective as an R -module. Then the Frobenius module $\mathbb{D}(M)$ of Construction 11.2.7 belongs to $D_{\text{fgu}}^b(R[F])$. Condition (b) of Definition 11.3.4 follows from Corollary 11.2.6

(note that the derived pullback $R^{1/p} \otimes_R^L \mathbb{D}(M)$ agrees with $\varphi_R^* \mathbb{D}(M)$, since $\mathbb{D}(M)$ is flat over R by virtue of Remark 11.2.10). Condition (a) of Definition 11.3.4 follows from the exact sequence $0 \rightarrow R[F] \otimes_R M^\vee \rightarrow R[F] \otimes_R M^\vee \rightarrow \mathbb{D}(M) \rightarrow 0$ of Proposition 11.2.5.

We now study the relationship between the collection of derived finitely generated unit objects of $D(R[F])$ and the collection of finitely generated unit objects of Mod_R^{Fr} . We begin with a simple observation which is valid for any \mathbf{F}_p -algebra R :

Proposition 11.3.7. *Let R be a commutative \mathbf{F}_p -algebra and let M be a nonzero object of $D_{\text{figu}}^b(R[F])$. Then:*

- (1) *There exists a largest integer n for which the Frobenius module $H^n(M)$ is nonzero.*
- (2) *For the integer n of (1), the Frobenius module $H^n(M)$ is finitely generated unit.*

Proof. Without loss of generality, we can assume that M is a bounded cochain complex of finitely generated projective left $R[F]$ -modules. Assertion (1) is immediate. To prove (2), we first note that we can arrange (replacing M by a quasi-isomorphic complex if necessary) that $M^m = 0$ for $m > n$; in this case, we have $H^n(M) = \text{coker}(M^{n-1} \rightarrow M^n)$, which guarantees that $H^n(M)$ is finitely generated as a left $R[F]$ -module. The spectral sequence

$$\text{Tor}_s^R(R^{1/p}, H^t(M)) \Rightarrow H^{t-s}(R^{1/p} \otimes_R^L M)$$

supplies an isomorphism $H^n(R^{1/p} \otimes_R^L M) \simeq \varphi_R^* H^n(M)$, so that condition (b) of Definition 11.1.3 follows from condition (b) of Definition 11.3.4. \square

Corollary 11.3.8. *Let R be a commutative \mathbf{F}_p -algebra and let M be a nonzero object of $D_{\text{figu}}^b(R[F])$, and let n be an integer for which the cohomology groups $H^m(M)$ vanish for $m > n$. Then there exists an object $N \in \text{Mod}_R^{\text{Fr}}$ which is finitely generated and projective as an R -module and a map $f : \mathbb{D}(N)[-n] \rightarrow M$ in $D(R[F])$ for which the induced map $\mathbb{D}(N) \rightarrow H^n(M)$ is surjective.*

Proof. Applying Proposition 11.3.7, we conclude that $H^n(M)$ is a finitely generated unit. Using Proposition 11.2.11, we can choose an object $N \in \text{Mod}_R^{\text{Fr}}$ which is finitely generated and projective as an R -module and a surjection of Frobenius modules $g_0 : \mathbb{D}(N) \rightarrow H^n(M)$. It follows from Proposition 11.2.9 that the map $\text{Hom}_{D(R[F])}(\mathbb{D}(N)[-n], M) \rightarrow \text{Hom}_{R[F]}(\mathbb{D}(N), H^n(M))$ is surjective, so we can lift g_0 to a morphism $g : \mathbb{D}(N)[-n] \rightarrow M$ in the derived category $D(R[F])$. \square

When the \mathbf{F}_p -algebra R is sufficiently nice, there is a very close relationship between Definitions 11.1.3 and 11.3.4:

Proposition 11.3.9. *Let R be a regular Noetherian \mathbf{F}_p -algebra. Then an object $M \in D(R[F])$ belongs to $D_{\text{fgu}}^b(R[F])$ if and only if it satisfies the following conditions:*

- (1) *For every integer n , the cohomology group $H^n(M)$ is a finitely generated unit Frobenius module (in the sense of Definition 11.1.3).*
- (2) *The cohomology groups $H^n(M)$ vanish for $n \ll 0$ and for $n \gg 0$.*

The proof of Proposition 11.3.9 will require a few preliminary remarks. We begin with a standard observation (see for example [11, 5.94]):

Lemma 11.3.10. *Let R be a regular Noetherian ring and let M be a finitely generated R -module. Then M has finite projective dimension as an R -module.*

Remark 11.3.11. In the situation of Lemma 11.3.10, it is not necessarily true that every R -module has finite projective dimension: this holds if and only if R has finite Krull dimension.

Proof of Lemma 11.3.10. We define finitely generated R -modules $\{M(n)\}_{n \geq 0}$ by recursion as follows: set $M(0) = M$, and for $n > 0$ let $M(n)$ denote the kernel of some surjection $R^k \rightarrow M(n-1)$. For each $n \geq 0$, let $U(n) \subseteq \text{Spec}(R)$ denote the set of prime ideals $\mathfrak{p} \subseteq R$ for which the localization $M(n)_{\mathfrak{p}}$ is a projective R -module. Then $U(n)$ is an open subset of $\text{Spec}(R)$: more precisely, it is the largest open subset on which the coherent sheaf associated to $M(n)$ is locally free. Note that a point \mathfrak{p} belongs to $U(n)$ if and only if the localization $M_{\mathfrak{p}}$ has projective dimension $\leq n$ as an $R_{\mathfrak{p}}$ -module. Since R is regular, the set $U(n)$ contains every prime ideal of height $\leq n$. We therefore have $\bigcup_{n \geq 0} U(n) = \text{Spec}(R)$. Since the spectrum $\text{Spec}(R)$ is quasi-compact, we must have $U(n) = \text{Spec}(R)$ for some $n \gg 0$, which guarantees that M has projective dimension $\leq n$. \square

Lemma 11.3.12. *Let R be a regular Noetherian \mathbf{F}_p -algebra and let M be a finitely generated unit Frobenius module over R . Then M has finite projective dimension as a left $R[F]$ -module.*

Proof. Using Corollary 11.2.12, we can assume that M is the unitalization M_0^u , where M_0 is a finitely generated R -module equipped with a map $\alpha : M_0 \rightarrow \varphi_R^* M_0$. Invoking Lemma 11.3.10, we deduce that there exists an integer $n \gg 0$ such that M_0 has finite projective dimension $\leq n$ as an R -module. Note that $R[F]$ is isomorphic to the direct sum $\bigoplus_{m \geq 0} R^{1/p^m}$ as a

right R -module, and is therefore flat over R since the Frobenius map φ_R is flat. It follows that the tensor product $R[F] \otimes_R M_0$ has projective dimension $\leq n$ as a left $R[F]$ -module. Using the exact sequence $0 \rightarrow R[F] \otimes_R M_0 \rightarrow R[F] \otimes_R M_0 \rightarrow M_0^u \rightarrow 0$ of Proposition 11.2.5, we see that $M_0^u \simeq M$ has projective dimension $\leq n + 1$ as a left $R[F]$ -module. \square

Proof of Proposition 11.3.9. Let R be a commutative \mathbf{F}_p -algebra and let $M \in D(R[F])$. We first show that if M satisfies conditions (1) and (2), then M is derived finitely generated unit. Since $D_{\text{fgu}}^b(R[F])$ is a triangulated subcategory of $D(R[F])$, we may assume without loss of generality that M is a finitely generated unit Frobenius module, regarded as a cochain complex concentrated in a single degree. Then the map $\psi_M : \varphi_R^* M \rightarrow M$ is an isomorphism. Since R is a regular Noetherian \mathbf{F}_p -algebra, the Frobenius morphism $\varphi_R : R \rightarrow R$ is flat; we may therefore identify $\varphi_R^* M$ with the derived pullback $R^{1/p} \otimes_R^L M$. It follows that M satisfies condition (b) of Definition 11.3.4. We now verify (a). Using Lemma 11.3.12, we see that there exists an integer $n \geq 1$ that M has projective dimension $\leq n$ as a left module over $R[F]$. We proceed by induction on n . Assume first that $n > 1$. Using Proposition 11.2.11, we can choose a short exact sequence of Frobenius modules

$$0 \rightarrow K \rightarrow \mathbb{D}(N) \rightarrow M \rightarrow 0$$

where $N \in \text{Mod}_R^{\text{Fr}}$ is finitely generated and projective as an R -module. Since $\mathbb{D}(N)$ has projective dimension ≤ 1 over $R[F]$ (Proposition 11.2.9), it follows that K has projective dimension $\leq n - 1$ as a left module over $R[F]$. Using Proposition 11.2.1, we deduce that K is a finitely generated unit Frobenius module. Applying our inductive hypothesis, we conclude that K belongs to $D_{\text{fgu}}^b(R[F])$. Since $\mathbb{D}(N)$ also belongs to $D_{\text{fgu}}^b(R[F])$ (Example 11.3.6), we conclude that M belongs to $D_{\text{fgu}}^b(R[F])$ as desired.

We now treat the case where M has projective dimension ≤ 1 over $R[F]$. Choose an exact sequence of Frobenius modules $0 \rightarrow Q \rightarrow P \rightarrow M \rightarrow 0$, where P is a finitely generated free left $R[F]$ -module. Our assumption that M has projective dimension ≤ 1 guarantees that Q is a projective $R[F]$ -module. Consequently, to verify condition (a) of Definition 11.3.4, it will suffice to show that Q is finitely generated as an $R[F]$ -module. Equivalently, it will suffice to show that the module M is finitely presented as an $R[F]$ -module. This follows from the exact sequence $0 \rightarrow K \rightarrow \mathbb{D}(N) \rightarrow M \rightarrow 0$ above, since $\mathbb{D}(N)$ is a finitely presented left $R[F]$ -module (Proposition 11.2.5) and K is a finitely generated left $R[F]$ -module (Proposition 11.2.1).

We now prove the converse. Suppose that M is an object of $D_{\text{fgu}}^b(R[F])$; we wish to show that M satisfies conditions (1) and (2). Condition (2) is

obvious (since M is quasi-isomorphic to a bounded chain complex of projective left modules over $R[F]$). Suppose that condition (1) fails: then there exists some largest integer n such that $H^n(M)$ is not a finitely generated unit Frobenius module. Form a distinguished triangle $M' \xrightarrow{f} M \rightarrow M'' \rightarrow M'[1]$, where f induces an isomorphism $H^k(M') \rightarrow H^k(M)$ for $k \leq n$, and the groups $H^k(M')$ vanish for $k > n$. Then M'' satisfies conditions (1) and (2), and therefore belongs to $D_{\text{fgu}}^b(R[F])$. It follows that M' also belongs to $D_{\text{fgu}}^b(R[F])$. This contradicts Proposition 11.3.7, since the top cohomology group $H^n(M') \simeq H^n(M)$ is not a finitely generated unit Frobenius module. \square

11.4. The Emerton-Kisin Correspondence

We now introduce a variant of Construction 2.3.1.

Construction 11.4.1. Let R be a commutative \mathbf{F}_p -algebra and let M be a Frobenius module over R . We define a functor

$$\text{Sol}_{\text{EK}}(M) : \text{CAlg}_R^{\text{ét}} \rightarrow \text{Mod}_{\mathbf{F}_p}$$

by the formula $\text{Sol}_{\text{EK}}(M)(A) = \text{Hom}_{R[F]}(M, A)$. It is not difficult to see that the functor $\text{Sol}_{\text{EK}}(M)$ is a sheaf for the étale topology, which is contravariantly functorial in M . We can therefore regard the functor $M \mapsto \text{Sol}_{\text{EK}}(M)$ as a functor of abelian categories $\text{Sol}_{\text{EK}} : (\text{Mod}_R^{\text{Fr}})^{\text{op}} \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$. We will refer to Sol_{EK} as the *Emerton-Kisin solution functor*.

Example 11.4.2. Let R be a commutative \mathbf{F}_p -algebra. Then the Emerton-Kisin solution functor Sol_{EK} carries the Frobenius module $R[F]$ to the structure sheaf of $\text{Spec}(R)$: that is, to the quasi-coherent sheaf \tilde{R} of Example 2.2.5.

Construction 11.4.3. Let R be a commutative \mathbf{F}_p -algebra. We let $D^-(R[F])$ denote the subcategory of $D(R[F])$ spanned by those cochain complexes M which are cohomologically bounded above: that is, which satisfy $H^n(M) \simeq 0$ for $n \gg 0$. Note that $D^-(R[F])$ contains the subcategory $D_{\text{fgu}}^b(R[F]) \subseteq D(R[F])$ of Definition 11.3.4.

Let $D_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ denote the derived category of the abelian category $\text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$. It follows immediately from the definitions that the Emerton-Kisin solution functor $\text{Sol}_{\text{EK}} : (\text{Mod}_R^{\text{Fr}})^{\text{op}} \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ is left exact. It therefore admits a right derived functor

$$\text{RSol}_{\text{EK}} : D^-(R[F])^{\text{op}} \rightarrow D_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p).$$

We can now formulate the main result:

Theorem 11.4.4. *Let R be a commutative \mathbf{F}_p -algebra. Then the functor $\mathrm{RSol}_{\mathrm{EK}} : D^-(R[F])^{\mathrm{op}} \rightarrow D_{\acute{e}t}(\mathrm{Spec}(R), \mathbf{F}_p)$ induces a fully faithful embedding*

$$D_{\mathrm{fgu}}^b(R[F])^{\mathrm{op}} \rightarrow D_{\acute{e}t}(\mathrm{Spec}(R), \mathbf{F}_p),$$

whose essential image is the subcategory $D_c^b(\mathrm{Spec}(R), \mathbf{F}_p) \subseteq D_{\acute{e}t}(\mathrm{Spec}(R), \mathbf{F}_p)$ spanned by those complexes of sheaves which are cohomologically bounded with constructible cohomology sheaves.

Remark 11.4.5. In the special case where R is a smooth algebra of finite type over a field k , Theorem 11.4.4 essentially follows from Theorem 11.3 of [6], applied to the affine scheme $X = \mathrm{Spec}(R)$. Beware that the functor $\mathrm{RSol}_{\mathrm{EK}}$ is not quite the same as the functor appearing in [6]: they differ by a cohomological shift by the dimension of X . Of course, this is an issue of normalization and has no effect on the conclusion of Theorem 11.4.4.

We will give a proof of Theorem 11.4.4 in §12 by developing a theory of duality for Frobenius modules, which will allow us to relate $\mathrm{RSol}_{\mathrm{EK}}$ to the solution functor Sol studied earlier in this paper (see §12.6).

12. Duality for Frobenius Modules

Let R be a commutative \mathbf{F}_p -algebra. Our goal in this section is to prove Theorem 11.4.4 by showing that the functor

$$\mathrm{RSol}_{\mathrm{EK}} : D_{\mathrm{fgu}}^b(R[F])^{\mathrm{op}} \rightarrow D_c^b(\mathrm{Spec}(R); \mathbf{F}_p)$$

is an equivalence of triangulated categories. Our strategy is to construct a commutative diagram of triangulated categories σ :

$$\begin{array}{ccc} & D_{\mathrm{hol}}^b(R[F]) & \\ \mathbb{D} \swarrow & & \searrow \mathrm{RSol} \\ D_{\mathrm{fgu}}^b(R[F])^{\mathrm{op}} & \xrightarrow{\mathrm{RSol}_{\mathrm{EK}}} & D_c^b(\mathrm{Spec}(R); \mathbf{F}_p), \end{array}$$

where $D_{\mathrm{hol}}^b(R[F])$ is the *holonomic* derived category of Frobenius modules, RSol is a derived version of the solution functor of Construction 2.3.1, and \mathbb{D} is a form of R -linear duality.

We begin in §12.1 with a general discussion of the derived category of Frobenius modules $D(R[F])$; in particular, we define the subcategory

$D_{\text{hol}}^b(R[F])$, the functor RSol , and show that it is an equivalence of categories (Corollary 12.1.7). This is essentially a formal consequence of the analogous equivalence at the level of abelian categories (Theorem 7.4.1), since we have already shown that the Riemann-Hilbert correspondence is compatible with the formation of Ext-groups (Corollary 7.2.3).

Most of this section is devoted to the study of the duality functor \mathbb{D} . In §12.3, we introduce the notion of a *weak dual* for an object of the derived category $D(R[F])$ (Definition 12.3.4). The weak dual of an object $M \in D(R[F])$ depends functorially on M , provided that it exists: in other words, the formation of weak duals determines a *partially defined* (contravariant) functor from the derived category $D(R[F])$ to itself. We have already met this functor in a special case: if M is a Frobenius module which is finitely generated and projective as an R -module, then the weak dual of M coincides with the Frobenius module $\mathbb{D}(M)$ given by Construction 11.2.7. This follows from a universal property of Construction 11.2.7, which we establish in §12.2 (Proposition 12.2.1). In §12.5, we exploit this fact to show that every object of $D_{\text{hol}}^b(R[F])$ admits a weak dual (Proposition 12.5.1); the proof uses a characterization of the holonomic derived category which we establish in §12.4 (Theorem 12.4.1). It follows that the construction $M \mapsto \mathbb{D}(M)$ determines a functor $\mathbb{D}(M) : D_{\text{hol}}^b(R[F]) \rightarrow D(R[F])^{\text{op}}$, which we prove to be fully faithful with essential image $D_{\text{fgu}}^b(R[F])$ (Theorem 12.5.4). In §12.6 we show that the diagram σ commutes (up to canonical isomorphism), thereby completing the proof of Theorem 11.4.4.

12.1. The Derived Riemann-Hilbert Correspondence

Our first goal is to extend the equivalence $\text{Sol} : \text{Mod}_R^{\text{hol}} \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ of Theorem 7.4.1 to the level of derived categories. We begin by establishing some notation.

Notation 12.1.1. Let R be a commutative \mathbf{F}_p -algebra. We define subcategories

$$D_{\text{hol}}(R[F]) \subseteq D_{\text{alg}}(R[F]) \subseteq D_{\text{perf}}(R[F]) \subseteq D(R[F])$$

as follows: an object $M \in D(R[F])$ belongs to the subcategory $D_{\text{perf}}(R[F])$ (respectively $D_{\text{alg}}(\text{Mod}_R^{\text{Fr}})$, $D_{\text{hol}}(R[F])$) if each cohomology group $H^i(M)$ is perfect (respectively algebraic, holonomic) when regarded as a Frobenius module over R . It follows from Remark 3.2.2, Proposition 4.2.4, and Corollary 4.3.3, we see that $D_{\text{perf}}(R[F])$, $D_{\text{alg}}(R[F])$, and $D_{\text{hol}}(R[F])$ are triangulated subcategories of $D(R[F])$.

We let $D^+(R[F])$ denote the full subcategory of $D(R[F])$ spanned by those objects M which are (cohomologically) bounded below: that is, for which the cohomology groups $H^i(M)$ vanish for $i \ll 0$. We let $D^b(R[F])$ denote the full subcategory of $D(R[F])$ spanned by those objects which are bounded above and below: that is, for which the cohomology groups $H^i(M)$ vanish both for $i \ll 0$ and $i \gg 0$. Similarly, we have full subcategories

$$D_{\text{hol}}^+(R[F]) \subseteq D_{\text{alg}}^+(R[F]) \subseteq D_{\text{perf}}^+(R[F]) \subseteq D^+(R[F])$$

$$D_{\text{hol}}^b(R[F]) \subseteq D_{\text{alg}}^b(R[F]) \subseteq D_{\text{perf}}^b(R[F]) \subseteq D^b(R[F])$$

which are defined in the obvious way.

For any commutative \mathbf{F}_p -algebra R , the inclusion functor $\text{Mod}_R^{\text{perf}} \hookrightarrow \text{Mod}_R^{\text{Fr}}$ is exact, and therefore extends to a functor of derived categories $D(\text{Mod}_R^{\text{perf}}) \rightarrow D(R[F])$.

Proposition 12.1.2. *Let R be a commutative \mathbf{F}_p -algebra. Then the forgetful functor $D(\text{Mod}_R^{\text{perf}}) \rightarrow D(R[F])$ is a fully faithful embedding, whose essential image is the full subcategory $D_{\text{perf}}(R[F]) \subseteq D(R[F])$.*

Proof. The inclusion functor $\text{Mod}_R^{\text{perf}} \hookrightarrow \text{Mod}_R^{\text{Fr}}$ has a left adjoint, given by the perfection functor $M \mapsto M^{1/p^\infty}$ of Notation 3.2.3. This functor is exact, and therefore extends to a functor of derived categories $F : D(R[F]) \rightarrow D(\text{Mod}_R^{\text{perf}})$ which is left adjoint to the forgetful functor. It now suffices to observe that the counit map $(F \circ G)(M) \rightarrow M$ is an isomorphism for every object $M \in D(\text{Mod}_R^{\text{perf}})$, and that the unit map $N \rightarrow (G \circ F)(N)$ is an isomorphism precisely when N belongs to the full subcategory $D_{\text{perf}}(R[F]) \subseteq D(R[F])$ (since both of these assertions can be checked at the level of cohomology). \square

We now wish to compare the derived categories of Notation 12.1.1 with suitable derived categories of étale sheaves.

Notation 12.1.3. Let R be any commutative ring. We let $D_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ denote the derived category of the abelian category $\text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ of p -torsion étale sheaves on $\text{Spec}(R)$. We define full subcategories

$$D_c^b(\text{Spec}(R), \mathbf{F}_p) \subseteq D_{\text{ét}}^+(\text{Spec}(R), \mathbf{F}_p) \subseteq D_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$$

as follows:

- An object $\mathcal{F} \in D_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ belongs to $D_{\text{ét}}^+(\text{Spec}(R), \mathbf{F}_p)$ if and only if the cohomology sheaves $H^n(\mathcal{F})$ vanish for $n \ll 0$.

- An object $\mathcal{F} \in D_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ belongs to $D_c^b(\text{Spec}(R), \mathbf{F}_p)$ if and only if the cohomology sheaves $H^n(\mathcal{F})$ are constructible for all n and vanish for $|n| \gg 0$.

If R is a commutative \mathbf{F}_p -algebra, then the Riemann-Hilbert functor $\text{RH} : \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p) \rightarrow \text{Mod}_R^{\text{perf}} \subseteq \text{Mod}_R^{\text{Fr}}$ is exact, and therefore extends to a functor of derived categories

$$\text{RH} : D_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p) \rightarrow D(\text{Mod}_R^{\text{perf}}) \simeq D_{\text{perf}}(R[F]).$$

This functor is t-exact, and therefore restricts to a functor $D_{\text{ét}}^+(\text{Spec}(R), \mathbf{F}_p) \rightarrow D^+(\text{Mod}_R^{\text{perf}}) \simeq D_{\text{perf}}^+(R[F])$. This restriction admits a right adjoint

$$\text{RSol} : D_{\text{perf}}^+(R[F]) \simeq D^+(\text{Mod}_R^{\text{perf}}) \rightarrow D_{\text{ét}}^+(\text{Spec}(R), \mathbf{F}_p),$$

given by the total right derived functor of $\text{Sol} : \text{Mod}_R^{\text{perf}} \rightarrow \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$.

Remark 12.1.4. For every object $M \in D_{\text{perf}}^+(R[F])$, we have a hypercohomology spectral sequence $\text{Sol}^s(H^t(M)) \Rightarrow H^{s+t}(\text{RSol}(M))$. Note that the groups $\text{Sol}^s(H^t(M))$ vanish for $s \geq 2$ (Proposition 7.2.1 and Theorem 2.4.3, or Proposition 9.4.1), so this spectral sequence degenerates to yield short exact sequences

$$0 \rightarrow \text{Sol}^1(H^{n-1}(M)) \rightarrow H^n \text{RSol}(M) \rightarrow \text{Sol}(H^n(M)) \rightarrow 0.$$

If M belongs to the subcategory $D_{\text{alg}}^+(R[F]) \subseteq D_{\text{perf}}^+(R[F])$, then the sheaves $\text{Sol}^1(H^{n-1}(M))$ vanish (Proposition 9.5.6); we therefore obtain isomorphisms $H^*(\text{RSol}(M)) \simeq \text{Sol}(H^*(M))$.

Theorem 12.1.5. *Let R be a commutative \mathbf{F}_p -algebra. Then the functor $\text{RH} : D_{\text{ét}}^+(\text{Spec}(R), \mathbf{F}_p) \rightarrow D^+(R[F])$ is a fully faithful embedding, whose essential image is the full subcategory $D_{\text{alg}}^+(R[F]) \subseteq D^+(R[F])$.*

Proof. Since the Riemann-Hilbert functor $\text{RH} : \text{Shv}_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p) \rightarrow \text{Mod}_R^{\text{perf}}$ is exact at the level of abelian categories, its extension to the level of derived categories is t-exact: that is, we have canonical isomorphisms $H^*(\text{RH}(\mathcal{F})) \simeq \text{RH}(H^*(\mathcal{F}))$ for each $\mathcal{F} \in D_{\text{ét}}^+(\text{Spec}(R), \mathbf{F}_p)$. It follows from Theorem 6.1.1 that the functor RH carries $D_{\text{ét}}^+(\text{Spec}(R), \mathbf{F}_p)$ into $D_{\text{alg}}^+(R[F])$. Combining this observation with Remark 12.1.4, we obtain isomorphisms

$$H^*((\text{RSol} \circ \text{RH})(\mathcal{F})) \simeq \text{Sol}(H^*(\text{RH}(\mathcal{F}))) \simeq (\text{Sol} \circ \text{RH})(H^*(\mathcal{F})).$$

It follows from Proposition 7.2.1 that the unit map $\mathcal{F} \rightarrow (\mathrm{RSol} \circ \mathrm{RH})(\mathcal{F})$ is an isomorphism: that is, the derived Riemann-Hilbert functor is fully faithful. To complete the proof, it will suffice to show that for every object $M \in D_{\mathrm{alg}}^+(R[F])$, the counit map $(\mathrm{RH} \circ \mathrm{RSol})(M) \rightarrow M$ is an isomorphism. Applying Remark 12.1.4 again, we obtain isomorphisms

$$H^*((\mathrm{RH} \circ \mathrm{RSol})(M)) \simeq \mathrm{RH}(H^*(\mathrm{RSol}(M))) \simeq (\mathrm{RH} \circ \mathrm{Sol})(H^*(M)),$$

so that the desired result follows from Theorem 2.4.3. \square

Remark 12.1.6. If R is a Noetherian \mathbf{F}_p -algebra of finite Krull dimension, then one can show that the category of étale sheaves $\mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(R), \mathbf{F}_p)$ has finite injective dimension. In this case, it is not hard to see that Theorem 12.1.5 can be extended to yield an equivalence $\mathrm{RH} : D_{\mathrm{ét}}(\mathrm{Spec}(R), \mathbf{F}_p) \rightarrow D(R[F])$ of *unbounded* derived categories. We do not know if this holds in general.

Combining Theorem 12.1.5 with Theorem 7.4.1, we obtain the following:

Corollary 12.1.7. *Let R be a commutative \mathbf{F}_p -algebra. Then the Riemann-Hilbert functor $\mathrm{RH} : \mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(R), \mathbf{F}_p) \rightarrow \mathrm{Mod}_R^{\mathrm{Fr}}$ induces an equivalence of triangulated categories $D_c^b(\mathrm{Spec}(R), \mathbf{F}_p) \rightarrow D_{\mathrm{hol}}^b(R[F])$; an inverse equivalence is given by applying the derived solution functor RSol .*

Theorem 12.1.5 also implies that a slightly weaker version Proposition 12.1.2 holds for algebraic Frobenius modules:

Corollary 12.1.8. *Let R be a commutative \mathbf{F}_p -algebra. Then the inclusion functor $\mathrm{Mod}_R^{\mathrm{alg}} \hookrightarrow \mathrm{Mod}_R^{\mathrm{Fr}}$ extends to a fully faithful embedding of derived categories $D^+(\mathrm{Mod}_R^{\mathrm{alg}}) \rightarrow D^+(R[F])$, whose essential image is the full subcategory $D_{\mathrm{alg}}^+(\mathrm{Mod}_R^{\mathrm{Fr}}) \subseteq D^+(R[F])$.*

Proof. Since the Riemann-Hilbert functor $\mathrm{RH} : \mathrm{Shv}_{\mathrm{ét}}(\mathrm{Spec}(R), \mathbf{F}_p) \rightarrow \mathrm{Mod}_R^{\mathrm{alg}}$ is an equivalence of categories (Theorem 1.0.2), Corollary 12.1.8 is a reformulation of Theorem 12.1.5. \square

12.2. Duality for R -Projective Frobenius Modules

Let R be a commutative \mathbf{F}_p -algebra and let $M \in \mathrm{Mod}_R^{\mathrm{Fr}}$ be finitely generated and projective as an R -module. In §11.2, we introduced a Frobenius module $\mathbb{D}(M)$, given by the direct limit of the sequence

$$M^\vee \xrightarrow{\psi_M^\vee} \varphi_R^* M^\vee \xrightarrow{\varphi_R^* \psi_M^\vee} \varphi_R^{2*} M^\vee \rightarrow \dots$$

As the notation suggests, we can think of $\mathbb{D}(M)$ as a kind of dual of M in the setting of Frobenius modules. Our goal in this section is to make this idea precise. We begin by observing that there is a canonical map

$$c : R \rightarrow M \otimes_R M^\vee \rightarrow M \otimes_R \mathbb{D}(M).$$

It is not hard to see that c is a map of Frobenius modules (where we regard the tensor product $M \otimes_R \mathbb{D}(M)$ as a Frobenius module via Construction 8.1.1). Moreover, it enjoys the following universal property:

Proposition 12.2.1. *Let R be a commutative \mathbf{F}_p -algebra and let M and N be Frobenius modules over R , where M is finitely generated and projective as an R -module. Then composition with the map $c : R \rightarrow M \otimes_R \mathbb{D}(M)$ induces isomorphisms $\mathrm{Ext}_{R[F]}^n(\mathbb{D}(M), N) \rightarrow \mathrm{Ext}_{R[F]}^n(R, M \otimes_R N)$.*

Proof. Using Proposition 11.2.5 and Remark 11.3.3, we see that both sides can be computed as the cohomology groups of the two-term chain complex

$$M \otimes_R N \xrightarrow{\mathrm{id} - \varphi_M \otimes \varphi_N} M \otimes_R N.$$

□

Remark 12.2.2. The abelian groups $\mathrm{Ext}_{R[F]}^n(\mathbb{D}(M), N) \simeq \mathrm{Ext}_{R[F]}^n(R, M \otimes_R N)$ of Proposition 12.2.1 vanish for $n \geq 2$.

We also have the following dual version of Proposition 12.2.1:

Proposition 12.2.3. *Let R be a commutative \mathbf{F}_p -algebra and let M and N be Frobenius modules over R , where M is finitely generated and projective as an R -module. If N is perfect, then composition with the map $c : R \rightarrow M \otimes_R \mathbb{D}(M)$ induces isomorphisms $\mathrm{Ext}_{R[F]}^n(M, N) \rightarrow \mathrm{Ext}_{R[F]}^n(R, N \otimes_R \mathbb{D}(M))$.*

Proof. Using Remark 11.3.3, we can identify $\mathrm{Ext}_{R[F]}^*(R, N \otimes_R \mathbb{D}(M))$ with the direct limit of the diagram

$$\mathrm{Ext}_{R[F]}^*(M, N) \rightarrow \mathrm{Ext}_{R[F]}^*(\varphi_R^* M, N) \rightarrow \mathrm{Ext}_{R[F]}^*(\varphi_R^{2*} M, N) \rightarrow \cdots,$$

where the transition maps are given by precomposition with the map $\psi_M : \varphi_R^* M \rightarrow M$ of Notation 11.1.1. It will therefore suffice to show that each of the transition maps $\mathrm{Ext}_{R[F]}^*(\varphi_R^{k*} M, N) \rightarrow \mathrm{Ext}_{R[F]}^*(\varphi_R^{(k+1)*} M, N)$ is an isomorphism. This follows from the assumption that N is perfect, since the map ψ_M induces an isomorphism of perfections $(\varphi_R^{k*} M)^{1/p^\infty} \rightarrow (\varphi_R^{(k+1)*} M)^{1/p^\infty}$ (Remark 11.1.2). □

12.3. Weak Duality in $D(R[F])$

We now introduce some language to place Proposition 12.2.1 in a more general context. First, we need a bit of notation.

Construction 12.3.1 (Derived Tensor Products). Let R be a commutative \mathbf{F}_p -algebra. Then we can identify $D(R[F])$ with the category whose objects are K -projective cochain complexes of left $R[F]$ -modules (in the sense of Spaltenstein, see [15, Tag 070G] for a summary), and whose morphisms are homotopy classes of chain maps. Using Example 8.1.2, it is not hard to show that if M^\bullet and N^\bullet are K -projective cochain complexes, then the tensor product $M^\bullet \otimes_R N^\bullet$ is also K -projective (where we regard the tensor product as a chain complex of left $R[F]$ -modules via Construction 8.1.1). This construction gives rise to a functor

$$\otimes_R^L : D(R[F]) \times D(R[F]) \rightarrow D(R[F])$$

which we will refer to as the *derived tensor product*.

Remark 12.3.2. Let R be a commutative \mathbf{F}_p -algebra. Then the forgetful functor $D(R[F]) \rightarrow D(R)$ is compatible with derived tensor products.

Remark 12.3.3. Let R be a commutative \mathbf{F}_p -algebra and let M and N be Frobenius modules over R , which we regard as objects of $D(R[F])$. Then we have canonical isomorphisms $H^n(M \otimes_R^L N) \simeq \mathrm{Tor}_{-n}^R(M, N)$ in the category of Frobenius modules. More generally, if M and N are arbitrary object of $D(R[F])$, we have a convergent spectral sequence

$$\bigoplus_{i+j=t} \mathrm{Tor}_s^R(H^i(M), H^j(N)) \Rightarrow H^{t-s}(M \otimes_R^L N).$$

Definition 12.3.4. Let R be a commutative \mathbf{F}_p -algebra and let M and M' be objects of the derived category $D(R[F])$. We will say that a morphism $c : R \rightarrow M \otimes_R^L M'$ exhibits M' as a weak dual of M if, for every object $N \in D(R[F])$, composition with c induces a bijection

$$\mathrm{Hom}_{D(R[F])}(M', N) \rightarrow \mathrm{Hom}_{D(R[F])}(R, M \otimes_R^L N).$$

Proposition 12.3.5. Let R be a commutative \mathbf{F}_p -algebra and let $M \in \mathrm{Mod}_R^{\mathrm{Fr}}$ be a projective R -module of finite rank. Then the map $c : R \rightarrow M \otimes_R \mathbb{D}(M)$ of Proposition 12.2.1 exhibits $\mathbb{D}(M)$ as a weak dual of M .

Proof. We first observe that $M \otimes_R \mathbb{D}(M)$ can be identified with the derived tensor product $M \otimes_R^L \mathbb{D}(M)$ (since both M and $\mathbb{D}(M)$ are flat R -modules). We wish to show that, for every object $N \in D(R[F])$, composition with c induces an isomorphism $\mathrm{Hom}_{D(R[F])}(\mathbb{D}(M), N) \rightarrow \mathrm{Hom}_{D(R[F])}(R, M \otimes_R^L N)$. Using the fact that $\mathbb{D}(M)$ has finite projective dimension as an $R[F]$ -module (Proposition 11.2.9), we can reduce to the case where N is concentrated in a single degree, in which case the desired result is a translation of Proposition 12.2.1. \square

Notation 12.3.6. Let R be a commutative \mathbf{F}_p -algebra and let $M \in D(R[F])$. It follows immediately from the definitions that if there exists a morphism $c : R \rightarrow M \otimes_R^L M'$ which exhibits M' as a weak dual of M , then the object M' (and the morphism c) are well-defined up to unique isomorphism (in the derived category $D(R[F])$). In this case, we will say that M is *weakly dualizable* and denote its weak dual M' by $\mathbb{D}(M)$. Note that, by virtue of Proposition 12.3.5, this notation is consistent with that of Construction 11.2.7.

Warning 12.3.7. In the situation Definition 12.3.4, the roles of M and M' are not symmetric. A morphism $c : R \rightarrow M \otimes_R^L M'$ which exhibits M' as a weak dual of M generally does not exhibit M as a weak dual of M' (see Example 12.3.10). This asymmetry already appeared in §12.2: note that in the statement of Proposition 12.2.3 we required the Frobenius module N to be perfect, but no corresponding hypothesis was needed in the statement of Proposition 12.2.1.

Proposition 12.3.8. *Let R be a commutative \mathbf{F}_p -algebra and let $c : R \rightarrow M \otimes_R^L M'$ be a morphism in $D(R[F])$ which exhibits M' as a weak dual of M . Then the composite map*

$$R \xrightarrow{c} M \otimes_R^L M' \rightarrow M^{1/p^\infty} \otimes_R^L M'$$

exhibits M' as a weak dual of the perfection M^{1/p^∞} .

We will deduce Proposition 12.3.8 from the following variant of Proposition 3.2.9:

Lemma 12.3.9. *Let R be a commutative \mathbf{F}_p -algebra and let $f : M \rightarrow M'$ be a morphism in $D(R[F])$ which induces an isomorphism $M^{1/p^\infty} \rightarrow M'^{1/p^\infty}$. Then the induced map*

$$\mathrm{Hom}_{D(R[F])}(R, M) \rightarrow \mathrm{Hom}_{D(R[F])}(R, M')$$

is an isomorphism.

Proof. Let N denote the cone of the morphism $M \rightarrow M'$; we will show that $\mathrm{Hom}_{D(R[F])}(R, N[k])$ vanishes for every integer k . By virtue of Remark 11.3.3, it will suffice to show that the map $\mathrm{id} - \varphi_N : H^*(N) \rightarrow H^*(N)$ is an isomorphism. This is clear: the assumption that f induces an equivalence $M^{1/p^\infty} \simeq M'^{1/p^\infty}$ guarantees that N^{1/p^∞} vanishes, so that the action of φ_N is locally nilpotent on $H^*(N)$. \square

Proof of Proposition 12.3.8. Let $c : R \rightarrow M \otimes_R^L M'$ be a morphism in $D(R[F])$ which exhibits M' as a weak dual of M , and let N be any object of $D(R[F])$. Then the composite map

$$\begin{aligned} \mathrm{Hom}_{D(R[F])}(M', N) &\rightarrow \mathrm{Hom}_{D(R[F])}(R, M \otimes_R^L N) \\ &\rightarrow \mathrm{Hom}_{D(R[F])}(R, M^{1/p^\infty} \otimes_R^L N) \end{aligned}$$

is an isomorphism, since the left map is an isomorphism (by virtue of our assumption that M' is a weak dual of M) and the right map is an isomorphism (Lemma 12.3.9). Allowing N to vary, we deduce that M' is also a weak dual of M^{1/p^∞} . \square

Example 12.3.10. Let R be a commutative \mathbf{F}_p -algebra. Then the canonical isomorphism $R \simeq R \otimes_R^L R$ exhibits R as a weak dual of itself. It follows from Proposition 12.3.8 that unit map $u : R \rightarrow R^{1/p^\infty} \simeq R^{1/p^\infty} \otimes_R^L R$ also exhibits R as a weak dual of R^{1/p^∞} . However, u cannot exhibit R^{1/p^∞} as a weak dual of R (unless R is perfect), since the weak dual of R is determined uniquely up to isomorphism.

We conclude this section with another application of Lemma 12.3.9:

Proposition 12.3.11. *Let R be a commutative \mathbf{F}_p -algebra and let $c : R \rightarrow M \otimes_R^L M'$ be a morphism in $D(R[F])$ which exhibits M' as a weak dual of M . Then M' belongs to $D_{\mathrm{fgu}}^b(R[F])$.*

Proof. From the isomorphism $\mathrm{Hom}_{D(R[F])}(M', \bullet) \simeq \mathrm{Hom}_{D(R[F])}(R, M \otimes_R^L \bullet)$ (and the compactness of R as an object of $D(R[F])$), we conclude that M' is a compact object of $D(R[F])$. It will therefore suffice to show that the canonical map $\psi_{M'} : R^{1/p} \otimes_R^L M' \rightarrow M'$ is an isomorphism. Note that $\psi_{M'}$ can be regarded as a morphism in $D(R[F])$; it will therefore suffice to show that for each $N \in D(R[F])$, composition with $\psi_{M'}$ induces an isomorphism

$$\begin{aligned} \theta : \mathrm{Hom}_{D(R[F])}(M', N) &\rightarrow \mathrm{Hom}_{D(R[F])}(R^{1/p} \otimes_R^L M', N) \\ &\simeq \mathrm{Hom}_{D(R[F])}(M', N^{1/p}). \end{aligned}$$

Invoking the universal property of M' , we can identify θ with the natural map $\mathrm{Hom}_{D(R[F])}(R, M \otimes_R^L N) \rightarrow \mathrm{Hom}_{D(R[F])}(R, M \otimes_R N^{1/p})$ (induced by the Frobenius map $\varphi_N : N \rightarrow N^{1/p}$). This map is an isomorphism, since the induced map $M \otimes_R^L N \rightarrow M \otimes_R^L N^{1/p}$ induces an isomorphism of perfection (Lemma 12.3.9). \square

12.4. Presentations of Holonomic Complexes

Let R be a commutative \mathbf{F}_p -algebra and let M be a Frobenius module over R . By definition, M is holonomic if and only if there exists an isomorphism $M \simeq M_0^{1/p^\infty}$, where $M_0 \in \mathrm{Mod}_R^{\mathrm{Fr}}$ is finitely presented as an R -module. Our goal in this section is to prove an analogous statement for objects of the derived category $D(R[F])$:

Theorem 12.4.1. *Let R be a commutative \mathbf{F}_p -algebra and let M be an object of $D(R[F])$. The following conditions are equivalent:*

- (1) *The complex M belongs to the subcategory $D_{\mathrm{hol}}^b(R[F]) \subseteq D(R[F])$: that is, it is cohomologically bounded with holonomic cohomologies.*
- (2) *There exists an isomorphism $M \simeq M_0^{1/p^\infty}$ in the category $D(R[F])$, where $M_0 \in D(R[F])$ has the property that its image in $D(R)$ is compact.*

The proof of Theorem 12.4.1 will require some preliminaries. We first study condition (2) of Theorem 12.4.1. Note that an object $M \in D(R[F])$ has compact image in $D(R)$ if and only if it is quasi-isomorphic to a bounded cochain complex N^* of finitely generated projective R -modules. We now show that, in this situation, we can arrange that N^* is also a cochain complex of Frobenius modules:

Lemma 12.4.2. *Let R be a commutative \mathbf{F}_p -algebra and let M be an object of the derived category $D(R[F])$. The following conditions are equivalent:*

- (1) *The object M is isomorphic (in $D(R[F])$) to a bounded cochain complex of Frobenius modules, each of which is projective of finite rank as an R -module.*
- (2) *The image of M in $D(R)$ is compact: that is, it is isomorphic to a bounded cochain complex of projective R -modules of finite rank.*

Proof. The implication (1) \Rightarrow (2) is clear. Conversely, suppose that (2) is satisfied; we will prove (1). Assume that, as an object of $D(R)$, the complex M is quasi-isomorphic to a finite cochain complex of finitely generated

projective R -modules concentrated in degrees $\{a - n, a - n + 1, \dots, a\}$. Replacing M by a shift if necessary, we can assume $a = 0$. We claim that, as an object of $D(R[F])$, the complex M is quasi-isomorphic to a finite cochain complex of Frobenius modules, which are finitely generated and projective over R , also concentrated in degrees $\{-n, -n + 1, \dots, 0\}$. We proceed by induction on n . If $n = 0$, then the cohomology groups $H^i(M)$ vanish for $i \neq 0$ and $H^0(M)$ is a projective R -module of finite rank. In this case, the desired result follows from the observation that M is isomorphic to $H^0(M)$ as an object of $D(R[F])$. Let us therefore suppose that $n > 0$, and set $N = H^0(M)$. Then N is finitely presented as an R -module, so that N^{1/p^∞} is a holonomic Frobenius module over R . Choose elements x_1, x_2, \dots, x_k which generate N as an R -module. It follows from Proposition 4.2.1 that the image of each x_i in N^{1/p^∞} is annihilated by some element $P_i \in R[F]$ of the form $F^{m_i} + c_{1,i}F^{m_i-1} + \dots + c_{m_i,i}$. Replacing P_i by $F^a P_i$ for $a \gg 0$, we may assume that $P_i(x_i) = 0$. Choose a cocycle $\bar{x}_i \in M^0$ representing x_i , so that we can write $P_i(\bar{x}_i) = dy_i$ for some elements $y_i \in M^{-1}$. The elements \bar{x}_i and y_i determine a map of cochain complexes $f : M' \rightarrow M$, where M' is the two-term complex

$$\dots \rightarrow 0 \rightarrow R[F]^k \xrightarrow{(P_1, \dots, P_k)} R[F]^k \rightarrow 0 \rightarrow \dots$$

Note that M' is isomorphic, as an object of $D(R[F])$, to the Frobenius module $K = \bigoplus_i R[F]/R[F]P_i$, which is projective of finite rank as an R -module. Extend f to a distinguished triangle $Q \xrightarrow{g} M' \xrightarrow{f} M \rightarrow Q[1]$ in $D(R[F])$. Then, as an object of $D(R)$, the complex Q is quasi-isomorphic to a chain complex of finitely generated projective R -modules concentrated in degrees $\{1 - n, \dots, 0\}$. Applying our inductive hypothesis, we may assume that each Q^i is a projective R -module of finite rank and that Q^i vanishes unless $-n < i \leq 0$. Then g determines a map of Frobenius modules $Q^0 \rightarrow K$, and M is quasi-isomorphic to the cochain complex of Frobenius modules

$$\dots \rightarrow 0 \rightarrow Q^{-n+1} \rightarrow Q^{-n+2} \rightarrow \dots \rightarrow Q^0 \rightarrow K \rightarrow 0 \rightarrow \dots$$

□

Remark 12.4.3. Let R be a commutative \mathbf{F}_p -algebra and let $M \in D(R[F])$ be an object whose image in $D(R)$ is compact. Then M is also compact as an object of $D(R[F])$: this follows immediately from Remark 11.3.3. However, the converse is false: the Frobenius module $R[F]$ is compact as an object of $D(R[F])$, but its image in $D(R)$ is not compact unless $R \simeq 0$.

We now apply Lemma 12.4.2 to give a simple characterization of the holonomic derived category $D_{\text{hol}}^b(R[F])$.

Lemma 12.4.4. *Let R be a commutative \mathbf{F}_p -algebra. Then $D_{\text{hol}}^b(R[F])$ is the smallest triangulated subcategory of $D(R[F])$ which contains every object of the form M^{1/p^∞} , where $M \in \text{Mod}_R^{\text{Fr}}$ is finitely generated and projective as an R -module.*

Proof. Let \mathcal{C} be a triangulated subcategory of $D(R[F])$ which contains every object of the form M^{1/p^∞} , where $M \in \text{Mod}_R^{\text{Fr}}$ is finitely generated and projective over R . We wish to show that \mathcal{C} contains every object of $D_{\text{hol}}^b(R[F])$. Using our assumption that \mathcal{C} is a triangulated subcategory, we are reduced to showing that \mathcal{C} contains every holonomic Frobenius module N over R (regarded as a chain complex concentrated in degree zero). Using Proposition 4.1.3, we can assume that N has the form $(R \otimes_{R_0} N')^{1/p^\infty}$, where R_0 is a finitely generated subring of R and N' is a holonomic Frobenius module over R_0 . Choose a surjection $A \rightarrow R_0$, where A is a polynomial ring over \mathbf{F}_p . Then N' is also holonomic when regarded as a Frobenius module over A (Remark 5.3.2). Choose an isomorphism $N' \simeq N'_0^{1/p^\infty}$, where N'_0 is a Frobenius module over A which is finitely generated as an A -module. Since A is a regular Noetherian ring, the A -module N'_0 admits a finite resolution by projective A -modules of finite rank. It follows from Lemma 12.4.2 that N'_0 admits a finite resolution

$$\cdots \rightarrow P_3 \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow N'_0 \rightarrow 0$$

in the category of Frobenius modules over A , where each P_k is projective of finite rank as an A -module. Applying Corollary 3.5.2, we see that $(R \otimes_A P_\bullet)^{1/p^\infty}$ is a finite resolution of N by objects of Mod_R^{Fr} which belong to \mathcal{C} . Since \mathcal{C} is a triangulated subcategory of $D(R[F])$, we deduce that N also belongs to \mathcal{C} . □

Proof of Theorem 12.4.1. Let R be a commutative \mathbf{F}_p -algebra and let \mathcal{C} denote the full subcategory of $D(R[F])$ spanned by those objects which are isomorphic to M_0^{1/p^∞} , for some $M_0 \in D(R[F])$ having compact image in $D(R)$. We wish to show that $\mathcal{C} = D_{\text{hol}}^b(R[F])$. We first show that \mathcal{C} is contained in $D_{\text{hol}}^b(R[F])$. Let $M_0 \in D(R[F])$ have compact image in $D(R)$; we wish to show that M_0^{1/p^∞} belongs to $D_{\text{hol}}^b(R[F])$. Using Lemma 12.4.2, we can assume that M_0 is a bounded cochain complex consisting of Frobenius modules which are finitely generated and projective over R . Since $D_{\text{hol}}^b(R[F])$ is a triangulated subcategory of $D(R[F])$, we can reduce to the case where

M_0 is a finitely generated projective R -module, concentrated in degree zero. In this case, the inclusion is clear (since M_0^{1/p^∞} is a holonomic Frobenius module over R).

We now show that $D_{\text{hol}}^b(R[F])$ is contained in \mathcal{C} . By virtue of Lemma 12.4.4, it will suffice to show that \mathcal{C} is a triangulated subcategory of $D(R[F])$. It is clear that \mathcal{C} contains zero objects of $D(R[F])$ and is closed under shifts; it will therefore suffice to show that it contains the cone of any morphism $f : M \rightarrow N$ where M and N belong to \mathcal{C} . Write $M = M_0^{1/p^\infty}$ and $N = N_0^{1/p^\infty}$, where M_0 and N_0 are objects of $D(R[F])$ having compact image in $D(R)$. Using Lemma 12.4.2, we can further assume that M_0 is a cochain complex of Frobenius modules which are finitely generated and projective over R . Note that N can be identified with the homotopy colimit of the diagram

$$N \xrightarrow{\varphi_N} N^{1/p} \xrightarrow{\varphi_N} N^{1/p^2} \rightarrow \dots$$

Since M_0 is a compact object of $D(R[F])$ (Remark 12.4.3), the composite map $M_0 \rightarrow M \xrightarrow{f} N$ factors through some map $f' : M_0 \rightarrow N_0^{1/p^n}$ for $n \gg 0$. Then f' is adjoint to a map $f'' : \varphi_R^{n*} M_0 \rightarrow N_0$, where $\varphi_R^{n*} M_0$ is the cochain complex obtained from M_0 by applying the pullback functor φ_R^{n*} degreewise. Note that $\varphi_R^{n*} M_0$ is also a bounded cochain complex of finitely generated projective R -modules, and therefore has compact image in $D(R)$. Let C_0 be a cone of f'' . Using Remark 11.1.2 (and the exactness of the functor $K \mapsto K^{1/p^\infty}$), we see that the cone of f can be identified with C_0^{1/p^∞} , and therefore belongs to \mathcal{C} as desired. \square

Remark 12.4.5. With a bit more effort, one can prove the following stronger version of Theorem 12.4.1: the construction $M \mapsto M^{1/p^\infty}$ induces an equivalence of triangulated categories $\mathcal{C} / \mathcal{C}_0 \simeq D_{\text{hol}}^b(R[F])$, where \mathcal{C} denotes the triangulated subcategory of $D(R[F])$ spanned by those objects having compact image in $D(R)$, $\mathcal{C}_0 \subseteq \mathcal{C}$ is the triangulated subcategory spanned by those objects $M \in \mathcal{C}$ satisfying $M^{1/p^\infty} \simeq 0$, and $\mathcal{C} / \mathcal{C}_0$ denotes the Verdier quotient of \mathcal{C} by \mathcal{C}_0 . Since we will not need this fact, the proof is left to the reader.

12.5. The Duality Functor

We now return to the study of the duality construction $M \mapsto \mathbb{D}(M)$ of §12.3.

Proposition 12.5.1. *Let R be a commutative \mathbf{F}_p -algebra and let M be an object of $D_{\text{hol}}^b(R[F])$. Then M is weakly dualizable (in the sense of Notation 12.3.6).*

Proof. Using Theorem 12.4.1, we can assume $M = M_0^{1/p^\infty}$, where $M_0 \in D(R[F])$ has compact image in $D(R)$. By virtue of Lemma 12.4.2, we may assume that M_0 is a bounded cochain complex of finitely generated projective R -modules. Let M_0^\vee denote its R -linear dual (which we also regard as a cochain complex of finitely generated projective R -modules) and let $\mathbb{D}(M_0)$ denote the cochain complex of Frobenius modules obtained by applying Construction 11.2.2 termwise. Let c denote the composite map

$$R \rightarrow M_0 \otimes_R M_0^\vee \rightarrow M_0 \otimes_R \mathbb{D}(M_0).$$

A simple calculation shows that c is a morphism of (cochain complexes of) Frobenius modules. Note that the tensor product $M_0 \otimes_R \mathbb{D}(M_0)$ is equivalent to the derived tensor product $M_0 \otimes_R^L \mathbb{D}(M_0)$ (since both M_0 and $\mathbb{D}(M_0)$ are bounded cochain complexes of flat R -modules). We claim that c exhibits $\mathbb{D}(M_0)$ as a weak dual of M_0 in the derived category $D(R[F])$. In other words, we claim that for every object $N \in D(R[F])$, composition with c induces a bijection

$$\mathrm{Hom}_{D(R[F])}(\mathbb{D}(M_0), N) \rightarrow \mathrm{Hom}_{D(R[F])}(R, M_0 \otimes_R^L N).$$

To prove this, we can proceed by induction on the length of the cochain complex M_0 and thereby reduce to the case where M_0 is concentrated in a single degree, which follows from Proposition 12.3.5. Applying Proposition 12.3.8, we deduce that the composite map

$$R \xrightarrow{c} M_0 \otimes_R^L \mathbb{D}(M_0) \rightarrow M_0^{1/p^\infty} \otimes_R^L \mathbb{D}(M_0) \simeq M \otimes_R^L \mathbb{D}(M_0)$$

exhibits $\mathbb{D}(M_0)$ as a weak dual of M , so that M is weakly dualizable as desired. \square

Recall that a morphism $c : R \rightarrow M \otimes_R^L M'$ which exhibits M' as a weak dual of M need not exhibit M as a weak dual of M' . However, holonomic Frobenius complexes do enjoy the following weak form of biduality.

Notation 12.5.2. Let R be a commutative \mathbf{F}_p -algebra. We let $D_{\mathrm{perf}}(R[F])$ denote the full subcategory of $D(R[F])$ spanned by those cochain complexes M whose cohomology groups $H^*(M)$ are perfect Frobenius modules.

Proposition 12.5.3. *Let R be a commutative \mathbf{F}_p -algebra and let M be an object of $D_{\mathrm{hol}}^b(R[F])$ with weak dual $\mathbb{D}(M)$. Then, for every object $N \in D_{\mathrm{perf}}(R[F])$, composition with the canonical map $c : R \rightarrow M \otimes_R^L \mathbb{D}(M)$ induces an isomorphism*

$$\mathrm{Hom}_{D(R[F])}(M, N) \rightarrow \mathrm{Hom}_{D(R[F])}(R, N \otimes_R^L \mathbb{D}(M)).$$

Proof. Let us say that an object $M \in D_{\text{hol}}^b(R[F])$ is *good* if, for every object $N \in D_{\text{perf}}(R[F])$, the canonical map $\text{Hom}_{D(R[F])}(M, N) \rightarrow \text{Hom}_{D(R[F])}(R, N \otimes_R^L \mathbb{D}(M))$ is an isomorphism. We wish to show that every object of $M \in D_{\text{hol}}^b(R[F])$ is good. It is easy to see that the good objects of $D_{\text{hol}}^b(R[F])$ span a triangulated subcategory. By virtue of Lemma 12.4.4, it will suffice to show that every object of the form M_0^{1/p^∞} is good, where $M_0 \in \text{Mod}_R^{\text{Fr}}$ is finitely generated and projective as an R -module. In this case, for each $N \in D_{\text{perf}}(R[F])$, we have a commutative diagram

$$\begin{array}{ccc} \text{Hom}_{D(R[F])}(M, N) & \longrightarrow & \text{Hom}_{D(R[F])}(R, N \otimes_R^L \mathbb{D}(M)) \\ \downarrow & & \downarrow \\ \text{Hom}_{D(R[F])}(M_0, N) & \xrightarrow{\theta_N} & \text{Hom}_{D(R[F])}(R, N \otimes_R^L \mathbb{D}(M_0)); \end{array}$$

here the right vertical map is bijective by virtue of Proposition 12.3.8, and the left vertical map is bijective by virtue of our assumption that N is perfect. It will therefore suffice to show that the map θ_N is an isomorphism for every perfect object $N \in D_{\text{perf}}(R[F])$. Using the fact that M_0 and R admit finite resolutions by projective left $R[F]$ -modules (Remark 12.4.3), we can reduce to the situation where N is concentrated in a single degree. In this case, the desired result follows from Proposition 12.2.3. \square

We are now ready to prove the main result of this section:

Theorem 12.5.4. *Let R be a commutative \mathbf{F}_p -algebra. Then the construction $M \mapsto \mathbb{D}(M)$ induces an equivalence of categories $D_{\text{hol}}^b(R[F]) \rightarrow D_{\text{fgu}}^b(R[F])^{\text{op}}$.*

Proof. It follows from Propositions 12.5.1 and 12.3.11 that the duality functor $\mathbb{D} : D_{\text{hol}}^b(R[F]) \rightarrow D_{\text{fgu}}^b(R[F])^{\text{op}}$ is well-defined. We next claim that it is fully faithful. Let M and N be objects of $D_{\text{hol}}^b(R[F])$; we wish to show that the canonical map

$$\theta : \text{Hom}_{D(R[F])}(M, N) \rightarrow \text{Hom}_{D(R[F])}(\mathbb{D}(N), \mathbb{D}(M)).$$

Using the definition of $\mathbb{D}(N)$, we can identify the codomain of θ with the set $\text{Hom}_{D(R[F])}(R, N \otimes_R^L \mathbb{D}(M))$. Under this identification, θ corresponds to the comparison map of Proposition 12.5.3, which is an isomorphism because N is perfect.

Let \mathcal{C} denote the essential image of the weak duality functor $\mathbb{D} : D_{\text{hol}}^b(R[F]) \rightarrow D_{\text{fgu}}^b(R[F])^{\text{op}}$, so that \mathcal{C} is a triangulated subcategory of

$D_{\text{fgu}}^b(R[F])$. We will complete the proof by showing that every object $N \in D_{\text{fgu}}^b(R[F])$ belongs to \mathcal{C} . We will deduce this from the following assertion:

(*) There exists a diagram

$$\cdots \rightarrow N(2) \rightarrow N(1) \rightarrow N(0) \rightarrow N(-1) \rightarrow N(-2) \rightarrow \cdots \rightarrow N$$

in the derived category $D(R[F])$, where each $N(k)$ belongs to \mathcal{C} and each of the maps $H^n(N(k)) \rightarrow H^n(N)$ is an isomorphism for $n > k$ and a surjection for $n = k$.

Assume (*) for the moment. Then N can be identified with the homotopy colimit of the diagram $\{N(k)\}_{k \in \mathbf{Z}}$ in the triangulated category $D(R[F])$. Since N is a compact object of $D(R[F])$, it follows that the identity map $\text{id}_N : N \rightarrow N$ factors through $N(k)$ for some integer k : that is, N is a direct summand of $N(k)$. Consequently, to prove that N belongs to \mathcal{C} , it will suffice to show that the category \mathcal{C} is idempotent complete. Using the equivalence $\mathbb{D} : D_{\text{hol}}^b(R[F]) \rightarrow \mathcal{C}^{\text{op}}$, we are reduced to proving that the category $D_{\text{hol}}^b(R[F])$ is idempotent complete, which is clear (since any direct summand of a holonomic Frobenius module over R is itself holonomic; see Corollary 4.2.2).

It remains to prove (*). We will construct the objects $N(k)$ by descending induction on k , taking $N(k) = 0$ for $k \gg 0$. To carry out the induction, it will suffice to prove the following:

(*') Let $f : N(k+1) \rightarrow N$ be a morphism in $D(R[F])$, where $N(k+1) \in \mathcal{C}$ and the induced map $H^n(N(k+1)) \rightarrow H^n(N)$ is an isomorphism for $n > k+1$ and a surjection for $n = k+1$. Then the morphism f factors as a composition $N(k+1) \xrightarrow{f'} N(k) \xrightarrow{f''} N$, where $N(k) \in \mathcal{C}$ and the map $H^n(N(k)) \rightarrow H^n(N)$ is an isomorphism for $n > k$ and a surjection for $n = k$.

To prove (*'), let C denote the cone of f , so that C belongs to $D_{\text{fgu}}^b(R[F])$ and the cohomology groups $H^n(C)$ vanish for $n > k$. Using Corollary 11.3.8, we can choose an object $M \in \text{Mod}_R^{\text{Fr}}$ which is finitely generated and projective as an R -module and a map $g : \mathbb{D}(M)[-n] \rightarrow C$ which induces a surjection $\mathbb{D}(M) \rightarrow H^n(C)$. Invoking the octahedral axiom, we conclude that f factors as a composition $N(k+1) \xrightarrow{f'} N(k) \xrightarrow{f''} N$, where the cone of f' is isomorphic to $\mathbb{D}(M)[-k]$ (which guarantees that $N(k)$ belongs to \mathcal{C}) and the cone of f'' is isomorphic to the cone of g (and therefore has vanishing cohomology in degrees $\geq k$). □

12.6. Comparison of Solution Functors

We will deduce Theorem 11.4.4 from the following comparison result:

Theorem 12.6.1. *Let R be a commutative \mathbf{F}_p -algebra. Then the diagram of categories*

$$\begin{array}{ccc}
 & D_{\text{hol}}^b(R[F]) & \\
 \mathbb{D} \swarrow & & \searrow \text{RSol} \\
 D_{\text{fgu}}^b(R[F])^{\text{op}} & \xrightarrow{\text{RSol}_{\text{EK}}} & D_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)
 \end{array}$$

commutes up canonical isomorphism. Here RSol denotes the derived solution functor of §12.1, \mathbb{D} is the duality functor of Theorem 12.5.4, and RSol_{EK} is the derived Emerton-Kisin solution functor of Construction 11.4.3.

Proof of Theorem 11.4.4 from Theorem 12.6.1. Theorem 12.5.4 asserts that the functor $\mathbb{D} : D_{\text{hol}}^b(R[F]) \rightarrow D_{\text{fgu}}^b(R[F])^{\text{op}}$ is an equivalence of categories, and Corollary 12.1.7 asserts that the functor $\text{RSol} : D_{\text{hol}}^b(R[F]) \rightarrow D_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ is a fully faithful embedding whose essential image is the constructible derived category $D_c^b(\text{Spec}(R), \mathbf{F}_p) \subseteq D_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$. Using the commutative diagram of Theorem 12.6.1, we deduce that $\text{RSol}_{\text{EK}} : D_{\text{fgu}}^b(R[F])^{\text{op}} \rightarrow D_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ is also a fully faithful embedding with essential image $D_c^b(\text{Spec}(R), \mathbf{F}_p)$. \square

The proof of Theorem 12.6.1 will require some auxiliary constructions. We begin by introducing a slight modification of the derived solution functor RSol .

Construction 12.6.2. Let R be a commutative \mathbf{F}_p -algebra and let $M = M^*$ be a cochain complex of Frobenius modules. We let \widetilde{M} denote the associated cochain complex of quasi-coherent sheaves on $\text{Spec}(R)$ (Example 2.2.5), so that the Frobenius morphism φ_M determines an endomorphism of \widetilde{M} , which we will denote by $\varphi_{\widetilde{M}}$. We let $\text{Sol}'(M)$ denote the cochain complex of étale sheaves on $\text{Spec}(R)$ given by the shifted mapping cone $\text{cn}(\text{id} - \varphi_{\widetilde{M}})[-1]$. It is clear that the construction $M \mapsto \text{Sol}'(M)$ respects quasi-isomorphisms and therefore determines a functor of derived categories $\text{Sol}' : D(R[F]) \rightarrow D_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$. By construction, we have a distinguished triangle

$$\text{Sol}'(M) \rightarrow \widetilde{M} \xrightarrow{\text{id} - \varphi_{\widetilde{M}}} \widetilde{M} \rightarrow \text{Sol}'(M)[1],$$

depending functorially on M .

Remark 12.6.3. In the special case where M^* is a bounded below cochain complex of injective objects of $\text{Mod}_R^{\text{perf}}$, we can identify $\text{RSol}(M)$ with the kernel (formed in the category of chain complexes of étale sheaves) of the map $\text{id} - \varphi_{\widetilde{M}} : \widetilde{M} \rightarrow \widetilde{M}$. We therefore obtain a canonical map $\text{RSol}(M) \rightarrow \text{Sol}'(M)$, and Lemma 7.1.2 guarantees that this map is a quasi-isomorphism (even at the level of presheaves). It follows that the functor $\text{RSol} : D_{\text{perf}}^+(R[F]) \rightarrow D_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ is canonically isomorphic to the restriction $\text{Sol}'|_{D_{\text{perf}}^+(R[F])}$.

Construction 12.6.4. Let R be a commutative \mathbf{F}_p -algebra and let P denote the two-term cochain complex

$$\dots \rightarrow 0 \rightarrow R[F] \xrightarrow{1-F} R[F] \rightarrow 0 \rightarrow \dots,$$

which we regard as a projective representative for R in the derived category $D(R[F])$. Let M' be a bounded above cochain complex of projective left $R[F]$ -modules, let M be an arbitrary cochain complex of left $R[F]$ -modules, and suppose we are given a morphism of cochain complexes $\bar{c} : P \rightarrow M \otimes_R M'$, which represents a morphism c from R to $M \otimes_R^L M'$ in the derived category $D(R[F])$. Note that we can identify $\text{RSol}_{\text{EK}}(M')$ and $\text{Sol}'(M)$ with the cochain complexes of étale sheaves given concretely by the formulae

$$\text{RSol}_{\text{EK}}(M')(A) = \text{Hom}_{R[F]}(M', A) \quad \text{Sol}'(M)(A) = \text{Hom}_{R[F]}(P, M \otimes_R A).$$

It follows that \bar{c} determines a map of cochain complexes

$$\begin{aligned} \text{RSol}_{\text{EK}}(M') &= \text{Hom}_{R[F]}(M', \bullet) \\ &\rightarrow \text{Hom}_{R[F]}(M \otimes_R M', M \otimes_R \bullet) \\ &\xrightarrow{\circ \bar{c}} \text{Hom}_{R[F]}(P, M \otimes_R \bullet) \\ &= \text{Sol}'(M). \end{aligned}$$

Note that the chain homotopy class of this map depends only on the chain homotopy class of \bar{c} . We therefore obtain a morphism $\gamma_c : \text{RSol}_{\text{EK}}(M') \rightarrow \text{Sol}'(M)$ in the derived category $D_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$ which depends only on the map $c : R \rightarrow R \rightarrow M \otimes_R^L M'$ in $D(R[F])$.

Proof of Theorem 12.6.1 . By virtue of Remark 12.6.3, it will suffice to show that the functors

$$\text{Sol}', \text{RSol}_{\text{EK}} \circ \mathbb{D} : D_{\text{hol}}^b(R[F]) \rightarrow D_{\text{ét}}(\text{Spec}(R), \mathbf{F}_p)$$

are naturally isomorphic. Fix an object $M \in D_{\text{hol}}^b(R[F])$ and let $c : R \rightarrow M \otimes_R^L \mathbb{D}(M)$ be a morphism in $D(R[F])$ which exhibits $\mathbb{D}(M)$ as a weak

dual of M . Applying Construction 12.6.4, we obtain a morphism $\gamma_c : (\mathrm{RSol}_{\mathrm{EK}} \circ \mathbb{D})(M) \rightarrow \mathrm{Sol}'(M)$ in the derived category $D_{\mathrm{\acute{e}t}}(\mathrm{Spec}(R), \mathbf{F}_p)$. It is not difficult to see that this morphism depends functorially on M , and therefore determines a natural transformation of functors $\gamma : \mathrm{RSol}_{\mathrm{EK}} \circ \mathbb{D} \rightarrow \mathrm{Sol}'$. To complete the proof, it will suffice to show that this natural transformation is invertible: that is, γ_c is a quasi-isomorphism for each $M \in D_{\mathrm{hol}}^b(R[F])$. To prove this, we may assume without loss of generality that $\mathbb{D}(M)$ is represented by a bounded above cochain complex of projective left $R[F]$ -modules and that c is represented by a morphism of cochain complexes $\bar{c} : P \rightarrow M \otimes_R \mathbb{D}(M)$, so that γ_c is represented by the map of cochain complexes of étale sheaves

$$\mathrm{Hom}_{R[F]}(\mathbb{D}(M), \bullet) \rightarrow \mathrm{Hom}_{R[F]}(P, M \otimes_R \bullet)$$

appearing in Construction 12.6.4. We wish to show that this map is a quasi-isomorphism of étale sheaves. In fact, we claim that it is already a quasi-isomorphism of presheaves: that is, for every étale R -algebra A , the map of complexes $\mathrm{Hom}_{R[F]}(\mathbb{D}(M), A) \rightarrow \mathrm{Hom}_{R[F]}(P, M \otimes_R A)$ is a quasi-isomorphism. This is a special case of our assumption that c exhibits $\mathbb{D}(M)$ as a weak dual of M . \square

References

- [1] B. Bhatt and P. Scholze. *Projectivity of the Witt vector affine Grassmannian*, Invent. Math. 209(2) (2017), 329–423.
- [2] M. Blickle and G. Böckle. *Cartier modules: finiteness results*, J. Reine Angew. Math. 661 (2011), 85–123. arXiv:0909.2531.
- [3] M. Blickle and G. Böckle. *Cartier crystals*, arXiv:1309.1035.
- [4] G. Böckle and R. Pink. *Cohomological theory of crystals over function fields*, volume 9 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2009.
- [5] A. Chambert-Loir. *Cohomologie cristalline: un survol*, Exposition. Math. 16(4) (1998), 333–382.
- [6] M. Emerton and M. Kisin. *The Riemann-Hilbert correspondence for unit F -crystals*, Astérisque (293):vi+257, 2004.
- [7] E. Freitag and R. Kiehl. *Étale cohomology and the Weil conjecture*, volume 13 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin,

1988. Translated from the German by Betty S. Waterhouse and William C. Waterhouse, With an historical introduction by J. A. Dieudonné.
- [8] O. Gabber. *Notes on some t -structures*, In *Geometric aspects of Dwork theory. Vol. I, II*, pages 711–734. Walter de Gruyter, Berlin, 2004.
- [9] R. Hotta, K. Takeuchi, and T. Tanisaki. *D-modules, perverse sheaves, and representation theory*, volume 236 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 2008. Translated from the 1995 Japanese edition by Takeuchi.
- [10] N. M. Katz *p -adic properties of modular schemes and modular forms*, pages 69–190. Lecture Notes in Mathematics, Vol. 350, 1973.
- [11] T. Y. Lam. *Lectures on Modules and Rings*, Springer-Verlag New York, 1999.
- [12] G. Lyubeznik. *F-modules: applications to local cohomology and D-modules in characteristic $p > 0$* , J. Reine Angew. Math. 491 (1997), 65–130.
- [13] S. Ohkawa. *Riemann-Hilbert correspondence for unit F-crystals on embeddable algebraic varieties*, arXiv:[1601.01525](https://arxiv.org/abs/1601.01525).
- [14] T. Schedlmeier. *Cartier crystals and perverse constructible étale p -torsion sheaves*, arXiv:[1603.07696](https://arxiv.org/abs/1603.07696).
- [15] The Stacks Project.
- [16] W. van der Kallen. *Descent for the K-theory of polynomial rings*, Math. Z. 191:3 (1986), 405–415.
- [17] J. L. Verdier. *Des catégories dérivées des catégories abéliennes*, Astérisque, vol. 239.

BHARGAV BHATT
 DEPARTMENT OF MATHEMATICS
 UNIVERSITY OF MICHIGAN
 ANN ARBOR, MI 48105
E-mail address: bhargav.bhatt@gmail.com

JACOB LURIE
 DEPARTMENT OF MATHEMATICS
 HARVARD UNIVERSITY
 CAMBRIDGE, MA 02138
E-mail address: lurie@math.harvard.edu

RECEIVED NOVEMBER 28, 2017