

Non-concavity of the Robin ground state

BEN ANDREWS*, JULIE CLUTTERBUCK†, AND DANIEL HAUER

On a convex bounded Euclidean domain, the ground state for the Laplacian with Neumann boundary conditions is a constant, while the Dirichlet ground state is log-concave. The Robin eigenvalue problem can be considered as interpolating between the Dirichlet and Neumann cases, so it seems natural that the Robin ground state should have similar concavity properties. The aim of this paper is to show that this is false by analyzing the perturbation problem from the Neumann case. First, we classify all convex polyhedral domains on which the first variation of the ground state with respect to the Robin parameter at zero is not a concave function. Then, we conclude from this that the Robin ground state is not log-concave (and indeed even has some superlevel sets which are non-convex) for small Robin parameter on polyhedral convex domains outside a special class, and hence also on convex domains with smooth boundary which approximate these in Hausdorff distance.

AMS 2000 SUBJECT CLASSIFICATIONS: 35J25, 35B65, 35J15, 47A75.

KEYWORDS AND PHRASES: Eigenfunction, Eigenvalue problem, Robin boundary condition, Concavity, Quasiconvexity.

1. Introduction and main results

The Laplacian eigenvalue problem on a bounded convex domain $\Omega \subset \mathbb{R}^n$ is to find a function u and a constant λ satisfying

$$(1.1) \quad -\Delta u = \lambda u \text{ in } \Omega,$$

subject to one of the following boundary conditions:

$$(1.2) \quad \begin{array}{ll} \text{Dirichlet:} & u = 0 \text{ on } \partial\Omega, \\ \text{Neumann:} & D_\nu u = 0 \text{ on } \partial\Omega, \\ \text{or Robin:} & D_\nu u + \alpha u = 0 \text{ on } \partial\Omega. \end{array}$$

*The research of the first author was supported by grants DP120102462 and FL150100126 of the Australian Research Council.

†The research of the second author was supported by grant FT1301013 of the Australian Research Council.

Here ν is the outward pointing unit normal to Ω , and α is a real constant. For each of these problems, there exists a non-decreasing sequence of eigenvalues

$$\lambda_0 < \lambda_1 \leq \dots \rightarrow \infty.$$

Our main interest in this paper is in the first Robin eigenvalue $\lambda_0^R(\alpha)$ for $\alpha > 0$, and the corresponding *ground state* u_α which is (up to scaling) the unique eigenfunction with definite sign (which we take to be positive). The Robin problem (1.1)-(1.2) with $\alpha > 0$ is often regarded as interpolating between the Dirichlet and Neumann cases: if we consider α as a parameter, the Neumann case corresponds to $\alpha = 0$ and the Dirichlet case to the limit as $\alpha \rightarrow \infty$. Moreover, if we write the eigenvalues for each boundary condition as $\lambda_j^D, \lambda_j^N, \lambda_j^R(\alpha)$, then the j th Robin eigenvalue $\lambda_j^R(\alpha)$ is monotone in α , so in particular we have the following monotonicity property:

$$\lambda_j^N \leq \lambda_j^R(\alpha) \leq \lambda_j^D \quad \text{for all } \alpha \geq 0.$$

We are particularly concerned with the shape of the first eigenfunction u_α . In the Neumann case, the first eigenfunction u_0 is constant. In the Dirichlet case, the first eigenfunction u_∞ is *log-concave* (that is, $\log u_\infty$ is concave) [6]. Explicit eigenfunctions on rectangular domains show that this cannot be improved to concavity of the eigenfunction itself.

In the Dirichlet case, the log-concavity of the first eigenfunction is a key step in proving the lower bound on the fundamental gap $\lambda_1^D - \lambda_0^D$ [21, 25, 1]. Our investigation of the concavity properties of the ground state was motivated by possible applications to such a lower bound for the Robin case: indeed, in those cases where the first Robin eigenfunction is log-concave, a similar proof as in the Dirichlet case applies, implying the (non-sharp) inequality

$$\lambda_1^R(\alpha) - \lambda_0^R(\alpha) \geq \frac{\pi^2}{D^2},$$

where D is the diameter of Ω and $\alpha > 0$. We describe this result in Section 2.

For some domains, the Robin eigenfunction u_α can be found explicitly and is log-concave. For example, on a ball $\Omega = B_R$ of radius $R > 0$, the first eigenfunction u_α is a rotationally symmetric function $u_\alpha(r)$ satisfying

$$u_\alpha'' + \frac{d-1}{r}u_\alpha' + \lambda_1^R(\alpha)u_\alpha = 0 \quad \text{on } (0, R), \quad \text{with } u_\alpha'(0) = 0.$$

Defining $v = (\log u_\alpha)'$, we have

$$(1.3) \quad v' = \frac{u_\alpha''}{u_\alpha} - \left(\frac{u_\alpha'}{u_\alpha}\right)^2 = -\frac{d-1}{r}v - \lambda_1^R(\alpha) - v^2 < -\frac{d-1}{r}v \quad \text{on } (0, R).$$

By (1.3), $\{r^{d-1}v\}'(r) < 0$ on $(0, R)$ and since $v(0) = 0$, we have that $v < 0$ on $(0, R)$. Letting $w = v'$, we find that

$$w' = -\left(\frac{d}{r} + 2v\right)w - \frac{\lambda_1^R(\alpha) + v^2}{r} < -\left(\frac{d}{r} + 2v\right)w \quad \text{on } (0, R).$$

Since $w < 0$ for small $r \in (0, R)$, the last differential inequality implies that $w < 0$ on $[0, R)$. The eigenvalues of the Hessian of $\log u_\alpha$ are $(\log u_\alpha)'' = w < 0$ and $\frac{(\log u_\alpha)'}{r} = \frac{v}{r} < 0$, so u_α is log-concave.

Another easily computed example is that of rectangular domains given by products of intervals, where separation of variables produces the first eigenfunction as a product of concave trigonometric functions, which is therefore log-concave.

One might expect then that in general, the first Robin eigenfunction u_α with $\alpha > 0$ on a convex domain is log-concave, a question raised by Smits [22]. In this paper we show that this is not the case: there exist convex domains, and small values of $\alpha > 0$, for which the first Robin eigenfunction u_α fails to be log-concave and has some non-convex superlevel sets.

Our first two main results are concerned with convex *polyhedral* domains Ω in \mathbb{R}^d , $d \geq 1$, by which we mean open bounded domains given by the intersection of finitely many open half-spaces:

$$\Omega = \bigcap_{i=1}^m \left\{ x \in \mathbb{R}^d \mid x \cdot \nu_i < b_i \right\},$$

where ν_1, \dots, ν_m are unit vectors and b_1, \dots, b_m are constants. Here, we can assume without loss of generality that the description is minimal, meaning that omitting any one of the half-spaces from the intersection results in a strictly larger set. The *faces* of Ω are given by

$$\Sigma_i = \left\{ x \in \overline{\Omega} \mid x \cdot \nu_i = b_i \right\}$$

for $i = 1, \dots, m$, each of which is itself a convex polyhedral subset of the affine subspace $\{x \in \mathbb{R}^d \mid \nu_i \cdot x = b_i\}$, where ν_i denotes the outer unit normal to Ω on the face Σ_i .

We introduce a special subclass of polyhedral domains, with terminology borrowed from [4]:

Definition 1.1. A convex polyhedral domain Ω in \mathbb{R}^d is a *circumsolid* if there exists a ball $B_R(y_0) \subset \overline{\Omega}$ touching every face of Ω (that is, $\partial B_R(y_0) \cap \Sigma_i$

contains exactly one point for every $i \in \{1, \dots, m\}$). Equivalently, Ω has the form

$$\Omega = \bigcap_{i=1}^m \left\{ x \in \mathbb{R}^d \mid (x - y_0) \cdot \nu_i < R \right\}.$$

We say that a convex polyhedron Ω is a *product of circumsolids* if there is a decomposition of \mathbb{R}^d into orthogonal subspaces E_1, \dots, E_k , and circumsolids $\Omega_i \subset E_i$ for $i = 1, \dots, k$ such that

$$\Omega = \left\{ x \in \mathbb{R}^d \mid \pi_i(x) \in \Omega_i \text{ for } i = 1, \dots, k \right\}$$

where π_i is the orthogonal projection from \mathbb{R}^d onto E_i for each i . Trivially, circumsolids are products of circumsolids.

For an open convex set Ω in \mathbb{R}^d , the *tangent cone* Γ_x to Ω at $x \in \overline{\Omega}$ is

$$\Gamma_x := \left\{ r(y - x) \mid y \in \Omega, r > 0 \right\} = \bigcup_{r>0} r(\Omega - x).$$

If Ω is a convex polyhedron, then the tangent cone Γ_x to Ω at $x \in \overline{\Omega}$ is

$$\Gamma_x = \bigcap_{i \in \mathcal{I}(x)} \left\{ y \in \mathbb{R}^d \mid y \cdot \nu_i < 0 \right\}$$

with index set

$$(1.4) \quad \mathcal{I}(x) := \left\{ i \in \{1, \dots, m\} \mid x \cdot \nu_i = b_i \right\},$$

where $\mathcal{I}(x) = \emptyset$ with $\Gamma_x = \mathbb{R}^d$ if $x \in \Omega$. In fact, Γ_x is a cone over the subset $A_x = \Gamma_x \cap \mathbb{S}^{d-1}$ of the unit sphere. In particular, Γ_x is the intersection of finitely many half-spaces with the origin in their common boundary. We call such a set a *polyhedral cone*.

Definition 1.2. For a convex polyhedral domain Ω , we say that a point $x \in \partial\Omega$ has *consistent normals* if the outward unit normals $\{\nu_i \mid i \in \mathcal{I}(x)\}$ to the tangent cone Γ_x are such that there exists a solution $\gamma \in \mathbb{R}^d$ of the system of equations

$$\gamma \cdot \nu_i = -1, \quad i \in \mathcal{I}(x).$$

Otherwise we say that x has *inconsistent normals*.




Figure 1: Planar circumsolid examples: Regular triangle, regular pentagon, skew quadrilateral.

An example of a convex polyhedral domain Ω with inconsistent normals given in Figure 8. Consistency of the normals at x is equivalent to the statement that the points $\{\nu_i \mid i \in \mathcal{I}(x)\}$ lie in a hyperplane disjoint from the origin, or to the statement that the tangent cone Γ_x is an (unbounded) circumsolid (see Proposition 9.3).

1.1. Examples of polyhedral domains

In one dimension any interval is a circumsolid. Planar examples include all regular polygons, such as the triangle and pentagon in Figure 1. However circumsolids can be non-symmetric, such as the skew quadrilateral in Figure 1.

Every triangle is a circumsolid (Figure 2). The same is not true for quadrilaterals: For the trapezium shown in Figure 3 only a specific spacing between the ends (marked with a dashed line) results in a circumsolid; a very long trapezium is not a circumsolid.




Figure 2: Skew triangle.

In higher dimensions any affine simplex is a circumsolid: For any $d + 1$ points x_0, \dots, x_d in \mathbb{R}^d which do not lie in a $(d - 1)$ -dimensional subspace, the tetrahedron $\{\sum_{i=0}^d \lambda_i x_i \mid \lambda_i \geq 0, \sum_i \lambda_i = 1\}$ is a circumsolid (Figure 4).




Figure 3: Trapezium.




Figure 4: Tetrahedron.

However, truncating one of the vertices as in Figure 6 does not produce a circumsolid unless the plane of truncation is chosen to match the inscribed sphere. Other examples of three-dimensional circumsolids include the platonic solids and other Archimedean solids (see for example Figure 5).

In the plane, the only domains which are nontrivial products of circumsolids are rectangles (products of intervals in orthogonal one-dimensional subspaces). In three dimensions, rectangular prisms (products of three intervals) are products of circumsolids, as are prisms over planar circumsolids, such as the example in Figure 7.

We note that if Ω is a product of circumsolids, then every boundary point has consistent normals, since we can define γ by

$$\pi_i(\gamma) = -\frac{1}{R_i} \pi_i(y_0^i - x) \quad \text{for } i = 1, \dots, k,$$

where y_0^i and R_i are the centre and radius of the circumsolid $\Omega_i \subset E_i$ for each i . In the plane, every boundary point of a convex polygon has consistent normals. Figure 8 is an example of a convex polyhedron in \mathbb{R}^3 with vertex x_0 having inconsistent normals.

The family of products of circumsolids is a special subclass of convex polyhedral domains Ω in \mathbb{R}^d which is directly related to the concavity prop-




Figure 5: Regular dodecahedron.




Figure 6: Tetrahedron with a flat top.




Figure 7: Prism over a regular pentagon.

erties of solutions v to the Neumann problem

$$(1.5) \quad \begin{cases} \Delta v + \mu = 0 & \text{in } \Omega, \\ D_\nu v = -1 & \text{on } \partial\Omega, \end{cases}$$

for some constant μ . Indeed, we show in Corollary 8.3 (see Section 8) that




Figure 8: Tetrahedron with non-horizontal sliced tip.

a weak solution v of (1.5) on a convex polyhedral domain Ω is concave if and only if Ω is a product of circumsolids. Thus, if Ω is a convex polyhedral domain *outside* this class, then v is not concave, and furthermore v has some non-convex superlevel sets. Since the Robin problem (1.1)-(1.2) for small positive α can be treated as a perturbation of (1.5), we conclude that the Robin ground state u_α is not a log-concave function on Ω for small $\alpha > 0$.

1.2. Main results of this paper

The following theorems are the main results of this paper.

Theorem 1.3. *Let Ω be a convex polyhedral domain in \mathbb{R}^d , $d \geq 2$, which is not a product of circumsolids. Then for sufficiently small $\alpha > 0$, the first Robin eigenfunction u_α is not log-concave.*

In our next theorem, we provide sufficient conditions implying that the first Robin eigenfunction admits non-convex superlevel sets.

Theorem 1.4. *Let Ω be a convex polyhedral domain in \mathbb{R}^d . If $d = 2$ and Ω is not a product of circumsolids, then the first Robin eigenfunction u_α admits non-convex superlevel sets for sufficiently small $\alpha > 0$. The same conclusion holds if $d \geq 3$ and Ω has boundary points with inconsistent normals.*

We stress that although Theorem 1.3 is stated for polyhedral domains, one cannot hope to avoid such non-concavity results by imposing more regularity on the boundary.

Corollary 1.5. *Let Ω_0 be a convex polyhedral domain in \mathbb{R}^d , $d \geq 2$, which is not a product of circumsolids. Then for any sufficiently small $\alpha > 0$, for*

any convex domain Ω which is sufficiently close to Ω_0 in Hausdorff distance, the first Robin eigenfunction u_α on Ω is not log-concave.

For $\alpha < 0$, the first Robin eigenvalue λ_α is negative, and the methods used to prove Theorem 1.3 and Corollary 1.5 also lead to the following result.

Theorem 1.6. *Let Ω be a convex polyhedral domain in \mathbb{R}^d , $d \geq 2$, which is not a product of circumsolids. Then for sufficiently small $\alpha < 0$, the first Robin eigenfunction u_α is not log-convex. Moreover, for any convex domain Ω_0 which is sufficiently close to Ω in Hausdorff distance, the first Robin eigenfunction on Ω_0 is not log-convex.*

As mentioned before, our approach to Theorem 1.3 is to treat the Robin problem (1.1)-(1.2) for small positive α as a perturbation from the Neumann case $\alpha = 0$. To be more precise, we show in Section 3 that the function $v = \frac{du_\alpha}{d\alpha} \Big|_{\alpha=0}$ is a weak solution of (1.5) for some μ . Thus, the concavity properties of u_α for small α relate directly to the concavity properties of v , so we proceed to investigate the latter, in the particular case of polyhedral domains. We deduce Theorem 1.3 from the statement that the solution v of (1.5) on a convex polyhedral domain Ω is concave precisely when Ω is a product of circumsolids.

Our argument proceeds as follows: After some preliminary material on the perturbation problem in Section 3, we prove in Section 4 the surprising result that every C^2 solution of (1.5) on a polyhedral domain is a quadratic function (see Theorem 4.3). In section 5 we relate this to semi-concave solutions, by showing that any concave solution of (1.5) is C^2 up to the boundary (see Theorem 5.1). This involves expanding the solution in terms of homogeneous harmonic functions about any boundary point, and requires in particular the interesting observation that any degree two homogeneous harmonic function with bounded second derivatives and with Neumann boundary condition on a polyhedral cone in \mathbb{R}^d is a quadratic function.

To the best of our knowledge, concavity conditions have not been used so far in the existing literature to improve the regularity of weak solutions to elliptic problems on polyhedral domains. Thus the regularity results Theorem 5.1 and Theorem 5.3 are new and may be of independent interest.

In Section 8 we prove that those polyhedral domains on which a quadratic function solves the equation (1.5) are products of circumsolids. This completes the preliminaries needed to prove our main Theorem 1.3 in Section 9.

In conclusion, we want to emphasize that the negative log-concavity statements obtained in Theorem 1.3 and Corollary 1.5 do not (necessarily) imply that the fundamental gap for the Robin case cannot hold. However,

any strategy that depends on the log-concavity of the first ground state (as in Section 2) will not obtain the result for all convex domains. Thus, we discuss in Section 10 some interesting observations and open problems for future progress.

Acknowledgments

We warmly thank the reviewers for their meticulous suggestions, which have improved the exposition in this paper.

2. Motivation: log-concavity and the fundamental gap

In the case of Dirichlet boundary data, the log-concavity of the first eigenfunction is a key step in proving the lower bound of the gap between the two smallest eigenvalues [1]. In the case that the first Robin eigenfunction is log-concave, then a similar bound holds. Here we note that we can include a potential, and since we impose the strong hypothesis that the first eigenfunction is log-concave, we do not need to assume that the potential is convex. In the Dirichlet case, the sharp bound is found by making a more quantitative log-concavity estimate (in essence, showing that the ground state in the general case is at least as log-concave as it is for the one-dimensional case); in this case, as we merely assume log-concavity, the lower bound is smaller by a factor of three.

Theorem 2.1. *Let λ_0 and λ_1 be the two smallest eigenvalues for the eigenvalue problem*

$$-\Delta u + Vu = \lambda u \text{ in } \Omega,$$

with Robin boundary conditions (1.2) on a bounded convex domain Ω with diameter D , and potential $V \in L^\infty(\Omega)$. If the ground state u_0 associated to λ_0 is log-concave, then

$$(2.1) \quad \lambda_1 - \lambda_0 \geq \frac{\pi^2}{D^2}.$$

Remark 2.2. It is worth mentioning that the assumption $V \in L^\infty(\Omega)$ in Theorem 2.1 is only needed to ensure that the Schrödinger operator $H = -\Delta u + Vu$ admits a discrete spectrum.

Proof. Let u_0 and u_1 be the eigenfunctions associated to λ_0 and λ_1 respectively. Since u_0 is positive on Ω , we can set

$$v(x, t) := \frac{e^{-\lambda_1 t} u_1(x)}{e^{-\lambda_0 t} u_0(x)}$$

which solves the parabolic equation

$$(2.2) \quad \frac{\partial v}{\partial t} = \Delta v + 2D \log u_0 \cdot Dv \quad \text{on } \Omega \times (0, +\infty).$$

On the lateral boundary $\partial\Omega \times (0, +\infty)$, the normal derivative of v disappears:

$$D_\nu v = \frac{e^{-\lambda_1 t}}{e^{-\lambda_0 t}} \left(\frac{D_\nu u_1}{u_0} - \frac{u_1 D_\nu u_0}{u_0^2} \right) = v(-\alpha + \alpha) = 0.$$

By hypothesis, u_0 is log-concave, so the drift term in (2.2) given by $X := 2D \log u_0$ satisfies the *modulus of contraction* inequality

$$\left(X(y, t) - X(x, t) \right) \cdot \frac{y - x}{|y - x|} \leq 0$$

corresponding to the modulus of contraction $\omega \equiv 0$. Therefore by [1, Theorem 2.1], for some large constant $C > 0$, the function

$$\varphi(s, t) := C e^{-\frac{\pi^2}{D^2} t} \sin\left(\frac{\pi s}{D}\right) \quad \text{for every } s \in [0, D/2], t \geq 0,$$

is a *modulus of continuity* for v , that is,

$$v(y, t) - v(x, t) \leq 2\varphi\left(\frac{y-x}{|y-x|}, t\right) \quad \text{for every } x, y \in \overline{\Omega}, t \geq 0,$$

where $\frac{\pi^2}{D^2}$ is the second (or the difference of the second and first) Neumann eigenvalue on the interval. From this, we can deduce that

$$e^{-(\lambda_1 - \lambda_0)t} \operatorname{osc}_{\overline{\Omega}} \left(\frac{u_1}{u_0} \right) \leq C e^{-\frac{\pi^2}{D^2} t} \quad \text{for all } t \geq 0,$$

which can only hold if inequality (2.1) holds. This completes the proof of Theorem 2.1. \square

The above argument follows the approach used in the Dirichlet case [1]. A similar result would follow using the gradient estimate approach of [21, 25].

The resulting estimate is sharp in the case $\alpha = 0$, where it is the Payne-Weinberger inequality for the first nontrivial Neumann eigenvalue [19, 26]. Otherwise, it is not sharp, as can be seen from the one dimensional case, where the eigenvalues can be computed. It is appealing to conjecture that the sharp lower bound for given α and D should correspond to the gap for the corresponding one-dimensional problem, which would result in an estimate which depends on α and increases from $\frac{\pi^2}{D^2}$ to $\frac{3\pi^2}{D^2}$ as α increases from 0 towards infinity. However, our main theorem (Theorem 1.3, that the ground state is in general not log-concave) means that a sharp result must necessarily be proved by rather different means.

3. The Robin eigenvalue problem and perturbations

In the first proposition of this section, we summarize some properties of the first Robin eigenvalue λ_α and the corresponding eigenfunction u_α . All statements, except for (iv) concerning the C^1 -dependence respect to the topology induced by the Hölder-norm $C^{0,\beta}(\Omega)$, are quite well-known; see, for example, [16, Theorem 1.3.1] or [13]. In fact, they follow from an application of Kato’s perturbation theory for linear operators (more precisely, see [15, Theorem 2.6 of Chapter 8.2]). The remaining part follows from [18, Theorem 3.14]. We omit the proofs here, but we provide them in the online preprint version [2].

Proposition 3.1. *Let Ω be a connected bounded Lipschitz domain in \mathbb{R}^d . Then*

- (i) *For every $\alpha \in \mathbb{R}$, there is a first Robin eigenvalue $\lambda_\alpha \in \mathbb{R}$ with a positive eigenfunction $u_\alpha \in H^1(\Omega)$.*
- (ii) *For every $\alpha \in \mathbb{R}$, the first Robin eigenvalue λ_α is simple.*
- (iii) *The function $\alpha \mapsto \lambda_\alpha$ is differentiable, with derivative given by*

$$\dot{\lambda}_\alpha = \frac{\int_{\partial\Omega} u_\alpha^2 \, d\mathcal{H}}{\int_\Omega u_\alpha^2 \, dx} \geq 0.$$

- (iv) *The first Robin eigenfunction u_α (normalised to have $\frac{1}{|\Omega|} \int_\Omega u_\alpha^2 \, dx = 1$) is C^1 -dependent on α in $H^1(\Omega)$ and in $C^{0,\beta}(\Omega)$ for some $\beta \in (0, 1)$. More precisely, u_α is continuously dependent on α in $H^1(\Omega)$ and in $C^{0,\beta}(\Omega)$, and if for $\alpha_0 \in \mathbb{R}$, v is the unique solution, orthogonal to u_{α_0} in $L^2(\Omega)$, of*

$$(3.1) \quad \begin{cases} \Delta v + \lambda_{\alpha_0} v = -\dot{\lambda}_{\alpha_0} u_{\alpha_0} & \text{in } \Omega, \\ D_\nu v + \alpha_0 v = -u_{\alpha_0} & \text{on } \partial\Omega, \end{cases}$$

then $u_\alpha = u_{\alpha_0} + v(\alpha - \alpha_0) + o(\alpha - \alpha_0)$ for every α in a neighbourhood of α_0 , where $o(\alpha - \alpha_0)/(\alpha - \alpha_0) \rightarrow 0$ in $H^1(\Omega) \cap C^{0,\beta}(\Omega)$ as $\alpha \rightarrow \alpha_0$.

Next, we state a convergence result on Robin problems on varying domains. For this, we need to recall the definition of the *Hausdorff complementary topology on open sets* (cf [7, Section 2]). For closed subsets F_1, F_2 in \mathbb{R}^d , the *Hausdorff metric* $d_{\mathcal{H}}$ is defined by

$$d_{\mathcal{H}}(F_1, F_2) = \max \left\{ \sup_{x \in F_1} \text{dist}(x, F_2), \sup_{x \in F_2} \text{dist}(x, F_1) \right\},$$

where $\text{dist}(x, F_i) := \inf_{y \in F_i} |x - y|$, with the conventions $\text{dist}(x, \emptyset) = \infty$ and $\sup \emptyset = 0$, so that $d_{\mathcal{H}}(\emptyset, F) = 0$ if $F = \emptyset$ and $d_{\mathcal{H}}(\emptyset, F) = \infty$ otherwise. Let $\Omega^c = \mathbb{R}^d \setminus \Omega$ be the complement of Ω . Now, a sequence $(\Omega_n)_{n \geq 1}$ of open sets Ω_n in \mathbb{R}^d converges to the open set Ω in \mathbb{R}^d in the Hausdorff complementary topology if for every closed ball B in \mathbb{R}^d , one has that $d_{\mathcal{H}}(B \cap \Omega_n^c, B \cap \Omega^c) \rightarrow 0$ as $n \rightarrow \infty$. We write this as $\Omega_n \rightarrow \Omega$ in \mathcal{H}^c .

The next preliminary result is an application of [7, Theorem 3.4 and Corollary 3.5] and the convergence (3.2) with respect to the Hölder-norm follows from [18, Theorem 3.14].

Proposition 3.2. *For $d \geq 1$, let $D \subseteq \mathbb{R}^d$ be an open and bounded set, and let Ω and Ω_n be open domains with a Lipschitz continuous boundary satisfying $\Omega, \Omega_n \subset\subset D$. Let*

$$\Omega_n \rightarrow \Omega \text{ in } \mathcal{H}^c, \quad |\Omega_n| \rightarrow |\Omega|, \quad \mathcal{H}^{d-1}(\partial\Omega_n) \rightarrow \mathcal{H}^{d-1}(\partial\Omega)$$

as $n \rightarrow +\infty$. In particular, for $\alpha > 0$, let $\lambda_{\alpha,n}$ and λ_α be the first Robin eigenvalue on Ω_n and Ω , and let $u_{\alpha,n}$ and u_α be the first positive Robin eigenfunctions with unit $L^2(\Omega)$ -norm. Then

$$\begin{aligned} \lambda_{\alpha,n} &\rightarrow \lambda_\alpha \quad \text{as } n \rightarrow +\infty, \\ u_{\alpha,n} \mathbf{1}_{\Omega_n} &\rightarrow u_\alpha \mathbf{1}_\Omega \quad \text{in } L^2(D) \text{ as } n \rightarrow +\infty, \\ Du_{\alpha,n} \mathbf{1}_{\Omega_n} &\rightarrow Du_\alpha \mathbf{1}_\Omega \quad \text{in } L^2(D) \text{ as } n \rightarrow +\infty. \end{aligned}$$

Furthermore, there are $\gamma \in (0, 1)$ and $C > 0$ such that

$$\|u_{\alpha,n}\|_{C^{0,\gamma}(\overline{\Omega_n})} \leq C \quad \text{for all } n \geq 1,$$

and for every non-empty set $B \subseteq \bigcap_{n \geq n_0} \overline{\Omega_n}$, $n_0 \geq 1$, and $0 \leq \hat{\gamma} < \gamma$, there is a subsequence $(u_{\alpha,k_n})_{k_n \geq 1}$ of $(u_{\alpha,n})_{n \geq 1}$ such that

$$(3.2) \quad u_{\alpha,k_n} \rightarrow u_\alpha \quad \text{in } C^{0,\hat{\gamma}}(B) \text{ as } n \rightarrow +\infty.$$

4. Regular solutions are quadratic

When $\alpha_0 = 0$, the perturbation problem (3.1) reduces to equation (1.5), with the constant μ computed by integrating the first equation over Ω and applying the boundary condition, yielding $\mu = \mathcal{H}^{d-1}(\partial\Omega)/\mathcal{H}^d(\Omega)$.

In this and the next several sections we consider a class of problems generalising (1.5), under the assumption that Ω is a convex polyhedral domain in \mathbb{R}^d for $d \geq 2$.

Remark 4.1. A special feature of polyhedral domains is that for every $x \in \overline{\Omega}$ there exists $r > 0$ for which $B_r(x) \cap \Omega = x + (B_r(0) \cap \Gamma_x)$; in other words, near x , Ω is a cone.

We now establish a version of the strong maximum principle on general open cones Γ with a Lipschitz boundary. In this paper, we apply Proposition 4.2 on cones with a polyhedral structure.

Proposition 4.2. *Let Γ be an open cone with Lipschitz boundary and vertex at the origin in \mathbb{R}^d , and $R > 0$. Then for every weak solution $w \in H^1(B_R(0) \cap \Gamma)$ of problem*

$$(4.1) \quad \begin{cases} \Delta w = 0 & \text{on } B_R(0) \cap \Gamma, \\ D_\nu w = 0 & \text{on } B_R(0) \cap \partial\Gamma, \end{cases}$$

the following statements hold.

1. The mean value

$$\overline{w}(r) = \int_A w(rz) \, d\mathcal{H}^{d-1}(z)$$

of $w(r \cdot)$ over the set $A = \Gamma \cap \mathbb{S}^{d-1}$, $r \in [0, R)$, is a solution of the 2nd-order linear ODE

$$(4.2) \quad \frac{1}{r^{d-1}} \left\{ r^{d-1} \overline{w}_r \right\}_r = 0 \quad \text{on } (0, R)$$

and hence, \overline{w} is given by

$$(4.3) \quad \overline{w}(r) = \begin{cases} c_1 \mathbb{1} + c_2 \log r & \text{if } d = 2, \\ c_1 \mathbb{1} + c_2 \frac{1}{r^{d-2}} & \text{if } d \geq 3, \end{cases}$$

for constants $c_1, c_2 \in \mathbb{R}$.

2. If $w(0) = 0$ and $w \leq 0$ on $B_R(0) \cap \Gamma$, then $w \equiv 0$ on $B_R(0) \cap \Gamma$.

Our proof uses a similar idea as in the proof of [11, Proposition 6.1, see also Lemma 6.2].

Proof. We begin by recalling that the mean operator $\bar{\cdot} : L^2(A) \rightarrow \mathbb{R}$ assigning $u \mapsto \bar{u}$ is the linear projection of the function u to the space of constant functions on A , which we simply denote by \mathbb{R} . But \mathbb{R} also coincides with the eigenspace $\ker(-\Delta^{\mathbb{S}^{d-1}})$ of the first eigenvalue of the negative Neumann Laplace-Beltrami operator $-\Delta^{\mathbb{S}^{d-1}}$ on A . Thus, for $r \in [0, R)$, $w(r\cdot) - \bar{w}(r)$ belongs to the orthogonal complement of $\ker(-\Delta^{\mathbb{S}^{d-1}})$ which is $L_0^2(A) := \{u \in L^2(A) \mid \bar{u} = 0\}$, the space of functions $u \in L^2(A)$ with zero mean over A . Since the space $L_0^2(A)$ is invariant under the Neumann Laplace-Beltrami operator $\Delta^{\mathbb{S}^{d-1}}$ on A , one has that

$$(4.4) \quad 0 = \int_A \Delta^{\mathbb{S}^{d-1}}(w(rz) - \bar{w}(r)) \, d\mathcal{H}^{d-1}(z) = \int_A \Delta^{\mathbb{S}^{d-1}} w(rz) \, d\mathcal{H}^{d-1}(z)$$

for every $r \in [0, R)$. By hypothesis, w is harmonic in $B_R(0) \cap \Gamma$, which in polar coordinates is

$$0 = \Delta w = \frac{1}{r^{d-1}} \left\{ r^{d-1} w_r \right\}_r + \Delta^{\mathbb{S}^{d-1}} w \quad \text{on } B_R(0) \cap \Gamma.$$

Now, taking the average over A of the last equation and using (4.4), one sees that \bar{w} is a solution of the 2nd-order linear ODE (4.2). Thus, there are constants $c_1, c_2 \in \mathbb{R}$ such that (4.3) holds, completing the proof of claim (1).

To prove claim (2), we first note that by elliptic regularity theory [18], each weak solution w of (4.1) is continuous on $\overline{B_{R/2}(0) \cap \Gamma}$ and hence, $w(0)$ and $\bar{w}(0)$ exist. In particular, if $w(0) = 0$ then $\bar{w}(0) = 0$ and hence in (4.3) we have $c_1 = c_2 = 0$. Thus, $w(r\cdot)$ has zero mean for all $r \in [0, R)$. But, if in addition, $w \leq 0$ on $B_R(0) \cap \Gamma$, then $w(r\cdot)$ has zero mean if and only if $w \equiv 0$ on $B_R(0) \cap \Gamma$. This completes the proof of this proposition. \square

Although we are mostly interested in the perturbation problem (1.5), the results of this section and the next also apply for a somewhat larger class: We consider (weak) solutions v of the problem

$$(4.5) \quad \begin{cases} \Delta v + \mu = 0 & \text{in } \Omega, \\ D_{\nu_i} v + \gamma_i = 0 & \text{on } \Sigma_i, \end{cases}$$

where μ and $\gamma_1, \dots, \gamma_m$ are constants. We observe (by integration of the first equation over Ω and application of the boundary condition on each face

Σ_i) that these constants necessarily satisfy the relation

$$\sum_{i=1}^m \gamma_i \mathcal{H}^{d-1}(\Sigma_i) = \mu \mathcal{H}^d(\Omega).$$

The main result of this section is the following:

Theorem 4.3. *Let Ω be a polyhedral domain in \mathbb{R}^d with faces $\Sigma_1, \dots, \Sigma_m$ and v be a solution of (4.5). If $v \in C^2(\overline{\Omega})$ then v is quadratic; that is, there are constants $a_{ij}, b_i, c \in \mathbb{R}$ such that*

$$v(x) = \sum_{i,j=1}^d a_{ij} x_i x_j + \sum_{i=1}^d b_i x_i + c \quad \text{for every } x \in \overline{\Omega}.$$

Our strategy to prove Theorem 4.3 is to show that there exists a subspace E in \mathbb{R}^d on which the Hessian function $(x, e) \mapsto D^2v|_x(e, e)$ is constant for all unit vectors $e \in E$ and $x \in \Omega$. It will follow from this that $v(x)$ is a multiple of the squared length of the E component of x , plus another function depending only on the E^\perp component, where E^\perp denotes the orthogonal complement of E in \mathbb{R}^d . This reduces the original problem to a similar problem on the lower-dimensional space E^\perp , enabling an induction on dimension to establish the result.

Accordingly, we proceed by induction: For $d = 1$, a polyhedral domain is simply an interval, and every solution to (4.5) is a quadratic function, so the statement of Theorem 4.3 holds in this case. Now, assume that the statement of Theorem 4.3 holds for every polyhedral domain in \mathbb{R}^j for $j < d$, and let Ω be a polyhedral domain in \mathbb{R}^d and $v \in C^2(\overline{\Omega})$ be a solution of (4.5) on Ω . Since $v \in C^2(\overline{\Omega})$, there exists $(x_0, e_1) \in \overline{\Omega} \times \mathbb{S}^{d-1}$ such that

$$\Lambda := \max_{x \in \overline{\Omega}, e \in \mathbb{S}^{d-1}} D^2v|_x(e, e) = D^2v|_{x_0}(e_1, e_1).$$

Lemma 4.4. *Suppose that v is a C^2 function on an open subset B of $\overline{\Omega}$, where Ω is a polyhedral domain in \mathbb{R}^d . For $j \in \{1, \dots, m\}$, let ν_j be the outward pointing unit normal vector on face Σ_j and suppose*

$$(4.6) \quad D_{\nu_j} v + \gamma_j = 0 \quad \text{on } \overline{\Sigma_j} \cap B.$$

Then for every tangent vector e parallel to Σ_j one has

$$(4.7) \quad D^2v|_x(e, \nu_j) = 0 \quad \text{for every } x \in \overline{\Sigma_j} \cap B.$$

In particular, ν_j is an eigenvector for the Hessian $D^2v|_x$ for each $x \in \overline{\Sigma}_j \cap B$.

In the following, if Σ_j is a face of the polyhedral domain Ω then we write $T\Sigma_j$ for the *tangent space* of Σ_j ; or more precisely, it is the linear vector space spanned by the linear independent vectors tangent to Σ_j .

Proof. On polyhedra, the normal vector ν_j is constant on face Σ_j . Differentiating the boundary condition (4.6) in the direction of any tangent vector $e \in T\Sigma_j$ yields (4.7). Since \mathbb{R}^d can be decomposed as a direct sum of the tangent space $T\Sigma_j$ and the normal vector ν_j , (4.7) implies that ν_j is an eigenvector for the Hessian $D^2v|_x$ for $x \in \overline{\Sigma}_j \cap B$. \square

Our second lemma captures in slightly greater generality the dimension-reduction argument outlined above:

Lemma 4.5. *Suppose that v is a C^2 solution of (4.5) on a convex open subset B of $\overline{\Omega}$, where Ω is a polyhedral domain in \mathbb{R}^d . If there exists (x_0, e_1) in $B \times \mathbb{S}^{d-1}$ such that*

$$(4.8) \quad D^2v|_{x_0}(e_1, e_1) = \Lambda := \sup_{(x,e) \in B \times \mathbb{S}^{d-1}} D^2v|_x(e, e),$$

then there exists a subspace E of positive dimension in \mathbb{R}^d such that

$$B \cap \Omega = \left\{ x \in B \mid \pi_E(x) \in \Omega^E, \pi_{E^\perp}(x) \in \Omega^\perp \right\},$$

where E^\perp is the orthogonal complement of E , π_E and π_{E^\perp} are the orthogonal projections onto E and E^\perp , and $\Omega^E = \pi_E(\Omega)$ and $\Omega^\perp = \pi_{E^\perp}(\Omega)$ are polyhedral domains in E and E^\perp respectively. Furthermore,

$$(4.9) \quad v(x) = \frac{\Lambda}{2} |\pi_E(x - x_0)|^2 + Dv|_{x_0}(\pi_E(x - x_0)) + g(\pi_{E^\perp}(x))$$

for all $x \in B$, where g is a C^2 solution of an equation of the form (4.5) on $\pi_{E^\perp}(B) \subseteq \overline{\Omega^\perp} \subseteq E^\perp$.

Proof. Without loss of generality, we can assume that we have chosen $x_0 \in B$ so that the dimension of the eigenspace of $H_v(x_0)$ with eigenvalue Λ is maximized. We begin by defining u to be the part of v without its *quadratic approximation* about x_0 :

$$(4.10) \quad u(x) := v(x) - v(x_0) - Dv|_{x_0}(x - x_0) - \frac{1}{2} D^2v|_{x_0}(x - x_0, x - x_0)$$

for every $x \in B$. Then u has the following properties:

$$(4.11) \quad \begin{aligned} u(x_0) &= 0, \quad Du(x_0) = 0, \quad D^2u|_{x_0} = 0; \\ Du|_x &= Dv|_x - Dv|_{x_0} - D^2v|_{x_0}(x - x_0, \cdot) \quad \text{for every } x \in B; \end{aligned}$$

$$(4.12) \quad \Delta u(x) = \Delta v(x) - \Delta v(x_0) = -\mu + \mu = 0 \quad \text{for every } x \in B \cap \Omega;$$

$$(4.13) \quad D_{\nu_j}u(x) = 0 \quad \text{for all } x \in \Sigma_j \cap B \quad \text{if } j \in \mathcal{I}(x_0),$$

where the index set $\mathcal{I}(x_0)$ is given by (1.4). To see that (4.13) holds, first note that this is trivially satisfied if $x_0 \notin \partial\Omega$, since then $\mathcal{I}(x_0)$ is empty. If $x_0 \in \partial\Omega$, then by Lemma 4.4, for every $j \in \mathcal{I}(x_0)$, v satisfies (4.7). If $x \in \Sigma_j \cap B$, then both x and x_0 lie in the same face Σ_j and so $(x - x_0) \in T\Sigma_j$. By taking $e = x - x_0$ and using (4.11) and (4.7), one has

$$D_{\nu_j}u(x) = D_{\nu_j}v(x) - D_{\nu_j}v(x_0) - D^2v|_{x_0}(e, \nu_j) = \gamma_j - \gamma_j = 0$$

for all $x \in \Sigma_j \cap B$. Now, let E be the eigenspace of $D^2v|_{x_0}$ corresponding to its largest eigenvalue Λ . Then, $e_1 \in E \cap \mathbb{S}^{d-1}$. We choose an orthonormal basis $\{e_1, e_2, \dots, e_k\}$ of E , $1 \leq k \leq d$, and set

$$f(x) := \text{tr}_E(D^2u|_x) = \sum_{i=1}^k D^2u|_x(e_i, e_i) \quad \text{for each } x \in B.$$

Then f has the following properties:

$$(4.14) \quad f(x) = \sum_{i=1}^k (D^2v|_x - D^2v|_{x_0})(e_i, e_i) \quad \text{for all } x \in B \text{ by (4.10);}$$

$$f(x_0) = 0 \quad \text{by the above;}$$

$$\Delta f(x) = 0 \quad \text{for every } x \in \Omega \cap B \text{ by (4.12);}$$

$$(4.15) \quad f(x) \leq 0 \quad \text{for every } x \in B;$$

$$(4.16) \quad D_{\nu_j}f(x) = 0 \quad \text{for every } x \in B \cap \Sigma_j, \text{ if } j \in \mathcal{I}(x_0).$$

To see that (4.15) holds, note that by (4.8),

$$D^2u|_x(\xi, \xi) = D^2v|_x(\xi, \xi) - D^2v|_{x_0}(\xi, \xi) = D^2v|_x(\xi, \xi) - \Lambda \leq 0$$

for all $\xi \in E \cap \mathbb{S}^{d-1}$ and $x \in B$. To show (4.16), fix $j \in \mathcal{I}(x_0)$. Then by Lemma 4.4 applied to v , the normal ν_j is an eigenvector of $D^2v|_x$ for $x \in \overline{\Sigma}_j$. On the interior of the face Σ_j , $v \in C^3(\Sigma_j)$ (since u extends by even

reflection in Σ_j as a harmonic function) and so we can differentiate (4.7) again to find

$$(4.17) \quad D^3v|_x(e, e, \nu_j) = 0 \quad \text{for every } e \in T\Sigma_j \text{ and } x \in \Sigma_j.$$

Since the normal ν_j is an eigenvector of $D^2v|_{x_0}$, and all eigenspaces of the matrix $D^2v|_{x_0}$ are orthogonal, the eigenvector ν_j is either in E or belongs to the orthogonal space E^\perp . If $\nu_j \in E^\perp$, then e_i is orthogonal to ν_j and so is in $T\Sigma_j$ for each $i \in \{1, \dots, k\}$. Then (4.17) implies

$$D_{\nu_j}f(x) = D_{\nu_j} \left(\sum_{i=1}^k D^2v|_x(e_i, e_i) \right) = 0$$

for every $x \in B \cap \Sigma_j$. On the other hand, if $\nu_j \in E$, then

$$\begin{aligned} D_{\nu_j}f(x) &= D_{\nu_j} \left(\sum_{i=1}^k D^2u|_x(e_i, e_i) \right) = D_{\nu_j} \left(\Delta u - \sum_{i=k+1}^d D^2u|_x(e_i, e_i) \right) \\ &= D_{\nu_j} \left(0 - \sum_{i=k+1}^d D^2u|_x(e_i, e_i) \right) = 0 \end{aligned}$$

for every $x \in B \cap \Sigma_j$, where $\{e_{k+1}, \dots, e_d\}$ is a basis for $E^\perp \subseteq \nu_j^\perp = T\Sigma_j$, and we again use (4.17).

By Remark 4.1, the set $\Omega \cap B \cap B_r(x_0)$ coincides with $x_0 + (\Gamma_{x_0} \cap B_r(0))$ for sufficiently small $r > 0$. Equations (4.14)-(4.16) (and that fact that f is continuous on B since $v \in C^2(B)$) allow us to apply Proposition 4.2 to the function $\tilde{f}(z) = f(x_0 + rz)$ on $B_1(0) \cap \Gamma_{x_0}$ to infer that f is identically zero on a neighbourhood of x_0 . We conclude that the set where f vanishes is a non-empty, open, and closed subset of B , hence equal to B . It follows from (4.10) that $\text{tr}_E D^2v \equiv k\Lambda$ on B . Since $D^2v \leq \Lambda I$ on B , this implies that $D^2v(e_i, e_i) = \Lambda$ on B for all $i = 1, \dots, k$ and so,

$$(4.18) \quad D^2v|_x(e, e) = \Lambda \quad \text{for all } x \in B \text{ and } e \in \mathbb{S}^{d-1} \cap E.$$

In particular E is contained in the Λ -eigenspace of $D^2v|_x$ for every $x \in B$. Since we chose $x_0 \in B$ such that k is the maximal dimension of the Λ -eigenspace of $D^2v|_x$ over all $x \in B$, we can conclude that E is the Λ -eigenspace of $D^2v|_x$ for every $x \in B$. It then also follows that

$$(4.19) \quad D^2v|_x(e, \hat{e}) = 0 \quad \text{for all } x \in B, e \in E, \text{ and } \hat{e} \in E^\perp.$$

Now, writing $x = \pi_E(x) + \pi_{E^\perp}(x)$, integrating (4.18) along directions in E yields

$$v(x) = v(\pi_E(x_0) + \pi_{E^\perp}(x)) + Dv(\pi_E(x_0)) + \pi_{E^\perp}(x) \pi_E(x - x_0) + \frac{\Lambda}{2} |\pi_E(x - x_0)|^2.$$

By (4.19), differentiating $Dv(\pi_E(x_0) + \pi_{E^\perp}(x))$ in a direction tangent to E^\perp gives zero, so $Dv(\pi_E(x_0) + \pi_{E^\perp}(x))$ is independent of $\pi_{E^\perp}(x)$ and in particular is equal to $Dv(x_0)$. Defining $g(\pi_{E^\perp}(x)) = v(\pi_E(x_0) + \pi_{E^\perp}(x))$ shows that v is of the form (4.9).

If $k = \dim(E) = d$ then E^\perp is trivial and there is nothing further to prove. Otherwise it follows that g is a C^2 function on $\pi_{E^\perp}(B) \subset \overline{\Omega^\perp}$, and we have

$$0 = \Delta v + \mu = \Delta g + k\Lambda + \mu$$

and for $\nu_i \in E^\perp$ we have

$$0 = D_{\nu_i} v + \gamma_i = D_{\nu_i} g + \gamma_i.$$

That is, g is a C^2 solution of an equation of the form (4.5) on the open subset $\pi_{E^\perp}(B)$ of $\overline{\Omega^\perp} \subseteq E^\perp$. By Lemma 4.4, ν_j is an eigenvector of $H_v(x)$ at every point $x \in \Sigma_j \cap B$, and hence the normals ν_j are either in E or E^\perp . Then we can write

$$\begin{aligned} \Omega \cap B &= \bigcap_{i=1}^m \left\{ x \in B \mid x \cdot \nu_i < b_i \right\} \\ &= \bigcap_{i: \nu_i \in E} \left\{ x \in B \mid x \cdot \nu_i < b_i \right\} \cap \bigcap_{i: \nu_i \in E^\perp} \left\{ x \in B \mid x \cdot \nu_i < b_i \right\} \\ &= \bigcap_{i: \nu_i \in E} \left\{ x \in B \mid \pi_E(x) \cdot \nu_i < b_i \right\} \cap \bigcap_{i: \nu_i \in E^\perp} \left\{ x \in B \mid \pi_{E^\perp}(x) \cdot \nu_i < b_i \right\} \\ &= \left\{ x \in B \mid \pi_E(x) \in \Omega^E, \pi_{E^\perp}(x) \in \Omega^\perp \right\}, \end{aligned}$$

where

$$\Omega^E = \bigcap_{i: \nu_i \in E} \left\{ x \in E \mid x \cdot \nu_i < b_i \right\} \text{ and } \Omega^\perp = \bigcap_{i: \nu_i \in E^\perp} \left\{ x \in E^\perp \mid x \cdot \nu_i < b_i \right\}.$$

This completes the proof of Lemma 4.5. □

Now, we can give the proof of Theorem 4.3:

Proof of Theorem 4.3. By Lemma 4.5 (applied with $B = \bar{\Omega}$), we have that v is of the form (4.9) for some solution g of (4.5) on Ω^\perp . If $k = \dim(E) = d$ then v is quadratic and there is nothing further to prove. Otherwise the function g is a C^2 solution of an equation of the form (4.5) on Ω^\perp in \mathbb{R}^{d-k} . By the inductive hypothesis, g is a quadratic function, and therefore v is also quadratic. This completes the induction and the proof of Theorem 4.3. \square

5. Tame domains

Our aim over the next several sections is to prove that concave solutions of (4.5) are twice continuously differentiable up to the boundary. The result of the previous section then implies that such solutions are quadratic functions.

Recall that a function f is *semi-concave* if there exists $C \in \mathbb{R}$ such that the function $x \mapsto f(x) - C|x|^2$ is concave.

Over the course of the next three sections we will prove the following:

Theorem 5.1. *Let Ω be a polyhedral domain in \mathbb{R}^d with faces $\Sigma_1, \dots, \Sigma_m$, and v be a weak solution of problem (4.5) for some $\mu, \gamma_1, \dots, \gamma_m \in \mathbb{R}$. If v is semi-concave in Ω , then $v \in C^2(\bar{\Omega})$.*

The main difficulty in proving that $v \in C^2(\bar{\Omega})$ is to understand the behaviour of v at points on the boundary $\partial\Omega$, particularly where two or more of the faces Σ_i intersect. We begin by using the series expansion (5.8) to understand the behaviour of v near a boundary point x_0 in terms of homogeneous Neumann harmonic functions on the tangent cone Γ_{x_0} . A crucial step in our argument will be to prove the result that homogeneous degree two Neumann harmonic functions must be quadratic if they have bounded second derivatives. We will accomplish this in the next section. In the rest of this section we will establish that this result is sufficient to prove regularity.

Definition 5.2. For given vectors $\nu_1, \dots, \nu_m \in \mathbb{R}^d$, a polyhedral cone

$$\Gamma = \bigcap_{i=1}^m \left\{ x \in \mathbb{R}^d \mid x \cdot \nu_i < 0 \right\}$$

is called *tame* if every degree two homogeneous harmonic function $v \in C^{1,1}(\bar{\Gamma})$ with homogeneous Neumann boundary condition on $\partial\Gamma$ is quadratic. If Ω is a polyhedral domain in \mathbb{R}^d and B is a relatively open subset of $\bar{\Omega}$, then B is called *tame* if the tangent cone Γ_x is tame for every $x \in B$.

The significance of tameness for our argument is captured by the following preliminary theorem which is the main result of this section.

Theorem 5.3. *Let Ω be a polyhedral domain in \mathbb{R}^d and B a relatively open tame subset of $\overline{\Omega}$. Then every weak solution $w \in C^{1,1}(B) \cap H^1(B)$ of problem*

$$(5.1) \quad \begin{cases} \Delta w = 0 & \text{on } \Omega \cap B, \\ D_\nu w = 0 & \text{on } \partial\Omega \cap B \end{cases}$$

is in $C^2(B)$.

Proof of Theorem 5.3. Since any harmonic function is of the class C^∞ at an interior point (here, $\Omega \cap B$), it remains to consider the case when $x_0 \in \partial\Omega \cap B$. We first establish that the harmonic function w is twice differentiable at each point $x_0 \in \partial\Omega \cap B$, using the decomposition (5.8). Since the restriction of B to a sufficiently small ball about x_0 agrees with a translate of the tangent cone to Ω at x_0 , it is sufficient to consider a Neumann harmonic function defined on a ball about the origin in a tame cone Γ . In other words, we study weak solutions w of problem (4.1) for $R = 1$.

We begin by setting $A = S^{d-1} \cap \Gamma$. Then the set $B_1(0) \cap \Gamma$ can be described by the *polar coordinate map*

$$(r, z) \in (0, 1) \times A \mapsto rz \in B_1(0) \cap \Gamma.$$

Since the set A is a Lipschitz domain in \mathbb{S}^{d-1} , there is a complete $L^2(A)$ -orthonormal set of eigenfunctions $\{\varphi_i\}_{i=0}^\infty$ for the Neumann Laplacian on A , with associated eigenvalues λ_i which we arrange in non-decreasing order with $\lambda_0 = 0$. Let $w \in H^1(B_1(0) \cap \Gamma)$. Then for every $r \in (0, 1)$, $w \in H^1(B_r(0) \cap \Gamma)$, the trace $w(r, \cdot)$ of w exists in $L^2(A)$. Using this, we see that w can be rewritten in polar coordinates as

$$(5.2) \quad w(rz) = \sum_{i=0}^\infty w_i(r) \varphi_i(z) \quad \text{for every } (r, z) \in [0, 1) \times A.$$

Here, for every $r \in (0, 1)$, the series (5.2) converges in $L^2(A)$ as the Fourier series of the trace of $w(r, \cdot)$ in $L^2(A)$ with i th Fourier coefficient

$$(5.3) \quad w_i(r) := (w(r \cdot), \varphi_i)_{L^2(A)}$$

for every $i \geq 1$. In order to continue the proof of Proposition 4.2, we first establish some properties of the series decomposition (5.2) of the weak solution w of (4.1). This is done in the next two statements.

Lemma 5.4. *Let Γ be an open cone with Lipschitz boundary and vertex at the origin in \mathbb{R}^d , and let $w \in H^1(B_1(0) \cap \Gamma)$ be a weak solution of Neumann problem (4.1). Then for all $i \geq 1$,*

$$(5.4) \quad f_i := \lim_{r \rightarrow 1} w_i(r)$$

exists, and furthermore the series $\sum_{i=0}^\infty \sqrt{1 + \lambda_i} f_i^2$ converges with

$$(5.5) \quad \sum_{i=0}^\infty \sqrt{1 + \lambda_i} f_i^2 \leq C \|w\|_{H^1(B_1(0) \cap \Gamma)}^2.$$

Proof. Due the classical regularity theory of elliptic equations and by possible dilation, there is no loss of generality in assuming that every weak solution $w \in H^1(B_1(0) \cap \Gamma)$ of Neumann problem (4.1) is continuous on $\overline{B_1(0) \cap \Gamma}$. Thus, the limit (5.4) exists for every $i \geq 1$. Since the gradient $\nabla w \in L^2(B_1(0) \cap \Gamma)$, one has that

$$w'_i(r) = \frac{dw_i}{dr}(r) = \int_A \nabla w(rz) \cdot z \varphi_i(z) \, d\mathcal{H}^{d-1}(z)$$

for every $r \in (0, 1)$. Thus the $H^1(B_1(0) \cap \Gamma)$ -norm of w can be written as

$$(5.6) \quad \begin{aligned} \|w\|_{H^1(B_1(0) \cap \Gamma)}^2 &= \int_{B_1(0) \cap \Gamma} \{w^2 + |\nabla w|^2\} \, dx \\ &= \sum_{i,j=0}^\infty \left\{ \int_0^1 (w_i w_j + w'_i w'_j) r^{d-1} \, dr \int_A \varphi_i \varphi_j \, d\mathcal{H}^{d-1} \right\} \\ &\quad + \sum_{i,j=1}^\infty \left\{ \int_0^1 w_i w_j r^{d-3} \, dr \int_A \nabla \varphi_i \cdot \nabla \varphi_j \, d\mathcal{H}^{d-1} \right\} \\ &= \sum_{i=0}^\infty \int_0^1 \left(\left(1 + \frac{\lambda_i}{r^2}\right) w_i^2 + (w'_i)^2 \right) r^{d-1} \, dr, \end{aligned}$$

where

$$\int_A \nabla \varphi_i \cdot \nabla \varphi_j \, d\mathcal{H}^{d-1} = - \int_A \Delta \varphi_i \varphi_j \, d\mathcal{H}^{d-1} = \lambda_i \int_A \varphi_i \varphi_j \, d\mathcal{H}^{d-1} = \lambda_i \delta_{ij}.$$

For $\delta \in (0, 1)$, consider the mapping $g : [\delta, 1) \rightarrow \mathbb{R}$ defined by

$$g(r) = \sum_{i=0}^{\infty} \sqrt{1 + \lambda_i} w_i^2(r) \quad \text{for every } r \in [\delta, 1),$$

and for every $N \geq 0$,

$$g_N(r) = \sum_{i=0}^N \sqrt{1 + \lambda_i} w_i^2(r) \quad \text{for every } r \in [\delta, 1).$$

Then, by (5.6), $g_N(r) \rightarrow g(r)$ as $N \rightarrow \infty$ for every $r \in [\delta, 1)$ and

$$\begin{aligned} |g_N(r_2) - g_N(r_1)| &\leq \sum_{i=0}^N \sqrt{1 + \lambda_i} |w_i^2(r_2) - w_i^2(r_1)| \\ &\leq \sum_{i=0}^N \sqrt{1 + \lambda_i} \int_{r_1}^{r_2} 2|w_i(r) w_i'(r)| \, dr \\ &\leq \sum_{i=0}^N \int_{r_1}^{r_2} \left(1 + \frac{\lambda_i}{r^2}\right) w_i^2 + \sum_{i=0}^{\infty} (w_i')^2 \, dr \\ &\leq C_\delta \sum_{i=0}^{\infty} \int_{r_1}^{r_2} \left(\left(1 + \frac{\lambda_i}{r^2}\right) w_i^2 + (w_i')^2 \right) r^{d-1} \, dr. \end{aligned}$$

Then, sending $N \rightarrow +\infty$ in the above inequality yields

$$(5.7) \quad |g(r_2) - g(r_1)| \leq C_\delta \sum_{i=0}^{\infty} \int_{r_1}^{r_2} \left(\left(1 + \frac{\lambda_i}{r^2}\right) w_i^2 + (w_i')^2 \right) r^{d-1} \, dr,$$

for every $0 < \delta < r_1 < r_2 < 1$. By (5.6), the right hand side in the last estimate of (5.7) tends to zero as $r_1, r_2 \rightarrow 1^-$. Hence, the Cauchy criterion implies that

$$\lim_{r \rightarrow 1^-} g(r) = \sum_{i=0}^{\infty} \sqrt{1 + \lambda_i} f_i^2 \quad \text{exists,}$$

where f_i is defined by (5.4). This shows that the function g is absolutely continuous on $[\delta, 1]$ for every $\delta \in (0, 1)$. By the mean value theorem for integrals, there is an $r_\delta \in (\delta, 1)$ satisfying

$$g(r_\delta) = \frac{1}{1-\delta} \int_\delta^1 g(r) \, dr = \frac{1}{1-\delta} \sum_{i=0}^{\infty} \int_\delta^1 \sqrt{1 + \lambda_i} w_i^2 \, dr \leq \frac{C_\delta}{1-\delta} \|w\|_{H^1(B_1(0) \cap \Gamma)}^2,$$

where we also used (5.7) and (5.6). Using this together with (5.7), one finds

$$g(r) = g(r) - g(r_\delta) + g(r_\delta) \leq C \|w\|_{H^1(B_1(0) \cap \Gamma)}^2$$

for some $C > 0$ independent of $r \in (\delta, 1)$. Sending $r \rightarrow 1$, we find (5.5). \square

Due to Lemma 5.4, every weak solution w of (4.1) has the following series expansion.

Proposition 5.5. *Let Γ be an open cone with Lipschitz boundary and vertex at the origin in \mathbb{R}^d , and follow the notation of Lemma 5.4. Then every weak solution $w \in H^1(B_1(0) \cap \Gamma)$ of (4.1) satisfies*

$$(5.8) \quad w(rz) = \sum_{i=0}^{\infty} f_i r^{\beta_i} \varphi_i(z) \quad \text{for every } z \in A \text{ and } r \in (0, 1),$$

where for every integer $i \geq 0$, $\beta_i > 0$ and a zero of the quadratic equation

$$(5.9) \quad \beta_i^2 + (d - 2)\beta_i - \lambda_i = 0.$$

There exists $\gamma_r \in (0, 1)$ such that the convergence of the series in (5.8) holds in $H^1(B_1(0) \cap \Gamma) \cap C^{\gamma_r}(\overline{B_r(0) \cap \Gamma})$ for every $0 < r < 1$.

Proof. For every $i \in \mathbb{N}_0$, we define

$$\psi_i(rz) := r^{\beta_i} \varphi_i(z) \quad \text{for every } rz \in B_1(0) \cap \Gamma.$$

Then ψ_i is harmonic on $B_1(0) \cap \Gamma$ since

$$\begin{aligned} \Delta(r^{\beta_i} \varphi_i(z)) &= r^{\beta_i-2} \Delta_{\mathbb{S}^{d-1}} \varphi_i + \frac{d-1}{r} \frac{\partial r^{\beta_i}}{\partial r} \varphi_i + \frac{\partial^2 r^{\beta_i}}{\partial r^2} \varphi_i \\ &= r^{\beta_i-2} (-\lambda_i + (d-2)\beta_i + \beta_i^2) \varphi_i \\ &= 0 \end{aligned}$$

by (5.9) and the fact that φ_i satisfies

$$\Delta_{\mathbb{S}^{d-1}} \varphi_i + \lambda_i \varphi_i = 0 \quad \text{on } A.$$

Furthermore, ψ_i satisfies Neumann boundary conditions on $B_1(0) \cap \partial\Gamma$, since φ_i satisfies Neumann conditions on ∂A . Thus, each ψ_i is a weak solution of (4.1).

Now, let $\tilde{w} : B_1(0) \cap \Gamma \rightarrow \mathbb{R}$ be given by

$$\tilde{w}(rz) := \sum_{i=0}^{\infty} f_i \psi_i(rz) = \sum_{i=0}^{\infty} f_i r^{\beta_i} \varphi_i(z) \quad \text{for every } rz \in B_1(0) \cap \Gamma,$$

where f_i is given by (5.4). Next, we show that the infinite series of \tilde{w} converges in $H^1(B^1(0) \cap \Gamma)$. For this, let \tilde{w}^N be the partial sum of \tilde{w} given by

$$\tilde{w}^N(rz) = \sum_{i=0}^N f_i r^{\beta_i} \varphi_i(z) \quad \text{for every } rz \in B_1(0) \cap \Gamma.$$

For integers $1 \leq M < N$, applying (5.6) to $\tilde{w}^N - \tilde{w}^M = \sum_{i=M+1}^N f_i r^{\beta_i} \varphi_i$, we find

$$\begin{aligned} \|\tilde{w}^N - \tilde{w}^M\|_{H^1(B_1(0) \cap \Gamma)}^2 &= \sum_{i=M+1}^N \int_0^1 \left(f_i^2 r^{2\beta_i+d-1} + \beta_i^2 f_i^2 r^{2\beta_i+d-3} \right) dr \\ &= \sum_{i=M+1}^N \left(\frac{1}{2\beta_i+d} + \frac{\beta_i^2}{2\beta_i+d-2} \right) f_i^2 \\ &\leq C \sum_{i=M+1}^N (\beta_i + 1) f_i^2 \\ &\leq C \sum_{i=M+1}^N \sqrt{1 + \lambda_i} f_i^2. \end{aligned}$$

Lemma 5.4 implies that the infinite series $\sum_{i=0}^{\infty} \sqrt{1 + \lambda_i} f_i^2$ is convergent, and so there is $\tilde{w} \in H^1(B_1(0) \cap \Gamma)$ such that $\tilde{w}^N \rightarrow \tilde{w}$ in $H^1(B_1(0) \cap \Gamma)$. Since every partial sum \tilde{w}^N is a weak solution of (4.1), the limit function \tilde{w} is also a weak solution of (4.1) and has L^2 -trace

$$\sum_{i=0}^{\infty} f_i \varphi_i \quad \text{on } A.$$

Since the same is true for w , we have $w = \tilde{w}$, proving that (5.8) holds in $H^1(B_1(0) \cap \Gamma)$. To obtain convergence of the series (5.2) in $C^{\gamma_r}(\overline{B_r(0) \cap \Gamma})$ for every $0 < r < 1$ with some $\gamma_r \in (0, 1)$, we employ a reflection argument in a small neighbourhood U of each boundary point of $B_r(0) \cap \partial\Gamma$ as in [18] and use the interior Hölder-regularity result [14, Theorem 8.24]. Further, we

can cover $\overline{B_r(0) \cap \Gamma} \setminus \partial\Gamma$ by finitely many balls and apply again the interior Hölder-regularity to w . Summarising, we see that for every $0 < r < 1$, there is a $\gamma_r \in (0, 1)$ such that the series (5.2) converges in $C_r^\gamma(\overline{B_r(0) \cap \Gamma})$. \square

Lemma 5.6. *Let Γ be a tame polyhedral cone in \mathbb{R}^d with outer unit face normals ν_1, \dots, ν_m , and let $B = B_1(0) \cap \overline{\Gamma}$, where $B_1(0)$ is the open unit ball in \mathbb{R}^d . Then there exist constants $C > 0$ and $\gamma \in (0, 1)$ depending only on Γ such that for every weak solution $w \in C^{1,1}(B) \cap H^1(B)$ of (5.1), there exists a linear functional $L : \mathbb{R}^d \rightarrow \mathbb{R}$ with $\{\nu_1, \dots, \nu_m\} \subseteq \ker(L)$ and a symmetric bilinear form $\mathbf{a} : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ with trace $\text{tr}(\mathbf{a}) := \sum_{i=1}^d a(e_i, e_i) = 0$ such that the following estimate holds:*

$$(5.10) \quad \left| w(x) - w(0) - L(x) - \frac{1}{2} \mathbf{a}(x, x) \right| \leq C \|w\|_{L^\infty(B \cap \Gamma)} |x|^{2+\gamma}$$

for every $x \in B_{1/2}(0) \cap \Gamma$. Consequently w has derivatives up to second order at $x = 0$, with $Dw|_0 = L$ and $D^2w|_0 = \mathbf{a}$.

Proof of Lemma 5.6. We only need to consider the case $d \geq 2$. By Proposition 5.5, w has the series decomposition (5.2). Since in the series (5.2), $\varphi_0 \equiv 1$ and $\beta_0 = 0$, we have $w(0) = f_0$. Thus, writing in polar coordinates $x = rz$ for $r > 0$ and $z \in \mathbb{S}^{d-1}$,

$$w(rz) = w(0) + \sum_{i>1} f_i r^{\beta_i} \varphi_i(z) \quad \text{for every } rz \in B \cap \Gamma.$$

The second derivatives $D^2\psi_i$ of $\psi_i(x) := |x|^{\beta_i} \varphi_i(x/|x|)$ are homogeneous of degree $(\beta_i - 2)$. In particular, for every i with $\beta_i < 2$, $D^2\psi_i$ is unbounded as $r = |x|$ approaches zero, except in the case where $\beta_i = 1$ and ψ_i is a linear function. Since $w \in C^{1,1}(B)$, the only non-zero ψ_i with $0 < \beta_i < 2$ are those with $\beta_i = 1$, and these form a linear function L . Those ψ_i satisfy homogeneous Neumann boundary conditions on $B \cap \partial\Gamma$, implying that $L(\nu_i) = 0$ for every $i = 1, \dots, m$. Now, defining

$$v(rz) := \sum_{\beta_i=2} f_i r^2 \varphi_i(z) \quad \text{for every } rz \in B \cap \Gamma,$$

one has that

$$(5.11) \quad w(rz) = w(0) + L(rz) + v(rz) + \sum_{\beta_i>2} f_i r^{\beta_i} \varphi_i(z)$$

for every $rz \in B \cap \Gamma$. The function v is harmonic and homogeneous of degree 2, satisfies $D_\nu v = 0$ on $\partial\Gamma$ and has bounded second derivatives since they

are given by limits of second derivatives of $w \in C^{1,1}(B)$ as $r \rightarrow 0^+$. Thus $v \in C^{1,1}(\bar{\Gamma})$. Since Γ is tame, v is a quadratic homogeneous function. Thus, there is a symmetric bilinear form \mathbf{a} on \mathbb{R}^d such that

$$v(x) = \frac{1}{2}\mathbf{a}(x, x) \quad \text{for every } x \in \bar{\Gamma}.$$

By the harmonicity of v , it follows that

$$0 = \Delta v(x) = \text{tr}(\mathbf{a}) \quad \text{for every } x \in B \cap \Gamma,$$

and since v satisfies homogeneous Neumann boundary conditions,

$$0 = D_{\nu_i} v(x) = Dv|_x(\nu_i) = \mathbf{a}(x, \nu_i) \quad \text{for every } x \in \Sigma_i \text{ and } i = 1, \dots, m.$$

Differentiating the last equality in any direction $e \in T_x \Sigma_i, = (\nu_i)^\perp$ gives

$$0 = \mathbf{a}(e, \nu_i) \quad \text{for every } e \perp \nu_i,$$

showing that ν_i is an eigenvector of \mathbf{a} .

Next, set $\beta_* = \min\{\beta_i > 2 : f_i \neq 0\}$. Then we need to show that the reminder term

$$(5.12) \quad S(rz) := \sum_{\beta_i > 2} f_i r^{\beta_i} \varphi_i(z) = r^{\beta_*} \sum_{\beta_i > 2} f_i r^{\beta_i - \beta_*} \varphi_i(z)$$

on the right-hand side in (5.11) converges pointwise for every $x = rz \in B \cap \Gamma$. To see this, we note that by the definition of f_i ((5.3)–(5.4)) and since $\|\varphi\|_{L^2(A)} = 1$, one has that

$$(5.13) \quad |f_i| \leq \|w\|_{L^\infty(B \cap \Gamma)} \quad \text{for every } i \geq 1,$$

and by (5.9), there are $i_0 \in \mathbb{N}$ and $M_1, M_2 >$ such that

$$(5.14) \quad M_1 \sqrt{\lambda_i} \leq \beta_i \leq M_2 \sqrt{\lambda_i} \quad \text{for all } i \geq i_0.$$

Thus, by [9, Corollary 1], there is a constant $C_1 = C_1(d) > 0$ such that

$$(5.15) \quad \|\varphi_i\|_{L^\infty(A)} \leq C \lambda_i^{\frac{d-1}{4}} \leq \frac{C}{M_1} \beta_i^{\frac{d-1}{2}} \quad \text{for every } i \geq 1.$$

Further, by [20, Theorem 4.12] and (5.14), there are $i_* \in \mathbb{N}$ and $C = C(d, |A|) > 0$ such that

$$(5.16) \quad \beta_i \geq M_1 \sqrt{\lambda_i} \geq C i^{\frac{1}{d-1}} \quad \text{for all } i \geq i_*,$$

where we denote the constant $M_1 C(d, |A|)$ again by C . Now, let $0 < r < 1$. Then there is an $N(r) > 2$ such that the function $f(\beta) := \beta^{\frac{d-1}{2}} r^{\beta-\beta_*}$ is monotonically decreasing on $[N(r), +\infty)$. If we choose the index i_* in (5.16) a bit larger such that $C i^{\frac{1}{d-1}} \geq N(r)$ for all $i \geq i_*$, then by applying (5.13), (5.15) and (5.16) to (5.12), one sees that

$$\begin{aligned} \left| \sum_{\beta_i > 2} f_i r^{\beta_i - \beta_*} \varphi_i(z) \right| &\leq C_1 \|w\|_{L^\infty(B \cap \Gamma)} \sum_{i \geq i_*} \beta_i^{\frac{d-1}{2}} r^{\beta_i - \beta_*} \\ &\leq C_1 \|w\|_{L^\infty(B \cap \Gamma)} \sum_{i \geq i_*} \sqrt{i} r^{C i^{\frac{1}{d-1}} - \beta_*}, \end{aligned}$$

where infinite series $\sum_{i \geq i_*} \sqrt{i} r^{C i^{\frac{1}{d-1}} - \beta_*}$ is finite by the classical integral test. This shows that the series S in (5.12) converges pointwise on $B \cap \Gamma$, and uniformly on $B_{1/2}(0) \cap \Gamma$. In particular, S is bounded on $B_{1/2}(0) \cap \Gamma$ by $C_{1/2} \|w\|_{L^\infty(B \cap \Gamma)}$ for some constant $C_{1/2} > 0$. Applying this to (5.11) and noting that $\beta_* > 2$ yields the desired estimate (5.10). The fact that $Dw(0) = L$ and $D^2w(0) = \mathbf{a}$ follows from this estimate. \square

Continuation of the Proof of Theorem 5.3. The remaining difficulty in the proof of Theorem 5.3 is to confirm continuity of the second derivative. As before in Lemma 5.6, it suffices to consider a Neumann harmonic function w on a cone, and to establish the continuity of the second derivative at the origin. Accordingly, we fix a point x_0 in $\partial\Omega \cap B$, and $r_0 > 0$ sufficiently small to ensure that

$$\Omega \cap B_{r_0}(x_0) = \left\{ x_0 + x \mid x \in \Gamma_{x_0}, |x| < r_0 \right\},$$

where Γ_{x_0} is the tangent cone to Ω at x_0 . To show that the second derivatives of w are continuous at x_0 , it is sufficient to show that the Neumann harmonic function

$$\hat{w}(x) = \frac{u(x_0 + r_0 x)}{\|w\|_{L^\infty(B_{r_0}(x_0) \cap \Gamma_{x_0})}} \quad \text{for every } x \in B \cap \Gamma$$

has continuous second derivative at the origin, where $B = B_1(0)$ is the open unit ball and Γ a polyhedral cone with vertex at the origin.

Now, we label parts of Γ according to the number of faces which intersect. Recall the faces of Ω are Σ_i with outward unit normal vectors ν_i for every

$i = 1, \dots, m$. Then

$$\Gamma^{(k)} := \bigcup_{\substack{S \subset \{1, \dots, m\} \\ |S|=k}} \left(\bigcap_{i \notin S} \{x \mid x \cdot \nu_i \leq 0\} \right) \cap \left(\bigcap_{j \in S} \{x \mid x \cdot \nu_j = 0\} \right)$$

denotes the set of all $x \in \bar{\Gamma}$ where k faces intersect. Thus $\Gamma^{(0)} = \bar{\Gamma}$, $\Gamma^{(1)} = \partial\Gamma$, and $0 \in \Gamma^{(m)}$.

We now proceed by (decreasing) induction on k , starting with $k = m$:

Proposition 5.7. *Let Γ be a tame polyhedral cone in \mathbb{R}^d . Then there exist constants $C > 0$ and $\gamma \in (0, 1)$ depending only on Γ such that for every weak solution $w \in C^{1,1}(B_1(0) \cap \bar{\Gamma}) \cap H^1(B_1(0) \cap \bar{\Gamma})$ of (5.1),*

$$(5.17) \quad \begin{aligned} |w(y) - w(x) - Dw|_x(y - x) - \frac{1}{2}D^2w|_x(y - x, y - x)| \\ \leq C \|w\|_{L^\infty(B_1 \cap \Gamma)} |y - x|^{2+\gamma} \end{aligned}$$

for every $x \in B_{1/2}(0) \cap \Gamma^{(m)}$ and $y \in B_1(0) \cap \bar{\Gamma}$.

For the proof of Proposition 5.7 we will use the following auxiliary result, which will be also useful several times later.

Lemma 5.8. *Let \mathbf{a} be a symmetric bilinear form and L a linear functional on \mathbb{R}^d , and let $c \in \mathbb{R}$. Define*

$$q(x) = \mathbf{a}(x, x) + L(x) + c \quad \text{for every } x \in \mathbb{R}^d.$$

If for $r > 0$ and $M \geq 0$, one has that $\sup_{x \in \bar{B}_r(0)} |q(x)| \leq M$, then $|c| \leq M$, $\|L\| \leq 2M/r$, and the eigenvalues λ_i of \mathbf{a} satisfy $|\lambda_i| \leq 2M/r^2$.

Proof. Choosing $x = 0$ gives $|c| \leq M$, implying that $|\mathbf{a}(x, x) + L(x)| \leq 2M$ for all $x \in \bar{B}_r(0)$. Further, for $x \in \bar{B}_r(0)$, we have (by replacing x by $-x$) that $|\mathbf{a}(x, x) - L(x)| \leq 2M$, and hence (taking sums and differences) $|a(x, x)| \leq 2M$ and $|L(x)| \leq 2M$. Thus, $|\lambda_i| \leq 2M/r^2$ follows by choosing x/r to be a normalised eigenvector of \mathbf{a} , and $\|L\| \leq 2M/r$ follows by choosing $x \in \partial B_r(0)$ with $L(x) = \|L\| |x|$. \square

In order to apply the lemma above, we need a suitable ball. This is provided by the following:

Lemma 5.9. *Let Ω be a bounded open convex set in \mathbb{R}^d . Then there exist $\sigma > 0$ and $R > 0$ such that for every $x \in \bar{\Omega}$ and every $r \in (0, R)$, there exists $\hat{x} \in \Omega$ such that the open ball $B_{\sigma r}(\hat{x})$ is contained in $B_r(x) \cap \Omega$.*

Proof. Let ρ_- be the inradius and x_- an incentre of Ω , and let ρ_+ be the circumradius of Ω . Then, for $R = 2\rho_+$ (so that Ω is included in $B_R(x)$ for any $x \in \overline{\Omega}$) and $\sigma = \frac{\rho_-}{R}$, one has that

$$(5.18) \quad B_{\sigma R}(x_-) = B_{\rho_-}(x_-) \subseteq \Omega = \Omega \cap B_R(x)$$

for any $x \in \overline{\Omega}$. Now, for fixed $x \in \overline{\Omega}$ and $r \in (0, R)$, let

$$T_\lambda(y) = x + \lambda(y - x) \quad \text{for } y \in \mathbb{R}^d \text{ and } \lambda = \frac{r}{R} \in (0, 1).$$

Since $T_\lambda(y) = (1 - \lambda)x + \lambda y$, convexity of Ω implies that $T_\lambda(\Omega) \subseteq \Omega$. Thus, by (5.18) and since $T_\lambda(B_R(x)) = B_r(x)$, one has that

$$B_{\sigma r}(x_-) = T_\lambda(B_{\sigma R}(x_-)) \subseteq T_\lambda(\Omega \cap B_R(x))$$

and

$$T_\lambda(\Omega \cap B_R(x)) = T_\lambda(\Omega) \cap T_\lambda(B_R(x)) \subseteq \Omega \cap B_r(x),$$

as claimed. □

With these preliminaries, we can prove the base case of our (decreasing) induction.

Proof of Proposition 5.7. For $x \in B_{1/2}(0) \cap \Gamma^{(m)}$, the tangent cone Γ_x to Ω at x agrees with Γ at the origin. Thus, we can apply Lemma 5.6 to the function

$$w^x(\hat{x}) = w\left(x + \frac{\hat{x}}{2}\right) \quad \text{for every } \hat{x} \in B_1 \cap \Gamma$$

and obtain that

$$|w^x(\hat{x}) - w^x(0) - Dw^x|_0(\hat{x}) - \frac{1}{2}D^2w^x|_0(\hat{x}, \hat{x})| \leq C \|w\|_{L^\infty(B_1 \cap \Gamma)} |\hat{x}|^{2+\gamma}$$

for all $\hat{x} \in B_{1/2}(0) \cap \Gamma_x$. Now, setting $\hat{x} = 2(y - x)$ for $y \in B_{1/4}(x) \cap \Gamma$ and using the definition of w^x we obtain that estimate (5.17) holds for all $y \in B_{1/4}(x) \cap \Gamma$. To derive the same inequality for $y \in B_1(0) \setminus B_{1/4}(x)$, we first derive bounds on the size of $Dw|_x$ and $D^2w|_x$, using Lemma 5.8: by Lemma 5.9 applied to $\Omega = B \cap \Gamma$ and $r = 1/4$, there are $\sigma > 0$ and $x_* \in B \cap \Gamma$ such that the open ball $B_{\sigma/4}(x_*)$ is contained in $B_{1/4}(x) \cap \Gamma$. Due to estimate (5.17) and since w is bounded on $\overline{B_{\sigma/4}(x_*)}$, there is a $C > 0$ such that

$$\sup_{y \in \overline{B_{\sigma/4}(x_*)}} \left| Dw|_x(y - x) + \frac{1}{2}D^2w|_x(y - x, y - x) \right| \leq C \|w\|_{L^\infty(B_1 \cap \Gamma)}.$$

For $y \in \overline{B}_{\sigma/4}(x_*)$, setting $\hat{x} = y - x_*$, this shows that the quadratic function

$$q(\hat{x}) := Dw|_x(\hat{x} + x_* - x) + \frac{1}{2}D^2w|_x(\hat{x} + x - x_*, \hat{x} + x - x_*)$$

is bounded on $\overline{B}_{\sigma/4}(0)$ and hence by Lemma 5.8, the coefficients of q are bounded. Moreover, the quadratic part of q gives that the eigenvalues $\lambda_i(x)$ of $D^2w|_x$ satisfy $|\lambda_i(x)| \leq 32C/\sigma^2$. Since $D^2w^x|_0 = \frac{1}{4}D^2w|_x$ and $D^2w^x|_0$ is symmetric by Lemma 5.6, the Hessian $D^2w|_x$ is symmetric and so, the bound on $\lambda_i(x)$ implies that

$$(5.19) \quad \|D^2w|_x\| \leq 32C \frac{\|w\|_{L^\infty(B_1 \cap \Gamma)}}{\sigma^2}.$$

Further, the linear part of q gives that

$$\|Dw|_x + D^2w|_x(x_* - x)\| \leq 8C \frac{\|w\|_{L^\infty(B_1 \cap \Gamma)}}{\sigma}$$

and since (5.19) and $|x - x_*| < 1/4$, this yields that

$$\|Dw|_x\| \leq 16C \frac{\|w\|_{L^\infty(B_1 \cap \Gamma)}}{\sigma}.$$

Now, if $y \in B_1(0) \setminus B_{1/4}(x)$, then we have $\frac{1}{4} \leq |y - x| \leq \frac{3}{2}$, and so, the bounds on $w(y)$, $w(x)$, $Dw|_x$, $D^2w|_x$, $|y - x|$ and $|y - x|^{-1}$ show that

$$\begin{aligned} |w(y) - w(x) - Dw|_x(y - x) - \frac{1}{2}D^2w|_x(y - x, y - x)| \\ \leq C \|w\|_{L^\infty(B_1 \cap \Gamma)} \\ \leq C \|w\|_{L^\infty(B_1 \cap \Gamma)} |y - x|^{2+\gamma}, \end{aligned}$$

as required. □

Next, we establish the inductive step:

Proposition 5.10. *Let Γ be a tame polyhedral cone in \mathbb{R}^d . If a weak solution $w \in C^{1,1}(B_1(0) \cap \overline{\Gamma}) \cap H^1(B_1(0) \cap \overline{\Gamma})$ of (5.1) satisfies*

$$(5.20) \quad |w(y) - w(x) - Dw|_x(y - x) - \frac{1}{2}D^2w|_x(y - x, y - x)| \lesssim |y - x|^{2+\gamma}$$

for every $x \in B_{1/2}(0) \cap \Gamma^{(k)}$, $y \in B_1(0) \cap \overline{\Gamma}$ and some $\gamma \in (0, 1)$, then w also satisfies (5.20) for all $x \in B_{1/2}(0) \cap \Gamma^{(k-1)}$ and $y \in B_1(0) \cap \overline{\Gamma}$.

To prove this proposition, we intend to apply Lemma 5.6 about $x \in (B_{1/2}(0) \cap \Gamma^{(k-1)}) \setminus \Gamma^{(k)}$. In order to do this we need to estimate the *cone radius*

$$(5.21) \quad \rho(x) := \sup \left\{ r > 0 \mid B_r(x) \cap \bar{\Gamma} = x + (B_r(0) \cap \bar{\Gamma}_x) \right\},$$

where Γ is a polyhedral cone in \mathbb{R}^d with vertex at the origin and Γ_x the tangent cone to Γ at $x \in \partial\Gamma \setminus \{0\}$. This is supplied by the following result.

Lemma 5.11. *Let Γ be a tame polyhedral cone in \mathbb{R}^d . Then there exists a $\sigma > 0$ such that*

$$(5.22) \quad \rho(x) \geq \sigma d(x, \Gamma^{(k)}) \quad \text{for all } x \in \Gamma^{(k-1)} \setminus \Gamma^{(k)}.$$

To show that Lemma 5.11 holds, we need to introduce the following definition and the subsequent lemma.

Definition 5.12. A convex cone Γ in \mathbb{R}^d admits a linear factor E if there exists a linear subspace E of \mathbb{R}^d of positive dimension with orthogonal complement E^\perp in \mathbb{R}^d and a convex cone $\tilde{\Gamma}$ in E^\perp such that

$$\Gamma = \left\{ x \in \mathbb{R}^d \mid \pi_{E^\perp}(x) \in \tilde{\Gamma} \right\},$$

where π_{E^\perp} is the orthogonal projection onto E^\perp . In this situation, we write $\Gamma = \tilde{\Gamma} \oplus E$.

The following observation is used in the inductive step of our argument, and will also be used again later.

Lemma 5.13. *Let Γ be a polyhedral cone in \mathbb{R}^d with vertex at the origin and outer unit face normals ν_1, \dots, ν_m . Let $x_0 \in \partial\Gamma \setminus \{0\}$. Then the tangent cone Γ_{x_0} to Γ at x_0 has a linear factor $\mathbb{R}x_0$, and so had the form $\Gamma_{x_0} = \tilde{\Gamma} \oplus \mathbb{R}x_0$, where $\tilde{\Gamma}$ is the polyhedral cone in the $(d - 1)$ -dimensional subspace $(\mathbb{R}x_0)^\perp$ of \mathbb{R}^d defined by*

$$(5.23) \quad \tilde{\Gamma} = \bigcap_{i \in \mathcal{I}(x_0)} \left\{ x \in (\mathbb{R}x_0)^\perp \mid x \cdot \nu_i < 0 \right\},$$

where $\mathcal{I}(x_0) := \left\{ i \in \{1, \dots, m\} \mid x_0 \cdot \nu_i = 0 \right\}$.

Proof. Since

$$\Gamma_{x_0} = \bigcap_{i \in \mathcal{I}(x_0)} \left\{ x \in \mathbb{R}^d \mid x \cdot \nu_i < 0 \right\},$$

and $i \in \mathcal{I}(x_0)$ implies $\nu_i \cdot x_0 = 0$, we have that $\nu_i \in (\mathbb{R}x_0)^\perp$ for all $i \in \mathcal{I}(x_0)$. Therefore $\Gamma_{x_0} = \tilde{\Gamma} \oplus \mathbb{R}x_0$, where $\tilde{\Gamma}$ is given by (5.23). \square

Proof of Lemma 5.11. If there is no such $\sigma > 0$ such that (5.22) holds, then there exists a sequence $(x_n)_{n \geq 1}$ of points $x_n \in \Gamma^{(k-1)} \setminus \Gamma^{(k)}$ such that

$$(5.24) \quad \frac{\rho(x_n)}{d(x_n, \Gamma^{(k)})} \rightarrow 0.$$

Since both $\rho(\cdot)$ and $d(\cdot, \Gamma^{(k)})$ are homogeneous of degree one, we can scale x_n so that $x_n \in \mathbb{S}^{d-1} \cap \bar{\Gamma}$.

We first exclude the possibility that there are $\alpha > 0$ and a subsequence $(x_{n'})_{n' \geq 1}$ of $(x_n)_{n \geq 1}$ such that $d(x_{n'}, \Gamma^{(k)}) \geq \alpha$ for all $n' \geq 1$. Otherwise, for such a subsequence $(x_{n'})_{n' \geq 1}$ of $(x_n)_{n \geq 1}$, one has that $\rho(x_{n'}) \rightarrow 0$. Since $x_{n'} \in \mathbb{S}^{d-1} \cap (\Gamma^{(k-1)} \setminus \Gamma^{(k)})$, we can extract another subsequence of $(x_{n'})_{n' \geq 1}$ which we denote, for simplicity, again by $(x_{n'})_{n' \geq 1}$ such that $x_{n'}$ converges to a point $\bar{x} \in \mathbb{S}^{d-1} \cap \Gamma^{(k-1)} \setminus \Gamma^{(k)}$. Label the faces so that $\bar{x} \cdot \nu_i$ is in non-increasing order. Then, since $\bar{x} \in \Gamma^{(k-1)} \setminus \Gamma^{(k)}$, we have $\bar{x} \cdot \nu_i = 0$ for $i = 1, \dots, k-1$ and $\bar{x} \cdot \nu_k < 0$. Since the function $x \mapsto x \cdot \nu_i$ is continuous, any point x in $\Gamma^{(k-1)} \setminus \Gamma^{(k)}$ sufficiently close to \bar{x} also satisfies $x \cdot \nu_i = 0$ for $i = 1, \dots, k-1$ and $x \cdot \nu_i < \frac{1}{2}\bar{x} \cdot \nu_k < 0$ for $i \geq k$. It follows that

$$\Gamma_x = \Gamma_{\bar{x}} = \bigcap_{i=1}^{k-1} \left\{ z \mid z \cdot \nu_i < 0 \right\},$$

so the tangent cone is constant and hence the cone radius ρ is continuous on $\Gamma^{(k-1)}$ near \bar{x} . In particular, we have that $\rho(x_{n'})$ is bounded below, contradicting the fact that $\rho(x_{n'}) \rightarrow 0$.

The remaining possibility is that $d(x_n, \Gamma^{(k)})$ converges to zero. Passing to a subsequence, we have convergence to a point $\bar{x} \in \mathbb{S}^{d-1} \cap \Gamma^{(k)}$. In particular for n sufficiently large $x_n \in B_{\rho(\bar{x})}(\bar{x}) \cap \bar{\Gamma}$.

In Lemma 5.13, we have observed that since $\bar{x} \neq 0$, the tangent cone $\Gamma_{\bar{x}}$ is the product $\Gamma_{\bar{x}} = \tilde{\Gamma} \oplus \mathbb{R}\bar{x}$, where $\tilde{\Gamma}$ is a polyhedral cone in the $(d-1)$ -dimensional subspace $(\mathbb{R}\bar{x})^\perp$. Thus, it follows that $\rho(x_n)$ and $d(x_n, \Gamma^{(k)})$ are invariant under translation in the \bar{x} -direction and homogeneous of degree

one under rescaling about \bar{x} . Therefore, we can replace x_n by

$$\tilde{x}_n = \frac{\left(x_n - \frac{x_n \cdot \bar{x}}{|\bar{x}|^2} \bar{x}\right)}{\left|x_n - \frac{x_n \cdot \bar{x}}{|\bar{x}|^2} \bar{x}\right|} \in \left(\tilde{\Gamma} \times \{0\}\right) \cap \mathbb{S}^{d-1}$$

and still have a sequence $(\tilde{x}_n)_{n \geq 1}$ satisfying $\tilde{x}_n \in \tilde{\Gamma} \cap (\Gamma^{(k-1)} \setminus \Gamma^{(k)})$ and (5.24), where x_n is replaced by \tilde{x}_n .

Now, we repeat the above argument inductively, with Γ replaced by $\tilde{\Gamma}$. At each application, the dimension of the cone reduces by one, which is impossible since Γ is finite-dimensional. This contradicts our assumption that there is no positive σ satisfying the statement of Lemma 5.11, so the proof of the Lemma is complete. \square

Now, we can complete the proof of the inductive step.

Proof of Proposition 5.10. Fix $x \in (B_{1/2}(0) \cap \Gamma^{(k-1)}) \setminus \Gamma^{(k)}$. Let $\tilde{x} \in \Gamma^{(k)}$ be the closest point to x in $\Gamma^{(k)}$ satisfying $|x - \tilde{x}| < 1/2$. We claim that $\tilde{x} \in B_{1/2}(0)$. As $\lambda \tilde{x}$ is in $\Gamma^{(k)}$ for $\lambda > 0$, $g(\lambda) := |x - \lambda \tilde{x}|^2$ is minimised at $\lambda = 1$, and so $0 = g'(1) = -2(x - \tilde{x}) \cdot \tilde{x}$. Since $x - \tilde{x}$ and \tilde{x} are orthogonal,

$$|x|^2 = |x - \tilde{x} + \tilde{x}|^2 = |x - \tilde{x}|^2 + |\tilde{x}|^2 \geq |\tilde{x}|^2$$

and since $|x| < 1/2$, it follows that $|\tilde{x}| < 1/2$ as claimed. Hence, by hypothesis, w satisfies (5.20) at \tilde{x} . More precisely,

$$(5.25) \quad |w(y) - w(\tilde{x}) - Dw|_{\tilde{x}}(y - \tilde{x}) - \frac{1}{2}D^2w|_{\tilde{x}}(y - \tilde{x}, y - \tilde{x})| \leq C|y - \tilde{x}|^{2+\gamma}$$

for all $y \in B_1(0) \cap \bar{\Gamma}$ for some constant $C > 0$ and $\gamma \in (0, 1)$. To make use of this, we define

$$\tilde{w}(y) := w(y) - w(\tilde{x}) - Dw|_{\tilde{x}}(y - \tilde{x}) - \frac{1}{2}D^2w|_{\tilde{x}}(y - \tilde{x}, y - \tilde{x})$$

for every $y \in B_1(0) \cap \bar{\Gamma}$. Then \tilde{w} is a weak solution of (5.1) on $B_1(0) \cap \Gamma$ and by (5.25),

$$(5.26) \quad |\tilde{w}(y)| \leq C|y - \tilde{x}|^{2+\gamma} \quad \text{for } y \in B_1(0) \cap \bar{\Gamma}.$$

To proceed, we will apply Lemma 5.6 about x . But first note that by $\tilde{x} \in \Gamma^{(k)}$, after a possible re-ordering, we may assume without loss of generality that $\tilde{x} \cdot \nu_i = 0$ for all $i = 1, \dots, k$ and since $x \in \Gamma^{(k-1)} \setminus \Gamma^{(k)}$, there must be an

$1 \leq i_0 \leq k$ such that $x \cdot \nu_{i_0} < 0$. Now, let $\rho(x)$ be the cone radius around x given by (5.21) and we claim that

$$(5.27) \quad \rho(x) \leq |x - \tilde{x}|.$$

If $\rho(x) > |x - \tilde{x}|$, then there is an $\varepsilon > 0$ such that

$$x + (B_{(1+\varepsilon)|x-\tilde{x}|}(0) \cap \bar{\Gamma}_x) = B_{(1+\varepsilon)|x-\tilde{x}|}(x) \cap \bar{\Gamma}$$

and since $\tilde{x} \in B_{(1+\varepsilon)|x-\tilde{x}|}(x) \cap \bar{\Gamma}$, there is a $v \in B_{|x-\tilde{x}|}(0) \cap \bar{\Gamma}_x$ such that $v = \tilde{x} - x$. Then $x + (1+\varepsilon)v \in x + (B_{(1+\varepsilon)|x-\tilde{x}|}(0) \cap \bar{\Gamma}_x)$ and hence, $x + (1+\varepsilon)v \in \bar{\Gamma}$. However,

$$(x + (1 + \varepsilon)v) \cdot \nu_{i_0} = x \cdot \nu_{i_0} + (1 + \varepsilon)(\tilde{x} - x) \cdot \nu_{i_0} = -\varepsilon x \cdot \nu_{i_0} > 0,$$

which contradicts the definition of Γ , proving our claim (5.27). Since $|x - \tilde{x}| < 1/2$,

$$\hat{w}(y) := \tilde{w}(x + y\rho(x)) \quad \text{for } y \in B_1(0) \cap \Gamma_x$$

is a well-defined function. Moreover, \hat{w} is a weak solution of (5.1) on $B_1(0) \cap \Gamma_x$. Hence, by Lemma 5.6, there is a $\gamma \in (0, 1)$ and a $C > 0$ such that

$$|\hat{w}(y) - \hat{w}(0) - D\hat{w}|_0 y - \frac{1}{2}D^2\hat{w}|_0(y, y)| \leq C\|\hat{w}\|_{L^\infty(B_1(0) \cap \Gamma_x)}|y|^{2+\gamma}$$

for $y \in B_{1/2}(0) \cap \Gamma_x$. Note, by (5.26) and using (5.27),

$$(5.28) \quad \sup_{B_1(0) \cap \bar{\Gamma}_x} \hat{w} = \sup_{B_{\rho(x)}(x) \cap \bar{\Gamma}} \tilde{w} \leq \sup_{B_{2|x-\tilde{x}|}(\tilde{x}) \cap \bar{\Gamma}} \tilde{w} \leq C|x - \tilde{x}|^{2+\gamma}.$$

Combining the last two estimates then gives

$$|\hat{w}(y) - \hat{w}(0) - D\hat{w}|_0(y) - \frac{1}{2}D^2\hat{w}|_0(y, y)| \leq C|y|^{2+\gamma}|x - \tilde{x}|^{2+\gamma}$$

for $y \in B_{1/2}(0) \cap \Gamma_x$. By the definition of \hat{w} , this gives

$$\begin{aligned} |\tilde{w}(y) - \tilde{w}(x) - D\tilde{w}|_x(y - x) - \frac{1}{2}D^2\tilde{w}|_x(y - x, y - x)| \\ \leq C \left(\frac{|y - x|}{\rho(x)} \right)^{2+\gamma} |x - \tilde{x}|^{2+\gamma} \end{aligned}$$

for every $|y - x| < \frac{1}{2}\rho(x)$. Since by Lemma 5.11, there is a $\sigma > 0$ such that

$$(5.29) \quad \rho(x) \geq \sigma|x - \tilde{x}|,$$

we can conclude from the last estimate that

$$(5.30) \quad |\tilde{w}(y) - \tilde{w}(x) - D\tilde{w}|_x(y-x) - \frac{1}{2}D^2\tilde{w}|_x(y-x, y-x)| \leq C|y-x|^{2+\gamma}$$

for every $|y-x| < \frac{1}{2}\rho(x)$. From this, we deduce bounds on $D\tilde{w}|_x$ and $D^2\tilde{w}|_x$: By Lemma 5.9 applied to $\Omega = B_{1/2}(x) \cap \bar{\Gamma}$, there are $x_* \in B_{1/2}(x) \cap \bar{\Gamma}$ and $\sigma_* > 0$ such that the open ball $B_{\sigma_*\rho(x)}(x_*)$ is contained in $B_{\rho(x)}(x) \cap \bar{\Gamma}$. By (5.28), we have

$$|\tilde{w}(x)| + |\tilde{w}(y)| \leq C|x - \tilde{x}|^{2+\gamma} \quad \text{for every } y \in B_{\sigma_*\rho(x)}(x_*)$$

and so, by (5.30),

$$|D\tilde{w}|_x(y-x) + \frac{1}{2}D^2\tilde{w}|_x(y-x, y-x)| \leq C|x - \tilde{x}|^{2+\gamma}$$

for every $y \in B_{\sigma_*\rho(x)}(x_*)$. Moreover, from the previous application of Lemma 5.6 to \hat{w} , we know that the Hessian $D^2\hat{w}|_0 = \rho^{-2}(x)D^2\tilde{w}|_x$ is symmetric. Thus Lemma 5.8 yields that

$$(5.31) \quad \begin{aligned} & \|D\tilde{w}|_x(x_* - x) + \frac{1}{2}D^2\tilde{w}|_x(x_* - x)\| \leq C|x - \tilde{x}|^{2+\gamma} \\ & \|D\tilde{w}|_x + D^2\tilde{w}|_x(x_* - x)\| \leq \frac{2C|x - \tilde{x}|^{2+\gamma}}{\sigma_*\rho(x)} \leq C|x - \tilde{x}|^{1+\gamma}, \end{aligned}$$

$$(5.32) \quad \|D^2\tilde{w}|_x\| \leq 2\frac{4C|x - \tilde{x}|^{2+\gamma}}{\sigma_*^2\rho^2(x)} \leq C|x - \tilde{x}|^\gamma,$$

where we used the estimate (5.29) in the second inequalities of both (5.31) and (5.32). Since $|x - x_*| \leq C|x - \tilde{x}|$, inequality (5.31) implies that

$$(5.33) \quad \|D\tilde{w}|_x\| \leq C|x - \tilde{x}|^{1+\gamma}.$$

Next, we establish estimate (5.30) for $y \in (B_1(0) \setminus B_{\rho(x)/2}(x)) \cap \bar{\Gamma}$: On this set, we have $|x - \tilde{x}| + |y - \tilde{x}| \leq C|y - x|$ due to (5.29) and since $\rho(x)/2 \leq |y - x|$. Thus, by (5.26), (5.33), and (5.32),

$$\begin{aligned} & |\tilde{w}(y) - \tilde{w}(x) - D\tilde{w}|_x(y-x) - \frac{1}{2}D^2\tilde{w}|_x(y-x, y-x)| \\ & \leq |\tilde{w}(y)| + |\tilde{w}(x)| + \|D\tilde{w}|_x\||y-x| + \frac{1}{2}\|D^2\tilde{w}|_x\||y-x|^2 \\ & \leq C|y - \tilde{x}|^{2+\gamma} + C|x - \tilde{x}|^{2+\gamma} + C|x - \tilde{x}|^{1+\gamma}|y-x| + C|x - \tilde{x}|^\gamma|y-x|^2 \\ & \leq C|y-x|^{2+\gamma}, \end{aligned}$$

as required. This shows that estimate (5.30) holds for all $y \in B_1(0) \cap \bar{\Gamma}$. Finally, we note that \tilde{w} and w differ by a quadratic function, so

$$(5.34) \quad \begin{aligned} \tilde{w}(y) - \tilde{w}(x) - D\tilde{w}|_x(y-x) - \frac{1}{2}D^2\tilde{w}|_x(y-x, y-x) \\ = w(y) - w(x) - Dw|_x(y-x) - \frac{1}{2}D^2w|_x(y-x). \end{aligned}$$

Therefore inequality (5.20) holds for all $y \in B_1(0) \cap \bar{\Gamma}$ and $x \in B_{1/2}(0) \cap \Gamma^{(k-1)}$, and the proof of Proposition 5.10 is complete. \square

Completion of the Proof of Theorem 5.3. Now, Proposition 5.7 and Proposition 5.10 allow us to establish estimate (5.17) for all points $x \in B_{1/2}(0) \cap \bar{\Gamma}$ and all points $y \in B_1(0) \cap \bar{\Gamma}$, by (decreasing) induction on k : Due to Proposition 5.7, estimate (5.17) holds for $x \in \Gamma^{(m)}$, and by Proposition 5.10 if estimate (5.17) holds for $x \in \Gamma^{(k)}$ then it also holds for $x \in \Gamma^{(k-1)}$. Therefore, by induction, estimate (5.17) holds for all $x \in B_{1/2}(0) \cap \Gamma^{(0)} = B_{1/2}(0) \cap \bar{\Gamma}$. This allows us to complete the proof of Theorem 5.3 by proving that D^2w is continuous at the origin. So we must prove that $D^2w|_x$ approaches $D^2w|_0$ as $x \in B_{1/2}(0) \cap \bar{\Gamma}$ approaches zero. To do this, we apply estimate (5.17) about $x \in B_{1/2}(0) \cap \bar{\Gamma}$: Let

$$\tilde{w}(y) := w(y) - w(0) - Dw|_0(y) - \frac{1}{2}D^2w|_0(y, y)$$

for every $y \in B \cap \bar{\Gamma}$. Then by estimate (5.17),

$$(5.35) \quad |\tilde{w}(y)| \leq C \|w\|_{L^\infty(B_1 \cap \Gamma)} |y|^{2+\gamma} \quad \text{for every } y \in B_1(0) \cap \bar{\Gamma}.$$

Note, $D\tilde{w}|_y = Dw|_y - Dw|_0 - D^2w|_0(y, \cdot)$ and $D^2\tilde{w}|_y = D^2w|_y - D^2w|_0$. Hence, (5.34) holds and so, we can apply estimate (5.17) to w . Then, one finds

$$\begin{aligned} |\tilde{w}(y) - \tilde{w}(x) - D\tilde{w}|_x(y-x) - \frac{1}{2}D^2\tilde{w}|_x(y-x, y-x)| \\ \leq C \|w\|_{L^\infty(B_1 \cap \Gamma)} |y-x|^{2+\gamma} \end{aligned}$$

for every $x \in B_{1/2}(0) \cap \bar{\Gamma}$ and $y \in B_1(0) \cap \bar{\Gamma}$. From this together with (5.35), we deduce that

$$\sup_{y \in \bar{B}_{|x|}(x)} \left| D\tilde{w}|_x(y-x) + \frac{1}{2}D^2\tilde{w}|_x(y-x, y-x) \right| \leq C|x|^{2+\gamma}$$

for every $x \in B_{1/2}(0) \cap \bar{\Gamma}$. By Lemma 5.9 there is a ball of radius comparable to $|x|$ in $B_{|x|}(x) \cap \bar{\Gamma}$, and applying Lemma 5.8 on this ball gives that

$$|D\tilde{w}|_x(x) + \frac{1}{2}D^2\tilde{w}|_x(x, x)| \leq C|x|^{2+\gamma}, \quad \|D\tilde{w}|_x + D^2\tilde{w}|_x(x, \cdot)\| \leq C|x|^{1+\gamma},$$

and

$$(5.36) \quad \|D^2\tilde{w}|_x\| \leq C|x|^\gamma$$

for every $x \in B_{1/2}(0) \cap \bar{\Gamma}$. Since $D^2\tilde{w}|_x = D^2w|_x - D^2w|_0$, inequality (5.36) can be rewritten as

$$\|D^2w|_x - D^2w|_0\| \leq C|x|^\gamma \quad \text{for every } x \in B_{1/2}(0) \cap \bar{\Gamma},$$

proving that harmonic functions on a tame cone $B_1 \cap \Gamma$ satisfying homogeneous Neumann boundary condition on $B_1 \cap \partial\Gamma$ are $C^{2,\gamma}$. This completes the proof of Theorem 5.3. \square

6. Polyhedral cones are tame

Next, we prove the following, making the tameness hypothesis in Theorem 5.3 redundant.

Theorem 6.1. *Every polyhedral cone Γ in \mathbb{R}^d is tame.*

Proof. The proof uses an induction on the dimension $d \geq 1$, and uses the regularity results for tame domains established in the previous section. Our argument here is similar to that used in the proof of Proposition 4.3, in that we apply a strong maximum principle to the Hessian of the function. The homogeneity of the function allows us to consider points $x_0 \in \partial\Gamma$, which are not near the vertex of the cone, and this is the basis of the induction on dimension: We observe that by Lemma 5.13, the tangent cone is a direct product of a lower-dimensional cone with a line: $\Gamma_{x_0} = \tilde{\Gamma} \oplus \mathbb{R}x_0$, where $\tilde{\Gamma}$ is a polyhedral cone in the subspace $(\mathbb{R}x_0)^\perp$. To proceed, we need to understand the relationship between homogeneous harmonic functions on Γ_{x_0} and those on $\tilde{\Gamma}$:

Lemma 6.2. *Any homogeneous degree 2 Neumann harmonic function u on $\tilde{\Gamma} \oplus \mathbb{R}x_0$ has the form*

$$(6.1) \quad u(x + sx_0) = \tilde{u}(x) + s\tilde{v}(x) + C \left(s^2|x_0|^2 - \frac{1}{d-1}|x|^2 \right)$$

for $x \in \tilde{\Gamma}$, $s \in \mathbb{R}$, where \tilde{u} is a homogeneous degree 2 Neumann harmonic function on $\tilde{\Gamma}$, \tilde{v} is a homogeneous degree 1 Neumann harmonic function on $\tilde{\Gamma}$, and C is constant.

Proof. Without loss of generality, we may assume that $|x_0| = 1$. We choose an orthonormal basis for \mathbb{R}^d so that $x_0 = e_d$. Denote $A = (\tilde{\Gamma} \oplus \mathbb{R}e_d) \cap \mathbb{S}^{d-1}$, and $\tilde{A} = \tilde{\Gamma} \cap \mathbb{S}^{d-2}$. Then homogeneous degree 2 harmonic Neumann functions on $\tilde{\Gamma} \oplus \mathbb{R}e_d$ are determined by their restriction to A which is a Neumann eigenfunction. The corresponding eigenvalue is determined by the relation (5.9) which produces $\lambda_i = 2d$ when $\beta_i = 2$ (cf [8, Chapter 2.4]).

In the case $d = 2$, the cone $\tilde{\Gamma}$ cannot be $\mathbb{R}e_1$, since then Γ would be \mathbb{R}^2 , contradicting $x_0 \in \partial\Gamma$. Therefore $\tilde{\Gamma}$ is a ray in the direction of $\pm e_1$, and the cone $\tilde{\Gamma} \oplus \mathbb{R}x_0$ is congruent to the half-space $H = \{x > 0\}$ in \mathbb{R}^2 .

Any Neumann harmonic function u on H extends by even reflection to an entire harmonic function on \mathbb{R}^2 , which is therefore C^∞ . In particular a homogeneous degree 2 Neumann harmonic function on H is C^2 at the origin and therefore agrees with the degree 2 Taylor polynomial, since the second derivatives are homogeneous of degree zero, which must equal $C(x^2 - y^2)$. In this case, (6.1) is satisfied with $\tilde{v} \equiv \tilde{u} \equiv 0$.

Now, consider the case $d \geq 3$. We will construct eigenfunctions on A from eigenfunctions on \tilde{A} using separation of variables: We parametrise points of A by the map

$$\Phi : \tilde{A} \times \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \rightarrow \mathbb{S}^{d-1} \text{ given by } \Phi(z, \theta) = (\cos \theta) z + (\sin \theta) e_d,$$

for every $z \in \tilde{A}$ and $\theta \in [-\frac{\pi}{2}, \frac{\pi}{2}]$. The following construction is quite general (producing a basis of eigenfunctions on warped product spaces in terms of eigenfunctions on the warping factors), but we describe it here only in our specific situation.

The metric g induced by the map Φ on $\tilde{A} \times [-\pi/2, \pi/2]$ is


$$g = \cos^2 \theta \bar{g} + d\theta^2,$$

where \bar{g} is the metric on \mathbb{S}^{d-2} . The Laplacian in these coordinates is

$$\Delta^{\mathbb{S}^{d-1}} = \frac{1}{\cos^2 \theta} \Delta^{\mathbb{S}^{d-2}} - (d-2) \tan \theta \partial_\theta + \partial_\theta^2.$$

If φ is an eigenfunction on \tilde{A} satisfying $\Delta^{\mathbb{S}^{d-2}} \varphi + \mu \varphi = 0$ on \tilde{A} , then the function $f(\theta)\varphi(z)$ satisfies the eigenvalue equation

$$(6.2) \quad \mathcal{L}_\mu f := -f'' + (d-2) \tan \theta f' + \frac{\mu}{\cos^2 \theta} f = \lambda f \quad \text{on } A$$



with eigenvalue λ . Then $f\varphi$ is a Neumann eigenfunction of \mathcal{L}_μ on A provided φ satisfies homogeneous Neumann boundary conditions on $\partial\tilde{A}$ and $f\varphi$ extends continuously to the poles $\theta = \pm\frac{\pi}{2}$ of A . If φ is constant on \tilde{A} (corresponding to $\mu = 0$) then this amounts simply to the requirement that f extends continuously to $[-\pi/2, \pi/2]$, but if φ is non-constant (corresponding to $\mu > 0$) then continuity of $f\varphi$ at the poles amounts to the requirement that f has limit zero at $\pm\frac{\pi}{2}$. We note that the endpoints $\pm\pi/2$ are regular singular points of the ODE (6.2). Thus, a solution f to (6.2) satisfies the asymptotics

$$f(\theta) \sim \left(C_1(\theta + \pi/2)^{-\frac{d-3}{2} - \sqrt{(\frac{d-3}{2})^2 + \mu}} + C_2(\theta + \pi/2)^{-\frac{d-3}{2} + \sqrt{(\frac{d-3}{2})^2 + \mu}} \right)$$

as $\theta \rightarrow -\pi/2$, and

$$f(\theta) \sim \left(C_3(\pi/2 - \theta)^{-\frac{d-3}{2} - \sqrt{(\frac{d-3}{2})^2 + \mu}} + C_4(\pi/2 - \theta)^{-\frac{d-3}{2} + \sqrt{(\frac{d-3}{2})^2 + \mu}} \right)$$

as $\theta \rightarrow \pi/2$. Continuity requires that f has limit zero at $\pm\frac{\pi}{2}$, hence $C_1 = 0$ and $C_3 = 0$.

The operator \mathcal{L}_μ is essentially self-adjoint on $L^2((-\frac{\pi}{2}, \frac{\pi}{2}), (\cos \theta)^{d-2}d\theta)$. Accordingly, for any μ there is an increasing sequence of values $\lambda_{\mu,j}$ approaching infinity such that there is a solution $f_{\mu,j}$ of equation (6.2) satisfying the required endpoint conditions. These form a complete orthonormal basis for $L^2((-\frac{\pi}{2}, \frac{\pi}{2}), (\cos \theta)^{d-2}d\theta)$. We claim that if $\{\varphi_i\}_{i=0}^\infty$ is a complete orthonormal basis of Neumann eigenfunctions on \tilde{A} with eigenvalues μ_i , then the resulting collection of eigenfunctions $\{f_{\mu_i,j}(\theta)\varphi_i(z)\}$ forms a complete orthonormal basis of Neumann eigenfunctions on A . To see this, suppose that

g is a function in $L^2(\tilde{A} \times [-\pi/2, \pi/2], d\omega_{\tilde{g}}(\cos \theta)^{d-2} d\theta)$ which is orthogonal to $f_{\mu_i,j}(\theta)\varphi_i(z)$ for all i and j . That is, we have

$$\int_{-\pi/2}^{\pi/2} \int_{\tilde{A}} g(z, \theta)\varphi_i(z) d\omega_{\tilde{g}}(z) f_{\mu_i,j}(\theta)(\cos \theta)^{d-2} d\theta = 0$$

for all i and j . Fix i , and let $g_i(\theta) = \int_{\tilde{A}} g(z, \theta)\varphi_i(z) d\omega_{\tilde{g}}(z)$. Then g_i is orthogonal to $f_{\mu_i,j}$ in $L^2((-\frac{\pi}{2}, \frac{\pi}{2}), (\cos \theta)^{d-2} d\theta)$ for every j , and so vanishes almost everywhere. It follows that

$$g_i(\theta) = 0 \quad \text{for a.e. } \theta \in (-\frac{\pi}{2}, \frac{\pi}{2}) \text{ and every } i.$$

But this means that for a.e. $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, $g(\theta, \cdot)$ is orthogonal to φ_i in $L^2(\tilde{A}, d\omega_{\tilde{g}})$ for every i and hence, $g(\theta, z) = 0$ for almost all $z \in \tilde{A}$. This proves that $g = 0$ almost everywhere, proving completeness.

It follows that an eigenfunction on A with eigenvalue $\lambda = 2d$ is a finite linear combination of terms of the form $f_{\mu_i,j}(\theta)\varphi_i(z)$ for which $\lambda_{\mu_i,j} = 2d$.

Lemma 6.3. *For $\lambda = 2d$, solutions f_μ of (6.2) with the required boundary conditions*

$$f_\mu \sim C^\pm (\pi/2 - |\theta|)^{-\frac{d-3}{2} + \sqrt{(\frac{d-3}{2})^2 + \mu}} \text{ as } \theta \rightarrow \pm\pi/2$$

exist only for $\mu = 0$, $\mu = d - 2$ and $\mu = 2(d - 1)$, and these are given by $f_0(\theta) = \sin^2 \theta - \frac{1}{d-1} \cos^2 \theta$, $f_{d-2}(\theta) = \sin \theta \cos \theta$, and $f_{2(d-1)}(\theta) = \cos^2 \theta$.

Proof. The particular solutions given are constructed from homogeneous degree two spherical harmonics (harmonic polynomials on \mathbb{R}^d). These arise from the above construction in the case $\tilde{A} = \mathbb{S}^{d-2}$, and so give rise to solutions of (6.2). On \mathbb{S}^{d-1} , we have $x_d = \sin \theta$ and $|x| = \cos \theta$, where $x = (x_1, \dots, x_{d-1})$.

Therefore, the harmonic function

$$x_d^2 - \frac{1}{d-1}|x|^2 \quad \text{restricts to} \quad f_0(\theta) = \sin^2 \theta - \frac{1}{d-1} \cos^2 \theta \text{ on } \mathbb{S}^{d-1}.$$

The restriction of this to \mathbb{S}^{d-2} is constant, hence an eigenfunction with eigenvalue $\mu = 0$ on \mathbb{S}^{d-2} . It follows that $\mathcal{L}_0 f_0 + 2df_0 = 0$.

The harmonic function

$$x_d x_1 \quad \text{restricts to} \quad \sin \theta \cos \theta \frac{x_1}{|x|} = f_{d-2}(\theta)\varphi \left(\frac{x}{|x|} \right) \text{ on } \mathbb{S}^{d-1},$$

where $\varphi(x) = x_1$ is a homogeneous degree one harmonic function on \mathbb{R}^{d-1} , hence an eigenfunction of the Laplacian on \mathbb{S}^{d-2} with eigenvalue $\mu = d - 2$. It follows that $\mathcal{L}_{d-2}f_{d-2} + 2df_{d-2} = 0$.

Finally, the harmonic function

$$x_2^2 - x_1^2 \text{ restricts to } f_{2(d-1)}(\theta)\varphi\left(\frac{x}{|x|}\right) \text{ on } \mathbb{S}^{d-1},$$

where $f_{2(d-1)}(\theta) = \cos^2 \theta$ and $\varphi(x) = x_2^2 - x_1^2$, which is the restriction to \mathbb{S}^{d-2} of a degree 2 homogeneous harmonic function on \mathbb{R}^{d-1} , hence an eigenfunction of the Laplacian on \mathbb{S}^{d-2} with eigenvalue $\mu = 2(d - 1)$. It follows that $\mathcal{L}_{2(d-1)}f_{2(d-1)} + 2df_{2(d-1)} = 0$, as required. These formulae can be checked by explicit computation.

The harder part of the proof is to show that these are the only solutions of (6.2) with the required boundary conditions. It is convenient to perform a transformation of equation (6.2) to de-singularise the endpoints at $\pm\pi/2$. To do this we introduce the new variable s by

$$\tanh(s/2) = \tan(\theta/2),$$

so that $s \in \mathbb{R}$ increases over the entire real line as θ increases from $-\pi/2$ to $\pi/2$. This choice implies that $\frac{d\theta}{ds} = \cos \theta$, and we have the identities $\cos \theta = \frac{1}{\cosh s}$, $\sin \theta = \tanh(s)$ and $\tan \theta = \sinh s$. The equation (6.2) transforms to

$$0 = f_{ss} - (d - 3) \tanh s f_s + \left(\frac{2d}{\cosh^2 s} - \mu\right) f.$$

Defining $f = (\cosh s)^{\frac{d-3}{2}} g$ then produces the equation

$$(6.3) \quad 0 = g_{ss} + \left(\frac{(d+1)(d+3)}{4 \cosh^2 s} - \left(\frac{d-3}{2}\right)^2 - \mu\right) g.$$

The behaviour at $\theta = \pm\pi/2$ translates to the condition that g is asymptotic to $C_2 e^{s\sqrt{(\frac{d-3}{2})^2 + \mu}}$ as $s \rightarrow -\infty$ and to $C_4 e^{-s\sqrt{(\frac{d-3}{2})^2 + \mu}}$ as $s \rightarrow \infty$.

Next, we consider the Riccati equation associated to the ODE (6.2), which is the first order ODE satisfied by the function $q = \frac{g_s}{g}$:

$$\partial_s q = \frac{g_{ss}}{g} - \left(\frac{g_s}{g}\right)^2$$

$$= \mu + \left(\frac{d-3}{2}\right)^2 - \frac{(d+1)(d+3)}{4 \cosh^2 s} - q^2.$$

The boundary conditions then become the requirement that

$$\lim_{s \rightarrow -\infty} q(s) = \sqrt{\left(\frac{d-3}{2}\right)^2 + \mu}$$

and

$$\lim_{s \rightarrow +\infty} q(s) = -\sqrt{\left(\frac{d-3}{2}\right)^2 + \mu}.$$

The function q approaches infinity whenever the value of g crosses zero. We remove these singularities by defining a new variable σ which gives (twice) the angle from the positive x axis of the point $(g(s), g_s(s))$, so that $\tan(\sigma/2) = g_s(s)/g(s) = q$. This is defined only modulo 2π , but a continuous choice of σ exists and is uniquely defined up to an integer multiple of 2π . It follows from the definition that $\tan(\sigma/2) = q$, and we deduce that

$$(6.4) \quad \sigma_s = (1 + \cos \sigma) \left(\mu + 1 + \left(\frac{d-3}{2}\right)^2 - \frac{(d+1)(d+3)}{4 \cosh^2 s} \right) - 2.$$

From the asymptotic conditions on q , our construction requires a solution σ such that

$$\sigma(s) \rightarrow \sigma_-(\mu) := 2 \arctan \left(\sqrt{\left(\frac{d-3}{2}\right)^2 + \mu} \right)$$

as $s \rightarrow -\infty$, and $\sigma(s) \rightarrow \sigma_+(\mu)$ modulo $2\pi\mathbb{Z}$ as $s \rightarrow \infty$, where

$$\sigma_+(\mu) := -2 \arctan \left(\sqrt{\left(\frac{d-3}{2}\right)^2 + \mu} \right).$$

For each μ there is a unique solution $\sigma_\mu(s)$ of (6.4) with $\sigma_\mu(s) \rightarrow \sigma_-(\mu)$ as $s \rightarrow -\infty$ (arising from the solutions of (6.2) with the required asymptotics near $\theta = -\pi/2$ provided by the theory of regular singular points). It remains to find those values of μ for which σ_μ has the required behaviour as $s \rightarrow \infty$.

The crucial property we require is monotonicity of $\sigma_\mu(s)$ with respect to μ for each s :

Suppose $\mu_2 > \mu_1 \geq 0$. Then we observe that $\sigma_{\mu_1}(x)$ satisfies

$$\begin{aligned} \partial_s \sigma_{\mu_1} &= (1 + \cos \sigma) \left(\mu_1 + 1 + \left(\frac{d-3}{2} \right)^2 - \frac{(d+1)(d+3)}{4 \cosh^2 s} \right) - 2 \\ &\leq (1 + \cos \sigma) \left(\mu_2 + 1 + \left(\frac{d-3}{2} \right)^2 - \frac{(d+1)(d+3)}{4 \cosh^2 s} \right) - 2, \end{aligned}$$

so that solutions of (6.4) for $\mu = \mu_2$ cannot cross σ_{μ_1} from above. But now for s sufficiently negative we have $\sigma_{\mu_1}(s)$ as close as desired to $\sigma_-(\mu_1)$, while $\sigma_{\mu_2}(s)$ is as close as desired to $\sigma_-(\mu_2)$, and we have $\sigma_-(\mu_1) < \sigma_-(\mu_2)$. That is, we have $\sigma_{\mu_1}(s) < \sigma_{\mu_2}(s)$ for s sufficiently negative, and the comparison principle implies that this remains true for all $s \in \mathbb{R}$. This proves that $\sigma_\mu(s)$ is strictly increasing in $\mu \geq 0$ for any fixed s . The limit $\bar{\sigma}_\mu := \lim_{s \rightarrow \infty} \sigma(\mu, s)$ therefore also exists and is (weakly) increasing in μ , although it can (and will) be discontinuous.

Our construction produces a solution f_μ with the required boundary behaviour precisely when $\bar{\sigma}_\mu - \sigma_+(\mu) = 2\pi k$ for some $k \in \mathbb{Z}$. Since $\bar{\sigma}_\mu$ is increasing in μ and $\sigma_+(\mu)$ is strictly decreasing in μ , we have that $\bar{\sigma}_\mu - \sigma_+(\mu)$ is strictly increasing in μ , and hence each integer k can arise for at most one value of μ . We note from (6.4) that $\sigma_\mu(s)$ is strictly decreasing at any point where it takes values which are an odd multiple of π (corresponding to points where $g(s) = 0$), and hence the value of k can be computed as the number of points where the corresponding solution g of (6.3) equals zero.

The three solutions constructed above allow us to compute $\bar{\sigma}_\mu - \sigma_+(\mu)$ for these three specific values of μ : For $\mu = 0$, the solution $f_0 = \sin^2 \theta - \frac{1}{d-1} \cos^2 \theta$ gives rise to

$$g = (\cosh s)^{-\frac{d-3}{2}} \left(1 - \frac{d}{d-1} \frac{1}{\cosh^2 s} \right),$$

which has two crossings of zero, so that we have $\bar{\sigma}_0 - \sigma_+(0) = -4\pi$. For $\mu = d - 2$, the solution $f_{d-2} = \sin \theta \cos \theta$ gives $g = (\cosh s)^{-\frac{d+1}{2}} \sinh s$ which has a single crossing of zero and so, we have $\bar{\sigma}_{d-2} - \sigma_+(d-2) = -2\pi$. Finally, for $\mu = 2(d - 1)$, the solution $f_{2(d-1)} = \cos^2 \theta$ produces $g = (\cosh(s))^{-\frac{d+1}{2}}$, which has no zero crossings, and hence $\bar{\sigma}_{2(d-1)} - \sigma_+(2(d-1)) = 0$. Since the $\bar{\sigma}_\mu - \sigma_+(\mu)$ is strictly increasing, there can be no other values of μ between 0 and $2(d-1)$ for which $\bar{\sigma} - \sigma_+ \in 2\pi\mathbb{Z}$. For $\mu > 2(d-1)$ we have $\bar{\sigma}_\mu - \sigma_+(\mu) > 0$, and we observe that the line $\sigma = \pi$ cannot be crossed by solutions of (6.4) from below, so that we can never have $\bar{\sigma}_\mu - \sigma_+(\mu) = 2\pi k$ for k a positive

integer. This completes the proof that only the values $\mu = 0, d - 2, 2(d - 1)$ are possible. \square

Finally, we complete the proof of Lemma 6.2: The argument above shows that a Neumann eigenfunction on A with eigenvalue $2d$ has the form

$$f_0(\theta)\varphi_0(z) + f_{d-2}(\theta)\varphi_{d-2}(z) + f_{2(d-1)}(\theta)\varphi_{2(d-1)}(z)$$

where f_0, f_{d-2} and $f_{2(d-1)}$ are given in Lemma 6.3, and φ_0, φ_{d-2} and $\varphi_{2(d-1)}$ are Neumann eigenfunctions with the corresponding eigenvalues on $\tilde{A} \subset S^{d-2}$. In particular, φ_0 is a constant, φ_{d-2} is the restriction to \tilde{A} of a Neumann homogeneous degree 1 harmonic function \tilde{v} on $\tilde{\Gamma} \subset \mathbb{R}^{d-1}$, and $\varphi_{2(d-1)}$ is the restriction to \tilde{A} of a Neumann homogeneous degree 2 harmonic function \tilde{u} on $\tilde{\Gamma}$.

The homogeneous degree 2 Neumann harmonic function u is then given by extending this eigenfunction on A using the homogeneity:

$$\begin{aligned} u(x + sx_0) &= |x + sx_0|^2 \left(\cos^2 \theta \tilde{u} \left(\frac{x}{|x|} \right) + \sin \theta \cos \theta \tilde{v} \left(\frac{x}{|x|} \right) \right. \\ &\quad \left. + \varphi_0 \left(\sin^2 \theta - \frac{1}{d-1} \cos^2 \theta \right) \right) \\ &= |x + sx_0|^2 \left(\frac{|x|^2}{|x + sx_0|^2} \frac{1}{|x|^2} \tilde{u}(x) + \frac{s|x|}{|x + sx_0|^2} \frac{1}{|x|} \tilde{v}(x) \right. \\ &\quad \left. + \varphi_0 \left(\frac{s^2}{|x + sx_0|^2} - \frac{1}{d-1} \frac{|x|^2}{|x + sx_0|^2} \right) \right) \\ &= \tilde{u}(x) + s\tilde{v}(x) + \varphi_0 \left(s^2 - \frac{1}{d-1} |x|^2 \right) \end{aligned}$$

where we used $\sin^2 \theta = \frac{s^2}{s^2 + |x|^2}$ and $\cos^2 \theta = \frac{|x|^2}{s^2 + |x|^2}$, the expressions for f_0, f_{d-2} and $f_{2(d-1)}$ from Lemma 6.3, and the homogeneity of \tilde{v} and \tilde{u} . \square

Remark 6.4. The proof above applies with minor modifications to prove that for any positive integer k , the values of μ which can give rise to an eigenfunction on A with eigenvalue $\lambda = k^2 + (d - 2)k$ (corresponding to the restriction of a harmonic function on $\tilde{\Gamma} \times \mathbb{R}$ which is homogeneous of degree k) are precisely $\mu = j^2 + (d - 3)j$ for $j = 0, \dots, k$ (corresponding to eigenfunctions on \tilde{A} given by the restriction of a harmonic function on $\tilde{\Gamma}$ which is homogeneous of an integer degree no greater than k).

Lemma 6.5. *If $\tilde{\Gamma}$ is a tame cone in a $(d - 1)$ -dimensional subspace $E = (x_0)^\perp$ of \mathbb{R}^d , then $\tilde{\Gamma} \oplus \mathbb{R}x_0$ is a tame cone in \mathbb{R}^d .*

Proof. Suppose u is a homogeneous degree two Neumann harmonic function on $\tilde{\Gamma} \oplus \mathbb{R}x_0$, with bounded second derivatives. By Lemma 6.2 we can write

$$u(x + sx_0) = \tilde{u}(x) + s\tilde{v}(x) + C \left(s^2|x_0|^2 - \frac{1}{d-1}|x|^2 \right) \quad \text{for every } x \in \tilde{\Gamma},$$

where \tilde{u} is a homogeneous degree 2 Neumann harmonic function on $\tilde{\Gamma}$, \tilde{v} is a homogeneous degree 1 Neumann harmonic function on $\tilde{\Gamma}$, and C is constant. The last term has bounded second derivatives, so the sum of the other two terms must also. Fixing $s = 0$ we conclude that \tilde{u} has bounded second derivatives, and hence is quadratic function since $\tilde{\Gamma}$ is tame. Fixing $s = 1$ we conclude that \tilde{v} also has bounded second derivatives. But the second derivatives of a homogeneous degree one function are homogeneous of degree -1 , and hence are unbounded unless they are zero. Therefore \tilde{v} is a linear function, and we conclude that u is a quadratic function. \square

Now, we complete the proof of Theorem 6.1. We apply an induction on dimension. Suppose that u is a homogeneous degree 2 Neumann harmonic function on Γ with bounded second derivatives. We must show that u is a quadratic function.

First, for $d = 1$ then every Neumann harmonic function is constant, so every homogeneous degree 2 Neumann harmonic function vanishes and hence is a quadratic function.

Now suppose that every polyhedral cone in \mathbb{R}^p is tame for $1 \leq p < d$, and let Γ be a polyhedral cone in \mathbb{R}^d . We observe that by Lemma 5.13, for every $x_0 \in \partial\Gamma \setminus \{0\}$ the tangent cone Γ_{x_0} is a product of a cone $\tilde{\Gamma}$ in $(x_0)^\perp$ with $\mathbb{R}x_0$. By the induction hypothesis, $\tilde{\Gamma}$ is tame, and hence by Lemma 6.5 we conclude that Γ_{x_0} is tame. That is, $\bar{\Gamma} \setminus \{0\}$ is a tame domain. It follows from Theorem 5.3 that u is C^2 on $\bar{\Gamma} \setminus \{0\}$.

Since the second derivatives of u are bounded, there exists a sequence $(x_k)_{k \geq 1}$ of points x_k in Γ and a sequence $(e_k)_{k \geq 1}$ of $e_k \in \mathbb{S}^{d-1}$ such that

$$e_k^T D^2 u(x_k) e_k \rightarrow C_2 := \sup_{(x,e) \in \Gamma \times \mathbb{S}^{d-1}} e^T D^2 u(x) e \quad \text{as } k \rightarrow +\infty.$$

The second derivatives of a homogeneous degree 2 function are homogeneous of degree zero, so we can replace $(x_k)_{k \geq 1}$ by $(\tilde{x}_k)_{k \geq 1}$ given by $\tilde{x}_k = \frac{x_k}{|x_k|} \in \mathbb{S}^{d-1} \cap \Gamma$, and conclude that $e_k^T D^2 u(\tilde{x}_k) e_k \rightarrow C_2$ as $k \rightarrow +\infty$. By compactness, (\tilde{x}_k, e_k) converges for a subsequence of k to $(\bar{x}, e) \in (\mathbb{S}^{d-1} \cap \bar{\Gamma}) \times \mathbb{S}^{d-1}$. Since u is C^2 at \bar{x} , we have that $D^2 u|_{\bar{x}}(\bar{e}, \bar{e}) = C_2$.

Now we apply Lemma 4.5 with $B = \bar{\Gamma} \setminus \{0\}$, and deduce that $\Gamma = \Gamma^E \times \Gamma^\perp$, where Γ^E is a polyhedral cone in a subspace E of \mathbb{R}^d of positive dimension K , and Γ^\perp is a polyhedral cone in E^\perp , and we have

$$u(x) = \Lambda |\pi_E(x)|^2 + g(\pi_{E^\perp}(x)).$$

If $K = \dim E = d$ then since u is harmonic we have $\Lambda = 0$ and u vanishes. Otherwise we write

$$u(x) = K\Lambda \left(\frac{1}{K} |\pi_E(x)|^2 - \frac{1}{d-K} |\pi_{E^\perp}(x)|^2 \right) + \tilde{g}(\pi_{E^\perp}(x)).$$

The first term is harmonic, and u is harmonic, so the last term \tilde{g} is also harmonic. Furthermore, since u is homogeneous of degree 2, so is \tilde{g} , and \tilde{g} also satisfies zero Neumann boundary conditions on Γ^\perp since u and the first term do. Finally, \tilde{g} has bounded second derivatives since u does. Therefore by the induction hypothesis, \tilde{g} is a quadratic function, and so u is quadratic and Γ is tame. This completes the induction and the proof of Theorem 6.1. \square

7. Concave implies regular

The results of the previous two sections allow us to complete the proof of the main regularity result, Theorem 5.1. We begin with the following observation.

Lemma 7.1. *Let Ω be a bounded domain in \mathbb{R}^d with a continuous boundary $\partial\Omega$. For $\mu \in \mathbb{R}$, let $v \in H^1_{loc}(\Omega)$ be weak solution of $\Delta v + \mu = 0$ on Ω . If v is semi-concave on Ω , then v belongs to $C^{1,1}(\bar{\Omega})$.*

Proof. Note, that due to classical regularity theory of second order elliptic equations (cf [14, Corollary 8.11]), $v \in C^\infty(\Omega)$. By assumption, there is constant $C \in \mathbb{R}$ such that $D^2v|_x \leq CI$ for every $x \in \Omega$. Given any $x \in \Omega$ and any unit vector e , choose an orthonormal basis $\{e_1, \dots, e_d\}$ with $e = e_d$. Then

$$D^2v|_x(e, e) = \Delta v(x) - \sum_{i=1}^{d-1} D^2v|_x(e_i, e_i) \geq \mu - C(d-1)$$

for every $x \in \Omega$. Thus D^2v is also bounded from below. It follows that Dv is Lipschitz with bounded Lipschitz constant, and so extends continuously to $\bar{\Omega}$ as a Lipschitz function. \square

We are now ready to prove Theorem 5.1.

Proof of Theorem 5.1. We prove that v is C^2 on a neighbourhood of any point $x_0 \in \partial\Omega$. Choose $r > 0$ sufficiently small such that

$$(7.1) \quad B_r(x_0) \cap \Omega = x_0 + r(B_1(0) \cap \bar{\Gamma}_{x_0})$$

and set

$$w(x) = v(x_0 + rx) - Dv|_{x_0}(rx) + \frac{\mu}{2d}r^2|x|^2 \quad \text{for every } x \in B_1 \cap \bar{\Gamma}_{x_0}.$$

Then w is well-defined on $B_1 \cap \bar{\Gamma}_{x_0}$, with $\Delta w = 0$ on $B_1 \cap \Gamma_{x_0}$, and

$$D_\nu w|_x = rD_\nu v|_{x_0+rx} - rD_\nu v|_{x_0} + \frac{\mu}{d}r^2 x \cdot \nu = 0 \quad \text{for } x \in B_1 \cap \partial\Gamma_{x_0},$$

since both x_0 and $x_0 + rx$ are in Σ_i , so $D_{\nu_i} v|_{x_0+rx} = D_{\nu_i} v|_{x_0} = -\gamma_i$. We also use that x is normal to ν_i . This shows that w is a weak solution of (5.1). By hypothesis, there is a constant $C \in \mathbb{R}$ such that $D^2v \leq C$ on Ω , and so

$$D^2w|_x(e, e) = r^2D^2v|_{x_0+rx}(e, e) + \frac{\mu}{d}r^2|e|^2 \leq \left(r^2C + \frac{\mu}{d}r^2\right) |e|^2$$

for every $e \in \mathbb{R}^d$ and $x \in B_1 \cap \Gamma_{x_0}$, showing that w is semi-concave on $B_1 \cap \Gamma_{x_0}$. Thus, by Lemma 7.1, w is in $C^{1,1}(\overline{B_1 \cap \Gamma_{x_0}})$. By Theorem 6.1, $B = B_1(0) \cap \bar{\Gamma}_{x_0}$ is tame and hence by Theorem 5.3, $w \in C^2(B)$. Since x_0 is arbitrary, $w \in C^2(\bar{\Omega})$. \square

The results of Theorem 5.1 and Theorem 4.3 imply the following:

Corollary 7.2. *Let Ω be a convex polyhedral domain in \mathbb{R}^d with faces $\Sigma_1, \dots, \Sigma_m$, and for given $\mu, \gamma_1, \dots, \gamma_m \in \mathbb{R}$, let v be a weak solution of problem (4.5). If v is semi-concave, then v is a quadratic function.*

8. Quadratic solutions and circumsolids

In this section we determine precisely which are the domains on which the solution of (1.5) (or, more generally, (4.5)) is a quadratic function:

Proposition 8.1. *Let v be a quadratic function on \mathbb{R}^d , and let E_1, \dots, E_k be the eigenspaces of the Hessian of v with eigenvalues $\lambda_1, \dots, \lambda_k$. Then v satisfies an equation of the form (4.5) on a convex polyhedral domain Ω if and only if $\Omega = \{x \in \mathbb{R}^d \mid \pi_{E_i}(x) \in \Omega_i\}$, where Ω_i is a polyhedral domain in E_i for each i . Furthermore, v satisfies equation (1.5) if and only if $\lambda_i < 0$ and Ω_i is a circumsolid in E_i with center at the maximum of $v|_{E_i}$ and radius equal to $-1/\lambda_i$ for each i (see Definition 1.1).*

Proof. For a quadratic function, the Hessian $D^2v|_x$ is constant. Accordingly we denote the Hessian by A and let E_1, \dots, E_k be the eigenspaces of A , so that we have $v(x) = \frac{1}{2} \sum_{i=1}^k \lambda_i |\pi_i(x)|^2 + b \cdot x + c$, where π_i is the orthogonal projection onto E_i , where $\lambda_1, \dots, \lambda_k$ are the eigenvalues of A , and $b \in \mathbb{R}^d$ and $c \in \mathbb{R}$ are constants.

First we show that v satisfies (4.5) on a polyhedral domain Ω if and only if Ω is a product of polyhedral domains $\Omega_i \subset E_i$: If Ω has this form then

$$\begin{aligned} \Omega &= \bigcap_{i=1}^k \{x \mid \pi_i(x) \in \Omega_i\} \\ &= \bigcap_{i=1}^k \bigcap_{j=1}^{m_i} \left\{x \mid \pi_i(x) \cdot \nu_j^i \leq b_j^i\right\} \\ &= \bigcap_{i,j} \left\{x \mid x \cdot \nu_j^i \leq b_j^i\right\}, \end{aligned}$$

where $\Omega_i = \bigcap_{j=1}^{m_i} \{x \in E_i \mid x \cdot \nu_j^i \leq b_j^i\}$ for each i . Thus the normals to the faces of Ω are ν_j^i for $1 \leq i \leq k$ and $1 \leq j \leq m_i$, corresponding to the face $\Sigma_i^j = \overline{\Omega} \cap \{x \mid x \cdot \nu_j^i = b_j^i\}$. The derivative of v is given by

$$Dv|_x(e) = \sum_{p=1}^k \lambda_p \pi_p(x) \cdot e + b \cdot e,$$

so on the face Σ_i^j we have

$$D_{\nu_j^i} v|_x = \sum_{p=1}^k \lambda_p \pi_p(x) \cdot \nu_j^i + b \cdot e = \lambda_i x \cdot \nu_j^i + b \cdot e = \lambda_i b_j^i + b \cdot e,$$

which is constant on the face. Also we have $\Delta v = \sum_{i=1}^k \dim(E_i) \lambda_i$ which is constant, and so v is a solution of an equation of the form (4.5) on Ω .

The converse statement follows from the argument of Lemma 4.5: Equation (4.7) shows that each normal vector ν_i to a face of Ω is an eigenvector of A , and so lies in E_j for some j . This allows us to write

$$\begin{aligned} \Omega &= \bigcap_i \left\{x \in \mathbb{R}^d \mid x \cdot \nu_i < b_i\right\} \\ &= \bigcap_{j=1}^k \bigcap_{\nu_i \in E_j} \left\{x \in \mathbb{R}^d \mid x \cdot \nu_i < b_i\right\} \end{aligned}$$

$$\begin{aligned} &= \bigcap_{j=1}^k \bigcap_{\nu_i \in E_j} \left\{ x \in \mathbb{R}^d \mid \pi_j(x) \cdot \nu_i < b_i \right\} \\ &= \bigcap_{j=1}^k \left\{ x \in \mathbb{R}^d \mid \pi_j(x) \in \Omega_j \right\} \end{aligned}$$

where $\Omega_j = \bigcap_{i: \nu_i \in E_j} \{x \in E_j \mid x \cdot \nu_i < b_i\}$.

Now we specialise to the case of equation (1.5): First suppose v is strictly concave, so that $\lambda_i < 0$ for $i = 1, \dots, k$. Then we have

$$(8.1) \quad v = \frac{1}{2} \sum_{i=1}^k \lambda_i |\pi_i(x)|^2 + b \cdot x + c = \frac{1}{2} \sum_{i=1}^k \lambda_i \left| \pi_i(x) - \frac{1}{\lambda_i} \pi_i(b) \right|^2 + \tilde{c}$$

for some constant \tilde{c} . Hence $\frac{1}{\lambda_i} \pi_i(b)$ is the maximum point of v restricted to E_i . The condition that Ω_i is a circumsolid in E_i with centre at the maximum of $v|_{E_i}$ and radius $-1/(2\lambda_i)$ is that

$$\Omega_i = \bigcap_{j=1}^{m_i} \left\{ x \in E_i \mid \left(x - \frac{\pi_i(b)}{\lambda_i} \right) \cdot \nu_j^i < -\frac{1}{\lambda_i} \right\}.$$

In this case, we have for x in the face $\Sigma_j^i = \{x \mid (x - \frac{\pi_i(b)}{\lambda_i}) \cdot \nu_j^i = -\frac{1}{\lambda_i}\}$ that

$$\begin{aligned} D_{\nu_j^i} v(x) &= \sum_p \lambda_p \left(\pi_p(x) - \frac{\pi_p(b)}{\lambda_p} \right) \cdot \nu_j^i \\ &= \lambda_i \left(\pi_i(x) - \frac{\pi_i(b)}{\lambda_i} \right) \cdot \nu_j^i \\ &= \lambda_i \frac{-1}{\lambda_i} = -1 \end{aligned}$$

as required. Conversely, if we suppose that the boundary condition in (1.5) holds, then we can show that $\lambda_i < 0$ for every i as follows. We have

$$Dv|_x(e) = \sum_{p=1}^k \lambda_p \pi_p(x) \cdot e + b \cdot e.$$

Integrating over Ω_i and using the divergence theorem gives

$$-|\partial\Omega_i| = \int_{\partial\Omega_i} \nu_j^i \cdot Dv = \int_{\Omega_i} \Delta^{E_i} v = \dim(E_i) \lambda_i |\Omega_i|,$$

so that $\lambda_i < 0$ and v is strictly concave. Therefore v has the form (8.1), and the boundary condition gives

$$-1 = Dv|_x \cdot \nu_j^i = \lambda_i \left(\pi_i(x) - \frac{\pi_i(b)}{\lambda_i} \right) \cdot \nu_j^i$$

so that Ω_i is a circumsolid in E_i with radius $\frac{1}{\lambda_i}$ and centre at $\frac{\pi_i(b)}{\lambda_i}$. □

Corollary 8.2. *For a convex polyhedral domain Ω , there is a quadratic function v solving the elliptic boundary-value problem (1.5) if and only if Ω is a product of circumsolids.*

Proof. Proposition 8.1 shows that if Ω has a quadratic solution of (1.5) then Ω is a product of circumsolids. Conversely, suppose Ω is a product of circumsolids. Then there is a decomposition $\mathbb{R}^d = E_1 \oplus \dots \oplus E_k$ of \mathbb{R}^d into orthogonal subspaces E_1, \dots, E_k and

$$\Omega = \bigcap_{i=1}^k \left\{ x \in \mathbb{R}^d \mid \pi_i(x) \in \Omega_i \right\},$$

where

$$\Omega_i := \bigcap_{j=1}^{m_i} \left\{ x \in E_i \mid (x - p_i) \cdot \nu_j^i < R_i \right\}$$

for some $p_i \in E_i$ and $R_i > 0$. The above calculations show that

$$v(x) = -\frac{1}{2} \sum_{i=1}^k \frac{|\pi_i(x) - p_i|^2}{R_i} \quad \text{for every } x \in \Omega,$$

is a solution of (1.5) on Ω . □

Summarising, let v be a weak solution of (1.5) on a convex polyhedral domain Ω in \mathbb{R}^d . Then by Lemma 7.1, if v is semi-concave then $v \in C^{1,1}(\overline{\Omega})$. Note that for every boundary point $x_0 \in \partial\Omega$ and $r > 0$ small enough, v can be written as $v(x_0 + r \cdot) = w + q$ on $B_1(0) \cap \overline{\Gamma}_{x_0}$ for a quadratic function q and a weak solution w of (5.1). Then Theorem 5.3 and Theorem 6.1 state that $v \in C^{1,1}(\overline{\Omega})$ implies v is in $C^2(\overline{\Omega})$ and according to Theorem 4.3, the latter yields that v is quadratic. By Proposition 8.1, v then needs to be concave. Combining this together with Corollary 8.2, we can state the following characterisation.

Corollary 8.3. *Let v be a weak solution of (1.5) on a convex polyhedral domain Ω in \mathbb{R}^d . Then the following statements are equivalent.*

- (i) v is semi-concave;
- (ii) v is in $C^{1,1}(\overline{\Omega})$;
- (iii) v is in $C^2(\overline{\Omega})$;
- (iv) v is quadratic;
- (v) v is concave;
- (vi) Ω is a product of circumsolids.

9. Proof of the main results

In this section, we complete the proofs of our main results: Theorem 1.3, Theorem 1.4, and Corollary 1.5.

Proof of Theorem 1.3. Suppose Ω is polyhedral domain in \mathbb{R}^d that is not a product of circumsolids. We first show that for all $\alpha > 0$ small enough, the first Robin eigenfunction u_α is not log-concave. Set $v_\alpha = \log u_\alpha$. Then $v_0 \equiv 0$ and so, by Proposition 3.1, v_α can be expanded as

$$v_\alpha = \alpha v + f^\alpha,$$

where f^α belongs to $o(\alpha)$ in $C^{0,\beta}(\overline{\Omega})$ for all $\alpha > 0$ small enough, $\beta \in (0, 1)$, and v is a solution of the Neumann problem (1.5) for $\mu = \frac{d\lambda_\alpha}{d\alpha}|_{\alpha=0}$. Now, by Corollary 8.3, v is not concave on $\overline{\Omega}$. Thus, there exist $x, y \in \overline{\Omega}$ and $t \in (0, 1)$ such that

$$\varepsilon := tv(x) + (1-t)v(y) - v(tx + (1-t)y) > 0.$$

On the other hand, for every $\delta > 0$, there is an $\alpha_0 > 0$ such that $\|f^\alpha\|_\infty \leq \delta\alpha$ for all $0 < \alpha \leq \alpha_0$. Set $\delta = \varepsilon/4$, and let α be less than the corresponding α_0 . Then

$$\begin{aligned} & tv_\alpha(x) + (1-t)v_\alpha(y) - v_\alpha(tx + (1-t)y) \\ &= \alpha [tv(x) + (1-t)v(y) - v(tx + (1-t)y)] \\ &\quad + tf^\alpha(x) + (1-t)f^\alpha(y) - f^\alpha(tx + (1-t)y) \\ &\geq \alpha\varepsilon - 3\delta\alpha > 0, \end{aligned}$$

so v_α is not concave for any $\alpha < \alpha_0$, proving Theorem 1.3. □

Next we consider the convexity of superlevel sets $\{x \mid u_\alpha(x) > c\}$. We first establish two preliminary results. The first is a Lichnérowicz-Obata type result for the first non-trivial Neumann eigenvalue on a convex subset of the sphere \mathbb{S}^{d-1} , which extends partially the result of [12] by allowing non-smooth boundary, resulting in a larger class of equality cases.

Theorem 9.1. *For $d \geq 3$, let A be a convex open subset of the sphere \mathbb{S}^{d-1} . Then the first nontrivial eigenvalue*

$$\lambda_1(A) = \inf_{\varphi \in C^\infty(\bar{A}): \int_A \varphi d\mathcal{H}^{d-1} = 0} \frac{\int_A |D\varphi|^2 d\mathcal{H}^{d-1}}{\int_A |\varphi|^2 d\mathcal{H}^{d-1}}$$

of the Neumann Laplacian on A satisfies $\lambda_1 \geq d-1$. Moreover, $\lambda_1(A) = d-1$ if and only if the cone $\Gamma = \{x = rz \in \mathbb{R}^d \mid z \in A\}$ in \mathbb{R}^d has a linear factor, so that (after an orthogonal transformation) $\Gamma = \tilde{\Gamma} \times \mathbb{R}$ for some convex cone $\tilde{\Gamma}$ in \mathbb{R}^{d-1} . In this case, the corresponding eigenfunction is the restriction to \mathbb{S}^{d-1} of the linear function $L(x, y) = y$ for $(x, y) \in \mathbb{R}^{d-1} \times \mathbb{R}$.

Proof. First suppose that A has smooth boundary. Then for any $u \in H^1(A)$ and $f \in H^3(A)$ with $D_\nu f = 0$ on ∂A , the following Reilly-type formula holds:

$$\begin{aligned} & \int_A (\Delta f - (d-1)u)^2 d\mathcal{H}^{d-1} - \int_A |\nabla^2 f - ug|^2 d\mathcal{H}^{d-1} \\ (9.1) \quad & - (d-2) \int_A |\nabla f + \nabla u|^2 d\mathcal{H}^{d-1} - \int_{\partial A} h(\bar{\nabla} f, \bar{\nabla} f) d\mathcal{H}^{d-1} \\ & = (d-2) \left[(d-1) \int_A u^2 d\mathcal{H}^{d-1} - \int_A |\nabla u|^2 d\mathcal{H}^{d-1} \right], \end{aligned}$$

where ∇ is the covariant derivative on S^{d-1} , h is the second fundamental form of ∂A , and $\bar{\nabla} f$ is the gradient vector of the restriction of f to ∂A . This is proved by integration by parts and application of the curvature identity (the proof due to the first author for the situation without boundary is described in [10, Theorem B.18]).

In particular, given $u \in H^1(A)$ with $\int_A u d\mathcal{H}^{d-1} = 0$, let f be a solution of the problem

$$(9.2) \quad \begin{cases} \Delta f = (d-1)u & \text{on } A; \\ D_\nu f = 0 & \text{on } \partial A. \end{cases}$$

With this choice of f , the first term on the left hand-side in (9.1) vanishes,

and the remaining terms are non-positive, so the right-hand side is non-positive, proving the following Poincaré inequality

$$\int_A |\nabla u|^2 \, d\mathcal{H}^{d-1} \geq (d - 1) \int_A u^2 \, d\mathcal{H}^{d-1}$$

for all $u \in H^1(A)$ with $\int_A u \, d\mathcal{H}^{d-1} = 0$, implying that $\lambda_1(A) \geq d - 1$.

Now, consider the general case, where the boundary of A may not be smooth. Suppose that $\{A_n\}$ is a sequence of convex domains in S^n with smooth boundary, which converge in Hausdorff distance to A (these can be constructed by smoothing level sets of the distance to ∂A , for example). Let $\{u_n\}$ be the corresponding sequence of first eigenfunctions, normalised to $\int_{A_n} u_n^2 \, d\mathcal{H}^{d-1} = 1$. The solution of (9.2) is then given by $f_n = -\frac{d-1}{\lambda_1(A_n)} u_n$. As $n \rightarrow \infty$ we have $\lambda_1(A_n) \rightarrow \lambda_1(A)$, so $\lambda_1(A) \geq d - 1$.

Suppose that equality holds. Then, we can find a subsequence along which u_n converges weakly in H^1 to the first eigenfunction u on A , and the interior regularity estimates imply that u_n converges to u in $C^\infty(B)$ for any compact subset B of A . The right-hand side of (9.1) is equal to $(d - 1) - \lambda_1(A_n)$, which converges to zero as $n \rightarrow \infty$. The first term on the left hand-side in (9.1) is equal to zero for every n , and the last term on the left is non-positive by the convexity of A_n . Thus, on any compact subset B of A , we have

$$\begin{aligned} & \int_B |\nabla^2 f_n - (d - 1)u_n|^2 \, d\mathcal{H}^{d-1} + (d - 2) \int_B |\nabla f_n + \nabla u_n|^2 \, d\mathcal{H}^{d-1} \\ & \leq \int_{A_n} |\nabla^2 f_n - (d - 1)u_n|^2 \, d\mathcal{H}^{d-1} + (d - 2) \int_{A_n} |\nabla f_n + \nabla u_n|^2 \, d\mathcal{H}^{d-1} \\ & \leq (d - 2)(\lambda_1(A_n) - (d - 1)). \end{aligned}$$

Since f_n converges to $-u$ in $C^\infty(B)$, one has that

$$\begin{aligned} & \int_B |\nabla^2 f_n - (d - 1)u_n|^2 \, d\mathcal{H}^{d-1} + (d - 2) \int_B |\nabla f_n + \nabla u_n|^2 \, d\mathcal{H}^{d-1} \\ & \rightarrow \int_B |\nabla^2 u + ug|^2 \, d\mathcal{H}^{d-1} \end{aligned}$$

as $n \rightarrow +\infty$, while $(\lambda_1(A_n) - (d - 1)) \rightarrow 0$ as $n \rightarrow \infty$. Therefore we have $\nabla^2 u + ug = 0$ at every point of B , and hence at every point of A since B is an arbitrary compact subset of A .

It follows that u is the restriction of a linear function on \mathbb{R}^d to \mathbb{S}^{d-1} . To see this, define $e(z) := u(z)z + \nabla_i u(z)g^{ij}\partial_j z \in \mathbb{R}^d$. Then we have

$$\begin{aligned} \partial_k e &= \partial_k u z + u \partial_k z + \nabla_k \nabla_i u g^{ij} \partial_j z + \nabla_i u g^{ij} (-g_{ki} z) \\ &= (\nabla^2 u + u g)_{ki} g^{ij} \partial_j z = 0, \end{aligned}$$

so that e is constant on A . Finally, we have $u(z) = e(z) \cdot z$, which is a linear function. The claimed structure of A now follows from the Neumann condition $D_\nu u = 0$. \square

This result has an immediate consequence, which is important in our proof of Theorem 1.4.

Lemma 9.2. *Let Γ be a polyhedral convex cone in \mathbb{R}^d with vertex at the origin. Then there is a harmonic function \hat{w} on Γ which is homogenous of degree one and satisfies $D_\nu \hat{w} = -1$ on $\partial\Gamma$.*

Proof. Set $A := \Gamma \cap \mathbb{S}^{d-1}$. First, consider the case when Γ does not admit a linear factor. Then by Theorem 9.1, $d-1$ is in the resolvent set $\rho(-\Delta_{|A}^{\mathbb{S}^{d-1}})$ of the operator $-\Delta_{|A}^{\mathbb{S}^{d-1}}$ equipped with homogeneous Neumann boundary conditions and realised in $L^2(A)$. Therefore, there exists a unique weak solution $\tilde{\varphi}$ of

$$(9.3) \quad \begin{cases} \Delta^{\mathbb{S}^{d-1}} \tilde{\varphi} + (d-1)\tilde{\varphi} &= 0 \text{ on } A, \\ D_\nu \tilde{\varphi} &= -1 \text{ on } \partial A. \end{cases}$$

It follows that the function

$$\hat{w}(rz) := r \tilde{\varphi}(z) \quad \text{for every } r \in [0, 1] \text{ and } z \in A$$

is harmonic on Γ , homogeneous of degree one, and satisfies $D_\nu \hat{w} = -1$ on $\partial\Gamma$.

Now, suppose Γ has a linear factor, so that there is a $k \in \{1, \dots, d-1\}$ such that $A = (\mathbb{R}^k \oplus \tilde{\Gamma}) \cap \mathbb{S}^{d-1}$ for a polyhedral cone $\tilde{\Gamma}$ in \mathbb{R}^{d-k} with no linear factors. In particular, if $k = d-1$ then $\tilde{\Gamma} = (0, +\infty)$, and then $\tilde{\varphi}(z) = z$ is a solution of (9.3) on $\tilde{\Gamma}$ and so,

$$\hat{w}(x_1, \dots, x_{d-1}, z) := z \quad \text{for every } (x_1, \dots, x_{d-1}, z) \in \Gamma$$

is harmonic on Γ , homogeneous of degree one, and satisfies $D_\nu \hat{w} = -1$ on $\partial\Gamma$. Otherwise we have $1 \leq k \leq d-2$ and $\tilde{\Gamma}$ is a convex polyhedral cone with

has no linear factor. Then by the first case, there is a harmonic function \tilde{w} on $\tilde{\Gamma}$ which is homogeneous of degree one and satisfies $D_\nu \tilde{w} = -1$ on $\partial \tilde{\Gamma}$. Then the function

$$\hat{w}(x_1, \dots, x_k, z) := \tilde{w}(z) \quad \text{for every } (x_1, \dots, x_k, z) \in \mathbb{R}^{d-k} \times \tilde{\Gamma} = \Gamma$$

is a harmonic on Γ , homogeneous of degree one, and satisfies $D_\nu \hat{w} = -1$ on $\partial \Gamma$. We note that the solution space is in general of dimension k , since we can add an arbitrary linear function on the linear factors. \square

Further, in dimension $d \geq 3$, we will use the following characterisation of polyhedra with boundary points with inconsistent normals. We omit the proof of this result.

Proposition 9.3. *Let Ω be a convex polyhedral domain in \mathbb{R}^d , $d \geq 2$, with outer unit face normals ν_1, \dots, ν_m . For each point $x \in \partial \Omega$, let $\mathcal{I}(x)$ be the index set (1.4) of faces touching x . Then the following statements are equivalent.*

1. *x has inconsistent normals: That is, there is no $\gamma \in \mathbb{R}^d$ satisfying*

$$\nu_i \cdot \gamma = -1 \quad \text{for every } i \in \mathcal{I}(x).$$

2. *If \hat{w} is a function on Γ_x which is harmonic and homogeneous of degree one and satisfies*

$$D_{\nu_i} \hat{w} = -1 \quad \text{on } \Sigma_i \text{ for all } i \in \mathcal{I}(x),$$

then \hat{w} is not a linear function.

3. *The tangent cone Γ_x to Ω is not a circumsolid.*

We now proceed to the proof of Theorem 1.4:

Proof of Theorem 1.4. Under the assumptions of Theorem 1.4, we first prove that the function v satisfying the Neumann problem (1.5) has some non-convex superlevel sets.

Let $x_0 \in \partial \Omega$ and $\Gamma = \Gamma_{x_0}$, and choose $r > 0$ small enough so that (7.1) holds. We define

$$\tilde{v}(x) := v(x_0 + x) + \frac{\mu}{2d}|x|^2 \quad \text{for } x \in B_r(0) \cap \Gamma.$$

Then \tilde{v} is harmonic on $B_r \cap \Gamma$ and satisfies $D_\nu \tilde{v} = -1$ on $B_r \cap \partial \Gamma_{x_0}$. By Lemma 9.2, there is a harmonic function \hat{w} on Γ which is homogenous of

degree one and satisfies $D_\nu \hat{w} = -1$ on $\partial\Gamma$. Then the function

$$w(x) := \tilde{v}(x) - \hat{w}(x) \quad \text{for every } x \in B_r(0) \cap \Gamma$$

is a weak solution of the Neumann problem (4.1) on $B_r(0) \cap \Gamma$. Proposition 5.5 applied to a suitable dilation of w gives the series expansion (5.8). Therefore, v can be written as

$$(9.4) \quad v(x_0 + x) = -\frac{\mu}{2d}|x|^2 + \hat{w}(x) + \sum_{i=0}^{\infty} f_i \psi_i(x) \quad \text{for every } x \in B_r \cap \Gamma,$$

where ψ_i is the harmonic function on Γ given by

$$\psi_i(x) := s^{\beta_i} \varphi_i(z) \quad \text{for every } x = sz \text{ with } s > 0 \text{ and } z \in A.$$

Here $A = \Gamma \cap \mathbb{S}^{d-1}$, $\{\varphi_i\}_{i=0}^{\infty}$ is an orthonormal basis of $L^2(A)$ consisting of eigenfunctions φ_i of the Neumann-Laplacian $\Delta^{\mathbb{S}^{d-1}}$ on A , and β_i are given by (5.9). Further, $\beta_0 = 0$ (corresponding to $\lambda_0 = 0$), and the remaining β_i are estimated by Theorem 9.1 and (5.9), so that $\beta_i \geq 1$ for $i \geq 1$. Moreover, there is no loss of generality in assuming that each $\beta_i > 1$, since $\tilde{w}(x) := \sum_{i:\beta_i=1} f_i \psi_i(x)$ is harmonic on Γ , of homogeneous degree one, and satisfies $D_\nu \tilde{w} = 0$ on $\partial\Gamma$, and so \tilde{w} can be included in \hat{w} . Summarising, we can write

$$(9.5) \quad v(x_0 + x) = v(x_0) - \frac{\mu}{2d}|x|^2 + \hat{w}(x) + \sum_{i=1}^{\infty} f_i \psi_i(x)$$

for every $x \in B_r \cap \Gamma$, where the non-vanishing terms in the sum all have exponent $\beta_i > 1$.

Before continuing the proof of Theorem 1.4, we observe that the proof of Theorem 5.3 applies almost without change to prove the following generalisation:

Theorem 9.4. *Let Ω be a polyhedral domain in \mathbb{R}^d , and B a relatively open subset of Ω . Let $w \in H^1(B)$ be a weak solution of problem (5.1). For each $x_0 \in B \cap \partial\Omega$, choose $r(x_0) > 0$ small enough so that (7.1) holds, so that by Proposition 5.5 w is given by the expansion*

$$w(x_0 + x) = \sum_{i=0}^{\infty} f_i(x_0) \psi_i^{x_0}(x) \quad \text{for every } x \in B_{r(x_0)}(0) \cap \Gamma_{x_0},$$

where $\psi_i^{x_0}$ is the harmonic function on Γ_{x_0} given by $\psi_i^{x_0}(x) = |x|^{\beta_i(x_0)} \varphi_i^{x_0}\left(\frac{x}{|x|}\right)$, $A_{x_0} = \Gamma_{x_0} \cap \mathbb{S}^{d-1}$, $\{\varphi_i^{x_0}\}_{i=0}^\infty$ is an orthonormal basis of $L^2(A_{x_0})$ consisting of eigenfunctions $\varphi_i^{x_0}$ of the Neumann-Laplacian $\Delta^{\mathbb{S}^{d-1}}$ on A_{x_0} , and $\beta_i(x_0)$ are given by (5.9). If $f_i(x_0) \neq 0$ only for those i with $\beta_i(x_0) \geq 2$ for every $x_0 \in B \cap \partial\Omega$, then $w \in C^2(B)$.

Continuation of the Proof of Theorem 1.4.

The case of inconsistent normals: In the case where Ω has a boundary point x_0 where the normal vectors are inconsistent, we have by Proposition 9.3 that \hat{w} is not a linear function. It follows that \hat{w} does not have convex superlevel sets: Choosing any point $z \in A$ where $\hat{w}(z) \neq 0$ and $D^2\hat{w}|_z \neq 0$, we have that z is a null eigenvector of $D^2\hat{w}$ (since \hat{w} is homogeneous of degree one), and that the trace of $D^2\hat{w}|_z$ on the orthogonal subspace $(\mathbb{R}z)^\perp$ is zero (since \hat{w} is harmonic). It follows that $D^2\hat{w}|_z$ has an eigenvector $\xi \in (\mathbb{R}z)^\perp$ with positive eigenvalue, so that $D^2\hat{w}|_z(\xi, \xi) > 0$.

Now let $\eta = \xi - \frac{D\hat{w}|_z(\xi)}{\hat{w}(z)}z$. Then we have

$$D\hat{w}|_z(\eta) = D\hat{w}|_z(\xi) - \frac{D\hat{w}|_z(\xi)}{\hat{w}(z)}D\hat{w}|_z(z) = 0,$$

since $D\hat{w}|_z(z) = \hat{w}(z)$ by the homogeneity of \hat{w} . Also, we have

$$D^2\hat{w}|_z(\eta, \eta) = D^2\hat{w}|_z(\xi, \xi) > 0,$$

since z is a null eigenvector of $D^2\hat{w}|_z$. It follows that the superlevel set $S = \{x \mid \hat{w}(x) > \hat{w}(z)\}$ is not convex near z , since for small $s \neq 0$ we have $\hat{w}(z \pm s\eta) > \hat{w}(z)$ and hence $z \pm s\eta \in S$, but $z \notin S$. Since \hat{w} is homogeneous, the superlevel sets $S_\lambda = \{x \mid \hat{w}(x) > \lambda\hat{w}(z)\}$ are also non-convex near λz , for any $\lambda > 0$.

Now we conclude that v also has some non-convex superlevel sets: By the non-convexity and openness of S , there exist points x_1 and x_2 in Γ such that $x_1, x_2 \in S$ but $\frac{x_1+x_2}{2} \notin \bar{S}$. It follows that there exists $\varepsilon > 0$ such that $\hat{w}(x_i) > \hat{w}(z) + \varepsilon$ for $i = 1, 2$, but $\hat{w}\left(\frac{x_1+x_2}{2}\right) < \hat{w}(z) - \varepsilon$. Now we use the expression (9.4) to write

$$\begin{aligned} v(x_0 + \lambda x_j) &= v(x_0) + \lambda\hat{w}(x_j) - \frac{\mu}{2d}\lambda^2|x_j|^2 + \sum_{i>1} f_i\lambda^{\beta_i}\psi_i(x_j) \\ &= v(x_0) + \lambda\left(\hat{w}(x_j) - \frac{\mu\lambda}{2d}|x_j|^2 + \sum_{i>1} \lambda^{\beta_i-1}\psi_i(x_j)\right) \end{aligned}$$

$$\begin{aligned} &> v(x_0) + \lambda(\hat{w}(x_j) - \varepsilon) \\ &> v(x_0) + \lambda\hat{w}(z) \end{aligned}$$

for $j = 1, 2$, for $\lambda > 0$ sufficiently small. Here we used the fact that the sum $\sum_{i>1} \lambda^{\beta_i-1} \psi_i(x_j)$ converges to zero as λ approaches zero, which follows as in the proof of Lemma 5.6. Similarly, we have

$$v\left(x_0 + \lambda \frac{x_1 + x_2}{2}\right) < v(x_0) + \lambda\hat{w}(z)$$

for $\lambda > 0$ sufficiently small. This proves that the superlevel set $\{x \mid v(x) > v(x_0) + \lambda\hat{w}(z)\}$ is not convex.

The case of consistent normals: Now we consider the case where the normals are consistent at every point. By Proposition 9.3 this implies that for every x_0 , the function \hat{w} on Γ_{x_0} provided by Lemma 9.2 is linear. If for every x_0 the non-zero terms in the expansion of the Neumann harmonic function $w(x) = \sum_{i=1}^{\infty} f_i \psi_i(x)$ had exponent $\beta_i \geq 2$ for every x_0 , then by Proposition 9.4 w is C^2 near x_0 and hence (9.5) implies that v is also C^2 near x_0 .

However, if we assume that Ω is not a product of circumsolids, then by Corollary 8.3, we have that v is not C^2 and so there must be some $x_0 \in \partial\Omega$ such that the first nontrivial term in the sum in (9.5) has exponent β_i between 1 and 2: Precisely, we can assume (by choosing a new basis for the corresponding eigenspace if necessary) that

$$v(x_0 + x) = v(x_0) - \frac{\mu}{2d}|x|^2 + f_1\psi_1(x) + \sum_{i>1}^{\infty} f_i \psi_i(x) + \hat{w}(x)$$

for every $x \in B_r \cap \Gamma$, where $f_1 > 0$, $1 < \beta_1 < 2$, and $\beta_i > \beta_1$ for $i > 1$. Since ψ_i is homogeneous of order β_i and $\hat{w}(x) = x \cdot \gamma$, we have

$$(9.6) \quad Dv|_{x_0+\lambda x}(\xi) = \gamma \cdot \xi - \frac{\mu\lambda}{d} x \cdot \xi + \sum_{i \geq 1}^{\infty} f_i \lambda^{\beta_i-1} D\psi_i|_x(\xi)$$

$$(9.7) \quad \begin{aligned} D^2v|_{x_0+\lambda x}(\xi, \eta) &= -\frac{\mu}{d} \xi \cdot \eta + f_1 \lambda^{\beta_1-2} D^2\psi_1|_x(\xi, \eta) + \\ &+ \sum_{i>1}^{\infty} f_i \lambda^{\beta_i-2} D^2\psi_i|_x(\xi, \eta) \end{aligned}$$

for every $x \in B_r(0) \cap \Gamma$, $\lambda \in (0, 1)$, and $\xi, \eta \in \mathbb{R}^d$.

To see that v has a non-convex superlevel set, it suffices to show that there exists x with $x_0 + x \in \Omega$, and $\xi \in \mathbb{R}^d$, such that

$$Dv|_{x_0+x}(\xi) = 0 \quad \text{and} \quad D^2v|_{x_0+x}(\xi, \xi) > 0.$$

We note that as λ approaches zero, the right-hand side of (9.6) is dominated by the first term since the remaining terms are homogeneous of positive degree in λ , while the right-hand side of (9.7) is dominated by the first non-trivial term in the sum since this is homogeneous of degree $\beta_1 - 2 < 0$ in λ .

This motivates the following lemma:

Lemma 9.5. *Suppose that the restriction of ψ_1 to the hyperplanar section $L := \{x \in \Gamma \mid \gamma \cdot x = |\gamma|\}$ of $\Gamma = \Gamma_{x_0}$ is not concave. Then v has a non-convex superlevel set.*

Proof. Since the restriction of ψ_1 to L is not concave, there exists $x \in L$ and $\xi_0 \perp \gamma$ such that $D^2\psi_1|_x(\xi_0, \xi_0) > 0$. The expression (9.6) then implies

$$Dv|_{x_0+\lambda x}(\xi_0) = O(\lambda^{\beta_1-1}), \quad \text{and} \quad Dv|_{x_0+\lambda x}(x) = |\gamma| + O(\lambda^{\beta_1-1}),$$

for $\lambda \rightarrow 0+$, from which it follows that $Dv|_{x_0+\lambda x}(\xi_0 + c(\lambda)x) = 0$ for some $c(\lambda) = O(\lambda^{\beta_1-1})$ as $\lambda \rightarrow 0+$. Then we have by (9.7) that

$$D^2v|_{x_0+\lambda x}(\xi_0 + c(\lambda)x, \xi_0 + c(\lambda)x) = \lambda^{\beta_1-2} (f_1 D^2\psi_1|_x(\xi_0, \xi_0) + O(\lambda^\sigma)),$$

where $\sigma = \min\{\beta_1 - 1, 2 - \beta_2, \beta_2 - \beta_1\}$. Thus and since $D^2\psi_1|_x(\xi_0, \xi_0) > 0$, we have that $D^2v|_{x_0+\lambda x}(\xi_0 + c(\lambda)x, \xi_0 + c(\lambda)x) > 0$ for $\lambda > 0$ sufficiently small, proving that v has a non-convex superlevel set. \square

Remark 9.6. We are unable to establish the hypothesis of Lemma 9.5 for dimensions $d \geq 3$, but note here that this would be sufficient to prove that v has a non-convex superlevel set whenever Ω is not a product of circumsolids, substantially strengthening the result of Theorem 1.4.

The case $d = 2$: We can establish the hypothesis of Lemma 9.5 in the case $d = 2$, as follows: In this case the tangent cone Γ_{x_0} at any boundary point $x_0 \in \partial\Omega$ is a sector with opening angle $\theta_0 \leq \pi$. The case $\theta_0 = \pi$ cannot arise, since in that case the homogeneous Neumann harmonic functions on the half-plane Γ_{x_0} are spherical harmonics with integer degree of homogeneity, so one cannot have $\beta_1 \in (1, 2)$. Therefore $\theta_0 \in (0, \pi)$.

Let γ be the inward-pointing bisector of this sector of length $1/\sin(\frac{\theta_0}{2})$. Then we have $\nu_i \cdot \gamma = -1$ for $i = 1, 2$, where ν_1 and ν_2 are the outer unit

normal vectors to the two faces of Ω which meet at x_0 . The homogenous degree one harmonic function of Lemma 9.2 is then given by $\hat{w}(x) = \gamma \cdot x$. In particular \hat{w} is linear, so we are in the situation where all boundary points have consistent normals. The corresponding eigenfunctions are given by

$$\psi_i(r(\cos \theta) e_1 + r(\sin \theta) e_2) = \begin{cases} r^{\frac{i\pi}{\theta_0}} \cos\left(\frac{i\pi}{\theta_0} \theta\right), & i \text{ even;} \\ r^{\frac{i\pi}{\theta_0}} \sin\left(\frac{i\pi}{\theta_0} \theta\right), & i \text{ odd,} \end{cases}$$

for every $\theta \in \left(-\frac{\theta_0}{2}, \frac{\theta_0}{2}\right)$. For every non-negative integer i , ψ_i has degree of homogeneity $\beta_i = \frac{i\pi}{\theta_0}$. Here, $e_1 = \frac{\gamma}{|\gamma|}$, and e_2 is a unit vector orthogonal to γ .




Figure 9: The case $d = 2$.

The only possibilities which can give rise to $1 < \beta_i < 2$ are where $\theta_0 \in (\pi/2, \pi)$ and $i = 1$. In this case ψ_1 is odd in θ , and hence is an odd function when restricted to the line L (see Figure 9). Since an odd concave function is necessarily a multiple of the identity function, the only possibility in which ψ_1 has a concave restriction to L is when

$$\psi_1(e_1 + ye_2) = cy,$$

which implies by homogeneity that

$$\psi_1(xe_1 + ye_2) = cx^{\beta_1-1}y.$$

However a direct computation shows that this is harmonic only in the cases $\beta_1 = 1$ or $\beta_1 = 2$, which are impossible. This proves Lemma 9.5 for the case $d = 2$, so we have established that v has a non-convex superlevel set whenever Ω is not a product of circumsolids. We note that for $d = 2$ this applies except when Ω is either a circumsolid or a rectangle.

Now we complete the proof of Theorem 1.4, by proving that the Robin eigenfunction u_α also has some non-convex superlevel sets for sufficiently small $\alpha > 0$:

By Proposition 3.1, for all sufficiently small $\alpha > 0$, the first Robin eigenfunction u_α is given by

$$(9.8) \quad u_\alpha = \mathbf{1} + \alpha v + f^\alpha$$

where f^α is $o(\alpha)$ in $C^{0,\beta}(\overline{\Omega})$ for some $\beta \in (0, 1)$.

We have proved that v has some non-convex superlevel sets, which means that there exist points x_1 and x_2 in Ω , a number $c \in \mathbb{R}$, and $\varepsilon > 0$ such that $v(x_i) > c + \varepsilon$ for $i = 1, 2$, but $v\left(\frac{x_1+x_2}{2}\right) < c - \varepsilon$. But then we have by (9.8) for α sufficiently small that

$$u_\alpha(x_i) = 1 + \alpha v(x_i) + o(\alpha) > 1 + \alpha c + \alpha \varepsilon + o(\alpha) > 1 + \alpha c$$

for $i = 1, 2$, while

$$u_\alpha\left(\frac{x_1 + x_2}{2}\right) = 1 + \alpha v\left(\frac{x_1 + x_2}{2}\right) + o(\alpha) < 1 + \alpha c - \alpha \varepsilon + o(\alpha) < 1 + \alpha c.$$

It follows that the superlevel set $\{x \mid u_\alpha(x) > 1 + \alpha c\}$ is not convex for sufficiently small $\alpha > 0$. □

It remains to give the proof of Corollary 1.5.

Proof of Corollary 1.5. It suffices to show the following: If Ω is a convex domain for which the Robin ground state $u_\alpha(\Omega)$ is not log-convex (or has a non-convex superlevel set) for some α , and $\{\Omega_n\}$ is a sequence of convex domains which approach Ω in Hausdorff distance, then the Robin eigenfunction $u_{\alpha,n}$ of Ω_n is not log-concave (respectively, has a non-convex superlevel set) for sufficiently large n .

We apply Proposition 3.2, which applies since the volume and perimeter of convex sets are continuous with respect to Hausdorff distance. In particular, by (3.2) the eigenfunctions $u_{n,\alpha}$ converge uniformly to u_α on any subset which is contained in $\overline{\Omega}_n$ for all large n .

Under the assumption that u_α is not log-concave on Ω , there exist points x_1 and x_2 in Ω such that $\frac{1}{2}(\log u_\alpha(x_1) + \log u_\alpha(x_2)) > \log u_\alpha\left(\frac{x_1+x_2}{2}\right)$, or equivalently $u_\alpha(x_1)u_\alpha(x_2) > u_\alpha\left(\frac{x_1+x_2}{2}\right)^2$. For sufficiently large n the points x_1, x_2 and $\frac{x_1+x_2}{2}$ are all contained in Ω_n , and hence we have

$$u_{\alpha,n}(x_1)u_{\alpha,n}(x_2) - u_{\alpha,n}\left(\frac{x_1+x_2}{2}\right)^2 \rightarrow u_\alpha(x_1)u_\alpha(x_2) - u_\alpha\left(\frac{x_1+x_2}{2}\right)^2 > 0$$

as $n \rightarrow \infty$, and hence the left-hand side is positive for sufficiently large n , proving that $u_{\alpha,n}$ is not log-concave for n large.

Similarly, under the assumption that u_α has a non-convex superlevel set, there exist points x_1, x_2 in Ω and $c \in \mathbb{R}$ such that $u_\alpha(x_i) > c$ for $i = 1, 2$, while $u_\alpha\left(\frac{x_1+x_2}{2}\right) < c$. As before the convergence of $u_{\alpha,n}$ to u_α at the points x_1, x_2 and $\frac{x_1+x_2}{2}$ guarantees that $u_{\alpha,n}(x_i) > c$ for $i = 1, 2$ and $u_{\alpha,n}\left(\frac{x_1+x_2}{2}\right) < c$ for n sufficiently large, proving that $u_{\alpha,n}$ has a non-convex superlevel set. \square

10. Discussion and conjectures

We conclude this paper by formulating some observations and conjectures.

We recall that the Dirichlet eigenvalue problem corresponds to the limiting case $\alpha \rightarrow +\infty$ in which it is well-known (cf [6]) that the first eigenfunction is log-concave. Thus, our first conjecture is naturally:

1. Conjecture. *For a given bounded convex domain Ω , there is an $\alpha_0 > 0$ such that for all $\alpha \geq \alpha_0$, the first Robin eigenfunction u_α is log-concave.*

Furthermore, it would be interesting to know whether the threshold α_0 depends on the dimension $d \geq 2$ and whether it can be independent of the domain Ω .

Let Ω be a convex polyhedral domain that is not the product of circum-solids. In order to prove in dimensions $d \geq 3$ that the first Robin eigenfunction u_α has non-convex superlevel sets without imposing the stronger hypothesis Ω has inconsistent normals at some boundary point, our proof of Theorem 1.4 shows that one needs to study the second case when the harmonic function \hat{w} given by Lemma 9.2 is linear. The linear case in dimension $d = 2$ is much simpler to treat than the $(d - 1)$ -dimensional hyperplane $\mathcal{H} := \{x \in \mathbb{R}^d \mid x \cdot \gamma = |\gamma|\}$ reduces to a line segment L . Nevertheless, we are convinced that the following conjecture holds.

2. Conjecture. *If Ω is a convex polyhedron in \mathbb{R}^d for $d \geq 3$ which is not a product of circumsolids, then for sufficiently small $\alpha > 0$, the first Robin eigenfunction u_α has non-convex superlevel sets.*

Our argument shows that it would be sufficient to establish Lemma 9.5 whenever Γ is a polyhedral convex cone which is a circumsolid about the point γ , and ψ_1 is a homogeneous harmonic function with Neumann boundary conditions on Γ with degree of homogeneity between 1 and 2 (see Remark 9.6).

Our initial motivation for the work undertaken in this paper was to establish the **fundamental gap conjecture for Robin eigenvalues**:

Let Ω be a bounded convex domain in \mathbb{R}^d of diameter D , V be a weakly convex potential, and for $\alpha > 0$, let $\lambda_i(\alpha)$ be the Robin eigenvalues on the interval $(-\frac{D}{2}, \frac{D}{2})$. Then for $\alpha > 0$, the Robin eigenvalues $\lambda_i^V(\alpha)$ of the Schrödinger operator $-\Delta + V$ satisfy

$$\lambda_1^V(\alpha) - \lambda_0^V(\alpha) \geq \lambda_1(\alpha) - \lambda_0(\alpha).$$

In the Dirichlet case this conjecture was first observed by van den Berg [23] and then later independently suggested by Ashbaugh and Benguria [5], and Yau [24]. The complete proof of the fundamental gap conjecture in this case was given in [1]. Theorem 2.1 is a first attempt to prove the fundamental gap conjecture for Robin eigenvalues, but provides non-optimal lower bounds. But due to our main Theorem 1.3, it is clear that this conjecture can only be proved by methods avoiding the log-concavity of the first Robin eigenfunction. To the best of our knowledge, only Lavine's work [17] provides a proof of the fundamental gap conjecture which does not use the log-concavity of the first eigenfunction. That paper concerns the Dirichlet and Neumann case on a bounded interval. With this in mind, we conclude with the following question:

Open problem. *How can one prove the fundamental gap conjecture for Robin eigenvalues without using the log-concavity of the first eigenfunction?*

In the recent work [3], we adapted Lavine's method on a bounded interval to show that the fundamental gap conjecture for Robin eigenvalues of the linear Schrödinger operator holds. In dimension $d \geq 2$ the above problem still remains open.

References

- [1] Ben Andrews and Julie Clutterbuck. Proof of the fundamental gap conjecture. *J. Amer. Math. Soc.*, 24(3):899–916, 2011.
- [2] Ben Andrews, Julie Clutterbuck, and Daniel Hauer. Non-concavity of Robin eigenfunctions. arXiv:1711.02779, November 2017.
- [3] Ben Andrews, Julie Clutterbuck, and Daniel Hauer. The fundamental gap for a one-dimensional Schrödinger operator with Robin boundary conditions. arXiv:2002.06900, 2020.
- [4] Tom M. Apostol and Mamikon A. Mnatsakanian. Solids circumscribing spheres. *Amer. Math. Monthly*, 113(6):521–540, 2006.
- [5] Mark S. Ashbaugh and Rafael Benguria. Optimal lower bound for the gap between the first two eigenvalues of one-dimensional Schrödinger operators with symmetric single-well potentials. *Proc. Amer. Math. Soc.*, 105(2):419–424, 1989.
- [6] Herm Jan Brascamp and Elliott H. Lieb. On extensions of the Brunn-Minkowski and Prékopa-Leindler theorems, including inequalities for log concave functions, and with an application to the diffusion equation. *J. Functional Analysis*, 22(4):366–389, 1976.
- [7] Dorin Bucur, Alessandro Giacomini, and Paola Trebeschi. The Robin-Laplacian problem on varying domains. *Calc. Var. Partial Differential Equations*, 55(6):Paper No. 133, 29, 2016.
- [8] Isaac Chavel. *Eigenvalues in Riemannian geometry*, volume 115 of *Pure and Applied Mathematics*. Academic Press, Inc., Orlando, FL, 1984.
- [9] Shiu Yuen Cheng and Peter Li. Heat kernel estimates and lower bound of eigenvalues. *Comment. Math. Helv.*, 56(3):327–338, 1981.
- [10] Bennett Chow, Peng Lu, and Lei Ni. *Hamilton’s Ricci flow*, volume 77 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI; Science Press Beijing, New York, 2006.
- [11] Daniel Daners, Daniel Hauer, and E. N. Dancer. Uniform convergence of solutions to elliptic equations on domains with shrinking holes. *Adv. Differential Equations*, 20(5-6):463–494, 2015.
- [12] José F. Escobar. Uniqueness theorems on conformal deformation of metrics, Sobolev inequalities, and an eigenvalue estimate. *Comm. Pure Appl. Math.*, 43(7):857–883, 1990.

- [13] Alexey Filinovskiy. On the eigenvalues of a Robin problem with a large parameter. *Math. Bohem.*, 139(2):341–352, 2014.
- [14] David Gilbarg and Neil S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [15] Tosio Kato. *Perturbation theory for linear operators*. Classics in Mathematics. Springer-Verlag, Berlin, 1995. Reprint of the 1980 edition.
- [16] J. B. Kennedy. *On the Isoperimetric Problem for the Laplacian with Robin and Wentzell Boundary Conditions*. PhD thesis, School of Mathematics and Statistics, University of Sydney, NSW 2006, Australia, 2010.
- [17] Richard Lavine. The eigenvalue gap for one-dimensional convex potentials. *Proc. Amer. Math. Soc.*, 121(3):815–821, 1994.
- [18] Robin Nittka. Regularity of solutions of linear second order elliptic and parabolic boundary value problems on Lipschitz domains. *J. Differential Equations*, 251(4-5):860–880, 2011.
- [19] L. E. Payne and H. F. Weinberger. An optimal Poincaré inequality for convex domains. *Arch. Rational Mech. Anal.*, 5:286–292, 1960.
- [20] R. Schoen and S.-T. Yau. *Lectures on differential geometry*. Conference Proceedings and Lecture Notes in Geometry and Topology, I. International Press, Cambridge, MA, 1994. Lecture notes prepared by Wei Yue Ding, Kung Ching Chang [Gong Qing Zhang], Jia Qing Zhong and Yi Chao Xu, Translated from the Chinese by Ding and S. Y. Cheng, With a preface translated from the Chinese by Kaising Tso.
- [21] I. M. Singer, Bun Wong, Shing-Tung Yau, and Stephen S.-T. Yau. An estimate of the gap of the first two eigenvalues in the Schrödinger operator. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 12(2):319–333, 1985.
- [22] Robert G. Smits. Spectral gaps and rates to equilibrium for diffusions in convex domains. *The Michigan Mathematical Journal*, 43(1):141–157, 1996.
- [23] M. van den Berg. On condensation in the free-boson gas and the spectrum of the Laplacian. *J. Statist. Phys.*, 31(3):623–637, 1983.
- [24] Shing-Tung Yau. *Nonlinear analysis in geometry*, volume 33 of *Monographies de L'Enseignement Mathématique [Monographs of L'Enseignement Mathématique]*. L'Enseignement Mathématique, Geneva, 1986. Série des Conférences de l'Union Mathématique Internationale [Lecture Series of the International Mathematics Union], 8.

- [25] Qi Huang Yu and Jia Qing Zhong. Lower bounds of the gap between the first and second eigenvalues of the Schrödinger operator. *Trans. Amer. Math. Soc.*, 294(1):341–349, 1986.
- [26] Jia Qing Zhong and Hong Cang Yang. On the estimate of the first eigenvalue of a compact Riemannian manifold. *Sci. Sinica Ser. A*, 27(12):1265–1273, 1984.

BEN ANDREWS
MATHEMATICAL SCIENCES INSTITUTE
AUSTRALIAN NATIONAL UNIVERSITY
ACT 2601
AUSTRALIA
YAU MATHEMATICAL SCIENCES CENTER
TSINGHUA UNIVERSITY
BEIJING 100084
CHINA
E-mail address: Ben.Andrews@anu.edu.au

JULIE CLUTTERBUCK
SCHOOL OF MATHEMATICS
MONASH UNIVERSITY
VIC 3800
AUSTRALIA
E-mail address: julie.clutterbuck@monash.edu

DANIEL HAUER
SCHOOL OF MATHEMATICS AND STATISTICS
THE UNIVERSITY OF SYDNEY
NSW 2006
AUSTRALIA
E-mail address: daniel.hauer@sydney.edu.au

RECEIVED FEBRUARY 20, 2019