

FAST COMMUNICATION

A SIMPLE DERIVATION OF BV BOUNDS FOR INHOMOGENEOUS  
RELAXATION SYSTEMS\*

BENOÎT PERTHAME<sup>†</sup>, NICOLAS SEGUIN<sup>‡</sup>, AND MAGALI TOURNUS<sup>§</sup>

**Abstract.** We consider relaxation systems of transport equations with heterogeneous source terms and with boundary conditions, whose limits are scalar conservation laws. Classical bounds, and in particular BV estimates, fail in this context. They are the simplest and the most standard way to prove compactness and convergence.

We provide a novel, simple method to obtain partial BV regularity and strong compactness in this framework. The standard notion of entropy is not convenient either and we also indicate another, closely related, notion. We give two examples motivated by renal flows which consist of  $2 \times 2$  and  $3 \times 3$  relaxation systems with 2-velocities, but the method is more general.

**Key words.** Hyperbolic relaxation; spatial heterogeneity; entropy condition; boundary conditions; strong compactness.

**AMS subject classifications.** 35L03, 35L60, 35B40, 35Q92.

1. Introduction

The usual framework of hyperbolic relaxation [3, 4, 16] concerns the convergence of a general hyperbolic system with stiff source terms toward a conservation law, when the relaxation parameter  $\epsilon$  goes to zero. More specifically, Jin and Xin [12] introduced a  $2 \times 2$  linear hyperbolic system with stiff source term that approximates any given conservation law. The problem of interest is to prove the convergence of the microscopic quantities depending on  $\epsilon$  toward the macroscopic quantities. A problem entering the framework of hyperbolic relaxation is motivated by very simplified models of kidney physiology [20, 21, 22] and fits the framework of Jin and Xin with two major differences: boundary conditions and spatial dependence of the source term. The type of boundary condition and the spatial dependence constitute the main novelty of the present study.

The system represents two solute concentrations  $u_\epsilon(x, t)$  and  $v_\epsilon(x, t)$ , and is written, for  $t \geq 0$  and  $x \in [0, L]$ ,

$$\begin{cases} \frac{\partial u_\epsilon}{\partial t}(x, t) + \frac{\partial u_\epsilon}{\partial x}(x, t) = \frac{1}{\epsilon} [h(v_\epsilon(x, t), x) - u_\epsilon(x, t)], \\ \frac{\partial v_\epsilon}{\partial t}(x, t) - \frac{\partial v_\epsilon}{\partial x}(x, t) = \frac{1}{\epsilon} [u_\epsilon(x, t) - h(v_\epsilon(x, t), x)], \\ u_\epsilon(0, t) = u_0, \quad v_\epsilon(L, t) = \alpha u_\epsilon(L, t), \quad \alpha \in (0, 1), \\ u_\epsilon(x, 0) = u^0(x) > 0, \quad v_\epsilon(x, 0) = v^0(x) > 0. \end{cases} \quad (1.1)$$

\*Received: April 5, 2014; accepted (in revised form): June 21, 2014. Communicated by Shi Jin.

<sup>†</sup>Sorbonne Universités, UPMC Univ Paris 06, CNRS UMR 7598, Laboratoire Jacques-Louis Lions, F-75005, Paris, France (benoit.perthame@upmc.fr).

<sup>‡</sup>Sorbonne Universités, UPMC Univ Paris 06, CNRS UMR 7598, Laboratoire Jacques-Louis Lions, F-75005, Paris, France (nicolas.seguin@upmc.fr).

<sup>§</sup>Sorbonne Universités, UPMC Univ Paris 06, CNRS UMR 7598, Laboratoire Jacques-Louis Lions, F-75005, Paris, France

Present address: Department of Mathematics, Pennsylvania State University, University Park, Pennsylvania 16802, USA (magali.tournus@ann.jussieu.fr).

The question of interest here is to understand the behavior of  $u_\epsilon$  and  $v_\epsilon$  when  $\epsilon$  vanishes. Another question, addressed in [19], is to explain the urine concentration mechanism due to the ‘pumps’, on the cell membranes, represented by the nonlinearity  $h(v, x)$ .

We make the following hypotheses:

$$h(0, x) = 0, \quad 1 < \beta \leq \frac{\partial h}{\partial v}(v, x) \leq \mu, \tag{1.2}$$

$$\sup_v \int_0^L \left| \frac{\partial h}{\partial x}(v, x) \right| dx \leq C, \quad h(\cdot, x) \text{ is not locally affine.} \tag{1.3}$$

Here  $C$ ,  $\beta$ , and  $\mu$  are positive constants. Also, we use the following bounds on the initial data:

$$u^0, v^0 \in L^\infty(0, L), \quad \frac{d}{dx}u^0 \in L^1(0, L), \quad \frac{d}{dx}v^0 \in L^1(0, L), \tag{1.4}$$

and, the well-prepared initial data condition

$$u^0(x) = h(v^0(x), x), \quad x \in [0, L]. \tag{1.5}$$

It is simple and standard to find the formal limit as  $\epsilon$  vanishes. Adding the two equations of (1.1), we find the conservation law

$$\frac{\partial [u_\epsilon + v_\epsilon](x, t)}{\partial t} + \frac{\partial [u_\epsilon - v_\epsilon](x, t)}{\partial x} = 0. \tag{1.6}$$

We expect that in the limit

$$u(x, t) = h(v(x, t), x),$$

and this identifies the limiting quasilinear conservation law

$$\frac{\partial [h(v(x, t), x) + v(x, t)]}{\partial t} + \frac{\partial [h(v(x, t), x) - v(x, t)]}{\partial x} = 0, \tag{1.7}$$

and the boundary conditions on  $\rho(x, t) = h(v(x, t), x) + v(x, t)$ , namely  $\rho(0, t) = u_0 + h^{-1}(u_0, 0)$  and  $\rho(L, t)$  is free and can be identified via the standard argument of Bardos, le Roux, and Nédélec [2], using the adapted Kruřkov entropies that we introduce in Section 3.

The justification poses particular difficulties due to the boundary condition and  $x$ -dependent flux.

One difficulty is that, in order to justify the above limit, an entropy identity is needed for two reasons: firstly to prove that equilibrium is reached [23], and, secondly, because at the limit  $\epsilon = 0$  a quasilinear equation arises, and shocks can be produced. Therefore, an entropy formulation is needed to uniquely define the solutions. When applied to this conservation equation, usual convex entropies as in either [6] or Kruřkov [14] contain a term with a derivative of  $h$  with respect to  $x$ . When the regularity of  $h$  is limited to BV,  $h_x$  is a measure and the weak entropy formulation is not well defined. One of our goals is to present a convenient notion of entropy with  $x$ -dependent relaxation in order to get rid of this problem. We use an entropy formulation introduced in [1] and adapted to scalar conservation laws with spatial heterogeneities.

Another difficulty is that, in order to prove the validity of the limit, strong compactness is also needed. Several methods have been developed in this goal. Uniform

bounds in the space of functions with bounded variation (BV for short) is the strongest method and the most standard. The homogeneous version of the system (1.1) on  $\mathbb{R}^n$  is treated in [13], where an  $L^1$  contraction property is used to establish BV bounds in  $x$ .

A similar approach is used in [17] where the authors deal with a similar homogeneous system with boundary conditions; the result is extended to a system with a source term in [24]. The authors of [10] also derive BV estimates for an inhomogeneous system that relaxes to a homogeneous system. More recently, and for the purpose of internal layers, the authors of [5] are interested in a similar setting and also study the limit when the relaxation time goes to zero only on a sub-domain of the space. More elaborate tools are the compensated compactness method [15] or the kinetic formulation and averaging lemmas [9, 18] (in particular extending the measure valued solutions of DiPerna [8]). These methods are weaker, because they give only convergence but no bounds, and thus apply to more general situations than BV bounds. Because of the spatial  $x$ -dependence of  $h$ , the BV framework is not available as of now for (1.1). Whereas time BV estimates follow immediately from the equations, compactness in the spatial direction cannot be obtained in this way. We propose a new method to prove spatial compactness. It does not use BV bounds on each component, but gives BV bounds in  $x$  for a single quantity and can be applied to spatially heterogeneous systems. We are going to prove the following:

PROPOSITION 1.1. *We make assumptions (1.2), (1.3), (1.4), (1.5), and fix a time  $T$ . Then*

- (i)  $u_\epsilon$  and  $v_\epsilon$  are uniformly bounded in  $L^\infty([0, L] \times [0, T])$ ,
- (ii)  $\frac{\partial}{\partial t} u_\epsilon$ ,  $\frac{\partial}{\partial t} v_\epsilon$ , and  $\frac{\partial}{\partial x} [u_\epsilon - v_\epsilon]$  are bounded in  $L^\infty([0, T]; L^1(0, L))$ ,
- (iii) there exists  $v \in L^\infty([0, L] \times [0, T])$  such that

$$u_\epsilon(x, t) \xrightarrow{\epsilon \rightarrow 0} h(v(x, t), x), \quad v_\epsilon(x, t) \xrightarrow{\epsilon \rightarrow 0} v(x, t), \text{ a.e.} \tag{1.8}$$

- (iv) the equation (1.7) is satisfied and entropy inequalities hold (see Section 3).

However, it is not correct that  $\frac{\partial}{\partial x} u_\epsilon$  and  $\frac{\partial}{\partial x} v_\epsilon$  are separately bounded in  $L^\infty([0, T]; L^1(0, L))$ .

**2.  $L^\infty$  bound**

We first prove uniform estimates. Unlike the homogeneous case,  $L^\infty$  bounds are not always true and the most general existence theory is in  $L^1$  (see [7]). Here we build particular sub and supersolutions of (1.1) which are uniformly bounded in  $\epsilon$ .

LEMMA 2.1. *The solution of (1.1) satisfies the uniform estimate*

$$\|u_\epsilon\|_{L^\infty([0, L] \times [0, T])} \leq K(\beta, u_0, u^0, v^0), \quad \|v_\epsilon\|_{L^\infty([0, L] \times [0, T])} \leq K(\beta, u_0, u^0, v^0). \tag{2.1}$$

*Proof.* To obtain an  $L^\infty$  bound on the time-dependent solution, we follow the approach of [1] and use the comparison principle with appropriate supersolution. Indeed, because of the  $x$ -dependence of  $h$ , constant functions are not supersolutions of the stationary problem. We introduce the stationary version of (1.1)

$$\begin{cases} \frac{dU_\epsilon}{dx}(x) = \frac{1}{\epsilon} [h(V_\epsilon(x), x) - U_\epsilon(x)], \\ -\frac{dV_\epsilon}{dx}(x) = \frac{1}{\epsilon} [U_\epsilon(x) - h(V_\epsilon(x), x)], \\ U_\epsilon(0) = U_0 > 0, \quad V_\epsilon(L) = \alpha U_\epsilon(L). \end{cases} \tag{2.2}$$

We are going to prove that there exists a smooth supersolution  $(U_\epsilon, V_\epsilon)$  of the stationary problem (2.2), and a constant  $K(U_0, \beta) > 0$  such that

$$0 \leq U_\epsilon(x) \leq K(U_0, \beta), \quad 0 \leq V_\epsilon \leq K(U_0, \beta). \tag{2.3}$$

This will conclude the proof of Lemma 2.1 because a solution of (2.2) where  $U_0 \geq u_0$ , is a supersolution of (1.1), and the comparison principle gives  $0 \leq u_\epsilon \leq U_\epsilon$  and  $0 \leq v_\epsilon \leq V_\epsilon$ .

Because it has been proved in [21] that (2.2) admits a unique solution which lies in  $BV([0, L] \times [0, T])$  (using a fixed point argument for the existence and contraction for the uniqueness), it remains to prove that a solution of (2.2) with  $U_0 \geq u_0$  is uniformly bounded in  $\epsilon$ .

Adding the two lines of (2.2), we obtain a quantity which does not depend on  $x$ ,

$$U_\epsilon(x) - V_\epsilon(x) =: K_\epsilon. \tag{2.4}$$

Using the boundary values, we find uniform bounds on  $K_\epsilon$

$$K_\epsilon = U_0 - V_\epsilon(0) \leq U_0, \quad K_\epsilon = U_\epsilon(L) - V_\epsilon(L) = (1 - \alpha)U_\epsilon(L) \geq 0, \tag{2.5}$$

and thus

$$0 \leq K_\epsilon \leq U_0, \quad U_\epsilon(L) \leq \frac{U_0}{1 - \alpha}. \tag{2.6}$$

Hence, we just have to prove that  $U_\epsilon$  is uniformly bounded in  $L^\infty$ , knowing that  $U_0$  and  $U_\epsilon(L)$  are uniformly bounded in  $\mathbb{R}$ . For that, we use the maximum principle assuming  $\mathcal{C}^1$  regularity (one can easily justify it for a regularized function  $h(v, x)$  and pass to the limit). Indeed, if  $U_\epsilon$  reaches its maximal value on the boundary, the result follows from (2.6). If  $U_\epsilon$  reaches its maximal value at  $x_0 \in [0, L]$ , then

$$0 = \frac{dU_\epsilon}{dx}(x_0) = \frac{1}{\epsilon} \left( h(U_\epsilon(x_0) - K_\epsilon, x_0) - U_\epsilon(x_0) \right), \tag{2.7}$$

and thus by assumption (1.2),

$$U_\epsilon(x_0) = h(U_\epsilon(x_0) - K_\epsilon, x_0) \geq \beta U_\epsilon(x_0) - \beta K_\epsilon. \tag{2.8}$$

Finally, (2.3) is proved because the above inequality gives

$$U_\epsilon(x_0) \leq \frac{\beta}{\beta - 1} K_\epsilon. \tag{2.9}$$

□

### 3. Adapted (heterogeneous) entropies

As explained in the introduction, entropies are useful to derive additional bounds and to characterize the limit as  $\epsilon$  vanishes, see [4, 11, 23]. We are going to use specific entropies adapted to spatial dependence.

We recall the usual approach which is to define the following:

**DEFINITION 3.1 (Entropy pair).** *We call an entropy pair for the system (1.1) a couple of functions  $(S, \Sigma)$  in  $BV([0, 1], \mathcal{C}(\mathbb{R}))$  which satisfy*

$$(i) S(., x) \text{ and } \Sigma(., x) \text{ are convex,} \quad (ii) \frac{\partial S}{\partial u}(h(v, x), x) = \frac{\partial \Sigma}{\partial v}(v, x). \tag{3.1}$$

For such entropy pairs, it is immediate to check that

$$\begin{aligned} & \frac{\partial}{\partial t} [S(u_\epsilon, x) + \Sigma(v_\epsilon, x)] + \frac{\partial}{\partial x} [S(u_\epsilon, x) - \Sigma(v_\epsilon, x)] \\ &= \frac{1}{\epsilon} \left( \frac{\partial S}{\partial v}(u_\epsilon, x) - \frac{\partial \Sigma}{\partial v}(v_\epsilon, x) \right) (h(v_\epsilon, x) - u_\epsilon) + \frac{\partial S}{\partial x}(u_\epsilon, x) - \frac{\partial \Sigma}{\partial x}(v_\epsilon, x). \end{aligned}$$

Now, using (3.1)(ii), we can write

$$\begin{aligned} & \frac{\partial}{\partial t} [S(u_\epsilon, x) + \Sigma(v_\epsilon, x)] + \frac{\partial}{\partial x} [S(u_\epsilon, x) - \Sigma(v_\epsilon, x)] \\ &= \frac{1}{\epsilon} \left( \frac{\partial S}{\partial v}(u_\epsilon, x) - \frac{\partial S}{\partial v}(h(v_\epsilon, x), x) \right) (h(v_\epsilon, x) - u_\epsilon) + \frac{\partial S}{\partial x}(u_\epsilon, x) - \frac{\partial \Sigma}{\partial x}(v_\epsilon, x) \\ &\leq \frac{\partial S}{\partial x}(u_\epsilon, x) - \frac{\partial \Sigma}{\partial x}(v_\epsilon, x), \end{aligned}$$

because  $S$  is convex with respect to its first variable and thus  $\frac{\partial S}{\partial v}$  is non decreasing with respect to its first variable.

The shortcoming of Definition 3.1 is that the above right hand side is not always well defined for BV functions  $u_\epsilon$  and  $v_\epsilon$ . Indeed, being given  $S(u, x)$ , we compute

$$\Sigma(v, x) = \int_0^v S'_u(h(\bar{v}, x), x) d\bar{v},$$

and the expression for the  $x$ -derivative

$$\frac{\partial \Sigma}{\partial x}(v, x) = \int_0^v [S''_{uu}(h(\bar{v}, x), x) h_x(\bar{v}, x) + \dots] d\bar{v}$$

does not make intrinsic sense.

However, this entropy inequality is enough to prove that equilibrium is reached. Because the expressions  $\frac{\partial S}{\partial x}(u_\epsilon, x)$  and  $\frac{\partial \Sigma}{\partial x}(v_\epsilon, x)$  are bounded thanks to our assumptions on  $h$ , we may choose

$$S(u) = \frac{u^2}{2}, \quad \Sigma(v, x) = \int_0^v h(\bar{v}, x) d\bar{v},$$

and, using the entropy dissipation, with assumption (1.3), we find after integration of equality (3) in  $(x, t)$ , that for all  $T > 0$  and some constant  $C(T)$  which does not depend on  $\epsilon$ , it holds

$$\frac{1}{\epsilon} \int_0^T \int_0^L |u_\epsilon(x, t) - h(v_\epsilon(x, t), x)|^2 dx dt \leq C(T).$$

Therefore, we arrive at the conclusion

PROPOSITION 3.2. *For  $u_\epsilon$  and  $v_\epsilon$  solutions of (1.1), we have the convergence*

$$u_\epsilon - h(v_\epsilon, x) \xrightarrow[\epsilon \rightarrow 0]{} 0, \quad L^2\left([0, L] \times [0, T]\right). \tag{3.2}$$

This convergence result toward equilibrium is an intermediate step in the proof of Proposition 1.1.

We wish to go further and avoid the two terms containing  $x$ -derivatives. For this goal, we define adapted (heterogeneous) entropies by imposing more restrictions on the spatial dependence of the entropy pair.

**DEFINITION 3.3** (An adapted (heterogeneous) entropy family). *The pair of continuous functions  $(S, \Sigma)$ , is an adapted (heterogeneous) entropy for the system (1.1) if it satisfies the conditions of Definition 3.1 and if*

$$-\frac{\partial S}{\partial x}(h(v, x), x) + \frac{\partial \Sigma}{\partial x}(v, x) = 0, \quad \forall v \geq 0. \tag{3.3}$$

An example of such a family of entropies parametrized by  $p \in \mathbb{R}$  are the adapted Kruřkov entropies

$$S_p(u, x) = |u - h(k_p(x), x)|, \quad \Sigma_p(v, x) = |v - k_p(x)|, \tag{3.4}$$

where  $(k_p)_{p \in \mathbb{R}}$  is the family of stationary solutions of the limit Equation (1.7), which is equivalent to saying

$$h(k_p(x), x) - k_p(x) = p.$$

With this choice of an entropy pair, the above entropy inequality then reduces to

$$\frac{\partial}{\partial t} [S_p(u_\epsilon, x) + \Sigma_p(v_\epsilon, x)] + \frac{\partial}{\partial x} [S_p(u_\epsilon, x) - \Sigma_p(v_\epsilon, x)] \leq 0.$$

As a consequence, thanks to the strong compactness proven in next part, in the limit  $\epsilon \rightarrow 0$ , the quasilinear conservation law (1.7) comes with the family of adapted Kruřkov entropies. Indeed, if we define

$$\rho(x, t) := h(v(x, t), x) + v(x, t), \quad A(\rho, t) := h(v(x, t), x) - v(x, t), \tag{3.5}$$

then, as in [1] and because  $h$  is increasing, the following entropy inequality holds in the sense of distributions

$$\frac{\partial}{\partial t} \left| \rho(x, t) - \left( k_p(x) + h(k_p(x), x) \right) \right| + \frac{\partial}{\partial x} \left| A(\rho, x) - A\left( k_p(x) + h(k_p(x), x), x \right) \right| \leq 0. \tag{3.6}$$

**4. BV bounds**

For well prepared initial conditions (1.5), we present here our method to prove BV bounds for appropriate quantities and strong compactness for  $u_\epsilon$  and  $v_\epsilon$ , that are points (ii), (iii) of Proposition 1.1.

**1st step. A bound on the time derivative at  $t = 0$ .** Our first statement is

$$\int_0^L \left| \frac{\partial u_\epsilon}{\partial t}(x, 0) \right| dx \leq K_1(u^0), \quad \int_0^L \left| \frac{\partial v_\epsilon}{\partial t}(x, 0) \right| dx \leq K_2(v^0). \tag{4.1}$$

Indeed, because initial conditions are at equilibrium, we have  $\frac{\partial v_\epsilon}{\partial t}(x, 0) - \frac{\partial v_\epsilon}{\partial x}(x, 0) = 0$ .

We multiply this equality by  $\text{sign} \left( \frac{\partial}{\partial t} v_\epsilon \right) (x, 0)$  and integrate over  $[0, L]$  to get

$$\int_0^L \left| \frac{\partial v_\epsilon}{\partial t}(x, 0) \right| dx = \int_0^L \left| \frac{\partial v_\epsilon}{\partial x}(x, 0) \right| dx \leq K_2(v^0), \tag{4.2}$$

and the above inequality follows from assumption (1.4). This gives the first inequality of estimates (4.1). The same argument applies for  $u_\epsilon$ .

**2nd step. The time BV estimate.** To prove (2.1), we differentiate each line of (1.1) with respect to time and we multiply it respectively by  $\text{sign}\left(\frac{\partial}{\partial t}u_\epsilon\right)$  and  $\text{sign}\left(\frac{\partial}{\partial t}v_\epsilon\right)$ , and integrate in  $x$ . Adding the two lines, we obtain

$$\begin{aligned} \frac{d}{dt} \int_0^L [|\frac{\partial}{\partial t}u_\epsilon| + |\frac{\partial}{\partial t}v_\epsilon|](x,t)dx &\leq |\frac{\partial}{\partial t}u_0| - |\frac{\partial}{\partial t}u_\epsilon(L,t)| + |\frac{\partial}{\partial t}v_\epsilon(L,t)| - |\frac{\partial}{\partial t}v_\epsilon(0,t)| \\ &= -\frac{1}{2}|\frac{\partial}{\partial t}u_\epsilon(L,t)| - |\frac{\partial}{\partial t}v_\epsilon(0,t)| \leq 0, \end{aligned} \tag{4.3}$$

which implies, using estimate (4.1),

$$\int_0^L [|\frac{\partial}{\partial t}u_\epsilon| + |\frac{\partial}{\partial t}v_\epsilon|](x,t)dx \leq \int_0^L [|\frac{\partial}{\partial t}u_\epsilon| + |\frac{\partial}{\partial t}v_\epsilon|](x,0)dx \leq K_1(u^0) + K_2(v^0). \tag{4.4}$$

**3rd step. BV bound in  $x$ .** We complete the proof of Proposition 1.1 (ii). Because of the space dependence of  $h$ , we cannot apply the same arguments for  $x$ -derivatives and build a single BV quantity. We add the two lines of (1.1) and obtain

$$\left(\frac{\partial}{\partial x}(u_\epsilon - v_\epsilon)\right)(x,t) = -\left(\frac{\partial}{\partial t}(u_\epsilon + v_\epsilon)\right)(x,t).$$

Using (2.1), we thus conclude that, for all  $t \geq 0$ ,

$$\int_0^L \left| \frac{\partial}{\partial x}(u_\epsilon - v_\epsilon) \right|(x,t)dx \leq K_1(u^0) + K_2(v^0).$$

**4th step. Compactness.** Therefore we can conclude that  $(u_\epsilon - v_\epsilon)$  is compact in  $L^1([0,L] \times [0,T])$ . Now, thanks to Proposition (3.2),  $(h(v_\epsilon, \cdot) - u_\epsilon)$  is compact in  $L^1([0,L] \times [0,T])$ . A combination of these two last compact embeddings gives us that

$$h(v_\epsilon, \cdot) - v_\epsilon \text{ is compact in } L^1([0,L] \times [0,T]). \tag{4.5}$$

Therefore, there is a function  $Q \in L^\infty(0,L)$  such that, after extraction of a subsequence,

$$h(v_\epsilon, \cdot) - v_\epsilon \xrightarrow{\epsilon \rightarrow 0} Q(x),$$

because  $v \mapsto h(v,x) - v$  is one-to-one, thanks to assumptions (1.2). In the same way, we conclude that the sequence  $v_\epsilon$  converges. Gathering the informations above, Proposition 1.1 (iii) is proved.

Then we can pass to the limit and obtain the last statement, Proposition 1.1 (iv).

**5. Extension to a more specific relaxation system**

The method we have developed so far can be extended to a more realistic problem arising in kidney physiology, which motivated this study. The system introduced and

studied in [21] is written, for  $t \geq 0$  and  $x \in [0, L]$ ,

$$\begin{cases} \frac{\partial C_\epsilon^1}{\partial t}(x, t) + \frac{\partial C_\epsilon^1}{\partial x}(x, t) = \frac{1}{3\epsilon} \left[ C_\epsilon^2(x, t) + h(C_\epsilon^3(x, t), x) - 2C_\epsilon^1(x, t) \right], \\ \frac{\partial C_\epsilon^2}{\partial t}(x, t) + \frac{\partial C_\epsilon^2}{\partial x}(x, t) = \frac{1}{3\epsilon} \left[ C_\epsilon^1(x, t) + h(C_\epsilon^3(x, t), x) - 2C_\epsilon^2(x, t) \right], \\ \frac{\partial C_\epsilon^3}{\partial t}(x, t) - \frac{\partial C_\epsilon^3}{\partial x}(x, t) = \frac{1}{3\epsilon} \left[ C_\epsilon^1(x, t) + C_\epsilon^2(x, t) - 2h(C_\epsilon^3(x, t), x) \right], \\ C_\epsilon^1(0, t) = C_0^1, \quad C_\epsilon^2(0, t) = C_0^2, \quad C_\epsilon^3(L, t) = C_\epsilon^2(L, t), \quad t > 0, \end{cases} \tag{5.1}$$

Again, we want to prove uniform BV bounds for a small parameter  $\epsilon$ , which measures the ratio between ionic exchanges and flow along the tubules.

We make the same assumptions (1.2) (the condition  $1 < \beta \leq \frac{\partial h}{\partial v}(v, x)$  can be relaxed to  $1 \leq \frac{\partial h}{\partial v}(v, x)$ ) and (1.3), and the same hypotheses on the initial conditions, namely that they belong to BV and are at equilibrium, which means

$$C^1 = C^2 = h(C^3, x).$$

Following (3.5), the conservative quantity  $\rho$  and the flux  $B$  are defined by

$$\rho(x, t) := 2h(C^3(x, t), x) + C^3(x, t), \quad B(\rho, t) := 2h(C^3(x, t), x) - C^3(x, t). \tag{5.2}$$

For  $p \in \mathbb{R}$ , we define uniquely the steady state  $k_p$  by

$$B(k_p(x), x) = p. \tag{5.3}$$

**THEOREM 5.1 (Limit  $\epsilon \rightarrow 0$ ).** *The functions  $C_\epsilon^i$ ,  $i = 1, 2, 3$  converge almost everywhere to bounded functions  $C^i$  and the quantity  $\rho(x, t)$  is an entropy solution to*

$$\begin{cases} \frac{\partial}{\partial t} \rho(x, t) + \frac{\partial}{\partial x} B(\rho(x, t), x) = 0, & t > 0, \quad x \in [0, L], \\ \rho(0, t) = C_0^1 + C_0^2 + h^{-1}\left(\frac{C_0^1 + C_0^2}{2}, 0\right), & t > 0, \\ \rho(x, 0) = \rho^0(x), \quad \rho^0(x) := C^0(x) + 2h(C^0(x), x), & x \in [0, L]. \end{cases} \tag{5.4}$$

The entropy formulation of the conservation law for adapted (heterogeneous) entropies is written, following [1] again,

$$\frac{\partial}{\partial t} \left| \rho(x, t) - \left( 2k_p(x) + h(k_p(x), x) \right) \right| + \frac{\partial}{\partial x} \left| B(\rho, x) - B\left( 2k_p(x) + h(k_p(x), x), x \right) \right| \leq 0.$$

This family of inequalities is enough to prove uniqueness as in [17] (see [20] for details).

The boundary conditions are understood in the following sense:

**Boundary condition at  $x = 0$ :** for all  $k_p$  such that  $k_p(0) + 2h(k_p(0), 0) \in I\left(\rho(0, t), h^{-1}\left(\frac{C_0^1 + C_0^2}{2}, 0\right) + C_0^1 + C_0^2\right)$ , we have

$$\text{sign} \left( \rho(0, t) - h^{-1}\left(\frac{C_0^1 + C_0^2}{2}, 0\right) - C_0^1 - C_0^2 \right) \left( B(\rho(0, t), 0) - [2h(k_p(0), 0) - k_p(0)] \right) \leq 0, \tag{5.5}$$



Boundary condition at  $x=L$ : for all  $k_p$  such that  $k_p(L) + 2h(k_p(L), L) \in I\left(\rho(L, t), 2w_L + h^{-1}(w_L, L)\right)$ , we have

$$\text{sign}\left(\rho(L, t) - 2w_L - h^{-1}(w_L, L)\right)\left(B(\rho(L, t), L) - [2h(k_p(L), L) - k_p(L)]\right) \geq 0, \quad (5.6)$$

where  $w_L(t) := \lim_{\epsilon \rightarrow 0} C_\epsilon^1(L, t)$ , and where  $I(a, b)$  denotes the interval  $(\min(a, b), \max(a, b))$ .

Following the arguments we gave for the  $2 \times 2$  system, which we do not repeat, we can prove BV bounds in several steps

- (i)  $C_\epsilon^1$ ,  $C_\epsilon^2$  and  $C_\epsilon^3$  are bounded in  $L^\infty((0, \infty) \times (0, L))$ ,
- (ii)  $C_\epsilon^2(x, t) + h(C_\epsilon^3(x, t), x) - 2C_\epsilon^1(x, t) \xrightarrow{\epsilon \rightarrow 0} 0$ ,  $C_\epsilon^1(x, t) + h(C_\epsilon^3(x, t), x) - 2C_\epsilon^2(x, t) \xrightarrow{\epsilon \rightarrow 0} 0$  in  $L^2((0, \infty) \times (0, L))$ ,
- (iii)  $\frac{\partial C_\epsilon^1}{\partial t}$ ,  $\frac{\partial C_\epsilon^2}{\partial t}$ ,  $\frac{\partial C_\epsilon^3}{\partial t}$  are bounded in  $L^\infty((0, \infty; L^1(0, L)))$ ,
- (iv)  $\frac{\partial C_\epsilon^1}{\partial x} + \frac{\partial C_\epsilon^2}{\partial x} + \frac{\partial C_\epsilon^3}{\partial x}$  is bounded in  $L^\infty((0, \infty; L^1(0, L)))$ .

These statements prove the convergence result in Theorem 5.1.

**Acknowledgement.** Funding for this study was provided by the program EMERGENCE (EME 0918) of the Université Pierre et Marie Curie (Paris Univ. 6).

#### REFERENCES

- [1] E. Audusse and B. Perthame, *Uniqueness for scalar conservation laws with discontinuous flux via adapted entropies*, Proc. Roy. Soc. Edinburgh Sect. A, 135, 253–265, 2005.
- [2] C. Bardos, A.Y. le Roux, and J.-C. Nédélec, *First order quasilinear equations with boundary conditions*, Comm. Part. Diff. Eqs., 4, 1017–1034, 1979.
- [3] F. Bouchut, *Nonlinear Stability of Finite Volume Methods for Hyperbolic Conservation Laws and Well-Balanced Schemes for Sources*, Birkhäuser Verlag, 2004.
- [4] G.Q. Chen, C.D. Levermore, and T.-P. Liu, *Hyperbolic conservation laws with stiff relaxation terms and entropy*, Comm. Pure Appl. Math., 47, 787–830, 1994.
- [5] F. Coquel, S. Jin, J.G. Liu, and L. Wang, *Well-posedness and singular limit of a semilinear hyperbolic relaxation system with a two-scale discontinuous relaxation rate*, Arch. Rational Mech. Anal., 214, 1051–1084, 2014.
- [6] C.M. Dafermos, *Hyperbolic conservation laws in continuum physics*, in Grundlehren der Mathematischen Wissenschaften, Fundamental Principles of Mathematical Sciences, Third Edition, Springer-Verlag, 2010.
- [7] A.-L. Dalibard, *Kinetic formulation for a parabolic conservation law. Application to homogenization*, SIAM J. Math. Anal., 39, 891–915 (electronic), 2007.
- [8] R.J. DiPerna, *Measure-valued solutions to conservation laws*, Arch. Rational Mech. Anal., 88, 223–270, 1985.
- [9] F. Golse, P.-L. Lions, B. Perthame, and R. Sentis, *Regularity of the moments of the solution of a transport equation*, J. Funct. Anal., 76, 110–125, 1988.
- [10] L. Gosse and G. Toscani, *Space localization and well-balanced schemes for discrete kinetic models in diffusive regimes*, SIAM J. Numer. Anal., 41, 641–658 (electronic), 2003.
- [11] F. James, *Convergence results for some conservation laws with a reflux boundary condition and a relaxation term arising in chemical engineering*, SIAM J. Math. Anal., 29, 1200–1223 (electronic), 1998.
- [12] S. Jin and Z.P. Xin, *The relaxation schemes for systems of conservation laws in arbitrary space dimensions*, Comm. Pure Appl. Math., 48, 235–276, 1995.
- [13] M.A. Katsoulakis and A.E. Tzavaras, *Contractive relaxation systems and the scalar multidimensional conservation law*, Comm. Part. Diff. Eqs., 22, 195–233, 1997.
- [14] S.N. Kružkov, *First order quasilinear equations with several independent variables*, Mat. Sb. (N.S.), 81(123), 228–255, 1970.
- [15] F. Murat, *Compacité par compensation*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 5, 489–507, 1978.

- [16] R. Natalini, *Recent results on hyperbolic relaxation problems*, in Analysis of systems of conservation laws (Aachen, 1997), Chapman & Hall/CRC Monogr. Surv. Pure Appl. Math., Chapman & Hall/CRC, Boca Raton, FL, 99, 128–198, 1999.
- [17] R. Natalini and A. Terracina, *Convergence of a relaxation approximation to a boundary value problem for conservation laws*, Comm. Part. Diff. Eqs., 26, 1235–1252, 2001.
- [18] B. Perthame, *Kinetic formulation of conservation laws*, Oxford Lecture Series in Mathematics and its Applications, Oxford University Press, Oxford, 21, 2002.
- [19] M. Tournus, *An asymptotic study to explain the role of active transport in models with counter-current exchangers*, SeMA J., 59, 19–35, 2012.
- [20] M. Tournus, *Modèles d'échanges ioniques dans le rein: théorie, analyse asymptotique et applications numériques*, PhD thesis, July 2013.
- [21] M. Tournus, A. Edwards, N. Seguin, and B. Perthame, *Analysis of a simplified model of the urine concentration mechanism*, Netw. Heterog. Media, 7, 989–1018, 2012.
- [22] M. Tournus, N. Seguin, B. Perthame, S. R. Thomas, and A. Edwards, *A model of calcium transport along the rat nephron*, American Journal of Physiology-Renal Physiology, 305, F979–F994, 2013.
- [23] A.E. Tzavaras, *Relative entropy in hyperbolic relaxation*, Commun. Math. Sci., 3, 119–132, 2005.
- [24] S.Y. Zhang and Y.G. Wang, *Well-posedness and asymptotics for initial boundary value problems of linear relaxation systems in one space variable*, Z. Anal. Anwendungen, 23, 607–630, 2004.