

SPECTRAL RESULTS FOR PERTURBED VARIATIONAL EIGENVALUE PROBLEMS AND THEIR APPLICATIONS TO COMPRESSED PDES*

ÖMER FARUK TEKİN[†], KE YIN[‡], AND FARZIN BAREKAT[§]

Abstract. We consider the solutions to a modification of the Courant’s minimax characterization of the Dirichlet eigenfunctions of second-order linear symmetric elliptic operators in a bounded domain Ω in \mathbb{R}^d . In particular, we perturb the objective functional by an arbitrary bounded penalty term. Without perturbation, it is well known that the Courant minimax principle yields the eigenfunctions, which form an orthonormal basis for $L^2(\Omega)$. We prove that the solutions of the perturbed problem still form an orthonormal basis in the case of $d = 1$ and $d = 2$, provided that the perturbation is sufficiently small in the latter case. As an application, we prove completeness results for compressed plane waves and compressed modes, which are the solutions to analogous variational problems with perturbations being an L^1 -regularization term. The completeness theory for these functions sets a foundation for finding a computationally efficient basis for the representation of the solution of differential equations.

Key words. Variational eigenvalue problems, spectral methods, L^1 -regularization.

AMS subject classifications. 35P10, 49G20, 49R05.

1. Introduction

For a second-order symmetric elliptic operator L on a bounded domain $\Omega \subset \mathbb{R}^d$, consider the following Dirichlet eigenvalue problem:

$$\begin{aligned}Lu &= \lambda u \text{ in } \Omega, \\u &= 0 \text{ on } \partial\Omega.\end{aligned}$$

The spectral theorem for the second-order elliptic operators asserts that the Dirichlet eigenfunctions of L form an orthonormal basis for $L^2(\Omega)$. Furthermore, the eigenvalues and eigenfunctions can be characterized via a hierarchical variational procedure involving the minimization of the functional $B[u, u] = \langle Lu, u \rangle_{L^2}$, known as the Courant variational method (see e.g. [6, 8]).

In this paper, we consider the Courant variational problem under a non-negative perturbation and prove for it an analogue of the spectral theorem. We treat this problem in a general Hilbert space setting, where we consider an arbitrary self-adjoint operator T whose inverse is compact. Namely, we rather minimize the functional

$$\mathcal{J}(u) := P(u) + \langle Tu, u \rangle,$$

in the domain of definition for T , where P is a non-negative penalty term. We verify that the spectral theorem still holds, as long as the eigenvalues of the original problem grow sufficiently fast. In particular, the growth condition holds when T is the weak realization of an elliptic operator on a bounded domain Ω lying inside \mathbb{R} or \mathbb{R}^2 .

*Received: June 29, 2014; accepted (in revised form): July 7, 2015. Communicated by Lexing Ying. This research was supported by NSF DMS 0914561 and 1118971. Research of Barekat was also supported by an NSERC PGS-D award.

[†]Mathematics Department, University of California at Los Angeles, Los Angeles, CA 90095-1555 USA (omerftekin@math.ucla.edu).

[‡]Mathematics Department, University of California at Los Angeles, Los Angeles, CA 90095-1555 USA (kyin@math.ucla.edu).

[§]Mathematics Department, University of California at Los Angeles, Los Angeles, CA 90095-1555 USA (fbarekat@math.ucla.edu).

As an application, we consider the spatially localized (“sparse”) modes introduced in [9, 10]. In [5, 7], it was proven that perturbing the variational quantity of certain elliptic and parabolic PDEs would result in compact support. In [9, 10], the authors consider variational problems corresponding to Schrödinger’s equation or Laplace’s equation, whose functional is modified by an L^1 -regularization term. Namely, their construction involves the minimization of the functional

$$\mathcal{J}(u) := \frac{1}{\mu} \|u\|_1 + \langle Lu, u \rangle_{L^2},$$

where $L = -\frac{1}{2}\Delta$, or $L = -\frac{1}{2}\Delta + V(x)$. By minimizing the above functional subject to the orthogonality constraint, so-called “compressed modes” are obtained by the authors in [9]. In a different formulation, where shift-orthogonality is imposed in addition to orthogonality, the minimizers are called the “compressed plane waves” [10]. Numerical experiments suggest that the regularization parameter μ for the L^1 term is used to balance between the consistency with the original problem and the sparsity of the solution. The consistency and the sparsity of compressed modes as μ varies have been verified in [1–3].

Our results yield further consistency results for compressed modes and compressed plane waves, as we conclude that the analogous spectral properties are preserved under perturbation. Such “completeness” conjectures are proposed in [9, 10]. In [9], the authors predicted that the compressed modes approximate the true eigenstates with high precision, whereas in [10], they conjectured that the compressed plane waves form a basis in $L^2(\Omega)$, where Ω is some bounded rectangular domain in \mathbb{R}^d . We will provide an affirmative answer to these conjectures in dimensions one and two, and we will propose Conjecture 2.2, which is phrased in a general Hilbert space setting. We note that this conjecture implies the above-mentioned conjectures.

This paper is organized as follows. In Section 2, we study the theory of completeness for the solutions of the perturbed Courant–Fisher variational problem for arbitrary linear operators defined on a Hilbert space. We prove the completeness in theorems 2.7 and 2.8 under linear and super-linear growth conditions on the eigenvalues of the original operator. Without the completeness, we still have Theorem 2.9, which yields an estimation of the deviation from the true solutions. Lemma 2.5 is the essential tool in the proof of these results. In sections 3 and 4, we apply the theory developed in Section 2 to the second-order linear symmetric elliptic operators.

2. Perturbed variational problems associated to linear operators

The proof of the spectral theorem for elliptic operators relies on the fact that the “inverse” of the elliptic differential operator is compact, hence the following spectral theorem for symmetric compact operators holds:

THEOREM 2.1 (Spectral Theorem for Compact Operators). *Let \mathcal{H} be a Hilbert space and $K : \mathcal{H} \rightarrow \mathcal{H}$ be a linear symmetric compact operator. Then,*

1. K has real eigenvalues ν_k , and $\nu_k \rightarrow 0$ as $k \rightarrow \infty$ and
2. The (normalized) eigenvectors $\{\phi_k\}_{k=1}^\infty$, with $K\phi_k = \nu_k\phi_k$, form a complete orthonormal system in \mathcal{H} .

As a consequence of this version of the spectral theorem, the inverse T , of a positive compact operator K satisfies the following spectral theorem:

THEOREM 2.2 (Spectral theorem for inverse-compact operators). *Let \mathcal{H} be a Hilbert space and $K : \mathcal{H} \rightarrow \mathcal{H}$ be a linear bijective symmetric compact operator that is also bounded below. Then, $T = K^{-1}$ satisfies the following properties:*

1. T has real eigenvalues λ_k , with $\{\lambda_k\}_{k=1}^\infty$ in increasing order, and $\lambda_k \rightarrow +\infty$ as $k \rightarrow \infty$,
2. The (normalized) eigenvectors $\{\phi_k\}_{k=1}^\infty$, with $T\phi_k = \lambda_k\phi_k$, form a complete orthonormal system.

REMARK 2.1. Notice that the operator T defined in Theorem 2.2 is unbounded, so T must have a domain of definition, $\mathcal{D}(T)$, for which it is self-adjoint. We consider the following natural choice of the domain of definition:

$$\mathcal{D}(T) = \left\{ \alpha = \sum_{n \in \mathbb{N}} \hat{\alpha}_n \phi_n \in \mathcal{H} \mid \sum_{n \in \mathbb{N}} \lambda_n \hat{\alpha}_n \phi_n \in \mathcal{H}, \text{ or equivalently, } \sum_{n \in \mathbb{N}} \lambda_n^2 |\hat{\alpha}_n|^2 < \infty \right\},$$

in which there will be no ambiguity of the definition of T .

We will now work with the operators T that can be represented as the inverse of some bijective symmetric compact operator. The eigenvalues and eigenvectors of T can be characterized via the following Courant–Fisher variational formulae (see e.g. [6]):

$$\begin{aligned} \lambda_1 &= \min_{\substack{u \in \mathcal{D}(T) \\ \|u\|=1}} \langle Tu, u \rangle, \\ \phi_1 &= \operatorname{argmin}_{\substack{u \in \mathcal{D}(T) \\ \|u\|=1}} \langle Tu, u \rangle, \\ \lambda_k &= \min_{\substack{u \in \mathcal{D}(T) \\ u \in \{\phi_1, \dots, \phi_{k-1}\}^\perp \\ \|u\|=1}} \langle Tu, u \rangle, \\ \phi_k &= \operatorname{argmin}_{\substack{u \in \mathcal{D}(T) \\ u \in \{\phi_1, \dots, \phi_{k-1}\}^\perp \\ \|u\|=1}} \langle Tu, u \rangle. \end{aligned}$$

We consider a similar variational problem, where we perturb the objective functional $\langle Tu, u \rangle$. Strictly speaking, we define

$$J[u] = \langle Tu, u \rangle + P(u),$$

where $P : \mathcal{H} \rightarrow \mathbb{R}$ is a non-negative penalty term. We view the term $J[u]$ as the “energy” of the element u , and we run a progressive energy-minimization procedure as in the Courant–Fisher formulae. In other words, we define

$$\begin{aligned} \zeta_1 &= \operatorname{argmin}_{\substack{u \in \mathcal{D}(T) \\ \|u\|=1}} J[u], \\ \zeta_k &= \operatorname{argmin}_{\substack{u \in \mathcal{D}(T) \\ u \in \{\zeta_1, \dots, \zeta_{k-1}\}^\perp \\ \|u\|=1}} J[u]. \end{aligned} \tag{2.1}$$

In case of a non-uniqueness in the minimization above, we define ζ_k to be one of the solutions to the corresponding minimization problem. To ensure the existence of ζ_k ’s, we impose that P is bounded and lower semi-continuous with respect to the norm-convergence, in the sense that

$$P(u) \leq C\|u\|,$$

$$\|u_n - u\| \rightarrow 0 \implies P(u) \leq \liminf P(u_n). \tag{2.2}$$

The smallest constant C satisfying the boundedness of P is the functional norm of P and is denoted by $\|P\|$.

We require that the eigenvectors of T , i.e. $\{\phi_n\}_{n \in \mathbb{N}}$, form a complete orthonormal system in \mathcal{H} . We conjecture that as long as the perturbation satisfies the existence criteria (2.2), such a spectral result still holds:

CONJECTURE 2.2. *The set $\{\zeta_n\}_{n \in \mathbb{N}}$ obtained via the variational procedure (2.1) forms a complete orthonormal system in \mathcal{H} .*

This section mainly focuses on verifying Conjecture 2.2 under certain growth assumptions on the eigenvalues λ_n . In order to verify this conjecture, one needs to show that

$$\phi_k \in \overline{\text{span}\{\zeta_n\}_{n \in \mathbb{N}}} \quad \forall k \in \mathbb{N}, \tag{2.3}$$

where $\text{span } E$ denotes the space consisting of the finite linear combinations of the elements in E . The following Hilbert theoretic result quantifies the relation (2.3).

LEMMA 2.3. *Let $\{e_n\}_{n \in \mathbb{N}}$ be a maximal orthonormal system in a Hilbert space \mathcal{H} . Let $\{f_n\}_{n \in \mathbb{N}}$ be some orthonormal system in \mathcal{H} . Assume that each f_n has the expansion*

$$f_n = \sum_{k \in \mathbb{N}} a_{n,k} e_k, \quad a_{n,k} \in \mathbb{C}.$$

Then, for each $k \in \mathbb{N}$,

$$d(e_k, \text{span}\{f_n\})^2 = 1 - \sum_{n \in \mathbb{N}} |a_{n,k}|^2,$$

where $d(e, M)$ denotes the distance between some $e \in \mathcal{H}$ and some linear subspace M of \mathcal{H} .

Proof. Let w be the projection of e_k onto $\text{span}\{f_n\}_{n \in \mathbb{N}}$. Then, since $\{f_n\}_{n \in \mathbb{N}}$ is an orthonormal system, w is given by

$$\begin{aligned} w &= \sum_{n \in \mathbb{N}} \langle e_k, f_n \rangle f_n \\ &= \sum_{n \in \mathbb{N}} \langle e_k, \sum_j a_{n,j} e_j \rangle f_n \\ &= \sum_{n \in \mathbb{N}} \overline{a_{n,k}} f_n. \end{aligned}$$

Hence, we can compute the size of w :

$$\|w\|^2 = \sum_{n \in \mathbb{N}} |a_{n,k}|^2. \tag{2.4}$$

Note by the property of the projection that $e_k - w \perp w$. Therefore, by the Pythagorean identity, we have

$$\|e_k\|^2 = \|e_k - w\|^2 + \|w\|^2. \tag{2.5}$$

Notice also that w , being the projection of e_k onto $\text{span}\{f_n\}_{n \in \mathbb{N}}$, is the closest point to e_k inside $\text{span}\{f_n\}_{n \in \mathbb{N}}$, so that

$$d(e_k, \text{span}\{f_n\})^2 = \|e_k - w\|^2. \tag{2.6}$$

Combining (2.4)–(2.6), we get

$$d(e_k, \text{span}\{f_n\})^2 = \|e_k\|^2 - \|w\|^2 = 1 - \sum_{n \in \mathbb{N}} |a_{n,k}|^2,$$

as desired. □

COROLLARY 2.4. *Let $\{e_n\}_{n \in \mathbb{N}}$, $\{f_n\}_{n \in \mathbb{N}}$, $a_{n,k}$ be as in Lemma 2.3. Then, for each $k \in \mathbb{N}$,*

$$\sum_{n \in \mathbb{N}} |a_{n,k}|^2 \leq 1 \tag{2.7}$$

and

$$e_k \in \overline{\text{span}\{f_n\}} \iff \sum_{n \in \mathbb{N}} |a_{n,k}|^2 = 1.$$

The following lemma yields an estimate for the elements that are lying inside the orthogonal complement of any arbitrary orthonormal system, in terms of the deviation of their functional values from the sum of the eigenvalues corresponding to the true eigenstates. We denote the deviation of the first N elements by $F(N)$.

LEMMA 2.5. *Let ϕ_n be the eigenvectors of the operator T , with the corresponding eigenvalues $\{\lambda_n\}_{n \in \mathbb{N}}$ being in increasing order. Let $\{e_n\}_{n \in \mathbb{N}}$ be an orthonormal system satisfying*

$$\sum_{n=1}^N \langle Te_n, e_n \rangle \leq F(N) + \sum_{n=1}^N \lambda_n \quad \forall N \in \mathbb{N}. \tag{2.8}$$

Suppose that there exists $f \in \{e_n\}_{n \in \mathbb{N}}^\perp$, $\|f\| = 1$, with the expansion

$$f = \sum_{n \in \mathbb{N}} f_n \phi_n, \quad f_n \in \mathbb{C}.$$

Then, we have

$$\sum_{n=1}^N |f_n|^2 (\lambda_{N+1} - \lambda_n) \leq F(N), \quad \forall N \in \mathbb{N}. \tag{2.9}$$

Proof. Let the $\{a_{n,k}\}_{n,k \in \mathbb{N}}$ denote the coefficients with e_n expanded in the basis $\{\phi_k\}_{k \in \mathbb{N}}$, i.e.

$$e_n = \sum_{k \in \mathbb{N}} a_{n,k} \phi_k.$$

Applying the result (2.7) of Lemma 2.3 to the orthonormal systems $\{e_n\}_{n \in \mathbb{N}} \cup \{f\}$ and $\{\phi_n\}_{n \in \mathbb{N}}$, for each $k \in \mathbb{N}$, we get

$$\sum_{n \in \mathbb{N}} |a_{n,k}|^2 \leq 1 - |f_k|^2. \tag{2.10}$$

By the bilinearity of the inner product,

$$\langle Te_n, e_n \rangle = \sum_{k=1}^{\infty} |a_{n,k}|^2 \lambda_k.$$

Hence,

$$\sum_{n=1}^N \langle Te_n, e_n \rangle = \sum_{k=1}^{\infty} \left(\sum_{n=1}^N |a_{n,k}|^2 \right) \lambda_k. \tag{2.11}$$

Since the e_n 's have norm 1, in the expression (2.11), the sum of the coefficients of λ_k over k equals N . Given that the λ_k 's are in the increasing order, the expression (2.11) is minimized when the coefficients of λ_k are maximized for small k . Having the constraint (2.10), we get

$$\sum_{n=1}^N \langle Te_n, e_n \rangle = \sum_{k=1}^{\infty} \left(\sum_{n=1}^N |a_{n,k}|^2 \right) \lambda_k \geq \sum_{k=1}^N (1 - |f_k|^2) \lambda_k + \lambda_{N+1} \sum_{k=1}^N |f_k|^2.$$

Combining this with the inequality (2.8), we get

$$\sum_{n=1}^N \lambda_n + F(N) \geq \sum_{n=1}^N \langle Te_n, e_n \rangle \geq \sum_{n=1}^N (1 - |f_n|^2) \lambda_n + \lambda_{N+1} \sum_{n=1}^N |f_n|^2,$$

which implies

$$\sum_{n=1}^N |f_n|^2 (\lambda_{N+1} - \lambda_n) \leq F(N),$$

as desired. □

Lemma 2.5 will be essential for proving the completeness result. The estimate (2.9) provides us an understanding of the elements lying inside the orthogonal complement in terms of $F(N)$ and the eigenvalues of T . If the estimate (2.9) is incompatible with the growth of eigenvalues, then we deduce that the orthogonal complement is empty and, hence, the orthonormal system is maximal.

The next lemma provides an estimate for the deviation of the functional values of ζ_n from the eigenvalues corresponding to the true eigenstates, so that Lemma 2.5 is applicable.

LEMMA 2.6. *Let $\{\zeta_n\}_{n \in \mathbb{N}}$ be the solutions to the variational procedure (2.1). Let $\{\lambda_n\}_{n \in \mathbb{N}}$ be the eigenvalues of T , in increasing order. Then,*

$$J[\zeta_n] \leq \lambda_n + \|P\|.$$

In particular, we have

$$\sum_{n=1}^N \langle T\zeta_n, \zeta_n \rangle \leq \sum_{n=1}^N J[\zeta_n] \leq \sum_{n=1}^N \lambda_n + \|P\|N. \tag{2.12}$$

Proof. Let $\{a_{n,k}\}_{n,k \in \mathbb{N}}$ denote the coefficients with ζ_n expanded into the basis $\{\phi_k\}_{k \in \mathbb{N}}$:

$$\zeta_n = \sum_{k \in \mathbb{N}} a_{n,k} \phi_k.$$

Let $n \in \mathbb{N}$ be fixed. For integers j with $1 \leq j \leq n - 1$, define

$$\eta_j = \sum_{k=1}^n a_{j,k} \phi_k.$$

Clearly, $\{\eta_1, \eta_2, \dots, \eta_{n-1}\} \subset \text{span}\{\phi_1, \phi_2, \dots, \phi_n\}$. Furthermore,

$$\dim \text{span}\{\phi_1, \phi_2, \dots, \phi_n\} = n,$$

i.e. the space $\text{span}\{\phi_1, \phi_2, \dots, \phi_n\}$ has dimension larger than the cardinality of $\{\eta_1, \eta_2, \dots, \eta_{n-1}\}$, so that we can find $\eta \in \text{span}\{\phi_1, \phi_2, \dots, \phi_n\}$, $\eta \neq 0$, such that

$$\eta \perp \eta_j, \quad \forall j : 1 \leq j \leq n - 1.$$

Now, since η_j 's and η lie inside $\text{span}\{\phi_1, \phi_2, \dots, \phi_n\}$, we get

$$\langle \eta, \zeta_j \rangle = \langle \eta, \eta_j \rangle = 0, \quad \forall j : 1 \leq j \leq n - 1,$$

so that

$$\eta \perp \zeta_j, \quad \forall j : 1 \leq j \leq n - 1.$$

By rescaling, we may assume $\|\eta\| = 1$, so that η lies precisely in the class of functions where we look for a minimizer to determine ζ_n . The function η is in the solution set of the variational problem (2.1) at the n th step. Hence,

$$J[\zeta_n] \leq J[\eta] = \langle T\eta, \eta \rangle + P(\eta). \tag{2.13}$$

Note first by the boundedness of P that

$$P(\eta) \leq \|P\| \|\eta\| = \|P\|. \tag{2.14}$$

Suppose η has the expansion

$$\eta = \sum_{k=1}^n b_k \phi_k.$$

Since, $\|\eta\| = 1$, we have $\sum_{k=1}^n |b_k|^2 = 1$. Furthermore, by the bilinearity of the inner product,

$$\langle T\eta, \eta \rangle = \sum_{k=1}^n |b_k|^2 \lambda_k.$$

We also have that λ_k 's are in increasing order, so that

$$\langle T\eta, \eta \rangle = \sum_{k=1}^n |b_k|^2 \lambda_k \leq \lambda_n \sum_{k=1}^n |b_k|^2 = \lambda_n. \tag{2.15}$$

Combining (2.13)–(2.15), we obtain

$$J[\zeta_n] \leq \lambda_n + \|P\|. \tag{2.16}$$

Summing up the inequality (2.16) for $n = 1, 2, \dots, N$ and combining with the non-negativity of P , we verify (2.12). \square

The following theorem provides the completeness of the orthonormal system $\{\zeta_n\}_{n \in \mathbb{N}}$, provided the eigenvalues satisfy the super-linear growth.

THEOREM 2.7. *Suppose the eigenvalues of T satisfy*

$$\lim_{n \rightarrow \infty} \frac{\lambda_n}{n} = \infty. \tag{2.17}$$

Then, $\{\zeta_n\}_{n \in \mathbb{N}}$, which is defined by the variational procedure (2.1), forms a complete orthonormal system in \mathcal{H} .

Proof. Lemma 2.6 implies

$$\sum_{n=1}^N \langle T\zeta_n, \zeta_n \rangle \leq \sum_{n=1}^N \lambda_n + \|P\|N,$$

so that Lemma 2.5 is applicable to the orthonormal system $\{\zeta_n\}_{n \in \mathbb{N}}$ with the function $F(N) = \|P\|N$. That is, assuming the existence of an $f \in \{\zeta_n\}_{n \in \mathbb{N}}^\perp$, $\|f\| = 1$ with the expansion

$$f = \sum f_n \phi_n, \quad f_n \in \mathbb{C},$$

we obtain the estimate

$$\sum_{n=1}^N |f_n|^2 (\lambda_{N+1} - \lambda_n) \leq \|P\|N, \quad \forall N \in \mathbb{N}. \tag{2.18}$$

This last inequality implies (assuming $f_n \neq 0$)

$$\lambda_{N+1} - \lambda_n \leq \frac{\|P\|N}{|f_n|^2}, \quad \forall N \in \mathbb{N},$$

which yields a contradiction by violating the growth condition (2.17) on λ_{N+1} as we pass to limit as $N \rightarrow \infty$. Therefore, there is no non-zero function $f \in \{\zeta_n\}_{n \in \mathbb{N}}^\perp$, so that the orthonormal system $\{\zeta_n\}_{n \in \mathbb{N}}$ is complete, as desired. \square

By trading the magnitude of the penalty term with the growth of λ_n , we can generalize the Theorem 2.7 so that it holds under a weaker growth condition on λ_n .

THEOREM 2.8. *Suppose that the eigenvalues λ_n grow linearly in the sense that they satisfy*

$$\lambda_n = Mn + o(n), \quad \text{as } n \rightarrow \infty \tag{2.19}$$

for some constant M . Suppose also that the penalty term P satisfies the following bound:

$$\|P\| < M. \tag{2.20}$$

Then, $\{\zeta_n\}_{n \in \mathbb{N}}$ forms a complete orthonormal system in \mathcal{H} .

Proof. Proceeding similarly to the proof of Theorem 2.7, we get the inequality (2.18), namely that if f is a function with unit norm, lying in the orthogonal complement of $\{\zeta_n\}_{n \in \mathbb{N}}$, then we have

$$\sum_{n=1}^N |f_n|^2 (\lambda_{N+1} - \lambda_n) \leq \|P\|N, \quad \forall N \in \mathbb{N}. \tag{2.21}$$

Now, for $m < N$, by the monotonicity of λ_k , we can lower-bound the LHS of (2.21) by

$$(\lambda_{N+1} - \lambda_m) \sum_{n=1}^m |f_n|^2,$$

so that

$$\frac{\lambda_{N+1} - \lambda_m}{N} \sum_{n=1}^m |f_n|^2 \leq \|P\|, \quad \forall N \in \mathbb{N}, \forall m : 0 < m < N. \tag{2.22}$$

Taking the limit in (2.22) as $N \rightarrow \infty$, with the aid of the growth condition (2.19), we obtain

$$M \sum_{n=1}^m |f_n|^2 \leq \|P\|, \quad \forall m \in \mathbb{N}.$$

Now, taking the limit as $m \rightarrow \infty$ and keeping in mind that $\|f\| = 1$, we obtain

$$M \leq \|P\|,$$

contradicting (2.20). □

The following theorem establishes how close the functions ζ_n approximate the subspaces generated by the first few eigenvectors of the operator T .

THEOREM 2.9. *Let V_m be the subspace generated by the functions $\{\zeta_1, \zeta_2, \dots, \zeta_m\}$. Then, for any $n \leq m$, we have*

$$\sum_{k=1}^n d(\phi_k, V_m)^2 \leq \frac{m\|P\|}{\lambda_{m+1} - \lambda_n},$$

provided $\lambda_{m+1} \neq \lambda_n$.

Proof. Recall from Lemma 2.6 that

$$\sum_{j=1}^m \langle T\zeta_j, \zeta_j \rangle \leq \sum_{j=1}^m J[\zeta_j] \leq m\|P\| + \sum_{j=1}^m \lambda_j. \tag{2.23}$$

Recall by the bilinearity of the inner product that

$$\langle T\zeta_j, \zeta_j \rangle = \sum_{k=1}^{\infty} |a_{j,k}|^2 \lambda_k. \tag{2.24}$$

Combining (2.24) and (2.23) yields that

$$\sum_{j=1}^m \sum_{k=1}^{\infty} |a_{j,k}|^2 \lambda_k \leq m \|P\| + \sum_{j=1}^m \lambda_j. \tag{2.25}$$

Rearranging (2.25), we obtain

$$\sum_{k=m+1}^{\infty} \sum_{j=1}^m |a_{j,k}|^2 \lambda_k - \sum_{k=1}^m \left(1 - \sum_{j=1}^m |a_{j,k}|^2 \right) \lambda_k \leq m \|P\|.$$

Bounding the terms λ_k from below with $k > m$ by λ_{m+1} in the last expression, we obtain

$$\lambda_{m+1} \sum_{k=m+1}^{\infty} \sum_{j=1}^m |a_{j,k}|^2 - \sum_{k=1}^m \left(1 - \sum_{j=1}^m |a_{j,k}|^2 \right) \lambda_k \leq m \|P\|. \tag{2.26}$$

Since $\sum_{k=1}^{\infty} |a_{j,k}|^2 = 1$ for $j = 1, 2, \dots, m$, we conclude that

$$\sum_{k=m+1}^{\infty} \sum_{j=1}^m |a_{j,k}|^2 = \sum_{k=1}^m \left(1 - \sum_{j=1}^m |a_{j,k}|^2 \right).$$

Substituting this into the inequality (2.26) and rearranging further, we obtain

$$\sum_{k=1}^m \left(1 - \sum_{j=1}^m |a_{j,k}|^2 \right) (\lambda_{m+1} - \lambda_k) \leq m \|P\|. \tag{2.27}$$

Notice by Lemma 2.3 that the coefficient in front of $\lambda_{m+1} - \lambda_k$ in (2.27) is equal to $d(\phi_k, V_m)^2$, so that (2.27) becomes

$$\sum_{k=1}^m d(\phi_k, V_m)^2 (\lambda_{m+1} - \lambda_k) \leq m \|P\|.$$

Exploiting the monotonicity of λ_k once more, we obtain

$$\sum_{k=1}^n d(\phi_k, V_m)^2 (\lambda_{m+1} - \lambda_n) \leq \sum_{k=1}^m d(\phi_k, V_m)^2 (\lambda_{m+1} - \lambda_k) \leq m \|P\|,$$

implying

$$\sum_{k=1}^n d(\phi_k, V_m)^2 \leq \frac{m \|P\|}{\lambda_{m+1} - \lambda_n},$$

as desired. □

3. Perturbed variational problems associated to elliptic operators

The results of Section 2 can now be applied to second-order linear symmetric elliptic operators. Let L be a second-order linear symmetric elliptic operator defined on a

bounded domain Ω in \mathbb{R}^d . For simplicity, we will consider symmetric elliptic operators with principal parts $-\Delta$, i.e. the Schrödinger’s operator given by

$$Lu = -\Delta u + cu,$$

where $c : \Omega \rightarrow \mathbb{R}$ is a bounded measurable function.

As noted in the beginning of Section 2, according to the spectral theorem for second-order linear symmetric elliptic operators, L satisfies the following properties:

- (1) L has real (Dirichlet) eigenvalues, λ_k , with $\{\lambda_k\}_{k=1}^\infty$ in increasing order, and $\lambda_k \rightarrow +\infty$ as $k \rightarrow \infty$ and
- (2) the (normalized) eigenfunctions $\{\phi_k\}_{k=1}^\infty$, with $L\phi_k = \lambda_k\phi_k$, form a complete orthonormal system in $L^2(\Omega)$.

Furthermore, the Courant–Fisher principle applies to L , so that the eigenvalues and eigenfunctions of L is found by the following variational formulae:

$$\begin{aligned} \lambda_1 &= \min_{\substack{u \in H_0^1(\Omega) \\ \|u\|_2=1}} B[u, u], \\ \phi_1 &= \operatorname{argmin}_{\substack{u \in H_0^1(\Omega) \\ \|u\|_2=1}} B[u, u], \\ \lambda_k &= \min_{\substack{u \in H_0^1(\Omega) \\ u \in \{\phi_1, \dots, \phi_{k-1}\}^\perp \\ \|u\|_2=1}} B[u, u], \\ \phi_k &= \operatorname{argmin}_{\substack{u \in H_0^1(\Omega) \\ u \in \{\phi_1, \dots, \phi_{k-1}\}^\perp \\ \|u\|_2=1}} B[u, u], \end{aligned}$$

where

$$B[u, v] = \langle Lu, v \rangle = \int_\Omega \nabla u \cdot \overline{\nabla v} + cu\overline{v} \, dx$$

is the bilinear form associated to L .

We proceed similarly to as in Section 2, where we perturb the functional $B[u, u]$. This time, we restrict ourselves to the penalty terms given by a constant multiple of the L^1 norm. Namely, we consider the energy functional

$$J_\mu[u] = B[u, u] + \frac{1}{\mu} \|u\|_{L^1}$$

and analogously define the functions $\{\psi_k\}_{k \in \mathbb{N}}$ via

$$\begin{aligned} \psi_1 &= \operatorname{argmin}_{\substack{u \in H_0^1(\Omega) \\ \|u\|_2=1}} J_\mu[u], \\ \psi_k &= \operatorname{argmin}_{\substack{u \in H_0^1(\Omega) \\ u \in \{\psi_1, \dots, \psi_{k-1}\}^\perp \\ \|u\|_2=1}} J_\mu[u]. \end{aligned} \tag{3.1}$$

We call these functions $\{\psi_k\}_{k \in \mathbb{N}}$ “compressed modes of second type” (CM-II) by analogy to the “compressed modes” (CM) defined in [9]. The main difference between

CM and CM-II is that CMs are defined as the minimizers of a joint energy sum under orthogonality constraints, whereas CM-IIs marginally minimize the energy functional under a progressive orthogonality constraint. Nevertheless, CM-IIs $\{\psi_1, \psi_2, \dots, \psi_m\}$, being an orthonormal sequence, are in the solution set of the associated variational problem for CMs, so that the theory for CM-IIs can be used to prove the analogous results for CMs.

The following lemma establishes the existence of $\{\psi_k\}_{k \in \mathbb{N}}$ by verifying the existence criteria (2.2) for the L^1 penalty term in the definition of J_μ .

LEMMA 3.1. *Let $P : L^2(\Omega) \rightarrow \mathbb{R}$ be defined by $P(u) = \frac{1}{\mu} \|u\|_{L^1}$. Then, P satisfies the criteria in (2.2). Furthermore,*

$$\|P\| = \frac{|\Omega|^{\frac{1}{2}}}{\mu}.$$

Proof. By the Cauchy–Schwarz inequality,

$$\frac{1}{\mu} \|u\|_{L^1} \leq \frac{1}{\mu} \|u\|_{L^2} \|\mathbf{1}_\Omega\|_{L^2} = \frac{|\Omega|^{\frac{1}{2}}}{\mu} \|u\|_{L^2}, \tag{3.2}$$

with equality when u is a non-zero constant function. Therefore, P is bounded with functional norm $\frac{|\Omega|^{\frac{1}{2}}}{\mu}$.

As for the lower semi-continuity, consider a sequence $u_n \in L^2(\Omega)$ converging to some $u \in L^2(\Omega)$ in L^2 . The inequality (3.2) implies that u_n converges to u also in L^1 , so that $P(u) = \lim P(u_n)$, as desired. \square

Now, theorems 2.7, 2.8, and 2.9, can be derived for $\{\psi_k\}_{k \in \mathbb{N}}$. Theorems 2.7 and 2.8 hold when the eigenvalues grow super-linearly and linearly, respectively. Weyl’s law yields the exact asymptotic behavior of the eigenvalues of a second-order linear symmetric elliptic operator:

THEOREM 3.2 (Weyl’s law (see e.g. [8])). *Let L be a second-order linear elliptic operator on a bounded domain $\Omega \subset \mathbb{R}^d$ of the form*

$$Lu = -\Delta u + cu,$$

where $c : \Omega \rightarrow \mathbb{R}$ is a bounded measurable function. Let $\{\lambda_n\}_{n \in \mathbb{N}}$ be the eigenvalues of L , arranged in increasing order. Then,

$$\lambda_n = \frac{(2\pi)^d}{\omega_d |\Omega|} n^{2/d} + o(n^{2/d}), \text{ as } n \rightarrow \infty,$$

where ω_d denotes the volume of the unit ball in \mathbb{R}^d .

Therefore, we can deduce from Weyl’s law that super-linear and linear growth conditions on eigenvalues holds true precisely in dimensions 1 and 2, so that we have the following corollaries as direct consequences of theorems 2.7, 2.8, and 2.9:

COROLLARY 3.3 (Corollary to Theorem 2.7). *Let $L = -\Delta + c(x)$ be defined on a bounded open interval in \mathbb{R} , where c is a bounded measurable function. Then, the associated compressed modes of second type (CM-II), $\{\psi_n\}_{n \in \mathbb{N}}$, which is defined by the variational procedure (3.1), forms a complete orthonormal system in $L^2(\Omega)$.*

COROLLARY 3.4 (Corollary to Theorem 2.8). *Let $L = -\Delta + c(x)$ be defined on a bounded open interval in \mathbb{R} , where c is a bounded measurable function. Suppose also that the penalty parameter μ satisfies the following bound:*

$$\mu > \frac{1}{4\pi|\Omega|^{\frac{3}{2}}}.$$

Then, the associated compressed modes of second type (CM-II), $\{\psi_n\}_{n \in \mathbb{N}}$, which is defined by the variational procedure (3.1), forms a complete orthonormal system in $L^2(\Omega)$.

COROLLARY 3.5 (Corollary to Theorem 2.9). *Let $\{\phi_k\}_{k \in \mathbb{N}}$ be the (Dirichlet) eigenfunctions of the operator $L = -\Delta + c(x)$, defined on a bounded domain $\Omega \subset \mathbb{R}^n$. Let $\{\lambda_k\}_{k \in \mathbb{N}}$ be the associated eigenvalues. Let $\{\psi_n\}_{n \in \mathbb{N}}$ be the functions defined by the variational procedure (3.1), and let V_m be the subspace generated by the functions $\{\psi_1, \psi_2, \dots, \psi_m\}$. Then, for any $n \leq m$, we have*

$$\sum_{k=1}^n d(\phi_k, V_m)^2 \leq \frac{m|\Omega|^{\frac{1}{2}}}{\mu(\lambda_{m+1} - \lambda_n)}, \tag{3.3}$$

provided that $\lambda_{m+1} \neq \lambda_n$.

REMARK 3.1. Corollaries 3.3–3.5 holds true (with modified inequalities) for second-order linear symmetric elliptic operators with appropriate coercivity properties, so that the eigenvalues grow at the same order as $-\Delta$.

The following theorem establishes that the elements that lie in the orthogonal complement of $\{\psi_n\}_{n \in \mathbb{N}}$ cannot lie inside the space $H_0^1(\Omega)$. In other words, the orthogonal complement consists of highly irregular functions.

THEOREM 3.6. *Let $\Omega \subset \mathbb{R}^d$ and $\{\psi_n\}_{n \in \mathbb{N}}$ be the solutions to the variational procedure (2.1). Then,*

$$\{\psi_n\}_{n \in \mathbb{N}}^\perp \cap H_0^1(\Omega) = \{0\}.$$

Proof. Assume to the contrary that there is a non-zero $f \in \{\psi_n\}^\perp \cap H_0^1(\Omega)$. We may normalize f such that $\|f\|_2 = 1$. Since $f \in \{\psi_1, \dots, \psi_{n-1}\}^\perp$ for all n , f is in the class of functions where we look for a minimizer to obtain ψ_n , and hence, is in the solution set of the variational problem (2.1) at n th step. As ψ_n is the actual solution to the corresponding minimization problem, we have

$$J_\mu[\psi_n] \leq J_\mu[f]. \tag{3.4}$$

We now prove that

$$\lim_{n \rightarrow \infty} J_\mu[\psi_n] = \infty, \tag{3.5}$$

which together with (3.4) implies

$$J_\mu[f] = \infty. \tag{3.6}$$

Recall that

$$\sum_{n=1}^N B[\psi_n, \psi_n] = \sum_{k=1}^\infty \left(\sum_{n=1}^N |a_{n,k}|^2 \right) \lambda_k, \tag{3.7}$$

where the coefficients in front of λ_k have magnitude less than or equal to 1 for each k and have their sum over k is N . Since λ_k 's are in increasing order, the expression (3.7) is minimized when the coefficients of λ_k are maximized for small k . Therefore,

$$\sum_{n=1}^N B[\psi_n, \psi_n] \geq \sum_{k=1}^N \lambda_k,$$

and, since P is non-negative,

$$\sum_{n=1}^N J_\mu[\psi_n] = \sum_{n=1}^N B[\psi_n, \psi_n] + P(\psi_n) \geq \sum_{n=1}^N B[\psi_n, \psi_n] \geq \sum_{k=1}^N \lambda_k. \tag{3.8}$$

We know that $\lim_{n \rightarrow \infty} \lambda_n = \infty$ and that both λ_n 's and $J_\mu[\psi_n]$'s are in increasing order. Therefore, the inequality (3.8) can hold only if (3.5) holds. Hence, we verify (3.6), i.e.

$$J_\mu[f] = B[f, f] + P(f) = \infty. \tag{3.9}$$

Now, since P is bounded, the expression (3.9) yields

$$B[f, f] = \infty. \tag{3.10}$$

On the other hand, as B is the bilinear form associated to a second-order elliptic operator, it is bounded in the sense that

$$|B[u, v]| \leq C \|u\|_{H^1(\Omega)} \|v\|_{H^1(\Omega)}. \tag{3.11}$$

Combining (3.10) and (3.11) applied to $u = v = f$, we get

$$\|f\|_{H^1(\Omega)} = \infty,$$

i.e. $f \notin H^1(\Omega)$, contradicting the assumption that $f \in \{\psi_n\}^\perp \cap H_0^1(\Omega)$. □

3.1. Scaling properties. In this subsection, our aim is to provide the scaling invariance between μ and Ω . Let $\kappa\Omega$ denote the usual scaling of the domain Ω by a positive real number κ . Let \tilde{L} on $\kappa\Omega$ be given by

$$\tilde{L}u = -\Delta u + \tilde{c}u = -\Delta u + \kappa^{-2}c\left(\frac{x}{\kappa}\right)u.$$

If the operator L on Ω has orthonormal eigenpairs $\{(\phi_n(x), \lambda_n)\}_{n=1}^\infty$, then

$$\{(\kappa^{-d/2}\phi_n\left(\frac{x}{\kappa}\right), \kappa^{-2}\lambda_n)\}_{n=1}^\infty$$

forms the orthonormal set of eigenpairs for \tilde{L} . Let $\{\psi_1, \psi_2, \dots\}$ be the compressed modes of second type (CM-II), corresponding to the operator L . For each $n \in \mathbb{N}$, define $\tilde{\psi}_n(x) = \kappa^{-d/2}\psi_n\left(\frac{x}{\kappa}\right)$. Then, clearly $\{\tilde{\psi}_1, \tilde{\psi}_2, \dots\}$ forms an orthonormal system. Moreover,

$$\begin{aligned} \tilde{B}[\tilde{\psi}_n, \tilde{\psi}_n] &= \langle \tilde{L}\tilde{\psi}_n, \tilde{\psi}_n \rangle = \int_{\kappa\Omega} |\nabla \tilde{\psi}_n|^2 dx + \int_{\kappa\Omega} \tilde{c}\tilde{\psi}_n^2 dx \\ &= \kappa^{-2} \int_{\Omega} |\nabla \psi_n|^2 dx + \kappa^{-2} \int_{\Omega} c\psi_n^2 dx \end{aligned}$$

$$= \kappa^{-2} B[\psi_n, \psi_n]$$

and

$$\|\tilde{\psi}_n\|_1 = \int_{\kappa\Omega} |\tilde{\psi}_n| dx = \kappa^{d/2} \int_{\kappa\Omega} |\psi_n| dx = \kappa^{d/2} \|\psi_n\|_1.$$

Hence, $\{\tilde{\psi}_1, \tilde{\psi}_2, \dots\}$ is the corresponding CM-II for the operator \tilde{L} on $\kappa\Omega$, with respect to the energy functional

$$\tilde{J}_{\tilde{\mu}}[u] = \tilde{B}[u, u] + \frac{1}{\tilde{\mu}} \|u\|_{L^1},$$

where

$$\tilde{\mu} = \frac{\mu}{\kappa^{2+d/2}}.$$

Notice that this last scaling relation is consistent with the scaling properties of the inequalities in Corollary 3.4 and Corollary 3.5.

4. Applications

We now establish the analogues of theorems 2.7–2.9 for the compressed modes and the compressed plane waves. We first provide the precise definitions of Compressed Modes (CMs) and Compressed Plane Waves (CPWs) as introduced in [9, 10], and establish their connections to the theory we developed in Section 2. Then we proceed with the verification of the analogous theorems.

4.1. Compressed modes. CMs are defined via the following minimization procedure:

$$\Psi^{(m)} = \{\psi_1^{(m)}, \dots, \psi_m^{(m)}\} = \underset{h_1, h_2, \dots, h_m}{\operatorname{argmin}} \sum_{i=1}^m J_\mu[h_i] \quad \text{s.t.} \quad \langle h_j, h_k \rangle = \delta_{jk}, \quad (4.1)$$

where

$$J_\mu[u] = \frac{1}{\mu} \|u\|_{L^1} + \langle u, \left(-\frac{1}{2}\Delta + V\right) u \rangle = \frac{1}{\mu} \|u\|_{L^1} + \frac{1}{2} \|\nabla u\|_{L^2}^2 + \int_\Omega V u^2 dx, \quad (4.2)$$

where V is a bounded measurable real-valued function defined on Ω . Here, the quantity $\langle u, \left(-\frac{1}{2}\Delta + V\right) u \rangle$ corresponds to the bilinear form associated to the elliptic operator $-\frac{1}{2}\Delta + V$. We denote the eigenvalues and eigenfunctions of $-\frac{1}{2}\Delta + V$ by λ_n and ϕ_n , with the λ_n 's in increasing order, as usual.

As noted earlier, CMs of second type $\{\psi_1, \psi_2, \dots, \psi_m\}$ defined by the variational procedure (3.1), being an orthonormal sequence, are in the solution set of the minimization problem (4.1), so that

$$\sum_{i=1}^m J_\mu[\psi_i^{(m)}] \leq \sum_{i=1}^m J_\mu[\psi_i].$$

Combining this with the estimate (2.12), we obtain

$$\sum_{i=1}^m J_\mu[\psi_i^{(m)}] \leq m \frac{|\Omega|^{\frac{1}{2}}}{\mu} + \sum_{j=1}^m \lambda_j. \quad (4.3)$$

The proof of Theorem 2.9 relies essentially on the estimation (2.23) and the orthonormality of the sequence $\{\psi_1, \dots, \psi_m\}$. We still have the orthonormality and the estimation (4.3) analogous to (2.23). Hence, the following corollary holds.

COROLLARY 4.1 (Corollary to Theorem 2.9). *Let V_m be the subspace generated by the compressed modes $\Psi = \{\psi_1^{(m)}, \psi_2^{(m)}, \dots, \psi_m^{(m)}\}$. Then, for any $n \leq m$, we have*

$$\sum_{k=1}^n d(\phi_k, V_m)^2 \leq \frac{m|\Omega|^{\frac{1}{2}}}{\mu(\lambda_{m+1} - \lambda_n)},$$

provided that $\lambda_{m+1} \neq \lambda_n$.

From Corollary 4.1, we deduce the following approximation result:

COROLLARY 4.2. *Let $\{\phi_k\}_{k \in \mathbb{N}}$ be the (Dirichlet) eigenfunctions of a second-order linear symmetric elliptic operator L , defined on a bounded domain $\Omega \subset \mathbb{R}$. Given any fixed parameter μ , the first m compressed modes up to a linear transformation, denoted by $\{\xi_1^{(m)}, \dots, \xi_m^{(m)}\}$, satisfy*

$$\lim_{m \rightarrow \infty} \|\phi_k - \xi_k^{(m)}\|_2 = 0, \quad \forall k \in \mathbb{N}.$$

Proof. Let $\xi_k^{(m)}$ denote the projection of $\phi_k^{(m)}$ onto the linear subspace spanned by $\{\psi_1^{(m)}, \dots, \psi_m^{(m)}\}$, which we denote by V_m . Then, clearly, $\xi_k^{(m)}$ is a linear combination of $\{\psi_1^{(m)}, \dots, \psi_m^{(m)}\}$. Furthermore, as a property of the projection, we have

$$d(\phi_k, V_m) = \|\phi_k^{(m)} - \xi_k^{(m)}\|_2,$$

so that Corollary 4.1 implies

$$\sum_{k=1}^n \|\phi_k - \xi_k^{(m)}\|_2^2 \leq \frac{m|\Omega|^{\frac{1}{2}}}{\mu(\lambda_{m+1} - \lambda_n)}. \tag{4.4}$$

As Ω is a bounded domain inside \mathbb{R} , By Weyl’s law, we know that λ_m grows quadratically in m . Hence, passing to the limit in (4.4) as $m \rightarrow \infty$, we conclude that the summands in the LHS of (4.4) decay to zero, i.e.

$$\lim_{m \rightarrow \infty} \|\phi_k - \xi_k^{(m)}\|_2 = 0, \tag{4.5}$$

as desired. □

Corollary 4.2 can be viewed as a completeness result, since (4.5) yields that any eigenfunction ϕ_k is well approximated by its projection $\xi_k^{(m)}$ onto V_m . In a sense, V_m ’s trace the full space, as $m \rightarrow \infty$.

4.2. Compressed plane waves. The construction of CPWs is closely related to that of compressed modes, where both involve minimizing a certain functional. The difference is that compressed plane waves have multi-resolution capabilities, which is achieved by adding the shift-orthogonality constraints.

Let $\mathbf{w} = (w_1, \dots, w_d) \in \mathbb{R}_+^d$ be a basis of a d -dimensional lattice, and let Ω be a rectangular box with

$$\Omega = [0, n_1 w_1] \times \dots \times [0, n_d w_d], \quad (n_1, \dots, n_d) \in \mathbb{N}^d.$$

Define the lattice

$$\Gamma_{\mathbf{w}} = \{\mathbf{jw} := (j_1w_1, \dots, j_dw_d) \mid 0 \leq j_1 < n_1, \dots, 0 \leq j_d < n_d\}.$$

The first n basic compressed plane waves (BCPWs), $\{\psi^k\}_{k=1}^n$, are defined via

$$\begin{aligned} \psi^1 &= \underset{\psi}{\operatorname{argmin}} J_{\mu}[\psi] \text{ s.t. } \langle \psi(\mathbf{x}), \psi(\mathbf{x} - \mathbf{jw}) \rangle = \delta_{\mathbf{j}, \mathbf{0}} \quad \forall \mathbf{jw} \in \Gamma_{\mathbf{w}}; \\ \psi^k &= \underset{\psi}{\operatorname{argmin}} J_{\mu}[\psi] \text{ s.t. } \begin{cases} \langle \psi(\mathbf{x}), \psi(\mathbf{x} - \mathbf{jw}) \rangle = \delta_{\mathbf{j}, \mathbf{0}} \quad \forall \mathbf{jw} \in \Gamma_{\mathbf{w}} \\ \langle \psi(\mathbf{x}), \psi^i(\mathbf{x} - \mathbf{jw}) \rangle = 0 \quad \forall i : 0 < i < k, \end{cases} \end{aligned}$$

where the functional J_{μ} is defined by

$$J_{\mu}[u] = \frac{1}{\mu} \|u\|_{L^1} + \langle u, -\frac{1}{2} \Delta u \rangle = \frac{1}{\mu} \|u\|_{L^1} + \frac{1}{2} \|\nabla u\|_{L^2}.$$

Notice that this functional is a special case of the functional (4.2), with $V \equiv 0$.

The translations of the BCPWs on the lattice $\Gamma_{\mathbf{w}}$ produce all CPWs. Unlike CMs that are solved in a single minimization problem, the compressed plane waves are constructed hierarchically. This is similar to the shift-orthogonality wavelets, but a distinction of CPWs is that it is adapted to the Laplace’s operator.

The existence of CPWs essentially follows from the observation that shift orthogonality (i.e. the constraints in the definition of BCPWs) is preserved under L^2 -limits, so that any minimizing sequence has a subsequential limit, which still satisfies the shift orthogonality properties.

The following theorem (see for example [4]) characterizes any orthonormal sequence of shift orthogonal functions.

THEOREM 4.3. *Let $\Omega \subset \mathbb{R}^d$, and let the lattice $\Gamma_{\mathbf{w}}$ be defined as above. Let $\{\xi^k\}_{k=1}^{\infty}$ be an orthonormal sequence of shift orthogonal functions. Then, the (Hilbert) space $\mathcal{H} = L^2(\Omega)$ can be written as a direct sum*

$$\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2 \oplus \dots \oplus \mathcal{H}_N,$$

where each \mathcal{H}_k is the Hilbert space spanned by some eigenfunctions for the Laplace’s equation in the rectangular box Ω , with the property that, if ξ^k has the decomposition

$$\sqrt{N} \xi^k = e_1^k + e_2^k + \dots + e_N^k, \quad e_j^k \in \mathcal{H}_j,$$

then $E = \{e_j^k \mid j = 1, 2, \dots, N; k \in \mathbb{N}\}$ forms an orthonormal system in \mathcal{H} . Furthermore, $N = n_1 n_2 \dots n_d = |\Gamma_{\mathbf{w}}|$.

A detailed discussion of Theorem 4.3, with a characterization of the Hilbert spaces, \mathcal{H}_k is given in Appendix A.

REMARK 4.1. For a fixed k , both $\{\xi_j^k \mid \mathbf{jw} \in \Gamma_{\mathbf{w}}\}$ and $\{e_j^k \mid j = 1, 2, \dots, N\}$ form an orthonormal system and have the same cardinality. Hence, their linear spans agree. Therefore,

$$\operatorname{span}\{\xi_j^k \mid \mathbf{jw} \in \Gamma_{\mathbf{w}}; k = 1, \dots, M\} = \operatorname{span}\{e_j^k \mid j = 1, \dots, N; k = 1, \dots, M\}$$

for any $M \in \mathbb{N} \cup \{\infty\}$.

With this remark and Theorem 4.3 in mind, instead of working with the CPWs $\{\psi_j^k\}$, it is natural to switch to $\{e_j^k\}$ for completeness results. We define

$$J_\infty[u] = \frac{1}{2} \|\nabla u\|_{L^2(\Omega)}^2.$$

Then, we can write

$$J_\mu[u] = J_\infty[u] + \frac{1}{\mu} \|u\|_{L^1}.$$

As the Hilbert spaces \mathcal{H}_j are the span of some eigenfunctions of the Laplacian, the functional J_∞ satisfies the following linearity property:

$$J_\infty[e_1 + e_2 + \dots + e_N] = J_\infty[e_1] + J_\infty[e_2] + \dots + J_\infty[e_N], \quad e_j \in \mathcal{H}_j, \quad j = 1, 2, \dots, N.$$

Now, the minimization procedure for $\{e_j^k\}$ becomes

$$\begin{aligned} \{e_1^1, e_2^1, \dots, e_N^1\} &= \underset{\substack{f_j \in \mathcal{H}_j \\ \|f_j\|_2=1}}{\operatorname{argmin}} J_\infty[f_1] + \dots + J_\infty[f_N] + \frac{1}{\mu} \|f_1 + \dots + f_N\|_1, \\ \{e_1^k, e_2^k, \dots, e_N^k\} &= \underset{\substack{f_j \in \{e_j^1, \dots, e_j^{k-1}\}^\perp \cap \mathcal{H}_j \\ \|f_j\|_2=1}}{\operatorname{argmin}} J_\infty[f_1] + \dots + J_\infty[f_N] + \frac{1}{\mu} \|f_1 + \dots + f_N\|_1. \end{aligned}$$

This is analogous to the variational procedure (2.1), except that, at each step in the minimization, we obtain multiple functions. Nevertheless, we might view one particular e_j^k , say e_1^k for simplicity, as the solution to the following minimization problem over \mathcal{H}_1 :

$$e_1^k = \underset{\substack{f \in \{e_1^1, \dots, e_1^{k-1}\}^\perp \cap \mathcal{H}_1 \\ \|f\|_2=1}}{\operatorname{argmin}} J_\infty[f] + J_\infty[e_2^k] + \dots + J_\infty[e_N^k] + \frac{1}{\mu} \|f + e_2^k + \dots + e_N^k\|_1.$$

We still have the boundedness of the penalty term $P(f) = \frac{1}{\mu} \|f + e_2^k + \dots + e_N^k\|_1$, as

$$\begin{aligned} \frac{1}{\mu} \|f + e_2^k + \dots + e_N^k\|_1 &\leq \frac{1}{\mu} (\|f\|_1 + \|e_2^k\|_1 + \dots + \|e_N^k\|_1) \\ (\text{Cauchy-Schwarz}) &\leq \frac{|\Omega|^{\frac{1}{2}}}{\mu} (\|f\|_2 + \|e_2^k\|_2 + \dots + \|e_N^k\|_2) \\ &= \frac{N|\Omega|^{\frac{1}{2}}}{\mu}, \end{aligned}$$

i.e.

$$\|P\| \leq \frac{N|\Omega|^{\frac{1}{2}}}{\mu}. \tag{4.6}$$

Therefore, $\{e_j^k\}_{k \in \mathbb{N}}$ could be viewed as the solutions to an analogue of the variational procedure (2.1) in the Hilbert space \mathcal{H}_j , with the linear functional being the restriction of $-\Delta$ to \mathcal{H}_j .

Next, let's enumerate the eigenfunctions forming each Hilbert space \mathcal{H}_j as follows

$$\begin{aligned} \mathcal{H}_j &= \text{span}\{\phi_j^k | k = 1, 2, \dots\}, \\ -\frac{1}{2}\Delta\phi_j^k &= \lambda_j^k \phi_j^k \\ \lambda_j^1 &\leq \lambda_j^2 \leq \dots \end{aligned} \tag{4.7}$$

The following theorem establishes an analogue of Weyl's law.

THEOREM 4.4. *Let λ_j^k be defined as in (4.7). Then,*

$$\lambda_j^k = \frac{N(2\pi)^d}{2\omega_d|\Omega|}k^{2/d} + o(k^{2/d}), \text{ as } k \rightarrow \infty,$$

where ω_d denotes the volume of the unit ball in \mathbb{R}^d .

A discussion of Theorem 4.4 can be found in Appendix A.

We verified that the functions $\{e_j^k\}$ are obtained via a variational procedure analogous to (2.1). We also noted in Remark 4.1 that the spans of $\{e_j^k\}$ and CPWs agree. Therefore, the theory developed in Section 2 applies to CPWs, so that we obtain the following corollaries as direct consequences of theorems 2.7, 2.8, and 2.9.

COROLLARY 4.5 (Corollary to Theorem 2.7). *Let Ω be a bounded interval in \mathbb{R} that is an integer multiple of some lattice $\Gamma_{\mathbf{w}}$. Then, for any parameter μ , the set of compressed plane waves $\{\psi_j^k\}$ defined on Ω forms a complete orthonormal system in $L^2(\Omega)$.*

Proof. Notice that $\{e_j^k\}_{k \in \mathbb{N}} \subset \mathcal{H}_j$ are obtained as the solutions to a variational problem in \mathcal{H}_j , analogous to the variational procedure (2.1). Furthermore, since Ω lies inside \mathbb{R} , by Theorem 4.4, the corresponding eigenvalues grow super-linearly. Therefore, Theorem 2.7 applies so that $\{e_j^k\}_{k \in \mathbb{N}}$ forms a complete orthonormal system in \mathcal{H}_j for each $j = 1, 2, \dots, N$. Finally, by Remark 4.1, $\{\psi_j^k\}$ is a complete orthonormal system in \mathcal{H} . □

COROLLARY 4.6 (Corollary to Theorem 2.8). *Let Ω be a rectangular domain inside \mathbb{R}^2 that is an integer multiple of some lattice $\Gamma_{\mathbf{w}}$. Then, for any parameter μ satisfying*

$$\mu > \frac{|\Omega|^{\frac{3}{2}}}{2\pi}, \tag{4.8}$$

the set of compressed plane waves $\{\psi_j^k\}$ defined on Ω forms a complete orthonormal system in $L^2(\Omega)$.

Proof. Since Ω lies inside \mathbb{R}^2 , notice by Theorem 4.4 that the corresponding eigenvalues grow linearly, with the linearity constant $\frac{2\pi N}{|\Omega|}$. The proof proceeds analogous to the proof of Corollary 4.5, except that we apply Theorem 2.8. The bound for the penalty term is provided in (4.6), so that Theorem 2.8 holds precisely when the bound (4.8) is satisfied. □

COROLLARY 4.7 (Corollary to Theorem 2.9). *Let V_j^m be the subspace generated by the functions $\{e_j^1, e_j^2, \dots, e_j^m\}$. Then, for any $n \leq m$, we have*

$$\sum_{k=1}^n d(\phi_j^k, V_j^m)^2 \leq \frac{mN|\Omega|^{\frac{1}{2}}}{\mu(\lambda_j^{m+1} - \lambda_j^n)},$$

provided that $\lambda_j^{m+1} \neq \lambda_j^n$. Defining V^m via

$$V^m = \text{span} \{ \psi_j^k \mid \mathbf{j} \in \mathbb{Z}^d, k = 1, 2, \dots, m \},$$

we further have

$$\sum_{k \leq m, j \leq N} d(\phi_j^k, V^m)^2 = \frac{mN|\Omega|^{\frac{1}{2}}}{\mu} \sum_{j \leq N} \frac{1}{\lambda_j^{m+1} - \lambda_j^n}.$$

Appendix A. We will provide an explicit characterization of the Hilbert spaces \mathcal{H}_j in Theorem 4.3. The eigenfunctions of the Laplace’s operator in a rectangular domain $\Omega = [0, n_1w_1] \times \dots \times [0, n_dw_d]$ is given by

$$\phi_{m_1, \dots, m_d}(\mathbf{x}) = e^{2\pi i \left(\frac{m_1x_1}{n_1w_1} + \frac{m_2x_2}{n_2w_2} + \dots + \frac{m_dx_d}{n_dw_d} \right)},$$

where $(m_1, m_2, \dots, m_d) \in \mathbb{Z}^d$. Hence, if we form the lattice

$$\Pi_{\mathbf{w}} = \left\{ \left(\frac{m_1}{n_1w_1}, \frac{m_2}{n_2w_2}, \dots, \frac{m_d}{n_dw_d} \right) \mid (m_1, m_2, \dots, m_d) \in \mathbb{Z}^d \right\},$$

then each of the eigenfunctions of the Laplace’s operator in the domain Ω can be represented as

$$\phi_v(\mathbf{x}) = e^{2\pi i v \cdot \mathbf{x}}, \quad v \in \Pi_{\mathbf{w}},$$

with the corresponding eigenvalue $\lambda_v = 4\pi^2|v|^2$. Now, we define

$$\Lambda_{\mathbf{w}} = \left\{ \left(\frac{m_1}{n_1w_1}, \frac{m_2}{n_2w_2}, \dots, \frac{m_d}{n_dw_d} \right) \mid 0 \leq m_1 < n_1, 0 \leq m_2 < n_2, \dots, 0 \leq m_d < n_d \right\}.$$

Each $\rho \in \Lambda_{\mathbf{w}}$ has a natural periodic extension in $\Pi_{\mathbf{w}}$ with respect to $\Gamma_{\mathbf{w}}$. For each $\rho \in \Lambda_{\mathbf{w}}$, we denote such extension by Σ_{ρ} . Now, the family of Hilbert spaces \mathcal{H}_j in Theorem 4.3 consists of the Hilbert spaces

$$\mathcal{H}_{\rho} = \text{span}\{\phi_v \mid v \in \Sigma_{\rho}\}.$$

Since $\Lambda_{\mathbf{w}}$, and $\Gamma_{\mathbf{w}}$ each has cardinality $n_1n_2 \dots n_d$; N , the cardinality of the family of Hilbert spaces $\{\mathcal{H}_j\}_{j=1}^N$, satisfies $N = n_1n_2 \dots n_d = |\Gamma_{\mathbf{w}}|$, as asserted in Theorem 4.3.

We have already observed that the eigenvalue corresponding to ϕ_v is $\lambda_v = 4\pi^2|v|^2$. Weyl’s law in the rectangular domain case can be viewed as the growth of the size of the distance between lattice points and the origin. Therefore, with all these lattice characterization of the eigenfunctions, it is not hard to see that the growth of the eigenvalues corresponding to the eigenfunctions in each of the Hilbert spaces \mathcal{H}_j are given precisely as in Theorem 4.4.

As an illustration, we can consider $\Omega = [0, 2] \times [0, 3] \subset \mathbb{R}^2$, and $\mathbf{w} = (1, 1)$. Then, $\Gamma_{\mathbf{w}}$ becomes

$$\Gamma_{\mathbf{w}} = \{(0, 0), (0, 1), (0, 2), (1, 0), (1, 1), (1, 2)\}.$$

The eigenfunctions for the Laplace’s equation in Ω are given by

$$\phi_{m,n}(x, y) = e^{2\pi i \left(\frac{mx}{2} + \frac{ny}{3} \right)},$$

so that

$$\Pi_{\mathbf{w}} = \left\{ \left(\frac{m}{2}, \frac{n}{3} \right) \mid m, n \in \mathbb{Z} \right\}.$$

Now, the finite lattice $\Lambda_{\mathbf{w}}$ becomes

$$\Lambda_{\mathbf{w}} = \left\{ (0, 0), \left(0, \frac{1}{3} \right), \left(0, \frac{2}{3} \right), \left(\frac{1}{2}, 0 \right), \left(\frac{1}{2}, \frac{1}{3} \right), \left(\frac{1}{2}, \frac{2}{3} \right) \right\}.$$

Finally, the decomposition given in Theorem 4.3 becomes

$$L^2(\Omega) = \mathcal{H}_1 \oplus \mathcal{H}_2 \oplus \mathcal{H}_3 \oplus \mathcal{H}_4 \oplus \mathcal{H}_5 \oplus \mathcal{H}_6,$$

where

$$\begin{aligned} \mathcal{H}_1 &= \text{span}\{\phi_{2k,3l}\}_{k,l \in \mathbb{Z}}, & \mathcal{H}_2 &= \text{span}\{\phi_{2k,3l+1}\}_{k,l \in \mathbb{Z}}, & \mathcal{H}_3 &= \text{span}\{\phi_{2k,3l+2}\}_{k,l \in \mathbb{Z}}, \\ \mathcal{H}_4 &= \text{span}\{\phi_{2k+1,3l}\}_{k,l \in \mathbb{Z}}, & \mathcal{H}_5 &= \text{span}\{\phi_{2k+1,3l+1}\}_{k,l \in \mathbb{Z}}, & \mathcal{H}_6 &= \text{span}\{\phi_{2k+1,3l+2}\}_{k,l \in \mathbb{Z}}. \end{aligned}$$

Acknowledgement. The authors would like to thank Stanley Osher and Russel Caffisch for their helpful comments throughout the development of the paper.

REFERENCES

- [1] Farzin Barekat, *Applications of Stochastic Simulation and Compressed Sensing to Large Systems*, PhD thesis, University of California, Los Angeles, 2014.
- [2] Farzin Barekat, *On the consistency of compressed modes for variational problems associated with the Schrödinger operator*, SIAM J. Math. Anal., 46(5), 3568–3577, 2014.
- [3] Farzin Barekat, Russel Caffisch, and Stanley Osher, *On the support of compressed modes*, UCLA CAM Reports, 14-14, 2014.
- [4] Farzin Barekat, Ke Yin, Russel E Caffisch, Stanley J Osher, Rongjie Lai, and Vidvuds Ozolins, *Compressed Wannier modes found from an l_1 regularized energy functional*, arXiv preprint arXiv:1403.6883, 2014.
- [5] Haim Brézis, *Solutions with compact support of variational inequalities*, Russian Mathematical Surveys, 29(2), 103–108, 1974.
- [6] Haim Brézis, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Springer, 2011.
- [7] Haim Brézis, Avner Friedman, et al., *Estimates on the support of solutions of parabolic variational inequalities*, Illinois Journal of Mathematics, 20(1), 82–97, 1976.
- [8] Lawrence Evans, *Partial Differential Equations*, American Mathematical Soc., 2012.
- [9] Vidvuds Ozoliņš, Rongjie Lai, Russel Caffisch, and Stanley Osher, *Compressed modes for variational problems in mathematics and physics*, Proceedings of the National Academy of Sciences, 110(46), 18368–18373, 2013.
- [10] Vidvuds Ozoliņš, Rongjie Lai, Russel Caffisch, and Stanley Osher, *Compressed plane waves yield a compactly supported multiresolution basis for the Laplace operator*, Proceedings of the National Academy of Sciences, 111(5), 1691–1696, 2014.