

On q -analogs of some families of multiple harmonic sums and multiple zeta star value identities

KH. HESSAMI PILEHROOD, T. HESSAMI PILEHROOD,
AND JIANQIANG ZHAO

In recent years, there has been intensive research on the \mathbb{Q} -linear relations between multiple zeta (star) values. In this paper, we prove many families of identities involving the q -analog of these values, from which we can always recover the corresponding classical identities by taking $q \rightarrow 1$. The main results of the paper (Theorems 1.4 and 5.4) are the duality relations between multiple zeta star values and Euler sums and their q -analogs, which are generalizations of the Two-one formula and some multiple harmonic sum identities and their q -analogs proved by the authors recently. Such duality relations lead to a proof of the conjecture by Ihara *et al.* that the Hoffman \star -elements $\zeta^*(s_1, \dots, s_r)$ with $s_i \in \{2, 3\}$ span the vector space generated by multiple zeta values over \mathbb{Q} .

1. Introduction

Multiple harmonic sums (MHS) are nested generalizations of harmonic sums and multiple zeta values (MZV) are the limits of MHS when the number of terms in the sum goes to infinity. In recent years, MHS, MZV and their generalizations have been found to be intimately related to Feynman integrals in perturbative quantum field theory [2, 5, 19] in physics as well as to Hopf and Lie algebras, combinatorics (double shuffle relations) [12–14], algebraic geometry [4, 8, 9], and even modular forms [6] in mathematics.

We now recall their basic setup. In order to unify MHS, MZV and their alternating versions we first define a sort of double cover of the set $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ where \mathbb{N} is the set of positive integers.

2010 Mathematics Subject Classification: 11M32, 11B65.

Key words and phrases: multiple harmonic sums, multiple zeta values, multiple zeta star values, Euler sums.

Definition 1.1. Let $\mathbb{D}_0 := \mathbb{N}_0 \cup \bar{\mathbb{N}}_0$ and $\mathbb{D} := \mathbb{N} \cup \bar{\mathbb{N}}$ be the sets of *signed nonnegative* and *signed positive numbers*, respectively, where

$$\bar{\mathbb{N}}_0 = \{\bar{k} : k \in \mathbb{N}_0\} \quad \text{and} \quad \bar{\mathbb{N}} = \{\bar{k} : k \in \mathbb{N}\}.$$

In some sense, \bar{k} is k dressed by a negative sign, but \bar{k} is not a negative number. Define for all $k \in \mathbb{N}_0$ the absolute value function $|\cdot|$ on \mathbb{D}_0 by $|k| = |\bar{k}| = k$ and the sign function by $\operatorname{sgn}(k) = 1$ and $\operatorname{sgn}(\bar{k}) = -1$. We make \mathbb{D}_0 a semi-group by defining a commutative and associative binary operation \oplus (called *O-plus*) as follows: for all $a, b \in \mathbb{D}_0$

$$(1) \quad a \oplus b = \begin{cases} \overline{|a| + |b|}, & \text{if only one of } a \text{ or } b \text{ is in } \mathbb{N}_0; \\ a + b, & \text{if } a, b \in \mathbb{N}_0; \\ |a| + |b|, & \text{if } a, b \in \bar{\mathbb{N}}_0. \end{cases}$$

For $\mathbf{s} = (s_1, \dots, s_m) \in \mathbb{D}^m$, we define the (alternating) multiple harmonic sums by

$$H_n(\mathbf{s}) := \sum_{n \geq k_1 > \dots > k_m \geq 1} \prod_{j=1}^m \frac{\operatorname{sgn}(s_j)^{k_j}}{k_j^{|s_j|}},$$

$$\text{and} \quad H_n^*(\mathbf{s}) := \sum_{n \geq k_1 \geq \dots \geq k_m \geq 1} \prod_{j=1}^m \frac{\operatorname{sgn}(s_j)^{k_j}}{k_j^{|s_j|}}.$$

Correspondingly, we can define the (alternating) Euler sums by

$$(2) \quad \zeta(\mathbf{s}) := \sum_{k_1 > \dots > k_m \geq 1} \prod_{j=1}^m \frac{\operatorname{sgn}(s_j)^{k_j}}{k_j^{|s_j|}},$$

$$\text{and} \quad \zeta^*(\mathbf{s}) := \sum_{k_1 \geq \dots \geq k_m \geq 1} \prod_{j=1}^m \frac{\operatorname{sgn}(s_j)^{k_j}}{k_j^{|s_j|}}$$

where $s_1 \neq 1$ in order for the series to converge. If $\mathbf{s} \in \mathbb{N}^m$ then $\zeta(\mathbf{s})$ is called a *multiple zeta value* (MZV) and $\zeta^*(\mathbf{s})$ a *multiple zeta star value* (MZSV). We call $\ell(\mathbf{s}) = m$ the *length* (or *depth*) and $|\mathbf{s}| = |s_1| + \dots + |s_m|$ the *weight* of the string \mathbf{s} . One of the central themes in the study of Euler sums, MZV and MZSV is to find as many \mathbb{Q} -linear relations between these values as possible. Conjecturally, nontrivial relations can exist only among MZV and MZSV of the same weight. Following Ganois [7], we define the Euler \sharp sums

by

$$\zeta^\sharp(\mathbf{s}) := \sum_{\mathbf{p} = s_1 \circ s_2 \circ \dots \circ s_d} 2^{\ell(\mathbf{p})} \zeta(\mathbf{p}),$$

where \mathbf{p} runs through all indices of the form $(s_1 \circ s_2 \circ \dots \circ s_d)$ with “ \circ ” being either the symbol “,” or the O-plus “ \oplus ”. Similar notation for MZV has already appeared in the literature before, see, for example, [20]. In [17, 23] Linebarger and the third author obtained many families of identities involving both MHS and MZSV after getting inspiration from [11]. In particular, the third author proved the following so-called Two-one formula conjectured by Ohno and Zudilin in [18]:

Theorem 1.2. ([23, Theorem 1.3]) *Let $r \in \mathbb{N}$ and $\mathbf{s} = (\{2\}^{a_1}, 1, \dots, \{2\}^{a_r}, 1)$ where $a_1 \in \mathbb{N}$ and $a_j \in \mathbb{N}_0$ for all $j \geq 2$. Then we have*

$$\zeta^*(\mathbf{s}) = \zeta^\sharp(2a_1 + 1, \dots, 2a_r + 1).$$

One particularly well-behaved q -analog of the multiple zeta functions is defined in [21] by the third author, generalizing the Riemann q -zeta function studied by Kaneko et al. [16]. There, again, it is very important to understand the relations between their special values, see [3] for some relevant results. Recently, the first two authors proved a q -analog of the Two-one formula in [10]. Our original goal of this paper was to provide further analogs of the identities contained in [17, 23]. However, we have achieved much more because we can now actually treat arbitrary q -MZSV and express it in terms of q -analog Euler sums (of the non-star version). By taking $q \rightarrow 1$ we obtain the following result.

Theorem 1.3. *Let $\mathbf{s} = (s_1, \dots, s_d) \in \mathbb{N}^d$ with $s_1 > 1$. Set $\iota_{\mathbf{s}} = 1$ if $s_d = 1$ and $\iota_{\mathbf{s}} = -1$ if $s_d > 1$. Suppose there exists $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_m) \in \mathbb{D}^m$ (determined uniquely by \mathbf{s}) such that there is an expansion of the form*

$$\zeta^*(\mathbf{s}) = \iota_{\mathbf{s}} \zeta^\sharp(\lambda_1, \lambda_2, \dots, \lambda_m).$$

Then we have

(i) *For any positive integer a ,*

$$\zeta^*(\{2\}^a, \mathbf{s}) = \iota_{\mathbf{s}} \zeta^\sharp(2a \oplus \lambda_1, \lambda_2, \dots, \lambda_m).$$

(ii) *For any positive integers a and l ,*

$$\zeta^*(\{2\}^a, \{1\}^l, \mathbf{s}) = \iota_{\mathbf{s}} \zeta^\sharp(2a + 1, \{1\}^{l-1}, \lambda_1, \lambda_2, \dots, \lambda_m).$$

(iii) For any positive integers $c \geq 3$ and l ,

$$\zeta^*(c, \{1\}^l, \mathbf{s}) = \iota_{\mathbf{s}} \zeta^\sharp(\overline{2}, \{1\}^{c-3}, \overline{2}, \{1\}^{l-1}, \lambda_1, \lambda_2, \dots, \lambda_m).$$

(iv) For any integers $b \geq 0$ and $c \geq 3$,

$$\zeta^*(\{2\}^b, c, \mathbf{s}) = \iota_{\mathbf{s}} \zeta^\sharp(\overline{2b+2}, \{1\}^{c-3}, \lambda_1 \oplus \overline{1}, \lambda_2, \dots, \lambda_m).$$

This provides very elegant simplifications when \mathbf{s} contains many 2's in it. The following identity is an illuminating example: for any $a \in \mathbb{N}$ and $b, c, d \in \mathbb{N}_0$, we have

$$\begin{aligned} (3) \quad & \zeta^*(\{2\}^a, 1, \{2\}^b, 1, \{2\}^c, 3, \{2\}^d, 1) \\ &= 2\zeta(2a + 2b + 2c + 2d + 6) + 4\zeta(2a + 1, 2b + 2c + 2d + 5) \\ &\quad + 4\zeta(2a + 2b + 2, 2c + 2d + 4) + 4\zeta(\overline{2a + 2b + 2c + 4}, \overline{2d + 2}) \\ &\quad + 8\zeta(2a + 1, 2b + 1, 2c + 2d + 4) + 8\zeta(2a + 1, \overline{2b + 2c + 3}, \overline{2d + 2}) \\ &\quad + 8\zeta(2a + 2b + 2, \overline{2c + 2}, \overline{2d + 2}) + 16\zeta(2a + 1, 2b + 1, \overline{2c + 2}, \overline{2d + 2}). \end{aligned}$$

We also verified this identity numerically for $1 \leq a \leq 2$ and $0 \leq b, c, d \leq 2$ with EZ-face [1] with errors bounded by 10^{-50} .

According to Theorem 1.3, we can treat arbitrary MZSV by building up from the three base cases: $\zeta^*(\{2\}^a)$, $\zeta^*(\{2\}^a, 1)$ ($a \geq 1$) and $\zeta^*(\{2\}^b, c)$ ($b \geq 0$ and $c \geq 3$) treated in [11]. Here is the general statement.

Theorem 1.4. Let $a_0, a_j \in \mathbb{N}_0$, $c_j \in \mathbb{N}$ and $c_j \neq 2$ for all $j = 1, \dots, d$. Assume $a_0 > 0$ or $c_1 \geq 3$. Set $\delta(c) = 1$ if $c = 1$ and $\delta(c) = 0$ if $c \geq 3$. Moreover, put $\{1\}^n = \{1\}^{\max(n, 0)}$. Then we have

$$(4) \quad \zeta^*(\{2\}^{a_0}, c_1, \{2\}^{a_1}, \dots, c_d, \{2\}^{a_d}) = \pm \zeta^\sharp(B_0, \{1\}^{c_1-3}, B_1, \dots, \{1\}^{c_d-3}, B_d).$$

Here the leading sign \pm is + if and only if $a_d = 0$ and $c_d = 1$, and

$$B_j = \begin{cases} A_j, & \text{if } A_j \text{ is odd;} \\ \overline{A_j}, & \text{if } A_j > 0 \text{ and even;} \\ \text{vacuous,} & \text{if } A_j = 0, \end{cases}$$

where

$$A_j = \begin{cases} 2a_0 + 2 - \delta(c_1), & \text{if } j = 0; \\ 2a_d + 1 - \delta(c_d), & \text{if } j = d; \\ 2a_j + 3 - \delta(c_j) - \delta(c_{j+1}), & \text{if } 0 < j < d. \end{cases}$$

Formula (4) can be considered as a general duality relation which expresses arbitrary multiple zeta star value in terms of Euler \sharp sums. It generalizes the Two-one formula and many other 2-c-2-c, 2-1-2-c, 2-c-2-1 formulas with $c \geq 3$, proved by the third author [17, 23]. A q -analog of Theorem 1.4 is given in Section 5, which is Theorem 5.4.

In her Ph.D. thesis, Glanois [7] studied motivic versions of multiple Euler \sharp sums and proved that the motivic versions of

$$\zeta^\sharp(\overline{2a_0+2}, 2a_1+3, \dots, 2a_{d-1}+3, 2a_d+1), \quad \text{with } a_i \geq 0,$$

form a graded basis of the space of motivic multiple zeta values. As a consequence, by application of the period map, she obtained the following important result.

Theorem 1.5 (Glanois). *Each multiple zeta value is a \mathbb{Q} -linear combination of elements of the same weight in*

$$\{\zeta^\sharp(\overline{2a_0+2}, 2a_1+3, \dots, 2a_{d-1}+3, 2a_d+1), \quad a_i \geq 0\}.$$

Note that Ihara *et al.* [15] conjectured that the Hoffman \star -elements $\zeta^*(s_1, \dots, s_d)$ with $s_i \in \{2, 3\}$ form a basis of the space of MZVs over \mathbb{Q} . Taking into account the Two-three formula, which is a consequence of identity (4) with $c_1 = \dots = c_d = 3$,

$$\begin{aligned} & \zeta^*(\{2\}^{a_0}, 3, \{2\}^{a_1}, \dots, 3, \{2\}^{a_d}) \\ &= -\zeta^\sharp(\overline{2a_0+2}, 2a_1+3, \dots, 2a_{d-1}+3, 2a_d+1), \end{aligned}$$

and combining it with Theorem 1.5, we get the following statement confirming the conjecture of Ihara *et al.*

Corollary 1.6. *Every multiple zeta value of weight w is a \mathbb{Q} -linear combination of the Hoffman \star -elements $\zeta^*(s_1, \dots, s_d)$ with $s_i \in \{2, 3\}$ and $\sum s_i = w$.*

In her Ph.D. thesis, Glanois also conjectures that the motivic version of Theorem 1.4 should hold (see [7, Conjecture 4.5.1]), whose proof should follow from Theorem 1.4 and a Galois descent argument used first by Brown in [4] to prove that all the periods of mixed motives unramified over \mathbb{Z} are $\mathbb{Q}[\frac{1}{2\pi i}]$ -linear combinations of MZVs. The motivic version of Theorem 1.4 would imply the motivic version of the Conjecture of Ihara *et al.* for the space of motivic multiple zeta values.

Acknowledgement. KHP and THP gratefully acknowledge support from the Research Immersion Fellowships of the Fields Institute. JZ is partly supported by NSF grant DMS-1162116 and the Severo Ochoa Excellence Program at ICMAT. He also would like to thank the Max-Planck Institute for Mathematics, the Morningside Center of Mathematics in Beijing, the Kavli Institute for Theoretical Physics China and National Taiwan University for their hospitality where part of this work was done.

2. Preliminaries and notations

In this section, we first fix some notation. Throughout the paper let m and n denote nonnegative integers and q a real number with $0 < q < 1$. For any real number a , put

$$(a)_0 := (a; q)_0 := 1, \quad (a)_n := (a; q)_n := \prod_{k=0}^{n-1} (1 - aq^k), \quad n \geq 1.$$

As a convention, throughout the paper we always use $[]$ to denote q -analog objects. For example, the q -analog of a positive integer n is given by

$$[n] = [n]_q := \sum_{k=0}^{n-1} q^k = \frac{1 - q^n}{1 - q},$$

and the Gaussian q -binomial coefficient

$$\begin{bmatrix} n \\ m \end{bmatrix} := \begin{cases} \frac{(q)_n}{(q)_m (q)_{n-m}}, & \text{if } 0 \leq m \leq n, \\ 0, & \text{otherwise.} \end{cases}$$

For $m \in \mathbb{N}_0$ and $\mathbf{s} = (s_1, \dots, s_m) \in \mathbb{D}_0^m$, we set $\mathbf{s} = \emptyset$ if $m = 0$ and define the q -analogs of multiple harmonic (star) sums (q -MHS)

$$\begin{aligned} H_n[\mathbf{s}] &:= \sum_{n \geq k_1 > \dots > k_m \geq 1} \prod_{j=1}^m \frac{\operatorname{sgn}(s_j)^{k_j} q^{k_j}}{[k_j]^{s_j}}, \\ \text{and} \quad H_n^*[\mathbf{s}] &:= \sum_{n \geq k_1 \geq \dots \geq k_m \geq 1} \prod_{j=1}^m \frac{\operatorname{sgn}(s_j)^{k_j} q^{k_j}}{[k_j]^{s_j}}, \end{aligned}$$

with the convention that $H_n[\mathbf{s}] = 0$ if $n < m$, and $H_n^*[\emptyset] = H_n[\emptyset] = 1$ for all $n \geq 0$. Notice that we allow s_j to be 0 or $\bar{0}$ in these q -MHS.

Now we fix a symbol θ and define for any $r \in \mathbb{Z} \cup \{\theta\}$ and $k \in \mathbb{N}$ the function

$$Q(r, k) := \begin{cases} rk(k-1)/2, & \text{if } r > 0; \\ rk(k-1)/2 - k, & \text{if } r \leq 0; \\ 0, & \text{if } r = \theta. \end{cases}$$

For $\mathbf{s} = (s_1, \dots, s_m) \in \mathbb{D}_0^m$, $\mathbf{t} = (t_1, \dots, t_m) \in (\mathbb{N}_0)^m$, and $\mathbf{r} = (r_1, \dots, r_m) \in (\mathbb{Z} \cup \{\theta\})^m$, we define the mollified companion of $H_n[\mathbf{s}]$ by

$$(5) \quad \mathcal{H}_n[\mathbf{s}; \mathbf{t}; \mathbf{r}] := \sum_{n \geq k_1 > \dots > k_m \geq 1} \frac{\begin{bmatrix} n \\ k_1 \end{bmatrix}}{\begin{bmatrix} n+k_1 \\ k_1 \end{bmatrix}} \prod_{j=1}^m \frac{q^{t_j k_j + Q(r_j, k_j)} (1 + q^{k_j})}{\operatorname{sgn}(s_j)^{k_j} [k_j]^{s_j}}.$$

We call $[\mathbf{s}; \mathbf{t}; \mathbf{r}]$ an *admissible triple of mollifiers* if the limit $\lim_{n \rightarrow \infty} \mathcal{H}_n[\mathbf{s}; \mathbf{t}; \mathbf{r}]$ exists.

Definition 2.1. Let \square -plus \boxplus be a binary operation on $\mathbb{Z} \cup \{\theta\}$ such that

- $\theta \boxplus a = a \boxplus \theta = a$ for all $a \in \mathbb{Z} \cup \{\theta\}$,
- $a \boxplus b = a + b$ for all $a, b \in \mathbb{Z}$ with $(a, b) \neq (1, -1), (-1, 1)$, and
- $1 \boxplus (-1) = \theta$ and $(-1) \boxplus 1 = 0$.

Lemma 2.2. Let $r \in \{\theta\} \cup \mathbb{Z} \setminus \{0\}$, $d \in \{0, -1\}$. Then for any $k \in \mathbb{Z}$

$$(6) \quad Q(r, k) + Q(1, k) = Q(r \boxplus 1, k),$$

$$(7) \quad k^2 + Q(d, k) = Q(2 \boxplus d, k).$$

Moreover, the projection

$$\begin{aligned} \pi : (\mathbb{Z} \cup \{\theta\}, \boxplus) &\longrightarrow (\mathbb{Z}, +) \\ a &\longmapsto a \quad \forall a \in \mathbb{Z}, \\ \theta &\longmapsto 0, \end{aligned}$$

is a homomorphism of semi-groups and its restriction to \mathbb{Z}^* is injective.

Proof. Clear. □

For an admissible triple of mollifiers $[\mathbf{s}; \mathbf{t}; \mathbf{r}]$, we define

$$\{s_1 \circ \cdots \circ s_m; t_1 \circ \cdots \circ t_m; r_1 \circ \cdots \circ r_m\}$$

to be the set of triples of strings produced by replacing every \circ in \mathbf{s} by either comma “,” or O-plus “ \oplus ”, replacing every \circ in \mathbf{t} by either comma “,” or the usual plus “+”, and replacing every \circ in \mathbf{r} by either comma “,” or \square -plus “ \boxplus ”. Moreover, the commas should be at the same positions for all \mathbf{s} , \mathbf{t} and \mathbf{r} . Now we set

$$\mathcal{H}_n^\sharp[\mathbf{s}; \mathbf{t}; \mathbf{r}] := \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \{s_1 \circ \cdots \circ s_m; t_1 \circ \cdots \circ t_m; r_1 \circ \cdots \circ r_m\}} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}].$$

In the above notation, the Two-one formulas for q -MHS obtained in [10] have the form

$$(8) \quad H_n^*[\{2\}^a] = -\mathcal{H}_n[\overline{2a}; a; 1]$$

and, for $a_{\ell+1} \neq 0$

$$(9) \quad \begin{aligned} & H_n^*[\{2\}^{a_1}, 1, \dots, \{2\}^{a_\ell}, 1] \\ &= \mathcal{H}_n^\sharp[2a_1 + 1, \dots, 2a_\ell + 1; a_1 + 1, \dots, a_\ell + 1; 2, \{0\}^{\ell-1}], \\ & H_n^*[\{2\}^{a_1}, 1, \dots, \{2\}^{a_\ell}, 1, \{2\}^{a_{\ell+1}}] \\ &= -\mathcal{H}_n^\sharp[2a_1 + 1, \dots, 2a_\ell + 1, \overline{2a_{\ell+1}}; \\ & \quad a_1 + 1, \dots, a_\ell + 1, a_{\ell+1}; 2, \{0\}^{\ell-1}, -1]. \end{aligned}$$

Finally, we define the q -analog of *multiple zeta values*, q -MZV for short, and q -analog of *multiple zeta star values*, or q -MZSV, as

$$\zeta[\mathbf{s}] := \sum_{k_1 > \dots > k_m \geq 1} \prod_{j=1}^m \frac{\operatorname{sgn}(s_j)^{k_j} q^{k_j}}{[k_j]^{s_j}},$$

$$\text{and } \zeta^*[\mathbf{s}] := \sum_{k_1 \geq \dots \geq k_m \geq 1} \prod_{j=1}^m \frac{\operatorname{sgn}(s_j)^{k_j} q^{k_j}}{[k_j]^{s_j}},$$

respectively. The *mollified companion* of $\zeta[\mathbf{s}]$ associated with the admissible triple of mollifiers $[\mathbf{s}; \mathbf{t}; \mathbf{r}]$ is defined by

$$\begin{aligned} \mathfrak{z}[\mathbf{s}; \mathbf{t}; \mathbf{r}] &:= \mathfrak{z}[s_1, \dots, s_m; t_1, \dots, t_m; r_1, \dots, r_m] \\ &= \sum_{k_1 > \dots > k_m \geq 1} \prod_{j=1}^m \frac{\operatorname{sgn}(s_j)^{k_j} q^{t_j k_j + Q(r_j, k_j)} (1 + q^{k_j})}{[k_j]^{s_j}}, \end{aligned}$$

and its \sharp -version is defined by

$$\mathfrak{z}^\sharp[\mathbf{s}; \mathbf{t}; \mathbf{r}] := \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \{s_1 \circ \dots \circ s_m; t_1 \circ \dots \circ t_m; r_1 \circ \dots \circ r_m\}} \mathfrak{z}[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}].$$

If $m = 0$, we put $\zeta^*[\emptyset] = \mathfrak{z}[\emptyset; \emptyset; \emptyset] = 1$. Throughout the paper the triples of mollifiers $[\mathbf{s}; \mathbf{t}; \mathbf{r}]$ are chosen in such a way that the above multiple series always converges. Notice that in [10] the mollified companions of $\zeta[\mathbf{s}]$ are defined similarly.

Although our primary goal is to prove q -MZSV identities, throughout the paper we will always work with binomial identities for q -MHS first. To obtain the corresponding q -MZSV identities, we need the next result.

Lemma 2.3. ([10, Lemma 4.1]) *Let $0 < q < 1$, $c, c_1, c_2 \in \mathbb{R}$, $c > 0$, and let R_k be a sequence of real numbers satisfying $|R_k| < k^{c_1} q^{c_2 k}$ for all $k = 1, 2, \dots$. Then*

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n q^{ck^2} \left(1 - \frac{\begin{bmatrix} n \\ k \end{bmatrix}}{\begin{bmatrix} n+k \\ k \end{bmatrix}} \right) R_k = 0.$$

3. q -binomial identities

The following two combinatorial identities have been proved by the first two authors using q -WZ method.

Lemma 3.1. ([10, Lemma 2.1]) *For integers $n \geq 1$ and $l \geq 0$, we have*

$$(10) \quad \sum_{k=l+1}^n (1 + q^k) \frac{\begin{bmatrix} n \\ k \end{bmatrix}}{\begin{bmatrix} n+k \\ k \end{bmatrix}} (-1)^k q^{k(k-1)/2} = \frac{\begin{bmatrix} l \\ n \end{bmatrix}}{\begin{bmatrix} l \\ l \end{bmatrix}} \frac{\begin{bmatrix} n \\ l \end{bmatrix}}{\begin{bmatrix} n+l \\ l \end{bmatrix}} (-1)^l q^{l(l-1)/2},$$

$$(11) \quad \sum_{k=l+1}^n (1 + q^k) \frac{\begin{bmatrix} k \\ n \end{bmatrix}}{\begin{bmatrix} n+k \\ k \end{bmatrix}} q^{k(k-1)} = ([n] - [l]) \frac{\begin{bmatrix} n \\ l \end{bmatrix}}{\begin{bmatrix} n+l \\ l \end{bmatrix}} q^{l^2}.$$

The next lemma is the q -analog of [23, Lemma 2.1].

Lemma 3.2. *Let $a \in \mathbb{D}_0$, $b \in \mathbb{N}_0$, $c \in \mathbb{N}$, $r \in \{\theta\} \cup \mathbb{Z} \setminus \{0\}$, and $[\mathbf{x}; \mathbf{y}; \mathbf{z}]$ an admissible triple of mollifiers. Then for any positive integer n ,*

$$\frac{1}{[n]^c} \mathcal{H}_n[a, \mathbf{x}; b, \mathbf{y}; r \boxplus 1, \mathbf{z}] = \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \{\bar{0} \circ 1 \circ (c-1) \circ (a \oplus \bar{1}); 0 \circ c \circ b; 1 \circ \theta \circ (c-1) \circ r\}} \mathcal{H}_n[\mathbf{p}, \mathbf{x}; \tilde{\mathbf{p}}, \mathbf{y}; \tilde{\tilde{\mathbf{p}}}, \mathbf{z}].$$

Proof. We prove the lemma by induction on c . Suppose strings $\mathbf{x}, \mathbf{y}, \mathbf{z}$ have length m . Set

$$(12) \quad A_{n,k} = (-1)^k (1 + q^k) q^{k(k-1)/2} \frac{\begin{bmatrix} n \\ k \end{bmatrix}}{\begin{bmatrix} n+k \\ k \end{bmatrix}}.$$

Then by (10),

$$(13) \quad \frac{(1 + q^l)}{[l]} \sum_{k=l+1}^n A_{n,k} = \left(\frac{1}{[n]} - \frac{1}{[l]} \right) A_{n,l}$$

which, together with (6) yields

$$\begin{aligned} & \mathcal{H}_n[\bar{0}, a \oplus \bar{1}, \mathbf{x}; 0, b, \mathbf{y}; 1, r, \mathbf{z}] \\ &= \sum_{n \geq k > k_0 > k_1 > \dots > k_m \geq 1} \frac{A_{n,k} q^{bk_0 + Q(r, k_0)} (1 + q^{k_0})}{(-\text{sgn}(a))^{k_0} [k_0]^{|a|+1}} \prod_{j=1}^m \frac{q^{y_j k_j + Q(z_j, k_j)} (1 + q^{k_j})}{\text{sgn}(x_j)^{k_j} [k_j]^{|x_j|}} \\ &= \sum_{n \geq k_0 > k_1 > \dots > k_m \geq 1} \frac{q^{bk_0 + Q(r, k_0)} (1 + q^{k_0})}{(-\text{sgn}(a))^{k_0} [k_0]^{|a|+1}} \prod_{j=1}^m \frac{q^{y_j k_j + Q(z_j, k_j)} (1 + q^{k_j})}{\text{sgn}(x_j)^{k_j} [k_j]^{|x_j|}} \sum_{k=k_0+1}^n A_{n,k} \\ &= \frac{1}{[n]} \mathcal{H}_n[a, \mathbf{x}; b, \mathbf{y}; r \boxplus 1, \mathbf{z}] - \mathcal{H}_n[a \oplus 1, \mathbf{x}; b, \mathbf{y}; r \boxplus 1, \mathbf{z}]. \end{aligned}$$

This proves the lemma for $c = 1$. Now suppose $c > 1$. By the case $c = 1$ we have just proved,

$$\begin{aligned} \frac{1}{[n]^c} \mathcal{H}_n[a, \mathbf{x}; b, \mathbf{y}; r \boxplus 1, \mathbf{z}] &= \frac{1}{[n]^{c-1}} \left(\frac{1}{[n]} \mathcal{H}_n[a, \mathbf{x}; b, \mathbf{y}; r \boxplus 1, \mathbf{z}] \right) \\ &= \frac{1}{[n]^{c-1}} \mathcal{H}_n[a \oplus 1, \mathbf{x}; b, \mathbf{y}; r \boxplus 1, \mathbf{z}] \\ &\quad + \frac{1}{[n]^{c-1}} \mathcal{H}_n[\bar{0}, a \oplus \bar{1}, \mathbf{x}; 0, b, \mathbf{y}; 1, r, \mathbf{z}]. \end{aligned}$$

For the first summand, we now apply induction assumption using case $c - 1$ with a replaced by $a \oplus 1$. For the second summand, we apply Lemma 3.2 using case $c - 1$ with $a = \bar{0}$, $b = 0$, and $r = \theta$. Then we see the above is equal

to

$$\begin{aligned}
& \sum_{(\mathbf{p}, \tilde{\mathbf{p}}, \tilde{\tilde{\mathbf{p}}}) \in \{\bar{0} \circ 1^{\circ(c-2)} \circ ((a \oplus 1) \oplus \bar{1}); 0^{\circ(c-1)} \circ b; 1 \circ \theta^{\circ(c-2)} \circ r\}} \mathcal{H}_n[\mathbf{p}, \mathbf{x}; \tilde{\mathbf{p}}, \mathbf{y}; \tilde{\tilde{\mathbf{p}}}, \mathbf{z}] \\
& + \sum_{(\mathbf{p}, \tilde{\mathbf{p}}, \tilde{\tilde{\mathbf{p}}}) \in \{\bar{0} \circ 1^{\circ(c-1)}; 0^{\circ c}; 1 \circ \theta^{\circ(c-1)}\}} \mathcal{H}_n[\mathbf{p}, a \oplus \bar{1}, \mathbf{x}; \tilde{\mathbf{p}}, b, \mathbf{y}; \tilde{\tilde{\mathbf{p}}}, r, \mathbf{z}] \\
= & \sum_{(\mathbf{p}, \tilde{\mathbf{p}}, \tilde{\tilde{\mathbf{p}}}) \in \{\bar{0} \circ 1^{\circ(c-1)} \circ (a \oplus \bar{1}); 0^{\circ c} \circ b; 1 \circ \theta^{\circ(c-1)} \circ r\}} \mathcal{H}_n[\mathbf{p}, \mathbf{x}; \tilde{\mathbf{p}}, \mathbf{y}; \tilde{\tilde{\mathbf{p}}}, \mathbf{z}],
\end{aligned}$$

since $(a \oplus 1) \oplus \bar{1} = 1 \oplus (a \oplus \bar{1})$ for all $a \in \mathbb{D}_0$. We have now completed the proof of the lemma. \square

The next corollary is the degenerate case of the proceeding lemma.

Corollary 3.3. *For all $c \in \mathbb{N}_0$, we have*

$$\frac{1}{[n]^c} = -\mathcal{H}_n^\sharp[\bar{0}, \{1\}^c; \{0\}^{c+1}; 1, \{\theta\}^c].$$

Proof. The case $c = 0$ follows from (10) by setting $l = 0$. For $c \geq 1$, using the $c = 0$ case, we get

$$\frac{1}{[n]^c} = -\frac{1}{[n]^c} \mathcal{H}_n[\bar{0}; 0; 1] = - \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(c)} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}]$$

by taking $a = \bar{0}$, $b = 0$, $r = \theta$ and $\mathbf{x} = \mathbf{y} = \mathbf{z} = \emptyset$ in Lemma 3.2. Hence the corollary is proved. \square

4. MHS and MZSV identities: 2-c formula

In this section, we start with some q -MHS identities involving arguments of $(\{2\}^a, c)$ -type ($c \geq 3$). This provides one of the base cases upon which we may build general formulas of q -MZSV and MZSV whose arguments can be any admissible strings of positive integers.

Theorem 4.1. *Let $\mathbf{s} = (\{2\}^a, c)$ with $a, c \in \mathbb{N}_0$ and $c \geq 3$. Then*

$$(14) \quad H_n^*[\mathbf{s}] = -\mathcal{H}_n^\sharp[\overline{2a+2}, \{1\}^{c-2}; a+1, \{0\}^{c-2}; 1, \{\theta\}^{c-2}].$$

Proof. We proceed by induction on n . Set $\Pi(\mathbf{s}) = \{(\overline{2a+2}) \circ 1^{\circ(c-2)}; (a+1) \circ 0^{\circ(c-2)}; 1 \circ \theta^{\circ(c-2)}\}$. When $n = 1$, we have $H_1^\star(\{2\}^a, c) = q^{a+1}$. On the other hand,

$$\sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(\mathbf{s})} \mathcal{H}_1[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] = \mathcal{H}_1[\overline{2a+c}; a+1; 1] = -q^{a+1},$$

and therefore the formula is true. Suppose the statement is true for $n-1$. Then by definition

$$H_n^\star[\mathbf{s}] = \sum_{i=0}^a \frac{q^{n(a-i)}}{[n]^{2(a-i)}} H_{n-1}^\star[\{2\}^i, c] + \frac{q^{n(a+1)}}{[n]^{2a+c}}.$$

Applying inductive hypothesis, we obtain

$$(15) \quad H_n^\star[\mathbf{s}] = - \sum_{i=0}^a \frac{q^{n(a-i)}}{[n]^{2(a-i)}} \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(\{2\}^i, c)} \mathcal{H}_{n-1}[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] + \frac{q^{n(a+1)}}{[n]^{2a+c}}.$$

Set $\Pi(\mathbf{u}_{-1}) = \{0 \circ 1^{\circ(c-2)}; 0^{\circ(c-1)}; \theta^{\circ(c-1)}\}$. To save space, for any string $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_m)$, we write the substring $\boldsymbol{\lambda}_{\hat{1}} = (\lambda_2, \dots, \lambda_m)$. Then the inner sum in (15) becomes

$$\begin{aligned} & \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(\{2\}^i, c)} \mathcal{H}_{n-1}[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] \\ &= \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(\mathbf{u}_{-1})} \mathcal{H}_{n-1}[\overline{2i+2+p_1}, \mathbf{p}_{\hat{1}}; i+1+\tilde{p}_1, \tilde{\mathbf{p}}_{\hat{1}}; 1 \boxplus \tilde{\tilde{p}}_1, \tilde{\tilde{\mathbf{p}}}_{\hat{1}}] \\ &= \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(\mathbf{u}_{-1})} \sum_{n > k_1 > \dots > k_m \geq 1} \frac{q^{k_1(i+1+\tilde{p}_1)} A_{n-1, k_1}}{[k_1]^{2i+2+p_1}} \prod_{j=2}^m \frac{q^{\tilde{p}_j k_j} (1 + q^{k_j})}{[k_j]^{p_j}}, \end{aligned}$$

where $A_{n,k}$ is defined in (12). Plugging this into (15) and summing over i by the formula

$$(16) \quad A_{n-1, k} \sum_{i=0}^a \frac{[n]^{2i}}{[k]^{2i}} q^{(k-n)i} = A_{n, k} \left(\frac{[n]^{2a}}{[k]^{2a}} q^{(k-n)a} - \frac{[k]^2}{[n]^2} q^{n-k} \right),$$

we obtain

$$\begin{aligned} H_n^*[\mathbf{s}] = & - \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(\mathbf{u}_{-1})} \sum_{n \geq k_1 > \dots > k_m \geq 1} \frac{q^{k_1(\tilde{p}_1+a+1)} A_{n,k_1}}{[k_1]^{2a+2+p_1}} \prod_{j=2}^m \frac{q^{\tilde{p}_j k_j} (1+q^{k_j})}{[k_j]^{p_j}} \\ & + \frac{q^{n(a+1)}}{[n]^{2a+2}} \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(\mathbf{u}_{-1})} \sum_{n \geq k_1 > \dots > k_m \geq 1} \frac{q^{k_1 \tilde{p}_1} A_{n,k_1}}{[k_1]^{p_1}} \prod_{j=2}^m \frac{q^{\tilde{p}_j k_j} (1+q^{k_j})}{[k_j]^{p_j}} + \frac{q^{n(a+1)}}{[n]^{2a+c}}, \end{aligned}$$

which implies

$$H_n^*[\mathbf{s}] = - \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(\mathbf{s})} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] + \frac{q^{n(a+1)}}{[n]^{2a+2}} \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(\hat{\mathbf{u}}_{-1})} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] + \frac{q^{n(a+1)}}{[n]^{2a+c}},$$

where $\Pi(\hat{\mathbf{u}}_{-1}) = \{\bar{0} \circ 1^{\circ(c-2)}; 0^{\circ(c-1)}; 1 \circ \theta^{\circ(c-2)}\}$. Hence the theorem follows from Corollary 3.3 immediately by replacing c by $c-2$ there. \square

5. MHS and MZSV identities: general case

In this section, we prove some general rules which explain what to expect when we add strings $(\{2\}^a, \{1\}^l)$ or $(\{2\}^b, c)$ to a string of positive integer arguments. This allows us to extend expansion formulas from the three base cases (8), (9) and (14) to every string that contains an arbitrary number of repetitions of $(\{2\}^b, c)$, $(c, \{1\}^l)$ ($b \geq 0$, $c \geq 3$, $l \geq 1$) and $(\{2\}^a, \{1\}^l)$ ($a \geq 0$ except at the leading position when $a \geq 1$). For example, $(3, 1, 2, 7, 1, 1, 5, 2, 2, 4)$ can be written as $(\{2\}^0, 3, \{2\}^0, 1, \{2\}^1, 7, \{2\}^0, \{1\}^2, \{2\}^0, 5, \{2\}^2, 4)$.

For any string $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_m)$, we set $\boldsymbol{\lambda}^\circ = \lambda_1 \circ \dots \circ \lambda_m$ and $\boldsymbol{\lambda}_{\widehat{1}} = (\lambda_2, \dots, \lambda_m)$.

Theorem 5.1. *Let $n \in \mathbb{N}$ and $\mathbf{s} = (s_1, \dots, s_d)$ be a string of positive integers. Set $\iota_{\mathbf{s}} = 1$ if $s_d = 1$ and $\iota_{\mathbf{s}} = -1$ if $s_d > 1$. Suppose \mathbf{s} uniquely determines the triple of mollifiers $[\boldsymbol{\lambda}; \tilde{\boldsymbol{\lambda}}; \tilde{\tilde{\boldsymbol{\lambda}}}] = [\lambda_1, \dots, \lambda_m; \tilde{\lambda}_1, \dots, \tilde{\lambda}_m; \tilde{\tilde{\lambda}}_1, \dots, \tilde{\tilde{\lambda}}_m]$ satisfying*

$$(17) \quad \tilde{\tilde{\lambda}}_1 \boxplus \dots \boxplus \tilde{\tilde{\lambda}}_j \in \{1, 2\} \quad \forall j \geq 1,$$

such that there is an expansion of the form

$$H_n^*[\mathbf{s}] = \iota_{\mathbf{s}} \mathcal{H}_n^\sharp[\boldsymbol{\lambda}; \tilde{\boldsymbol{\lambda}}; \tilde{\tilde{\boldsymbol{\lambda}}}].$$

Then for any integers $a, b \geq 0$ and $c \geq 3$, we have

$$(18) \quad H_n^{\star}[\{2\}^a, \mathbf{s}] = \iota_{\mathbf{s}} \mathcal{H}_n^{\sharp}[2a \oplus \lambda_1, \boldsymbol{\lambda}_{\hat{1}}; a + \tilde{\lambda}_1, \tilde{\boldsymbol{\lambda}}_{\hat{1}}; \tilde{\boldsymbol{\lambda}}],$$

$$(19) \quad H_n^{\star}[\{2\}^a, 1, \mathbf{s}] = \iota_{\mathbf{s}} \mathcal{H}_n^{\sharp}[2a + 1, \boldsymbol{\lambda}; a + 1, \tilde{\boldsymbol{\lambda}}; 2, \tilde{\lambda}_1 \boxplus -2, \tilde{\boldsymbol{\lambda}}_{\hat{1}}],$$

$$(20) \quad H_n^{\star}[\{2\}^b, c, \mathbf{s}] = \iota_{\mathbf{s}} \mathcal{H}_n^{\sharp}[\overline{2b+2}, \{1\}^{c-3}, \lambda_1 \oplus \bar{1}, \boldsymbol{\lambda}_{\hat{1}}; \\ b+1, \{0\}^{c-3}, \tilde{\boldsymbol{\lambda}}; 1, \{\theta\}^{c-3}, \tilde{\lambda}_1 \boxplus -1, \tilde{\boldsymbol{\lambda}}_{\hat{1}}].$$

Moreover, in all the index sets appearing on the right hand side above, the third components still satisfy (17).

Notice that condition (17) essentially guarantees that all the triples of mollifiers considered in the paper are admissible.

Proof. Set $\Pi(\mathbf{s}) = \{\boldsymbol{\lambda}^{\circ}; \tilde{\boldsymbol{\lambda}}^{\circ}; \tilde{\boldsymbol{\lambda}}^{\circ}\}$. The proof of the identities is by induction on $n+a$ or $n+b$. When $n=1$ the theorem is clear. Assume formulas (18) and (19) are true for all $a+n \leq N$ where $N \geq 2$. Suppose now we have $n \geq 2$ and $n+a = N+1$. Set

$$\begin{aligned} \Pi_{\mathbf{2}^a} &= \{(2a \oplus \lambda_1) \circ \boldsymbol{\lambda}_{\hat{1}}^{\circ}; (a + \tilde{\lambda}_1) \circ \tilde{\boldsymbol{\lambda}}_{\hat{1}}^{\circ}; \tilde{\boldsymbol{\lambda}}^{\circ}\}, \\ \Pi_{\mathbf{2}^a 1} &= \{(2a+1) \circ \boldsymbol{\lambda}^{\circ}; (a+1) \circ \tilde{\boldsymbol{\lambda}}^{\circ}; 2 \circ (\tilde{\lambda}_1 \boxplus -2) \circ (\tilde{\boldsymbol{\lambda}}_{\hat{1}}^{\circ})^{\circ}\}, \\ \Pi_{\mathbf{2}^b c} &= \{(\overline{2b+2}) \circ 1^{\circ(c-3)} \circ (\lambda_1 \oplus \bar{1}) \circ \boldsymbol{\lambda}_{\hat{1}}^{\circ}; \\ &\quad (b+1) \circ 0^{\circ(c-3)} \circ \tilde{\boldsymbol{\lambda}}^{\circ}; 1 \circ \theta^{\circ(c-3)} \circ (\tilde{\lambda}_1 \boxplus -1) \circ (\tilde{\boldsymbol{\lambda}}_{\hat{1}}^{\circ})^{\circ}\}. \end{aligned}$$

We start proving the first identity. By definition, we have

$$H_n^{\star}[\{2\}^a, \mathbf{s}] = \sum_{i=1}^a \frac{q^{n(a-i)}}{[n]^{2a-2i}} H_{n-1}^{\star}[\{2\}^i, \mathbf{s}] + \frac{q^{na}}{[n]^{2a}} H_n^{\star}[\mathbf{s}].$$

Applying induction assumption, we obtain

$$\begin{aligned} (21) \quad \iota_{\mathbf{s}} H_n^{\star}[\{2\}^a, \mathbf{s}] &= \sum_{i=1}^a \frac{q^{n(a-i)}}{[n]^{2a-2i}} \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\mathbf{p}}) \in \Pi_{\mathbf{2}^i}} \mathcal{H}_{n-1}[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\mathbf{p}}] \\ &\quad + \frac{q^{na}}{[n]^{2a}} \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\mathbf{p}}) \in \Pi(\mathbf{s})} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\mathbf{p}}]. \end{aligned}$$

Expanding the inner sum

$$\begin{aligned}
& \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_{2^i}} \mathcal{H}_{n-1}[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] \\
&= \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(\mathbf{s})} \mathcal{H}_{n-1}[2i \oplus p_1, \mathbf{p}_1; i + \tilde{p}_1, \tilde{\mathbf{p}}_1; \tilde{\tilde{\mathbf{p}}}] \\
&= \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(\mathbf{s})} \sum_{n > k_1 > \dots > k_r \geq 1} \frac{\begin{bmatrix} n-1 \\ k_1 \end{bmatrix}}{\begin{bmatrix} n-1+k_1 \\ k_1 \end{bmatrix}} \frac{q^{ik_1}}{[k_1]^{2i}} \prod_{j=1}^r \frac{q^{\tilde{p}_j k_j + Q(\tilde{\tilde{\mathbf{p}}}_j, k_j)} (1 + q^{k_j})}{\text{sgn}(p_j)^{k_j} [k_j]^{|p_j|}}
\end{aligned}$$

and summing over i in (21), we obtain

$$\begin{aligned}
\iota_{\mathbf{s}} H_n^*[\{2\}^a, \mathbf{s}] &= \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(\mathbf{s})} \sum_{n \geq k_1 > \dots > k_r \geq 1} \frac{\begin{bmatrix} n \\ k_1 \end{bmatrix}}{\begin{bmatrix} n+k_1 \\ k_1 \end{bmatrix}} \left(\frac{q^{ak_1}}{[k_1]^{2a}} - \frac{q^{an}}{[n]^{2a}} \right) \\
&\quad \times \prod_{j=1}^r \frac{q^{\tilde{p}_j k_j + Q(\tilde{\tilde{\mathbf{p}}}_j, k_j)} (1 + q^{k_j})}{\text{sgn}(p_j)^{k_j} [k_j]^{|p_j|}} \\
&\quad + \frac{q^{na}}{[n]^{2a}} \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(\mathbf{s})} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}],
\end{aligned}$$

which implies (18) by definition and straightforward cancellation. Similarly, for the second identity, we have by definition

$$H_n^*[\{2\}^a, 1, \mathbf{s}] = \sum_{i=0}^a \frac{q^{n(a-i)}}{[n]^{2a-2i}} H_{n-1}^*[\{2\}^i, 1, \mathbf{s}] + \frac{q^{n(a+1)}}{[n]^{2a+1}} H_n^*[\mathbf{s}].$$

Applying induction assumption, we obtain

$$\begin{aligned}
(22) \quad \iota_{\mathbf{s}} H_n^*[\{2\}^a, 1, \mathbf{s}] &= \sum_{i=0}^a \frac{q^{n(a-i)}}{[n]^{2a-2i}} \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_{2^i}} \mathcal{H}_{n-1}[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] \\
&\quad + \frac{q^{n(a+1)}}{[n]^{2a+1}} \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(\mathbf{s})} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}].
\end{aligned}$$

Setting $\Pi_0 = \{\boldsymbol{\lambda}^\circ; \tilde{\boldsymbol{\lambda}}^\circ; (\tilde{\lambda}_1 \boxplus -2) \circ (\tilde{\boldsymbol{\lambda}}_{\hat{1}}^\circ)\}$, we have

$$\begin{aligned}
& \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_{2^{i_1}}} \mathcal{H}_{n-1}[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] \\
&= \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_0} \mathcal{H}_{n-1}[2i+1, \mathbf{p}; i+1, \tilde{\mathbf{p}}; 2, \tilde{\tilde{\mathbf{p}}}] \\
&\quad + \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_0} \mathcal{H}_{n-1}[(2i+1) \oplus p_1, \mathbf{p}_{\hat{1}}; i+1+\tilde{p}_1, \tilde{\mathbf{p}}_{\hat{1}}; 2 \boxplus \tilde{\tilde{p}}_1, \tilde{\tilde{\mathbf{p}}}_{\hat{1}}] \\
&= \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_0} \left(\sum_{n > k_0 > k_1 > \dots > k_m \geq 1} \frac{\begin{bmatrix} n-1 \\ k_0 \end{bmatrix}}{\begin{bmatrix} n-1+k_0 \\ k_0 \end{bmatrix}} \frac{(1+q^{k_0})q^{(i+1)k_0+k_0(k_0-1)}}{[k_0]^{2i+1}} \right. \\
&\quad \times \prod_{j=1}^m \frac{q^{\tilde{p}_j k_j + Q(\tilde{\tilde{p}}_j, k_j)} (1+q^{k_j})}{\operatorname{sgn}(p_j)^{k_j} [k_j]^{|p_j|}} \\
&\quad + \sum_{n > k_1 > \dots > k_m \geq 1} \frac{\begin{bmatrix} n-1 \\ k_1 \end{bmatrix}}{\begin{bmatrix} n-1+k_1 \\ k_1 \end{bmatrix}} \frac{(1+q^{k_1})q^{(i+1+\tilde{p}_1)k_1+Q(2 \boxplus \tilde{\tilde{p}}_1, k_1)}}{\operatorname{sgn}(p_1)^{k_1} [k_1]^{2i+1+|p_1|}} \\
&\quad \left. \times \prod_{j=2}^m \frac{q^{\tilde{p}_j k_j + Q(\tilde{\tilde{p}}_j, k_j)} (1+q^{k_j})}{\operatorname{sgn}(p_j)^{k_j} [k_j]^{|p_j|}} \right).
\end{aligned}$$

Substituting the above expression into (22) and summing over i by (16), we obtain

$$\begin{aligned}
& \iota_s H_n^\star[\{2\}^a, 1, \mathbf{s}] - \frac{q^{n(a+1)}}{[n]^{2a+1}} \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(\mathbf{s})} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] \\
&= \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_0} \left(\sum_{n \geq k_0 > k_1 > \dots > k_m \geq 1} \frac{\begin{bmatrix} n \\ k_0 \end{bmatrix}}{\begin{bmatrix} n+k_0 \\ k_0 \end{bmatrix}} \frac{(1+q^{k_0})q^{(a+1)k_0+k_0(k_0-1)}}{[k_0]^{2a+1}} \right. \\
&\quad \times \prod_{j=1}^m \frac{q^{\tilde{p}_j k_j + Q(\tilde{\tilde{p}}_j, k_j)} (1+q^{k_j})}{\operatorname{sgn}(p_j)^{k_j} [k_j]^{|p_j|}} \\
&\quad - \frac{q^{n(a+1)}}{[n]^{2a+2}} \sum_{n \geq k_0 > k_1 > \dots > k_m \geq 1} \frac{\begin{bmatrix} n \\ k_0 \end{bmatrix}}{\begin{bmatrix} n+k_0 \\ k_0 \end{bmatrix}} (1+q^{k_0})q^{k_0(k_0-1)} [k_0] \\
&\quad \left. \times \prod_{j=1}^m \frac{q^{\tilde{p}_j k_j + Q(\tilde{\tilde{p}}_j, k_j)} (1+q^{k_j})}{\operatorname{sgn}(p_j)^{k_j} [k_j]^{|p_j|}} \right)
\end{aligned}$$

$$\begin{aligned}
& + \sum_{n \geq k_1 > \dots > k_m \geq 1} \frac{\left[\begin{smallmatrix} n \\ k_1 \end{smallmatrix} \right]}{\left[\begin{smallmatrix} n+k_1 \\ k_1 \end{smallmatrix} \right]} \frac{(1+q^{k_1})q^{(a+1+\tilde{p}_1)k_1+Q(2\boxplus\tilde{p}_1,k_1)}}{\operatorname{sgn}(p_1)^{k_1}[k_1]^{2a+1+|p_1|}} \\
& \quad \times \prod_{j=2}^m \frac{q^{\tilde{p}_j k_j + Q(\tilde{p}_j, k_j)}(1+q^{k_j})}{\operatorname{sgn}(p_j)^{k_j}[k_j]^{|p_j|}} \\
& - \frac{q^{n(a+1)}}{[n]^{2a+2}} \sum_{n \geq k_1 > \dots > k_m \geq 1} \frac{\left[\begin{smallmatrix} n \\ k_1 \end{smallmatrix} \right]}{\left[\begin{smallmatrix} n+k_1 \\ k_1 \end{smallmatrix} \right]} \frac{(1+q^{k_1})q^{\tilde{p}_1 k_1 + Q(2\boxplus\tilde{p}_1,k_1)}}{\operatorname{sgn}(p_1)^{k_1}[k_1]^{(|p_1|-1)}} \\
& \quad \times \prod_{j=2}^m \frac{q^{\tilde{p}_j k_j + Q(\tilde{p}_j, k_j)}(1+q^{k_j})}{\operatorname{sgn}(p_j)^{k_j}[k_j]^{|p_j|}} \Big).
\end{aligned}$$

Noticing that the first and third sums on the right-hand side of the above add up to

$$\sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_{2^{a_1}}} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}],$$

we have

$$\begin{aligned}
& \iota_s H_n^\star[\{2\}^a, 1, s] - \frac{q^{n(a+1)}}{[n]^{2a+1}} \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(s)} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] \\
& = \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_{2^{a_1}}} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] \\
& \quad - \frac{q^{n(a+1)}}{[n]^{2a+2}} \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_0} \sum_{n \geq k_1 > \dots > k_m \geq 1} \left(\prod_{j=1}^m \frac{q^{\tilde{p}_j k_j + Q(\tilde{p}_j, k_j)}(1+q^{k_j})}{\operatorname{sgn}(p_j)^{k_j}[k_j]^{|p_j|}} \right. \\
& \quad \times \sum_{k_0=k_1+1}^n \frac{\left[\begin{smallmatrix} n \\ k_0 \end{smallmatrix} \right]}{\left[\begin{smallmatrix} n+k_0 \\ k_0 \end{smallmatrix} \right]} \frac{(1+q^{k_0})q^{k_0(k_0-1)}}{[k_0]^{-1}} \\
& \quad + \frac{\left[\begin{smallmatrix} n \\ k_1 \end{smallmatrix} \right]}{\left[\begin{smallmatrix} n+k_1 \\ k_1 \end{smallmatrix} \right]} \frac{(1+q^{k_1})q^{\tilde{p}_1 k_1 + Q(2\boxplus\tilde{p}_1,k_1)}}{\operatorname{sgn}(p_1)^{k_1}[k_1]^{(|p_1|-1)}} \\
& \quad \times \left. \prod_{j=2}^m \frac{q^{\tilde{p}_j k_j + Q(\tilde{p}_j, k_j)}(1+q^{k_j})}{\operatorname{sgn}(p_j)^{k_j}[k_j]^{|p_j|}} \right).
\end{aligned}$$

Summing the multiple sum in the above over k_0 by (11) and noticing that for $(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_0$, by (17), the first component \tilde{p}_1 can take only values -1

and 0, we obtain with the help of (7) that

$$\iota_s H_n^{\star}[\{2\}^a, 1, s] = \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_{2^a 1}} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}].$$

This proves identity (19) by induction.

Finally, to prove (20), we proceed by induction on $n + b$. Assume formula (20) is true for all $b + n \leq N$. Now suppose $b + n = N + 1$. By definition, we have

$$H_n^{\star}[\{2\}^b, c, s] = \sum_{i=0}^b \frac{q^{n(b-i)}}{[n]^{2b-2i}} H_{n-1}^{\star}[\{2\}^i, c, s] + \frac{q^{n(b+1)}}{[n]^{2b+c}} H_n^{\star}[s].$$

By the induction assumption, we see that

$$(23) \quad \begin{aligned} \iota_s H_n^{\star}[\{2\}^b, c, s] &= \sum_{i=0}^b \frac{q^{n(b-i)}}{[n]^{2b-2i}} \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_{2^i c}} \mathcal{H}_{n-1}[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] \\ &\quad + \frac{q^{n(b+1)}}{[n]^{2b+c}} \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(s)} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] . \end{aligned}$$

Setting

$$\Pi_1 = \{0 \circ 1^{\circ(c-3)} \circ (\lambda_1 \oplus \bar{1}) \circ \boldsymbol{\lambda}_1^{\circ}; 0^{\circ(c-2)} \circ \tilde{\boldsymbol{\lambda}}^{\circ}; \theta^{\circ(c-2)} \circ (\tilde{\lambda}_1 \boxplus -1) \circ (\tilde{\lambda}_1^{\circ})^{\circ}\},$$

we have

$$\begin{aligned} &\sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_{2^i c}} \mathcal{H}_{n-1}[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] \\ &= \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_1} \mathcal{H}_{n-1}[2i+2 \oplus p_1, \mathbf{p}_1; i+1+\tilde{p}_1, \tilde{\mathbf{p}}_1; 1 \boxplus \tilde{\tilde{p}}_1, \tilde{\tilde{\mathbf{p}}}_1] \\ &= \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_1} \sum_{n > k_1 > \dots > k_r \geq 1} \frac{\binom{n-1}{k_1}}{\binom{n-1+k_1}{k_1}} \frac{q^{k_1(i+1+\tilde{p}_1)+Q(1 \boxplus \tilde{\tilde{p}}_1, k_1)}}{(-\text{sgn}(p_1))^{k_1} [k_1]^{2i+2+|p_1|}} \\ &\quad \times \prod_{j=2}^r \frac{q^{\tilde{p}_j k_j + Q(\tilde{\tilde{p}}_j, k_j)} (1 + q^{k_j})}{\text{sgn}(p_j)^{k_j} [k_j]^{|p_j|}}. \end{aligned}$$

Plugging this into (23) and summing over i by (16), we obtain

$$\begin{aligned} \iota_s H_n^{\star}[\{2\}^b, c, s] &= \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_1} \mathcal{H}_n[(\overline{2b+2}) \oplus p_1, \mathbf{p}_{\widehat{1}}; b+1+\tilde{p}_1, \tilde{\mathbf{p}}_{\widehat{1}}; 1 \boxplus \tilde{\tilde{p}}_1, \tilde{\tilde{\mathbf{p}}}_{\widehat{1}}] \\ &+ \frac{q^{n(b+1)}}{[n]^{2b+c}} \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(s)} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] - \frac{q^{n(b+1)}}{[n]^{2b+2}} \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_1} \mathcal{H}_n[\bar{p}_1, \mathbf{p}_{\widehat{1}}; \tilde{\mathbf{p}}; 1 \boxplus \tilde{\tilde{\mathbf{p}}}_{\widehat{1}}], \end{aligned}$$

which implies

$$\begin{aligned} (24) \quad \iota_s H_n^{\star}[\{2\}^b, c, s] &= \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_{2^b c}} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] - \frac{q^{n(b+1)}}{[n]^{2b+2}} \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_2} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] \\ &+ \frac{q^{n(b+1)}}{[n]^{2b+c}} \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(s)} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}], \end{aligned}$$

where

$$\begin{aligned} \Pi_2 = \{ &\bar{0} \circ 1^{\circ(c-3)} \circ (\lambda_1 \oplus \bar{1}) \circ \lambda_{\widehat{1}}^{\circ}; 0^{\circ(c-2)} \circ \tilde{\lambda}_{\widehat{1}}^{\circ}; \\ &1 \circ \theta^{\circ(c-3)} \circ (\tilde{\tilde{\lambda}}_1 \boxplus (-1)) \circ (\tilde{\tilde{\lambda}}_{\widehat{1}})^{\circ} \}. \end{aligned}$$

Expanding the second sum from (24), we have

$$\begin{aligned} &\sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_2} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] \\ &= \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_3} \sum_{\substack{\mathbf{w} = \bar{0} \circ 1^{\circ(c-3)} \circ (p_1 \oplus \bar{1}) \\ \tilde{\mathbf{w}} = 0^{\circ(c-2)} \circ \tilde{p}_1; \tilde{\tilde{\mathbf{w}}} = 1 \circ \theta^{\circ(c-3)} \circ \tilde{\tilde{p}}_1}} \mathcal{H}_n[\mathbf{w}, \mathbf{p}_{\widehat{1}}; \tilde{\mathbf{w}}, \tilde{\mathbf{p}}_{\widehat{1}}; \tilde{\tilde{\mathbf{w}}}, \tilde{\tilde{\mathbf{p}}}_{\widehat{1}}], \end{aligned}$$

where $\Pi_3 = \{\lambda^{\circ}; \tilde{\lambda}^{\circ}; (\tilde{\lambda}_1 \boxplus (-1)) \circ (\tilde{\lambda}_{\widehat{1}})^{\circ}\}$. Applying Lemma 3.2 to the inner sum with $a = p_1$, $b = \tilde{p}_1$, $r = \tilde{\tilde{p}}_1$, c replaced by $c-2$, and $\mathbf{x} = \mathbf{p}_{\widehat{1}}$, $\mathbf{y} = \tilde{\mathbf{p}}_{\widehat{1}}$, $\mathbf{z} = \tilde{\tilde{\mathbf{p}}}_{\widehat{1}}$, we obtain

$$\begin{aligned} (25) \quad \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_2} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] &= \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi_3} \frac{1}{[n]^{c-2}} \mathcal{H}_n[p_1, \mathbf{p}_{\widehat{1}}; \tilde{\mathbf{p}}; \tilde{\tilde{p}}_1 \boxplus 1, \tilde{\tilde{\mathbf{p}}}_{\widehat{1}}] \\ &= \frac{1}{[n]^{c-2}} \sum_{(\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}) \in \Pi(s)} \mathcal{H}_n[\mathbf{p}; \tilde{\mathbf{p}}; \tilde{\tilde{\mathbf{p}}}] . \end{aligned}$$

To justify the last equality above, we need to show that for the components of $\tilde{\lambda}$ satisfying (17) we have $\tilde{\lambda}_1 \boxplus (-1) \boxplus 1 = \tilde{\lambda}_1$ and for any $j \geq 2$,

$$\tilde{\lambda}_1 \boxplus (-1) \boxplus \tilde{\lambda}_2 \boxplus \cdots \boxplus \tilde{\lambda}_j \boxplus 1 = \tilde{\lambda}_1 \boxplus \tilde{\lambda}_2 \boxplus \cdots \boxplus \tilde{\lambda}_j.$$

These can be proved by using the projection π of Lemma 2.2 and the fact that $\pi(\tilde{\lambda}_1 \boxplus (-1) \boxplus \cdots \boxplus \tilde{\lambda}_j \boxplus 1) = \pi(\tilde{\lambda}_1 \boxplus \cdots \boxplus \tilde{\lambda}_j) \in \{1, 2\}$ by (17).

Now by (24) and (25), we see that (20) is true when $n + b = N + 1$. We have completed the proof of the theorem. \square

Repeatedly applying the theorem, we quickly find

Corollary 5.2. *Keep the same notation as in Theorem 5.1. Then for any integers $a, b \geq 0$, $l \geq 1$ and $c \geq 3$, we have*

$$(26) \quad H_n^*[\{2\}^a, \{1\}^l, \mathbf{s}] = \iota_{\mathbf{s}} \mathcal{H}_n^\# [2a+1, \{1\}^{l-1}, \boldsymbol{\lambda}; a+1, \{1\}^{l-1}, \tilde{\boldsymbol{\lambda}}; \\ 2, \{0\}^{l-1}, \tilde{\lambda}_1 \boxplus -2, \tilde{\boldsymbol{\lambda}}_1]$$

and

$$H_n^*[\{2\}^b, c, \{2\}^a, \{1\}^l, \mathbf{s}] = \iota_{\mathbf{s}} \mathcal{H}_n^\# [\overline{2b+2}, \{1\}^{c-3}, \overline{2a+2}, \{1\}^{l-1}, \boldsymbol{\lambda}; \\ b+1, \{0\}^{c-3}, a+1, \{1\}^{l-1}, \tilde{\boldsymbol{\lambda}}; \\ 1, \{\theta\}^{c-3}, 1, \{0\}^{l-1}, \tilde{\lambda}_1 \boxplus -2, \tilde{\boldsymbol{\lambda}}_1].$$

Proof. Repeatedly applying (19) by attaching $(2^{a_j}, 1)$, $j = 1, \dots, l$ and then setting $a_1 = \dots = a_{l-1} = 0$ and $a_l = a$, we can quickly verify (26). The corollary follows by applying (20) to (26). \square

We may take limit $n \rightarrow \infty$ in (18) of Theorem 5.1 and Corollary 5.2 to obtain identities for q -MZSV.

Theorem 5.3. *Let $\mathbf{s} = (s_1, \dots, s_d) \in \mathbb{N}^d$. Set $\iota_{\mathbf{s}} = 1$ if $s_d = 1$ and $\iota_{\mathbf{s}} = -1$ if $s_d > 1$. Suppose \mathbf{s} uniquely determines $[\boldsymbol{\lambda}; \tilde{\boldsymbol{\lambda}}; \tilde{\boldsymbol{\lambda}}]$ satisfying (17) such that $\zeta^*[\mathbf{s}] = \iota_{\mathbf{s}} \mathfrak{z}^\#[\boldsymbol{\lambda}; \tilde{\boldsymbol{\lambda}}; \tilde{\boldsymbol{\lambda}}]$. Then for any integers $a, b \geq 0$, $l \geq 1$ and $c \geq 3$, we have*

$$\zeta^*[\{2\}^a, \mathbf{s}] = \iota_{\mathbf{s}} \mathfrak{z}^\# [2a \oplus \lambda_1, \boldsymbol{\lambda}_1; a + \tilde{\lambda}_1, \tilde{\boldsymbol{\lambda}}_1; \tilde{\boldsymbol{\lambda}}], \\ \zeta^*[\{2\}^a, \{1\}^l, \mathbf{s}] = \iota_{\mathbf{s}} \mathfrak{z}^\# [2a+1, \{1\}^{l-1}, \boldsymbol{\lambda}; a+1, \{1\}^{l-1}, \tilde{\boldsymbol{\lambda}}; \\ 2, \{0\}^{l-1}, \tilde{\lambda}_1 \boxplus -2, \tilde{\boldsymbol{\lambda}}_1],$$

$$\begin{aligned}\zeta^*[\{2\}^b, c, \mathbf{s}] &= \iota_{\mathbf{s}} \mathfrak{z}^\# [\overline{2b+2}, \{1\}^{c-3}, \lambda_1 \oplus \bar{1}, \tilde{\boldsymbol{\lambda}}_1; b+1, \{0\}^{c-3}, \tilde{\boldsymbol{\lambda}}; \\ &\quad 1, \{\theta\}^{c-3}, \tilde{\tilde{\boldsymbol{\lambda}}}_1 \boxplus -1, \tilde{\tilde{\boldsymbol{\lambda}}}_1], \\ \zeta^*[c, \{1\}^l, \mathbf{s}] &= \iota_{\mathbf{s}} \mathfrak{z}^\# [\overline{2}, \{1\}^{c-3}, \overline{2}, \{1\}^{l-1}, \boldsymbol{\lambda}; 1, \{0\}^{c-3}, \{1\}^l, \tilde{\boldsymbol{\lambda}}; \\ &\quad 1, \{\theta\}^{c-3}, 1, \{0\}^{l-1}, \tilde{\tilde{\boldsymbol{\lambda}}}_1 \boxplus -2, \tilde{\tilde{\boldsymbol{\lambda}}}_1].\end{aligned}$$

Proof. The first three equations are straight-forward. The last one can be obtained by applying the middle two equations successively after setting $a = b = 0$. \square

By letting $q \rightarrow 1$ in Theorem 5.3 we can immediately prove Theorem 1.3 which gives the corresponding general rule for classical MZSV. Of course, to guarantee convergence we need to restrict $a \geq 1$ there.

From Theorem 5.3, we can obtain a general formula for arbitrary q -MZSV.

Theorem 5.4. *Let $a_0, a_j \in \mathbb{N}_0$, $c_j \in \mathbb{N}$ and $c_j \neq 2$ for all $j = 1, \dots, d$. Set $\delta(c) = 1$ if $c = 1$ and $\delta(c) = 0$ if $c \geq 3$. Moreover, put $\{\alpha\}^n = \{\alpha\}^{\max(n, 0)}$. Then we have*

$$\begin{aligned}\zeta^*[\{2\}^{a_0}, c_1, \{2\}^{a_1}, \dots, c_d, \{2\}^{a_d}] &= \pm \mathfrak{z}^\# [B_0, \{1\}^{c_1-3}, B_1, \dots, \{1\}^{c_d-3}, B_d; \\ &\quad \tilde{B}_0, \{0\}^{c_1-3}, \tilde{B}_1, \dots, \{0\}^{c_d-3}, \tilde{B}_d; \\ &\quad \tilde{\tilde{B}}_0, \{\theta\}^{c_1-3}, \tilde{\tilde{B}}_1, \dots, \{\theta\}^{c_d-3}, \tilde{\tilde{B}}_d].\end{aligned}$$

Here the leading sign \pm is $+$ if and only if $a_d = 0$ and $c_d = 1$,

$$\begin{aligned}B_j &= \begin{cases} A_j, & \text{if } A_j \text{ is odd;} \\ \overline{A_j}, & \text{if } A_j \text{ is even,} \end{cases} \\ \text{where } A_j &= \begin{cases} 2a_0 + 2 - \delta(c_1), & \text{if } j = 0; \\ 2a_d + 1 - \delta(c_d), & \text{if } j = d; \\ 2a_j + 3 - \delta(c_j) - \delta(c_{j+1}), & \text{if } 0 < j < d, \end{cases} \\ \tilde{B}_j &= \begin{cases} a_j + 1, & \text{if } 0 \leq j < d; \\ a_d, & \text{if } j = d, \end{cases} \\ \text{and } \tilde{\tilde{B}}_j &= \begin{cases} 1 + \delta(c_1), & \text{if } j = 0; \\ (1 - \delta(c_d)) \boxplus (-1), & \text{if } j = d; \\ (1 - \delta(c_j)) \boxplus (\delta(c_{j+1}) - 1), & \text{if } 0 < j < d. \end{cases}\end{aligned}$$

Moreover, if $a_d = 0$ and $c_d = 1$, then $B_d, \tilde{B}_d, \tilde{\tilde{B}}_d$ are vacuous.

Proof. The theorem can be proved easily by induction on d using Theorem 5.3. We leave the details to the interested reader. \square

By letting $q \rightarrow 1$ in Theorem 5.4, we get Theorem 1.4 which gives the corresponding result for classical MZSV. It is clear that to ensure convergence we need to assume that $a_0 > 0$ or $c_1 \geq 3$.

6. Some applications

The first application gives us the general **2-c-2** ($c \geq 3$) formula. Here the underline means the $(\{2\}^a, c)$ -type string may be repeated an arbitrary number of times where a and c may change in each repetition.

Theorem 6.1. Suppose $\ell \in \mathbb{N}_0$. Let $\mathbf{s} = (\{2\}^{a_1}, c_1, \dots, \{2\}^{a_\ell}, c_\ell, \{2\}^{a_{\ell+1}})$ with $a_j, c_j \in \mathbb{N}_0$ and $c_j \geq 3$ for all $j \geq 1$. Then

$$(27) \quad \begin{aligned} H_n^{\star}[\mathbf{s}] = -\mathcal{H}_n^{\sharp}[\overline{2a_1+2}, \{1\}^{c_1-3}, 2a_2+3, \{1\}^{c_2-3}, \\ \dots, 2a_\ell+3, \{1\}^{c_\ell-3}, 2a_{\ell+1}+1; \\ a_1+1, \{0\}^{c_1-3}, \dots, a_\ell+1, \{0\}^{c_\ell-3}, a_{\ell+1}; \\ 1, \{\theta\}^{c_1+\dots+c_\ell-2\ell}]. \end{aligned}$$

Proof. If $a_{\ell+1} = 0$, then starting from Theorem 4.1 for $H_n^{\star}[\{2\}^{a_\ell}, c_\ell]$ and repeatedly applying (20), we get the above identity. Otherwise, starting from (8) and repeatedly applying the attaching rule (20) we can arrive at (27) immediately. \square

By applying Lemma 2.3 to Theorem 6.1 we immediately get

Corollary 6.2. With the same notation as in Theorem 6.1, we have

$$\zeta^{\star}[\mathbf{s}] = -\zeta^{\sharp}[\overline{2a_1+2}, \{1\}^{c_1-3}, 2a_2+3, \{1\}^{c_2-3}, \dots, 2a_\ell+3, \{1\}^{c_\ell-3}, 2a_{\ell+1}+1; \\ a_1+1, \{0\}^{c_1-3}, \dots, a_\ell+1, \{0\}^{c_\ell-3}, a_{\ell+1}; 1, \{\theta\}^{c_1+\dots+c_\ell-2\ell}].$$

In particular, if $c_1 = c_2 = \dots = c_\ell = 3$, we get a q -analog of the Two-three formula:

$$\zeta^{\star}[\mathbf{s}] = \sum_{\substack{\mathbf{p}=(2a_1+2)\circ(2a_2+3)\circ\dots\circ(2a_\ell+3)\circ(2a_{\ell+1}+1) \\ \tilde{\mathbf{p}}=(a_1+1)\circ\dots\circ(a_\ell+1)\circ(a_{\ell+1})}} \sum_{k_1>\dots>k_m\geq 1} (-1)^{k_1-1} q^{\frac{k_1(k_1-1)}{2}} \prod_{j=1}^m \frac{q^{\tilde{p}_j k_j} (1+q^{k_j})}{[k_j]^{p_j}}.$$

Remark 6.3. When $q \rightarrow 1$ one can recover all the MZSV identities contained in [17].

Now starting from (9) and repeatedly and alternatively applying the attaching rules (19) and (20) we can find the following:

Theorem 6.4. *Suppose $\ell \in \mathbb{N}_0$, $n \in \mathbb{N}$ and $a_0, a_j, b_j, c_j - 3 \in \mathbb{N}_0$ for all $j \geq 1$. Consider the following two possible types of compositions:*

(**2-c-2-1**) :

$$\begin{aligned} \mathbf{s} &= (\{2\}^{b_1}, c_1, \{2\}^{a_1}, 1, \dots, \{2\}^{b_\ell}, c_\ell, \{2\}^{a_\ell}, 1), \quad \ell \in \mathbb{N}, \\ \mathbf{s}' &= (\overline{2b_1 + 2}, \{1\}^{c_1-3}, \overline{2a_1 + 2}, \dots, \overline{2b_\ell + 2}, \{1\}^{c_\ell-3}, \overline{2a_\ell + 2}; \\ &\quad b_1 + 1, \{0\}^{c_1-3}, a_1 + 1, \dots, b_\ell + 1, \{0\}^{c_\ell-3}, a_\ell + 1; \\ &\quad 1, \underbrace{\{\theta\}^{c_1-3}, 1, -1, \{\theta\}^{c_2-3}, 1, \dots, -1, \{\theta\}^{c_\ell-3}, 1}_{\text{appear only if } \ell > 1}. \end{aligned}$$

(**2-1-2-c-2-1**) :

$$\begin{aligned} \mathbf{s} &= (\{2\}^{a_0}, 1, \{2\}^{b_1}, c_1, \{2\}^{a_1}, 1, \dots, \{2\}^{b_\ell}, c_\ell, \{2\}^{a_\ell}, 1), \quad \ell \in \mathbb{N}_0, \\ \mathbf{s}' &= (2a_0 + 1, \overline{2b_1 + 2}, \{1\}^{c_1-3}, \overline{2a_1 + 2}, \dots, \overline{2b_\ell + 2}, \{1\}^{c_\ell-3}, \overline{2a_\ell + 2}; \\ &\quad a_0 + 1, b_1 + 1, \{0\}^{c_1-3}, a_1 + 1, \dots, b_\ell + 1, \{0\}^{c_\ell-3}, a_\ell + 1; \\ &\quad 2, \underbrace{-1, \{\theta\}^{c_1-3}, 1, \dots, -1, \{\theta\}^{c_\ell-3}, 1}_{\text{appear only if } \ell > 0}. \end{aligned}$$

Then in each case we have

$$H_n^{\star}[\mathbf{s}] = \mathcal{H}_n^{\sharp}[\mathbf{s}'].$$

Corollary 6.5. *With the same notation as in Theorem 6.4, we have*

$$\zeta^{\star}[\mathbf{s}] = \mathfrak{z}^{\sharp}[\mathbf{s}'].$$

For example, taking $\ell = 1$ and $c_1 = 3$, we get (cf. [23, (26)] and the identity after it)

$$\begin{aligned} \zeta^{\star}[\{2\}^b, 3, \{2\}^a, 1] &= \mathfrak{z}[2a + 2b + 4; a + b + 2; 2] \\ &\quad + \mathfrak{z}[\overline{2b + 2}, \overline{2a + 2}; b + 1, a + 1; 1, 1] \end{aligned}$$

and

$$\begin{aligned} (28) \quad &\zeta^{\star}[\{2\}^{a_0}, 1, \{2\}^b, 3, \{2\}^{a_1}, 1] \\ &= \mathfrak{z}[2(a_0 + b + a_1) + 5; a_0 + b + a_1 + 3; 2] \\ &\quad + \mathfrak{z}[2a_0 + 1, 2a_1 + 2b + 4; a_0 + 1, a_1 + b + 2; 2, 0] \\ &\quad + \mathfrak{z}[\overline{2a_0 + 2b + 3}, \overline{2a_1 + 2}; a_0 + b + 2, a_1 + 1; 1, 1] \\ &\quad + \mathfrak{z}[2a_0 + 1, \overline{2b + 2}, \overline{2a_1 + 2}; a_0 + 1, b + 1, a_1 + 1; 2, -1, 1]. \end{aligned}$$

We can also get the following identity which is the q -analog of [23, Theorem 6.1(i)].

Corollary 6.6. *Let a, b be two nonnegative integers. Then*

$$\begin{aligned} & \zeta^*[\{2\}^a, 3, \{2\}^b, 1] + \zeta^*[\{2\}^b, 3, \{2\}^a, 1] \\ &= \zeta^*[\{2\}^{a+1}] \zeta^*[\{2\}^{b+1}] \\ &\quad + (1-q)\mathfrak{z}[2a+2b+3; a+b+2; 2]. \end{aligned}$$

Proof. By taking $n \rightarrow \infty$ in (8) and using Lemma 2.3 we get

$$\zeta^*[\{2\}^{a+1}] = \mathfrak{z}[2a+2; a+1; 1] = \sum_{k=1}^{\infty} \frac{q^{(a+1)k+Q(1,k)}(1+q^k)}{[k]^{2a+2}}.$$

Thus

$$\begin{aligned} & \zeta^*[\{2\}^a, 3, \{2\}^b, 1] + \zeta^*[\{2\}^b, 3, \{2\}^a, 1] - \zeta^*[\{2\}^{a+1}] \zeta^*[\{2\}^{b+1}] \\ &= \sum_{k=1}^{\infty} \frac{q^{(a+b+2)k+Q(2,k)}(2(1+q^k) - (1+q^k)^2)}{[k]^{2a+2b+4}} \\ &= \sum_{k=1}^{\infty} \frac{q^{(a+b+2)k+Q(2,k)}(1+q^k)(1-q^k)}{[k]^{2a+2b+4}} \\ &= (1-q)\mathfrak{z}[2a+2b+3; a+b+2; 2] \end{aligned}$$

as desired. \square

If we start with (8) and repeatedly and alternatively apply the attaching rules (19) and (20) we can get:

Theorem 6.7. *Suppose $\ell \in \mathbb{N}_0$, $n, a_{\ell+1} \in \mathbb{N}$, and $a_0, a_j, b_j, c_j - 3 \in \mathbb{N}_0$ for all $1 \leq j \leq \ell$. Consider the following two possible types of compositions:*

$$\begin{aligned} & (\underline{\textbf{2-c-2-1-2}}) : \\ & \mathbf{s} = (\{2\}^{b_1}, c_1, \{2\}^{a_1}, 1, \dots, \{2\}^{b_\ell}, c_\ell, \{2\}^{a_\ell}, 1, \{2\}^{a_{\ell+1}}), \quad \ell \in \mathbb{N}, \\ & \mathbf{s}' = (\overline{2b_1+2}, \{1\}^{c_1-3}, \overline{2a_1+2}, \dots, \overline{2b_\ell+2}, \{1\}^{c_\ell-3}, \overline{2a_\ell+2}, \overline{2a_{\ell+1}}; \\ & \quad b_1+1, \{0\}^{c_1-3}, a_1+1, \dots, b_\ell+1, \{0\}^{c_\ell-3}, a_\ell+1, a_{\ell+1}; \\ & \quad 1, \{\theta\}^{c_1-3}, 1, \underbrace{-1, \{\theta\}^{c_2-3}, 1, \dots, -1, \{\theta\}^{c_\ell-3}, 1, -1}_{\text{appear only if } \ell > 1}). \end{aligned}$$

(2-1-2-c-2-1-2) :

$$\begin{aligned} \mathbf{s} &= (\{2\}^{a_0}, 1, \{2\}^{b_1}, c_1, \{2\}^{a_1}, 1, \dots, \{2\}^{b_\ell}, c_\ell, \{2\}^{a_\ell}, 1, \{2\}^{a_{\ell+1}}), \quad \ell \in \mathbb{N}_0, \\ \mathbf{s}' &= (2a_0 + 1, \overline{2b_1 + 2}, \{1\}^{c_1 - 3}, \overline{2a_1 + 2}, \dots, \overline{2b_\ell + 2}, \{1\}^{c_\ell - 3}, \overline{2a_\ell + 2}, \overline{2a_{\ell+1}}; \\ &\quad a_0 + 1, b_1 + 1, \{0\}^{c_1 - 3}, a_1 + 1, \dots, b_\ell + 1, \{0\}^{c_\ell - 3}, a_\ell + 1, a_{\ell+1}; \\ &\quad 2, \underbrace{-1, \{\theta\}^{c_1 - 3}, 1, \dots, -1, \{\theta\}^{c_\ell - 3}, 1, -1}_{\text{appear only if } \ell > 0}. \end{aligned}$$

Then in each case we have

$$H_n^{\star}[\mathbf{s}] = -\mathcal{H}_n^{\sharp}[\mathbf{s}'].$$

By taking $n \rightarrow \infty$ we have

Corollary 6.8. *Let notation be the same as in Theorem 6.7. Then*

$$\zeta^{\star}[\mathbf{s}] = -\mathfrak{z}^{\sharp}[\mathbf{s}'].$$

For example, taking $\ell = 1$ and $c_1 = 3$, we get in case **(2-c-2-1-2)**

$$\begin{aligned} \zeta^{\star}[\{2\}^b, 3, \{2\}^{a_1}, 1, \{2\}^{a_2}] &= -\mathfrak{z}[\overline{2a_1 + 2b + 2a_2 + 4}; a_1 + b + a_2 + 2; 1] \\ &\quad - \mathfrak{z}[\overline{2b + 2}, 2a_1 + 2a_2 + 2; b + 1, a_1 + a_2 + 1; 1, \theta] \\ &\quad - \mathfrak{z}[2a_1 + 2b + 4, \overline{2a_2}; a_1 + b + 2, a_2; 2, -1] \\ &\quad - \mathfrak{z}[\overline{2b + 2}, \overline{2a_1 + 2}, \overline{2a_2}; b + 1, a_1 + 1, a_2; 1, 1, -1]. \end{aligned}$$

By taking $q \rightarrow 1$ this yields the identity on the bottom of [23, p. 12].

As a non-trivial example of Theorem 5.3 we may attach a string of type $(2^a, 1)$ to the front of the already treated type $(\{2\}^b, 1, \{2\}^c, 3, \{2\}^d, 1)$ given by (28) and get the following q -MZSV identity: for any nonnegative integers a, b, c, d

$$\begin{aligned} (29) \quad \zeta^{\star}[\{2\}^a, 1, \{2\}^b, 1, \{2\}^c, 3, \{2\}^d, 1] \\ &= \mathfrak{z}[2a + 2b + 2c + 2d + 6; a + b + c + d + 4; 2] \\ &\quad + \mathfrak{z}[2a + 1, 2b + 2c + 2d + 5; a + 1, b + c + d + 3; 2, 0] \\ &\quad + \mathfrak{z}[2a + 2b + 2, 2c + 2d + 4; a + b + 2, c + d + 2; 2, 0] \\ &\quad + \mathfrak{z}[\overline{2a + 2b + 2c + 4}, \overline{2d + 2}; a + b + c + 3, d + 1; 1, 1] \\ &\quad + \mathfrak{z}[2a + 1, 2b + 1, 2c + 2d + 4; a + 1, b + 1, c + d + 2; 2, 0, 0] \\ &\quad + \mathfrak{z}[2a + 1, \overline{2b + 2c + 3}, \overline{2d + 2}; a + 1, b + c + 2, d + 1; 2, -1, 1] \\ &\quad + \mathfrak{z}[2a + 2b + 2, \overline{2c + 2}, \overline{2d + 2}; a + b + 2, c + 1, d + 1; 2, -1, 1] \\ &\quad + \mathfrak{z}[2a + 1, 2b + 1, \overline{2c + 2}, \overline{2d + 2}; a + 1, b + 1, c + 1, d + 1; 2, 0, -1, 1]. \end{aligned}$$

By taking $q \rightarrow 1$ in (29) we discover the classical MZSV identity (3) in the introduction, which has not been proved before.

References

- [1] J. M. Borwein, P. Lisonek, and P. Irvine, *An interface for evaluation of Euler sums*, available online at <http://wayback.cecm.sfu.ca/projects/EZFace>.
- [2] D. J. Broadhurst, *Conjectured enumeration of irreducible multiple zeta values, from knots and Feynman diagrams*, preprint, arXiv: [hep-th/9612012](https://arxiv.org/abs/hep-th/9612012).
- [3] D. M. Bradley, *Multiple q -zeta values*, J. of Algebra **283** (2005), 752–798.
- [4] F. Brown, *Mixed Tate motives over \mathbb{Z}* , Ann. Math. **175** (2012), no. 2, 949–976.
- [5] F. Brown and O. Schnetz, *Single-valued multiple polylogarithms and a proof of the zig-zag conjecture*, J. Number Theory **148** (2015), no. 3, 478–506.
- [6] H. Gangl, M. Kaneko, and D. Zagier, *Double zeta values and modular forms*, in: Automorphic Forms and Zeta Functions, S. Böcherer et al. eds., World Scientific, Hackensack, NJ (2006), 71–106.
- [7] C. Glanois, *Periods of the motivic fundamental groupoid of $\mathbb{P}^1 \setminus \{0, \mu_N, \infty\}$* , Ph.D. thesis, Université Pierre et Marie Curie, September 2015, available at <http://www2.mathematik.hu-berlin.de/~kreimer/publications/>.
- [8] A. B. Goncharov, *Multiple ζ -values, Galois groups, and geometry of modular varieties*, in: European Congress of Mathematics (Barcelona, 2000), Vol. I, Progr. Math. **201**, Birkhäuser, Basel (2001), 361–392.
- [9] A. B. Goncharov and Y. I. Manin, *Multiple ζ -motives and moduli spaces $M_{0,n}$* , Compositio Math. **140** (2004), 1–14.
- [10] Kh. Hessami Pilehrood and T. Hessami Pilehrood, *On q -analogues of two-one formulas for multiple harmonic sums and multiple zeta star values*, Monatsh. Math. **176** (2015), 275–291.
- [11] Kh. Hessami Pilehrood, T. Hessami Pilehrood, and R. Tauraso, *New properties of multiple harmonic sums modulo p and p -analogues of*

- Leshchiner's series*, Trans. Amer. Math. Soc. **366** (2014), no. 6, 3131–3159.
- [12] M. E. Hoffman, *The algebra of multiple harmonic series*, J. Alg. **194** (1997), 477–495.
 - [13] M. E. Hoffman, *The Hopf algebra structure of multiple harmonic sums*, Nuclear Phys. B (Proc. Suppl.) **135** (2004), 214–219.
 - [14] M. E. Hoffman, *Algebraic aspects of multiple zeta values*, in: Zeta Functions, Topology and Quantum Physics, T. Aoki et. al. (eds.), Developments in Math. **14**, Springer, New York (2005), 51–74.
 - [15] K. Ihara, J. Kajikawa, Y. Ohno, and J. Okuda, *Multiple zeta values vs. multiple zeta-star values*, J. Algebra **332** (2011), 187–208.
 - [16] M. Kaneko, N. Kurokawa, and M. Wakayama, *A variation of Euler's approach to values of the Riemann zeta function*, Kyushu J. Math. **57** (2003), 175–192.
 - [17] E. Linebarger and J. Zhao, *A family of multiple harmonic sum and multiple zeta star value identities*, Mathematika **61** (2015), no. 1, 63–71. [arXiv:1304.3927](https://arxiv.org/abs/1304.3927).
 - [18] Y. Ohno and W. Zudilin, *Zeta stars*, Commun. Number Theory Phys. **2** (2008), 325–347.
 - [19] J. A. M. Vermaseren, *Harmonic sums, Mellin transforms and integrals*, Internat. J. Modern Phys. A **14** (1999), no. 13, 2037–2076. [arXiv:hep-ph/9806280v1](https://arxiv.org/abs/hep-ph/9806280v1), (1998).
 - [20] S. Yamamoto, *Interpolation of multiple zeta and zeta-star values*, J. Algebra **385** (2013), 102–114.
 - [21] J. Zhao, *Multiple q -zeta functions and multiple q -polylogarithms*, Ramanujan J. **14** (2007), no. 2, 189–221. [arXiv:math/0304448](https://arxiv.org/abs/math/0304448).
 - [22] J. Zhao, *Wolstenholme Type Theorem for multiple harmonic sums*, Int. J. of Number Theory **4** (2008), no. 1, 73–106. [arXiv:math/0301252](https://arxiv.org/abs/math/0301252).
 - [23] J. Zhao, *Identity families of multiple harmonic sums and multiple zeta (star) values*, to appear: J. Math. Soc. Japan. [arXiv:1303.2227](https://arxiv.org/abs/1303.2227).

THE FIELDS INSTITUTE FOR RESEARCH IN MATHEMATICAL SCIENCES
222 COLLEGE ST, TORONTO, ON M5T 3J1, CANADA

E-mail address: hessamik@gmail.com

THE FIELDS INSTITUTE FOR RESEARCH IN MATHEMATICAL SCIENCES
222 COLLEGE ST, TORONTO, ON M5T 3J1, CANADA

E-mail address: hessamit@gmail.com

DEPARTMENT OF MATHEMATICS, THE BISHOP'S SCHOOL

LA JOLLA, CA 92037, USA

E-mail address: zhaoj@ihes.fr

RECEIVED MARCH 22, 2015