

ON THE COHOMOLOGY OF SOME HOPF ALGEBROIDS AND HATTORI-STONG THEOREMS

ANDREW BAKER

(*communicated by Hvedri Inassaridze*)

Abstract

We apply group cohomological methods to calculate the cohomology of $K(n)_*BP$ as a $K(n)_*K(n)$ -comodule, recovering recent results of Hovey and Sadofsky. As applications we determine the Chromatic Spectral Sequence for BP based on Johnson and Wilson's $E(n)$, showing the relationship to some generalizations of the classical Hattori-Stong Theorem and determine the change of Hopf algebroid spectral sequence associated with the natural map $BP \rightarrow E(n)$, extending calculations of Clarke for the Todd orientation $MU \rightarrow KU$.

Introduction

In an interesting paper [4], Hovey and Sadofsky showed amongst other things that the Chromatic Spectral Sequence associated to a p -local v_n -periodic ring spectrum such as Johnson and Wilson's $E(n)$ is obtained by taking the 0 to n th columns of the Chromatic Spectral Sequence associated with BP -theory. They also identified $K(n)_*BP$ as a comodule over the Hopf algebroid $K(n)_*K(n) = K(n)_*(E(n))$. In [2], we also analysed the structure of algebras of the form

$$R_*K(n) = R_* \otimes_{BP_*} BP_*K(n),$$

where R_* is a BP_* -algebra such as BP_* or $E(n)_*$, and used our results to give an alternative approach to such results.

Clarke [3] discussed the change of Hopf algebroid spectral sequence associated to the Todd orientation $\tau: MU \rightarrow KU$ and showed that

$$\mathrm{Ext}_{KU_*KU}^{1,*}(KU_*, KU_*MU) \neq 0,$$

despite the Hattori-Stong Theorem, which was taken as evidence that the KU -theory Adams Spectral Sequence might be trivial since

$$MU_* \cong \mathrm{Ext}_{KU_*KU}^{0,*}(KU_*, KU_*MU).$$

Of course, in the two decades since the publication of [3], our understanding of the target $\pi_*(L_{KU}MU)$ of this spectral sequence has improved and such a result no longer appears surprising.

In this paper we use methods of Galois cohomology to rederive certain cohomology groups over the Hopf algebra $K(n)_*K(n)$ and explore some consequences of our current knowledge of the Chromatic Spectral Sequence for

$$\mathrm{Ext}_{E(n)_*E(n)}^{*,*}(E(n)_*, E(n)_*BP),$$

Received 5 January 2000, revised 15 February 2000; published on 9 March 2000.

1991 Mathematics Subject Classification: Primary 55N20 55N22 55T15

Key words and phrases: Hopf algebroid, comodule, Brown-Peterson spectrum, Johnson-Wilson spectrum, Galois cohomology

© 2000, Andrew Baker. Permission to copy for private use granted.

and compute this completely, relating the result to a change of Hopf algebroid spectral sequence

$$\begin{aligned} E_2^{s,t,*} &= \text{Ext}_{BP_*BP}^{r,*}(BP_*, \text{Ext}_{E(n)_*E(n)}^{s,*}(E(n)_*, E(n)_*BP)) \\ &\implies \text{Ext}_{E(n)_*E(n)}^{s+t,*}(E(n)_*, E(n)_*). \end{aligned}$$

This generalizes Clarke's work and we show that there are always non-zero elements in

$$\text{Ext}_{E(n)_*E(n)}^{n,*}(E(n)_*, E(n)_*)$$

playing an analogous rôle to the elements he identified when $n = 1$. We also state some generalizations of the Hattori-Stong Theorem following from our analysis.

I would like to thank Ken Brown, Francis Clarke, Steve Cohen, Mark Hovey and Hal Sadofsky for their help and inspiration, and also the referee of an earlier version whose perceptive comments were invaluable.

1. Review of some results of Hovey and Sadofsky

Throughout this paper, we follow well established (but nevertheless unfortunate) fashion in writing

$$K(n)_*K(n) = K(n)_*(E(n)) \subsetneq K(n)_*(K(n)).$$

By [4, theorem B], we have a good description of the $K(n)_*K(n)$ -comodule structure of $K(n)_*BP$; note that this is not claimed to give a description of the $K(n)_*K(n)$ -comodule algebra structure.

Theorem 1.1. *There is an isomorphism of left $K(n)_*K(n)$ -comodules*

$$K(n)_*BP \cong K(n)_*E(n)[w_i : i \geq 1],$$

where $w_i \in K(n)_{2p^n(p^i-1)}BP$ is primitive with respect to the coaction.

$K(n)_*BP$ is an extended $K(n)_*K(n)$ -comodule, so for $s > 0$,

$$\text{Ext}_{K(n)_*K(n)}^{s,*}(K(n)_*, K(n)_*BP) = 0.$$

It is straightforward to deduce the following consequence.

Corollary 1.2. $E(n)_*BP/(p^\infty, v_1^\infty, \dots, v_{n-1}^\infty)$ is an extended left $E(n)_*E(n)$ -comodule,

$$\begin{aligned} E(n)_*BP/(p^\infty, v_1^\infty, \dots, v_{n-1}^\infty) \\ \cong E(n)_*E(n) \otimes_{E(n)_*} E(n)_*/(p^\infty, v_1^\infty, \dots, v_{n-1}^\infty)[\tilde{w}_i : i \geq 1], \end{aligned}$$

where \tilde{w}_i has degree $2p^n(p^i - 1)$ and is coaction primitive. Hence for $s > 0$,

$$\text{Ext}_{E(n)_*E(n)}^{s,*}(E(n)_*, E(n)_*BP/(p^\infty, v_1^\infty, \dots, v_{n-1}^\infty)) = 0.$$

In [4, theorem 5.1] the Chromatic Spectral Sequence $\text{CSS}(S; E(n))$ based on $E(n)$ for the sphere spectrum S was shown to be obtained by truncating that based on BP . There is an obvious generalization to the Chromatic Spectral Sequence $\text{CSS}(X; E(n))$ of a connective torsion free spectrum X based on $E(n)$ and converging to

$$\text{Ext}_{E(n)_*E(n)}^{*,*}(E(n)_*, E(n)_*X).$$

Recall that the E_1 -term of the Chromatic Spectral Sequence $\text{CSS}(X; BP)$ based on BP and converging to

$$\text{Ext}_{BP_*BP}^{*,*}(BP_*, BP_*X)$$

has the form

$$E_1^{n,t,*}(X; BP) = \text{Ext}_{E(n)_*E(n)}^{t,*}(E(n)_*, E(n)_*X/(p^\infty, v_1^\infty, \dots, v_{n-1}^\infty)). \quad (1.1)$$

Theorem 1.3. *In the E_1 -term of the spectral sequence $\text{CSS}(X; E(n))$,*

$$E_1^{k,t,*}(X; E(n)) = \begin{cases} E_1^{k,t,*}(X; BP) & \text{if } k \leq n, \\ 0 & \text{if } k > n, \end{cases}$$

and the differential $d_1^{k,t,*}$ with domain $E_1^{k,t,*}(X; E(n))$ is obtained from the corresponding differential of $\text{CSS}(X; BP)$ whenever $E_1^{k,t,*}(X; E(n)) \neq 0$.

When $X = BP$, Corollary 1.2 implies that the only possible non-zero differential in each of the spectral sequences $\text{CSS}(BP; E(n))$ and $\text{CSS}(BP; BP)$ is d_1 .

2. The cohomology of $K(n)_*BP$ as Galois cohomology

Our aim in this section is to reprove the results of Theorem 1.1 and Corollary 1.2 using our previous work [1, 2] and making use of continuous cohomology of profinite groups.

From now on, let R_* be an algebra over $\mathbb{F}_{p^n} \otimes BP_*$ annihilated by I_n and in which there is a unit u satisfying $v_n = u^{p^n - 1}$. In our applications to $K(n)_*BP$ we will take

$$R_* = \mathbb{F}_{p^n} \otimes (v_n^{-1}BP_*/I_n)[u]/(u^{p^n - 1} - v_n),$$

but for completeness we give the general version.

We may form the R_* -algebra

$$R_*K(n) = R_* \otimes_{BP_*} BP_*BP \otimes_{BP_*} K(n)_*.$$

From [2, Theorem A.3] we have

Theorem 2.1. *$R_*K(n)$ is a free R_* -module. Moreover, there is an exhaustive filtration by subalgebras*

$$R_* = R_*K(n)^{(1)} \subseteq R_*K(n)^{(2)} \subseteq \dots \subseteq R_*K(n)^{(k)} \subseteq \dots \subseteq R_*K(n)$$

in which each extension $R_*K(n)^{(k)} \rightarrow R_*K(n)^{(k+1)}$ is a free $R_*K(n)^{(k)}$ -module and a Galois extension of Artin-Schreier type with Galois group \mathbb{F}_{p^n} .

There is an action of the Morava stabiliser group \mathbb{S}_n on $R_*K(n)$ with respect to which

$$R_*K(n)^{(k)} = R_*K(n)_*^{\mathbb{S}_n^{[k]}},$$

the fixed point set of the closed subgroup

$$\mathbb{S}_n^{[k]} = \left\{ 1 + \sum_{k \leq r} \alpha_r S^r : \forall r, \alpha_r^{p^n} = \alpha_r \right\} \leq \mathbb{S}_n.$$

Here an extension $A \rightarrow B$ of commutative unital \mathbb{F}_q -algebras is of *Artin-Schreier type* if

$$B = A[t]/(t^q - t - a)$$

for some $a \in A$; its Galois group is $\text{Gal}(B/A) = \mathbb{F}_q \leq \text{Aut}_A(B)$ where $\gamma \in \mathbb{F}_q$ acts by $\gamma \cdot f(t) = f(t + \gamma)$. Notice that B is a free A -module of rank q and $B^{\text{Gal}(B/A)} = A$, however in general $\text{Gal}(B/A) \neq \text{Aut}_A(B)$ (consider for example the case where $a = 0$).

We will apply this result to calculate the continuous cohomology of $BP_*K(n)$ with respect to with action of \mathbb{S}_n , where this continuous cohomology is equivalent to the cohomology over the $K(n)_*$ -Hopf algebra $K(n)_*K(n)$. Further information the relationship of the action of \mathbb{S}_n and the $K(n)_*K(n)$ -comodule structure can be found in [1].

Theorem 2.2. *The continuous cohomology of $R_*K(n)$ with respect to \mathbb{S}_n is given by*

$$H_c^s(\mathbb{S}_n; R_*K(n)) = \begin{cases} R_* & \text{if } s = 0, \\ 0 & \text{if } s \neq 0. \end{cases}$$

In order to apply Galois cohomological machinery we will require two technical Galois theoretic results surely known to experts but for which we know of no convenient references.

Lemma 2.3 (Normal Basis Theorem for Artin-Schrier extensions). *Let A be a commutative unital \mathbb{F}_q -algebra. Let $B = A[t]/(f(t))$ be the Artin-Schrier extension associated with the polynomial $f(X) = X^q - X + a \in A[X]$. Then the elements $(t + \gamma)^{q-1}$ ($\gamma \in \mathbb{F}_q$) form an A -basis for B . Hence B is a free $A[\text{Gal}(B/A)]$ -module.*

Proof. We will show that in the \mathbb{F}_q -vector subspace spanned by the monomials t^k ($0 \leq k \leq q - 1$), the stated elements form a basis. Let $\omega \in \mathbb{F}_q^\times$ be a generator, so the non-zero elements of \mathbb{F}_q are expressible as powers of ω .

By the Binomial Theorem,

$$(t + \gamma)^{q-1} = \sum_{0 \leq r \leq q-1} \binom{q-1}{r} \gamma^r t^{q-1-r},$$

so it suffices to show the non-singularity of the matrix

$$\begin{bmatrix} 1 & 0 & 0 & \cdots & 0 \\ 1 & \binom{q-1}{1} & \binom{q-1}{2} & \cdots & 1 \\ 1 & \binom{q-1}{1} \omega & \binom{q-1}{2} \omega^2 & \cdots & 1 \\ 1 & \binom{q-1}{1} \omega^2 & \binom{q-1}{2} \omega^4 & \cdots & 1 \\ 1 & \binom{q-1}{1} \omega^3 & \binom{q-1}{2} \omega^6 & \cdots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & \binom{q-1}{1} \omega^{q-2} & \binom{q-1}{2} \omega^{2(q-2)} & \cdots & 1 \end{bmatrix}$$

whose determinant is

$$\begin{aligned} & \begin{vmatrix} 1 & 0 & 0 & \cdots & 0 \\ 1 & \binom{q-1}{1} & \binom{q-1}{2} & \cdots & 1 \\ 1 & \binom{q-1}{1} \omega & \binom{q-1}{2} \omega^2 & \cdots & 1 \\ 1 & \binom{q-1}{1} \omega^2 & \binom{q-1}{2} \omega^4 & \cdots & 1 \\ 1 & \binom{q-1}{1} \omega^3 & \binom{q-1}{2} \omega^6 & \cdots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & \binom{q-1}{1} \omega^{q-2} & \binom{q-1}{2} \omega^{2(q-2)} & \cdots & 1 \end{vmatrix} = \begin{vmatrix} 1 & 0 & 0 & \cdots & 0 \\ 1 & \binom{q-1}{1} & \binom{q-1}{2} & \cdots & 1 \\ 0 & \binom{q-1}{1} \omega & \binom{q-1}{2} \omega^2 & \cdots & 1 \\ 0 & \binom{q-1}{1} \omega^2 & \binom{q-1}{2} \omega^4 & \cdots & 1 \\ 0 & \binom{q-1}{1} \omega^3 & \binom{q-1}{2} \omega^6 & \cdots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \binom{q-1}{1} \omega^{q-2} & \binom{q-1}{2} \omega^{2(q-2)} & \cdots & 1 \end{vmatrix} \\ & = \begin{pmatrix} q-1 \\ 1 \end{pmatrix} \begin{pmatrix} q-1 \\ 2 \end{pmatrix} \cdots \begin{pmatrix} q-1 \\ q-2 \end{pmatrix} \begin{vmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 1 & \cdots & 1 \\ 0 & \omega & \omega^2 & \cdots & 1 \\ 0 & \omega^2 & \omega^4 & \cdots & 1 \\ 0 & \omega^3 & \omega^6 & \cdots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \omega^{q-2} & \omega^{2(q-2)} & \cdots & 1 \end{vmatrix} \\ & = \begin{pmatrix} q-1 \\ 1 \end{pmatrix} \begin{pmatrix} q-1 \\ 2 \end{pmatrix} \cdots \begin{pmatrix} q-1 \\ q-2 \end{pmatrix} \prod_{0 \leq r < s \leq q-2} (\omega^r - \omega^s). \end{aligned}$$

This is non-zero because the binomial coefficients satisfy

$$\binom{q-1}{r} \equiv \binom{p-1}{r_0} \binom{p-1}{r_1} \cdots \binom{p-1}{r_d} \pmod{p},$$

where $r = r_0 + r_1p + \dots + r_m p^m$ is the p -adic expansion of r , and the factors are never divisible by p . So the $(t + \gamma)^{q-1}$ form a basis in place of the t^k . As this is true over \mathbb{F}_q , it is also true over A . \square

The next result will allow an inductive calculation of cohomology in the situation of interest.

Lemma 2.4. *Let $A \subseteq B \subseteq C$ be a tower of commutative unital rings. Let $G \leq \text{Aut}_A(C)$ be finite and $N \triangleleft G$ with $C^G = A$, $C^N = B$, and $C \cong B[N]$ as a $B[N]$ -module. Then*

$$H^*(G; C) \cong H^*(G/N; B).$$

Proof. There is a Hochschild-Serre Spectral Sequence [7, chapter II §4],

$$E_2^{r,s} = H^r(G/N; H^s(N; C)) \implies H^{r+s}(G; C).$$

Since

$$H^*(N; C) \cong H^*(N; B[N]) = H^0(N; B[N]) \cong H^0(N; C) = B,$$

the E_2 -term is trivial except for $E_2^{r,0} = H^r(G/N; B)$, so the result follows. \square

Proof of Theorem 2.2. From the proof of [2, theorem A.3], for $k \geq 1$, $R_*K(n)^{(k)}$ is a free R_* -module with a basis consisting of the monomials

$$t_1^{s_1} t_2^{s_2} \dots t_{k-1}^{s_{k-1}}, \quad (0 \leq s_k \leq p^n - 1). \tag{2.1}$$

For each of these basis elements, the stabilizer is an open subgroup of \mathbb{S}_n since it contains an open subgroup of the form $\mathbb{S}_n^{[k]}$. Hence the action of \mathbb{S}_n on $R_*K(n)$ is *proper* in the sense that the stabilizer of each point is an open subgroup and this allows us to apply the cohomological machinery of Shatz [7] to $R_*K(n)$. In particular, [7, chapter 3 theorem 7] gives

$$H_c^*(\mathbb{S}_n; R_*K(n)) = \varinjlim_k H^*(\mathbb{S}_n/\mathbb{S}_n^{[k]}; R_*K(n)^{(k)}). \tag{2.2}$$

To complete the proof we will show by induction on $k \geq 1$ that

$$H^*(\mathbb{S}_n/\mathbb{S}_n^{[k]}; R_*K(n)^{(k)}) = H^0(\mathbb{S}_n/\mathbb{S}_n^{[k]}; R_*K(n)^{(k)}) = R_*.$$

For $k = 1$, the result is certainly true. Suppose that it holds for some $k = m \geq 1$. The extension $R_*K(n)^{(m)} \subseteq R_*K(n)^{(m+1)}$ satisfies the conditions of Lemma 2.3 and so we can also apply Lemma 2.4 with

$$A = R_*, \quad B = R_*K(n)^{(m)}, \quad C = R_*K(n)^{(m+1)}, \quad G = \mathbb{S}_n/\mathbb{S}_n^{[m+1]}, \quad N = \mathbb{S}_n^{[m]}/\mathbb{S}_n^{[m+1]}.$$

This yields

$$H^*(\mathbb{S}_n/\mathbb{S}_n^{[m+1]}; R_*K(n)^{(m+1)}) = H^*(\mathbb{S}_n/\mathbb{S}_n^{[m]}; R_*K(n)^{(m)}) = R_*,$$

which verifies the inductive step. \square

Remark. Since $\mathbb{S}_n/\mathbb{S}_n^{[k]}$ is a p -group, the vanishing of $H^s(\mathbb{S}_n/\mathbb{S}_n^{[k]}; R_*K(n)^{(k)})$ for $s > 0$ actually implies that $R_*K(n)^{(k)}$ is a projective $\mathbb{F}_q[\mathbb{S}_n/\mathbb{S}_n^{[k]}]$ -module and hence is free because this group algebra is local. Of course, if we knew an explicit permutation basis we could avoid the cohomological calculations.

Corollary 2.5.

$$\text{Ext}_{K(n)_*K(n)}^{s,*}(K(n)_*, K(n)_*BP) = \begin{cases} v_n^{-1}BP_*/I_n & \text{if } s = 0, \\ 0 & \text{if } s \neq 0. \end{cases}$$

Proof. We apply the Theorem in the case

$$R_* = \mathbb{F}_{p^n} \otimes (v_n^{-1}BP_*/I_n)[u]/(u^{p^n-1} - v_n).$$

A complication comes from the fact that we have extended $v_n^{-1}BP_*/I_n$ to R_* by tensoring with \mathbb{F}_{p^n} then adjoining a $(p^n - 1)$ st root of v_n . This is a Galois extension whose Galois group Γ_n is identified with $\mathbb{F}_{p^n}^\times \rtimes \mathbb{Z}/n$. Furthermore,

$$\text{Ext}_{K(n)_*K(n)}^{s,*}(K(n)_*, K(n)_*BP) = H_c^*(\mathbb{S}_n; R_*K(n))^{\Gamma_n}.$$

All of this is straightforward given the well understood identification of \mathbb{S}_n with the group of strict units in the maximal order \mathcal{O}_n of a certain p -adic division algebra \mathbb{D}_n . In fact there is a proper action on $R_*K(n)$ of a semi-direct product extension of \mathbb{S}_n , $\mathcal{O}_n^\times \rtimes \mathbb{Z}/n$, where the full group of units \mathcal{O}_n^\times can itself be viewed as an extension

$$1 \rightarrow \mathbb{S}_n \rightarrow \mathcal{O}_n^\times \rightarrow \mathbb{F}_{p^n}^\times \rightarrow 1.$$

□

We will use the notation $u_k = \eta_L(v_k)$, $v_k = \eta_R(v_k)$, where

$$E(n)_* \xrightarrow{\eta_L} E(n)_*BP \xleftarrow{\eta_R} BP_*$$

are the left and right units. As v_k is a coaction primitive, multiplication by v_k is a homomorphism of $E(n)_*E(n)$ -comodules $E(n)_*BP \rightarrow E(n)_*BP$. Hence $E(n)_*BP/(p^{r_0}, v_1^{r_1}, \dots, v_{n-1}^{r_{n-1}})$ is a quotient $E(n)_*E(n)$ -comodule of $E(n)_*BP$. As $I_n = (p, v_1, \dots, v_{n-1}) \triangleleft BP_*$ is an invariant ideal,

$$K(n)_*BP = E(n)_*BP/(p, v_1, \dots, v_{n-1}) = E(n)_*BP/(p, u_1, \dots, u_{n-1}).$$

Theorem 2.6. *Suppose that $1 \leq r_k \leq \infty$ for $k = 0, 1, \dots, n-1$. Then*

$$\begin{aligned} \text{Ext}_{E(n)_*E(n)}^{s,*}(E(n)_*, E(n)_*BP/(p^{r_0}, v_1^{r_1}, \dots, v_{n-1}^{r_{n-1}})) \\ = \begin{cases} v_n^{-1}BP_*/(p^{r_0}, v_1^{r_1}, \dots, v_{n-1}^{r_{n-1}}) & \text{if } s = 0, \\ 0 & \text{if } s \neq 0. \end{cases} \end{aligned}$$

Proof. Suppose that all of the exponents r_k are finite. We will prove the result by induction on the number

$$\text{excess}(r_0, r_1, \dots, r_{n-1}) = \sum_{k=0}^{n-1} (r_k - 1) \geq 0.$$

Corollary 2.5 implies that the result is true for the unique sequence $(1, 1, \dots, 1)$ of excess 0.

Now suppose that it holds whenever $\text{excess}(r_0, r_1, \dots, r_{n-1}) < m$. Then for at least one value of ℓ in the range $0 \leq \ell \leq (n-1)$ we have $r_\ell > 1$. Let $(r'_0, r'_1, \dots, r'_{n-1})$ be the exponent sequence for which

$$r'_k = \begin{cases} r_\ell - 1 & \text{if } k = \ell, \\ r_k & \text{if } k \neq \ell. \end{cases}$$

Then there is an exact sequence of $E(n)_*E(n)$ -comodules

$$\begin{aligned} 0 \rightarrow E(n)_*BP/(p^{r_0}, v_1^{r_1}, \dots, v_{\ell-1}^{r_{\ell-1}}, v_\ell, v_{\ell+1}^{r_{\ell+1}}, \dots, v_{n-1}^{r_{n-1}}) \rightarrow \\ E(n)_*BP/(p^{r_0}, v_1^{r_1}, \dots, v_{n-1}^{r_{n-1}}) \rightarrow E(n)_*BP/(p^{r'_0}, v_1^{r'_1}, \dots, v_{n-1}^{r'_{n-1}}) \rightarrow 0. \end{aligned}$$

Here the left-hand monomorphism is induced from the homomorphism of comodules

$$E(n)_*BP \rightarrow E(n)_*BP/(p^{r_0}, v_1^{r_1}, \dots, v_{n-1}^{r_{n-1}}); \quad x \mapsto v_\ell^{r_\ell-1}x.$$

The exponent sequences of the left and right hand terms each have excess less than m and so we obtain a long exact sequence in Ext yielding the result for

$$E(n)_*BP/(p^{r_0}, v_1^{r_1}, \dots, v_{n-1}^{r_{n-1}}).$$

Hence the result for sequences of arbitrary excess follows by induction.

When some of the r_k are infinite the result is proved by using appropriate colimits over finite exponent cases under obvious inclusions maps. \square

3. Some generalizations of the Hattori-Stong Theorem

Recall the well-known fact that

$$\text{Ext}_{BP_*BP}^{s,*}(BP_*, BP_*BP) = \begin{cases} BP_* & \text{if } s = 0, \\ 0 & \text{if } s \neq 0. \end{cases}$$

Using this we can deduce that $\text{CSS}(BP; BP)$ collapses at $E_2 = E_\infty$. This gives

Theorem 3.1 (Hattori-Stong Theorem in $E(n)$ -theory). *For each $n \geq 1$, the $E(n)$ -Hurewicz homomorphism $BP_* \rightarrow E(n)_*BP$ is a split monomorphism.*

The following observation will prove useful in the proof.

Lemma 3.2. *For $n \geq 1$,*

$$\text{Ext}_{E(n)_*E(n)}^{0,*}(E(n)_*, E(n)_*BP) \subseteq E(n)_*BP$$

is the kernel of the $\mathbb{Z}_{(p)}$ -homomorphism

$$\psi - 1 \otimes \text{Id}: E(n)_*BP \rightarrow E(n)_*E(n) \otimes_{E(n)_*} E(n)_*BP$$

in which the domain and codomain are torsion-free; hence it is an additive summand.

Proof of Theorem 3.1. There is an exact sequence

$$\begin{aligned} 0 \rightarrow \text{Ext}_{BP_*BP}^{0,*}(BP_*, BP_*BP) &\rightarrow \text{Ext}_{E(0)_*E(0)}^{0,*}(E(0)_*, E(0)_*BP) \\ &\xrightarrow{d_1} \text{Ext}_{E(1)_*E(1)}^{0,*}(E(1)_*, E(1)_*BP/(p^\infty)) \xrightarrow{d_1} \dots \end{aligned}$$

By the well known fact that

$$\text{Ext}_{E(0)_*E(0)}^{0,*}(E(0)_*, E(0)_*BP) = p^{-1}BP_*$$

this gives rise to an exact sequence

$$\begin{aligned} 0 \rightarrow \text{Ext}_{BP_*BP}^{0,*}(BP_*, BP_*BP) &\rightarrow p^{-1}BP_* \\ &\xrightarrow{d_1} \text{Ext}_{E(1)_*E(1)}^{0,*}(E(1)_*, E(1)_*BP/(p^\infty)) \xrightarrow{d_1} \dots \end{aligned}$$

From $\text{CSS}(BP; E(1))$ we obtain an exact sequence

$$0 \rightarrow BP_* \rightarrow p^{-1}BP_* \xrightarrow{d_1} \text{Ext}_{E(1)_*E(1)}^{0,*}(E(1)_*, E(1)_*BP/(p^\infty))$$

implying that the Hurewicz homomorphism $BP_*/(p^\infty) \rightarrow E(1)_*BP/(p^\infty)$ is a monomorphism. This in turn shows that the Hurewicz homomorphism $BP_* \rightarrow E(1)_*BP$ is indivisible, hence is a split monomorphism by Lemma 3.2.

For $n \geq 1$, $\text{CSS}(BP; E(n))$ gives rise to an exact sequence

$$\begin{aligned} 0 \rightarrow BP_* \rightarrow p^{-1}BP_* &\xrightarrow{d_1} \text{Ext}_{E(1)_*E(1)}^{0,*}(E(1)_*, E(1)_*BP/(p^\infty)) \xrightarrow{d_1} \dots \\ &\dots \xrightarrow{d_1} \text{Ext}_{E(n)_*E(n)}^{0,*}(E(n)_*, E(n)_*BP/(p^\infty, v_1^\infty, \dots, v_{n-1}^\infty)). \end{aligned}$$

From this we obtain

$$E_{\infty}^{0,0,*} \cong \text{Ext}_{E(n)_*E(n)}^{0,*}(E(n)_*, E(n)_*BP) = BP_*,$$

and together with Lemma 3.2, this proves the theorem. \square

Theorem 3.3. For $n \geq 1$,

$$\text{Ext}_{E(n)_*E(n)}^{s,*}(E(n)_*, E(n)_*BP) \cong \begin{cases} BP_* & \text{if } s = 0, \\ BP_*/(p^\infty, v_1^\infty, \dots, v_n^\infty) & \text{if } s = n, \\ 0 & \text{if } s \neq 0, n, \end{cases}$$

Proof. This is deduced using the exact sequence derived from $\text{CSS}(BP; E(n))$. It only remains to identify Ext^n which we do by induction on n .

When $n = 0$, $\text{CSS}(BP; E(0))$ has trivial E_1 -term

$$E_1^{0,t,*}(BP; E(0)) = \begin{cases} p^{-1}BP_* & \text{if } t = 0, \\ 0 & \text{if } t \neq 0. \end{cases}$$

This gives rise to the exact sequence

$$0 \rightarrow BP_* \rightarrow p^{-1}BP_* \rightarrow BP_*/(p^\infty) \rightarrow 0,$$

where $BP_*/(p^\infty)$ is identified with the image of the differential d_1 in $\text{CSS}(BP; E(1))$.

Consider the case $n = 1$ and $\text{CSS}(BP; E(1))$. Here the E_1 -term is

$$0 \rightarrow p^{-1}BP_* \xrightarrow{d_1} v_1^{-1}BP_*/(p^\infty) \rightarrow 0$$

in which d_1 is the evident composite

$$p^{-1}BP_* \rightarrow BP_*/(p^\infty) \rightarrow v_1^{-1}BP_*/(p^\infty).$$

Hence the E_1 -term fits into an exact sequence

$$0 \rightarrow BP_* \rightarrow p^{-1}BP_* \xrightarrow{d_1} v_1^{-1}BP_*/(p^\infty) \rightarrow BP_*/(p^\infty, v_1^\infty) \rightarrow 0$$

and the E_∞ -term is

$$E_{\infty}^{1,r,*}(BP; E(1)) = \begin{cases} BP_* & \text{if } r = 0, \\ BP_*/(p^\infty, v_1^\infty) & \text{if } r = 1, \\ 0 & \text{if } r \neq 0, 1. \end{cases}$$

If the stated result is correct for $n < m$, then for $n = m$ the E_1 -term is the complex

$$\begin{aligned} 0 \rightarrow p^{-1}BP_* \xrightarrow{d_1} v_1^{-1}BP_*/(p^\infty) \xrightarrow{d_1} \dots \\ \dots \xrightarrow{d_1} v_{n-1}^{-1}BP_*/(p^\infty, \dots, v_{n-2}^\infty) \xrightarrow{d_1} v_n^{-1}BP_*/(p^\infty, \dots, v_{n-1}^\infty) \rightarrow 0, \end{aligned}$$

in which the differentials are evident composites of the form

$$v_k^{-1}BP_*/(p^\infty, \dots, v_{k-1}^\infty) \rightarrow BP_*/(p^\infty, \dots, v_{k-1}^\infty, v_k^\infty) \rightarrow v_{k+1}^{-1}BP_*/(p^\infty, \dots, v_k^\infty).$$

From this it is clear that

$$E_{\infty}^{n,t,*}(BP; E(n)) = \begin{cases} BP_* & \text{if } t = 0, \\ BP_*/(p^\infty, v_1^\infty, \dots, v_n^\infty) & \text{if } t = n, \\ 0 & \text{if } t \neq 0, n. \end{cases}$$

This establishes the inductive step. \square

It is interesting to note that the case $n = 1$ gives the diagram of exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & BP_* & \longrightarrow & p^{-1}BP_* & \xrightarrow{d_1} & v_1^{-1}BP_*/(p^\infty) \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & BP_* & \longrightarrow & p^{-1}BP_* & \longrightarrow & E(1)_*BP/(p^\infty) \end{array}$$

This shows that the composite map

$$BP_*/(p^\infty) \longrightarrow v_1^{-1}BP_*/(p^\infty) \longrightarrow E(1)_*BP/(p^\infty)$$

is a monomorphism. As this map agrees with the $E(1)$ -theory Hurewicz map

$$BP_*/(p^\infty) \longrightarrow E(1)_*BP/(p^\infty),$$

the Hurewicz map $BP_* \longrightarrow E(1)_*BP$ is indivisible, hence is a split monomorphism. This is a reformulation of the classical Hattori-Stong Theorem.

One analogue of Hattori-Stong for $E(n)_*BP$ is that the $E(n)$ -theory Hurewicz map $BP_* \longrightarrow E(n)_*BP$ is indivisible; this follows from knowledge of the spectral sequence $CSS(BP; E(n))$. A more delicate consequence is that the homomorphism

$$BP_*/(p^\infty, \dots, v_{n-1}^\infty) \longrightarrow v_n^{-1}BP_*/(p^\infty, \dots, v_{n-1}^\infty) \longrightarrow E(n)_*BP/(p^\infty, \dots, v_{n-1}^\infty)$$

is a monomorphism, hence

$$BP_*/(p^\infty, \dots, v_{n-2}^\infty) \longrightarrow E(n)_*BP/(p^\infty, \dots, v_{n-2}^\infty)$$

is v_{n-1} -indivisible.

4. A change of Hopf algebroid spectral sequence

We will make use of a spectral sequence concerning a morphism $\varphi: (A, \Gamma) \longrightarrow (B, \Sigma)$ of Hopf algebroids over a commutative unital ring \mathbb{k} .

Let M be a left Γ -comodule and N a left Σ -comodule. We give $\Gamma \otimes_A B$ and $M \otimes_A B$ their natural right Σ -comodule structures. We also have the cotensor product analogue of the adjunction formula of [5]. We say that a A - B -bimodule L is a Γ - Σ -bicomodule if it admits left Γ -comodule and right Σ -comodule structures

$$\Gamma \otimes_A L \xleftarrow{\psi_L} L \xrightarrow{\psi_R} L \otimes_B \Sigma$$

which satisfy the compatibility condition

$$(1 \otimes \psi_R)\psi_L = (\psi_L \otimes 1)\psi_R.$$

If L is a Γ - Σ bicomodule, then by coassociativity

$$M \square_{\Gamma} (L \square_{\Sigma} N) \cong (M \square_{\Gamma} L) \square_{\Sigma} N,$$

implying

$$M \square_{\Gamma} \left((\Gamma \otimes_A B) \square_{\Sigma} N \right) \cong \left(M \square_{\Gamma} (\Gamma \otimes_A B) \right) \square_{\Sigma} N \cong (M \otimes_A B) \square_{\Sigma} N. \tag{4.1}$$

Theorem 4.1. *If M is a flat A -module, there is a spectral sequence*

$$E_2^{s,t,*}(M, N) = \text{Cotor}_{\Gamma}^{s,*}(M, \text{Cotor}_{\Sigma}^{t,*}(\Gamma \otimes_A B, N)) \implies \text{Cotor}_{\Sigma}^{s+t,*}(M \otimes_A B, N)$$

with differentials $d_r: E_2^{s,t,*}(M, N) \longrightarrow E_2^{s+r,t-r+1,*}(M, N)$.

Proof. This spectral sequence is constructed by Ravenel [6, theorem A1.3.11]. However, we feel it is illuminating to view it as a Grothendieck composite functor spectral sequence as discussed in [7, 8]. For completeness we outline this description.

It is a standing assumption that for a Hopf algebroid (A, Γ) , Γ is flat as a left or right A -module. Given a \mathbb{k} -algebra homomorphism $A \rightarrow B$, $\Gamma \otimes_A B$ is flat as a right B -module and $B \otimes_A \Gamma$ is flat as a left B -module. Also, if M is a flat A -module the functors $\text{Cotor}_\Gamma^s(M, _)$ ($s \geq 0$) is computable using resolutions of the second variable by weakly injective extended comodules.

Consider the following functors and their right derived functors:

$$G: N \rightsquigarrow (\Gamma \otimes_A B) \square_\Sigma N, \quad \text{R}^t G(_) = \text{Cotor}_\Sigma^{t,*}(\Gamma \otimes_A B, _),$$

from the category of left Σ -comodules to left Γ -comodules, and

$$F: L \rightsquigarrow M \square_\Gamma L, \quad \text{R}^s F(_) = \text{Cotor}_\Gamma^{s,*}(M, _),$$

from the category of left Γ -comodules to \mathbb{k} -modules. Then by Equation (4.1),

$$FG(_) \cong (M \otimes_A B) \square_\Sigma(_),$$

with right derived functors

$$\text{R}^n FG(_) = \text{Cotor}_\Sigma^{n,*}(M \otimes_A B, _).$$

For an extended Σ -comodule $\Sigma \otimes_B N_0$,

$$G(\Sigma \otimes_B N_0) = (\Gamma \otimes_A B) \square_\Sigma(\Sigma \otimes_B N_0) \cong (\Gamma \otimes_A B) \otimes_B N_0,$$

which is an extended Γ -comodule. So for $s > 0$,

$$\text{R}^s F(G(\Sigma \otimes_B N_0)) = 0.$$

This is the condition required to obtain the Grothendieck Spectral Sequence

$$E_2^{s,t}(M, N) = \text{R}^s F \circ \text{R}^t G(N) \implies \text{R}^{s+t} FG(N).$$

□

In the following corollary, Ext_Γ is taken on the category of *left* Γ -comodules, while Ext_Σ is taken on the category of *right* Σ -comodules.

Corollary 4.2. *There is a spectral sequence*

$$E_2^{s,t,*} = \text{Ext}_\Gamma^{s,*}(A, \text{Ext}_\Sigma^{t,*}(B, \Gamma \otimes_A B)) \implies \text{Ext}_\Sigma^{s+t,*}(B, B)$$

with differentials $d_r: E_2^{s,t,*} \rightarrow E_2^{s+r,t-r+1,*}$.

Proof. Recalling the standard equivalences of functors

$$\text{Ext}_\Gamma^{s,*}(A, _) \cong \text{Cotor}_\Gamma^{s,*}(A, _), \quad \text{Ext}_\Sigma^{s,*}(B, _) \cong \text{Cotor}_\Sigma^{s,*}(_, B),$$

this follows from the Theorem with $M = A$ (with its right coaction twisted into a left action) and $N = B$. □

As a particular case of this result, if $\Gamma \otimes_A B$ is a relatively injective right Σ -comodule then

$$\text{Ext}_\Sigma^{n,*}(B, B) \cong \text{Ext}^{n,*}(A, (\Gamma \otimes_A B) \square_\Sigma B) \cong \text{Ext}^{n,*}(A, (\Gamma \otimes_A B) \square_\Sigma B),$$

which is the *Shapiro Lemma* of Miller & Ravenel [5].

Clarke [3] pointed out that the morphism of Hopf algebroids

$$\tau: (MU_*, MU_*MU) \rightarrow (KU_*, KU_*KU)$$

induced by the Todd orientation $MU \rightarrow KU$ gives an example for which this assumption fails since $\text{Ext}_{KU_*KU}^{1,*}(KU_*, KU_*)$ has infinitely many elements surviving from

$$\text{Ext}_{MU_*MU}^{0,*}(MU_*, \text{Ext}_{KU_*KU}^{1,*}(KU_*, MU_*KU))$$

in the relevant spectral sequence. At each prime p this can also be seen by considering the morphism

$$\tau_1: (BP_*, BP_*BP) \rightarrow (E(1)_*, E(1)_*E(1)).$$

induced by the natural orientation $BP \rightarrow E(1)$. More generally, we will describe the relevant spectral sequence for each of the natural maps

$$\tau_n: (BP_*, BP_*BP) \rightarrow (E(n)_*, E(n)_*E(n)).$$

Let $\text{SS}(n)$ denote the spectral sequence of Corollary 4.2 associated with τ_n .

Theorem 4.3. *For $n \geq 1$, $\text{SS}(n)$ has*

$$E_2^{s,t,*} = \begin{cases} \text{Ext}_{BP_*BP}^{s,*}(BP_*, BP_*) & \text{if } t = 0, \\ \text{Ext}_{BP_*BP}^{s,*}(BP_*, BP_*/(p^\infty, v_1^\infty, \dots, v_n^\infty)) & \text{if } t = n, \\ 0 & \text{if } t \neq 0, n. \end{cases}$$

For $k > 0$ and $p \nmid k$,

$$E_2^{0,1,-2k(p-1)p^r} = \text{Ext}_{BP_*BP}^{0,-2k(p-1)p^r}(BP_*, BP_*/(p^\infty, v_1^\infty))$$

contains a summand isomorphic to \mathbb{Z}/p^{r+1} , having as generator the class represented in the usual chromatic notation by $\frac{1}{p^{r+1}v_1^{kp^r}}$. Such classes are non-vanishing infinite cycles which generate the cyclic groups in E_∞ observed by Clarke. There are of course other elements in this group, for example $\frac{v_2}{pv_1}$. Such elements are killed by the differential d_2 and in fact

$$d_2: \text{Ext}_{BP_*BP}^{s,*}(BP_*, BP_*/(p^\infty, v_1^\infty)) \rightarrow \text{Ext}_{BP_*BP}^{s+2,*}(BP_*, BP_*/(p^\infty, v_1^\infty))$$

is the composition of chromatic coboundaries

$$\begin{aligned} \text{Ext}_{BP_*BP}^{s,*}(BP_*, BP_*/(p^\infty, v_1^\infty)) &\xrightarrow{\delta_1} \text{Ext}_{BP_*BP}^{s+1,*}(BP_*, BP_*/(p^\infty)) \\ &\xrightarrow{\delta_0} \text{Ext}_{BP_*BP}^{s+2,*}(BP_*, BP_*). \end{aligned}$$

Thus we have

$$d_2 \left(\frac{v_2}{pv_1} \right) = \beta_1.$$

Similarly, when $n \geq 1$, the differential d_{n+1} is a composite of chromatic coboundaries

$$\begin{aligned} \text{Ext}_{BP_*BP}^{s,*}(BP_*, BP_*/(p^\infty, v_1^\infty, \dots, v_n^\infty)) &\xrightarrow{\delta_n} \\ \text{Ext}_{BP_*BP}^{s+1,*}(BP_*, BP_*/(p^\infty, v_1^\infty, \dots, v_{n-1}^\infty)) &\xrightarrow{\delta_{n-1}} \dots \\ \dots &\xrightarrow{\delta_0} \text{Ext}_{BP_*BP}^{s+n+1,*}(BP_*, BP_*). \end{aligned}$$

The elements

$$\frac{1}{p^{r_0}v_1^{r_1} \dots v_n^{r_n}} \in \text{Ext}_{BP_*BP}^{0,*}(BP_*, BP_*/(p^\infty, v_1^\infty, \dots, v_n^\infty)) \quad (r_j \geq 1),$$

are non-vanishing infinite cycles, corresponding to non-trivial elements of

$$\text{Ext}_{E(n)_*E(n)}^{n,*}(E(n)_*, E(n)_*).$$

The general picture is that there is an exact sequence of the form

$$0 \rightarrow F^{0,s,*} \rightarrow \text{Ext}_{BP_*BP}^{s,*}(BP_*, BP_*/(p^\infty, v_1^\infty, \dots, v_n^\infty)) \xrightarrow{\delta_0 \cdots \delta_n} \\ \text{Ext}_{BP_*BP}^{s+n+1,*}(BP_*, BP_*) \rightarrow F^{n,s+n+1,*} \rightarrow 0,$$

where the filtration groups $F^{k,s,*}$ fit into short exact sequences

$$0 \rightarrow F^{0,s,*} \rightarrow F^{n,s,*} \rightarrow \text{Ext}_{E(n)_*E(n)}^{s,*}(E(n)_*, E(n)_*) \rightarrow 0.$$

References

- [1] A. Baker, A version of the Landweber filtration theorem for v_n -periodic Hopf algebroids, *Osaka J. Math.* **32** (1995), 689–99.
- [2] A. Baker, On the Adams E_2 -term for elliptic cohomology, to appear in *Proceedings of the 1999 Boulder Conference* in *Contemp. Math.*
- [3] F. Clarke, On $\text{Ext}_{K_*(K)}(\pi_*(K), K_*(MU))$, *Proc. Amer. Math. Soc.* **74** (1979), 156–160.
- [4] M. Hovey & H. Sadofsky, Invertible spectra in the $E(n)$ -local stable homotopy category, *J. London Math. Soc.* **60** (1999), 284–302.
- [5] H. R. Miller & D. C. Ravenel, Morava stabilizer algebras and localization of Novikov's E_2 -term, *Duke Math. J.* **44** (1977), 433–47.
- [6] D. C. Ravenel, *Complex Cobordism and the Stable Homotopy Groups of Spheres*, Academic Press (1986).
- [7] S. Shatz, *Profinite Groups, Arithmetic and Geometry*, *Annals of Mathematics Studies* **67** (1972), Princeton University Press.
- [8] C. A. Weibel, *An Introduction to Homological Algebra*, Cambridge University Press (1994).

This article may be accessed via WWW at <http://www.rmi.acnet.ge/hha/> or by anonymous ftp at [ftp://ftp.rmi.acnet.ge/pub/hha/volumes/2000/n3/n3.\(dvi,ps,dvi.gz,ps.gz\)](ftp://ftp.rmi.acnet.ge/pub/hha/volumes/2000/n3/n3.(dvi,ps,dvi.gz,ps.gz))

Andrew Baker a.baker@maths.gla.ac.uk <http://www.maths.gla.ac.uk/~ajb>

Department of Mathematics
University of Glasgow
Glasgow G12 8QW
Scotland.