

PRESENTATION DEPTH AND THE LIPMAN-SATHAYE JACOBIAN THEOREM

MELVIN HOCHSTER

(communicated by Clas Löfwall)

Abstract

We give a version of the theorem of Lipman and Sathaye on Jacobian ideals but with substantially weaker hypotheses. Both their version and the result here are very useful in providing explicit test elements in tight closure theory. There are two separate ways in which the hypothesis in the theorem is weakened here: one is that the larger ring is not required to be a domain, although it will be reduced. Second, the regularity condition on the smaller ring is weakened to the point where one need not assume that it is Cohen-Macaulay. Instead, a condition on the ring homomorphism is imposed that may be viewed as a relative analogue of the Serre condition S_2 : a family of such conditions is introduced and studied here. The definition is made in terms of a presentation of an algebra, but is independent of the presentation.

To Jan–Erik Roos on his sixty–fifth birthday

1. Introduction

In this paper we shall give a version of the theorem of Lipman and Sathaye on Jacobian ideals proved in [LS] (their Theorem 2). The result presented here has substantially weaker hypotheses than in [LS]. Lipman and Sathaye used their theorem to prove the Briançon-Skoda theorem [BrS] (which was first proved using the analytic criterion for membership in an ideal in [Sk]; see also [LT] for background). Their result has also turned out to be *very* important in providing explicit test elements in tight closure theory (accounts of the latter are given in [HH1–8], [Ho2-3], [Hu], and [Br]; [HH8], in particular, treats the use of the Lipman-Sathaye theorem in constructing test elements).

In the main theorem, Theorem (2.1) of §2 (the reader may want to consult its statement at this point), there are two separate ways in which the hypothesis in [LS] on the map $R \rightarrow S$ is weakened: first, while the ring S is still required to be torsion-free and generically étale over R , S is not required to be a domain. Second,

The author was supported in part by a grant from the National Science Foundation.

Received March 5, 2001; published on July 12, 2002.

2000 Mathematics Subject Classification: 13B02, 13B21, 13C15.

Key words and phrases: Jacobian theorem, algebra presentation, depth, integral extension.

© 2002, Melvin Hochster. Permission to copy for private use granted.

the regularity condition on R is relaxed even to the point where we do not need to assume that R is Cohen-Macaulay: a condition on the map $R \rightarrow S$ is needed, but it is significantly weaker than assuming that R be Cohen-Macaulay. In this connection, we introduce in (1.2) what might be thought of as a relative analogue of the Serre condition S_i that is defined in terms of a presentation S over R , but later shown to be independent of the presentation: cf. Theorem (1.8). It turns out that we can replace the Cohen-Macaulay condition on R in the Lipman-Sathaye theorem by the condition that S be relatively S_2 presentable over R in the sense defined in §1. A key point is that when $R \rightarrow S$ has this property, it is preserved when we adjoin an element of the normalization of S : see Theorem (1.9). However, we still do need to assume that R is regular when localized at a prime lying under a height one prime of the normalization of S .

The proof of Theorem (2.1) here is modeled on the “down-to-earth” proof of Lipman-Sathaye, and we shall make frequent references to [LS]. One difference is that we need many sources of elements in “very general position,” particularly indeterminate linear combinations of generators of ideals. We are able to do this by replacing R at the outset by a localization $R(t)$ of a polynomial ring in infinitely many variables over R .

As mentioned above, the next section is devoted to developing the required theory of “presentation S_i conditions.” The main result is stated in Theorem (2.1) of §2. The proof is outlined in §2, and then the details of the various steps are supplied in §§3–5.

2. DEPTH PROPERTIES IN PRESENTATIONS

Definition-Discussion 2.1. In this section fix an integer $i \geq 1$, let R be a Noetherian domain that satisfies the Serre condition S_i (if $i = 1$, the fact that R is a domain guarantees this), and let S be a reduced algebraic R -torsion-free extension of R essentially of finite type over R . Thus, if \mathcal{K} is the fraction field of R , then $\mathcal{K} \otimes_R S$ is a finite product of finite algebraic field extensions of \mathcal{K} . In later sections we shall assume as well that S is *generically étale* over R , by which we mean that $\mathcal{K} \otimes_R S$ is a finite product of finite separable algebraic extensions of \mathcal{K} . We have that as R -algebras, $S \cong T/I$, where T is a localized polynomial ring in finitely many variables over R , and we shall refer to the corresponding surjection $T \twoheadrightarrow S$ as an *L-presentation* of S over R , where the “L-” is an indication that we are permitting T to be a localization of a polynomial ring. Informally we may also say that $S \cong T/I$ is an L-presentation of S . An L-presentation is a presentation if T is a polynomial ring in finitely many variables over R .

Remark 2.2. Notice that the minimal primes of I all have the same height in this situation: since the elements of R are nonzerodivisors in S , the situation is unaffected by localizing at the multiplicative system $R - \{0\}$, and thus we may assume that R is a field and S is a finite product of finite field extensions. Suppose that T is a localization of the polynomial ring $T_0 \subseteq T$. Then the minimal primes of I must correspond to maximal ideals of T_0 (or else S would be transcendental over R) and so all have the same height.

Definition 2.3. Let $R \rightarrow S$ be as in (1.0). We shall say that an L-presentation $\eta: T \twoheadrightarrow S$ with $\text{Ker } \eta = I$ is *relatively S_i* if, with $h = \text{ht } I$, for every prime ideal P of T containing I , $\text{depth } T_P \geq \min \{\text{ht } P, h + i\}$. (Note that this implies $\text{depth}_T I = \text{ht } I$.)

It will be convenient to have a name for the sort of ideal described above.

Definition 2.4. Let T be a Noetherian ring. An ideal I of T all of whose minimal primes have height h will be said to be of *CM_{+i} type* if for every prime ideal P containing I , $\text{depth } T_P \geq \min \{\text{ht } P, h + i\}$. (Again, note that this implies $\text{depth}_T I = \text{ht } I$.)

With the terminology of (1.3) and $R \rightarrow S$ as in (1.0), an L-presentation $S \cong T/I$ over R is relatively S_i if and only if I is of CM_{+i} type in T . Since R is a domain, T has no embedded primes, and (1.1) shows that the ideal I associated with the presentation has the property that all of its minimal primes have the same height. We shall say that an ideal all of whose minimal primes have height h has *pure height h* . Evidently, a Noetherian ring T is Cohen-Macaulay if and only if every ideal of pure height is of CM_{+i} type for all i .

Although we have given Definition (1.3) without restriction on T , in the sequel we shall almost always be assuming that T satisfies the Serre condition S_i when we discuss ideals of CM_{+i} type. Likewise, in the sequel, when discussing the relatively S_i condition in (1.2), we shall almost always be assuming, as indicated in (1.0), that R satisfies the Serre condition S_i .

Discussion 2.5 (behavior of depth and dimension under flat base change). For use in the proof of the very important Lemma (1.5) that follows, we note two well known facts: if $(T, P) \rightarrow (W, Q)$ is a flat local homomorphism of local rings, and $B = W/PB$ is the closed fiber, then $\dim W = \dim T + \dim B$ ([Mat], (13.B) Theorem 19 (2), p. 79), and $\text{depth } W = \text{depth } T + \text{depth } B$ ([Mat], (21.C) Corollary 1, p. 154). This implies at once, for example, that if T satisfies S_i and W is flat over T with Cohen-Macaulay fibers (or S_i fibers), then W satisfies S_i .

We shall make extensive use of the following alternative characterization of when an ideal I is of CM_{+i} type.

Lemma 2.6. *Let T be a Noetherian ring that is S_i . Let $I \subseteq T$ be an ideal of pure height h . Let f_1, \dots, f_r be generators of I . Let $t_{\mu j}$ be indeterminates over T , $1 \leq \mu \leq h$, $1 \leq j \leq r$, and let $g_\mu = \sum_{j=1}^r t_{\mu j} f_j$. Then the following are equivalent:*

- (1) *I has CM_{+i} type.*
- (2) *In some faithfully flat extension W of T , IW has height equal to its depth h , and contains a regular sequence of length h generating an ideal J such that W/J is S_i .*
- (3) *In the polynomial ring $W = T[t_{\mu j} : \mu, j]$, the elements g_1, \dots, g_h form a regular sequence generating an ideal J such that W/J is S_i .*
- (4) *In the localization W_U of the ring W from part (3) at some (resp., every) multiplicative system U consisting of polynomials whose coefficients generate the unit ideal, the elements g_1, \dots, g_h form a regular sequence generating an ideal J such that W/J is S_i .*

Proof. We shall prove (3) \Rightarrow (2) \Rightarrow (1) \Rightarrow (3). Note that each W_U as in (4) above is faithfully flat over T , and so we also have (3) \Rightarrow ((4) for all U) \Rightarrow ((4) for some U) \Rightarrow (2) as well.

(3) \Rightarrow (2) is obvious. To see that (2) \Rightarrow (1), note that, after a faithfully flat extension $T \rightarrow W$, the ideal IW has the same height as I and the same depth as I . Thus, I has height and depth h . Now let P be a prime ideal of R containing I of height greater than h . Then there is a minimal prime Q of PW lying over P , and $T_P \rightarrow W_Q$ will be faithfully flat with a zero-dimensional fiber. The depth and dimension of W_Q are the same as the depth and dimension of T_P : cf. the discussion preceding (1.5). When we kill J in W_Q , the depth and dimension drop by h , and we obtain an S_i ring. This shows that W_Q has the required depth, and, hence, so does R_P .

It remains to prove that (1) \Rightarrow (3). The fact that the elements g_i form a regular sequence is proved in [Ho1]. Let Q be a prime containing $J = (g_1, \dots, g_h)$ in $W = T[t_{\mu j} : \mu, j]$. First suppose that Q contains I . We must show that the depth of W_Q is at least the smaller of $\text{ht } Q$, $h + i$, since this will give the correct depth once we kill J . Then Q lies over a prime P of T that contains I . Then the depth of R_P is at least $\min\{\text{ht } P, h + i\}$, and by [Mat], (13.B) Theorem 19 (2), p. 79, we have $\text{ht } Q = \text{ht } P + \dim B$, where B is the fiber W_Q/PW_Q , and $\text{depth } W_Q = \text{depth } T_P + \text{depth } B$ by [Mat], (21.C) Corollary 1, p. 154.

But the fibers of $T \rightarrow W$ are Cohen-Macaulay (in fact, regular), and so

$$\begin{aligned} \text{depth } W_Q &\geq \min\{\text{ht } P, h + i\} + \text{depth } B = \min\{\text{ht } P, h + i\} + \\ \dim B &\geq \min\{\text{ht } P + \dim B, h + i + \dim B\} = \min\{\text{ht } Q, h + i + \dim B\} \geq \\ &\min\{\text{ht } Q, h + i\}, \text{ as required.} \end{aligned}$$

It remains only to consider the case where Q fails to contain I . Then one of the generators f_j of I is a unit in T_P , where P is the contraction of Q to T . Again, we pass to T_P , and renumber so that f_r is a unit. Then the equations g_μ that we are killing simply solve for the indeterminates $t_{\mu r}$ in terms of the others, and the quotient by J may be identified with $T_P[t_{\mu j} : 1 \leq \mu \leq h, 1 \leq j < r]$, a polynomial ring over T_P . Since this ring is S_i so is the polynomial ring, and the result follows. \square

We shall later need the following fact whose proof is really a variant of the argument that (1) \Rightarrow (3) in Lemma (1.5) above.

Lemma 2.7. *Let I be an ideal of pure height h in a Noetherian ring T such that T is S_i . Let u_1, \dots, u_h be a regular sequence in I such that $T/(u_1, \dots, u_h)T$ is S_i . Let f_1, \dots, f_r be generators of I . Let k be an integer, $1 \leq k \leq h$. Let $t_{\mu j}$ be indeterminates over T , $1 \leq \mu \leq k$, $1 \leq j \leq h$ and let $t'_{\mu j}$ also be indeterminates over T , $k < \mu \leq h$, $1 \leq j \leq r$. Let W be the ring obtained by adjoining all these independent indeterminates to T . Then the k elements $g_\mu = \sum_{j=1}^h u_j t_{\mu j}$, $1 \leq \mu \leq k$, together with the $h-k$ elements $g_\mu = \sum_{j=1}^r f_j t'_{\mu j}$, $k < \mu \leq h$, give a regular sequence in IW such that the quotient of W by the ideal they generate is S_i .*

Proof. Consider a prime Q of W that contains the g 's. The specified depth condition on $W_Q/(g_1, \dots, g_h)$ is equivalent to the condition that the depth of W_Q be at least the lesser of $\text{ht } Q, h + i$.

If Q contains all the elements u_1, \dots, u_h this follows because, after we kill this regular sequence, the quotient ring $W_Q/(u_1, \dots, u_h)$ is S_i , since it is flat over $T/(u_1, \dots, u_h)$ with regular fibers. If Q fails to contain all the u_h , we replace R by its localization at the contraction P of Q to T . We may assume that in each of the first k of the g_μ , one of the u_i is invertible, permitting us to use $g_\mu = 0$ to solve for one of the $t_{\mu j}$ in terms of the others. But then $P \not\supseteq I$ as well, and so in each of the last $n - k$ of the g_μ at least one of the f_j is invertible, permitting us to solve for one of the $t_{\mu j}$ in terms of the others. This means that the quotient by (g_1, \dots, g_h) may be identified with a polynomial ring over T_P , and the result follows from the fact that T is S_i . \square

Note that if R is Cohen-Macaulay then so is T , and it is then automatic that every L-presentation is relatively S_i for all i .

Our next results are aimed at showing that when S has a relatively S_i presentation over R , then every presentation of S over R is relatively S_i .

Lemma 2.8. *Let $R \rightarrow S$ be a homomorphism as in (1.0). Let $S \cong T/I$ be an L-presentation of S as an R -algebra that is relatively S_i , and let T_0 be a polynomial ring in finitely many variables over R contained in T such that T is a localization of T_0 .*

(a) *If U is a multiplicative system in T with image V in S then T_U/I_U is a relatively S_i L-presentation of S_V over R .*

(b) *(Descent of localizations.) Assume that the S_i locus is open in rings essentially of finite type over R (which holds if R is either a homomorphic image of a Cohen-Macaulay ring or if R is excellent). If $T = (T_0)_U$ for a multiplicative system U in T_0 , there is an element $u \in U$ such that if $T_1 = T_0[1/u]$, and $I_1 = I \cap T_1$, then $S_1 = T_1/I_1$ is a relatively S_i L-presentation.*

Proof. Part (a) is clear, since the set of localizations at primes containing I can only become smaller as we localize T . It is worth noting that the condition that I have pure height is needed here: otherwise, the height of I might increase after localization.

To prove part (b), note that localizing at finitely many elements is equivalent to localizing at their product. We may make several choices of u, T_1 , each time localizing at one more element to get additional conditions to hold. Choose generators f_1, \dots, f_r for I and initially choose u so that these will be in T_1 . Thus, we may assume that $I = I_1T$. Let $h = \text{ht } I$. Introduce hr new indeterminates $t_{\mu j}$ and define g_1, \dots, g_h as in part (3) of Lemma (1.5). Then we know that $U^{-1}T_1[t_{\mu j} : \mu, j]/(g_1, \dots, g_h)$ is S_i , by Lemma (1.5). This means that U meets the defining radical ideal of the non- S_i locus in the ring $T_1[t_{\mu j} : \mu, j]/(g_1, \dots, g_h)$ (our hypothesis guarantees that this locus is closed). Thus, we may choose $u_1 \in U$ meeting that radical ideal. Replacing T_1 by its localization at u_1 , we see that we can localize at just one element and get $T_1[t_{\mu j} : \mu, j]/(g_1, \dots, g_h)$ to be S_i (this is

a new choice of T_1), and so by part (3) of Lemma (1.5), $T_1 \twoheadrightarrow T_1/I_1$ is relatively S_i for this new choice of T_1 . \square

Theorem 2.9 ((independence of presentation)). *Let $R \twoheadrightarrow S$ be a homomorphism as in (1.0) and suppose that S has an L-presentation over R that is relatively S_i . Then every L-presentation of S over R is relatively S_i .*

Proof. Step 1. We first consider the case where S is finitely generated over R and we are considering only presentations of S over R , i.e., T is required to be a polynomial ring. Given two presentations $T \twoheadrightarrow S$, $T' \twoheadrightarrow S$, we can compare both with $T \otimes_R T' \twoheadrightarrow S$. By induction on the number of variables in T' , we reduce to considering a presentation $T \twoheadrightarrow S$ and another $T[X] \twoheadrightarrow S$, where X is a new indeterminate. Let s be the image of X and let $z \in T$ map to s . We may replace X by $X - z$, and so assume that X maps to 0 in S . Thus, if $I = \text{Ker}(T \twoheadrightarrow S)$, the kernel of the second presentation will be $IT[X] + XT[X]$. The primes of $T[X]$ containing the kernel are in bijective correspondence with the primes of T containing I , and the depth condition is obviously satisfied for T_P if and only if it is satisfied for $T[X]_Q$, where $Q = PT[X] + XT[X]$ is the corresponding prime, since $T[X]_Q/(X) \cong T_P$, and X is a nonzerodivisor.

Step 2. We next consider the case where S is finitely generated over R , but can be written $S = S_0[1/s]$ where $s \in S_0$. Let $T_0 \twoheadrightarrow S_0$ be a presentation of S_0 with kernel I_0 and $v \in T_0$ an element that maps to s . Then we have a surjection $T_0[1/v] \twoheadrightarrow S$, which is an L-presentation, and we also have a presentation $T_0[X] \twoheadrightarrow S$ extending $T_0 \twoheadrightarrow S_0$, where X maps to $1/s$. The kernel of the latter is $I_1 = I_0T_0[X] + (Xv - 1)$. Again, there is a bijection between the primes of $T_0[1/v]$ containing I_0 (or primes of T_0 containing I_0 and not v) and the primes of $T_0[X]$ containing I_1 , and the depths in the localizations at the latter primes are all one greater: we can see this, because we can start by killing $Xv - 1$, which gives the correspondence. Thus an L-presentation of this type is relatively S_i presentable if and only if the corresponding presentation is relatively S_i presentable.

Step 3. We now consider the general case. Let T_0 be a polynomial ring in finitely many variables over R and U a multiplicative system in T_0 such that $U^{-1}T_0 \twoheadrightarrow S$ is a relatively S_i L-presentation of S over R . By Lemma (1.7) (b) we can choose a localization T_1 of T_0 at one element and such that the restriction of $T \twoheadrightarrow S$ gives a surjection $T_1 \twoheadrightarrow S_1$ that is a relatively S_i L-presentation, and S is a localization of S_1 . By part (a), every localization S_2 of S_1 at one element has an L-presentation $T_2 \twoheadrightarrow S_2$ that is relatively S_i and such that T_2 is a localization of a polynomial ring at one element. It follows that S is a localization of a finitely generated R -subalgebra S_1 with the following property: every presentation of S_1 , and every presentation of any localization $S_1[1/u]$, is relatively S_i .

Now consider some other L-presentation $(U')^{-1}T' \twoheadrightarrow S$. Then we can choose a localization T'_1 of T' at one element of U' whose image S'_1 in S contains S_1 , and we can choose $w \in S_1 \subseteq S'_1$ such that $S'_1[1/w] = S_1[1/w]$, where w is invertible in S . Thus, $S'_1[1/w]$ has a presentation that is relatively S_i .

Suppose that $w = \tau/y$ where $\tau \in T'$ and $y \in U'$. Since w is invertible in S , we can choose $a \in T'$ and $z \in U'$ such that $a\tau/yz = 1$ in S , i.e., and then there exists $v \in U'$ such that $va\tau = vyz$. Then S'_1 localized at the image ζ of vyz contains $1/w$, since

that can be represented as a/z , and so $S_2 = S'_1[1/\zeta]$ has the property that every presentation is relatively S_i , since it is a localization of $S_1[1/w]$ at one element, and hence a localization of S_1 at one element. But $T'_1[1/vyz]$ is a localization of T' at an element of U' mapping onto S_2 . From Step 2 we know that this L-presentation is also relatively S_i (since the corresponding presentation will be), and then by (1.7) (a), $(U')^{-1}T' \twoheadrightarrow S$ is relatively S_i . \square

The following fact will be critical in the proof of the Lipman-Sathaye Jacobian theorem.

Theorem 2.10. *Let $R \rightarrow S$ be as in (1.0). Suppose that S is relatively S_i presentable over R . Let s be an element of the total quotient ring of S that is integral over S . Then $S[s]$ is relatively S_i presentable over R . Hence, any subring of the integral closure of S module-finite over R is relatively S_i presentable over R . Moreover, if $T \twoheadrightarrow R$ is an L-presentation with kernel I of height h such that g_1, \dots, g_h is a regular sequence in I with the property that $T/(g_1, \dots, g_h)$ is S_i , and we extend $T \twoheadrightarrow S$ to $T[X] \twoheadrightarrow S[s]$ by sending X to s , then there is a polynomial F monic in X in the kernel, and $T[X]/(g_1, \dots, g_h, F)$ is S_i .*

Proof. We first establish the last statement. Let $B = T/(g_1, \dots, g_h)$ and

$$C = T[X]/(g_1, \dots, g_h, F) \cong B[X]/(f)$$

where f is the image of F in $B[X]$ and is still monic in X . Clearly, $B[X]/(f)$ is flat (and even free) over B with zero-dimensional (hence, Cohen-Macaulay) fibers, and so is S_i .

The first part now follows. After tensoring with R, S, T $R[t_{\mu_j} : \mu, j] = R[t]$ we get such a regular sequence in $T[t]$ such that the quotient is S_i , by Lemma (1.5) (3). When we adjoin X the result of the paragraph above together with Lemma (1.5) (2) shows that we have a relatively S_i presentation of $S[s]$. \square

The following fact is used in Discussion (1.11) to show that there are many case where $R \rightarrow S$ as in (1.0) is such that S is relatively S_2 presentable over R , but R is not Cohen-Macaulay.

Proposition 2.11. *Let $R \rightarrow S$ be as in (1.0) (in particular, R satisfies S_i) and let R_1 be a domain flat over R with S_i fibers. Suppose that S is relatively S_i presentable over R . Then $R_1 \rightarrow R_1 \otimes_R S$ satisfies (1.0) and is relatively S_i presentable over R_1 .*

Proof. Let $T \twoheadrightarrow S$ be an L-presentation (with kernel I of pure height h) that is relatively S_i . After adjoining indeterminates t_{m_j} as in (1.5) we obtain a regular sequence g_1, \dots, g_h in the expansion of I such that $T[t]/(g_1, \dots, g_h)$ is S_i . The main point is that $R_1 \otimes_R T[t]/(g_1, \dots, g_h)$ is S_i . Cf. (1.4). \square

Discussion 2.12. We conclude this section with an example which shows that the property that S be relatively S_2 presentable over R is weaker than the assumption that R be Cohen-Macaulay.

One way to see this is to let R_0 be Cohen-Macaulay normal of finite type over an algebraically closed field K , and let B denote a finite type K -algebra that is

normal but not Cohen-Macaulay. Take any extension of S_0 of R_0 as in (1.0). This will be relatively S_2 -presentable simply because R_0 is Cohen-Macaulay. It is easy to see that $R = B \otimes_K R_0 \rightarrow B \otimes_K S_0 = S$ is still relatively S_2 presentable (apply (1.10) with $R = R_0$ and $R_1 = B \otimes_K R_0$), but R is not Cohen-Macaulay.

3. STATEMENT OF THE THEOREM AND A SKETCH OF THE PROOF

Throughout the rest of this paper R denotes a Noetherian domain with fraction field \mathcal{K} , and S denotes an algebra essentially of finite type over R (i.e., a localization at some multiplicative system of a finitely generated R -algebra) such that S is torsion-free and generically étale over R (by which we mean that $\mathcal{L} = \mathcal{K} \otimes_R S$ is a finite product of finite separable algebraic field extensions of \mathcal{K}). We shall denote by S' the integral closure of S in \mathcal{L} . We shall see that S' is module-finite over S if R is regular or excellent (cf. (2.7)).

If A and B are subsets of \mathcal{L} we denote by $A :_{\mathcal{L}} B$ the set $\{u \in \mathcal{L} : uB \subseteq A\}$. If T is a subring of \mathcal{L} and A is a T -module, then so is $A :_{\mathcal{L}} B$.

We shall write $J_{S/R}$ for the Jacobian ideal of S over R . If S is a finitely generated R -algebra, so that we may think of S as $R[X_1, \dots, X_n]/(f_1, \dots, f_m)$, then $J_{S/R}$ is the ideal of S generated by the images of the size n minors of the Jacobian matrix $(\partial f_j / \partial x_i)$ under the surjection $R[X] \rightarrow S$. This turns out to be independent of the presentation. Moreover, if $s \in S$, then $J_{S_s/R} = J_{S/R} S_s$. From this one sees that if S is essentially of finite type over R and one defines $J_{S/R}$ by choosing a finitely generated subalgebra S_0 of S such that $S = W^{-1} S_0$ for some multiplicative system W of S_0 , and takes $J_{S/R}$ to be $J_{S_0/R} S$, then $J_{S/R}$ is independent of the choices made. The result we aim to prove is:

Theorem 3.1 (Generalized Lipman-Sathaye Jacobian theorem). *Let R be a Noetherian domain with fraction field \mathcal{K} . Assume also that the S_2 locus is open in algebras essentially of finite type over R .¹ Let S be an extension algebra essentially of finite type over R such that S is torsion-free and generically étale over R . Suppose that for every maximal ideal \mathcal{M} of S with contraction \mathcal{P} to R , $R_{\mathcal{P}}$ is normal, and $S_{\mathcal{M}}$ has a relatively S_2 presentation over $R_{\mathcal{P}}$.² Let $\mathcal{L} = \mathcal{K} \otimes_R S$ and let S' be the integral closure of S in \mathcal{L} . Assume that S' is module-finite over S .³ Suppose that for every height one prime ideal Q of S' , $R_{Q \cap R}$ is regular. Then $S' :_{\mathcal{L}} J_{S'/R} \subseteq S :_{\mathcal{L}} J_{S/R}$.*

Overview of the Proof.

(2.2) Step 1: The local case suffices Note that it is enough to prove the result when S is replaced by its various localizations at maximal ideals. Thus, we may assume that S is local, although we shall only make this assumption at certain

¹This holds automatically if R is excellent or if R is a homomorphic image of a Cohen-Macaulay ring.

²This is automatic if R is normal and Cohen-Macaulay, or if R is normal and S has a relatively S_2 presentation over R .

³This is automatic if R is regular or excellent: cf. (2.7).

points in the proof. When S is local we may also replace R by its localization at the contraction of the maximal ideal of S , and so there is likewise no loss of generality in assuming that R is local and that $R \rightarrow S$ is local homomorphism (i.e., the maximal ideal of R maps into that of S). Because of this reduction to the local case, throughout the rest of the argument we assume that R is a normal local domain and, in particular, that R satisfies S_2 .

(2.3) Step 2: Adjoining indeterminates If $\mathcal{T} = \{t_\lambda : \lambda \in \Lambda\}$ is a family of indeterminates over a Noetherian ring R we denote by $R(\mathcal{T})$ or sometimes even $R(t)$ the localization of the polynomial ring $R[t_\lambda : \lambda \in \Lambda]$ at the multiplicative system consisting of all polynomials whose coefficients generate the unit ideal. This ring is Noetherian even when the family of indeterminates is infinite (cf. [HH7], Remark 2.18, especially the final, parenthetical paragraph) and faithfully flat over R with geometrically regular fibers, since it is a direct limit of smooth R -algebras and flat maps and happens to be Noetherian. To prove Theorem (2.2) it suffices to consider $R(t) \rightarrow R(t) \otimes_R S$ instead of $R \rightarrow S$. Note that the normalization of $R(t) \otimes_R S$ is $R(t) \otimes_R S'$: it is normal because it is a localization of a polynomial ring over S' . Note that the map $R(t) \rightarrow R(t) \otimes_R S$ is the direct limit of maps $R(\mathcal{T}_0) \rightarrow R(\mathcal{T}_0) \otimes_R S$ as \mathcal{T}_0 runs through finite subsets of \mathcal{T} .

Note that if R has the property that algebras essentially of finite type over R have open S_i locus, then so does $R(t)$. It suffices to see this for algebras of finite type over $R(t)$ and each of these arise from a finite type algebra S_0 over $R(\mathcal{T}_0)$ (where \mathcal{T}_0 is finite) by a base change $R(\mathcal{T}_0) \rightarrow R(t)$ that is faithfully flat with regular fibers. But then Q in $R(t) \otimes_{R(\mathcal{T}_0)} S_0$ with contraction P to S_0 is such that the localization at Q is S_i if and only if $(S_0)_P$ has property S_i , i.e., the locus that we want in $\text{Spec}(R(t) \otimes_{R(\mathcal{T}_0)} S_0)$ is the inverse image of an open set in $\text{Spec} S_0$ (since S_0 is essentially of finite type over R).

Note also that if $R \rightarrow S$ has the property that every height one prime of S' lies over a prime P in R such that R_P is regular, the same is true for $R(t) \rightarrow R(t) \otimes_R S$. Since the latter ring is flat over S , a height one prime there lies over either a height one or height 0 prime of S : in either case, that in turn lies over a prime P_0 in R such that R_{P_0} is regular (if the contraction to S is height 0, the prime lies over 0 in R). Thus, $R(t)_P$ will be a Noetherian localization of a polynomial ring over R_{P_0} , and, hence, regular.

Thus, we may replace $R \rightarrow S$ by $R(t) \rightarrow S(t)$ in proving the theorem, where the set of indeterminates is infinite. This will prove useful in constructing regular sequences in sufficiently general position.

(2.4) Step 3: Presenting S over R Let T denote a localization of $R[X_1, \dots, X_n]$ that maps onto S , and let I denote the kernel. Let U denote the complement in T of the set of minimal primes P_1, \dots, P_r of I in T . Since S is reduced, $I = \bigcap_{i=1}^r P_i$. Since S is a torsion-free R -module, the minimal primes of I do not meet R , and correspond to the minimal primes of $I(\mathcal{K} \otimes T)$. Since killing any of these minimal primes produces an algebraic extension of \mathcal{K} , they must correspond to maximal ideals of $\mathcal{K}[X_1, \dots, X_n]$, and it follows that the P_i all have the same height, which must be the same as the number of variables, n . Thus, $U^{-1}T$ is a semilocal regular ring in which each of the maximal ideals $\mathcal{M}_i = P_i U^{-1}T$ is generated by n elements.

(2.5) Step 4: Special sequences and the modules $W_{S/R}$ Call a sequence g_1, \dots, g_n of n elements of I *special* if it generates each of the \mathcal{M}_i , is a regular sequence in T , and all of the associated primes of the ideal it generates are of height n . We shall show that special sequences exist (this is really only using that S has a relatively S_1 presentation over R), and that there are sufficiently many of them that the images of the elements $\det(\partial g_j/\partial X_i)$ in S with g_1, \dots, g_n special generate the Jacobian ideal. Moreover, when g_1, \dots, g_n is special the image of $\det(\partial g_j/\partial X_i)$ in S is not a zerodivisor in S , and so has an inverse in \mathcal{L} . Given $\theta: T \rightarrow S$ and a special sequence g_1, \dots, g_n we define a map

$$\Phi: \frac{(g_1, \dots, g_n)T: {}_T I}{(g_1, \dots, g_n)T} \rightarrow \mathcal{L}$$

by sending the class of u to \bar{u}/γ , where \bar{u} is the image of u in S and γ is the image of $\det(\partial g_j/\partial X_i)$ in S . We shall show that Φ is injective. *A priori*, its image depends on the choice of $T \rightarrow S$ and on the choice of the special sequence g_1, \dots, g_n , but the image turns out to be independent of these choices, and so we have constructed a finitely generated canonically determined S -module $W_{S/R} \subseteq \mathcal{L}$.

(2.6) Step 5: The main idea of the argument It will turn out that, quite generally, $W_{S/R} \subseteq S: {}_{\mathcal{L}} J_{S/R}$. The result then follows from two further observations. The first is that when S is normal, this is an equality. (The idea is to show that $W_{S/R}$ is then S_2 , which forces Φ with its image restricted to $S: {}_{\mathcal{L}} J_{S/R}$ to be an isomorphism — this can be verified after localizing at height one primes of S . This is the only place in the proof where we use the hypothesis that S has a relatively S_2 presentation over R .) The second is that when one enlarges S to $S_1 = S[s_1]$ by adjoining one integral fraction $s_1 \in \mathcal{L}$ (so that $S \subseteq S_1 \subseteq S'$), then $W_{S_1/R} \subseteq W_{S/R}$. Repeated application of this fact yields that $W_{S'/R} \subseteq W_{S/R}$ and then we have

$$S': {}_{\mathcal{L}} J_{S'/R} = W_{S'/R} \subseteq W_{S/R} \subseteq S: {}_{\mathcal{L}} J_{S/R},$$

and we are done. In the sequel we shall systematically fill in the details of this outline.

We conclude this section with a result, essentially in [LS] but not stated in this generality, which shows that the normalization S' of S is module-finite over S under very mild hypotheses on R .

Fact 3.2 (finiteness of the normalization). Let S be torsion-free, generically étale, and essentially of finite type over a normal Noetherian domain R . Suppose that the completion of every local ring of R is reduced (which holds if R is either regular or excellent). Then the normalization S' of S over R is module-finite over S .⁴

⁴Sketch of proof. Let $S_0 \in S$ be finite type over R with $S = (S_0)_W$. Since localization commutes with normalization, we may replace S by S_0 and assume S is finite type over R . The integral closure of S is the product of the integral closures of the domains obtained by killing a minimal prime of S . Hence, we may assume S is a domain. Each of finitely many generators for S over R satisfies an algebraic equation over R with leading coefficient r_ν , say, and so with $r = \prod_\nu r_\nu \in R - \{0\}$, $S[1/r]$ is integral over $R[1/r]$. The integral closure of the normal domain $R[1/r]$ in the fraction field \mathcal{L} of $S[1/r]$ is the same as the normalization of $S[1/r]$, and is module-finite over $S[1/r]$ by Prop. (31.B) of Chapter 12 of [Mat]. We may enlarge S by adjoining finitely many elements of its normalization and so obtain a domain S such that $S[1/r]$ is normal for some $r \neq 0$. By Lemma 4

4. BEGINNING THE PROOF: EXISTENCE OF SPECIAL SEQUENCES

Throughout the rest of this paper R denotes a normal Noetherian domain with fraction field \mathcal{K} , and S denotes an algebra essentially of finite type over R (i.e., a localization at some multiplicative system of a finitely generated R -algebra) such that S is torsion-free and generically étale over R (by which we mean that $\mathcal{L} = \mathcal{K} \otimes_R S$ is a finite product of finite separable algebraic field extensions of \mathcal{K}). We shall denote by S' the integral closure of S in \mathcal{L} , and we shall assume that it is a finite module over S . We also assume that S is relatively S_1 presentable over R . (We shall not need that it is relatively S_2 presentable until §4.) Let $T \rightarrow S$ be an L -presentation with kernel I of pure height n . Recall that a regular sequence g_1, \dots, g_n of length n in I is *special* if g_1, \dots, g_n generate IT_P for every minimal prime P of I , and all minimal primes of $(g_1, \dots, g_n)T$ have height n . We construct many special sequences after replacing R, S, T by their tensor products over R with $R(T)$, where T is consists of infinitely many new indeterminates as in (2.3). We denote the results of tensoring R, S, T with $R(T) = R_1$ by R_1, S_1 , and T_1 , respectively. We write I_1 for the expansion of I to T_1 , so that $S_1 \cong T_1/I_1$. In later sections we shall change notation and write R, S, T for the rings denoted R_1, S_1, T_1 here.

Lemma 4.1. *With hypothesis as in the paragraph above, there are sufficiently many special sequences g_1, \dots, g_n in I_1 that the Jacobian ideal J_{S_1/R_1} is generated by the images of elements $\det(\partial g_j / \partial X_k)$ for g_1, \dots, g_n special (where the X_k are the indeterminates generating the polynomial ring of which T is the localization). Moreover, if T_1 is local, given any two special sequences there is a finite sequence (which we shall refer to as a chain) of special sequences with one of them as first term and the other as last term with the following property: given any two consecutive terms, one is obtained from the other by letting an invertible matrix over R_1 act, or else the two consecutive sequences differ in only one term.*

Proof. Pick generators f_1, \dots, f_r for I . Let $[t_{\mu j}]$ be an $r \times r$ matrix of indeterminates from the infinite set \mathcal{T} . The elements $g_\mu = \sum_{j=1}^r f_j t_{\mu j}$ give a new set of generators for I_1 , because the determinant of $[t_{\mu j}]$ is invertible in R_1 . By Lemma (1.5) applied with $i = 1$, any n of these generators form a regular sequence $W = T[t_{\mu j} : \mu j]$ such that the ideal they generate has no embedded primes (the last follows from the fact that the quotient is S_1). Because the map $W \rightarrow T_1$ is flat with Cohen-Macaulay fibers, the sequence retains these properties. Next, we want check that any n of these generate after localizing at a minimal prime P of I_1 . Since they generate I_1 ,

of Ch. 12 of [Mat], S has finite normalization if S_Q does for every maximal ideal Q of S . Choose $s \in S$ with $\mathcal{L} = \mathcal{K}[s]$. Let $R_1 = R[s]$ and $P = Q \cap R_1$. Then S_Q is a localization of $(R_1)_P[S]$, and S is generated over $(R_1)_P$ by elements of its fraction field. By a result of [Rees], if the completion of a local domain B is reduced, then the normalization of any algebra C finitely generated over B by fractions is module-finite over C . Thus, it suffices to show that the completion of $(R_1)_P$ is reduced. We may replace R by its localization at the contraction of P , and so we may assume that (R, m) is local with reduced completion. The completion of $(R_1)_P$ is one of the local rings of the completion of R_1 with respect to m . Thus, it suffices if this completion of R_1 is reduced. But this is $R_1 \otimes_R \hat{R} \subseteq \mathcal{L} \otimes_R \hat{R} \cong \mathcal{L} \otimes_{\mathcal{K}} (\mathcal{K} \otimes_R \hat{R})$, and the result follows because $\mathcal{K} \otimes_R \hat{R}$ is reduced and \mathcal{L}/\mathcal{K} is separable.

it is certainly true that we can choose n of them to generate $I_1(T_1)_P$. Given the symmetric roles played by the indeterminates, it must be true that any n generate. Since we may use all n elements subsets of this set of generators of I_1 , we obtain enough elements to generate the Jacobian ideal.

Now suppose that we have two special sequences in I_1 . They only involve finitely many of the indeterminates in \mathcal{T} . After replacing R by $R(\mathcal{T}_0)$ for a suitable finite subset \mathcal{T}_0 of \mathcal{T} , and making corresponding changes in S, T , we may assume without loss of generality that the two special sequences are in T .

We choose $n^2 + nr$ new indeterminates from \mathcal{T} to construct two $n \times n$ matrices α, β of indeterminates from \mathcal{T} as well as an $n \times r$ matrix γ . We replace one special sequence by letting α act on it: call it a_1, \dots, a_n . We replace the other by letting β act on it: call the result b_1, \dots, b_n . Also let γ act on the f_1, \dots, f_r to form n linear combinations, say g_1, \dots, g_n , of the f_1, \dots, f_r with indeterminate coefficients. If we replace the terms in a_1, \dots, a_n one at a time by the f 's, we make a chain from a_1, \dots, a_n to f_1, \dots, f_n . We can then reverse the process to make a chain from f_1, \dots, f_n to b_1, \dots, b_n , changing only one term at a time. The fact that each of the sequences formed is a regular sequence generating an ideal with no embedded primes is a consequence of Lemma (1.6). The proof that these sequence all generate $I_1(T_1)_P = P(T_1)P$ for each minimal prime P of I_1 reduces to studying the vector space $V = P(T_1)_P/P^2(T_1)_P$. One needs to know that given two sets of generators for this vector space, one can take general linear combinations of the first set (i.e., with indeterminate coefficients), say v_1, \dots, v_k of them, together with general linear combinations of the second, and if the number of these is $\dim V$, they will be a basis. The point is that the elements v_1, \dots, v_k form part of a basis. To decide whether the remaining elements complete the basis one may pass to $V/\text{Span}\{v_1, \dots, v_k\}$. Since we are working with elements that span this space, this is clear. \square

5. THE MAP Φ AND THE MODULES $W_{S/R}$

Our next main goal is to construct the maps Φ mentioned briefly in (2.5). Throughout this section we assume that R is a normal Noetherian domain, with fraction field \mathcal{K} , that S is a torsion-free generically étale R -algebra with total quotient ring \mathcal{L} (by hypothesis, a finite product of separable field extensions of \mathcal{K}), that T is a localization of the polynomial ring $R[X_1, \dots, X_n]$ that maps onto S , with kernel I , and that the minimal primes of I in T are P_1, \dots, P_r . We also assume that S is relatively S_1 presentable over R . However, in parts (c), (d) and (f) of Theorem (4.4), we shall need the stronger hypothesis that S is relatively S_2 presentable over R . Furthermore, we assume that $R \rightarrow S$ has arisen from $R_0 \rightarrow S_0$ satisfying the same conditions by tensoring with $R = R_0(\mathcal{T})$ for some infinite family \mathcal{T} of indeterminates, so that the results of Lemma (3.1) are applicable.

Lemma 5.1. *Let $g_1, \dots, g_n \in I$. Let X denote X_1, \dots, X_n . Then the images of the g_j in $R[X]_{P_\nu}$ generate $P_\nu R[X]_{P_\nu}$ if and only if $\det(\partial g_j / \partial X_i) \notin P_\nu$. (The same holds if we only assume that $g_1, \dots, g_n \in P_\nu$.) Hence, if g_1, \dots, g_n is a special sequence in I , then the image γ of $\det(\partial g_j / \partial X_i)$ is not a zerodivisor in S , and so represents an invertible element of the total quotient ring \mathcal{L} of S .*

Proof. The last statement clearly follows from the first, and we prove the first under the assumption that $g_1, \dots, g_n \in P = P_\nu$. Let \mathcal{M} be the maximal ideal corresponding to P in $\mathcal{K}[X]$. Then $PR[X]_P \cong \mathcal{K}[X]_{\mathcal{M}}$, and so this is really a statement about when elements of \mathcal{M} generate \mathcal{M} in $\mathcal{K}[X]$. For the rest of this proof we shall write \mathcal{L} (instead of, say, \mathcal{L}_ν) for the residue field $R[X]_P$, which is the same as the residue field of $\mathcal{K}[X]_{\mathcal{M}}$. *It is critical in what follows that \mathcal{L} is separable over \mathcal{K} .* Consider the universal \mathcal{K} -derivation $d : \mathcal{K}[X] \rightarrow \Omega_{\mathcal{K}[X]/\mathcal{K}}$, the module of Kähler differentials, which is the free $\mathcal{K}[X]$ -module generated by the elements dX_1, \dots, dX_n . Of course, if $f \in \mathcal{K}[X]$ then $df = \sum_{j=1}^n (\partial f / \partial x_j) dx_j$. The restriction of d to \mathcal{M} gives a \mathcal{K} -linear map $\mathcal{M} \rightarrow \Omega_{\mathcal{K}[X]/\mathcal{K}}$, and by the defining property of a derivation it sends $\mathcal{M}^2 \rightarrow \mathcal{M}\Omega_{\mathcal{K}[X]/\mathcal{K}}$. Thus, there is an induced map of \mathcal{K} -vector spaces

$$\delta : \mathcal{M}/\mathcal{M}^2 \rightarrow \mathcal{L} \otimes_{\mathcal{K}[X]} \Omega_{\mathcal{K}[X]/\mathcal{K}}.$$

Both modules are \mathcal{L} -vector spaces and it follows from the defining property of a derivation that δ is actually \mathcal{L} -linear. Since $\mathcal{K}[X]_{\mathcal{M}}$ is regular of dimension n , $\mathcal{M}/\mathcal{M}^2$ is an n -dimensional vector space over \mathcal{L} . The key point is that under the hypothesis that \mathcal{L} is separable over \mathcal{K} , the map δ is an isomorphism of \mathcal{L} -vector spaces. This is well known, but we give a very short proof. The map δ sends the elements represented by generators g_1, \dots, g_n for \mathcal{M} to the elements represented by the dg_j , and so it has a matrix which is the image of the matrix $(\partial g_j / \partial x_i)$ after mapping the entries to \mathcal{L} . Thus, δ is an isomorphism if and only if the Jacobian determinant $\det(\partial g_j / \partial x_i)$ has nonzero image in \mathcal{L} . But this determinant generates $J_{\mathcal{L}/\mathcal{K}}$, and so δ is an isomorphism if and only if the Jacobian ideal of \mathcal{L} over \mathcal{K} is \mathcal{L} . But we may use any presentation of \mathcal{L} over \mathcal{K} to calculate $J_{\mathcal{L}/\mathcal{K}}$, and so we may instead use $\mathcal{L} \cong \mathcal{K}[Z]/f(Z)$ where Z here represents just one variable and where f is a single separable polynomial. The Jacobian determinant is then the value of $f'(Z)$ in \mathcal{L} , which is not zero by virtue of the separability.

Thus, δ is an \mathcal{L} -isomorphism. Moreover, we have already seen that if g_1, \dots, g_n are generators of \mathcal{M} then the Jacobian determinant is not 0 in \mathcal{L} . But the converse is also clear, because if g_1, \dots, g_n are any elements of \mathcal{M} , they generate \mathcal{M} if and only if their images in $\mathcal{M}/\mathcal{M}^2$ span this vector space over \mathcal{L} , by Nakayama's lemma, and this will be the case if and only if their further images in $\mathcal{L} \otimes_{\mathcal{K}[X]} \Omega_{\mathcal{K}[X]/\mathcal{K}}$ span that vector space over \mathcal{L} , since δ is an isomorphism, i.e., if and only if the images of the dg_j span. But this is equivalent to the assertion that the images of the columns of the matrix $(\partial g_j / \partial x_i)$, after the entries are mapped to \mathcal{L} , span an n -dimensional space, i.e., to the nonvanishing of $\det(\partial g_j / \partial x_i)$ in \mathcal{L} , which is, of course, equivalent to its not being in P . □

(4.2) The definition of θ and $W_{S/R}$ We continue the conventions in the first paragraph of this section, but because we shall let both S and its presentation vary we shall write θ for the map $T \rightarrow S$ and we shall denote by g a special sequence g_1, \dots, g_n in I . We may then temporarily define

$$\Phi_{\theta,g} : \frac{(g_1, \dots, g_n)T : {}_T I}{(g_1, \dots, g_n)T} \rightarrow \mathcal{L}$$

by sending the class of u to \bar{u}/γ where \bar{u} is the image of u in \mathcal{L} , and γ is the image of $\det(\partial g_j/\partial x_i)$ in \mathcal{L} : the element γ is invertible in \mathcal{L} by Lemma (4.1). We shall often write Φ when θ and g are understood. We shall soon show that the image of Φ is contained in $S : \mathcal{L}J_{S/R}$. Once this is established we shall change the definition of Φ very slightly by restricting its range to be $S : \mathcal{L}J_{S/R} \subseteq \mathcal{L}$.

We note that $((g_1, \dots, g_n)T : {}_T I)/(g_1, \dots, g_n)T \cong \text{Hom}_T(T/I, T/(g_1, \dots, g_n)T)$.

We shall denote the image of $\Phi_{\theta,g}$ in \mathcal{L} by $W_{S/R}(\theta, g)$. However, we shall see just below that it is independent of the choices of θ and g , and once we know this we shall simply write it as $W_{S/R} \subseteq \mathcal{L}$.

Lemma 5.2. *The map $\Phi_{\theta,g}$ is injective, and its image in \mathcal{L} is independent of the choice of g , and of the choice of θ . Its image is contained in $S : \mathcal{L}J_{S/R}$.*

Proof. The domain of Φ may be thought of as the submodule of $T/(g_1, \dots, g_n)T$ consisting of elements killed by I . Since g_1, \dots, g_n is a special sequence, the associated primes of g_1, \dots, g_n are all minimal, and it suffices to show that the map is injective after localizing at each minimal prime of $(g_1, \dots, g_n)T$. If the prime does not contain I the domain of Φ becomes 0 after localization and there is nothing to prove. If the prime contains I it is one of the P_ν , and after localization the map becomes the composition of the isomorphism $T_{P_\nu}/P_\nu T_{P_\nu} \cong \mathcal{L}_{P_\nu}$ with the automorphism of \mathcal{L}_{P_ν} induced by multiplication by the image of γ , which is nonzero.

To prove for a fixed presentation that the map is independent of the choice of special sequence suppose that we have two special sequences that yield maps with different images. We can preserve the fact that the images are different while localizing at a suitable prime or even maximal ideal of T : S is replaced by its localization at a corresponding prime. Thus, there is no loss of generality in assuming that T and S are local. The sequences in question remain special as we localize. But we may now apply Lemma (3.1) to conclude the existence of a finite chain of special sequences joining the two that we are comparing such that any two consecutive sequences differ either in at most one spot, or by the action of an invertible matrix over R . Thus, we need only make the comparison when the two sequences differ in just one term, and since the sequences are permutable we may assume without loss of generality that one of them is g_1, \dots, g_n and the other is h_1, \dots, h_n where $h_j = g_j$ for $j \geq 2$. We set up an isomorphism

$$\sigma: \frac{(g_1, \dots, g_n)T : {}_T I}{(g_1, \dots, g_n)T} \cong \frac{(h_1, \dots, h_n)T : {}_T I}{(h_1, \dots, h_n)T}$$

as follows. If u is an element of $(g_1, \dots, g_n)T : {}_T I$ then since $uI \subseteq (g_1, \dots, g_n)T$ we may write $uh_1 = vg_1 + w$, where $w \in (g_2, \dots, g_n)T = (h_2, \dots, h_n)T$. We map the class of u to the class of v . Note that any such v is automatically in $(h_1, \dots, h_n)T : {}_T I$. (If $a \in I$ then $avg_1 + aw = auh_1$. Now, $au = bg_1 + w'$ where $w' \in (g_2, \dots, g_n)T$ and so $avg_1 + aw = bg_1h_1 + w'h_1$ and so $g_1(av - bh_1) = w'h_1 - aw \in (g_2, \dots, g_n)T$. Since g_1 is not a zerodivisor on $(g_2, \dots, g_n)T$, we have that $av - bh_1 \in (g_2, \dots, g_n)T$, and so $av \in (h_1, g_2, \dots, g_n)T = (h_1, \dots, h_n)T$, as required, for all $a \in I$.) Next note that the choice of v given u is unique modulo $(g_2, \dots, g_n)T$. Thus, we have defined a map from $(g_1, \dots, g_n)T : {}_T I$ to

$((h_1, \dots, h_n)T : {}_T I) / (h_1, \dots, h_n)T$. It is easy to see that this map kills $(g_1, \dots, g_n)T$ and so we have a map as required. It is also clear that if we similarly define a map

$$\frac{(h_1, \dots, h_n)T : {}_T I}{(h_1, \dots, h_n)T} \rightarrow \frac{(g_1, \dots, g_n)T : {}_T I}{(g_1, \dots, g_n)T}$$

it will be an inverse for the map already constructed.

To complete the proof of the independence of the image from the choice of special sequence we note that the following diagram commutes:

$$\begin{array}{ccc} \frac{(g_1, \dots, g_n)T : {}_T I}{(g_1, \dots, g_n)T} & \xrightarrow{\sigma} & \frac{(h_1, \dots, h_n)T : {}_T I}{(h_1, \dots, h_n)T} \\ \Phi_{\theta, g} \downarrow & & \downarrow \Phi_{\theta, h} \\ \mathcal{L} & \xrightarrow{\mathbf{1}_{\mathcal{L}}} & \mathcal{L} \end{array}$$

To see this, one simply needs to see that if

$$(*) \quad uh_1 - vg_1 = \sum_{j=2}^n t_j g_j$$

in T , then $\bar{u}/\gamma = \bar{v}/\eta$ in cL , where γ, η are the respective images of $\det(\partial g_j/\partial X_i)$ and $\det(\partial h_j/\partial X_i)$ in \mathcal{L} , i.e., that $u \det(\partial h_j/\partial X_i) \equiv v \det(\partial g_j/\partial X_i)$ modulo I . By differentiating $(*)$ with respect to each X_j in turn and using the fact that all the g_j and h_j are in I , we see that

$$u\nabla h_1 - v\nabla g_1 \equiv \sum_{j=2}^n t_j \nabla g_j \quad \text{modulo } I$$

which implies $(*)$.

For a given special sequence g it is obvious from the definition of $\Phi_{\theta, g}$ that γ multiplies the image of $\Phi_{\theta, g}$ into $S \subseteq \mathcal{L}$. Since the image is independent of the choice of special sequence, and since by Lemma (3.1) as the special sequence varies the values of γ generate $J_{S/R}$, it follows that the image is contained in $S : {}_{\mathcal{L}} J_{S/R}$.

It remains only to prove that the image of $\Phi_{T, g}$ is independent of the choice of $\theta : T \rightarrow S$ as well. We first consider the case of a finitely generated R -algebra S . The choice of a presentation is equivalent to the choice of a finite set of generators for S over R . We can compare the results from each of two different presentations with the result from their union, and so it suffices to see what happens when we enlarge a set of generators. By induction, it suffices to show that the image does not change when we enlarge a set of generators by one element, and so we may assume that we have $\theta : T = R[X_1, \dots, X_n] \rightarrow S$ and an extension of $\theta, \theta' : T[X_{n+1}] \rightarrow S$ by sending X_{n+1} to s . Let $T' = T[X_{n+1}]$. We can choose an element $F \in T$ such that F maps to s in S , and it follows easily that the kernel I' of θ' is $I + (X_{n+1} - F)$. It also follows easily that if $g = g_1, \dots, g_n$ is special in I then $g' = g_1, \dots, g_{n+1}$ with $g_{n+1} = X_{n+1} - F$ is a special sequence in I' . The larger (size $n + 1$) Jacobian matrix has the same determinant γ as the size n Jacobian matrix of g_1, \dots, g_n with

respect to X_1, \dots, X_n , and it is easy to check that there is an isomorphism

$$\tau: \frac{(g_1, \dots, g_n)T : TI}{(g_1, \dots, g_n)T} \cong \frac{(g_1, \dots, g_{n+1})T' : T'I'}{(g_1, \dots, g_{n+1})T'}$$

which is induced by the inclusion $(g_1, \dots, g_n)T : TI \subseteq (g_1, \dots, g_{n+1})T' : T'I'$. Since the Jacobian determinants are the same we have a commutative diagram

$$\begin{array}{ccc} \frac{(g_1, \dots, g_n)T : TI}{(g_1, \dots, g_n)T} & \xrightarrow{\tau} & \frac{(g_1, \dots, g_{n+1})T' : T'I'}{(g_1, \dots, g_{n+1})T'} \\ \Phi_{\theta, g} \downarrow & & \downarrow \Phi_{\theta', g'} \\ \mathcal{L} & \xrightarrow{1_{\mathcal{L}}} & \mathcal{L} \end{array}$$

and this yields that the images are the same.

We have now justified the notation $W_{S/R}$ when S is finitely generated over R . We leave it to the reader to verify that if s is a nonzerodivisor in S , then $W_{S[s^{-1}]/R} = (W_{S/R})_s$, and that $W_{S/R}(\theta)$ is independent of θ when S is essentially of finite type over R . \square

Proposition 5.3. *Proposition (4.4) Let S be generically étale, torsion-free and essentially of finite type over the Noetherian normal domain R . Assume that S is relatively S_1 presentable over R . Let $W = W_{S/R}$.*

- (a) *For any multiplicative system U in S , $W_{U^{-1}S/R} = U^{-1}W$.*
- (b) *W is torsion-free over S .*

For the remaining parts, assume that S is relatively S_2 presentable over R (automatic in part (e), since R is regular).

- (c) *For every prime ideal P of S , if u, v is part of a system of parameters for S_P then it is a regular sequence on W_P . (Thus, W is S_2 .)*
- (d) *If $W \subseteq W' \subseteq \mathcal{L}$ and $W_P = W'_P$ for all height one primes of S and for all minimal primes of S that are also maximal ideals, then $W = W'$.*
- (e) *If $R \rightarrow S$ is a local homomorphism of regular local rings then $J_{S/R}$ is principal and $W = S :_{\mathcal{L}} J_{S/R}$.*
- (f) *If S is normal and R_P is regular for every prime ideal P of R lying under a height one prime ideal Q of S , then $W = S :_{\mathcal{L}} J_{S/R}$.*

Proof. Part (a) is essentially the last part of (4.3), while (b) is evident from the fact that $W \subseteq \mathcal{L}$, by definition.

To prove (c) note that by (a) we may assume that S is local and that u, v is part of a system of parameters. We may choose a presentation $\theta: T \rightarrow S$ and think of W as $\cong ((g_1, \dots, g_n)T : TI)/(g_1, \dots, g_n)T$, where the sequence g_1, \dots, g_n is not only special, but has the additional property that $T/(g_1, \dots, g_n)$ is S_2 . Let $u_0, v_0 \in T$ be representatives of u, v . Then $u_0 + I$ cannot be contained in the union of the associated primes of (g_1, \dots, g_n) (these are the same as the minimal primes), or else it will be contained in one of them by [Kap], Theorem 124. Since this will contain I , it will be a minimal prime of I , and contradicts the statement that u is part of a system of parameters in $S = T/I$. Thus, we can replace u_0 by an element u_1 representing u such that g_1, \dots, g_n, u_1 is a regular sequence in T . Similarly, $v_0 + I$

cannot be contained in the union of the associated primes of $(g_1, \dots, g_n, u_1)T$, or else it is contained in one of them, say Q . Thinking modulo I , we see that, since Q contains u_0, v_0 , it has height at least two more than I . But then Q has depth at least $n + 2$, a contradiction, since it is supposedly an associated prime of an ideal generated by a regular sequence of length $n + 1$. Thus, we may choose u_1, v_1 in T representing u, v respectively and such that $g_1, \dots, g_n, u_1, v_1$ is a regular sequence. Clearly, u_1, v_1 form a regular sequence on $T/(g_1, \dots, g_n)T$. We claim they also form a regular sequence on the set of elements killed by I . It is clear that u_1 remains not a zerodivisor on this set. Suppose that $v_1z = u_1y$ where z, y are killed by I . Then $z = u_1x, y = -v_1x$ where, *a priori*, $x \in T/(g_1, \dots, g_n)T$. But $Iz = 0$ and so $Iu_1x = 0$, and since u_1 is not a zerodivisor on $T/(g_1, \dots, g_n)T$, it follows that $Ix = 0$ as well.

Part (d) is a standard consequence of what we proved in part (c), but we give a short argument. If $W \neq W'$ we can localize at a minimal prime of the support of W'/W and preserve the counterexample. By hypothesis, this prime cannot have height one (nor height 0, since if a height 0 prime is not maximal then we can localize at it in two steps: first localize at a height one prime that contains it). Thus, we may assume that S is local of height two or more, and that W'/W is a nonzero module of finite length. It follows that we can choose an element $x \in W' - W$ and part of a system of parameters u, v for S such that uz and vz are in W . The relations $v(uz) = u(vz)$ over W together with part (c) show that $uz \in uW$, and it follows that $z \in W$ after all, a contradiction.

To prove (e) note that when R is regular so is T , and so $T \rightarrow S$ will be a surjection of local rings. The kernel of such a surjection must be generated by part of a minimal set of generators for the maximal ideal of T . It follows that I is a prime and we have $I = (g_1, \dots, g_n)T$ is itself generated by a suitable special sequence. Then $J_{S/R}$ is generated by $\gamma = \det(\partial g_j / \partial X_i)$, and $((g_1, \dots, g_n)T : {}_T I) / (g_1, \dots, g_n)T = (I : {}_T I) / I = T / I = S$ and Φ sends 1 to $\frac{1}{\gamma}$, so that $W = S \frac{1}{\gamma}$, and one sees that $S : {}_{\mathcal{L}} J_{S/R} = S : {}_{\mathcal{L}} \gamma S = W$, as claimed.

To prove (f) it suffices by (d) to consider the problem after localizing at a height one or zero prime Q of S , and, without affecting the issue, one may also localize R at its contraction. If the prime of S has height 0, so does its contraction to R , and both rings become regular after localization. If the prime of S has height one, then, again, both rings become regular after localization, S because it is normal and R by hypothesis. In either case the result follows from part (e). \square

6. A CRITICAL LEMMA AND THE FINAL STEP OF THE PROOF

The following result of Lipman and Sathaye is critical in establishing that $W_{S/R}$ decreases as S is increased by adjoining integral fractions.

Lemma 6.1 (Lipman-Sathaye). *Let T be a commutative ring, Y an indeterminate, and J an ideal of $T[Y]$ such that J contains a monic polynomial h in Y of degree d , and such that J also contains an element of the form $\alpha Y - \beta$ where $\alpha, \beta \in T$ are such that $J : {}_{T[Y]} \alpha T[Y] = J$, i.e., such that α is not a zerodivisor mod-*

ulo J . Let $G \subseteq T$ be an ideal of T with $G \subseteq J$. Then for every element $v \in T[Y]$ such that $vJ \subseteq (h, G)T[Y]$ there is a $u \in T$ such that $u(J \cap T) \subseteq G$ and such that $v \equiv u \frac{\partial h}{\partial Y}$ modulo J .

Proof. See Lemma (3.17) on p. 216 of [LS]. □

We now use this to prove:

Theorem 6.2. *If S_1 is obtained from S by adjoining finitely many integral fractions of \mathcal{L} , then $W_{S_1/R} \subseteq W_{S/R}$.*

Proof. By induction on the number of fractions adjoined, it is obviously sufficient to prove this when $S_1 = S[\lambda]$, where λ is a single element of \mathcal{L} . Choose a presentation $\theta: T \rightarrow S$ and a special sequence g_1, \dots, g_n in the kernel I . Let Y be a new indeterminate and extend θ to a map $T[Y] \rightarrow S[\lambda]$ by sending Y to λ . Since λ is integral over S there is a monic polynomial $h = h(Y) \in T[Y]$ of degree say, d , in the kernel J of $T[Y] \rightarrow S[\lambda]$. If $\lambda \in S$ there is nothing to prove so that we may assume that $d \geq 2$. Since λ is in \mathcal{L} we may also choose α and β in T with α not a zerodivisor on I such that $\alpha Y - \beta$ is in the kernel. Consider the image of $h(Y)$ in $S[Y]$. There will be a certain subset of the minimal primes of S such that the image of λ is a multiple root of the image of h modulo those primes. If that set of primes is empty, we shall not alter h . If it is not empty choose an element of S that is not in any of those minimal primes but that is in the others, and represent it by an element $t \in T$. Then $h(Y) + t(\alpha Y - \beta)$ has the property that its image modulo any minimal prime of S has the image of λ as a simple root, and so we may assume, using this polynomial in place of the original choice of h , that h is a monic polynomial of degree $d \geq 2$ such that image of λ modulo every minimal prime of \mathcal{L} is a simple root of the image of h .

Because h is monic in Y , the sequence g_1, \dots, g_n, h is a regular sequence, and the Jacobian determinant with respect to X_1, \dots, X_n, Y is $\gamma \frac{\partial h}{\partial Y}$, where γ is $\det(\partial g_j / \partial X_i)$. Our choice of h implies that $\frac{\partial h}{\partial Y}$ has image that is not in any minimal prime of \mathcal{L} , and it follows, using Lemma (1.9), that g_1, \dots, g_n, h is a special sequence in J and can be used to calculate $W_{S[\lambda]/R}$. Let $v \in (g_1, \dots, g_n, h)T[Y] : T[Y]J$. We may now apply Lemma (5.1) with this $T, Y, J, v, \alpha, \beta$ and h , while taking $G = (g_1, \dots, g_n)T$. Note that $J \cap T = I$. Now, v gives rise to a typical element, the image of $v / (\gamma \frac{\partial h}{\partial Y})$ in \mathcal{L} , in $W_{S[\lambda]/R}$, and we want to show that this element is in $W_{S/R}$. Pick u as in Lemma (5.1). Then $u \in (g_1, \dots, g_n)T : T I$ and since $v \equiv u \frac{\partial h}{\partial Y}$ modulo J , this image is the same as the image of $(u \frac{\partial h}{\partial Y}) / (\gamma \frac{\partial h}{\partial Y}) = u/\gamma$, and so is in $W_{S/R}$, as required. □

The proof of the main theorem. Theorem (5.2) and Proposition (4.4f) justify the argument given much earlier in (2.6), and the proof of Theorem (2.1) is now complete. □

References

- [1] J. Briançon and H. Skoda, *Sur la clôture intégrale d'un idéal de germes de fonctions holomorphes en un point de C^n* , C.R. Acad. Sci. Paris Sér. A, **278**(1974), 949–951.
- [2] W. Bruns *Tight closure*, Bull. Amer. Math. Soc., **33**(1996), 447–458.
- [3] M. Hochster, *Properties of Noetherian rings stable under general grade reduction*, Arch. Math. **24**(1973), 393–396.
- [4] -----, *Tight closure in equal characteristic, big Cohen-Macaulay algebras, and solid closure* in Commutative Algebra: Syzygies, Multiplicities and Birational Algebra, Contemp. Math. **159**, Amer. Math. Soc., Providence, R. I., 1994, 173–196.
- [5] -----, *The notion of tight closure in equal characteristic zero* in Proc. of the CBMS Conference on Tight Closure and Its Applications (Fargo, North Dakota, July, 1995), Appendix to the notes on the main lectures by Craig Huneke, C.B.M.S. Regional Conference Series, Amer. Math. Soc., Providence, R.I., 1996.
- [6] M. Hochster and C. Huneke, *Tightly closed ideals*, Bull. Amer. Math. Soc. **18**(1988), 45–48.
- [7] -----, *Tight closure* in Commutative Algebra, Math. Sci. Research Inst. Publ., Springer-Verlag, New York-Berlin-Heidelberg, **15**(1989), 305–324.
- [8] -----, *Tight closure and strong F -regularity*, Mémoires de la Société Mathématique de France, numéro **38** (1989), 119–133.
- [9] -----, *Tight closure, invariant theory, and the Briançon-Skoda theorem*, J. Amer. Math. Soc. **3**(1990), 31–116.
- [10] -----, *Infinite integral extensions and big Cohen-Macaulay algebras*, Annals of Math. **135**(1992), 53–89.
- [11] -----, *F -regularity, test elements, and smooth base change*, Trans. Amer. Math. Soc. **346**(1994), 1–62.
- [12] -----, *Localization and test exponents for tight closure*, Michigan Math. J. **48**(2000), 305–329.
- [13] -----, *Tight closure in equal characteristic zero*, preliminary preprint.
- [14] C. Huneke, *Tight Closure and Its Applications*, Proc. of the CBMS Conference held at Fargo, North Dakota, July, 1995, C.B.M.S. Regional Conference Series, Amer. Math. Soc., Providence, R.I., 1996.
- [15] I. Kaplansky, *Commutative rings*, Revised Edition, The Univ. of Chicago Press, Chicago and London, 1974.
- [16] J. Lipman and A. Sathaye, *Jacobian ideals and a theorem of Briançon-Skoda*, Michigan Math. J. **28**(1981), 199–222.
- [17] J. Lipman and B. Teissier, *Pseudo-rational local rings and a theorem of Briançon-Skoda about integral closures of ideals*, Michigan Math. J. **28**(1981), 97–116.

- [18] H. Matsumura, *Commutative Algebra*, W.A. Benjamin, New York, 1970.
- [19] D. Rees, *A note on analytically unramified local rings*, J. London Math. Soc. **36**(1961), 24–28.
- [20] H. Skoda, *Applications des techniques L^2 a la théorie des idéaux d'une algèbre de fonctions holomorphes avec poids*, Ann. Scient. Ec. Norm. Sup. 4ème série, t. **5**(1972), 545–579.

This article may be accessed via WWW at <http://www.rmi.acnet.ge/hha/>
or by anonymous ftp at
[ftp://ftp.rmi.acnet.ge/pub/hha/volumes/2002/n2a13/v4n2a13.\(dvi,ps,pdf\)](ftp://ftp.rmi.acnet.ge/pub/hha/volumes/2002/n2a13/v4n2a13.(dvi,ps,pdf))

Melvin Hochster hochster@math.lsa.umich.edu
Department of Mathematics
University of Michigan
Ann Arbor, MI 48109–1109
USA