

HOMOTOPY COLIMITS OF DIAGRAMS OVER POSETS AND VARIATIONS ON A THEOREM OF THOMASON

XIMENA FERNÁNDEZ AND ELÍAS GABRIEL MINIAN

(communicated by Ronald Brown)

Abstract

We use a classical result of McCord and reduction methods of finite spaces to prove a generalization of Thomason’s theorem on homotopy colimits over posets. In particular, this allows us to characterize the homotopy colimits of diagrams of simplicial complexes in terms of the Grothendieck construction on the diagrams of their face posets. We also derive analogues of well known results on homotopy colimits in the combinatorial setting, including a cofinality theorem and a generalization of Quillen’s Theorem A for posets.

1. Introduction

Let \mathcal{C} be a small category and let $X: \mathcal{C} \rightarrow \text{CAT}$ be a functor to the category CAT of small categories. Recall that the *Grothendieck construction* on X , which is usually denoted by $\mathcal{C} \int X$, is the following category. The objects are the pairs (c, x) , where c is an object of \mathcal{C} and x is an object of $X(c)$, and the morphisms $(\alpha, \beta): (c, x) \rightarrow (c', x')$ are given by morphisms $\alpha: c \rightarrow c'$ in \mathcal{C} and $\beta: X(\alpha)(x) \rightarrow x'$ in $X(c')$. Thomason’s theorem [15] establishes the existence of a natural homotopy equivalence

$$\text{hocolim } \mathcal{N}X \rightarrow \mathcal{N}(\mathcal{C} \int X),$$

from the homotopy colimit of the nerve of X to the nerve of the Grothendieck construction.

In this article we focus our attention on homotopy colimits of diagrams of spaces indexed by finite partially ordered sets. The main idea is to use the interaction between the combinatorics and the topology of finite topological spaces to investigate homotopy colimits of diagrams of polyhedra. If P is a finite poset and $X: P \rightarrow \mathcal{P}_{<\infty}$ takes values in the category of finite posets, the Grothendieck construction $P \int X$ is also a finite poset and, in fact, it is very simple to characterize. On the other hand, any finite poset can be regarded as a finite topological space where the open subsets

Researchers of CONICET. Partially supported by grants ANPCyT PICT-2011-0812, CONICET PIP 112-201101-00746 and UBACyT 20020130100369.

Received February 20, 2015, revised February 5, 2016; published on October 26, 2016.

2010 Mathematics Subject Classification: 55U10, 55P15, 06A06, 18A30, 18B35.

Key words and phrases: Homotopy colimit, finite topological space, poset, Grothendieck construction, Quillen’s Theorem A.

Article available at <http://dx.doi.org/10.4310/HHA.2016.v18.n2.a13>

Copyright © 2016, International Press. Permission to copy for private use granted.

are its downsets (see, for example, [1, 3, 9, 10, 12]). We handle the Grothendieck construction on a diagram of finite posets as a finite topological space and use a local-to-global theorem of McCord [12, Thm 6] to derive analogues of well known results on homotopy colimits in the combinatorial setting and to prove a generalization of Thomason's theorem. This generalization allows us to apply combinatorial methods to investigate homotopy colimits of diagrams of polyhedra (indexed by finite posets). In [2] Barmak exhibited a very simple proof of Quillen's Theorem A for posets [13] (or equivalently, McCord's theorem for finite topological spaces [12]) using the *non-Hausdorff mapping cylinder* B_f of a poset map $f: X \rightarrow Y$. The non-Hausdorff mapping cylinder is a finite analogue of the classical mapping cylinder of a continuous function, and similarly to its classical version, B_f is the *non-Hausdorff homotopy colimit* (i.e., the Grothendieck construction) of the diagram of posets $X \xrightarrow{f} Y$, indexed by the poset $\mathbf{1}$ of two elements $0 < 1$.

In Section 2 we study non-Hausdorff homotopy colimits of P -diagrams from the finite space point of view. We use *reduction methods* to investigate their weak homotopy types. Quillen's Theorem A for posets follows immediately from Proposition 2.8 and Proposition 2.4 below, by applying the results to the poset $\mathbf{1}$.

The main result of the article is the following generalization of Thomason's theorem in the context of finite posets.

Theorem 1.1. *Let P be a finite poset. Let $K: P \rightarrow \mathcal{S}$ be a diagram of spaces and $X: P \rightarrow \mathcal{P}_{<\infty}$ be a diagram of finite posets. Let $\phi: K \rightarrow X$ be a diagram morphism (where X is viewed as a diagram of finite topological spaces) such that $\phi_p: K_p \rightarrow X_p$ is a weak homotopy equivalence for every $p \in P$. Then there exists a weak homotopy equivalence*

$$\hat{\phi}: \text{hocolim } K \rightarrow \underline{\text{hocolim}} X,$$

from the homotopy colimit of K to the non-Hausdorff homotopy colimit of X (viewed as a finite topological space).

As an immediate consequence of this result we derive a particular case of Thomason's theorem in the context of posets, and also a kind of converse of Thomason's theorem, which relates the homotopy colimit of a diagram of simplicial complexes with the non-Hausdorff homotopy colimit of the diagram of their face posets. In combination with the reduction methods of Section 2, this allows us to simplify the computation of homotopy colimits of diagrams of spaces.

It is well known that for any simplicial set T , there is a *natural* homotopy equivalence $sd T \rightarrow T$ from the (geometric realization of the) barycentric subdivision of T to T (see, for example, [11, Thm. 12.2.5]). By Bousfield–Kan's homotopy lemma [5], this implies that the homotopy colimit of a diagram of simplicial sets is homotopy equivalent to the homotopy colimit of the diagram of their barycentric subdivisions. On the other hand, any ordered simplicial complex K (i.e., a simplicial complex together with a partial ordering of its vertices that restricts to a total ordering on the vertices of each simplex) can be seen as a simplicial set K_s . Moreover, the simplicial set associated to its (geometric) barycentric subdivision $(K')_s$ is naturally isomorphic to $sd K_s$, the subdivision of the simplicial set K_s (see [11, Thm. 12.2.2]). This proves that the homotopy colimit of a diagram of ordered simplicial complexes (and ordered simplicial maps) is homotopy equivalent to the homotopy colimit of the diagram of

their barycentric subdivisions. However, in a general geometric situation, one has to deal with diagrams of unordered simplicial complexes, and in the unordered context there is no natural homotopy equivalence between K and its barycentric subdivision K' (although their underlying topological spaces are equal). Our methods turn out to be appropriate to circumvent this problem. As a corollary of our main theorem, we prove invariance of homotopy type under barycentric subdivision for homotopy colimits in the (general) unordered setting.

The paper concludes with some new results on homotopy colimits of diagrams of polyhedra, which are obtained by applying the main theorem together with the combinatorial methods introduced in Section 2.

2. The Grothendieck construction on posets and reduction methods

Given a finite poset X , we denote by $\mathcal{K}(X)$ its classifying space. $\mathcal{K}(X)$ is also called the order complex of X and it is the simplicial complex whose simplices are the non-empty chains of X . A finite poset can be seen as a finite topological space whose open subsets are the downsets (see, for example, [1, 3, 9, 10, 11, 12]). The topology of the finite space X is related to the topology of the classifying space $\mathcal{K}(X)$. This was studied by McCord in [12]. Concretely, there is a natural weak homotopy equivalence $\mu: \mathcal{K}(X) \rightarrow X$ (i.e., a continuous map which induces isomorphisms in all homotopy groups). μ is called *the McCord map* and it is defined as follows. Given $\alpha \in \mathcal{K}(X)$, write $\alpha = \sum_{i=1}^r t_i x_i$ with $\sum_{i=1}^r t_i = 1$ and $t_i > 0$, where $x_1 < x_2 < \dots < x_r$ is a chain of X , and define $\mu(\alpha) = x_1$.

Throughout this paper we will handle finite posets as finite topological spaces and use the weak equivalence $\mathcal{K}(X) \rightarrow X$. A function $f: X \rightarrow Y$ between finite posets (= finite spaces) is a poset map (i.e., it is order preserving) if and only if it is continuous. Note that a poset X is weakly contractible (= homotopically trivial) if and only if $\mathcal{K}(X)$ is a contractible polyhedron. More generally, two finite posets X and Y are weakly equivalent (denoted by $X \overset{we}{\simeq} Y$) if and only if $\mathcal{K}(X)$ and $\mathcal{K}(Y)$ are homotopy equivalent, and $f: X \rightarrow Y$ is a *weak equivalence* if and only if its realization is a homotopy equivalence. A finite poset is called a *finite model* of a CW-complex K if its classifying space $\mathcal{K}(X)$ is homotopy equivalent to K .

Given a finite simplicial complex K , its face poset will be denoted by $\mathcal{X}(K)$. This is the poset of simplices of K ordered by inclusion. A simplicial map $f: K \rightarrow L$ induces a poset map $\mathcal{X}(f): \mathcal{X}(K) \rightarrow \mathcal{X}(L)$. McCord defined in [12] a weak equivalence $\tilde{\nu}: K \rightarrow \mathcal{X}(K)$ using the McCord map of above and the identification of K with its barycentric subdivision K' . This weak equivalence is however not natural (only up to homotopy). Since we need naturality to apply Theorem 3.1 below, we will use the following variation of the McCord map $\tilde{\nu}$, which was introduced by Barmak in [1]. Given a simplicial complex K , we consider its face poset $\mathcal{X}(K)$ with the opposite order, denoted by $\mathcal{X}(K)^{op}$, and define a map $\nu: K \rightarrow \mathcal{X}(K)^{op}$ by $\nu(x) = \sigma$ if $x \in \overset{\circ}{\sigma}$. Here $\overset{\circ}{\sigma}$ denotes the interior of σ . By [1, Thm. 11.3.2] ν is a natural weak equivalence.

For any $x \in X$, let $U_x \subseteq X$ be the subposet of elements which are smaller than or equal to x and let $\hat{U}_x = U_x \setminus \{x\}$. Analogously, we denote by F_x the subposet of elements of X which are greater than or equal to x and let $\hat{F}_x = F_x \setminus \{x\}$. The

open subsets $\{U_x\}_{x \in X}$ form a basis for the topology of X . When a point x belongs to different posets X, Y , we write $U_x^X, U_x^Y, F_x^X, F_x^Y$ to distinguish whether the subposets are considered in X or in Y . For $x, y \in X$, we write $x \prec y$ if x is covered by y , i.e., if $x < y$ and there is no z such that $x < z < y$. A linear extension of a finite poset X is a total ordering x_1, \dots, x_n of its elements such that if $x_i \leq x_j$ in X then $i \leq j$.

In [1, 3, 4] various reduction methods were introduced. A reduction method allows us to study and handle the homotopy type of a polyhedron by combinatorial moves on their finite models. We describe here some of these methods, which will be used in the rest of this article. For a comprehensive exposition the reader may consult [1, 3, 4]. The first reduction method was introduced by Stong [14] (see also [9, 10]). Given a finite poset X , an element $x \in X$ is called an *up beat point* if \hat{F}_x has a minimum, i.e., there is a unique element $y \in X$ such that $x \prec y$. Analogously x is called a *down beat point* if \hat{U}_x has maximum (there is a unique y such that $y \prec x$). In both cases we say that x is *dominated* by y . If x is a beat point (up or down), $X \setminus \{x\} \subset X$ is a strong deformation retract (as finite topological spaces, and therefore at the level of classifying spaces). A finite poset X is contractible (= dismantlable) if and only if one can remove beat points from X , one by one, to obtain a one-point space (singleton). If X is contractible then it is weakly contractible, but the converse does not hold (see [1, Section 1.3] for more details). The notion of *collapse* in the context of posets was introduced by Barmak and Minian in [3] and it corresponds to Whitehead's classical notion of simplicial collapse. A point $x \in X$ is a *down weak point* if \hat{U}_x is contractible (i.e., dismantlable), and it is an *up weak point* if \hat{F}_x is contractible. An *elementary collapse* is the deletion of an up or down weak point. The inverse operation is called an *elementary expansion*. We say that X *collapses* to Y (or Y expands to X), and denote $X \searrow Y$, if there is a sequence of elementary collapses which starts in X and ends in Y . A poset X is said to be *collapsible* if it collapses to a point. Since any beat point is in particular a weak point, if X is a contractible poset then it is collapsible. In [1, Sections 4.2 and 4.3] and [3, Ex. 3.1] there are various examples of collapsible posets which are not contractible. Finally, we say that X is *simply equivalent* to Y if there exists a sequence of collapses and expansions that starts in X and ends in Y . This is denoted by $X \frown Y$. In [3, Thm. 3.10] it is proved that $X \frown Y$ if and only if $\mathcal{K}(X)$ and $\mathcal{K}(Y)$ are simply equivalent polyhedra (which, as customary, is denoted by $\mathcal{K}(X) \frown \mathcal{K}(Y)$). In particular, if $X \frown Y$ then they are weakly equivalent (and collapsible posets are homotopically trivial). Moreover, if $X \searrow Y$ then $\mathcal{K}(X) \searrow \mathcal{K}(Y)$. It can be shown that if \hat{U}_x or \hat{F}_x is a collapsible poset then $\mathcal{K}(X) \searrow \mathcal{K}(X \setminus \{x\})$ (see [4, Thm. 3.6]).

More generally, we say that a point $x \in X$ is a γ -point if \hat{U}_x or \hat{F}_x is homotopically trivial. It is proved in [4, Thm. 3.15] that in that case, $X \frown X \setminus \{x\}$. Therefore, if a finite poset X can be reduced to a point by removing γ -points, one by one, then it is homotopically trivial (i.e., its classifying space $\mathcal{K}(X)$ is contractible).

Working with diagrams over finite posets allows us to apply reduction methods to study their homotopy colimits. We use reduction methods to derive old and new results on homotopy colimits. We refer the reader to [5] and [16] for the basic definitions and results on homotopy colimits of spaces, and to [17] for applications of homotopy colimits to combinatorial problems.

Let P be a finite poset, viewed as a small category with a unique arrow $p \rightarrow q$ for each $p, q \in P$ such that $p \leq q$, and let X be a P -diagram of finite posets, i.e., a

functor from P to the category $\mathcal{P}_{<\infty}$ of finite posets. In this case the Grothendieck construction on X can be described as follows.

Definition 2.1 (The non-Hausdorff homotopy colimit of finite posets). Let $X: P \rightarrow \mathcal{P}_{<\infty}$ be a functor. The *non-Hausdorff homotopy colimit* of X , denoted by $\underline{\text{hocolim}} X$, is the following poset. The underlying set is the disjoint union $\coprod_{p \in P} X_p$. We keep the given ordering within X_p for all $p \in P$, and for every $x \in X_p$ and $y \in X_q$ such that $p \leq q$, we set $x \leq y$ in $\underline{\text{hocolim}} X$ if $f_{pq}(x) \leq y$ in X_q . Here $X_p = X(p)$ for each $p \in P$ and $f_{pq} = X(p \rightarrow q)$ for each $p \leq q$ in P .

By Thomason’s theorem we have a homotopy equivalence

$$\text{hocolim } \mathcal{K}X \simeq \mathcal{K}(\underline{\text{hocolim}} X).$$

In the context of finite posets, Thomason’s theorem can be deduced from a more general result. This will be proved in the next section.

Remark 2.2. At this point it is worth noting the difference between $\underline{\text{hocolim}} X$ and $\text{hocolim } X$ for a given diagram $X: P \rightarrow \mathcal{P}_{<\infty}$. The first one is the Grothendieck construction on X and it is a finite poset (which can be viewed as a finite topological space). The second one is the classical construction of homotopy colimit of a diagram of topological spaces (applied, in this case, to a diagram of finite topological spaces) and it is not a finite space. However, as an immediate consequence of our main result of next section, we will see that they are weakly equivalent spaces (when we view $\underline{\text{hocolim}} X$ as a finite topological space).

Example 2.3. Any map $f: X_0 \rightarrow X_1$ between finite posets can be viewed as a diagram $X: \mathbf{1} \rightarrow \mathcal{P}_{<\infty}$ where $\mathbf{1}$ is the poset of two elements $0 < 1$. Similarly, as in the topological context, we have $\underline{\text{hocolim}} X = B_f$, the non-Hausdorff mapping cylinder of f , introduced in [1].

It is a well known fact that the mapping cylinder of a continuous function $f: W \rightarrow Z$ is homotopy equivalent to Z . Similarly, if $f: X \rightarrow Y$ is a map of posets, the non-Hausdorff mapping cylinder B_f collapses to Y . This result can be viewed as a particular case of Corollary 2.5 below: if the indexing poset P has maximum p , the homotopy colimit of any P -diagram collapses to X_p . If $f: X \rightarrow Y$ is a poset map such that $f^{-1}(U_y)$ is weakly contractible for every $y \in Y$, then f is a weak equivalence (i.e., it induces a homotopy equivalence between the classifying spaces). This is Quillen’s Theorem A for posets [13] and McCord’s theorem for finite topological spaces [12]. In Proposition 2.8 we generalize this result for homotopy colimits of P -diagrams. Quillen’s Theorem A for posets follows immediately from Propositions 2.8 and 2.4 by applying the results to the poset $\mathbf{1}$.

Given a poset map $\phi: P \rightarrow Q$ and a Q -diagram X , we denote by ϕ^*X the P -diagram obtained by pulling back X along ϕ . Concretely, $\phi^*X = X\phi$. There is a canonical map $\underline{\text{hocolim}} \phi^*X \rightarrow \underline{\text{hocolim}} X$ induced by the identities $(\phi^*X)_p = X_{\phi(p)}$. If $i: Q' \rightarrow Q$ is a subposet and X is a Q -diagram, the restriction i^*X is denoted by $X|_{Q'}$. Note that in this case $\underline{\text{hocolim}} X|_{Q'}$ is a subposet of $\underline{\text{hocolim}} X$.

Proposition 2.4. *Let $X: P \rightarrow \mathcal{P}_{<\infty}$ be a P -diagram of finite posets. If $p \in P$ is an up beat point, then $\underline{\text{hocolim}} X \searrow \underline{\text{hocolim}} X|_{P \setminus \{p\}}$. In particular, they are weakly equivalent.*

Proof. Let x_0, x_1, \dots, x_n be a linear extension of X_p^{op} , the opposite poset of X_p . Let $Y_0 = \underline{\text{hocolim}} X$ and for each $0 \leq i \leq n$ define inductively $Y_{i+1} = Y_i \setminus \{x_i\}$. Let q be the minimum element of \hat{F}_p^P . Note that $\hat{F}_{x_i}^{Y_i} = F_{f_{pq}(x_i)}^{Y_i}$. This implies that x_i is an up beat point of Y_i for all $1 \leq i \leq n$ and therefore

$$\underline{\text{hocolim}} X = Y_0 \searrow Y_1 \searrow \dots \searrow Y_{n+1} = (\underline{\text{hocolim}} X) \setminus X_p = \underline{\text{hocolim}} X|_{P \setminus \{p\}}. \quad \square$$

Corollary 2.5. *Let $X: P \rightarrow \mathcal{P}_{<\infty}$ be P -diagram of finite posets. If P has a maximum element p , then $\underline{\text{hocolim}}(X) \searrow X_p$. In particular, they are weakly equivalent.*

Proof. Let $p = p_0, p_1, \dots, p_n$ be a linear extension of P^{op} . Since P has a maximum element, there is a sequence of collapses

$$P \searrow P \setminus \{p_1\} \searrow P \setminus \{p_1, p_2\} \cdots \searrow P \setminus \{p_1, p_2, \dots, p_n\} = \{p\},$$

where p_i is an up beat point of $P \setminus \{p_1, p_2, \dots, p_{i-1}\}$. By applying recursively Proposition 2.4, we have

$$\underline{\text{hocolim}} X \searrow (\underline{\text{hocolim}} X) \setminus X_{p_1} \searrow \dots \searrow (\underline{\text{hocolim}} X) \setminus \bigcup_{i=1}^n X_{p_i} = X_p. \quad \square$$

From this result and McCord's theorem [12, Thm. 6] we deduce the following analogue of Bousfield–Kan's homotopy lemma (cf. [5, 17]). Note that it can also be deduced from the original homotopy lemma for diagrams of spaces and Thomason's theorem.

Corollary 2.6 (Homotopy lemma). *Let P be a finite poset, let $X, Y: P \rightarrow \mathcal{P}_{<\infty}$ be P -diagrams of finite posets and $\alpha: X \rightarrow Y$ a morphism of diagrams. If $\alpha_p: X_p \rightarrow Y_p$ is a weak equivalence for every $p \in P$, then α induces a weak equivalence*

$$\underline{\text{hocolim}} X \underset{we}{\simeq} \underline{\text{hocolim}} Y.$$

Proof. For any $p \in P$ consider $\underline{\text{hocolim}} X|_{U_p}$ the homotopy colimit of the diagram restricted to U_p . Since p is the maximum of U_p , by Proposition 2.5 $\underline{\text{hocolim}} X|_{U_p} \searrow X_p$ and therefore α induces a weak equivalence $\underline{\text{hocolim}} X|_{U_p} \underset{we}{\simeq} \underline{\text{hocolim}} Y|_{U_p}$. Now the result follows from [12, Thm. 6] applied to the basis-like open cover $\{\underline{\text{hocolim}} Y|_{U_p}\}_{p \in P}$ of $\underline{\text{hocolim}} Y$. \square

Remark 2.7. Note that all the collapses in Proposition 2.4 and Corollary 2.5 are strong collapses, in the sense that all the points removed are beat points (not just weak points). This implies that if the finite space X_p in Corollary 2.5 is contractible (i.e., it is a dismantlable poset) then so is $\underline{\text{hocolim}} X$.

Proposition 2.8. *Let $X: P \rightarrow \mathcal{P}_{<\infty}$ be a P -diagram of finite posets. If p is a down beat point of P dominated by an element q and $f_{qp}^{-1}(U_x)$ is contractible for every $x \in X_p$, then $\underline{\text{hocolim}}(X) \searrow \underline{\text{hocolim}}(X|_{P \setminus \{p\}})$. In particular, they are weakly equivalent.*

Proof. Let x_0, x_1, \dots, x_n be a linear extension of X_p . Define $Y_0 = \underline{\text{hocolim}} X$, and inductively $Y_{i+1} = Y_i \setminus \{x_i\}$ for every $0 \leq i \leq n$. We will show that $Y_0 \searrow Y_1 \searrow \dots \searrow Y_{n+1} = (\underline{\text{hocolim}} X) \setminus X_p$.

Note that $\hat{U}_{x_i}^{Y_i} = \underline{\text{hocolim}} \tilde{X}^i$, where $\tilde{X}^i: \hat{U}_p^P \rightarrow \mathcal{P}_{<\infty}$ is the functor defined by $\tilde{X}^i(p') = f_{p'p}^{-1}(U_{x_i})$, for all $p' < p$ in P (where the transition maps are induced by the original transition maps).

Since \hat{U}_p^P has a maximum element q and $f_{qp}^{-1}(U_{x_i})$ is contractible, by Corollary 2.5 and Remark 2.7, $\hat{U}_{x_i}^{Y_i}$ is a contractible finite space. This proves that x_i is a weak point of Y_i . □

Proposition 2.9. *Let $X: P \rightarrow \mathcal{P}_{<\infty}$ be a P -diagram of finite posets. If p is a down beat point of P dominated by q and f_{qp} is a weak equivalence, then*

$$\underline{\text{hocolim}} X \simeq_{we} \underline{\text{hocolim}} X|_{P \setminus \{p\}}.$$

Proof. Since p is dominated by q , there is a well defined strong homotopy retraction $r: P \rightarrow P \setminus \{p\}$ which is the identity for any $p' \neq p$ and $r(p) = q$. Denote by $i: P \setminus \{p\} \rightarrow P$ the inclusion. There is a morphism of P -diagrams $\gamma: (ir)^*X \rightarrow X$ such that $\gamma_{p'}$ is the identity for every $p' \neq p$ and $\gamma_p = f_{qp}: ((ir)^*X)_p = X_q \rightarrow X_p$. By hypothesis and Corollary 2.6, γ induces a weak equivalence $\underline{\text{hocolim}}(ir)^*X \simeq_{we} \underline{\text{hocolim}} X$, and by Proposition 2.8, $\underline{\text{hocolim}}(ir)^*X \simeq_{we} \underline{\text{hocolim}} X|_{P \setminus \{p\}}$. □

Example 2.10. By the previous proposition one can immediately deduce that the non-Hausdorff homotopy pushout of the poset diagram

$$\begin{array}{ccc} X_0 & \xrightarrow{f_1} & X_1 \\ f_2 \downarrow & & \\ & & X_2 \end{array}$$

is weakly equivalent to X_0 provided the maps f_1 and f_2 are weak equivalences.

We prove now an analogue, in the context of posets, of a cofinality theorem of Bousfield and Kan. We need first a generalization of Proposition 2.4.

Proposition 2.11. *Let $X: P \rightarrow \mathcal{P}_{<\infty}$ be a P -diagram of finite posets. If p is a point of P such that \hat{F}_p is homotopically trivial, then $\underline{\text{hocolim}} X \simeq_{we} \underline{\text{hocolim}} X|_{P \setminus \{p\}}$.*

Proof. Let x_1, x_2, \dots, x_n be a linear extension of X_p^{op} . For each $1 \leq i \leq n$, we define the \hat{F}_p -diagram $\tilde{X}^i: \hat{F}_p \rightarrow \mathcal{P}_{<\infty}$ as follows. $\tilde{X}^i(p') = F_{f_{pp'}(x_i)}$ and $\tilde{X}^i(p' \rightarrow p'') = f_{p'p''|F_{f_{pp'}(x_i)}}: F_{f_{pp'}(x_i)} \rightarrow F_{f_{pp''}(x_i)}$.

Since $\tilde{X}^i(p')$ is contractible for all $p' \in \hat{F}_p$, then by Corollary 2.6 it follows that $\underline{\text{hocolim}} \tilde{X}^i \simeq_{we} \hat{F}_p$, and by hypothesis, the last one is weakly contractible. Therefore, since

$$\hat{F}_{x_i}(\underline{\text{hocolim}} X) \setminus \{x_1, x_2, \dots, x_{i-1}\} = \underline{\text{hocolim}} \tilde{X}^i,$$

which is homotopically trivial, we can remove the points x_1, \dots, x_n of X_p one by one, similarly as in the proof of Proposition 2.4, and the result follows. □

We give now a direct and simple proof of a cofinality theorem for posets. This is a particular case of a known result for functors between small categories which satisfy

the hypotheses of Quillen’s Theorem A (see [8, Prop. 3.2.23] and [6, Thm. 30.5]), and it is an analogue of Bousfield–Kan’s cofinality theorem in the combinatorial setting.

Theorem 2.12 (Cofinality theorem). *Let $\varphi: P \rightarrow Q$ be an order preserving map between finite posets. Let $X: Q \rightarrow \mathcal{P}_{<\infty}$ be a Q -diagram. If $\varphi^{-1}(F_q)$ is homotopically trivial for all $q \in Q$, then the canonical map $\mathop{\mathrm{hocolim}}\varphi^*X \rightarrow \mathop{\mathrm{hocolim}}X$ is a weak equivalence.*

Proof. Let R be the following poset. The underlying set is the disjoint union $Q \coprod P$. We keep the given ordering within Q and P and for every $q \in Q$ and $p \in P$ we set $q \leq p$ if there are $q' \in Q$ and $p' \in P$ such that $q \leq q'$, $p' \leq p$ and $\varphi(p') = q'$. Consider the following R -diagram $\tilde{X}: R \rightarrow \mathcal{P}_{<\infty}$. For each $q \in Q$ take $\tilde{X}(q) = X_q$, set $\tilde{X}(p) = X_{\varphi(p)}$ for $p \in P$, $\tilde{X}(p \rightarrow p') = X(\varphi(p) \rightarrow \varphi(p'))$, $\tilde{X}(q \rightarrow q') = X(q \rightarrow q')$, and $\tilde{X}(\varphi(p) \rightarrow p) = \mathrm{id}_{X_{\varphi(p)}}$. Note that the restriction of \tilde{X} to Q is the original diagram X and the restriction of \tilde{X} to P is φ^*X .

Take a linear extension q_1, q_2, \dots, q_m of Q^{op} . For each $1 \leq j \leq m$ we have

$$\hat{F}_{q_j}^{R \setminus \{q_1, q_2, \dots, q_{j-1}\}} = \varphi^{-1}(F_{q_j}),$$

which is homotopically trivial by hypothesis. By applying recursively Proposition 2.11, we get $\mathop{\mathrm{hocolim}}\tilde{X} \underset{we}{\simeq} \mathop{\mathrm{hocolim}}\varphi^*X$.

Similarly, take a linear extension p_1, p_2, \dots, p_n of P . For each $1 \leq i \leq n$ we have

$$\hat{U}_{p_i}^{R \setminus \{p_1, p_2, \dots, p_{i-1}\}} = \varphi(U_{p_i}) = U_{\varphi(p_i)}.$$

Therefore, p_i is a down beat point of $R \setminus \{p_1, p_2, \dots, p_{i-1}\}$ dominated by $\varphi(p_i)$, and $\tilde{X}(\varphi(p_i) \rightarrow p_i)$ is the identity. By Proposition 2.8 we have $\mathop{\mathrm{hocolim}}\tilde{X} \underset{we}{\simeq} \mathop{\mathrm{hocolim}}X$. □

3. Variations on Thomason’s theorem and applications

In this section we prove a result that relates the homotopy colimit of a diagram of spaces with the non-Hausdorff homotopy colimit of the diagram of their models. As a consequence we obtain an alternative and simple proof of a particular case of Thomason’s theorem in the context of posets. As another immediate consequence of the main result we deduce that the homotopy colimit of a diagram of finite simplicial complexes is weakly equivalent to the non-Hausdorff homotopy colimit of the diagram of their face posets. This implies that all the techniques developed in the previous section for non-Hausdorff homotopy colimits can be used for diagrams of polyhedra (indexed by finite posets) by means of the face poset functor.

We denote by \mathcal{S} the category of topological spaces and continuous maps. Sometimes we require a diagram of spaces $D: P \rightarrow \mathcal{S}$ to satisfy extra conditions (for instance D can be a diagram of simplicial complexes, finite topological spaces, etc.), however in all these cases the homotopy colimit $\mathop{\mathrm{hocolim}}D$ is taken in the category \mathcal{S} .

Theorem 3.1. *Let P be a finite poset. Let $K: P \rightarrow \mathcal{S}$ be a diagram of spaces and $X: P \rightarrow \mathcal{P}_{<\infty}$ be a diagram of finite posets. Let $\phi: K \rightarrow X$ be a diagram morphism (where X is viewed as a diagram of finite topological spaces) such that $\phi_p: K_p \rightarrow X_p$*

is a weak equivalence for every $p \in P$. Then there exists a weak equivalence

$$\hat{\phi}: \text{hocolim } K \rightarrow \underline{\text{hocolim}} X.$$

Proof. We define first the map $\hat{\phi}: \text{hocolim } K \rightarrow \underline{\text{hocolim}} X$. For every $p \leq p'$ denote by $f_{pp'} = K(p \rightarrow p')$ and $g_{pp'} = X(p \rightarrow p')$ the transition maps. Recall that $\text{hocolim } K$ can be constructed from the disjoint union $\coprod_{p \in P} K_p \times \mathcal{K}(F_p)$ by identifying the pairs $(\alpha, \beta) \in K_p \times \mathcal{K}(F_p)$ with $(\alpha', \beta') \in K_{p'} \times \mathcal{K}(F_{p'})$ if $f_{pp'}(\alpha) = \alpha'$ and $\beta = \beta' \in \mathcal{K}(F_{p'})$. We denote by \sim the equivalence relation generated by this identification.

For each $p \in P$, let $\mu_p: \mathcal{K}(F_p) \rightarrow F_p \subseteq X_p$ be the McCord map (defined at the beginning of the previous section). Given $(\alpha, \beta) \in K_p \times \mathcal{K}(F_p)$ we define

$$\hat{\phi}(\alpha, \beta) = \phi_{\mu_p(\beta)}(f_{p\mu_p(\beta)}(\alpha)) \in X_{\mu_p(\beta)} \subseteq \underline{\text{hocolim}} X.$$

It is easy to verify that $\hat{\phi}$ is well defined. In order to see that it is a continuous map, it suffices to prove that for each $y \in \underline{\text{hocolim}} X$, $\hat{\phi}^{-1}(U_y^{\underline{\text{hocolim}} X}) \cap (K_q \times \mathcal{K}(F_q))$ is open in $K_q \times \mathcal{K}(F_q)$ for every $q \in P$. Fix $y \in \underline{\text{hocolim}} X$ and let $p \in P$ such that $y \in X_p$. Note that $\hat{\phi}^{-1}(U_y^{\underline{\text{hocolim}} X}) \cap (K_q \times \mathcal{K}(F_q))$ is empty if $q \not\leq p$ and it is equal to $(g_{qp}\phi_q)^{-1}(U_y^{X_p}) \times \mu_q^{-1}(U_p^{F_q})$ if $q \leq p$. This proves that $\hat{\phi}$ is continuous.

In order to prove that $\hat{\phi}$ is a weak homotopy equivalence, we use McCord's theorem [12, Thm. 6] for the basis-like open cover $\{\underline{\text{hocolim}} X|_{U_p}\}_{p \in P}$ of $\underline{\text{hocolim}} X$. We have to see that

$$\hat{\phi}: \hat{\phi}^{-1}(\underline{\text{hocolim}} X|_{U_p}) \rightarrow \underline{\text{hocolim}} X|_{U_p}$$

is a weak equivalence for each p . Note that

$$\hat{\phi}^{-1}(\underline{\text{hocolim}} X|_{U_p}) = \coprod_{q \leq p} K_q \times \mu_q^{-1}(U_p^{F_q})/\sim$$

and that there is a commutative diagram

$$\begin{array}{ccc} K_p \times \{p\} & \xrightarrow{\phi_p} & X_p \\ \downarrow i & & \downarrow j \\ (\coprod_{q \leq p} K_q \times \mu_q^{-1}(U_p^{F_q})/\sim) & \xrightarrow{\phi} & \underline{\text{hocolim}} X|_{U_p}. \end{array}$$

The inclusion j is a weak equivalence by Corollary 2.5 and ϕ_p is a weak equivalence by hypothesis, thus we only need to check that the inclusion i is a homotopy equivalence.

Consider the retraction $r: (\coprod_{q \leq p} K_q \times \mu_q^{-1}(U_p^{F_q})/\sim) \rightarrow K_p \times \{p\}$ defined by $r(\alpha, \beta) = (f_{qp}(\alpha), p)$ for $(\alpha, \beta) \in K_q \times \mu_q^{-1}(U_p^{F_q})$. It is clear that $ri = 1$. We define a homotopy $H: ir \simeq 1$ as a composition of two linear homotopies. Any $\beta \in \mu_q^{-1}(U_p^{F_q})$ can be written as $\beta = t\beta_1 + (1-t)\beta_2$ with $0 < t \leq 1$, $\beta_1 \in \mathcal{K}(U_p)$ and $\beta_2 \in \mathcal{K}(X_p \setminus U_p)$. Take $H_1((\alpha, \beta), s) = (\alpha, (1-s)\beta + s\beta_1)$. Since $\beta_1 \in \mathcal{K}(U_p)$ which is a cone with apex p , we can define then $H_2((\alpha, \beta), s) = (\alpha, (1-s)\beta_1 + sp)$. \square

Remark 3.2. Under the hypotheses of Theorem 3.1, there is an alternative way to prove that $\text{hocolim } K$ and $\underline{\text{hocolim}} X$ are weakly equivalent by means of a zigzag of weak equivalences. First one can see that it suffices to reduce to the case $K = X$ and $\phi = 1_X$. Then, in order to prove that $\text{hocolim } X$ and $\underline{\text{hocolim}} X$ are weakly

equivalent one can use Thomason's theorem (to obtain a weak equivalence at the level of simplicial sets) and McCord's theorem (to come back to the context of topological spaces).¹ Theorem 3.1 exhibits an explicit and direct weak equivalence $\text{hocolim } K \rightarrow \underline{\text{hocolim}} X$.

As a first immediate corollary we obtain the following particular case of Thomason's theorem for posets.

Corollary 3.3. *Given a diagram of finite posets $X: P \rightarrow \mathcal{P}_{<\infty}$, there is a homotopy equivalence*

$$\text{hocolim } \mathcal{K}X \rightarrow \mathcal{K}(\underline{\text{hocolim}} X).$$

Proof. We apply Theorem 3.1 to the diagram morphism $\mu: \mathcal{K}X \rightarrow X$, where $\mu_p: \mathcal{K}(X_p) \rightarrow X_p$ is the McCord map. \square

Now we prove a kind of converse of Thomason's result, which relates the homotopy colimit of a diagram of simplicial complexes with the non-Hausdorff homotopy colimit of their face posets.

Corollary 3.4. *Let $K: P \rightarrow \mathcal{S}$ be a diagram of finite simplicial complexes. Then there is a weak equivalence*

$$\nu: \text{hocolim } K \rightarrow \underline{\text{hocolim}}(\mathcal{X}K)^{op},$$

from the homotopy colimit of K to the non-Hausdorff homotopy colimit of the diagram of the opposite of their face posets.

Proof. Apply Theorem 3.1 to the diagram morphism induced by the natural weak equivalences $\nu_p: K_p \rightarrow \mathcal{X}(K_p)^{op}$ defined in [1, Thm. 11.3.2] and the beginning of the previous section. \square

Since for every simplicial complex L , $\mathcal{K}(\mathcal{X}(L)^{op}) = \mathcal{K}(\mathcal{X}(L)) = L'$ (the barycentric subdivision of L), from Corollary 3.3 and Corollary 3.4 we deduce the following.

Corollary 3.5. *Let $K: P \rightarrow \mathcal{S}$ be a diagram of (unordered) finite simplicial complexes (and simplicial maps). Then $\text{hocolim } K$ and $\text{hocolim } K'$ are homotopy equivalent, where $K': P \rightarrow \mathcal{S}$ is the diagram of the barycentric subdivisions (of spaces and maps).*

Note that the homotopy equivalence of the last corollary cannot be deduced directly from a diagram map between K and K' since, although the underlying topological spaces of K_p and K'_p are equal and the transition maps f_{qp} and f'_{qp} are (linearly) homotopic, in the context of unordered simplicial complexes there is no *natural* homotopy equivalence from the barycentric subdivision functor to the identity functor. Moreover, in general the homotopies $f_{qp} \simeq f'_{qp}$ cannot be taken coherently.

Remark 3.6. From Theorem 3.1 one can easily deduce that if $X: P \rightarrow \mathcal{S}$ is a diagram of finite topological spaces, although $\text{hocolim } X$ and $\underline{\text{hocolim}} X$ are very different, there is a weak homotopy equivalence $\text{hocolim } X \rightarrow \underline{\text{hocolim}} X$ induced by the identity map.

¹We thank the referee for pointing this out to us.

Corollary 3.4 allows one to apply the results of Section 2 to homotopy colimits of diagrams of polyhedra, by means of the weak equivalence with the non-Hausdorff homotopy colimits of the opposite of the face posets. In particular, the following analogues of Propositions 2.11 and 2.9 are valid for diagrams of simplicial complexes.

Proposition 3.7. *Let $K: P \rightarrow \mathcal{S}$ be a P -diagram of finite simplicial complexes (and simplicial maps). If p is a point of P such that \hat{F}_p is homotopically trivial (in particular, if p is an up beat point or an up weak point), then $\text{hocolim } K \simeq \text{hocolim } K|_{P \setminus \{p\}}$.*

Proof. Consider the diagram $(\mathcal{X}K)^{op}: P \rightarrow \mathcal{P}_{<\infty}$, $(\mathcal{X}K)^{op}(p) = (\mathcal{X}(K_p))^{op}$, and apply Proposition 2.11 and Corollary 3.4. \square

Proposition 3.8. *Let $K: P \rightarrow \mathcal{S}$ be a P -diagram of finite simplicial complexes (and simplicial maps). If p is a down beat point of P dominated by q and f_{qp} is a homotopy equivalence, then*

$$\text{hocolim } K \simeq \text{hocolim } K|_{P \setminus \{p\}}.$$

Proof. Consider the diagram $(\mathcal{X}K)^{op}: P \rightarrow \mathcal{P}_{<\infty}$ and apply Proposition 2.9 and Corollary 3.4. \square

Suppose that $K: P \rightarrow \mathcal{S}$ is a diagram of finite simplicial complexes such that all the transition maps $f_{qp}: K_q \rightarrow K_p$ are homotopy equivalences (in particular, if P is connected, all K_p have the same homotopy type). In general, although the simplicial maps $f_{qp}: K_q \rightarrow K_p$ are homotopy equivalences, the homotopy type of K_p and the topology of $\mathcal{K}(P)$ do not determine the homotopy type of $\text{hocolim } K$. We will show that if the indexing poset P and the maps f_{qp} satisfy nice conditions, then $\text{hocolim } K \simeq K_p$ (for any $p \in P$).

Corollary 3.9. *Let $K: P \rightarrow \mathcal{S}$ be a P -diagram of finite simplicial complexes (and simplicial maps). If P is a contractible finite space (i.e., a dismantlable poset) and the transition maps f_{qp} are homotopy equivalences, then $\text{hocolim } K \simeq K_p$ (for any $p \in P$).*

Proof. Since P is a contractible finite space, there is a sequence p_1, \dots, p_n such that p_i is a beat point (up or down) of $P \setminus \{p_1, p_2, \dots, p_{i-1}\}$ and $P \setminus \{p_1, p_2, \dots, p_n\} = \{p\}$. Now apply recursively Propositions 3.7 and 3.8. \square

As we state at the beginning of the previous section, if P is contractible (as a finite space) then its classifying space $\mathcal{K}(P)$ is contractible but the converse does not hold. In [1, 3, 4] there are various examples of non-contractible finite spaces P with $\mathcal{K}(P)$ contractible. Suppose that the indexing poset P can be reduced to a single point by removing γ -points (recall that p is a γ -point if \hat{F}_p or \hat{U}_p is homotopically trivial). In that case the previous corollary is not longer valid since Proposition 3.8 works only for down beat points (i.e., when \hat{U}_p is a contractible finite space, not just homotopically trivial). However, one can impose extra conditions on the maps $f_{qp}: K_q \rightarrow K_p$ in order to extend Proposition 3.8 to γ -points, and Corollary 3.9 to a more general class of homotopically trivial posets. To this end we replace homotopy equivalences by *contractible mappings*. This class of maps was introduced by Cohen in [7]. A simplicial map $f: K \rightarrow L$ is called a contractible mapping if the preimage $f^{-1}(z)$ is contractible for every point z in the underlying space of L . In [7, Thm. 11.1] Cohen

proved that any contractible mapping $f: K \rightarrow L$ is a simple homotopy equivalence. In [2, Thm. 5.1] Barmak exhibited an alternative and simple proof of Cohen’s result. From the proof of [2, Thm. 5.1] one can deduce the following.

Proposition 3.10 (Barmak). *Let $f: K \rightarrow L$ be a contractible mapping and let $\mathcal{X}(f)^{op}: \mathcal{X}(K)^{op} \rightarrow \mathcal{X}(L)^{op}$ be the map induced in the opposite of their face posets. Then $(\mathcal{X}(f)^{op})^{-1}(U_\sigma)$ is homotopically trivial for every $\sigma \in \mathcal{X}(L)^{op}$.*


Corollary 3.11. *Let $K: P \rightarrow \mathcal{S}$ be a P -diagram of finite simplicial complexes (and simplicial maps). If p is a point of P such that \hat{U}_p is homotopically trivial and the transition maps f_{qp} are contractible mappings for every $q \leq p$, then $\text{hocolim } K \simeq \text{hocolim } K|_{P \setminus \{p\}}$.*

Proof. Consider the diagram $(\mathcal{X}K)^{op}: P \rightarrow \mathcal{P}_{<\infty}$ and follow the proof of Proposition 2.8, using that $(\mathcal{X}(f_{qp})^{op})^{-1}(U_\sigma)$ are homotopically trivial by Proposition 3.10. □

Corollary 3.11 in combination with Proposition 3.7 allows us to extend Corollary 3.9 to a more general class of (homotopically trivial) indexing posets, under the stronger assumption that the transition maps are contractible mappings.

Corollary 3.12. *Let $K: P \rightarrow \mathcal{S}$ be a P -diagram of finite simplicial complexes. If the indexing poset P can be reduced to a point by removing γ -points (in particular, if P is collapsible), and the transition maps f_{qp} are contractible mappings, then $\text{hocolim } K \simeq K_p$ (for any $p \in P$).*

Example 3.13. If the transition maps of the following diagram of simplicial complexes are contractible mappings, its homotopy colimit is homotopy equivalent to any of the K_p . This is because the indexing poset is a collapsible (but non-contractible) finite space.



Acknowledgments

We are grateful to Jonathan Barmak for many useful discussions and suggestions during the preparation of this article. We also would like to thank Volkmar Welker and Tim Porter for useful comments.

References

[1] Barmak, J.A. *Algebraic Topology of Finite Topological Spaces and Applications*. Lecture Notes in Math., Vol. 2032 (2011).

- [2] Barmak, J.A. *On Quillen's Theorem A for posets*. J. Combin. Theory Ser. A **118** (2011), 2445–2453.
- [3] Barmak, J.A.; Minian, E.G. *Simple homotopy types and finite spaces*. Adv. Math. **218** (2008), no. 1, 87–104.
- [4] Barmak, J.A.; Minian, E.G. *One-point reductions of finite spaces, h -regular CW-complexes and collapsibility*. Algebr. Geom. Topol. **8** (2008), no. 3, 1763–1780.
- [5] Bousfield, A. K.; Kan, D. M. *Homotopy Limits, Completions and Localizations*, Lecture Notes in Math., Vol. 304 (1972).
- [6] Chachólski, W.; Scherer, J. *Homotopy Theory of Diagrams*. Mem. Amer. Math. Soc., Vol. 155 (2002), no. 736, x+90 pp.
- [7] Cohen, M.M. *Simplicial structures and transverse cellularity*. Ann. of Math. **85** (1967), 218–245.
- [8] Maltiniotis, G. *La théorie de l'homotopie de Grothendieck*. Astérisque, Vol. 301 (2005), vi+140 pp.
- [9] May, J.P. *Finite topological spaces*. Notes for REU, Chicago (2003). Available at <http://www.math.uchicago.edu/~may/MISCMaster.html>.
- [10] May, J.P. *Finite spaces and simplicial complexes*. Notes for REU, Chicago (2003). Available at <http://www.math.uchicago.edu/~may/MISCMaster.html>.
- [11] May, J.P. *Finite Spaces and Larger Contexts*. Book in progress, Chicago (2012). Available at <http://math.uchicago.edu/~may/REU2012/>
- [12] McCord, M.C. *Singular homology groups and homotopy groups of finite topological spaces*. Duke Math. J. **33** (1966), 465–474.
- [13] Quillen, D. *Higher algebraic K-theory, I: Higher K-theories*. Lecture Notes in Math., Vol. 341 (1972), 85–147.
- [14] Stong, R.E. *Finite topological spaces*. Trans. Amer. Math. Soc. **123** (1966), 325–340.
- [15] Thomason, R. W. *Homotopy colimits in the category of small categories*. Math. Proc. Cambridge Philos. Soc. **85** (1979), no. 1, 91–109.
- [16] Vogt, R.M. *Homotopy limits and colimits*. Math. Z. **134** (1973), 11–52.
- [17] Welker, V.; Ziegler, G.; Živaljević, R. *Homotopy colimits – comparison lemmas for combinatorial applications*. J. Reine Angew. Math. **509** (1999), 117–149.

Ximena Fernández xfernand@dm.uba.ar

Elías Gabriel Minian gminian@dm.uba.ar

Departamento de Matemática – IMAS, FCEyN, Universidad de Buenos Aires, Buenos Aires, Argentina