

---

# Singularities and Diffeomorphisms

Dedicated to Blaine Lawson with admiration

by Tobias Holck Colding<sup>\*†</sup> and  
William P. Minicozzi II<sup>\*†</sup>

**Abstract.** Comparing and recognizing metrics can be extraordinarily difficult because of the group of diffeomorphisms. Two metrics, that could even be the same, could look completely different in different coordinates. This is the gauge problem. The general gauge problem is extremely subtle for non-compact spaces. Often it can be avoided if one uses some additional structure of the particular situation. However, in many problems there is no additional structure. Instead we solve the gauge problem directly in great generality.

The techniques and ideas apply to many problems. We use them to solve a well-known open problem in Ricci flow.

We solve the gauge problem by solving a nonlinear system of PDEs. The PDE produces a diffeomorphism that fixes an appropriate gauge in the spirit of the slice theorem for group actions. We then show optimal bounds for the displacement function of the diffeomorphism.

## 0. Introduction

Suppose we have two weighted manifolds  $(M_i, g_i, f_i)$  for  $i = 1, 2$  satisfying some PDE. Assume that on a large, but compact set, the manifolds  $M_i$ , metrics  $g_i$  and weights  $e^{-f_i}$  almost agree after identification by a diffeomorphism.

---

<sup>\*</sup> MIT, Dept. of Math., 77 Massachusetts Avenue, Cambridge, MA 02139-4307.

E-mail: [colding@math.mit.edu](mailto:colding@math.mit.edu)

E-mail: [minicozz@math.mit.edu](mailto:minicozz@math.mit.edu)

<sup>†</sup> The authors were partially supported by NSF DMS Grants 2104349 and 2005345.

- Are the manifolds, metrics and weights the same everywhere after some identification?

This is a common problem in many questions. The major obstacle for understanding this in general is the infinite dimensional gauge group of diffeomorphisms:

- Two metrics, that could even be the same, could look very different in different coordinates.

In some situations the gauge problem can be avoided if there is some additional structure. A classical example is the Killing-Hopf theorem that classifies constant curvature metrics. This classification uses that the curvature tensor is constant to construct a “canonical” isometry between the two spaces. In general, the gauge problem can be solved when there is strong asymptotic decay and circumvented when the space is characterized in a coordinate-free way, such as a large symmetry group, the vanishing of a special tensor, or a strong curvature condition.

In the problems we will be interested in, the manifold will be non-compact and we will not have any special structure. Thus, we will be forced to deal with the gauge problem head on. The flip side of this is that once we do that it gives new tools with broad applications.

### 0.1 Where Do Questions Like These Arise?

Problems about identifying spaces occur in many different situations. The one we are interested in here comes from Ricci flow. A one parameter family  $(M, g(t))$  of manifolds flows by the Ricci flow, [H], if

$$g_t = -2\text{Ric}_{g(t)},$$

where  $\text{Ric}_{g(t)}$  is the Ricci curvature of the evolving metric  $g(t)$  and  $g_t$  is the time derivative of the metric.

The key to understand Ricci flow is the singularities that form. The simplest singularity is a homothetically shrinking sphere that disappears (becomes extinct) at a point. The product of a sphere with  $\mathbf{R}$  gives a shrinking cylinder; this singularity, called a neck pinch, is much more complicated than the spherical extinction. In dimension three, spherical extinctions and neck pinches are essentially the only singularities. Adding another  $\mathbf{R}$  factor, gives the so-called bubble sheet singularity that is only recently partially understood. With each additional  $\mathbf{R}$  factor, the singularities become more complicated and the sets where they occur are larger.

A triple  $(M, g, f)$  of a manifold  $M$ , metric  $g$  and function  $f$  on  $M$  is a gradient shrinking Ricci soliton (or *shrinker*) if it satisfies

$$\text{Ric} + \text{Hess}_f = \frac{1}{2}g.$$

Shrinkers give special solutions of the Ricci flow that evolve by rescaling up to diffeomorphism and are singularity models. They arise as time-slices of limits of rescalings (magnifications) of the flow around a fixed future singular point in space-time. Such limits are said to be tangent flows at the singularity. Even when  $M$  is compact, the shrinker is typically non-compact and the convergence is on compact subsets. Shrinkers also arise in other important ways, such as blow-downs from  $-\infty$  for ancient flows with bounded entropy. Ancient flows are flows that have existed for all prior times; every tangent flow is ancient. Shrinkers are the key singularities in Ricci flow and will be our focus here.

Among shrinkers, cylinders are particularly important; they are the most prevalent. This is because the Almgren-Federer-White dimension reduction divides the singular set into strata whose dimension is the dimension of translation-invariance of the blowup. For Ricci flow, this suggests:

- Top strata of the singular set corresponds to points where the blowup is  $\mathbf{R}^{n-2} \times N^2$ .
- The next strata consists of points where the blowup is  $\mathbf{R}^{n-3} \times N^3$ .

The  $N$ 's are themselves shrinkers and have been classified in low dimensions by Cao-Chen-Zhu, Hamilton, Ivey, Naber, Ni-Wallach, Perelman. In dimensions two and three, they are  $N^2 = \mathbf{S}^2$  or  $\mathbf{RP}^2$  and  $N^3 = \mathbf{S}^3$  or  $\mathbf{S}^2 \times \mathbf{R}$  (plus quotients). The classification in dimension three relies on an equation for the 2-tensor  $\frac{\text{Ric}}{S}$  that fails in higher dimensions where there is no similar classification. In fact, there are huge families of shrinkers in higher dimensions. Combining dimension reduction

with the classification in low dimensions, we see that the most prevalent singularities are:

$$\mathbf{S}^2 \times \mathbf{R}^{n-2} \text{ followed by } \mathbf{S}^3 \times \mathbf{R}^{n-3} \text{ (and quotients).}$$

As one approaches a singularity in the flow and magnifies, one would like to know which singularity it is. Since most singularities are non-compact yet the evolving manifolds are closed, one only sees a compact piece of the singularity at each time as one approaches it. The next theorem recognizes the most prevalent singularities from just a compact piece.

**Theorem 0.1** ([CM1]). Cylindrical shrinkers  $\mathbf{S}^\ell \times \mathbf{R}^{n-\ell}$  are strongly rigid for any  $\ell$ .

The theorem holds for products of  $\mathbf{R}^{n-\ell}$  with quotients of  $\mathbf{S}^\ell$  and a large class of other positive Einstein manifolds; see [CM1] for details. *Strong rigidity means* that if another shrinker is close enough on a large compact set, then it must agree.

In most problems in geometric PDEs, it would be unthinkable to control an entire solution from just knowing roughly how it looks on a compact set. This is exactly what we do here. If one knew exactly how it looked like on a compact set, it would be much less surprising and essentially follow from unique continuation. The surprising thing here is that we only assume closeness and only on a compact set and this is enough to characterize the shrinker. This is an illustration of a *shrinker principle* which roughly says that “uniqueness radiates outwards”. Nothing like this is true for Einstein manifolds (or steady solitons), where gravitational instantons contain arbitrarily large arbitrarily Euclidean regions. The shrinker principle was originally discovered in mean curvature flow [CIM], [CM2]. It has been conjectured since that something similar holds for Ricci flow, but the gauge group has been one of the major obstacles. Using extrinsic coordinates, the gauge is circumvented in mean curvature flow.

Tangent flows are limits of a subsequence of rescalings at the singularity. A priori different subsequences might give different limits. Using Theorem 0.1, we get the following uniqueness:

**Theorem 0.2** ([CM1]). For a Ricci flow, if one tangent flow at a point in space-time is a cylinder, then all other tangent flows at that point are also cylinders.

Unlike most results in Ricci flow, these results hold for every  $n$  and  $\ell$ . Increasing the dimension of the Euclidean factor is a subtle problem (e.g. surgery, cylindrical estimates, and  $k$ -convexity estimates only allow small Euclidean factors). For general  $n$  and  $\ell$ , cylinders do not have a coordinate-free characterization. This is a major part of the difficulty, forcing us to address the gauge problem head on.

At singularities where the tangent flows are compact shrinkers the singularities are isolated in space-time. For compact shrinkers, rigidity was proven in dimension three by Hamilton in 1982 and by Huisken in 1985 for higher dimensional spheres.

Rigidity for necks  $S^{n-1} \times \mathbf{R}$  was proven independently by Li-Wang [LW]. They are able to circumvent the gauge problem by using that their Euclidean factor is a line. They do that, in part, by using tensors with special properties on the product of a sphere with a line to prove asymptotic structure and approximate symmetry. Once they have this, they are able to use again that their Euclidean factor is a line to apply Brendle's symmetry improvement to get  $O(n)$  symmetry and, finally, Kotschwar's classification of rotationally symmetric shrinkers.

## 0.2 Further Applications

Rigidity and uniqueness of blowups are fundamental questions in regularity theory that have many applications. For instance in mean curvature flow, they play a major role in understanding the singular set, proving optimal regularity, understanding solitons, classifying ancient solutions, and understanding low entropy flows.

# 1. What Is Needed for Rigidity?

We need to show that if two shrinkers are close on a large but compact set, then they agree identically everywhere. This will be done by iterating two estimates: extension and improvement. Extension shows that the shrinkers remain close even on a fixed larger scale, but with a loss in the estimates. Improvement recovers this loss and shows that they are even closer on the larger scale. Once we have this, we can iterate the argument to get estimates on larger and larger scales, eventually giving the strong rigidity. Estimates proving polynomial losses will be played off against estimates with exponential gains.

Four key ingredients in the proof of strong rigidity are:

1. Gauge fixing.
2. New polynomial growth estimates for PDEs.
3. Propagation of almost splitting.
4. Quadratic rigidity in the right gauge.

We will use new polynomial growth estimates as ingredients in both (1) and (3).

## 1.1 Gauge Fixing

Fix  $(M, g, f)$ . We are given a diffeomorphism from a large compact set in  $M$  to a second weighted space.

- The pull-back metric and weight are  $g + h$  and  $e^{-f-k}$ .
- $h$  and  $k$  are small on the compact set.

Composing with a diffeomorphism on  $M$  gives a different  $h$  and  $k$ . We want to mod out by this group action. This is gauge fixing.

One of the most interesting results of transformation groups is the existence of slices. A slice for the action of a group on a manifold is a submanifold which is transverse to the orbits near a given point. Ebin and Palais proved the existence of a slice for the diffeomorphism group of a compact manifold acting on the space of all Riemannian metrics. The slice can be thought of as the gauge fixing on the compact manifold. However, manifolds are not compact here.

In our setting, gauge fixing is choosing a diffeomorphism  $\Phi$  on  $M$  so the new  $h$  is orthogonal to the group action. This is a nonlinear PDE for  $\Phi$ . Orthogonality corresponds to making  $\operatorname{div}_f h = 0$ , where  $\operatorname{div}_f(h) = e^f \operatorname{div}(e^{-f} h) = \operatorname{div}(h) - h(\nabla f, \cdot)$ . The solution to the nonlinear PDE asks to find a diffeomorphism  $\Phi$  so that  $\tilde{h} = \Phi^*(g + h) - g$  satisfies

$$(1.1) \quad \operatorname{div}_f \tilde{h} = 0.$$

Terms involving  $\operatorname{div}_f$  come up again and again, so many quantities simplify in this gauge and having them drop out as they do when  $\operatorname{div}_f \tilde{h} = 0$  makes things possible to analyze.

We construct the diffeomorphism  $\Phi$  that solves this nonlinear PDE using an iteration scheme for the linearized operator  $\mathcal{P}$  on vector fields  $Y$ . Using optimal polynomial bounds on  $\mathcal{P}$ , we show sharp polynomial bounds for the displacement function of  $\Phi$

$$x \rightarrow \operatorname{dist}_g(x, \Phi(x)).$$

For applications, it is crucial that we only assume closeness on a compact set and, in particular, a priori the two shrinkers do not need to be diffeomorphic. This means that we cannot fix the gauge at the outset. Instead we need to apply our gauge fixing procedure iteratively to fix the gauge on larger and larger scales as we move outward and show closeness on larger and larger scales. To pull this off requires very strong estimates for the displacement which is what we show. Our optimal estimates show that the displacement of the gauge fixing diffeomorphism grows at a sharp polynomial rate. The bound is relative so closeness on the initial scale implies closeness at a larger scale.

On a shrinker  $(M, g, f)$  there is a natural gaussian  $L^2$  norm given by  $\|u\|_{L^2}^2 = \int_M u^2 e^{-f}$ . Diffeomorphisms near the identity are infinitesimally generated by integrating a vector field  $X$ . The infinitesimal change of the metric is given by the Lie derivative of the metric with respect to  $X$ . This is equal to  $-\frac{1}{2} \operatorname{div}_f^* X$ , where

$\operatorname{div}_f^*$  is the operator adjoint of  $\operatorname{div}_f$  with respect to the gaussian inner product. Thus, if we define the operator  $\mathcal{P}$  by

$$\mathcal{P}X = \operatorname{div}_f \circ \operatorname{div}_f^* X,$$

then the linearization of (1.1) is to find a vector field  $X$  with

$$\mathcal{P}X = \frac{1}{2} \operatorname{div}_f h.$$

A detailed analysis of  $\mathcal{P}$  and its properties play an important role in the gauge fixing.

The canonical second order elliptic “drift Laplacian” that is self-adjoint and positive definite with respect to the gaussian inner product is a generalized Ornstein-Uhlenbeck operator. On tensors  $u$  this operator is given by  $\mathcal{L}u = \Delta u - \nabla_{\nabla_f} u$ . Given a vector field  $X$  on a shrinker, the two operators  $\mathcal{L}$  and  $\mathcal{P}$  are related by the identity

$$-2\mathcal{P}X = \nabla \operatorname{div}_f X + \mathcal{L}X + \frac{1}{2}X.$$

The unweighted version of  $\mathcal{P}$  was used implicitly by Bochner to show that closed manifolds with negative Ricci curvature have no Killing fields. The unweighted operator was later used by Bochner and Yano, to show that the isometry group of such manifolds is finite. The unweighted operator also arises in general relativity and fluid dynamics. The weighted operator  $\mathcal{P}$  appears to have been largely overlooked. The relationship between  $\mathcal{P}$  and the unweighted version, used by Bochner, mirrors the relationship between the Ornstein-Uhlenbeck operator and the Laplacian.

## 1.2 New Polynomial Growth Estimates for PDEs

Surprisingly, in very general settings, we show the same polynomial growth bounds for  $\mathcal{L}$  that Laplace and Hermite observed on  $\mathbf{R}^n$  for the standard gaussian. Namely, suppose  $f$  is a proper  $C^n$  function and  $S \geq 0$  satisfies

$$(1.2) \quad \Delta f + S = \frac{n}{2},$$

$$(1.3) \quad |\nabla f|^2 + S = f.$$

**Theorem 1.4** ([CM1]). If (1.2) and (1.3) hold,  $u \in L^2(e^{-f})$  and  $\mathcal{L}u = -\lambda u$ , then  $u$  grows polynomially of degree at most  $2\lambda$ .

The conditions (1.2) and (1.3) hold for all shrinkers in both Ricci and mean curvature flow, giving a powerful new tool for a wide array of problems. These estimates are used in both the gauge fixing and the propagation of almost splitting. We use the relationship between  $\mathcal{L}$  and  $\mathcal{P}$  to translate these optimal bounds for  $\mathcal{L}$  into optimal bounds for  $\mathcal{P}$ .

## 1.3 Propagation of Almost Splitting

One of the important new ingredients is that a Ricci shrinker close to a product  $N \times \mathbf{R}^{n-\ell}$  on a large scale remains close on a fixed larger scale. The idea is that the initial closeness will imply that  $\mathcal{L}$  has eigenvalues that are exponentially close to  $\frac{1}{2}$ . The drift Bochner formula on a shrinker implies that every eigenvalue is at least  $\frac{1}{2}$  with equality only when it splits. We show that being close to  $\frac{1}{2}$  gives that the hessian is almost zero in  $L^2$ , which is very strong on the region where the weight  $f$  is small but says almost nothing further out. The crucial point is that the hessian can grow only polynomially, so the very small initial bound gives bounds much further out. Thus, the gradients of these eigenfunctions give the desired almost parallel vector fields and almost splitting.

If a shrinker is exponentially close to a cylinder on scale  $R$ , then almost splitting gives on scale  $(1+\epsilon)R$ :

- (A)  $n - \ell$  almost translations and a metric almost splitting.
- (B) The slices  $\{f = c\}$  are almost spherical.
- (C)  $f$  also almost splits  $f = f_0 + \frac{|x|^2}{4}$ .

Combining (A) and (B) gives a diffeomorphism on scale  $(1+\epsilon)R$  from the cylinder to the second shrinker and this diffeomorphism is close to an isometry. The condition (C) guarantees that the diffeomorphism also almost preserves  $f$ . As a result, we see that the shrinker looks cylindrical even on the larger scale. However, there is a loss in the estimates – it may look less cylindrical on the larger scale – that makes this impossible to iterate on its own.

## 1.4 Quadratic Rigidity

The propagation of almost splitting and gauge fixing give that the shrinker is close to a cylinder on a large set via a diffeomorphism that fixes the gauge. The last of the four key ingredients is an estimate for the difference in metrics that is small enough to be iterated. For this, it is essential that the gauge be right, or else it just isn’t true. The closeness cannot be seen via linear analysis. However, we show that there is a second order rigidity that gives the estimate; we call this quadratic rigidity.

To explain the estimate, let  $(M, g, f)$  be the cylinder and  $(M, g+h, f+k)$  the shrinker that is close on a large compact set. We need bounds on  $h$  and  $k$  that can be iterated. The linearization of the shrinker equation is

$$(1.5) \quad \frac{1}{2}Lh + \operatorname{Hess}_{\frac{1}{2}\operatorname{Tr}(h)-k} + \operatorname{div}_f^* \operatorname{div}_f h.$$

This linearization was derived by Cao-Hamilton-Ilmanen in their calculation of the second variation

operator for Perelman’s entropy. The operator  $L$  acts on 2-tensors by

$$Lh = \mathcal{L}h + 2R(h),$$

and  $R(h)$  is the natural action of the Riemann tensor.

Since  $(M, g + h, f + k)$  is also a shrinker, (1.5) must be at least quadratic in  $(h, k)$ . The last two terms in (1.5) are gauge terms - i.e., in the image of  $\operatorname{div}_f^*$  and there is no reason for these - or  $h$  - to be small if not in right gauge. In the *right gauge*, the difference  $h$  in the metrics satisfies the Jacobi equation  $Lh = 0$  up to higher order terms. This does not force  $h$  to be small since cylinders have non-trivial Jacobi fields that could potentially integrate to give nearby shrinkers. The second variation of the **shrinker equation** in the direction of a Jacobi field is given by the tensor

$$(1.6) \quad -2|\nabla u|^2 \operatorname{Ric} - 2Su \operatorname{Hess}_u - S\nabla u \otimes \nabla u.$$

Here  $S$  is scalar curvature and  $u$  is a quadratic Hermite polynomial that measures the projection of  $h$  onto Jacobi fields. The first order Taylor expansion will give that  $h$  is a Jacobi field to first order and, thus,  $|h|$  is

$|u|$  up to higher order. On the other hand, the second order Taylor expansion will imply that (1.6) vanishes to at least third order in  $h$ . Combining these, we see that the quadratic expression (1.6) is in fact at least cubic in  $u$ . When  $u$  is small, this implies that  $u$  and  $h$  vanish; we will have extra error terms so will get that  $h$  is exponentially small, giving the improvement that we needed to iterate.

## References

- [CIM] T.H. Colding, T. Ilmanen and W.P. Minicozzi II, *Rigidity of generic singularities of mean curvature flow*. Publ. Math. Inst. Hautes Etudes Sci. 121 (2015), 363–382.
- [CM1] T.H. Colding and W.P. Minicozzi II, *Singularities of Ricci flow and diffeomorphisms*, preprint.
- [CM2] T.H. Colding and W.P. Minicozzi II, *Uniqueness of blowups and Lojasiewicz inequalities*. Ann. of Math. (2) 182 (2015), no. 1, 221–285.
- [H] R. Hamilton, *The formation of singularities in the Ricci flow*. Surveys in differential geometry, Vol. II (Cambridge, MA, 1993), 7–136, Int. Press, Cambridge, MA, 1995.
- [LW] Y. Li and B. Wang, *Rigidity of the round cylinders in Ricci shrinkers*, preprint.