

Transversal designs and induced decompositions of graphs*

CSILLA BUJTÁS AND ZSOLT TUZA

Dedicated to Adrian Bondy on the occasion of his 70th birthday

We prove that for every complete multipartite graph F there exist very dense graphs G_n on n vertices, namely with as many as $\binom{n}{2} - cn$ edges for all n , for some constant $c = c(F)$, such that G_n can be decomposed into edge-disjoint *induced* subgraphs isomorphic to F . This result identifies and structurally explains a gap between the growth rates $O(n)$ and $\Omega(n^{3/2})$ on the minimum number of non-edges in graphs admitting an induced F -decomposition.

AMS 2000 SUBJECT CLASSIFICATIONS: Primary 05C35, 05C70; secondary 05C75.

KEYWORDS AND PHRASES: Induced subgraph, edge decomposition, complete multipartite graph, extremal graph theory.

1	Introduction	258
1.1	The problem to be studied	258
1.2	A short summary of earlier asymptotic results	259
1.3	Our main contributions	259
2	Edge decompositions of complete k-partite graphs	260
2.1	Transversal designs	260
2.2	Decompositions of complete k -partite graphs	261
3	Proof of the linear upper bound	262
4	Complete k-partite graphs and transversal designs	264
5	Concluding remarks	266
	References	268

arXiv: [1501.03518](https://arxiv.org/abs/1501.03518)

*Research supported in part by the Hungarian State and the European Union under the grant TAMOP-4.2.2.A-11/1/KONV-2012-0072.

1. Introduction

In this paper we study an extremal problem raised by Bondy and Szwarcfiter [3] in 2013, and extend the results proved in [3, 6, 9]. Throughout, we consider simple undirected graphs, with the central concept of induced decomposition, under which we mean edge-decomposition into induced subgraphs.

1.1. The problem to be studied

Definition 1. For two graphs F and G , an *induced F -decomposition* of G is a collection $\{F_1, \dots, F_\ell\}$ of induced subgraphs of G such that

- each F_i is isomorphic to F ($1 \leq i \leq \ell$),
- the subgraphs are pairwise edge-disjoint, and
- $\bigcup_{i=1}^{\ell} E(F_i) = E(G)$.

Clearly, for any fixed F different from K_2 , not all graphs G admit an induced F -decomposition. Especially, if F is not a complete graph and the induced decomposition exists, then G cannot be complete. Therefore, the following problem was proposed by Bondy and Szwarcfiter.

Problem 2 ([3]). *Given a non-complete graph F and a positive integer $n \geq |V(F)|$, determine the maximum number $\text{ex}^*(n, F)$ of edges in a graph G of order n which admits an induced F -decomposition.*

For the asymptotic¹ version of this problem, the following basic result was proved by Cohen and Tuza.

Theorem 3 ([6]). *For every graph F we have $\text{ex}^*(n, F) = \binom{n}{2} - o(n^2)$.*

Note that this theorem is analogous to a result of Frankl and Füredi from [8]; the latter one is discussed in a more general framework.

Theorem 3 suggests that in fact the ‘complementary’ function

$$\overline{\text{ex}}^*(n, F) = \binom{n}{2} - \text{ex}^*(n, F)$$

is the appropriate subject of study, as it is more informative when asymptotic results are desired.

¹For the asymptotic comparison of the growth orders of functions we use the standard notation O , o , Ω , Θ of complexity theory; cf. e.g. https://en.wikipedia.org/wiki/Big_O_notation#Family_of_Bachmann.E2.80.93Landau_notations.

1.2. A short summary of earlier asymptotic results

- If F is a complete graph then, by Wilson’s theorem [10], we infer that for each k the equality $\overline{\text{ex}^*}(n, K_k) = 0$ holds with infinitely many values of n . In general, $\overline{\text{ex}^*}(n, K_k) = O(n)$ holds for every $k \geq 2$.
- As observed in [3], if F' is obtained from F by extending it with an isolated vertex,

$$\overline{\text{ex}^*}(n, F) \leq \overline{\text{ex}^*}(n, F') \leq \overline{\text{ex}^*}(n - 1, F) + n - 1$$

holds.

- If F is isolate-free and non-complete then $\overline{\text{ex}^*}(n, F) = \Omega(n)$. This was proved recently by Halász and Tuza [9]. Particularly, this linear lower bound is asymptotically sharp; that is, $\overline{\text{ex}^*}(n, F) = \Theta(n)$ holds if F is a complete bipartite or tripartite graph which is non-complete [9].
- If F is isolate-free, non-complete and not a complete multipartite graph, then $\overline{\text{ex}^*}(n, F) = \Omega(n^{3/2})$ [9]. This bound was proved to be asymptotically sharp for $P_4, C_6, 2K_2$ and $K_4 - P_3$ in [3, 6, 9].² Moreover, as a consequence of results in [1], with Noga Alon we proved that $\overline{\text{ex}^*}(n, F) = \Theta(n^{3/2})$ is valid for every disconnected graph F whose each component is complete bipartite [2].

A more detailed summary of results (not only of the asymptotic ones) and open problems is given in [4].

Concerning the lower bound $\Omega(n)$ in the third point above, one can observe that the exclusion of isolated vertices is in fact irrelevant.

Proposition 4. *If each vertex of F has a non-neighbor (and, in particular, if F has an isolated vertex), then $\overline{\text{ex}^*}(n, F) \geq n/2$.*

Proof. Under the given assumption, if G admits an induced F -decomposition then every vertex of G , too, must have a non-neighbor. □

1.3. Our main contributions

The most important contribution of this paper is as follows.

Theorem 5. *If F is a complete multipartite graph, then $\overline{\text{ex}^*}(n, F) = O(n)$.*

²Here P_4 is the path on four vertices, C_6 is the 6-cycle, $2K_2$ is the graph with four vertices and two vertex-disjoint edges, and $K_4 - P_3$ is the “paw”, which is obtained from K_4 by deleting two edges incident with a common vertex.

Together with the earlier results cited above and with Proposition 4, Theorem 5 immediately implies the following consequences. The first one is the characterization of linear growth, while the second one identifies a gap in the exponents of possible growth functions for $\overline{\text{ex}}^*(n, F)$.

Corollary 6. *We have $\overline{\text{ex}}^*(n, F) = \Theta(n)$ if and only if either F is a complete graph plus at least one isolated vertex, or F is a complete multipartite graph having at least one non-edge, possibly together with some isolated vertices.*

Corollary 7. *If $1 < c < 3/2$, there exists no F satisfying $\overline{\text{ex}}^*(n, F) = \Theta(n^c)$.*

In some steps of the constructions we apply design-theoretic methods, in particular ‘transversal designs’, whose formal definition is given in Section 2. Using those structures the following result will be proved.

Theorem 8. *Let $F = K_{a_1, \dots, a_k}$, and suppose that there exists a transversal design $TD(k, a_i)$ for each $i = 1, 2, \dots, k$. Then the complete multipartite graph $F^* = K_{ma_1, \dots, ma_k}$ with $m = \prod_{i=1}^k a_i$ admits an induced F -decomposition. Moreover, such a decomposition can be encoded with m^2 integer sequences of length $2k$ whose i^{th} and $(k+i)^{\text{th}}$ terms vary between 1 and a_i .*

Note that every F -subgraph of F^* necessarily is an induced subgraph because both of them are complete k -partite.

After some preliminary observations in Section 2, we prove our main result, Theorem 5 in Section 3. The proof of Theorem 8 is given in Section 4. We close the paper with several open problems and conjectures.

2. Edge decompositions of complete k -partite graphs

In this section we make some preparations towards the proofs of Theorems 5 and 8. In the first subsection we recall some important known facts from the theory of transversal designs. In the second subsection we construct decompositions of some restricted classes of graphs — having the same number of partite classes that the fixed given graph F has — which will be applied later for the general results.

2.1. Transversal designs

In general, a *transversal design* has three parameters: the ‘order’ or ‘group-size’ n , the ‘blocksize’ k , and the ‘index’ λ . For our purpose, however, only

the case $\lambda = 1$ will be relevant, therefore we disregard the general definition. Hence, the combinatorial structure denoted $TD(k, n)$ is a triple $(X, \mathcal{G}, \mathcal{B})$, where

1. X is a set of kn elements;
2. $\mathcal{G} = (X_1, X_2, \dots, X_k)$ is a partition of X into k classes X_i (the groups), each of size n ;
3. \mathcal{B} is a collection of k -element subsets of X (the blocks);
4. every unordered pair of elements from X is contained either in exactly one group or in exactly one block, but not both.

The more widely known structures of Latin squares are arrangements of the numbers $1, 2, \dots, n$ in an $n \times n$ square in such a way that each number occurs precisely once in each row and also in each column. Two Latin squares $(a_{ij})_{1 \leq i, j \leq n}$ and $(b_{ij})_{1 \leq i, j \leq n}$ are said to be *orthogonal* if their position-wise concatenation $\{a_{ij}b_{ij} \mid 1 \leq i \leq n, 1 \leq j \leq n\}$ lists all the n^2 ordered pairs of $\{1, 2, \dots, n\} \times \{1, 2, \dots, n\}$.

We shall apply the following two facts, the first of which is not very hard to prove.

Lemma 9. (see Theorem 3.18 in [7]) *The existence of a $TD(k, n)$ is equivalent to the existence of $k - 2$ mutually orthogonal Latin squares of order n .*

Lemma 10. (Chowla, Erdős and Straus, [5]) *The maximum number of mutually orthogonal Latin squares of order n tends to infinity as $n \rightarrow \infty$.*

2.2. Decompositions of complete k -partite graphs

Next, we describe some steps which make it possible to apply transversal designs to the general decomposition problem. We begin with a definition.

Definition 11. Let $G = (V, E)$ and $F = (X, H)$ be complete k -partite graphs with vertex partitions $V = V_1 \cup \dots \cup V_k$ and $X = X_1 \cup \dots \cup X_k$. Assume further that p is an integer and that $|V_i| = p|X_i|$ for all $1 \leq i \leq k$; hence $|V| = p|X|$. An *embedded F -decomposition of G* is an (induced) F -decomposition with the further property that each V_i is partitioned into p sets,

$$V_i = V_{i,1} \cup \dots \cup V_{i,p}$$

and, for each copy of F in the decomposition, each partite set X_i coincides with some $V_{i,j}$ inside V_i .

Lemma 12. *Let k and a_1, \dots, a_k be fixed natural numbers. Then, for every sufficiently large integer p , the complete multipartite graph K_{pa_1, \dots, pa_k} has an embedded K_{a_1, \dots, a_k} -decomposition.*

Proof. Based on Lemma 10 we can choose a threshold value $p_0 = p_0(k)$ such that there exist at least $k - 2$ mutually orthogonal Latin squares of order p whenever $p > p_0$ holds. Then, by Lemma 9, there exists a transversal design $TD(k, p)$ of order p and blocksize k . Observe that $TD(k, p)$ is precisely an edge decomposition of the complete k -partite graph $K_{p, \dots, p}$ into copies of the complete graph K_k . Substituting mutually disjoint sets of cardinality a_i for all vertices in the i^{th} group of $TD(k, p)$ for $i = 1, 2, \dots, k$, each copy of K_k becomes a copy of K_{a_1, \dots, a_k} . Thus, an embedded K_{a_1, \dots, a_k} -decomposition of K_{pa_1, \dots, pa_k} is obtained. \square

Choose p_0 as in the proof above, and let $a_i^* = pa_i$ for $i = 1, 2, \dots, k$ with an arbitrarily fixed $p > p_0$. Then we obtain:

Corollary 13. *For every k -tuple (a_1, \dots, a_k) of positive integers there exists a k -tuple (a_1^*, \dots, a_k^*) such that*

- $K_{a_1^*, \dots, a_k^*}$ has an embedded K_{a_1, \dots, a_k} -decomposition;
- a transversal design $TD^i = TD(k, a_i^*)$ exists for each $i = 1, 2, \dots, k$.

The applicability of this approach is demonstrated by Theorem 8, which was stated in the Introduction and will be proved in Section 4.

3. Proof of the linear upper bound

In this section we prove Theorem 5.

Let $F = K_{a_1, \dots, a_k}$ be a fixed given graph. With reference to Lemma 12 we choose an integer $p > 1$ such that

- p is a multiple of $\prod_{i=1}^k a_i$,
- $F_p := K_{pa_1, \dots, pa_k}$ has an embedded F -decomposition.

The general form of Wilson’s theorem yields that there exists a modulus q and a residue r such that the complete graph K_{sq+r} has an F -decomposition whenever $s \geq s_0 = s_0(F)$ is sufficiently large. Note that, regarding the linear upper bound to be proved, we may assume that n is large. We write n in the form

$$n = (sq + r) \cdot p + t \quad (s \geq s_0, \quad 0 \leq t \leq pq - 1)$$

and define

$$n' = sq + r.$$

Step 1. Decompose $K_{n'}$ into edge-disjoint copies of F . Of course, those copies are non-induced subgraphs of $K_{n'}$ (unless F itself is a complete graph, which case we disregard).

Step 2. Replace each vertex of $K_{n'}$ with an independent set of cardinality p . That is, two vertices are adjacent if and only if they belong to different independent sets; in this way the graph

$$G' := K_{(sq+r) \cdot p} - (sq + r) \cdot K_p = K_{n-t} - \frac{n-t}{p} \cdot K_p$$

is obtained. Moreover, each copy of F in $K_{n'}$ becomes a copy of F_p in G' . Observe that, although the copies of F_p are still non-induced, the i^{th} vertex class of cardinality pa_i is the union of a_i independent sets of size p each.

Step 3. In any copy of F_p , if an independent set S of size p belongs to the i^{th} vertex class of F_p , then partition S into p/a_i disjoint sets. In this way the a_i independent p -element sets inside the i^{th} vertex class together yield a partition of that class into p independent sets of size a_i each.

Step 4. Denote by V_1, \dots, V_k the vertex classes in the copy of F_p , and by $V_{i,j}$ the sets in the partition obtained ($1 \leq i \leq k, 1 \leq j \leq p, |V_{i,j}| = a_i$ for all i and j). Then the embedded F -decomposition of F_p — guaranteed by Lemma 12 and by the choice of p — yields p^2 copies of F which are *induced* subgraphs of G' .

Step 5. Complete the construction with t isolated vertices, to obtain a graph G . It is clear that this G admits an induced F -decomposition.

It remains to observe that the complement of G has no more than a linear number of edges. There are $\binom{t}{2} + tn - t^2 < tn < pqn$ vertex pairs containing at least one of the t isolated vertices; and the number of non-edges in G' is exactly $\frac{n-t}{p} \cdot \binom{p}{2} < pn/2$. Thus,

$$\overline{\text{ex}^*}(n, F) \leq \binom{n}{2} - |E(G)| < (pq + p/2) \cdot n = O(n).$$

This completes the proof of Theorem 5.

4. Complete k -partite graphs and transversal designs

Here we prove Theorem 8. Recall that $F = K_{a_1, \dots, a_k}$, $F^* = K_{ma_1, \dots, ma_k}$, and $m = \prod_{i=1}^k a_i$. It is now assumed that there exists a transversal design $\text{TD}(k, a_i) = (X^i, \mathcal{G}^i, \mathcal{B}^i)$ for every $i = 1, 2, \dots, k$, which we shall refer to as TD^i or $\text{TD}^i(k, a_i)$. Those TD^i will become important in the last step of the construction.

In order to simplify notation throughout, we write the usual shorthand $[a_i] = \{1, 2, \dots, a_i\}$.

Notation. For $1 \leq i \leq k$, $1 \leq p \leq a_i$, and $1 \leq q \leq k$, the p^{th} element in the q^{th} group X_q^i of $\text{TD}^i(k, a_i)$ will be denoted by $\pi^i(p, q)$. (Inside each group the elements are numbered as $1, 2, \dots, a_i$.) Note that for all $1 \leq p, p' \leq a_i$ and all $1 \leq q, q' \leq k$ ($q \neq q'$), a block in \mathcal{B}^i containing the p^{th} element of the q^{th} group and the $(p')^{\text{th}}$ element of the $(q')^{\text{th}}$ group in $\text{TD}^i(k, a_i)$ exists and is unique.

Construction. Let the vertex set of F^* be

$$V = V^1 \cup \dots \cup V^k, \quad |V^i| = ma_i \text{ for } i = 1, \dots, k.$$

Perhaps the best way of thinking about V^i is to view it as a k -dimensional box composed of unit-cube cells containing a_i items each; in the j^{th} direction the box has size a_j . In this view we see a_i layers orthogonal to the i^{th} direction, each of those layers consists of m/a_i cells, and hence each layer contains precisely m items.

To state this in a more formal description, we partition each V^i into m disjoint sets of cardinality a_i (corresponding to the contents of the cells), and represent each of those subsets inside V^i with a vector

$$\mathbf{j} = (j_1, \dots, j_k) \in [a_1] \times \dots \times [a_k]$$

of length k , so that

$$V^i = \bigcup_{\mathbf{j} \in [a_1] \times \dots \times [a_k]} V_{\mathbf{j}}^i.$$

In this way a copy of F inside F^* is characterized by a k -tuple of vectors,

$$(\mathbf{j}^1, \dots, \mathbf{j}^k) \in ([a_1] \times \dots \times [a_k])^k,$$

where

$$\mathbf{j}^i = (j_1^i, \dots, j_k^i) \in [a_1] \times \dots \times [a_k]$$

for $i = 1, 2, \dots, k$. Let us refer to this $(\mathbf{j}^1, \dots, \mathbf{j}^k)$ as the *detailed representation* of the copy in question.

There are $(ma_i)(ma_{i'}) = m^2 a_i a_{i'}$ edges of F^* between V^i and $V^{i'}$, and each copy of F contains $a_i a_{i'}$ edges between those two partite sets. Thus, we have to define m^2 edge-disjoint copies of F inside F^* . The key point is how to shuffle the vertex sets of the m^2 copies to ensure that those subgraphs compose a partition of the edge set of F^* .

The copies of F will be encoded with vectors of length $2k$ (i.e., a suitable collection of k^2 -dimensional vectors will be encoded with $(2k)$ -dimensional ones), which we shall view as concatenations of two vectors

$$(\mathbf{b}, \mathbf{c}) \in ([a_1] \times \dots \times [a_k])^2$$

where

$$\begin{aligned} \mathbf{b} &= (b_1, \dots, b_k) \in [a_1] \times \dots \times [a_k], \\ \mathbf{c} &= (c_1, \dots, c_k) \in [a_1] \times \dots \times [a_k] \end{aligned}$$

are vectors of length k .

We identify a copy of F for any given (\mathbf{b}, \mathbf{c}) in the way detailed next.

- Generally speaking, the first k coordinates (i.e., those of \mathbf{b}) specify a k -tuple of layers (one layer in each of the boxes V^i) that means a copy of $K_{m, \dots, m}$; and the role of \mathbf{c} is similar, but “rotating” by 1 among the boxes, e.g. while the first coordinate of \mathbf{b} will provide a data for V^1 , the first coordinate of \mathbf{c} will be relevant for V^2 , hence specifying a second copy of $K_{m, \dots, m}$. After that, the transversal designs will be used for defining further $k - 2$ layers inside each V^i . Having all this at hand, exactly k layers are determined inside each box, and all the intersections of those layers in the different boxes together uniquely identify a copy of F associated with (\mathbf{b}, \mathbf{c}) .
- The i^{th} coordinate b_i of \mathbf{b} determines that the i^{th} partite set of F (inside V^i) must be one of those $V_{\mathbf{j}^i}$ which have $j_i = b_i$. This equivalently means that $j_i^i = b_i$ holds in the detailed representation. (Recall that \mathbf{j}^i has the form $\mathbf{j}^i = (j_1^i, \dots, j_k^i)$.)
- The i^{th} coordinate c_i of \mathbf{c} determines the i^{th} coordinate j_i^{i+1} of the next \mathbf{j}^{i+1} in the detailed representation. (For $i = k$ we mean $\mathbf{j}^{k+1} = \mathbf{j}^1$.)
- By the definition of the transversal designs $\text{TD}^i(k, a_i)$, there is a unique block $B^i = B^i(\mathbf{b}, \mathbf{c}) \in \mathcal{B}^i$ such that both $\pi^i(b_i, i) \in B^i$ and $\pi^i(c_i, i + 1) \in B^i$. (Also here, the successor $i + 1$ of $i = k$ is meant to be 1.) Then the i^{th} coordinate of the copy of F associated with

(\mathbf{b}, \mathbf{c}) in $V^{i'}$, that is $j_i^{i'}$ in the representation $(\mathbf{j}^1, \dots, \mathbf{j}^k)$, is defined as $j_i^{i'} = B^i \cap X_{i'}^i$ for all i' . This is our defining rule for $i' \notin \{i, i + 1\}$, but the formula is valid also for $i' \in \{i, i + 1\}$ because this was the way we selected B^i .

Verification of requirements. It is clear that precisely m^2 copies are defined in this way. Since the total number of edges in those copies is equal to that in F^* , the proof will be done if we show that each edge of F^* is contained in some copy of F . This is equivalent to saying that each edge $v_i v_{i'}$ ($v_i \in V^i, v_{i'} \in V^{i'}$) determines all the k^2 coordinates of some k -tuple of vectors $(\mathbf{j}^1, \dots, \mathbf{j}^k)$. The property is valid indeed, as can be seen from the following considerations.

The positions of v_i and $v_{i'}$ determine the representing vectors \mathbf{j}^i and $\mathbf{j}^{i'}$ of the a_i -element and $a_{i'}$ -element sets (cells) containing v_i and $v_{i'}$, respectively. This determines k pairs of coordinates:

$$(j_1^i, j_1^{i'}), (j_2^i, j_2^{i'}), \dots, (j_k^i, j_k^{i'}).$$

The pair $(j_\ell^i, j_\ell^{i'})$ uniquely determines the block $B^\ell = B^\ell(\mathbf{b}, \mathbf{c}) \in \mathcal{B}^\ell$ in $\text{TD}^\ell(k, a_\ell)$ for $\ell = 1, \dots, k$; this B^ℓ is the unique block containing the $(j_\ell^i)^{\text{th}}$ element of the i^{th} block and the $(j_\ell^{i'})^{\text{th}}$ element of the $(i')^{\text{th}}$ block in $\text{TD}^\ell(k, a_\ell)$. Thus, all the k^2 coordinates of the vectors $\mathbf{j}^1, \dots, \mathbf{j}^k$ can be computed from those k blocks by the rule $j_\ell^i = B^\ell \cap X_i^\ell$.

This completes the proof of the theorem.

5. Concluding remarks

In this paper we studied the growth of the number of edges in graphs which admit edge decompositions into induced subgraphs isomorphic to a given complete multipartite graph. We proved a linear upper bound on the number of non-edges. More explicitly, our construction yields the following.

Theorem 14. *If n satisfies the following three conditions:*

- $\binom{n}{2}$ is divisible by the number $\sum_{1 \leq i < i' \leq k} a_i a_{i'}$ of edges in F ,
- $n - 1$ is divisible by the greatest common divisor of the vertex degrees $\left(\sum_{i=1}^k a_i\right) - a_1, \left(\sum_{i=1}^k a_i\right) - a_2, \dots, \left(\sum_{i=1}^k a_i\right) - a_k$,
- $n > n_0$ for some threshold value $n_0 = n_0(a_1, \dots, a_k)$,

then the complete n -partite graph $K_{m, \dots, m}$ with $m = \prod_{i=1}^k a_i$ admits an edge decomposition into induced subgraphs isomorphic to F .

Comparison with the results from [9] identifies a gap in the exponent of n ; this can be expressed in the following way, where the first case is trivial and the second case is taken from Wilson's theorem.

Let F be a graph without isolated vertices.

- If $F = K_2$, then $\overline{\text{ex}^*}(n, F) = 0$.
- If F is a complete graph of order at least 3, then $\overline{\text{ex}^*}(n, F)$ oscillates between 0 and $\Theta(n)$, reaching both extremities on sets of positive upper density.
- If F is a complete multipartite graph, but not a complete graph, then $\overline{\text{ex}^*}(n, F) = \Theta(n)$.
- If F is not a complete multipartite graph, then $\overline{\text{ex}^*}(n, F) = \Omega(n^{3/2})$.

The next task could be to identify further gaps in the exponent of n describing the growth of $\overline{\text{ex}^*}(n, F)$, if there are any. We believe that not all reals (and not all rational numbers) between $3/2$ and 2 can occur as exponents for some F , and also that the gap between n and $n^{3/2}$ is not the only one. This is expressed in the following sequence of conjectures.

Conjecture 15. *There exist graphs F for which*

$$\lim_{n \rightarrow \infty} \frac{\overline{\text{ex}^*}(n, F)}{n^{3/2}} = \infty.$$

Conjecture 16. *For every non-complete graph F , without isolated vertices, there exists a constant $c = c(F)$ such that $\overline{\text{ex}^*}(n, F) = \Theta(n^c)$.*

Conjecture 17. *The constant $c = c(F)$ in Conjecture 16 is rational for every graph F .*

Conjecture 18. *Every non-complete graph F without isolated vertices satisfies the asymptotic equality $\overline{\text{ex}^*}(n, F) = \Theta(n^{2-1/t})$ where $t = t(F)$ is a positive integer.*

If Conjecture 18 is true, the next question is: Which structural property of F determines $t(F)$?

Conjecture 19. *For every non-complete graph F without isolated vertices, the quotient*

$$\frac{\overline{\text{ex}^*}(n, F)}{n^{2-1/t}}$$

tends to a positive limit as n gets large, for some positive integer $t = t(F)$.

If this is true, we would be interested in the exact value of the limit.

Problem 20. Verify Conjectures 18 and 19 for all graphs on at most six vertices.

The smallest open case for Conjecture 18 is $F = C_5$, while the one for Conjecture 19 is $F = 2K_2$.

Hypergraphs. It is very natural to extend the above questions to r -uniform hypergraphs ($r \geq 3$), but rather too little is known so far about the analogous function $\overline{\text{ex}}^*(n, \mathcal{F})$. Already the study of some very small examples may provide interesting pieces of information.

References

- [1] N. Alon, A. Moitra, and B. Sudakov (2013). Large induced matchings and their applications. *J. Europ. Math. Soc. (JEMS)* **15** 1575–1596. [MR3082238](#)
- [2] N. Alon and Zs. Tuza (2014). Unpublished, July 2014.
- [3] J. A. Bondy and J. L. Szwarcfiter (2013). Induced decompositions of graphs. *J. Graph Theory* **72** 462–477. [MR3021353](#)
- [4] Cs. Bujtás and Zs. Tuza (2014). Problems and results on induced subgraph decompositions of graphs. *Proc. ASCONIKK 2014: Extended Abstracts III. Future Internet Technologies*, ISBN 978-963-396-048-6, University of Pannonia, Veszprém, pp. 11–18.
- [5] P. Erdős, S. A. Chowla, and E. G. Straus (1960). On the maximal number of pairwise orthogonal latin squares of a given order. *Canad. J. Math.* **13** 204–208. [MR0122730](#)
- [6] N. Cohen and Zs. Tuza (2015). Induced decompositions of highly dense graphs. *J. Graph Theory* **78** 97–107. [MR3293078](#)
- [7] C. J. Colbourn and J. H. Dinitz, Eds. (2007). *Handbook of Combinatorial Designs*, 2nd ed. Chapman & Hall/CRC. [MR2246267](#)
- [8] P. Frankl and Z. Füredi (1987). Colored packing of sets. *Ann. Discrete Math.* **34** 165–178. [MR0920642](#)
- [9] V. Halász and Zs. Tuza (2014). Asymptotically optimal induced decompositions. *Applicable Anal. Discrete Math.* **8** 320–329. [MR3243239](#)
- [10] R. M. Wilson (1976). Decompositions of complete graphs into subgraphs isomorphic to a given graph. Proc. 5th British Combinatorial Conference, *Congr. Numer.* **XV** 647–659. [MR0396347](#)

CSILLA BUJTÁS

DEPARTMENT OF COMPUTER SCIENCE AND SYSTEMS TECHNOLOGY

UNIVERSITY OF PANNONIA

H-8200 VESZPRÉM

EGYETEM U. 10

HUNGARY

E-mail address: bujtas@dcs.uni-pannon.hu

ZSOLT TUZA

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS

HUNGARIAN ACADEMY OF SCIENCES

H-1053 BUDAPEST

REÁLTANODA U. 13-15

HUNGARY

AND

DEPARTMENT OF COMPUTER SCIENCE AND SYSTEMS TECHNOLOGY

UNIVERSITY OF PANNONIA

VESZPRÉM

HUNGARY

E-mail address: tuza@dcs.uni-pannon.hu

RECEIVED 14 JANUARY 2015