

Multiple bracket function, Stirling number, and Lah number identities

HASAN COSKUN

The author has constructed multiple analogues of several families of combinatorial numbers in a recent article, including the bracket symbol, and the Stirling numbers of the first and second kind. In the present paper, a multiple analogue of another sequence, the Lah numbers, is developed, and certain associated identities and significant properties of all these sequences are constructed.

AMS 2000 SUBJECT CLASSIFICATIONS: Primary 05A10, 11B65; secondary 33D67.

KEYWORDS AND PHRASES: Multiple combinatorial numbers, multiple binomial coefficients, multiple bracket function, multiple Stirling numbers, multiple Lah numbers, well poised rational Macdonald functions.

1. Introduction

The Stirling numbers of the first and the second kind have been studied and their properties have been investigated extensively in number theory, combinatorics and other areas. One dimensional generalizations of these numbers have also been subject of interest. An important class of generalizations is their one parameter q -extensions. Many have made significant contributions to the q -extensions investigating their properties and applications. We will give references to some of these important works in Section 5 below.

In a recent paper, the author took a major step from one dimensional generalizations to multiple analogues of combinatorial numbers by constructing elegant multiple qt -generalizations of Stirling numbers of the first and second kind, besides sequences of other combinatorial numbers including multiple binomial, Fibonacci, Bernoulli, Catalan, and Bell numbers [9]. In this paper, we focus on the multiple analogies of the factorial function, Stirling numbers of both kinds, and Lah numbers, and give interesting new identities they satisfy.

The multiple generalizations developed in [9] are given in terms of the qt -binomial coefficients constructed in the same paper. Its definition may be written in the general form as

$$\binom{z}{\mu}_{q,t} := \frac{q^{|\mu|} t^{2n(\mu)+(1-n)|\mu|}}{(qt^{n-1})_\mu} \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\mu_i - \mu_j}}{(qt^{j-i-1})_{\mu_i - \mu_j}} \right\} w_\mu(q^z t^{\delta(n)}; q, t)$$

where μ is a partition of at most n parts, $z \in \mathbb{C}^n$ and $q, t \in \mathbb{C}$. The w_μ function that enters the definition is a limiting case of the BC_n well-poised symmetric rational Macdonald function W_λ . Note that this definition makes sense even when μ is not an integer partition, but is a vector $\mu \in \mathbb{C}^n$. The qt -binomial coefficients are constructed independently by Kaneko [18] in a special case, and Okounkov [25] using difference operator methods for integer partitions. Lassalle constructed an equivalent set of qt -binomial coefficients independently [23], and Sahi has developed the non-symmetric version [28]. Rains [27] used Okounkov’s BC_n symmetric interpolation polynomials to define a three parameter qts -binomial coefficients. The most general definition (29) of multiple qt -binomial coefficients we constructed in [9] where $z \in \mathbb{C}^n$ (and $\mu \in \mathbb{C}^n$) are variables is essential in the development of the multiple brackets, and thus the Stirling and Lah numbers in this paper.

Please note that for each multiple identity proved in this paper, we included a statement about the corresponding one dimensional classical identity or its q -analogue for comparison, right before or after the result. The reader who is not familiar with the theory of multiple hypergeometric summation identities should review the corresponding one dimensional result before studying the details of the multiple generalization. This should make it easier to follow the construction.

2. Background

The (basic) q -Pochhammer symbol $(a; q)_\alpha$ may be defined formally for complex parameters $q, \alpha \in \mathbb{C}$ as

$$(1) \quad (a)_\alpha = (a; q)_\alpha := \frac{(a; q)_\infty}{(aq^\alpha; q)_\infty}$$

where the infinite product $(a; q)_\infty$ is defined by $(a; q)_\infty := \prod_{i=0}^\infty (1 - aq^i)$. Note that when $\alpha = m$ is a positive integer, the definition reduces to the finite product $(a; q)_m = \prod_{k=0}^{m-1} (1 - aq^k)$. An elliptic analogue is defined [11] by

$$(2) \quad (a; q, p)_m := \prod_{k=0}^{m-1} \theta(aq^m)$$

where $a \in \mathbb{C}$, m is a positive integer, and the normalized theta function $\theta(x)$ is given by

$$(3) \quad \theta(x) = \theta(x; p) := (x; p)_\infty (p/x; p)_\infty$$

for $x, p \in \mathbb{C}$ with $|p| < 1$. The definition is extended to negative m by setting $(a; q, p)_m = 1/(aq^m; q, p)_{-m}$. It is clear that when $p = 0$, the elliptic $(a; q, p)_m$ reduces to the basic (trigonometric) q -Pochhammer symbol (1).

For any partition $\lambda = (\lambda_1, \dots, \lambda_n)$ and $t \in \mathbb{C}$, define [30]

$$(4) \quad (a)_\lambda = (a; q, p, t)_\lambda := \prod_{i=1}^n (at^{1-i}; q, p)_{\lambda_i}.$$

Note that when $\lambda = (\lambda_1) = \lambda_1$ is a single part partition, then $(a; q, p, t)_\lambda = (a; q, p)_{\lambda_1} = (a)_{\lambda_1}$. For brevity of notation, we also use

$$(5) \quad (a_1, \dots, a_k)_\lambda = (a_1, \dots, a_k; q, p, t)_\lambda := (a_1)_\lambda \dots (a_k)_\lambda.$$

Recall that we use V to denote [8] the space of infinite lower-triangular matrices whose entries are rational functions over the field $\mathbb{F} = \mathbb{C}(q, p, t, r, a, b)$ which are indexed by partitions with respect to the partial inclusion ordering \subseteq defined by

$$(6) \quad \mu \subseteq \lambda \Leftrightarrow \mu_i \leq \lambda_i, \quad \forall i \geq 1.$$

The condition that a matrix $u \in V$ is lower triangular with respect to \subseteq can be stated in the form

$$(7) \quad u_{\lambda\mu} = 0, \quad \text{when } \mu \not\subseteq \lambda.$$

The multiplication operation defined by

$$(8) \quad (uv)_{\lambda\mu} := \sum_{\nu \subseteq \lambda \subseteq \mu} u_{\lambda\nu} v_{\nu\mu}$$

for matrices $u, v \in V$ makes V into an \mathbb{F} -algebra.

2.1. Well-poised Macdonald functions

The construction of our multiple Stirling numbers involves the elliptic well-poised Macdonald functions $W_{\lambda/\mu}$, and elliptic well-poised Jackson coefficients $\omega_{\lambda/\mu}$ on BC_n [8]. These remarkable families of symmetric rational functions are first introduced in the author's Ph.D. thesis [6] in the basic (trigonometric) case, and later in [7] in the more general elliptic form.

Let $\lambda = (\lambda_1, \dots, \lambda_n)$ and $\mu = (\mu_1, \dots, \mu_n)$ be partitions of at most n parts for a positive integer n such that the skew partition λ/μ is a hori-

zontal strip; i.e. $\lambda_1 \geq \mu_1 \geq \lambda_2 \geq \mu_2 \geq \dots \lambda_n \geq \mu_n \geq \lambda_{n+1} = \mu_{n+1} = 0$. Following [8], define

$$(9) \quad H_{\lambda/\mu}(q, p, t, b) := \prod_{1 \leq i < j \leq n} \left\{ \frac{(q^{\mu_i - \mu_{j-1}} t^{j-i})_{\mu_{j-1} - \lambda_j} (q^{\lambda_i + \lambda_j} t^{3-j-i} b)_{\mu_{j-1} - \lambda_j}}{(q^{\mu_i - \mu_{j-1} + 1} t^{j-i-1})_{\mu_{j-1} - \lambda_j} (q^{\lambda_i + \lambda_j + 1} t^{2-j-i} b)_{\mu_{j-1} - \lambda_j}} \cdot \frac{(q^{\lambda_i - \mu_{j-1} + 1} t^{j-i-1})_{\mu_{j-1} - \lambda_j}}{(q^{\lambda_i - \mu_{j-1}} t^{j-i})_{\mu_{j-1} - \lambda_j}} \right\} \cdot \prod_{1 \leq i < (j-1) \leq n} \frac{(q^{\mu_i + \lambda_j + 1} t^{1-j-i} b)_{\mu_{j-1} - \lambda_j}}{(q^{\mu_i + \lambda_j} t^{2-j-i} b)_{\mu_{j-1} - \lambda_j}}$$

and

$$(10) \quad W_{\lambda/\mu}(x; q, p, t, a, b) := H_{\lambda/\mu}(q, p, t, b) \cdot \frac{(x^{-1}, ax)_\lambda (qbx/t, qb/(axt))_\mu}{(x^{-1}, ax)_\mu (qbx, qb/(ax))_\lambda} \cdot \prod_{i=1}^n \left\{ \frac{\theta(bt^{1-2i} q^{2\mu_i})}{\theta(bt^{1-2i})} \frac{(bt^{1-2i})_{\mu_i + \lambda_{i+1}}}{(bqt^{-2i})_{\mu_i + \lambda_{i+1}}} \cdot t^{i(\mu_i - \lambda_{i+1})} \right\}$$

where $q, p, t, x, a, b \in \mathbb{C}$. Note that $W_{\lambda/\mu}(x; q, p, t, a, b)$ vanishes unless λ/μ is a horizontal strip. The function $W_{\lambda/\mu}(y, z_1, \dots, z_\ell; q, p, t, a, b)$ is extended to $\ell + 1$ variables $y, z_1, \dots, z_\ell \in \mathbb{C}$ through the following recursion formula

$$(11) \quad W_{\lambda/\mu}(y, z_1, z_2, \dots, z_\ell; q, p, t, a, b) = \sum_{\mu \prec \nu \prec \lambda} W_{\lambda/\nu}(yt^{-\ell}; q, p, t, at^{2\ell}, bt^\ell) W_{\nu/\mu}(z_1, \dots, z_\ell; q, p, t, a, b).$$

2.2. The limiting $w_{\lambda/\mu}$ function

The Macdonald functions W_λ are essentially equivalent to BC_n abelian functions constructed independently in [26]. The limiting cases defined above are closely related to the Macdonald polynomials [20], and interpolation Macdonald polynomials [25].

The following limiting the basic (the $p = 0$) case of the elliptic W functions will be used in our constructions below. The existence of these limits can be seen from (the $p = 0$ case of) the definition (10), the recursion formula (11) and the limit rule

$$(12) \quad \lim_{a \rightarrow 0} a^{|\mu|} (x/a)_\mu = (-1)^{|\mu|} x^{|\mu|} t^{-n(\mu)} q^{n(\mu')}$$

where $|\mu| = \sum_{i=1}^n \mu_i$ and $n(\mu) = \sum_{i=1}^n (i-1)\mu_i$, and $n(\mu') = \sum_{i=1}^n \binom{\mu_i}{2}$. We denote $H_{\lambda/\mu}(q, t) = H_{\lambda/\mu}(q, 0, t, 0)$, and for $x \in \mathbb{C}$ define

$$(13) \quad w_{\lambda/\mu}(x; q, t) := \lim_{s \rightarrow \infty} \left(s^{|\lambda|-|\mu|} \lim_{a \rightarrow 0} W_{\lambda/\mu}(x; q, t, a, as) \right) \\ = (-q/x)^{-|\lambda|+|\mu|} q^{-n(\lambda')+n(\mu')} H_{\lambda/\mu}(q, t) \frac{(x^{-1})_\lambda}{(x^{-1})_\mu}.$$

The recurrence formula for $w_{\lambda/\mu}$ function turns out to be

$$(14) \quad w_{\lambda/\mu}(y, z; q, t) = \sum_{\nu \prec \lambda} t^{\ell(|\lambda|-|\nu|)} w_{\lambda/\nu}(yt^{-\ell}; q, t) w_{\nu/\mu}(z; q, t).$$

Similarly, for $x \in \mathbb{C}$ define the dual function

$$(15) \quad \hat{w}_{\lambda/\mu}(x; q, t) := \lim_{s \rightarrow 0} \left(\lim_{a \rightarrow 0} W_{\lambda/\mu}(x; q, t, a, as) \right) \\ = t^{-n(\lambda)+|\mu|+n(\mu)} H_{\lambda/\mu}(q, t) \frac{(x^{-1})_\lambda}{(x^{-1})_\mu}.$$

The recurrence formula for the dual $\hat{w}_{\lambda/\mu}(x; q, t)$ may be written as

$$(16) \quad \hat{w}_{\lambda/\mu}(y, z; q, t) = \sum_{\mu \prec \nu \prec \lambda} \hat{w}_{\lambda/\nu}(yt^{-\ell}; q, t) \hat{w}_{\nu/\mu}(z; q, t)$$

for $y \in \mathbb{C}$ and $z \in \mathbb{C}^\ell$. We now derive some new basic properties in Corollary 2.1, and recall some existing properties in Remark 2.1 below for the w function and its dual.

Corollary 2.1. *Let μ be a partition of at most n parts, and $x = (x_1, \dots, x_n) \in \mathbb{C}^n$. Then*

(1) *The w_μ and its dual \hat{w}_μ are flipped versions of one another. That is,*

$$(17) \quad \hat{w}_\mu(x, q, t) = q^{-|\mu|} t^{-2n(\mu)+(n-1)|\mu|} w_\mu(1/x, 1/q, 1/t).$$

(2) *The limit $\lim_{q \rightarrow 1} w_\mu(xt^{\delta(n)}; q, t)$ exists when denominators do not vanish. For the particular case when $x = \lambda$ is a partition, we use the notation*

$$(18) \quad \bar{w}_\mu(1^\lambda t^{\delta(n)}; 1, t) := \lim_{q \rightarrow 1} (1 - q)^{-\mu_1} w_\mu(q^\lambda t^{\delta(n)}; q, t).$$

Proof. Both properties follow, by direct calculation, from the definition (13) of $w_{\lambda/\mu}$, the recurrence relation (14) for $w_{\lambda/\mu}$, and limit formula (12), and the flip formula

$$(19) \quad x^{|\mu|} (x^{-1}, q, t)_\mu = (-1)^{|\mu|} q^{n(\mu')} t^{-n(\mu)} (x; q^{-1}, t^{-1})_\mu.$$

The proof also uses the result that, in the limit

$$(20) \quad \lim_{q \rightarrow 1} H_{\lambda/\mu}(q, t) = \frac{(k)_m}{m!}$$

where k is the maximum of the list $k = \max\{\lambda_1 - \lambda_2, \lambda_2 - \lambda_3, \dots, \lambda_{n-1} - \lambda_n\}$, and m is the maximum of $m = \max\{\lambda_1 - \mu_1, \mu_1 - \lambda_2, \lambda_2 - \mu_2, \dots, \mu_{n-1} - \lambda_n\}$. If the second list has a negative number (which means that λ/μ is not a horizontal strip), then $H_{\lambda/\mu}(q, t) = 0$. \square

Remark 2.1. *We will need the following properties from [9] in what follows.*

(3) *If $z = xt^{\delta(n)}$, for some $x \in \mathbb{C}$, we get*

$$(21) \quad w_\mu(xt^{\delta(n)}; q, t) = (-1)^{|\mu|} x^{|\mu|} t^{n(\mu)} q^{-|\mu| - n(\mu')} (x^{-1})_\mu \prod_{1 \leq i < j \leq n} \frac{(t^{j-i+1})_{\mu_i - \mu_j}}{(t^{j-i})_{\mu_i - \mu_j}}$$

which, after flipping q and t and using the flip rule (19), may be written as

$$(22) \quad w_\mu(xt^{\delta(n)}; q, t) = q^{-|\mu|} (x; q^{-1}, t^{-1})_\mu \prod_{1 \leq i < j \leq n} \frac{(t^{j-i+1})_{\mu_i - \mu_j}}{(t^{j-i})_{\mu_i - \mu_j}}$$

(4) *The vanishing property of W functions implies that*

$$(23) \quad w_\mu(q^\lambda t^\delta; q, t) = 0$$

when $\mu \not\subseteq \lambda$, where \subseteq denotes the partial inclusion ordering.

(5) *Let λ be an n -part partition with $\lambda_n \neq 0$ and $0 \leq k \leq \lambda_n$ for some integer k , and let $z = (x_1, \dots, x_n) \in \mathbb{C}^n$. It was shown in [9] that*

$$(24) \quad w_{\bar{k}}(z; q, t) = q^{-nk} \prod_{i=1}^n (q^{1-k} x_i)_k$$

where $\bar{k} = (k, \dots, k)$ is an n -tuple with integer $k \geq 0$.

(6) *With the same notation as above, we also have*

$$(25) \quad w_\mu(q^\mu t^{\delta(n)}; q, t) = q^{-|\mu|} t^{(n-1)|\mu| - 2n(\mu)} (qt^{n-1})_\mu \prod_{1 \leq i < j \leq n} \frac{(qt^{j-i-1})_{\mu_i - \mu_j}}{(qt^{j-i})_{\mu_i - \mu_j}}.$$

2.3. The elliptic Jackson coefficients $\omega_{\lambda/\mu}$

We will also need the basic ($p = 0$) case of the elliptic Jackson coefficients $\omega_{\lambda/\mu}$ below. Let λ and μ be again partitions of at most n -parts such that λ/μ is a skew partition. Then the Jackson coefficients $\omega_{\lambda/\mu}$ are defined [7] by

$$\begin{aligned}
 (26) \quad \omega_{\lambda/\mu}(x; r, q, p, t; a, b) &:= \frac{(x^{-1}, ax)_{\lambda}}{(qbx, qb/ax)_{\lambda}} \frac{(qbr^{-1}x, qb/axr)_{\mu}}{(x^{-1}, ax)_{\mu}} \\
 &\cdot \frac{(r, br^{-1}t^{1-n})_{\mu}}{(qbr^{-2}, qt^{n-1})_{\mu}} \prod_{i=1}^n \left\{ \frac{\theta(br^{-1}t^{2-2i}q^{2\mu_i})}{\theta(br^{-1}t^{2-2i})} (qt^{2i-2})^{\mu_i} \right\} \\
 &\cdot \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\mu_i - \mu_j}}{(qt^{j-i-1})_{\mu_i - \mu_j}} \frac{(br^{-1}t^{3-i-j})_{\mu_i + \mu_j}}{(br^{-1}t^{2-i-j})_{\mu_i + \mu_j}} \right\} \\
 &\cdot W_{\mu}(q^{\lambda}t^{\delta(n)}; q, p, t, bt^{2-2n}, br^{-1}t^{1-n})
 \end{aligned}$$

where $x, r, q, p, t, a, b \in \mathbb{C}$, with an additional parameter r . Note that $W_{\lambda/\mu}$ vanishes unless λ/μ is a horizontal strip, whereas $\omega_{\lambda/\mu}$ is defined even when λ/μ is not a horizontal strip.

It is shown [7] that the multivariable $\omega_{\lambda/\mu}$ are symmetric functions constructed by the recursion formula

$$(27) \quad \omega_{\lambda/\mu}(y, z; r; a, b) := \sum_{\nu} \omega_{\lambda/\nu}(r^{-k}y; r; ar^{2k}, br^k) \omega_{\nu/\mu}(z; r; a, b)$$

where $y = (x_1, \dots, x_{n-k}) \in \mathbb{C}^{n-k}$ and $z = (x_{n-k+1}, \dots, x_n) \in \mathbb{C}^k$. Here and elsewhere we surpass the dimension parameters q, t, s for clarity of notation, and write $\omega_{\lambda/\mu}(x; r; a, b) = \omega_{\lambda/\mu}(x; r, q, p, t; a, b)$ unless otherwise specified. It is also shown [7] that the $\omega_{\lambda/\mu}$ are symmetric functions that satisfy the key result, called the cocycle identity

$$(28) \quad \omega_{\lambda/\mu}((sr)^{-1}; sr, as^2, bs) = \sum_{\nu} \omega_{\lambda/\nu}(s^{-1}; s, as^2, bs) \omega_{\nu/\mu}(r^{-1}; r, a, b)$$

where the summation index ν runs over all partitions.

2.4. The multiple qt -binomial coefficients

The multiple Stirling numbers we develop in this paper are closely connected with binomial coefficients just as in the one dimensional case. Recall that the multiple qt -binomial coefficient is defined [9] as

Definition 2.1. Let $z = (x_1, \dots, x_n) \in \mathbb{C}^n$ and μ be an n -part partition. Then the qt -binomial coefficient is defined by

$$(29) \quad \binom{z}{\mu}_{q,t} := \frac{q^{|\mu|} t^{2n(\mu)+(1-n)|\mu|}}{(qt^{n-1})_\mu} \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\mu_i - \mu_j}}{(qt^{j-i-1})_{\mu_i - \mu_j}} \right\} w_\mu(q^z t^{\delta(n)}; q, t)$$

where $q, t \in \mathbb{C}$. It should be noted that this definition makes sense even for $\mu \in \mathbb{C}^n$.

As pointed out in the Introduction, the qt -binomial coefficients are constructed independently by Kaneko [18] in a special case, and Okounkov [25] using difference operator methods for integer partitions $z = \lambda$ and μ . Lassalle constructed an equivalent set of qt -binomial coefficients independently [23], and Sahi has developed the non-symmetric version [28]. Rains [27] used Okounkov’s BC_n symmetric interpolation polynomials to define a three parameter qts -binomial coefficients which are equivalent to a limiting case of our $\omega_{\lambda/\mu}$ functions (26) up to rescaling.

Setting $t = q^\alpha$ and sending $q \rightarrow 1$ yields the multiple ordinary α -binomial coefficients. For $n = 1$, $z = (n)$ and $\mu = (k)$ for some integers m and k , the definition reduces to that of the one dimensional q -binomial coefficients

$$(30) \quad \binom{m}{k}_q := \frac{(q)_m}{(q)_{m-k}(q)_k}$$

which are also known as the Gaussian polynomials. These are studied extensively in the literature including but not limited to the works in [1, 2, 3, 5, 14, 15, 17, 21].

Using the definition above, we write [9] the terminating qt -binomial theorem in the form

$$(31) \quad (x)_\lambda = \sum_{\mu \subseteq \lambda} (-1)^{|\mu|} q^{n(\mu')} t^{-n(\mu)} \binom{\lambda}{\mu}_{q,t} x^{|\mu|}.$$

This special case and its nonsymmetric analogues are also proved independently in [25, 28], and studied in more recent works [22].

For the special case $z = \bar{x} = (x, \dots, x) \in \mathbb{C}^n$, we get

$$(32) \quad \binom{\bar{x}}{\mu}_{q,t} = \frac{t^{2n(\mu)+(1-n)|\mu|}}{(qt^{n-1})_\mu} (q^x; q^{-1}, t^{-1})_\mu \cdot \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\mu_i - \mu_j}}{(qt^{j-i-1})_{\mu_i - \mu_j}} \frac{(t^{j-i+1})_{\mu_i - \mu_j}}{(t^{j-i})_{\mu_i - \mu_j}} \right\}.$$

This is also established independently in [23].

Another special case occurs when μ is a rectangular partition, that is $\mu = \bar{k}$. Using (24) we get

$$(33) \quad \binom{z}{\bar{k}}_{q,t} = \prod_{i=1}^n \frac{(q^{1-k}q^{x_i}t^{n-i})_k}{(qt^{n-i})_k}$$

In the particular case for $k = 1$, the definition reduces to

$$(34) \quad \binom{z}{\bar{1}}_{q,t} = \prod_{i=1}^n \frac{(q^{x_i}t^{n-i})_1}{(qt^{n-i})_1} = \prod_{i=1}^n \frac{(1 - q^{x_i}t^{n-i})}{(1 - qt^{n-i})}$$

The last property we point out from [9] is that

$$(35) \quad \binom{\lambda}{\lambda}_{q,t} = \binom{\lambda}{\bar{0}}_{q,t} = 1$$

where λ is an n -part partition, and $\bar{0}$ is the n -part partition whose parts are all zero. In addition,

$$(36) \quad \binom{\lambda}{\mu}_{q,t} = 0$$

when $\mu \not\subseteq \lambda$ or $\bar{0} \not\subseteq \mu$.

2.5. The multiple qt -factorial function

We now recall another important extension [9] that generalizes the one dimensional q -bracket and q -factorial polynomial to the multiple case as follows.

Definition 2.2. Let μ be a partition of at most n parts, $z = (x_1, \dots, x_n) \in \mathbb{C}^n$ and $s \in \mathbb{C}^n$. Then

$$(37) \quad [z, s]_{\mu} = [z, s, n, q, t]_{\mu} := q^{|\mu|} \prod_{i=1}^n \left\{ \frac{1}{(1 - qt^{n-i})^{\mu_i}} \right\} \prod_{1 \leq i < j \leq n} \left\{ \frac{(t^{j-i})_{\mu_i - \mu_j}}{(t^{j-i+1})_{\mu_i - \mu_j}} \right\} w_{\mu}(sq^z t^{\delta^{(n)}}; q, t)$$

is called the qt -factorial (bracket) function. Note that the definition combines a multiplicative variable s , and an exponential variable z . Depending on the application we often set $z = \bar{0}$ and write $\langle s \rangle_{\mu} = [\bar{0}, s]_{\mu}$, or set $s = \bar{1}$ and write $[z]_{\mu} = [z, \bar{1}]_{\mu}$. Using the identity (24) in the special case when $\mu = \bar{1}$,

we define the qt -bracket as

$$(38) \quad [z] = [z, \bar{1}, n, q, t]_{\bar{1}} = \prod_{i=1}^n \frac{(1 - q^{x_i} t^{n-i})}{(1 - qt^{n-i})}$$

which is a multiple analogue of the classical q -number or q -bracket.

The definition (37) of $[z]_{\mu}$ may also be written as

$$(39) \quad [z]_{\mu} = t^{-2n(\mu) - (1-n)|\mu|} \prod_{i=1}^n \left\{ \frac{(qt^{n-i})_{\mu_i}}{(1 - qt^{n-i})^{\mu_i}} \right\} \cdot \prod_{1 \leq i < j \leq n} \left\{ \frac{(t^{j-i})_{\mu_i - \mu_j}}{(t^{j-i+1})_{\mu_i - \mu_j}} \frac{(qt^{j-i-1})_{\mu_i - \mu_j}}{(qt^{j-i})_{\mu_i - \mu_j}} \right\} \binom{z}{\mu}_{q,t}$$

which reduces to (38) when $\mu = \bar{1}$ by the identity (33).

Remark 2.2. The qt -factorial function satisfies the following properties:

(a) Let $z = (x, \dots, x) = \bar{x} \in \mathbb{C}^n$ for a single variable $x \in \mathbb{C}$, then the qt -factorial function $[\bar{x}]_{\mu}$ may be written as

$$(40) \quad [\bar{x}]_{\mu} = \prod_{i=1}^n \left\{ \frac{1}{(1 - qt^{n-i})^{\mu_i}} \right\} (q^x; q^{-1}, t^{-1})_{\mu} = \prod_{i=1}^n \left\{ \frac{(q^x t^{i-1}; q^{-1})_{\mu_i}}{(1 - qt^{n-i})^{\mu_i}} \right\}.$$

This definition reduces to the classical q -bracket in the one variable case.

(b) Note that $(x; 1/q, 1/t)_{\mu}$, with the reciprocals of q and t , corresponds to a multiple basic qt -analogue of the falling factorial $x_{\underline{n}} := x(x-1) \cdots (x-(n-1))$ as opposed to the rising factorial or the Pochhammer symbol.

(c) Setting $z = \mu$, and substituting the evaluation (25) in (37) above gives

$$(41) \quad [\mu]_{\mu} = t^{-2n(\mu) - (1-n)|\mu|} \prod_{i=1}^n \left\{ \frac{(qt^{n-i})_{\mu_i}}{(1 - qt^{n-i})^{\mu_i}} \right\} \cdot \prod_{1 \leq i < j \leq n} \left\{ \frac{(t^{j-i})_{\mu_i - \mu_j}}{(t^{j-i+1})_{\mu_i - \mu_j}} \frac{(qt^{j-i-1})_{\mu_i - \mu_j}}{(qt^{j-i})_{\mu_i - \mu_j}} \right\}$$

which is precisely the front factor in (39) as expected. Similar to the classical case, we may use the notation $\mu! = [\mu]_{\mu}$ and write

$$(42) \quad [z]_{\mu} = \mu! \binom{z}{\mu}_{q,t}.$$

Note that in the particular case when $\mu = \bar{k}$ is a rectangular partition, the $\mu!$ reduces to a product of one dimensional quotients for each part

$$(43) \quad \bar{k}! = \prod_{i=1}^n \left\{ \frac{(qt^{n-i})_k}{(1 - qt^{n-i})^k} \right\}.$$

(d) Recall that a multiple analogue $E_{q,t}(z)$ of the exponential function e^z was defined [8] by

$$(44) \quad E_{q,t}(z) := (-z)_{\infty^n} = \sum_{\mu \in P_n} \frac{z^{|\mu|} q^{n(\mu')} t^{n(\mu) + (1-n)|\mu|}}{(qt^{n-1})_{\mu}} \cdot \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\mu_i - \mu_j}}{(qt^{j-i-1})_{\mu_i - \mu_j}} \frac{(t^{j-i+1})_{\mu_i - \mu_j}}{(t^{j-i})_{\mu_i - \mu_j}} \right\}$$

where $z \in \mathbb{C}$, and P_n denotes the set of all partitions of length at most n , and $(z)_{\infty^n} = \prod_{i=1}^n (zt^{1-i})_{\infty}$. Using the $\mu!$ notation and setting $z = 1$ gives a multiple analogue of the number e :

$$(45) \quad E_{q,t}(1) := (-1)_{\infty^n} = \sum_{\mu \in P_n} \frac{q^{n(\mu')} t^{-n(\mu)}}{\prod_{i=1}^n (1 - qt^{n-i})^{\mu_i}} \cdot \frac{1}{\mu!}$$

similar to the classical case

$$e = \sum_{m=0}^{\infty} \frac{1}{m!}.$$

3. Recurrence relations for multiple factorial function

We treat the recurrence relations for the multiple factorial function in a separate section here. We derive three distinct recurrences for the multiple binomial coefficients, and for multiple factorials $[z]_{\lambda}$: A recurrence relation with respect to the variable z , another recurrence with respect to the index λ , and a third recurrence with respect to the weight $|\lambda|$ of the index partition.

3.1. The variable z

We can write the W -Jackson sum [8] in the form

$$(46) \quad W_{\lambda}(z; q, t, at^{-2n}, bt^{-n})$$

$$\begin{aligned}
 &= \frac{(s)_\lambda (as^{-1}t^{-n-1})_\lambda}{(qbs^{-1}t^{-1})_\lambda (qbt^n s/a)_\lambda} \cdot \prod_{1 \leq i < j \leq n} \left\{ \frac{(t^{j-i+1})_{\lambda_i - \lambda_j} (qbt^{-i-j+1})_{\lambda_i + \lambda_j}}{(t^{j-i})_{\lambda_i - \lambda_j} (qbt^{-i-j})_{\lambda_i + \lambda_j}} \right\} \\
 &\cdot \sum_{\mu \subseteq \lambda} \frac{(bs^{-1}t^{-n})_\mu (qbt^n/a)_\mu}{(qt^{n-1})_\mu (as^{-1}t^{-n-1})_\mu} \cdot \prod_{i=1}^n \left\{ \frac{(1 - bs^{-1}t^{1-2i}q^{2\mu_i})}{(1 - bs^{-1}t^{1-2i})} (qt^{2i-2})^{\mu_i} \right\} \\
 &\cdot \prod_{1 \leq i < j \leq n} \left\{ \frac{(t^{j-i})_{\mu_i - \mu_j} (qt^{j-i})_{\mu_i - \mu_j}}{(qt^{j-i-1})_{\mu_i - \mu_j} (t^{j-i+1})_{\mu_i - \mu_j}} \right. \\
 &\quad \left. \cdot \frac{(bs^{-1}qt^{-i-j})_{\mu_i + \mu_j} (bs^{-1}t^{-i-j+2})_{\mu_i + \mu_j}}{(bs^{-1}t^{-i-j+1})_{\mu_i + \mu_j} (qbs^{-1}t^{-i-j+1})_{\mu_i + \mu_j}} \right\} \\
 &\cdot W_\mu(q^\lambda t^{\delta(n)}; q, t, bt^{1-2n}, bs^{-1}t^{-n}) \cdot W_\mu(zs; q, t, as^{-2}t^{-2n}, bs^{-1}t^{-n})
 \end{aligned}$$

where $z \in \mathbb{C}^n$. This sum may be viewed as writing the W function on the left-hand side as a combination of the shifted W function on the right (shifted by a factor of $s \in \mathbb{C}$). Therefore, the left-hand side is independent of the parameter s .

If we set $b = ar$ in this identity and send $a \rightarrow 0$, multiply both sides by $(rt^n)^{|\lambda|}$ and send $r \rightarrow \infty$ using the limit rule (12), and we get

$$\begin{aligned}
 (47) \quad &w_\lambda(z; q, t) \\
 &= (-1)^{|\lambda|} q^{-|\lambda| - n(\lambda')} t^{n(\lambda)} s^{-|\lambda|} (s)_\lambda \cdot \prod_{1 \leq i < j \leq n} \left\{ \frac{(t^{j-i+1})_{\lambda_i - \lambda_j}}{(t^{j-i})_{\lambda_i - \lambda_j}} \right\} \\
 &\cdot \sum_{\mu \subseteq \lambda} \frac{(-1)^{|\mu|} q^{2|\mu|} t^{n(\mu)} q^{n(\mu')}}{(qt^{n-1})_\mu} \cdot \prod_{1 \leq i < j \leq n} \left\{ \frac{(t^{j-i})_{\mu_i - \mu_j} (qt^{j-i})_{\mu_i - \mu_j}}{(qt^{j-i-1})_{\mu_i - \mu_j} (t^{j-i+1})_{\mu_i - \mu_j}} \right\} \\
 &\cdot \left(\lim_{a \rightarrow 0} W_\mu(q^\lambda t^{\delta(n)}; q, t, at^{1-2n}, as^{-1}t^{-n}) \right) \cdot w_\mu(zs; q, t) s^{-|\mu|}.
 \end{aligned}$$

Shifting z by $s^{-1}z$ in (47) above, setting $s^{-1} = q^\alpha$ and $z = q^z t^{\delta(n)}$ and using the flip rule

$$(48) \quad x^{-|\mu|} (x; q, t)_\mu = (-1)^{|\mu|} q^{n(\mu')} t^{-n(\mu)} (x^{-1}; q^{-1}, t^{-1})_\mu$$

to simplify the front factors, and substituting the definition (29) into (47) gives

$$(49) \quad \binom{z + \alpha}{\lambda}_{q,t} = \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\lambda_i - \lambda_j} (t^{j-i+1})_{\lambda_i - \lambda_j}}{(qt^{j-i-1})_{\lambda_i - \lambda_j} (t^{j-i})_{\lambda_i - \lambda_j}} \right\}$$

$$\cdot \frac{t^{2n(\lambda)+(1-n)|\lambda|} (q^\alpha; q^{-1}, t^{-1})_\lambda}{(qt^{n-1})_\lambda} \cdot \sum_{\mu \subseteq \lambda} (-1)^{|\mu|} q^{(\alpha+1)|\mu|+n(\mu')} t^{-n(\mu)-(1-n)|\mu|}$$

$$\cdot \prod_{1 \leq i < j \leq n} \left\{ \frac{(t^{j-i})_{\mu_i - \mu_j}}{(t^{j-i-1})_{\mu_i - \mu_j}} \right\} \left(\lim_{a \rightarrow 0} W_\mu(q^\lambda t^{\delta(n)}; q, t, at^{1-2n}, aq^\alpha t^{-n}) \right) \binom{z}{\mu}_{q,t}.$$

This identity defines a general recurrence relation for the multiple binomial coefficients with respect to the z variable. Choosing $\alpha \in \mathbb{C}$ and $z \in \mathbb{C}^n$ properly yields interesting special cases. For an n partition λ , let λ^i denote $\lambda^i = \lambda + e^i$ for $e^i = (e_1^i, \dots, e_j^i, \dots, e_n^i)$ with $e_i^i = 1$, and $e_j^i = 0$ for $i \neq j$. Setting $z + \alpha = \lambda^i$ in (49), for example, gives a recurrence for

$$\binom{\lambda^i}{\lambda}_{q,t} = \binom{(\lambda_1, \dots, \lambda_{i-1}, \lambda_i + 1, \lambda_{i+1}, \dots, \lambda_n)}{(\lambda_1, \dots, \lambda_{i-1}, \lambda_i, \lambda_{i+1}, \dots, \lambda_n)}_{q,t}$$

for any $1 \leq i \leq n$.

If we further substitute (39) for the multiple binomial coefficients on both sides, we get a recurrence for the multiple bracket function again with respect to the z variable as follows.

$$(50) \quad \frac{\prod_{i=1}^n \{(1 - qt^{n-i})^{\lambda_i}\}}{(q^\alpha; q^{-1}, t^{-1})_\lambda} [z + \alpha]_\lambda$$

$$= \sum_{\mu \subseteq \lambda} \frac{(-1)^{|\mu|} q^{(\alpha+1)|\mu|} t^{n(\mu)} q^{n(\mu')}}{(qt^{n-1})_\mu} \cdot \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\mu_i - \mu_j}}{(qt^{j-i-1})_{\mu_i - \mu_j}} \right\}$$

$$\cdot \left(\lim_{a \rightarrow 0} W_\mu(q^\lambda t^{\delta(n)}; q, t, at^{1-2n}, aq^\alpha t^{-n}) \right) \prod_{i=1}^n \{(1 - qt^{n-i})^{\mu_i}\} [z]_\mu.$$

We may again specialize z and α to derive interesting special cases.

3.2. The index λ

We have shown [7] that for $0 < k \leq \lambda_n$

$$(51) \quad W_\lambda(z; q, t, a, b) = \prod_{1 \leq i < j \leq n} \frac{(qbt^{j-2i})_{2k}}{(qbt^{j-1-2i})_{2k}}$$

$$\cdot \prod_{i=1}^n \frac{(z_i^{-1})_k (az_i)_k}{(qbz_i)_k (qb/(az_i))_k} W_{\lambda - \bar{k}}(zq^{-k}; aq^{2k}, bq^{2k}, t, q)$$

where $\lambda - \bar{k}$ denotes $\lambda - k^n = (\lambda_1 - k, \dots, \lambda_n - k)$. We shift μ by \bar{k} , replace b by as , send $a \rightarrow 0$, multiply both sides by $s^{|\mu|+kn}$, and then send $s \rightarrow \infty$ in the last identity, and use the limit rule (12) and the definition (13) of w function to get

$$(52) \quad w_{\mu+\bar{k}}(z; q, t) = (-1)^{nk} q^{-nk-n\binom{k}{2}} \prod_{i=1}^n z_i^k (z_i^{-1})_k \cdot w_{\mu}(zq^{-k}; t, q).$$

We now expand the right-hand side of this identity using (47). Replace z by $s^{-1}z$, set $s = q^k$ and $z = q^z t^{\delta(n)}$, and substitute the definition (29) here. Then using the flip formula (48) and the calculation that $t^{2n(\lambda+\bar{k})+(1-n)|\lambda+\bar{k}|} = t^{2n(\lambda)+(1-n)|\lambda|}$, we write

$$(53) \quad \begin{aligned} \binom{z}{\lambda + \bar{k}}_{q,t} &= t^{2n(\lambda)+(1-n)|\lambda|} \prod_{i=1}^n (q^{z_i} t^{(n-i)}; q^{-1})_k \\ &\cdot \frac{(q^{-k}; q^{-1}, t^{-1})_{\lambda}}{(qt^{n-1})_{\lambda+\bar{k}}} \prod_{1 \leq i < j \leq n} \left\{ \frac{(t^{j-i+1})_{\lambda_i-\lambda_j}}{(t^{j-i})_{\lambda_i-\lambda_j}} \frac{(qt^{j-i})_{\lambda_i-\lambda_j}}{(qt^{j-i-1})_{\lambda_i-\lambda_j}} \right\} \\ &\cdot \sum_{\mu \subseteq \lambda} (-1)^{|\mu|} q^{(k+1)|\mu|+n(\mu')} t^{-n(\mu)-(1-n)|\mu|} \prod_{1 \leq i < j \leq n} \left\{ \frac{(t^{j-i})_{\mu_i-\mu_j}}{(t^{j-i+1})_{\mu_i-\mu_j}} \right\} \\ &\cdot \left(\lim_{a \rightarrow 0} W_{\mu}(q^{\lambda} t^{\delta(n)}; q, t, at^{1-2n}, aq^{-k}t^{-n}) \right) \binom{z}{\mu}_{q,t}. \end{aligned}$$

This is a general recurrence relation for the multiple binomial with respect to the index partition μ . If we replace the binomials by the corresponding bracket functions using (39), we get

$$(54) \quad \begin{aligned} [z]_{\lambda+\bar{k}} &= \prod_{i=1}^n \left\{ \frac{(q^{z_i} t^{(n-i)}; q^{-1})_k}{(1 - qt^{n-i})^k} \right\} \cdot \frac{(q^{-k}; q^{-1}, t^{-1})_{\lambda}}{\prod_{i=1}^n (1 - qt^{n-i})^{\lambda_i}} \\ &\cdot \sum_{\mu \subseteq \lambda} (-1)^{|\mu|} q^{(k+1)|\mu|+n(\mu')} t^{n(\mu)} \left(\lim_{a \rightarrow 0} W_{\mu}(q^{\lambda} t^{\delta(n)}; q, t, at^{1-2n}, aq^{-k}t^{-n}) \right) \\ &\cdot \prod_{i=1}^n \left\{ \frac{(1 - qt^{n-i})^{\mu_i}}{(qt^{n-i})^{\mu_i}} \right\} \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\mu_i-\mu_j}}{(qt^{j-i-1})_{\mu_i-\mu_j}} \right\} [z]_{\mu}. \end{aligned}$$

Specializing the variables z and the integer k gives interesting special cases as before.

3.3. The weight $|\lambda|$

Recall that a recurrence relation for the qt -binomial coefficients based on the weights of the indexing partitions is written [8] as follows: For an n -part partition λ and $k \leq |\lambda|$, we have

$$(55) \quad \sum_{\mu \vdash k} q^{n(\mu')} t^{-n(\mu)} \binom{\lambda^i}{\mu}_{qt} \\ = \sum_{\tau \vdash k} q^{n(\tau')} t^{-n(\tau)} \binom{\lambda}{\tau}_{qt} + \sum_{\nu \vdash (k-1)} q^{n(\nu')+\lambda_i} t^{-n(\nu)+1-i} \binom{\lambda}{\nu}_{qt}$$

where $\mu \subseteq \lambda^i = (\lambda_1, \dots, \lambda_i + 1, \dots, \lambda_n)$ when it is a partition, and $\tau, \nu \subseteq \lambda$. Here, $\mu \vdash k$ denotes that μ is a partition of the integer $k \geq 1$. Using (42), we write a recurrence formula for the qt -factorial function $[\lambda]_\mu$ in the following form:

$$(56) \quad \sum_{\mu \vdash k} q^{n(\mu')} t^{-n(\mu)} \mu! [\lambda^i]_\mu \\ = \sum_{\tau \vdash k} q^{n(\tau')} t^{-n(\tau)} \tau! [\lambda]_\tau + \sum_{\nu \vdash (k-1)} q^{n(\nu')+\lambda_i} t^{-n(\nu)+1-i} \mu! [\lambda]_\mu.$$

4. Summation formulas for multiple factorial functions

The cocycle identity (28) for the Jackson coefficients $\omega_{\lambda/\mu}$ may be written in explicit form as

$$(57) \quad \frac{(rs)_\lambda (asr^{-1})_\lambda}{(qbr^{-1})_\lambda (qbr/a)_\lambda} \frac{(qb)_\lambda (qb/a)_\lambda}{(s)_\lambda (as)_\lambda} \frac{(qb/as)_\mu (ar^{-1})_\mu}{(asr^{-1})_\mu (qb/a)_\mu} \\ \cdot W_\mu(t, bst^{2-2n}, br^{-1}t^{1-n}; q^\lambda t^{\delta(n)}; q) \\ = \sum_\nu q^{|\nu|} t^{2n(\nu)} \frac{(qb/as)_\nu (bt^{1-n})_\nu}{(as)_\nu (qt^{n-1})_\nu} \frac{(r)_\nu (ar^{-1})_\nu}{(qbr^{-1})_\nu (qbr/a)_\nu} \\ \cdot \prod_{i=1}^n \left\{ \frac{(1 - bt^{2-2i} q^{2\nu_i})}{(1 - bt^{2-2i})} \right\} \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\nu_i - \nu_j} (bt^{3-i-j})_{\nu_i + \nu_j}}{(qt^{j-i-1})_{\nu_i - \nu_j} (bt^{2-i-j})_{\nu_i + \nu_j}} \right\} \\ \cdot W_\nu(t, bst^{2-2n}, bt^{1-n}; q^\lambda t^{\delta(n)}; q) W_\mu(t, bt^{2-2n}, br^{-1}t^{1-n}; q^\nu t^{\delta(n)}; q).$$

A specialization of this result gave the elegant identity for qt -binomial coefficients: For n -part partitions ν and μ , we have [9]

$$(58) \quad t^{-n(\mu)} q^{n(\mu')} \frac{(-1)_\nu}{(-1)_\mu} \binom{\nu}{\mu}_{q,t} = \sum_{\mu \subseteq \lambda \subseteq \nu} t^{-n(\lambda)} q^{n(\lambda')} \binom{\nu}{\lambda}_{q,t} \binom{\lambda}{\mu}_{q,t}.$$

Setting $\mu = \bar{1}$ here gives

$$(59) \quad t^{-n(\mu)} q^{n(\mu')} \frac{(-1)_\nu}{(-1)_\mu} [\nu]_{q,t} = \sum_{\mu \subseteq \lambda \subseteq \nu} t^{-n(\lambda)} q^{n(\lambda')} \binom{\nu}{\lambda}_{q,t} [\lambda]_{q,t}$$

which is a multiple analogue of the identity

$$(60) \quad 2^{n-1} n = \sum_{k=1}^n \binom{n}{k} k$$

for the qt -bracket function (37).

We like to also write a multiple analogue of the identity that expresses the sum of first n integers as

$$(61) \quad \sum_{i=1}^n k = \binom{n+1}{2} = \frac{n(n+1)}{2}$$

for the multiple bracket. One dimensional q -analogues of this identity appeared in several papers [12, 31, 29] recently. For the multiple analogue, we first recall another special case of the cocycle identity written [8] in the form

$$(62) \quad \frac{(zs)_\nu}{(s)_\nu} \frac{(z)_\mu}{(zs)_\mu} s^{|\mu|} w_\mu(q^\nu t^{\delta(n)}; q, t) \\ = \sum_{\mu \subseteq \lambda \subseteq \nu} q^{|\lambda|} t^{2n(\lambda)} \frac{(z)_\lambda}{(qt^{n-1})_{\lambda_1}} \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\lambda_i - \lambda_j}}{(qt^{j-i-1})_{\lambda_i - \lambda_j}} \right\} \\ \cdot \left(\lim_{a \rightarrow 0} W_\lambda(q^\nu t^{\delta(n)}; q, t, at^{1-2n}, as^{-1}t^{-n}) \right) w_\mu(q^\lambda t^{\delta(n)}; q, t).$$

It follows from the recurrence (11) that the W function in the sum simplifies to

$$(63) \quad \left(\lim_{a \rightarrow 0} W_\lambda(q^\nu t^{\delta(n)}; q, t, at^{1-2n}, aq^{-1}t^{1-2n}) \right) = \prod_{1 \leq i < j \leq n} \frac{(t^{j-i+1})_{\lambda_i - \lambda_j}}{(t^{j-i})_{\lambda_i - \lambda_j}}$$

in the special case when $s = qt^{n-1}$. Note that this specialization removed the W function that terminated the sum from above. Using the definition of multiple bracket function (37) on the right and multiple binomial (29) on

the left, we may write

$$\begin{aligned}
 (64) \quad & t^{-2n(\mu)-2(1-n)|\mu|} (z)_\mu \frac{(zqt^{n-1})_\nu}{(qt^{n-1})_\nu} \frac{(qt^{n-1})_\mu}{(zqt^{n-1})_\mu} \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i-1})_{\mu_i-\mu_j}}{(qt^{j-i})_{\mu_i-\mu_j}} \right\} \binom{\nu}{\mu}_{q,t} \\
 &= \sum_{\mu \subseteq \lambda \subseteq \nu} \frac{(z)_\lambda}{(qt^{n-1})_\lambda} q^{|\lambda|} t^{2n(\lambda)} \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\lambda_i-\lambda_j}}{(qt^{j-i-1})_{\lambda_i-\lambda_j}} \frac{(t^{j-i+1})_{\lambda_i-\lambda_j}}{(t^{j-i})_{\lambda_i-\lambda_j}} \right\} \\
 &\quad \cdot q^{-|\mu|} \prod_{i=1}^n \{(1-qt^{n-i})^{\mu_i}\} \prod_{1 \leq i < j \leq n} \left\{ \frac{(t^{j-i+1})_{\mu_i-\mu_j}}{(t^{j-i})_{\mu_i-\mu_j}} \right\} [\lambda]_\mu.
 \end{aligned}$$

Setting $z = qt^{n-1}$ further removes the factor that terminates the sum from below as well, but we still get a finite sum

$$\begin{aligned}
 (65) \quad & q^{|\mu|} t^{-2n(\mu)-2(1-n)|\mu|} \frac{(q^2t^{2(n-1)})_\nu}{(qt^{n-1})_\nu} \frac{(qt^{n-1})_\mu^2}{(q^2t^{2(n-1)})_\mu} \prod_{i=1}^n \left\{ \frac{1}{(1-qt^{n-i})^{\mu_i}} \right\} \\
 &\quad \cdot \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i-1})_{\mu_i-\mu_j}}{(qt^{j-i})_{\mu_i-\mu_j}} \frac{(t^{j-i})_{\mu_i-\mu_j}}{(t^{j-i+1})_{\mu_i-\mu_j}} \right\} \binom{\nu}{\mu}_{q,t} \\
 &= \sum_{\mu \subseteq \lambda \subseteq \nu} q^{|\lambda|} t^{2n(\lambda)} \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\lambda_i-\lambda_j}}{(qt^{j-i-1})_{\lambda_i-\lambda_j}} \frac{(t^{j-i+1})_{\lambda_i-\lambda_j}}{(t^{j-i})_{\lambda_i-\lambda_j}} \right\} [\lambda]_\mu.
 \end{aligned}$$

In particular for the special case $\mu = \bar{1}$, we get a multiple analogue of (61) as

$$\begin{aligned}
 (66) \quad & q^n t^{(n-1)n} \frac{(q^2t^{2(n-1)})_\nu}{(qt^{n-1})_\nu} \prod_{i=1}^n \frac{(1-q^{\nu_i}t^{n-i})}{(1-q^2t^{2n-1-i})} \\
 &= \sum_{\bar{1} \subseteq \lambda \subseteq \nu} q^{|\lambda|} t^{2n(\lambda)} \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\lambda_i-\lambda_j}}{(qt^{j-i-1})_{\lambda_i-\lambda_j}} \frac{(t^{j-i+1})_{\lambda_i-\lambda_j}}{(t^{j-i})_{\lambda_i-\lambda_j}} \right\} [\lambda].
 \end{aligned}$$

Note that other specializations of z in (64) above, such as $z = 0$, gives different versions of this multiple identity. Using similar methods and the recurrences for multiple factorial functions, one may also write multiple analogues of the sums of powers of brackets as well.

5. Multiple basic and ordinary qt -Stirling numbers

In this section we review the definition and fundamental properties of the multiple Stirling numbers of the first and second kind indexed by partitions

[9]. The classical Stirling numbers of the first kind are defined to be the coefficients of the power functions x^k in the expansion of the falling factorial $x_{\underline{n}} = x(x - 1) \dots (x - n + 1)$ written as

$$(67) \quad x_{\underline{n}} = n! \binom{x}{n} = \sum_{k=0}^n s_1(n, k) x^k.$$

The classical Stirling numbers of the second kind are defined the other way around as the coefficients of the falling factorials in the expansion of the power functions. The q -analogue of these numbers are defined in [4] and their properties are studied in [24, 16, 19, 33] and the works of others.

First, we recall [9] the definition of the multiple qt -Stirling numbers generalizing the one dimensional q -analogues.

Definition 5.1. For an n part partition λ , the qt -Stirling numbers of the first kind $s_1(\lambda, \mu)$ are defined by

$$(68) \quad [\bar{x}]_{\lambda} = \sum_{\mu \subseteq \lambda} q^{-n(\lambda')} t^{2n(\mu) - (n-1)|\mu|} s_1(\lambda, \mu) \prod_{i=1}^{\mu_1} [\bar{x}^i]$$

where $x \in \mathbb{C}$, $\bar{x} = \{x, \dots, x\} \in \mathbb{C}^n$, and $\bar{x}^i = \{x, \dots, x, q, \dots, q\} \in \mathbb{C}^n$ with μ'_i copies of x for the dual partition μ' . That is,

$$(69) \quad \prod_{i=1}^{\mu_1} [\bar{x}^i] = \prod_{i=1}^n \frac{(1 - q^x t^{n-i})^{\mu_i}}{(1 - qt^{n-i})^{\mu_i}}.$$

Likewise, the qt -Stirling numbers of the second kind $s_2(\lambda, \mu)$ are defined by

$$(70) \quad \prod_{i=1}^n \frac{(1 - q^x t^{n-i})^{\lambda_i}}{(1 - qt^{n-i})^{\lambda_i}} = \prod_{i=1}^{\lambda_1} [\bar{x}^i] = \sum_{\mu \subseteq \lambda} q^{n(\mu')} t^{-2n(\nu) + (n-1)|\nu|} s_2(\lambda, \mu) [\bar{x}]_{\mu}.$$

Note that both the right and left hand sides are bases for the space. So, existence of the uniquely defined coefficients makes the multiple qt -Stirling numbers well defined.

We now recall the explicit formula for the qt -Stirling numbers given in [9], generalizing the one dimensional q -analogues. We will refer to certain identities derived in the proof several times later on. In addition, there are some notational changes adopted in this paper. Thus, we include the main steps of the proof here as well.

Theorem 5.1. For n -part partitions ν and μ , an explicit formula for the qt -Stirling numbers of first and second kind $s_1(\lambda, \mu)$ and $s_2(\lambda, \mu)$ are given by

$$(71) \quad s_1(\nu, \mu) = s_1(\nu, \mu, q, t) = \frac{q^{n(\nu')}t^{-2n(\mu)+2(n-1)|\mu|}}{\prod_{i=1}^n (1 - qt^{n-i})^{\nu_i - \mu_i}} f_\mu(q, t) \cdot \sum_{\mu \subseteq \lambda \subseteq \nu} u(\nu, \lambda, q, t) t^{(1-n)|\lambda|} \bar{w}_\mu(q^{-\lambda}t^{-\delta(n)}; 1, 1/t)$$

and

$$(72) \quad s_2(\nu, \mu) = s_2(\nu, \mu, q, t) := \frac{q^{-n(\mu')}t^{2n(\nu)+(1-n)|\nu|}}{\prod_{i=1}^n (1 - qt^{n-i})^{\nu_i - \mu_i}} \cdot \sum_{\mu \subseteq \lambda \subseteq \nu} (-1)^{|\lambda|} t^{n(\lambda)} \bar{w}_\lambda(q^\nu t^{\delta(n)}; 1, t) f_\lambda(q, t) v(\lambda, \mu, q, t)$$

where \hat{w}_μ and \bar{w}_μ are as defined above in Corollary 2.1, and $f(\mu)$, $u(\lambda, \mu)$ and $v(\lambda, \mu)$ are given by

$$(73) \quad f(\mu, q, t) := (1 - t)^{\mu_1 - \mu_n} \prod_{i=1}^{n-1} \frac{1}{(1 - t^{n-i})^{\mu_i}} \cdot \prod_{\substack{1 \leq i < j \leq n \\ j \neq i+1}} \left(\frac{1 - t^{j-i}}{1 - t^{j-i-1}} \right)^{\mu_i - \mu_j} \frac{1}{\mu_n!} \prod_{i=1}^{n-1} \frac{1}{(\mu_i - \mu_{i+1})!}$$

and

$$(74) \quad u(\lambda, \mu, q, t) := \frac{q^{|\mu|}t^{2n(\mu)}}{(qt^{n-1})_\mu} \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\mu_i - \mu_j}}{(qt^{j-i-1})_{\mu_i - \mu_j}} \right\} \hat{w}_\mu(q^\lambda t^{\delta(n)}; q, t)$$

and

$$(75) \quad v(\lambda, \mu, q, t) := (-1)^{|\mu|} q^{n(\mu')} t^{-n(\mu)} \binom{\lambda}{\mu}_{q,t} = \frac{(-1)^{|\mu|} q^{|\mu| + n(\mu')} t^{n(\mu) + (1-n)|\mu|}}{(qt^{n-1})_\mu} \cdot \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\mu_i - \mu_j}}{(qt^{j-i-1})_{\mu_i - \mu_j}} \right\} w_\mu(q^\lambda t^{\delta(n)}; q, t).$$

Proof. Simplify the identity (47) for the case $z = xt^{\delta(n)}$ using (22), and send $s \rightarrow \infty$ to get

$$(76) \quad (x; q^{-1}, t^{-1})_\lambda = \sum_{\mu \subseteq \lambda} u(\lambda, \mu) x^{|\mu|}$$

with the definition of $u(\lambda, \mu)$ above. Similarly, apply the shifts $a \rightarrow as^2$, $b \rightarrow bs$ and $x \rightarrow x/s$ in (46) at the beginning, and follow the same steps except send s to 0 to get

$$(77) \quad x^{|\lambda|} = \sum_{\mu \subseteq \lambda} v(\lambda, \mu) (x; q^{-1}, t^{-1})_\mu$$

where $v(\lambda, \mu)$ is defined as in the theorem. It is clear, by a change of basis argument, that

$$(78) \quad \delta_{\nu\lambda} = \sum_{\mu \subseteq \lambda \subseteq \nu} u(\nu, \lambda) v(\lambda, \mu).$$

Note that the left hand side of (77) does not depend on q or t . Now, flip the parameters $q \rightarrow 1/q, t \rightarrow 1/t$, take limit $q \rightarrow 1$, and multiply and divide the summand by $\prod_{i=1}^n 1/(1 - qt^{n-i})^{\mu_i}$ to get

$$(79) \quad x^{|\lambda|} = \sum_{\mu \subseteq \lambda} \prod_{i=1}^n (1 - qt^{n-i})^{\mu_i} \lim_{q \rightarrow 1} v(\lambda, \mu, 1/q, 1/t) \prod_{i=1}^{\mu_1} \langle \bar{x}^i \rangle$$

where $\langle \bar{x}^i \rangle$ defined as in (69) by replacing q^x by x . That is,

$$\prod_{i=1}^{\mu_1} \langle \bar{x}^i \rangle = \prod_{i=1}^n \frac{(1 - xt^{n-i})^{\mu_i}}{(1 - qt^{n-i})^{\mu_i}}.$$

Multiply and divide the summand in (76) by $t^{(n-1)|\mu|}$, substitute (79) into (76) for $(xt^{n-1})^{|\mu|}$, multiply both sides of this latter identity by $\prod_{i=1}^n 1/(1 - qt^{n-i})^{\nu_i}$ to get

$$(80) \quad \prod_{i=1}^n \left\{ \frac{1}{(1 - qt^{n-i})^{\nu_i}} \right\} (x; q^{-1}, t^{-1})_\nu = \sum_{\mu \subseteq \nu} \left(\prod_{i=1}^n \frac{(1 - qt^{n-i})^{\mu_i}}{(1 - qt^{n-i})^{\nu_i}} \right) \cdot \sum_{\mu \subseteq \lambda \subseteq \nu} u(\nu, \lambda, q, t) t^{-(n-1)|\lambda|} \left(\lim_{q \rightarrow 1} v(\lambda, \mu, 1/q, 1/t) \right) \prod_{i=1}^{\mu_1} \langle \bar{x}^i \rangle.$$

Multiplying and dividing the summand now by $q^{n(\nu')}t^{-2n(\mu)+(n-1)|\mu|}$ gives

$$(81) \quad \langle \bar{x} \rangle_\nu = \sum_{\mu \subseteq \nu} q^{-n(\nu')}t^{2n(\mu)-(n-1)|\mu|} s_1(\nu, \mu) \cdot \prod_{i=1}^n \frac{(1 - xt^{n-i})^{\mu_i}}{(1 - qt^{n-i})^{\mu_i}}$$

where $s_1(\nu, \mu)$ is as defined in the theorem. A similar sequence of calculations show that

$$(82) \quad \prod_{i=1}^n \frac{(1 - xt^{n-i})^{\nu_i}}{(1 - qt^{n-i})^{\nu_i}} = \sum_{\mu \subseteq \nu} \left(\prod_{i=1}^n \frac{(1 - qt^{n-i})^{\mu_i}}{(1 - qt^{n-i})^{\nu_i}} \right) \cdot \sum_{\mu \subseteq \lambda \subseteq \nu} \left(\lim_{q \rightarrow 1} u(\nu, \lambda, 1/q, 1/t) \right) t^{(n-1)|\lambda|} v(\lambda, \mu, q, t) \langle x \rangle_\mu.$$

Multiplying and dividing the summand now by $q^{-n(\mu')}t^{2n(\nu)+(1-n)|\nu|}$ gives the explicit formula for the qt -Stirling numbers of the second kind

$$(83) \quad \prod_{i=1}^n \frac{(1 - xt^{n-i})^{\nu_i}}{(1 - qt^{n-i})^{\nu_i}} = \sum_{\mu \subseteq \nu} q^{n(\mu')}t^{-2n(\nu)+(n-1)|\nu|} s_2(\nu, \mu) \cdot \langle \bar{x} \rangle_\mu.$$

Finally, substituting $x \rightarrow q^x$ completes the proof. We conclude by simplifying the flips and limits that entered the formulas above.

It follows immediately from (19) that, if

$$(84) \quad h(\mu, q, t) := \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})^{\mu_i - \mu_j}}{(qt^{j-i-1})^{\mu_i - \mu_j}} \right\} \text{ and } g(\mu, q, t) := (qt^{n-1})_\mu$$

then

$$(85) \quad h(\mu, 1/q, 1/t) = t^{2n(\mu)-(n-1)|\mu|} h(\mu, q, t)$$

and

$$(86) \quad g(\mu, 1/q, 1/t) = (-1)^{|\mu|} q^{-|\mu|-n(\mu')} t^{n(\mu)-(n-1)|\mu|} g(\mu, q, t).$$

Thus, flipping the parameters give

$$(87) \quad u(\lambda, \mu, 1/q, 1/t) = \frac{(-1)^{|\mu|} q^{|\mu|+n(\mu')} t^{n(\mu)-(n-1)|\mu|}}{(qt^{n-1})_\mu}$$

$$\cdot \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\mu_i - \mu_j}}{(qt^{j-i-1})_{\mu_i - \mu_j}} \right\} w_\mu(q^\lambda t^{\delta(n)}; q, t)$$

and

$$(88) \quad v(\lambda, \mu, 1/q, 1/t) = \frac{t^{(n-1)|\mu|}}{(qt^{n-1})_\mu} \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\mu_i - \mu_j}}{(qt^{j-i-1})_{\mu_i - \mu_j}} \right\} \cdot w_\mu(q^{-\lambda} t^{-\delta(n)}; 1/q, 1/t).$$

Using the limit rule (12), direct calculations give that

$$(89) \quad \lim_{q \rightarrow 1} (1 - q)^{\mu_1} \frac{1}{(qt^{n-1})_\mu} \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\mu_i - \mu_j}}{(qt^{j-i-1})_{\mu_i - \mu_j}} \right\} = f(\mu, q, t)$$

where the $f(\mu, q, t)$ factor was defined above. Therefore, multiplying and dividing the identities (87) and (88) by $(1 - q)^{\mu_1}$, and passing the limit as $q \rightarrow 1$ gives

$$(90) \quad \lim_{q \rightarrow 1} u(\lambda, \mu, 1/q, 1/t) = (-1)^{|\mu|} t^{n(\mu) - (n-1)|\mu|} \bar{w}_\mu(1^\lambda t^{\delta(n)}; 1, t) f(\mu, q, t).$$

where \bar{w}_μ is defined as above. Similarly,

$$(91) \quad \lim_{q \rightarrow 1} v(\lambda, \mu, 1/q, 1/t) = t^{(n-1)|\mu|} \bar{w}_\mu(1^{-\lambda} t^{-\delta(n)}; 1, 1/t) f(\mu, q, t)$$

which completes the proof. □

Remark 5.1. *It is worthwhile to list certain immediate properties of the multiple Stirling numbers of both kinds, some of which were already pointed out briefly in [9].*

(a) *Let P_n denote the set of all partitions with at most n parts. Note that by the vanishing property (23) of w_μ functions, both $u(\lambda, \mu, q, t)$ and $v(\lambda, \mu, q, t)$ coefficients vanish when $\mu \not\subseteq \lambda$ for all partitions $\mu, \lambda \in P_n$. This implies by the definitions (71) and (72) that the vanishing property*

$$s_1(\nu, \mu) = s_2(\nu, \mu) = 0$$

also holds for the multiple Stirling numbers when $\mu \not\subseteq \nu$. This result generalizes the one dimensional property that $s_2(n, m) = s_2(n, m) = 0$ for $m \geq n$.

(b) *The identities (76) and (77) show that the $(x; q^{-1}, t^{-1})_\mu$ form another basis for the space of polynomials with coefficients from the field $\mathbb{C}(q, t)$ of ratio-*

nal functions in parameters q and t . Therefore, the identities (81) and (83) indicate that the multiple Stirling numbers $s_1(\nu, \mu)$ and $s_2(\nu, \mu)$ are well defined as the unique connection coefficients between the two bases

$$(92) \quad t^{2n(\mu)-(n-1)|\mu|} \prod_{i=1}^n \frac{(1 - xt^{n-i})^{\mu_i}}{(1 - qt^{n-i})^{\mu_i}} \quad \text{and} \quad q^{n(\mu')} \langle \bar{x} \rangle_\mu$$

where

$$(93) \quad \langle \bar{x} \rangle_\mu = \prod_{i=1}^n \left\{ \frac{1}{(1 - qt^{n-i})^{\mu_i}} \right\} (x; q^{-1}, t^{-1})_\mu = \prod_{i=1}^n \left\{ \frac{(xt^{i-1}; q^{-1})_{\mu_i}}{(1 - qt^{n-i})^{\mu_i}} \right\}$$

for all $\mu \in P_n$.

(c) These explicit formulas in one dimensional case ($n = 1$) reduce essentially to those for Carlitz's q -Stirling numbers [4, 19]. Moreover, sending $q \rightarrow 1$ in that case yields the sequence of classical Stirling numbers for both types. The one dimensional form of the identities (81) and (83), for example, may be written in the form

$$(94) \quad \frac{(x; q^{-1})_v}{(1 - q)^v} = \sum_{m=0}^v q^{-\binom{v}{2}} s_1(v, m) \frac{(1 - x)^m}{(1 - q)^m}$$

and

$$(95) \quad \frac{(1 - x)^v}{(1 - q)^v} = \sum_{i=0}^v q^{\binom{m}{2}} s_2(v, m) \frac{(x; q^{-1})_m}{(1 - q)^m}$$

where

$$(96) \quad s_1(v, m) = s_1(v, m, q, t) = \frac{q^{\binom{v}{2}}}{(1 - q)^{v-m}} \sum_{j=m}^v \frac{(-1)^m j_m q^j (q^{-v})_j}{m! (q)_j}$$

and

$$(97) \quad s_2(v, m) = s_2(v, m, q, t) = \frac{q^{-\binom{m}{2}}}{(1 - q)^{v-m}} \sum_{j=m}^v \frac{(-1)^j v_j q^{jm} (q^{-j})_m}{j! (q)_m}$$

where $x_n = x(x - 1) \cdots (x - (n - 1))$ denotes the falling factorial as before. Setting $x \rightarrow q^x$ and sending $q \rightarrow 1$ in these results clearly gives the classical Stirling numbers (67) of both types.

(d) The matrix m with entries $m_{\lambda\mu} = s_1(\lambda, \mu)$ is invertible in the sense of V algebra defined at the beginning of Section 2, and its inverse is given by $m_{\lambda\mu}^{-1} = s_2(\lambda, \mu)$. More precisely, we have

$$(98) \quad \delta_{\nu\lambda} = \sum_{\mu \subseteq \lambda \subseteq \nu} s_1(\nu, \lambda) s_2(\lambda, \mu) = \sum_{\mu \subseteq \lambda \subseteq \nu} s_2(\nu, \lambda) s_1(\lambda, \mu)$$

which follows immediately from the inversion relation (78).

(e) Similar to the one dimensional case for the q -Stirling numbers, we have

$$(99) \quad s_1(\lambda, \lambda) = s_2(\lambda, \lambda) = 1$$

for an arbitrary n -part partition λ .

(f) Setting $t = q^\alpha$ and sending $q \rightarrow 1$ gives the multiple ordinary α -Stirling numbers of the first and second kind.

6. Summation identities for multiple Stirling numbers

We derive some additional new properties of the multiple Stirling numbers in this section.

(1) First note that $\lim_{x \rightarrow 0} \langle x \rangle_\mu = \prod_{i=1}^n 1/(1 - qt^{n-i})^{\mu_i}$ follows readily from the formula (40) written in the qt -angle bracket function $\langle \bar{x} \rangle_\mu$ above in (93). In the multiple case, setting $x = 0$ in (81) and (83) respectively gives

$$(100) \quad \prod_{i=1}^n \frac{1}{(1 - qt^{n-i})^{\nu_i}} = \sum_{\mu \subseteq \nu} q^{-n(\nu')} t^{2n(\mu) + (n-1)|\mu|} s_1(\nu, \mu) \prod_{i=1}^n \frac{1}{(1 - qt^{n-i})^{\mu_i}}$$

and

$$(101) \quad \prod_{i=1}^n \frac{1}{(1 - qt^{n-i})^{\nu_i}} = \sum_{\mu \subseteq \nu} q^{n(\mu')} t^{-2n(\nu) + (n-1)|\nu|} s_2(\nu, \mu) \prod_{i=1}^n \frac{1}{(1 - qt^{n-i})^{\mu_i}}.$$

The identities may be interpreted as giving the eigenvectors of certain operators. They appear to be new identities, even in the one dimensional case for $n = 1$ which may be obtained easily from (94) and (95) above.

(2) Note that, for an n -part partition ν with $\nu_n \neq 0$, the bracket function $\langle x \rangle_\nu$ has roots at $x = t^{1-j} q^{m_j}$ for $j = 1, \dots, n$, and $m_j = 0, \dots, \nu_j - 1$. The limit bracket function $\prod_{i=1}^n (1 - xt^{n-i})^{\mu_i} / (1 - qt^{n-i})^{\mu_i}$ has roots at $x = t^{1-j}$

for $j = 1, \dots, n$. Therefore, if we set $x = t^{1-j}q^{m_j}$ (for some $m_j < \nu_j$) in (83) we get

$$(102) \quad 0 = \sum_{\mu \subsetneq \nu} q^{-n(\nu')} t^{2n(\mu)-(n-1)|\mu|} \prod_{i=1}^n \frac{(1 - q^{m_j} t^{1-j+n-i})^{\mu_i}}{(1 - qt^{n-i})^{\mu_i}} s_1(\nu, \mu)$$

where the summation is over all partitions $\mu \subsetneq \nu$, that is all partitions $\mu \subseteq \nu$ such that $\mu_j \leq m_j$. This inequality follows from the vanishing property of the w functions (23).

In the particular case, setting $x = q$ in (81) gives

$$(103) \quad 0 = \sum_{\mu \subseteq \nu} q^{-n(\nu')} t^{2n(\mu)-(n-1)|\mu|} s_1(\lambda, \mu)$$

which is an analogue of $\sum_{k=0}^n s_1(n, k) = 0$ in the classical case.

In another special case, setting $x = t^{1-j}$ (i.e., $m_j = 0$) in (81) and (83) would amount to vanishing of all factorial functions $\langle x \rangle_\mu$ except the ones in whose index the j -th part (thus all parts $j + 1, \dots, n$ after j) are 0. That is, the factorial functions will be nonzero only for partitions such as $\mu = (\mu_1, \dots, \mu_{j-1}, 0, \dots, 0)$, and others will vanish. This is particularly interesting, for the substitution $x = t^{1-j}$ the limit factorial functions $\prod_{i=1}^n (1 - xt^{n-i})^{\mu_i} / (1 - qt^{n-i})^{\mu_i}$ also vanish except for partitions $\mu = (\mu_1, \dots, \mu_{n-j}, 0, \dots, 0)$.

Therefore, setting $x = t^{1-j}$ in (81) and (83) respectively gives

$$(104) \quad 0 = \sum_{\mu \subsetneq \nu} q^{-n(\nu')} t^{2n(\mu)-(n-1)|\mu|} s_1(\nu, \mu) \prod_{i=1}^n \frac{(1 - t^{1-j+n-i})^{\mu_i}}{(1 - qt^{n-i})^{\mu_i}}$$

where the sum is over all partitions $\mu \subsetneq \nu$ such that $\mu = (\mu_1, \dots, \mu_{j-1}, 0, \dots, 0)$. Likewise,

$$(105) \quad 0 = \sum_{\mu \subsetneq \nu} q^{n(\mu')} t^{-2n(\nu)+(n-1)|\nu|} s_2(\nu, \mu) \langle t^{1-j} \rangle_\mu$$

where the sum is over all partitions $\mu \subsetneq \nu$ such that $\mu = (\mu_1, \dots, \mu_{n-j}, 0, \dots, 0)$. The particular cases when $j = 1$ in (104) and $j = n$ in (105) show that the multiple Stirling numbers vanish when $\mu = \bar{0} = (0, \dots, 0) \in \mathbb{C}^n$ as in the classical case. That is,

$$s_1(\lambda, \bar{0}) = s_2(\lambda, \bar{0}) = 0$$

for any n -part partition λ with $\lambda_n \neq 0$.

(3) Note that setting $x = t^{1-j}q^{m_j}$ for $j = 1, \dots, n$, and $m_j \geq \nu_j$ in (81) and (83) will not vanish the bracket functions. In particular, setting $\nu = \bar{x} = \bar{k} = (k, \dots, k) \in \mathbb{Z}^n$ in (81) and using (43) gives

$$(106) \quad \bar{k}! = \prod_{i=1}^n \frac{(qt^{n-i})_k}{(1 - qt^{n-i})^k} \\ = \sum_{\mu \subseteq \bar{k}} q^{-n(\lambda')} t^{2n(\mu) - (n-1)|\mu|} s_1(\bar{k}, \mu) \prod_{i=1}^n \frac{(1 - q^k t^{n-i})^{\mu_i}}{(1 - qt^{n-i})^{\mu_i}}.$$

which is an analogue of the classical identity

$$k! = \sum_{m=0}^k s_1(k, m) k^m$$

for the special case when $\nu = \bar{k}$ is a rectangular partition.

(4) Recall that, $s_1(n, m) = -s_2(n, m) = -\binom{n}{2}$ when $n - m = 1$ for the classical Stirling numbers. Similarly, if the index partitions satisfy $|\lambda| - |\tilde{\lambda}| = 1$, we have that

$$s_1(\lambda, \tilde{\lambda}) = -s_2(\lambda, \tilde{\lambda})$$

exactly as in the one dimensional case.

The proof follows easily from the inversion (98) relation, and the observation that there are only two partitions between λ and $\tilde{\lambda}$ under the inclusion ordering, namely the two partitions themselves. That is,

$$(107) \quad 0 = \delta_{\lambda\tilde{\lambda}} = \sum_{\tilde{\lambda} \subseteq \mu \subseteq \lambda} s_1(\lambda, \mu) s_2(\mu, \tilde{\lambda})$$

which implies that $s_1(\lambda, \lambda) s_2(\lambda, \tilde{\lambda}) = -s_1(\lambda, \tilde{\lambda}) s_2(\tilde{\lambda}, \tilde{\lambda})$. That the diagonal entries of both type of multiple qt -Stirling numbers are 1 by (99) is now enough to conclude.

7. The qt -Lah numbers

The classical Lah numbers are defined to be the connection coefficients in the expansion

$$(108) \quad x^{\bar{n}} = \sum_{k=0}^n L(n, k) x_{\underline{k}}$$

where $x_{\underline{n}} = x(x-1)\dots(x-n+1)$ denotes the falling factorial as before, and $x^{\overline{n}} = x(x+1)\dots(x+n-1)$ the rising factorial. Various q -analogues of these numbers are developed in one dimensional case in [13, 32] and others.

We now give the definition of multiple qt -Lah numbers in terms of the multiple factorial function and its flipped version.

Definition 7.1. Let $[\bar{x}]^\lambda$ denote the multiple analogue of the rising factorial. That is,

$$(109) \quad [\bar{x}]^\lambda = [\bar{x}, q, t]^\lambda := [\bar{x}, q^{-1}, t^{-1}]_\lambda.$$

For an n part partition λ , the qt -Lah numbers $L(\lambda, \mu) = L(\lambda, \mu, q, t)$ are defined by

$$(110) \quad [\bar{x}]^\lambda = \sum_{\mu \subseteq \lambda} (-1)^{|\mu|} q^{-|\mu|+2n(\mu')} t^{-n(\mu)} L(\lambda, \mu, q, t) [\bar{x}]_\mu$$

where $x \in \mathbb{C}$, and $\bar{x} = \{x, \dots, x\} \in \mathbb{C}^n$ as before.

In one dimensional case, the Lah numbers admit some explicit representations. We show that the same is true for the multiple Lah numbers next.

Theorem 7.1. Let ν and μ be partitions with at most n -parts. Then

$$(111) \quad L(\nu, \mu) = (-1)^{-|\nu|+|\mu|} q^{-|\nu|+|\mu|-2n(\mu')} t^{n(\nu)+n(\mu)} \cdot \prod_{i=1}^n \left\{ (1 - qt^{n-i})^{-\nu_i+\mu_i} \right\} \sum_{\mu \subseteq \lambda \subseteq \nu} u(\nu, \lambda, q^{-1}, t^{-1}) v(\lambda, \mu, q, t)$$

where u and v factors are as defined in (74) and (75) above.

Proof. Multiply and divide the right hand side of (76) by $\prod_{i=1}^n (1 - qt^{n-i})^{\lambda_i}$, flip the parameters q and t , and use the definitions (93) and (109) to get

$$(112) \quad \langle x; q, t \rangle^\lambda = \sum_{\mu \subseteq \lambda} \prod_{i=1}^n \left\{ \frac{1}{(1 - q^{-1}t^{-(n-i)})^{\lambda_i}} \right\} u(\lambda, \mu, q^{-1}, t^{-1}) x^{|\mu|}.$$

Similarly, multiplying and dividing the right hand side of (77) by the factor $\prod_{i=1}^n (1 - qt^{n-i})^{\mu_i}$ gives

$$(113) \quad x^{|\lambda|} = \sum_{\mu \subseteq \lambda} v(\lambda, \mu, q, t) \prod_{i=1}^n \left\{ (1 - qt^{n-i})^{\mu_i} \right\} \langle x; q, t \rangle_\mu.$$

Combine the two to get

$$(114) \quad \langle x; q, t \rangle^\lambda = \sum_{\mu \subseteq \nu} \sum_{\mu \subseteq \lambda \subseteq \nu} \prod_{i=1}^n \left\{ \frac{1}{(1 - q^{-1}t^{-(n-i)})^{\nu_i}} \right\} u(\nu, \lambda, q^{-1}, t^{-1}) \\ \cdot v(\lambda, \mu, q, t) \prod_{i=1}^n \{(1 - qt^{n-i})^{\mu_i}\} \langle x; q, t \rangle_\mu.$$

Setting $x \rightarrow q^x$ and comparing the last identity to the definition of multiple Lah numbers (110) gives the desired result. \square

7.1. Properties of Lah numbers

We now establish a few fundamental properties of multiple Lah numbers starting with some special evaluations.

(1) Set $\nu = \mu = \lambda$ in (111) to write

$$(115) \quad L(\lambda, \lambda) = (-1)^{-|\lambda|+|\lambda|} q^{-|\lambda|+|\lambda|-2n(\lambda')} t^{n(\lambda)+n(\lambda)} \\ \cdot \prod_{i=1}^n \left\{ (1 - qt^{n-i})^{-\lambda_i + \lambda_i} \right\} u(\nu, \lambda, q^{-1}, t^{-1}) v(\lambda, \mu, q, t)$$

or

$$(116) \quad L(\lambda, \lambda) = q^{-2n(\lambda')} t^{2n(\lambda)} u(\lambda, \lambda, q^{-1}, t^{-1}) v(\lambda, \lambda, q, t).$$

It is clear from the definition (75) that

$$(117) \quad v(\lambda, \lambda, q, t) = (-1)^{|\lambda|} q^{n(\lambda')} t^{-n(\lambda)}.$$

Likewise, the identities and show that

$$(118) \quad \hat{w}_\lambda(q^\lambda t^{\delta(n)}; q, t) \\ = (-1)^{|\lambda|} t^{-n(\lambda)} q^{-|\lambda|-n(\lambda')} (qt^{n-1})_\lambda \prod_{1 \leq i < j \leq n} \frac{(qt^{j-i-1})_{\lambda_i - \lambda_j}}{(qt^{j-i})_{\lambda_i - \lambda_j}}.$$

Setting $\lambda = \mu$, and substituting the last evaluation in (74), and flipping the parameters q and t shows

$$(119) \quad u(\lambda, \lambda, 1/q, 1/t) = (-1)^{|\lambda|} t^{-n(\lambda)} q^{n(\lambda')}$$

Putting these into (116) now gives that $L(\lambda, \lambda) = 1$, as in the classical case.

(2) In the view of property (36), the definition (75), and the explicit formula (111), we see that $L(\lambda, \mu) = 0$ when $\mu \not\subseteq \lambda$.

(3) We now derive a multiple analogue of the closed formula

$$(120) \quad L(n, m) = \binom{n}{m} \frac{(n-1)!}{(m-1)!}.$$

We first write a generalization of the identity (58).

Lemma 7.1. *Let $r \in \mathbb{C}$, and ν is an n part partition. Then*

$$(121) \quad \begin{aligned} & (-1)^{|\mu|} r^{|\mu|} t^{-n(\mu)} q^{n(\mu')} \frac{(r)_\nu}{(r)_\mu} w_\mu(q^\nu t^{\delta(n)}; q, t) \\ &= \sum_{\substack{\lambda \\ \mu \subseteq \lambda \subseteq \nu}} \frac{(-1)^{|\lambda|} r^{|\lambda|} t^{n(\lambda)+(1-n)|\lambda|} q^{|\lambda|+n(\lambda')}}{(qt^{n-1})_\lambda} \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\lambda_i - \lambda_j}}{(qt^{j-i-1})_{\lambda_i - \lambda_j}} \right\} \\ & \quad \cdot w_\lambda(q^\nu t^{\delta(n)}; t, q) w_\mu(q^\lambda t^{\delta(n)}; t, q). \end{aligned}$$

Proof. Recall the following transformation identity for w_λ functions [8], which is obtained from a multiple analogue of Bailey’s $_{10}\phi_9$ transformation formula from [7].

$$(122) \quad \begin{aligned} & \frac{(s)_\nu}{(z)_\mu} (-1)^{|\mu|} z^{|\mu|} t^{-n(\mu)} q^{n(\mu')} \\ & \cdot \sum_{\substack{\lambda \\ \mu \subseteq \lambda \subseteq \nu}} \frac{(z)_\lambda}{(qt^{n-1})_\lambda} q^{|\lambda|} t^{2n(\lambda)} \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\lambda_i - \lambda_j}}{(qt^{j-i-1})_{\lambda_i - \lambda_j}} \right\} \\ & \cdot \left(\lim_{a \rightarrow 0} W_\lambda(q^\nu t^{\delta(n)}; q, t, at^{1-2n}, as^{-1}t^{-n}) \right) w_\mu(q^\lambda t^{\delta(n)}; t, q) \\ &= \sum_{\substack{\lambda \\ \mu \subseteq \lambda \subseteq \nu}} \frac{(-1)^{|\lambda|} z^{|\lambda|} s^{|\lambda|} t^{n(\lambda)+(1-n)|\lambda|} q^{|\lambda|+n(\lambda')}}{(qt^{n-1})_\lambda} \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\lambda_i - \lambda_j}}{(qt^{j-i-1})_{\lambda_i - \lambda_j}} \right\} \\ & \quad \cdot w_\lambda(q^\nu t^{\delta(n)}; t, q) w_\mu(q^\lambda t^{\delta(n)}; t, q). \end{aligned}$$

The sum on the left may be evaluated using the summation formula (62) to get the result where we set $r = zs$. □

This powerful result is more general than (58) because of the additional free variable r . If we set $r = t^{2n-2}$, we get

$$\begin{aligned}
 (123) \quad & (-1)^{|\mu|} t^{-n(\mu)} q^{n(\mu')} \frac{(t^{2(n-1)})_\nu}{(t^{2(n-1)})_\mu} \binom{\nu}{\mu}_{q,t} \\
 &= \sum_{\substack{\lambda \\ \mu \subseteq \lambda \subseteq \nu}} (-1)^{|\lambda|} t^{-n(\lambda)+2(n-1)|\lambda|} q^{n(\lambda')} \binom{\nu}{\lambda}_{q,t} \binom{\lambda}{\mu}_{q,t}
 \end{aligned}$$

which will be useful in writing our explicit formula that we give next.

Theorem 7.2. *With the notation as above,*

$$\begin{aligned}
 (124) \quad L(\nu, \mu) &= (-1)^{|\nu|+|\mu|} q^{-|\nu|+|\mu|} t^{n(\nu)-n(\mu)} \\
 &\quad \cdot \prod_{i=1}^n \{ (1 - qt^{n-i})^{-\nu_i+\mu_i} \} \frac{(t^{2(n-1)})_\nu}{(t^{2(n-1)})_\mu} \binom{\nu}{\mu}_{q,t}.
 \end{aligned}$$

Proof. Substituting (74) and (75) into the fomula (111) gives

$$\begin{aligned}
 (125) \quad L(\nu, \mu) &= (-1)^{|\nu|} q^{-|\nu|+|\mu|-n(\mu')} t^{n(\nu)} \prod_{i=1}^n \{ (1 - qt^{n-i})^{-\nu_i+\mu_i} \} \\
 &\quad \cdot \sum_{\mu \subseteq \lambda \subseteq \nu} \frac{q^{-|\lambda|} t^{-2n(\lambda)}}{(q^{-1}t^{-(n-1)}, q^{-1}, t^{-1})_\lambda} \prod_{1 \leq i < j \leq n} \left\{ \frac{(q^{-1}t^{-(j-i)})_{\lambda_i-\lambda_j}}{(q^{-1}t^{-(j-i-1)})_{\lambda_i-\lambda_j}} \right\} \\
 &\quad \cdot \hat{w}_\lambda(q^{-\nu}t^{-\delta(n)}; q^{-1}, t^{-1}) \binom{\lambda}{\mu}_{q,t}.
 \end{aligned}$$

By definition (17) we have

$$(126) \quad \hat{w}_\lambda(q^{-\nu}t^{-\delta(n)}; q^{-1}, t^{-1}) = q^{|\lambda|} t^{2n(\lambda)+(n-1)|\lambda|} w_\lambda(q^\nu t^{\delta(n)}; q, t)$$

and by the flip rule (48) we get

$$(127) \quad (q^{-1}t^{-(n-1)}, q^{-1}, t^{-1})_\lambda = (-1)^{|\lambda|} q^{-n(\lambda')-|\lambda|} t^{n(\lambda)+(1-n)|\lambda|} (qt^{n-1}; q, t)_\lambda.$$

Observe also that

$$(128) \quad \prod_{1 \leq i < j \leq n} \left\{ \frac{(q^{-1}t^{-(j-i)})_{\lambda_i-\lambda_j}}{(q^{-1}t^{-(j-i-1)})_{\lambda_i-\lambda_j}} \right\} = t^{2n(\lambda)+(1-n)|\lambda|} \prod_{1 \leq i < j \leq n} \left\{ \frac{(qt^{j-i})_{\lambda_i-\lambda_j}}{(qt^{j-i-1})_{\lambda_i-\lambda_j}} \right\}$$

Therefore, putting the multiple binomial (29) into (125) gives

$$(129) \quad L(\nu, \mu) = (-1)^{|\nu|} q^{-|\nu|+|\mu|-n(\mu')} t^{n(\nu)} \prod_{i=1}^n \{ (1 - qt^{n-i})^{-\nu_i + \mu_i} \} \\ \cdot \sum_{\mu \subseteq \lambda \subseteq \nu} (-1)^{|\lambda|} q^{n(\lambda')} t^{-n(\lambda)+2(n-1)|\lambda|} \binom{\nu}{\lambda}_{q,t} \binom{\lambda}{\mu}_{q,t}.$$

Finally, we use (123) to evaluate the sum which gives the desired result. \square

(4) The special evaluations we proved earlier, such as $L(\lambda, \lambda) = 1$ and $L(\lambda, \mu) = 0$ for $\mu \not\subseteq \lambda$ can be verified easily using this closed formula. Other special evaluations are also possible. For example, setting $\mu = \bar{1}$ gives

$$(130) \quad L(\nu, \bar{1}) = (-1)^{|\nu|+n} q^{-|\nu|+n} t^{n(\nu)-\binom{n}{2}} \\ \cdot (t^{2(n-1)})_\nu \prod_{i=1}^n \left\{ \frac{(1 - qt^{n-i})^{-\nu_i+1} (1 - q^{\nu_i} t^{n-i})}{(1 - t^{2n-1-i}) (1 - qt^{n-i})} \right\}$$

which is an analogue of

$$L(n, 1) = \binom{n}{1} \frac{(n-1)!}{(1-1)!} = n!.$$

This may be viewed as an alternative definition for $\nu!$.

(5) The final property we derive here is that the Lah numbers are self inverse of themselves in the sense that

$$(131) \quad x^{\bar{n}} = \sum_{k=0}^n L(n, k) x_{\underline{k}}, \quad \text{and} \quad x_{\underline{n}} = \sum_{k=0}^n (-1)^{n-k} L(n, k) x^{\bar{k}}.$$

Note that flipping the parameters q and t in the identity (110) gives the inverse result

$$(132) \quad [\bar{x}]_\lambda = \sum_{\mu \subseteq \lambda} (-1)^{|\mu|} q^{|\mu|-2n(\mu')} t^{n(\mu)} L(\lambda, \mu, t^{-1}, q^{-1}) [\bar{x}]^\mu$$

so that the matrices defined by the entries $L(\lambda, \mu, t, q)$ and $L(\lambda, \mu, t^{-1}, q^{-1})$ respectively are inverses of each other in the sense of Section 2.

8. Conclusion

We have derived several interesting identities for the multiple qt -factorial functions, multiple qt -Stirling numbers, and multiple qt -Lah numbers in the

present paper. We will construct additional properties such as new recurrence relations they satisfy, their explicit evaluations in various other special cases, their combinatorial interpretations (partially developed in [10]), other generating functions they satisfy, and their connections to different families of multiple combinatorial numbers in an upcoming article. The Stirling and Lah numbers have interesting connections to various branches in mathematics such as the one expressed in the classical Dobinski's formula. Such relations will also be formulated in that paper.

It should be noted that the multiple analogues developed in this paper are constructed mostly by taking a top-down approach, using the theory of multiple hypergeometric summation identities and well-poised Macdonald functions. The author is working on a book project on multiple combinatorial numbers where these combinatorial numbers as well as several other multiple sequences are reconstructed by taking a more elementary bottom-up approach.

References

- [1] G. E. Andrews, *q-Series: Their Development and Application in Analysis, Number Theory, Combinatorics, Physics, and Computer Algebra*, CBMS Regional Conference Series in Mathematics, 66 Amer. Math. Soc. 1986. [MR0858826](#)
- [2] G. E. Andrews, *Gaussian polynomials and finite Rogers-Ramanujan identities. Theory and applications of special functions*, Dev. Math., 13 (2005) 39–60. [MR2132458](#)
- [3] A. Berkovich, S. O. Warnaar, *Positivity preserving transformations for q-binomial coefficients*, Trans. Amer. Math. Soc. 357 6 (2005) 2291–2351. [MR2140441](#)
- [4] L. Carlitz, *q-Bernoulli numbers and polynomials*, Duke Math. J., 15 (1948), 987–1000. [MR0027288](#)
- [5] W. E. Clark, M. Ismail, *Binomial and q-binomial coefficient inequalities related to the Hamiltonicity of the Kneser graphs and their q-analogues*, J. Combin. Theory Ser. A 76 1 (1996), 83–98. [MR1405991](#)
- [6] H. Coskun, *A BC_n Bailey lemma and generalizations of Rogers-Ramanujan identities*, August 2003, Ph.D. thesis. [MR2705473](#)
- [7] H. Coskun, R. Gustafson, *The well-poised Macdonald functions W_λ and Jackson Coefficients ω_λ on BC_n* , Jack, Hall-Littlewood and Macdonald Polynomials, ICMS, AMS Contemporary Mathematics, 417 (2006), 127–155. [MR2284125](#)

- [8] H. Coskun, *An Elliptic BC_n Bailey Lemma, Multiple Rogers–Ramanujan Identities and Euler’s Pentagonal Number Theorems*, AMS Transactions, 360 (2008), 5397–5433. [MR2415079](#)
- [9] H. Coskun, *Multiple analogues of binomial coefficients and families of related special numbers*, Discrete Mathematics, 310 (2010) 17, 2280–2298. [MR2659180](#)
- [10] H. Coskun, *Combinatorial formulas for certain sequences of multiple numbers*, Ramanujan Journal, 2016, DOI: 10.1007/s11139-016-9843-8.
- [11] L. B. Frenkel, V. G. Turaev, *Elliptic solutions of the Yang-Baxter equation and modular hypergeometric functions*, The Arnold–Gelfand Mathematical Seminars, Birkhäuser, Boston, MA (1997), 171–204. [MR1429892](#)
- [12] K. C. Garrett, K. Hummel, *A combinatorial proof of the sum of q -cubes*, Electron. J. Comb. 11 (2004). [MR2034423](#)
- [13] A. Garsia, J. Remmel, *A combinatorial interpretation of q -derangement and q -Laguerre numbers*, (1980), European J. Combin. 1, 47–59. [MR0576766](#)
- [14] G. Gasper, M. Rahman, *Basic hypergeometric series, Encyclopedia of mathematics and its applications*, Vol 96, Cambridge University Press, Cambridge, (2004). [MR2128719](#)
- [15] F. Garvan, D. Stanton, *Sieved partition functions and q -binomial coefficients*, Math. Comp. 55 191 (1990), 299–311. [MR1023761](#)
- [16] H. W. Gould, *The q -Stirling numbers of first and second kinds*, Duke Math. J. 28 (1961), 281–289. [MR0122759](#)
- [17] H. W. Gould, *The bracket function, q -binomial coefficients, and some new Stirling number formulas*, Fibonacci Quart. 5 (1967), 401–422. [MR0225669](#)
- [18] J. Kaneko, *q -Selberg integrals and Macdonald polynomials*, Ann. Sci. ‘Ecole Norm. Sup. (4) 29 (1996), no. 5, 583–637. [MR1399617](#)
- [19] T. Kim, *q -Bernoulli numbers and polynomials associated with Gaussian binomial coefficients*, Russian Journal of Math. Physics 15 No. 1 (2008), 51–57. [MR2390694](#)
- [20] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*, 2nd ed., Oxford University Press, (1995). [MR1354144](#)

- [21] I. G. Macdonald, *An elementary proof of a q -binomial identity. q -series and partitions*, IMA Vol. Math. Appl. 18 (1989) 73–75. [MR1019844](#)
- [22] A. Lascoux, E. M. Rains, S. Ole Warnaar, *Nonsymmetric interpolation Macdonald polynomials and gl_n basic hypergeometric series*, Trans. Groups 14 (2009), 613–647. [MR2534801](#)
- [23] M. Lassalle, *Coefficients binomiaux generalises et polynomes de Macdonald*, J. Funct. Anal. 158 (1998), 289–324. [MR1648471](#)
- [24] S. C. Milne, *Restricted growth functions, rank row matchings of partition lattices, and q -Stirling numbers*. Adv. in Math. 43 (1982), no. 2, 173–196. [MR0644671](#)
- [25] A. Okounkov, *Binomial formula for Macdonald polynomials and applications*, Math. Res. Lett. 4 (1997), 533–553. [MR1470425](#)
- [26] E. Rains, *BC_n symmetric abelian functions*, Duke Math. Journal, Volume 135 (2006) 1, 99–180. [MR2259924](#)
- [27] E. Rains, *BC_n symmetric polynomials*, Transform. Groups 10 (2005) 1, 63–132. [MR2127341](#)
- [28] S. Sahi, *The binomial formula for nonsymmetric Macdonald polynomials*, Duke Math. J. 94 (1998), 465–477. [MR1639523](#)
- [29] M. Schlosser, *q -Analogues of the sums of consecutive integers, squares, cubes, quarts and quints*, The Electronic Journal of Combinatorics 11 (2004). [MR2097337](#)
- [30] S. O. Warnaar, *Summation and transformation formulas for elliptic hypergeometric series*, Constr. Approx. 18 (2002), 479–502. [MR1920282](#)
- [31] S. O. Warnaar, *On the q -analogue of the sum of cubes*, Electron. J. Comb. 11 (2004). [MR2114194](#)
- [32] C. Wagner, *Generalized Stirling and Lah numbers*, Discrete Math. (1996) 160, 199–218. [MR1417571](#)
- [33] J. Zeng, *The q -Stirling numbers, continued fractions and the q -Charlier and q -Laguerre polynomials*, J. Comput. Appl. Math. 57 (1995), no. 3, 413–424. [MR1335793](#)

HASAN COSKUN
DEPARTMENT OF MATHEMATICS
TEXAS A&M UNIVERSITY-COMMERCE
BINNION HALL, ROOM 314
COMMERCE, TX 75429
USA

E-mail address: hasan.coskun@tamuc.edu

URL: faculty.tamuc.edu/hcoskun

RECEIVED 13 MAY 2015