

ANALYTIC TEST CONFIGURATIONS AND GEODESIC RAYs

JULIUS ROSS AND DAVID WITT NYSTRÖM

Starting with the data of a curve of singularity types, we use the Legendre transform to construct weak geodesic rays in the space of locally bounded metrics on an ample line bundle L over a compact manifold. Using this we associate weak geodesics to suitable filtrations of the algebra of sections of L . In particular this works for the natural filtration coming from an algebraic test configuration, and we show how this recovers the weak geodesic ray of Phong–Sturm.

1. Introduction

Let $\mathcal{H}(L)$ be the space of smooth strictly positive Hermitian metrics on an ample holomorphic line bundle L over a compact manifold X . Then, by the work of Mabuchi [31], Semmes [43] and Donaldson [23], formally $\mathcal{H}(L)$ has the structure of an infinite-dimensional symmetric space with a canonical Riemannian metric. Thus, a natural way to study this space is through its geodesics, an approach that has been taken up by a number of authors (e.g., Berndtsson [11], Chen–Tian [18], Donaldson, Phong–Sturm [34, 35], Mabuchi, and Semmes among others).

In this paper, we give a general method for constructing weak geodesic rays in the space of locally bounded positive metrics on L . In the following we shall, in the standard way, identify a (positive) metric h on L with its (plurisubharmonic) potential $\phi = \log h$. Our initial data consist of a fixed smooth plurisubharmonic potential ϕ and a curve of singular plurisubharmonic potentials ψ_λ on L for $\lambda \in \mathbb{R}$. We are really only interested in the singularity type of ψ_λ , so we consider the equivalence class of ψ_λ under the relation $\psi_\lambda \sim \psi'_\lambda$ if $\psi_\lambda - \psi'_\lambda$ is bounded globally on X . We define the *maximal envelope* of this data to be

$$(1.1) \quad \phi_\lambda := \sup^* \{ \psi : \psi \leq \phi \text{ and } \psi \sim \psi_\lambda \}$$

where the supremum is over positive metrics ψ with the same singularity type as ψ_λ , and the star denotes the operation of taking the upper-semicontinuous regularization.

This equivalence relation on potentials was considered by Demailly–Peternell–Schneider [22] and is relevant to metrics with minimal singularities. The envelope in (1.1) was studied in a special case by Berman [6], and is the global analogue of a construction of Rashkovskii and Sigurdsson [32]. Observe that if ψ_λ is itself locally bounded then ϕ is a candidate for this envelope implying $\phi_\lambda = \phi$, and thus we are most interested in the case that ψ_λ is singular along some non-trivial subset of X . When ψ_λ has analytic singularities, it can be shown that ϕ_λ is the largest plurisubharmonic potential bounded above by ϕ with the same singularity type as ψ_λ (see Remark 4.6).

We shall call a curve ψ_λ of plurisubharmonic potentials a *test curve* if it is concave in λ , locally bounded for λ sufficiently negative and identically $-\infty$ for λ sufficiently large (we also make one further technical condition concerning the kind of singularity allowed; see Definition 5.1). A justification of this terminology will be given below, and as a simple example to have in mind, suppose that s is a holomorphic section of L scaled so $|s|_\phi^2 \leq 1$ and let

$$(1.2) \quad \psi_\lambda = \begin{cases} \phi & \lambda < 0, \\ (1 - \lambda)\phi + \lambda \ln |s|^2 & 0 \leq \lambda < 1, \\ -\infty & \lambda \geq 1, \end{cases}$$

so ψ_λ is decreasing and concave in λ . Geometrically then $\phi_\lambda = \phi$ for $\lambda \leq 0$ and for $\lambda \in (0, 1)$ it is the largest plurisubharmonic potential bounded above by ϕ with Lelong number at least λ along the divisor $Y = \{s = 0\}$.

Theorem 1.1. *Let ψ_λ be a test curve and ϕ be a smooth positive potential, and consider the Legendre transform of its maximal envelope ϕ_λ given by*

$$\widehat{\phi}_t := \sup_\lambda^* \{\phi_\lambda + \lambda t\} \quad \text{for } t \in [0, \infty).$$

Then $\widehat{\phi}_t$ is a weak geodesic ray in the space of locally bounded plurisubharmonic potentials on L that emanates from ϕ .

Looking back at (1.2) it is clear that specific examples of test curves are easy to produce, and thus this theorem gives a new construction of many weak geodesic rays emanating from ϕ . In fact, as we shall observe in Remark 6.4, essentially every weak geodesic ray can be produced in this way.

We recall precisely what is meant by a weak geodesic. Let $I \subset \mathbb{R}$ be an interval and consider the annulus $A := \{w \in \mathbb{C} : -\ln |w| \in I\}$ and $\pi: X \times A \rightarrow X$ be the projection. Then a curve of plurisubharmonic potentials ϕ_t for $t \in I$ can be identified with a rotation invariant potential $\Phi(x, w) := \phi_{-\log |w|}$ on π^*L . A simple calculation [43, 2.3, 2.4] reveals that if ϕ_t is a smooth curve of smooth potentials then the geodesic equation for ϕ_t is equivalent to the

homogeneous complex Monge–Ampère equation

$$(1.3) \quad MA(\Phi) := (dd^c\Phi)^{n+1} = 0 \quad \text{on } X \times A^\circ.$$

A curve $(\phi_t)_{t \in I}$ of locally bounded (not necessarily smooth) plurisubharmonic potentials given by $\Phi = \Phi(x, w)$ is said to be a *weak geodesic* if $dd^c\Phi$ is positive and solves (1.3) in the sense of currents. When $I = [0, \infty)$ (so A is the punctured unit disc) we say that ϕ_t is a *weak geodesic ray*.

The first step in our approach to Theorem 1.1 is showing that the Monge–Ampère measure of the maximal envelope ϕ_λ satisfies

$$(1.4) \quad MA(\phi_\lambda) = \mathbf{1}_{\{\phi_\lambda = \phi\}} MA(\phi_\lambda) \quad \text{for all } \lambda,$$

where $\mathbf{1}_S$ denotes the characteristic function of a set S . We say that a positive metric ϕ_λ bounded by ϕ and having property (1.4) is *maximal* with respect to ϕ (see Definition 4.5), and a test curve ϕ_λ where ϕ_λ is maximal with respect to ϕ for all λ is referred to as a *maximal test curve*. We shall show that the Legendre transform $\widehat{\phi}_t$ of a maximal test curve ϕ_λ is a subgeodesic, and also that Aubin–Mabuchi energy is linear in t , which together imply that it is a weak geodesic (in the smooth case this is well known and goes back to Mabuchi [30] and more generally can be deduced from a Theorem of Berman–Boucksom–Guedj–Zeriahi, see Lemma 3.11).

The well-known Yau–Tian–Donaldson conjecture states that for a smooth projective manifold it should be possible to detect the existence of a constant scalar curvature Kähler metric algebraically. Through ideas developed by many authors (e.g., Chen, Donaldson, Mabuchi, Tian) a general picture has emerged in which such metrics appear as critical points of certain energy functionals that are convex along smooth geodesics. The input from algebraic geometry arises through notion of an (algebraic) test configuration, originally due to Tian [46] and then extended by Donaldson [24], which, roughly speaking, is a one-parameter algebraic degeneration of the original projective manifold X .

In a series of papers, Phong–Sturm show how one can naturally associate a weak geodesic ray to a test configuration [34, 35, 37] (see also [2, 15–17, 44, 45] for other constructions of geodesic rays related to test configurations). We show how the geodesic of Phong–Sturm can be viewed as a particular case of the Legendre transform construction. For example, applying the Legendre transform construction to the example of the test curve (1.2) given above recovers the geodesic of Phong–Sturm that is associated to the test configuration given by the degeneration to the normal cone of the divisor Y .

Generalizing slightly, suppose that $\mathcal{F}_{k,\lambda}$, for $k \in \mathbb{N}, \lambda \in \mathbb{R}$ is a multiplicative filtration of the graded algebra $\oplus_k H^0(X, kL)$. Using the underlying smooth positive metric ϕ and some auxiliary volume form we have an L^2 -inner product on each $H^0(X, kL)$, and thus can consider the associated

Bergman metric

$$\phi_{k,\lambda} = \frac{1}{k} \ln \sum_{\alpha} |s_{\alpha}|^2$$

where $\{s_{\alpha}\}$ is an orthonormal basis for $\mathcal{F}_{k,\lambda k} \subset H^0(X, kL)$. So, for example, if Y is a divisor on X we can let $\mathcal{F}_{k,\lambda} = \{s \in H^0(kL) : \text{ord}_Y(s) \geq \lambda\}$ be the multiplicative filtration given by order of vanishing along Y , and then $\phi_{k,\lambda}$ is the potential associated to the partial Bergman kernel coming from sections of kL that vanish to at least order $k\lambda$ along Y .

Theorem 1.2. *Suppose that $\mathcal{F}_{k,\lambda}$ is left continuous and decreasing in λ and bounded (see (7.2)). Then there is a well-defined limit*

$$\phi_{\lambda}^{\mathcal{F}} = \lim_{k \rightarrow \infty}^* \phi_{k,\lambda}$$

where the star denotes taking the upper semicontinuous regularization after the limit. Furthermore this limit is maximal except possibly for one critical value of λ , and its Legendre transform is a weak geodesic ray.

In particular, this applies to a natural filtration associated to a test configuration, and thus we have associated a weak geodesic to any such test configuration. We prove that, in this case, we recover precisely Phong–Sturm geodesic, thus reproving the main result of [35]. Hence, one interpretation of Theorem 1.1 is that in the problem of finding weak geodesic rays, the algebraic data of a test configuration can be replaced with a suitable concave curve of singularity types which we thus refer to as an *analytic test configuration*.

The relationship between algebraic and analytic test configurations is analogous to the relationship between holomorphic and plurisubharmonic functions. Being given by a concave curve of singularity types, analytic test configurations are easier to produce and manipulate. For example, it is possible to interpolate between two analytic test configurations merely by taking the line between them, whereas it is not so clear what the analogous construction is for algebraic test configurations. With regard to the Yau–Tian–Donaldson conjecture it is now expected that the notion of algebraic test configuration needs to be extended in some way for it to detect a constant scalar curvature metric (see the examples of Apostolov–Calderbank–Gauduchon–Friedman [1] and also work of Székelyhidi [38], which considers general filtrations of the ring of sections). Thus, analytic test configurations provide a context for such extensions.

Interesting examples of analytic test configurations that do not correspond to algebraic ones appear for example in the setting of Arakelov geometry. In [12] Boucksom–Chen construct multiplicative filtrations of the section ring, which encodes arithmetic properties of the sections such as their adelic norm. This filtration then gives rise to an analytic test configuration whose geometrical significance remains unclear.

Another advantage of the notion of analytical test configurations is that they can be defined independent of the polarization, and even in the non-projective case (although we will not consider that further here). If we pick an arbitrary Kähler form ω we can consider a concave curve of singularity types $[\psi_\lambda]$ of ω -plurisubharmonic functions. Then choosing a polarization L we get an analytic test configuration $[\phi_\lambda]$ by letting ϕ_λ be the envelope of all positive singular metrics of L bounded from above by $\phi + \psi_\lambda$, where ϕ is some locally bounded metric of L . In fact, rather than choosing a polarization we could have picked an arbitrary big $(1, 1)$ -cohomology class and worked with that instead.

It should be stressed that in the problem of finding constant scalar curvature metrics it is important to have control of the regularity of geodesics under consideration. By using approximations to known regularity results of solutions of Monge–Ampère equations, Phong–Sturm prove that their weak geodesic is in fact $C^{1,\alpha}$ for $0 < \alpha < 1$ (see [37]). It is interesting to ask whether such regularity holds more generally, which is a topic we take up in [40].

Organization: We start in Section 2 with some motivation from convex analysis and toric geometry, and Section 3 contains preliminary material on the space of singular metrics, the Monge–Ampère measure and the Aubin–Mabuchi functional. The real work starts in Section 4 where we consider the maximal envelopes associated to a given singularity type. Along the way we prove a generalization of a theorem of Bedford–Taylor which says that such envelopes are maximal (Theorem 4.10). This is then extended to the case of a test curve of singularities, and in Section 6 we discuss the Legendre transform and prove Theorem 1.1.

Following these analytic results, we move on to the algebraic picture. In Section 7, we associate a test curve to a suitable filtration of the coordinate ring of (X, L) , and prove Theorem 1.2. We then recall how such filtrations arise from test configurations, and in Section 9 show how this agrees with the construction of Phong and Sturm.

2. Convex motivation

This section contains some motivation from convex analysis in the study of the homogeneous Monge–Ampère equation. Much of this material is standard; our main references are the two papers [41, 42] by Rubinstein–Zelditch. We shall presently see how solutions to this equation can be found using the Legendre transform in two different, but ultimately equivalent, ways. Although this is logically independent of the rest of the paper, the techniques used are similar and give an illustration in the toric setting.

Let $\text{Conv}(\mathbb{R}^n)$ denote the space of convex functions on \mathbb{R}^n . We take the convention that the function identically equal to $-\infty$ is in $\text{Conv}(\mathbb{R}^n)$.

Definition 2.1. Let ϕ be a C^2 convex function on an open subset of \mathbb{R}^n . The (real) Monge–Ampère measure of ϕ , denoted by $MA(\phi)$, is the Borel measure defined as

$$MA(\phi) := d\frac{\partial\phi}{\partial x_1} \wedge \dots \wedge d\frac{\partial\phi}{\partial x_{n+1}}.$$

The operator MA has an unique extension to a continuous operator on the cone of (finite-valued) convex functions (see [42] for references). If ϕ is C^2 then

$$(2.1) \quad MA(\phi) = \det(\nabla^2\phi)dx = (\nabla\phi)^*dx,$$

i.e., the Monge–Ampère measure is the pullback of the Lebesgue measure under the gradient map.

If $\phi \in \text{Conv}(\mathbb{R}^n)$, we say y is a *subgradient* of ϕ at x_0 if the function $x \mapsto \phi(x) - \phi(x_0) - y \cdot (x - x_0)$ is bounded from below, and we let

$$\Delta_\phi = \{y : y \text{ is a subgradient for some } x_0\}$$

be the set of all subgradients of ϕ . So, if ϕ is differentiable then Δ_ϕ is simply the image of $\nabla\phi$. One can easily check that Δ_ϕ is convex, that if $r > 0$ then $\Delta_{r\phi} = \Delta_\phi$ and $\Delta_{\phi+\psi} \subseteq \Delta_\phi + \Delta_\psi$.

When ϕ is C^2 it follows from equation (2.1) that the total mass of the Monge–Ampère measure $MA(\phi)$ equals the Lebesgue volume of Δ_ϕ . An important fact [42] is that this is true for all convex functions on \mathbb{R}^n with linear growth, i.e.,

$$(2.2) \quad \int_{\mathbb{R}^n} MA(\phi) = \text{vol}(\Delta_\phi).$$

We say two convex functions ϕ and ψ are *equivalent* if $|\phi - \psi|$ is bounded, and denote this by $\phi \sim \psi$. Since for two equivalent convex functions ϕ and ψ with linear growth we clearly have that

$$\Delta_\phi = \Delta_\psi,$$

it follows from (2.2) that

$$\int_{\mathbb{R}^n} MA(\phi) = \int_{\mathbb{R}^n} MA(\psi) \quad \text{whenever } \phi \sim \psi.$$

Definition 2.2. Let $\phi \in \text{Conv}(\mathbb{R}^n)$ and let $\dot{\phi}$ be a bounded continuous function on \mathbb{R}^n . A curve ϕ_t in $\text{Conv}(\mathbb{R}^n)$, $t \in [a, b]$, is said to solve the *Cauchy problem* for the homogeneous real Monge–Ampère equation, abbreviated as HRMA, with initial data $(\phi, \dot{\phi})$, if the function $\Phi(x, t) := \phi_t(x)$ is convex on $\mathbb{R}^n \times [a, b]$, and satisfies the equation

$$MA(\Phi) = 0 \quad \text{on the strip } \mathbb{R}^n \times (a, b),$$

with initial data

$$\phi_0 = \phi, \quad \frac{\partial}{\partial t}|_{t=0^+} \phi_t = \dot{\phi}.$$

Remark 2.3. This convex geometry has particular geometric significance when Δ is the moment polytope of a polarized toric manifold (X, L) . Then, through the notion of symplectic potentials, there is a correspondence between Hermitian metrics on L and convex functions on Δ (where due care is to be taken to ensure that a given convex function determines a smooth or positive metric on L) and the Cauchy problem for the HRMA translates to the Cauchy problem for finding geodesics in the space of Hermitian metrics on L . We refer the reader to [41, 42] for a detailed discussion of this idea.

Now let ϕ_0 and ϕ_1 be two equivalent convex functions with linear growth, and ϕ_t be the affine curve between them. The *energy* of ϕ_1 relative to ϕ_0 , denoted by $\mathcal{E}(\phi_1, \phi_0)$ is defined as

$$\mathcal{E}(\phi_1, \phi_0) := \int_{t=0}^1 \left(\int_{\mathbb{R}^n} (\phi_1 - \phi_0) MA(\phi_t) \right) dt.$$

We observe that by the linear growth assumption it follows that the relative energy $\mathcal{E}(\phi_1, \phi_0)$ is finite. This energy has a cocycle property, namely if ϕ_0 , ϕ_1 and ϕ_2 are equivalent with finite energy then

$$\mathcal{E}(\phi_2, \phi_0) = \mathcal{E}(\phi_2, \phi_1) + \mathcal{E}(\phi_1, \phi_0),$$

which is easily seen to be equivalent to the fact that

$$\frac{\partial}{\partial t} \mathcal{E}(\phi_t, \phi) = \int_{\mathbb{R}^n} \frac{\partial}{\partial t} \phi_t MA(\phi_t).$$

The energy along a smooth curve ϕ_t of convex functions with linear growth is related to the Monge–Ampère measure of $\Phi(x, t) := \phi_t(x)$ by the identity

$$(2.3) \quad \int_{\mathbb{R}^n \times [a, b]} MA(\Phi) = \frac{\partial}{\partial t}|_{t=b} \mathcal{E}(\phi_t, \phi_a) - \frac{\partial}{\partial t}|_{t=a} \mathcal{E}(\phi_t, \phi_a).$$

Thus a smooth curve ϕ_t of equivalent convex functions of linear growth solves the HRMA equation if and only if Φ is convex and the energy $\mathcal{E}(\phi_t, \phi_a)$ is linear in t .

As is noted in [42] this Cauchy problem is not always solvable. Nevertheless there is a standard way to produce solutions ϕ_t with $t \in [0, \infty)$ to the homogeneous Monge–Ampère equation with given starting point $\phi_0 = \phi$ using the Legendre transform. We give a brief account of this.

For simplicity assume from now on that ϕ is differentiable and strictly convex. Recall that the Legendre transform of ϕ is the function on Δ_ϕ defined as

$$\phi^*(y) := \sup_x \{x \cdot y - \phi(x)\}$$

(which we can also think of as being defined on the whole of \mathbb{R}^n , by being $-\infty$ outside of Δ_ϕ). Since ϕ^* is defined as the supremum of the linear functions $x \cdot y - \phi(x)$, it is convex. In fact, one can show that ϕ being differentiable and strictly convex essentially implies that ϕ^* is also differentiable and strictly convex (see [33, Theorem 1] for a precise statement that requires a further boundary condition).

For a given $y \in \Delta_\phi$, the function $x \cdot y - \phi(x)$ is strictly concave, and is maximized at the point where the gradient is zero. Thus

$$(2.4) \quad \phi^*(y) = x \cdot y - \phi(x), \quad \text{where} \quad \nabla\phi(x) = y,$$

and hence

$$\nabla\phi^*(y) = x, \quad \text{where} \quad \nabla\phi(x) = y.$$

The Legendre transform is an involution. For using the above formula $\nabla\phi^{**}(x) = y$ for x such that $\nabla\phi^*(y) = x$, which holds when $\nabla\phi(x) = y$, i.e.,

$$\nabla\phi^{**}(x) = \nabla\phi(x).$$

If $\nabla\phi(x) = y$, then $\phi^*(y) = x \cdot y - \phi(x)$, therefore

$$\phi^{**}(x) = x \cdot y - \phi^*(y) = x \cdot y - (x \cdot y - \phi(x)) = \phi(x),$$

as claimed.

Lemma 2.4. *If ϕ_t is a curve of convex functions, then for any point $y \in \Delta_{\phi_t}$*

$$\frac{\partial}{\partial t}\phi_t^*(y) = -\frac{\partial}{\partial t}\phi_t(x),$$

where x is the point such that $\nabla\phi(x) = y$.

Proof. Let x_t be the solution to the equation $\nabla\phi_t(x_t) = y$. By the implicit function theorem x_t varies smoothly with t . By equation (2.4) we know

$$\frac{\partial}{\partial t}\phi_t^*(y) = \frac{\partial}{\partial t}(x_t \cdot y - \phi_t(x_t)) = \frac{\partial}{\partial t}(x_t \cdot y - \phi(x_t)) - \frac{\partial}{\partial t}\phi_t(x_t).$$

Since $x_t \cdot y - \phi(x_t)$ is maximized at $x = x_t$ the derivative of that part vanishes at $t = 0$, so we get the lemma for $t = 0$, and similarly for all t . \square

This leads us to the following formula relating the energy with the Legendre transform,

Lemma 2.5. *We have*

$$(2.5) \quad \mathcal{E}(\phi_t, \phi) = \int_{\Delta_\phi} (\phi^* - \phi_t^*) dy.$$

Proof. We noted above that the derivative with respect to t of the left-hand side of (2.5) is equal to

$$\int_{\mathbb{R}^n} \frac{\partial}{\partial t}\phi_t MA(\phi_t).$$

On the other hand, differentiating the right-hand side yields

$$\begin{aligned} \frac{\partial}{\partial t} \int_{\Delta_\phi} (\phi^* - \phi_t^*) dy &= - \int_{\Delta_\phi} \frac{\partial}{\partial t} \phi_t^* dy = \int_{\Delta_\phi} \frac{\partial}{\partial t} \phi_t (\nabla \phi_t^{-1}(y)) dy \\ &= \int_{\mathbb{R}^n} \frac{\partial}{\partial t} \phi_t (\nabla \phi_t)^* dy = \int_{\mathbb{R}^n} \frac{\partial}{\partial t} \phi_t MA(\phi_t), \end{aligned}$$

where we used Lemma 2.4 and the fact that the pullback $(\nabla \phi_t)^* dy$ of the Lebesgue measure is $MA(\phi_t)$. Since both sides of the equation (2.5) is zero when $\phi_t = \phi$ and the derivatives coincide, we get that they must be equal for all t . \square

Now fix a smooth bounded strictly concave function u on Δ_ϕ and let

$$(2.6) \quad \tilde{\phi}_t := (\phi^* - tu)^*.$$

Proposition 2.6. *The curve $\tilde{\phi}_t$, $t \in [0, \infty)$ solves the HRMA equation with initial data*

$$\tilde{\phi}_0 = \phi \quad \text{and} \quad \frac{\partial}{\partial t} \Big|_{t=0+} \tilde{\phi}_t = u((\nabla \phi)^{-1}).$$

To see this note that from (2.5) it follows that

$$\mathcal{E}(\tilde{\phi}_t, \phi) = \int_{\Delta_\phi} (\phi^* - \tilde{\phi}_t^*) dy = \int_{\Delta_\phi} (\phi^* - \phi^* + tu) dy = t \int_{\Delta_\phi} u dy,$$

which is linear in t , and the initial conditions follow from (2.6) and Lemma 2.4. The convexity of $\tilde{\Phi}(t, x) = \tilde{\phi}_t(x)$ can of course be shown directly, but it also follows from another characterization of $\tilde{\phi}_t$ that also involves a Legendre transform, but in the t -coordinate instead of in the x -coordinates, which we now discuss.

Let A_λ be the subset of Δ_ϕ where u is greater than or equal to λ and let ϕ_λ be defined as

$$(2.7) \quad \phi_\lambda := \sup\{\psi \leq \phi : \psi \in \text{Conv}(\mathbb{R}^n), \Delta_\psi \subseteq A_\lambda\}.$$

Lemma 2.7. *The curve of functions ϕ_λ is concave in λ and*

$$\{\phi_\lambda = \phi\} = \{x : \nabla \phi(x) \in A_\lambda\}.$$

Proof. Let $\psi_i \leq \phi$ be such that $\Delta_{\psi_i} \subseteq A_{\lambda_i}$ with $i = 1, 2$. Let $0 < t < 1$. From our discussion above it follows that $t\psi_1 + (1-t)\psi_2 \leq \phi$ and

$$\Delta_{t\psi_1 + (1-t)\psi_2} \subseteq t\Delta_{\psi_1} + (1-t)\Delta_{\psi_2} \subseteq tA_{\lambda_1} + (1-t)A_{\lambda_2} \subseteq A_{t\lambda_1 + (1-t)\lambda_2},$$

where the last inclusion follows from the fact that u was assumed to be concave. For the second statement, it is easy to see that in fact ϕ_λ is equal to the supremum of affine functions $x \cdot y + C$ bounded by ϕ and y lying in A_λ . \square

Definition 2.8. For $t \geq 0$ let $\widehat{\phi}_t$ be defined as

$$\widehat{\phi}_t := \sup_{\lambda} \{\phi_{\lambda} + t\lambda\}.$$

Since for each λ the function $(x, t) \mapsto \phi_{\lambda}(x) + t\lambda$ is convex in all its variables, and the supremum of convex functions is convex, $\widehat{\Phi}(x, t) := \widehat{\phi}_t(x)$ is convex.

Proposition 2.9. *Recalling that $\tilde{\phi}_t = (\phi^* - tu)^*$ we have $\tilde{\phi}_t = \widehat{\phi}_t$. In particular, this proves that $\widehat{\Phi}$ is convex, thereby proving $\tilde{\phi}_t$ solves the HRMA equation (Proposition 2.6).*

Proof. We claim

$$(2.8) \quad \frac{\partial}{\partial t} \widehat{\phi}_t(x) = u(\nabla \widehat{\phi}_t(x)).$$

To see this first consider the right-derivative at $t = 0$. As we noted above, the gradient of a Legendre transform is the point where the maximum is attained, thus in this case

$$\frac{\partial}{\partial t} \Big|_{t=0^+} \widehat{\phi}_t(x) = \sup\{\lambda : \phi_{\lambda}(x) = \phi(x)\}.$$

By the second statement in Lemma 2.7 it follows that this supremum is equal to $u(\nabla \phi(x))$, and we are done for $t = 0$. On the other hand it is easy to see that

$$\widehat{\phi}_{t_1+t_2} = \widehat{\psi}_{t_2},$$

with $\psi := \widehat{\phi}_{t_1}$. Using this, equation (2.8) holds for all t . Thus, by Lemma 2.4 the Legendre transform of $\widehat{\phi}_t$ is equal to $\phi - tu$, so by the involution property of the Legendre transform $\widehat{\phi}_t$ coincides with $\tilde{\phi}_t$. \square

Now the above discussion can be reformulated as follows. Let ψ_{λ} be a curve in $\text{Conv}(\mathbb{R}^n)$. We say that ψ_{λ} is a *test curve* if there is a C such that

- (1) ψ_{λ} is concave in λ ,
- (2) $\psi_{\lambda} - \phi$ bounded for $\lambda < -C$ and
- (3) $\psi_{\lambda} \equiv -\infty$ for $\lambda > C$.

Given such a test curve let u be the function on Δ_{ϕ} defined by

$$u(y) := \sup\{\lambda : y \in \Delta_{\psi_{\lambda}}\}.$$

and observe that since ψ_{λ} is assumed to be concave the function u is also concave.

Thus, our definition of ϕ_{λ} above (2.7) is

$$(2.9) \quad \begin{aligned} \phi_{\lambda} &= \sup\{\psi \leq \phi : \psi \in \text{Conv}(\mathbb{R}^n), \Delta_{\psi} \subseteq A_{\lambda}\} \\ &= \sup\{\psi \leq \phi : \psi \in \text{Conv}(\mathbb{R}^n), \Delta_{\psi} \subseteq \{u \geq \lambda\}\}. \end{aligned}$$

Hence, we in fact have

$$(2.10) \quad \phi_\lambda := \sup\{\psi : \psi \leq \phi, \psi \leq \psi_\lambda + o(1), \psi \in \text{Conv}(\mathbb{R}^n)\}.$$

From Proposition 2.9, $\widehat{\phi}_t$ solves the homogeneous real Monge–Ampère equation. Thus in order to get solutions to the HRMA, instead of starting with a concave function u on Δ_ϕ we can just as well start with a test curve ψ_λ . It is this second reformulation (2.10) that extends more naturally to the context of positive metrics on line bundles.

Remark 2.10. This convex picture can be given some geometric context by considering toric manifolds (compare [5, Example 5.2]). Consider a complex toric manifold (X, L) of dimension n , so L carries a compatible torus invariant Hermitian metric ϕ_0 . Pick complex coordinates z_1, \dots, z_n on a dense complex torus $T = (\mathbb{C}^*)^n \subset X$ and let $t_i = \ln |z_i|^2$ so we consider t_1, \dots, t_n as coordinates on \mathbb{R}^n . Then $\phi_0(z)$ descends to a convex function on \mathbb{R}^n which, by abuse of notation, we denote by $\phi_0(t)$. Conversely any function ϕ on \mathbb{R}^n induces a metric on $L|_T$, which will be (strictly) positive as long as ϕ is (strictly) convex (one must make some additional hypothesis as to the behavior of $\phi(t)$ at infinity if one wishes to ensure that this induced metric extends to a smooth or locally bounded metric over all of X , but we will not consider that further here).

As an illustration, consider the simplest case of dimension 1 so $X = \mathbb{P}^1$. Fix a smooth strictly positive metric on L and assume the strictly convex potential $\phi(t)$ on \mathbb{R} satisfies $\phi'(0) = 0$ and $\lim_{t \rightarrow \infty} \phi'(t) = 1$. Then, for $\lambda \in (0, 1)$ define

$$\psi_\lambda(t) = \begin{cases} \lambda t & t \geq 0 \\ 0 & t \leq 0 \end{cases}$$

(this can be made into a test curve by extending it to be $\phi + C'$ for negative λ and to be $-\infty$ for $\lambda \geq 1$ similar to the example (1.2) in the Introduction). Now for $\lambda \in (0, 1)$ let t_λ be the point where $\phi'(t_\lambda) = \lambda$. Then one sees from (2.9) that $\phi_\lambda = \phi$ for $t \leq t_\lambda$ and $\phi_\lambda = \lambda(t - t_\lambda) + \phi(t_\lambda)$ for $t \geq t_\lambda$. One can check that ϕ_λ defines a singular metric on L with the same singularity type as ψ_λ . Observe that ϕ_λ is linear for $t > t_\lambda$ making this metric pluriharmonic (and thus maximal) over this set, and it is this maximal property that will be crucial in the generalization that follows.

3. Preliminary material

We collect here some preliminary material on the space of positive metrics, the (non-pluripolar) Monge–Ampère measure and the Aubin–Mabuchi energy functional. Most of this material is standard, and we give proofs only for those results for which we did not find a convenient reference.

3.1. The space of positive singular metrics. Let X be a complex projective manifold of complex dimension n , and let L be an ample line bundle on X . We start with some preliminaries on singular metrics, for which a convenient reference is [19]. A continuous (or smooth) Hermitian metric on L is a continuous (or smooth) choice of scalar product on the complex line L_p at each point p on the manifold. If f is a local holomorphic frame for L on U_f , then one writes

$$|f|_h^2 = h_f = e^{-\phi_f},$$

where ϕ_f is a continuous (or smooth) function on U_f . We will use the convention to let ϕ denote the metric $h = e^{-\phi}$, thus if ϕ is a metric on L , $k\phi$ is a metric on $kL := L^{\otimes k}$.

The curvature of a smooth metric is given by $dd^c\phi$ which is the $(1, 1)$ -form locally defined as $dd^c\phi_f$, where f is any local holomorphic frame. Here d^c is short-hand for the differential operator

$$\frac{i}{2\pi}(\bar{\partial} - \partial),$$

so $dd^c = i/\pi\partial\bar{\partial}$. A classic fact is that the curvature form of a smooth metric ϕ is a representative for the first Chern class of $c_1(L)$. The metric ϕ is said to be strictly positive if the curvature $dd^c\phi$ is strictly positive as a $(1, 1)$ -form, i.e., if for any local holomorphic frame f , the function ϕ_f is strictly plurisubharmonic. We let $\mathcal{H}(L)$ denote the space of smooth strictly positive (i.e., locally strictly plurisubharmonic) metrics on L , which is non-empty since we assumed that L was ample.

A positive singular metric ψ is a metric that can be written as $\psi := \phi + u$, where ϕ is a smooth metric and u is a globally defined $dd^c\phi$ -psh function, i.e., u is upper semicontinuous and $dd^c\psi := dd^c\phi + dd^c u$ is a positive $(1, 1)$ -current. For convenience we also allow $u \equiv -\infty$. We let $PSH(L)$ denote the space of positive singular metrics on L .

As an important example, if $\{s_i\}$ is a finite collection of holomorphic sections of kL , we get a positive metric $\psi := \frac{1}{k} \ln(\sum |s_i|^2)$ which is defined by letting for any local frame f ,

$$e^{-\psi_f} := \frac{|f|^2}{(\sum |s_i|^2)^{1/k}}.$$

We note that $PSH(L)$ is a convex set, since any convex combination of positive metrics yields a positive metric. Another important fact is that if $\psi_i \in PSH(L)$ for all $i \in I$ are uniformly bounded above by some fixed positive metric, then the upper semicontinuous regularization of the supremum denoted by $\sup^*\{\psi_i : i \in I\}$ lies in $PSH(L)$ as well. Similarly the upper semicontinuous regularization of the pointwise limit $\lim_i \psi_i$ (when defined) will be denoted by $\lim_i^* \psi_i$.

If ψ is in $PSH(L)$, then the translate $\psi + c$ where c is a real constant is also in $PSH(L)$. For any $\psi \in PSH(L)$, $dd^c\psi$ is a closed positive $(1, 1)$ -current, and from the dd^c lemma it follows that any closed positive current cohomologous with $dd^c\psi$ can be written as $dd^c\phi$ for some ϕ in $PSH(L)$. By the maximum principle this ϕ is uniquely determined up to translation.

If there exists a constant C such that $\psi \leq \phi + C$, we say that ψ is more singular than ϕ , and we will write this as

$$\psi \preceq \phi.$$

If both $\psi \succeq \phi$ and $\phi \succeq \psi$ we say that ψ and ϕ are *equivalent*, which we write as $\psi \sim \phi$. Following [13] an equivalence class $[\psi]$ is called a *singularity type*, and we use the notation $Sing(L)$ for the set of singularity types. If ψ is equivalent to an element in $\mathcal{H}(L)$ we say that ψ is *locally bounded*.

The *singularity locus* of a positive metric ψ is the set where ψ is minus infinity, i.e., the set where $\psi_f = -\infty$ when f is a local frame and the *unbounded locus* of ψ is the set where ψ is not locally bounded. Recall that a set is said to be *complete pluripolar* if it is locally the singularity locus of a plurisubharmonic function, while it is called pluripolar if you only have a local inclusion in the singularity set. Pluripolar sets have zero measure with respect to any smooth volume form (since this is true locally with respect to the Lebesgue measure [29, Corollary 2.9.10]). In [13] Boucksom–Eyssidieux–Guedj–Zeriahi give the following definition.

Definition 3.1. A positive metric ψ is said to have *small unbounded locus* if its unbounded locus is contained in a closed complete pluripolar subset of X .

We note that metrics of the form $\frac{1}{k} \ln(\sum |s_i|^2)$ have small unbounded locus, since they are locally bounded away from the algebraic set $\cup_i \{s_i = 0\}$ which is a closed pluripolar set.

3.2. Regularization of positive singular metrics. If f is a plurisubharmonic function on an open subset U of \mathbb{C}^n then using a convolution we can write f as the limit of a decreasing sequence of smooth plurisubharmonic functions on any relatively compact subset of U [29, Theorem 2.9.2].

If ψ is a positive singular metric, we can use a partition of unity with respect to some open cover U_{f_i} to patch together the smooth decreasing approximations of ψ_{f_i} . Thus any positive singular metric can be written as the pointwise limit of a decreasing sequence of smooth metrics, but of course because of the patching these smooth approximations will in general not be positive.

A fundamental result due to Demailly [20] is that any positive singular metric can be approximated by metrics of the form $k^{-1} \ln(\sum_i |s_i|^2)$, where s_i are sections of kL . Let $\mathcal{I}(\psi)$ denote the multiplier ideal sheaf of germs of holomorphic functions locally integrable against $e^{-\psi_f} dV$, where f is a local

frame for L and dV is an arbitrary volume form. We get a scalar product $(\cdot, \cdot)_{k\psi}$ on the space $H^0(kL \otimes \mathcal{I}(k\psi))$ by letting

$$\|s\|_{k\psi}^2 := \int_X |s|^2 e^{-k\psi} dV.$$

Let $\{s_i\}$ be an orthonormal basis for $H^0(kL \otimes \mathcal{I}(k\psi))$ and set

$$\psi_k := \frac{1}{k} \ln \left(\sum |s_i|^2 \right).$$

Theorem 3.2. *The sequence of metrics ψ_k converge pointwise to ψ as k tends to infinity, and there exists a constant C such that for large k ,*

$$\psi \leq \psi_k + \frac{C}{k}.$$

As a reference see [21, (A4)] (note in fact the results of Demailly are in fact much stronger than that stated here, and hold in greater generality [20]). When ψ is assumed to be smooth and strictly positive, a celebrated result by Bouche–Catlin–Tian–Zelditch [10, 14, 45, 50] on Bergman kernel asymptotics implies that the ψ_k in fact converge to ψ in any C^m norm.

Using a variation of this construction, Guedj–Zeriahi prove [27, Theorem 7.1] that any positive singular metric on an ample line bundle over a compact X is the pointwise limit of a decreasing sequence of smooth positive metrics.

3.3. Monge–Ampère measures. Let ψ_i , $1 \leq i \leq n$, be an n -tuple of positive metrics, so for each i , $dd^c\psi_i$ is a positive $(1, 1)$ -current. If all ψ_i are smooth one can consider the wedge product

$$(3.1) \quad dd^c\psi_1 \wedge \dots \wedge dd^c\psi_n,$$

which is a positive measure on X . The fundamental work of Bedford–Taylor shows that one can still take the wedge product of positive currents $dd^c\psi_i$ to get a positive measure as long as the metrics ψ_i are all locally bounded. The Monge–Ampère measure of a locally bounded positive metric ψ , is then defined as the positive measure

$$MA(\psi) := (dd^c\psi)^n.$$

This measure does not put any mass on pluripolar sets. We recall the following important continuity property, proved in [3].

Theorem 3.3 (Bedford–Taylor). *If $\psi_{i,k}$, $1 \leq i \leq n+2$, $k \in \mathbb{N}$, are sequences of locally bounded positive metrics such that each $\psi_{i,k}$ decreases to a locally bounded positive metric ψ_i , then the signed measures $(\psi_{1,k} - \psi_{2,k})dd^c\psi_{3,k} \wedge \dots \wedge dd^c\psi_{n+2,k}$ converge weakly to $(\psi_1 - \psi_2)dd^c\psi_3 \wedge \dots \wedge dd^c\psi_{n+2}$. If each sequence of locally bounded positive metrics $\psi_{i,k}$ instead increase pointwise a.e. to a positive metric ψ_i , then again the measures $(\psi_{1,k} - \psi_{2,k})dd^c\psi_{3,k} \wedge \dots \wedge dd^c\psi_{n+2,k}$ converge weakly to $(\psi_1 - \psi_2)dd^c\psi_3 \wedge \dots \wedge dd^c\psi_{n+2}$.*

Since the curvature form $dd^c\phi$ of any smooth metric ϕ is a representative of $c_1(L)$, we see that if ϕ_i is any n -tuple of smooth metrics then

$$(3.2) \quad \int_X dd^c\phi_1 \wedge \dots \wedge dd^c\phi_n = \int_X c_1(L)^n$$

which is just a topological invariant of L . Since any positive metric can be approximated from above in the manner of Theorem 3.3 by positive metrics that are smooth, we see that (3.2) still holds if the ϕ_i are merely assumed to be locally bounded instead of smooth.

Our proof that maximal envelopes are maximal (Theorem 4.10) is based on an approximation argument that requires some technical results concerning convergence of plurisubharmonic functions. Recall that a plurisubharmonic function is, by definition, upper semicontinuous, so if ψ is a positive metric then for each local frame f the function ψ_f is upper semicontinuous. The plurifine topology is defined as the coarsest topology in which all local plurisubharmonic functions are continuous; a basis for this topology is given by sets of the form $A \cap \{u > 0\}$, where A is open in the standard topology and u is a local plurisubharmonic function. This topology has the quasi-Lindelöf property [4, Thm 2.7], meaning that an arbitrary union of plurifine open sets differs from a countable subunion by at most a pluripolar set. Any basis set $A \cap \{u > 0\}$ is Borel, so it follows from the quasi-Lindelöf property that the plurifine open (and closed) sets lie in the completion of the Borel σ -algebra with respect to any Monge–Ampère measure [4, Prop 3.1].

Definition 3.4. A function f is said to be *quasi-continuous* on a set Ω if for every $\epsilon > 0$ there exists an open set U with capacity less than ϵ so that f is continuous on $\Omega \setminus U$.

We refer to [3, (1.3)] for the definition of capacity, and in [3, Thm 3.5] it is shown that plurisubharmonic functions are quasi-continuous.

If f_k is a sequence of non-negative continuous functions increasing to the characteristic function of an open set A then the characteristic function of a basis set $A \cap \{u > 0\}$ is the increasing limit of the non-negative quasi-continuous functions

$$kf_k(\max\{u, 0\} - \max\{u - 1/k, 0\}).$$

From this fact and the quasi-Lindelöf property it follows that the characteristic function of any plurifine open set differs from an increasing limit of non-negative quasi-continuous functions at most on a pluripolar set.

A fundamental property of the Bedford–Taylor product is that it is local in the plurifine topology, so if $\psi_i = \psi'_i$ for all i on some plurifine open set O then

$$\mathbf{1}_O dd^c\psi_1 \wedge \dots \wedge dd^c\psi_n = \mathbf{1}_O dd^c\psi'_1 \wedge \dots \wedge dd^c\psi'_n,$$

where $\mathbf{1}_O$ denotes the characteristic function of O . We also have that the convergence in Theorem 3.3 is local in this topology [4, Thm 3.2], i.e., we get convergence when testing against bounded quasi-continuous functions.

Lemma 3.5. *Let ψ_k be a sequence of locally bounded positive metrics that decreases pointwise (or increases a.e.) to a locally bounded positive metric ψ , and let O be a plurifine open set. Then*

$$\mathbf{1}_O MA(\psi) \leq \liminf_{k \rightarrow \infty} \mathbf{1}_O MA(\psi_k),$$

where the \liminf is to be understood in the weak sense, i.e., when testing against non-negative continuous functions.

Proof. Let u_i be a sequence of quasi-continuous functions increasing to $\mathbf{1}_O$ except on a pluripolar set. Let f be a non-negative continuous function. Since $u_i MA(\psi_k)$ converges weakly to $u_i MA(\psi)$, and $MA(\psi_k)$ does not put any mass on a pluripolar set,

$$(3.3) \quad \int_X f u_i MA(\psi) = \lim_{k \rightarrow \infty} \int_X f u_i MA(\psi_k) \leq \liminf_{k \rightarrow \infty} \int_O f MA(\psi_k).$$

Now u_i increases to the characteristic function of O except possibly on a pluripolar set, so letting i tend to infinity in (3.3) yields

$$\int_O f MA(\psi) \leq \liminf_{k \rightarrow \infty} \int_O f MA(\psi_k). \quad \square$$

For singular ψ_i there is a (non-pluripolar) product constructed by Boucksom–Eyssi-dieux–Guedj–Zeriahi [13], building on a local construction due to Bedford–Taylor [4]. Fix a locally bounded metric ϕ , and consider the auxiliary metrics $\psi_{i,k} := \max\{\psi_i, \phi - k\}$ for $k \in \mathbb{N}$, and the sets $O_k := \bigcap_i \{\psi_i > \phi - k\}$. The *non-pluripolar product* of the currents $dd^c \psi_i$, here denoted by $dd^c \psi_1 \wedge \dots \wedge dd^c \psi_n$ is defined as the limit

$$dd^c \psi_1 \wedge \dots \wedge dd^c \psi_n := \lim_{k \rightarrow \infty} \mathbf{1}_{O_k} dd^c \psi_{1,k} \wedge \dots \wedge dd^c \psi_{n,k}.$$

Since we are assuming that X is compact this limit is well defined [13, Prop. 1.6]. The (non-pluripolar) Monge–Ampère measure of a positive metric ψ is defined as $MA(\psi) := (dd^c \psi)^n$. Essentially by construction, the non-pluripolar product is local in the plurifine topology [13, Prop. 1.4], and is multilinear [13, Prop 4.4].

Clearly from the definition and (3.2), for any n -tuple of positive metrics ψ_i on L ,

$$\int_X dd^c \psi_1 \wedge \dots \wedge dd^c \psi_n \leq \int_X c_1(L)^n,$$

however the inequality may well be strict.

Combining Lemma 3.5 with the fact that the Monge–Ampère measure is local in the plurifine topology yields the following continuity result.

Lemma 3.6. *Let ψ_k be a sequence of positive metrics decreasing to a positive metric ψ , and let ϕ be some locally bounded positive metric. If O is a plurifine open set contained in $\{\psi > \phi - C\}$ for some constant C then*

$$(3.4) \quad \mathbf{1}_O MA(\psi) \leq \liminf_{k \rightarrow \infty} \mathbf{1}_O MA(\psi_k),$$

where again the \liminf is to be understood in the weak sense. If ψ_k instead is increasing a.e. to ψ , and O is a plurifine open set contained in $\{\psi_j > \phi - C\}$ for some natural number j and some constant C then once again

$$\mathbf{1}_O MA(\psi) \leq \liminf_{k \rightarrow \infty} \mathbf{1}_O MA(\psi_k).$$

Proof. First assume that ψ_k is decreasing to ψ . Let $\psi'_k := \max\{\psi_k, \phi - C\}$ and $\psi' := \max\{\psi, \phi - C\}$. From Lemma 3.5 it follows that

$$\mathbf{1}_O MA(\psi') \leq \liminf_{k \rightarrow \infty} \mathbf{1}_O MA(\psi'_k),$$

and since by assumption $\psi' = \psi$ and $\psi'_k = \psi_k$ on O the lemma follows from the locality of the non-pluripolar product. The case where ψ_k is increasing a.e. follows by the same reasoning. \square

In [13, Thm 1.16] it is shown that the non-pluripolar product, when restricted to metrics with small unbounded locus, has the following monotonicity property.

Theorem 3.7. *Let ψ_i, ψ'_i be two n -tuples of positive metrics with small unbounded locus, and suppose that for all i , ψ_i is more singular than ψ'_i . Then*

$$\int_X dd^c \psi_1 \wedge \dots \wedge dd^c \psi_n \leq \int_X dd^c \psi'_1 \wedge \dots \wedge dd^c \psi'_n.$$

There is also a comparison principle for metrics with small unbounded locus [13, Cor 2.3] and a domination principle [13, Cor 2.5]. When combined with the comparison principle, the proof of the domination principle in [13] in fact yields a slightly stronger version:

Theorem 3.8. *Let ϕ be a positive metric with small unbounded locus and suppose that there exists a positive metric ρ , more singular than ϕ , with small unbounded locus and such that $MA(\rho)$ dominates a volume form. If ψ is a positive metric more singular than ϕ and such that $\psi \leq \phi$ a.e. with respect to $MA(\phi)$, then it follows that $\psi \leq \phi$ on the whole of X .*

3.4. The Aubin–Mabuchi Energy. The Aubin–Mabuchi energy bifunctional maps any pair of equivalent positive metrics ψ_1 and ψ_2 to the number

$$\mathcal{E}(\psi_1, \psi_2) := \frac{1}{n+1} \sum_{i=0}^n \int_X (\psi_1 - \psi_2) (dd^c \psi_1)^i \wedge (dd^c \psi_2)^{n-i}.$$

Observe

$$\mathcal{E}(\psi + t, \psi) = t \int_X MA(\psi).$$

The Aubin–Mabuchi energy restricted to the class of locally bounded metrics has a cocycle property (see, for example, [7, Cor 4.2]), namely if ϕ_0 , ϕ_1 and ϕ_2 are locally bounded equivalent metrics then

$$\mathcal{E}(\phi_0, \phi_2) = \mathcal{E}(\phi_0, \phi_1) + \mathcal{E}(\phi_1, \phi_2).$$

In fact, the proof in [7] of the cocycle property extends to the case where the equivalent metrics are only assumed to have small unbounded locus, since the integration-by-parts formula of [13] used in the proof holds in that case.

This leads to an important monotonicity property. If ψ_0, ψ_1 and ψ_2 are equivalent with small unbounded locus, and $\psi_0 \geq \psi_1$, then

$$\mathcal{E}(\psi_0, \psi_2) \geq \mathcal{E}(\psi_1, \psi_2)$$

since $\mathcal{E}(\psi_0, \psi_2) = \mathcal{E}(\psi_0, \psi_1) + \mathcal{E}(\psi_1, \psi_2)$, and $\mathcal{E}(\psi_0, \psi_1) \geq 0$ as it is the integral of the positive function $\psi_0 - \psi_1$ against a positive measure.

We also record the following lemma, which comes from the locality of the non-pluripolar product in the plurifine topology.

Lemma 3.9. *Let $\psi_1 \sim \psi_2$ be such that $\psi_1 \geq \psi_2$. Let ψ'_1 and ψ'_2 be two other metrics such that $\psi'_1 \sim \psi'_2$ and assume that $\{\psi'_1 = \psi'_2\} = \{\psi_1 = \psi_2\}$ and that $\psi'_1 = \psi_1$ and $\psi'_2 = \psi_2$ on the set where $\psi_1 > \psi_2$. Then*

$$\mathcal{E}(\psi'_1, \psi'_2) = \mathcal{E}(\psi_1, \psi_2).$$

Following Phong–Sturm in [34] we can relate weak geodesics to the energy functional. Let $I \subset \mathbb{R}$ be an interval and consider the annulus $A := \{w \in \mathbb{C} : -\ln |w| \in I\}$ and $\pi: X \times A \rightarrow X$ be the projection.

Definition 3.10. A curve of positive metrics ϕ_t , for $t \in I$ is said to be a *weak subgeodesic* if there exists a locally bounded positive metric Φ on π^*L that is rotation invariant and whose restriction to $X \times \{w\}$ equals $\phi_{-\ln |w|}$. The curve ϕ_t is said to be a *weak geodesic* if it is a weak subgeodesic and furthermore Φ solves the HCMA equation, i.e.,

$$MA(\Phi) = 0$$

on $X \times A^\circ$. A weak geodesic ϕ_t defined for $0 \leq t < \infty$ will be called a *weak geodesic ray*.

As in the convex setting (2.3) there is a formula [8, Proposition 6.2] relating the Aubin–Mabuchi energy of a locally bounded subgeodesic ϕ_t with the Monge–Ampère measure of Φ , namely

$$(3.5) \quad dd_t^c \mathcal{E}(\phi_t, \phi_a) = \pi_*(MA(\Phi)),$$

where $\pi_*(MA(\Phi))$ denotes the push-forward of the measure $MA(\Phi)$ with respect to the projection π . From this we immediately get the following lemma.

Lemma 3.11. *A curve ϕ_t of locally bounded positive metrics defined for $t \in [0, \infty)$ is a weak geodesic ray if and only if it is a subgeodesic and for any $a \in (0, \infty)$ the Aubin–Mabuchi energy $\mathcal{E}(\phi_t, \phi_a)$ is linear in t .*

4. Envelopes and maximal metrics

In studying the Dirichlet problem for the HCMA equation it is often possible to give a solution as an envelope in some space of plurisubharmonic functions (or positive metrics). Such envelopes will be crucial in our setting as well.

Definition 4.1. If ϕ is a continuous metric, not necessarily positive, let $P\phi$ denote the envelope

$$P\phi := \sup\{\psi \leq \phi, \psi \in PSH(L)\}.$$

Since ϕ is assumed to be continuous it follows that the upper semicontinuous regularization of $P\phi$ is bounded from above by ϕ , and hence also by $P\phi$. Thus $P\phi$ is itself upper semicontinuous and so $P\phi \in PSH(L)$.

The next theorem is essentially just a reformulation of a local result of Bedford–Taylor [3, Corollary 9.2] in our global setting. It follows as a special case of [7, Proposition 1.10] (letting $K = X$).

Theorem 4.2. *If ϕ is a continuous metric then $P\phi = \phi$ a.e. with respect to $MA(P\phi)$.*

Recall that if A is a closed set and μ is a Borel measure we say that μ is said to be *concentrated* on A if $\mathbf{1}_A\mu = \mu$, or equivalently $\mu(A^c) = 0$. Thus another way of formulating Theorem 4.2 is to say that $MA(P\phi)$ is concentrated on $\{P\phi = \phi\}$. We now extend this result to more general envelopes that arise from the additional data of singularity type.

Definition 4.3. Given a positive metric $\psi \in PSH(L)$ let P_ψ denote the projection operator on $PSH(L)$ defined by

$$P_\psi\phi := \sup\{\psi' \leq \min\{\phi, \psi\}, \psi' \in PSH(L)\}.$$

We also define $P_{[\psi]}$ by

$$P_{[\psi]}\phi := \lim_{C \rightarrow \infty} P_{\psi+C}\phi = \sup\{\psi' \leq \phi, \psi' \sim \psi, \psi' \in PSH(L)\}.$$

Clearly $P_\psi\phi$ is monotone with respect to both ψ and ϕ . Since $\min\{\phi, \psi\}$ is upper semicontinuous, it follows that the upper semicontinuous regularization of $P_\psi\phi$ is still less than $\min\{\phi, \psi\}$, and thus $P_\psi\phi \in PSH(L)$. By this it follows that $P_\psi(P_\psi\phi) = P_\psi\phi$, i.e., that P_ψ is indeed a projection operator on $PSH(L)$. One also notes that the upper semicontinuous regularization of $P_{[\psi]}\phi$, lies in $PSH(L)$ and is bounded by ϕ .

Definition 4.4. The *maximal envelope* of ϕ with respect to the singularity type $[\psi]$ is defined to be

$$\phi_{[\psi]} := \text{usc}(P_{[\psi]}\phi),$$

where usc denotes the process of taking the upper-semicontinuous regularization.

Definition 4.5. If $\psi \in PSH(L)$, then ψ is said to be *maximal* with respect to a metric ϕ if $\psi \leq \phi$ and furthermore $\psi = \phi$ a.e. with respect to $MA(\psi)$. Similarly, if A is a measurable set, we say that ψ is maximal with respect to ϕ on A if $\psi \leq \phi$ and $\psi = \phi$ a.e. on A with respect to $MA(\psi)$.

Remark 4.6. The terminology is justified by a proof below that the maximal envelope of a continuous metric ϕ is maximal with respect to ϕ . As it is defined as a limit, it is not clear from the definition if $\phi_{[\psi]}$ is equivalent to ψ (this can be shown if ψ has algebraic singularities by passing to a suitable resolution, and we refer the reader to [40] for a further study of maximal envelopes). For this reason, the method in the proof of Theorem 4.2 in [7] does not directly apply, so instead we will use an approximation argument.

Our use of the word maximal is motivated by the following property:

Proposition 4.7. *Let ψ be maximal with respect to a metric ϕ . Suppose also that there exists a positive metric $\rho \succeq \psi$ with small unbounded locus and such that $MA(\rho)$ dominates a volume form. Then for any $\psi' \sim \psi$ with $\psi \leq \phi$ we have $\psi' \leq \psi$.*

Proof. Since $\psi' \leq \phi$, the maximality assumption yields $\psi' \leq \psi$ a.e. with respect to $MA(\psi)$, so the proposition thus follows from the domination principle (Theorem 3.8). \square

The next two lemmas are the main steps in showing that maximal envelopes are maximal.

Lemma 4.8. *Let ψ_k be a sequence of positive metrics increasing a.e. to a positive metric ψ , and assume that all ψ_k are maximal with respect to a fixed continuous metric ϕ on some plurifine open set O . Then ψ is maximal with respect to ϕ on O .*

Proof. Since ϕ was assumed to be continuous, $\psi \leq \phi$. Now, for all k

$$\{\psi_k = \phi\} \subseteq \{\psi = \phi\}$$

and thus by the maximality of ψ_k , we know $\mathbf{1}_O MA(\psi_k)$ is concentrated on $\{\psi = \phi\}$. Since $\psi \leq \phi$ we have that $\{\psi = \phi\} = \{\psi \geq \phi\}$, and since ϕ is continuous this is a closed set. Let C be a constant. The set $O \cap \{\psi_1 > \phi - C\}$ is plurifinely open, so by Lemma 3.6 it follows that

$$(4.1) \quad \mathbf{1}_O \mathbf{1}_{\{\psi_1 > \phi - C\}} MA(\psi) \leq \liminf_{k \rightarrow \infty} \mathbf{1}_O \mathbf{1}_{\{\psi_1 > \phi - C\}} MA(\psi_k).$$

It is easy to see that if μ_k is a sequence of measures all concentrated on a closed set A , and

$$\mu \leq \liminf_{k \rightarrow \infty} \mu_k$$

in the weak sense, then μ is also concentrated on A . It thus follows from (4.1) that $\mathbf{1}_O \mathbf{1}_{\{\psi_1 > \phi - C\}} MA(\psi)$ is concentrated on $\{\psi = \phi\}$. Since $MA(\psi)$ puts no mass on the pluripolar set $\{\psi_1 = -\infty\}$ the lemma follows by letting C tend to infinity. \square

Lemma 4.9. *Let $\psi \in PSH(L)$ and let ϕ be a continuous metric. Then the envelope $P_\psi \phi$ is maximal with respect to ϕ on the plurifine open set $\{\psi > \phi\}$.*

Proof. Clearly $P_\psi \phi \leq \phi$, so we have to show $P_\psi \phi$ is equal to ϕ almost everywhere on $\{\psi > \phi\}$ with respect to $MA(P_\psi \phi)$. Let ϕ_k be a sequence of continuous metrics decreasing pointwise to $\min\{\phi, \psi\}$, so that $\phi_k \leq \phi$ for all k and $\phi_k = \phi$ on the set $\{\psi > \phi\}$. For example let $\phi_k := \min\{\phi, \psi_k\}$ where ψ_k is a sequence of smooth metrics decreasing pointwise to ψ . From Theorem 4.2 it follows that $MA(P\phi_k)$ is concentrated on $\{P\phi_k = \phi_k\}$, and since $\phi_k = \phi$ when $\psi > \phi$ we see $\mathbf{1}_{\{\psi > \phi\}} MA(P\phi_k)$ is concentrated on $\{P\phi_k = \phi\}$. Now $P\phi_k$ is decreasing in k and $\lim_{k \rightarrow \infty} P\phi_k \leq \min\{\phi, \psi\}$. At the same time, for any $k \in \mathbb{N}$ we clearly have that $P_\psi \phi \leq P\phi_k$, which taken together means that

$$\lim_{k \rightarrow \infty} P\phi_k = P_\psi \phi.$$

Since $P\phi_k \leq \phi$ this implies that $\{P\phi_k = \phi\}$ is decreasing in k and

$$(4.2) \quad \{P_\psi \phi = \phi\} = \bigcap_{k \in \mathbb{Z}} \{P\phi_k = \phi\}.$$

Let O denote the plurifine open set $\{\psi > \phi\} \cap \{P_\psi \phi > \phi - C\}$. By Lemma 3.6 we know

$$\mathbf{1}_O MA(P_\psi \phi) \leq \liminf_{k \rightarrow \infty} \mathbf{1}_O MA(P\phi_k),$$

and thus we conclude that $\mathbf{1}_O MA(P_\psi \phi)$ is concentrated on $\{P\phi_k = \phi\}$ for any k , so by (4.2) we get that $\mathbf{1}_O MA(P_\psi \phi)$ is concentrated on $\{P_\psi \phi = \phi\}$. Since $MA(P_\psi \phi)$ puts no mass on the pluripolar set $\{P_\psi \phi = -\infty\}$, letting C tend to infinity yields the lemma. \square

Theorem 4.10. *Let $\psi \in PSH(L)$ and let ϕ be a continuous metric. Then $\phi_{[\psi]}$ is maximal with respect to ϕ , i.e., $\phi_{[\psi]} = \phi$ a.e. with respect to $MA(\phi_{[\psi]})$.*

Proof. $P_{[\psi]} \phi = \phi_{[\psi]}$ a.e., and since $P_{\psi+C} \phi$ increases to $P_{[\psi]} \phi$, it thus increases to $\phi_{[\psi]}$ a.e. By Lemma 4.9 we get that $P_{\psi+C} \phi$ is maximal with respect to ϕ on the plurifine open set $\{\psi > \phi - C\}$ and thus also on any set $\{\psi > \phi - C'\}$ whenever $C' \leq C$. From Lemma 4.8 it thus follows that $\phi_{[\psi]}$ is maximal with respect to ϕ on the set $\{\psi > \phi - C\}$ for any C . Since $MA(\phi_{[\psi]})$ puts no mass on $\{\psi = -\infty\}$ the theorem follows. \square

Example 4.11. Consider the case that s is a section of rL that vanishes along a divisor D , and set $\psi = \frac{1}{r} \ln |s|^2$. Then the maximal envelope $\phi_{[\psi]}$ is considered by Berman [6, Section 4], and equals

$$\sup^* \{ \psi' \leq \phi : \psi' \in PSH(L), \nu_D(\psi') \geq 1 \}$$

where ν_D denotes the Lelong number along D . This metric governs the Bergman kernel asymptotics of sections of kL for $k \gg 0$ that vanish along the divisor D . The more general case when ψ has analytic singularities is also considered in [6].

The maximal property gives the following bounds on the energy functional which will be crucial for our construction of weak geodesics (Theorem 6.8).

Proposition 4.12. *Suppose that ψ is maximal with respect to a positive metric ϕ with small unbounded locus, and let $t > 0$. Then*

$$(4.3) \quad t \int_X MA(\psi) \leq \mathcal{E}(\max\{\psi + t, \phi\}, \phi) \leq t \int_X MA(\phi).$$

Proof. Since by assumption $\psi \leq \phi$ we know $\max\{\psi + t, \phi\} \leq \phi + t$, so from the monotonicity of the Aubin–Mabuchi energy it follows that

$$\mathcal{E}(\max\{\psi + t, \phi\}, \phi) \leq \mathcal{E}(\phi + t, \phi) = t \int_X MA(\phi)$$

which gives the upper bound. For the lower bound, first choose an ϵ with $0 < \epsilon < t$. Again by monotonicity,

$$(4.4) \quad \mathcal{E}(\max\{\psi + t, \phi\}, \phi) \geq \mathcal{E}(\max\{\psi + t, \phi\}, \max\{\psi + \epsilon, \phi\}).$$

Now clearly

$$(4.5) \quad \mathcal{E}(\max\{\psi + t, \phi\}, \max\{\psi + \epsilon, \phi\}) \geq (t - \epsilon) \int_{\{\psi + \epsilon > \phi\}} MA(\psi).$$

By the assumption that ψ is maximal with respect to ϕ

$$\int_{\{\psi = \phi\}} MA(\psi) = \int_X MA(\psi)$$

and since $\{\psi = \phi\} \subseteq \{\psi + \epsilon > \phi\}$, the combination of (4.4) and (4.5) yields

$$\mathcal{E}(\max\{\psi + t, \phi\}, \phi) \geq (t - \epsilon) \int_X MA(\psi).$$

Since $\epsilon > 0$ was chosen arbitrarily the lower bound in (4.3) follows. \square

5. Test curves and analytic test configurations

Definition 5.1. A map $\lambda \mapsto \psi_\lambda$ from \mathbb{R} to $PSH(L)$ is called a *test curve* if there is a constant C such that

- (i) ψ_λ is equal to some locally bounded positive metric $\psi_{-\infty}$ for $\lambda < -C$,
- (ii) $\psi_\lambda \equiv -\infty$ for $\lambda > C$,
- (iii) ψ_λ has small unbounded locus whenever $\psi_\lambda \not\equiv -\infty$ and
- (iiii) ψ_λ is concave in λ .

Observe that since ψ_λ is concave and constant for λ sufficiently negative it is decreasing in λ . The set of test curves forms a convex set, by letting

$$\left(\sum r_i \gamma_i\right)(\lambda) := \sum r_i \gamma_i(\lambda).$$

It is also clear that any translate $\gamma_a(\lambda) := \gamma(\lambda - a)$ of a test curve γ is a new test curve.

We introduce the notation λ_c for the critical value of a test curve defined as

$$\lambda_c := \inf\{\lambda : \psi_\lambda \equiv -\infty\}.$$

For later use, we record here two continuity properties of test curves.

Lemma 5.2.

- (1) A test curve ψ_λ is left-continuous in λ as long as $\lambda < \lambda_c$.
- (2) Suppose that $\lambda < \lambda_c$ and λ_k is a decreasing sequence that tends to λ .
Then

$$(5.1) \quad \lim_{k \rightarrow \infty}^* \psi_{\lambda_k} = \psi_\lambda.$$

(So a test curve is right continuous modulo taking an upper semicontinuous regularization.)

Proof. For (1), let λ_k increase to some $\lambda < \lambda_c$, so we need to show that

$$\lim_{k \rightarrow \infty} \psi_{\lambda_k} = \psi_\lambda.$$

By our hypothesis there exists a λ' such that $\lambda < \lambda' < \lambda_c$, and thus $\psi_{\lambda'} \not\equiv -\infty$. Since $\psi_\lambda(x)$ is concave in λ it is continuous for all x such that $\psi_{\lambda'}(x) \not\equiv -\infty$. Thus, ψ_{λ_k} converges to ψ_λ at least away from a pluripolar set, i.e., a.e. with respect to a volume form. On the other hand we have that ψ_{λ_k} is decreasing in k , so the limit is a positive metric. Now if two positive metrics coincide a.e. with respect to a volume form it follows that they are equal everywhere, because this is true locally for plurisubharmonic function [29, Corollary 2.9.8].

The proof of (2) is essentially the same. This time λ_k is a decreasing sequence, so as $\lambda < \lambda_c$ we may as well assume that each $\lambda_k < \lambda'$ and so in particular $\psi_{\lambda_k} \not\equiv -\infty$. Then the ψ_{λ_k} form an increasing sequence so the left hand side of (5.1) is a positive metric. But for the same reason as above,

the limit $\lim_{k \rightarrow \infty} \psi_{\lambda_k}$ equals ψ_λ away from a pluripolar set, and thus the left and right hand side of (5.1) agree a.e. with respect to a volume form, and thus are equal everywhere. \square

Definition 5.3. A map γ from \mathbb{R} to $Sing(L)$ is called an *analytic test configuration* if it is the composition of a test curve with the natural projection of $PSH(L)$ to $Sing(L)$.

As with the set of test curves, the set of analytic test configurations is convex. We now extend the definition of the maximal envelope (Definition 4.4) to test curves.

Definition 5.4. Let ψ_λ be a test curve and ϕ an element in $\mathcal{H}(L)$. The *maximal envelope* of ϕ with respect to ψ_λ is the map

$$\lambda \mapsto \phi_\lambda := \phi_{[\psi_\lambda]} = \text{usc}(P_{[\psi_\lambda]}\phi).$$

It is easy to see that ϕ_λ only depends on ϕ and the analytic test configuration $[\psi_\lambda]$, since if $\psi'_\lambda \sim \psi_\lambda$ we trivially have $\phi_{[\psi_\lambda]} = \phi_{[\psi'_\lambda]}$. Observe also that since $\psi_{-\infty}$ is locally bounded, we have $\phi_\lambda = \phi$ for $\lambda < -C$.

Definition 5.5. We say that a test curve ψ_λ is *maximal* if for all λ the metric ψ_λ is maximal with respect to $\psi_{-\infty}$.

Since ψ_λ is decreasing in λ ,

$$\{\psi_{\lambda'} = \psi_\lambda\} \supseteq \{\psi_{\lambda'} = \psi_{-\infty}\} \quad \text{if } \lambda \leq \lambda'.$$

It follows that if ψ_λ is a maximal test curve, $\psi_{\lambda'}$ is maximal with respect to ψ_λ whenever $\lambda \leq \lambda'$. We shall show in the next section how the Legendre transform of a maximal test curve gives a weak geodesic ray, and end this section by showing how maximal envelopes give rise to maximal test curves.

Proposition 5.6. *The maximal envelope ϕ_λ is a maximal test curve.*

Proof. We first show it is a test curve. Pick a real number C . Let λ and λ' be two real numbers, and let $0 \leq t \leq 1$. By the concavity of ψ_λ ,

$$tP_{\psi_\lambda+C}\phi + (1-t)P_{\psi_{\lambda'}+C}\phi \leq t\psi_\lambda + (1-t)\psi_{\lambda'} + C \leq \psi_{t\lambda+(1-t)\lambda'} + C.$$

Thus from the definition of the projection operator,

$$tP_{\psi_\lambda+C}\phi + (1-t)P_{\psi_{\lambda'}+C}\phi \leq P_{\psi_{t\lambda+(1-t)\lambda'}+C}\phi,$$

which means that $P_{\psi_\lambda+C}\phi$ is concave in λ for all C . Since $P_{\psi_\lambda+C}\phi$ increases to $P_{[\psi_\lambda]}\phi$, and an increasing sequence of concave functions is concave, we get that $P_{[\psi_\lambda]}\phi$ is concave, and because of the monotonicity of the upper semi-continuous regularization it follows that $\text{usc}(P_{[\psi_\lambda]}\phi) = \phi_\lambda$ also is concave. The other properties of a test curve are immediate.

Clearly $\phi_{-\infty} = \phi$, so that ϕ_λ is maximal follows from Theorem 4.10. \square

6. The Legendre transform and geodesic rays

If f is a convex function in the real variable λ , the set of subgradients of f , denoted by Δ_f , is the set of $t \in \mathbb{R}$ such that $f(\lambda) - t\lambda$ is bounded from below. If f happens to be differentiable, then the set subgradients coincides with the image of the derivative of f . By convexity of f , the set of subgradients is convex, i.e., an interval. Recall that the Legendre transform of f , here denoted by \widehat{f} , is the function on Δ_f defined as

$$\widehat{f}(t) := \sup_{\lambda} \{t\lambda - f(\lambda)\}.$$

Since \widehat{f} is defined as the supremum of the linear functions $t\lambda - f(\lambda)$, it follows that \widehat{f} is convex.

If f is concave then of course $-f$ is convex, and one can define the Legendre transform of f , also denoted by \widehat{f} , as the Legendre transform of $-f$, i.e.,

$$\widehat{f}(t) := \sup_{\lambda} \{f(\lambda) + t\lambda\},$$

which is thus convex.

Definition 6.1. The *Legendre transform* of a test curve ψ_{λ} is

$$\widehat{\psi}_t := \sup_{\lambda \in \mathbb{R}}^* \{\psi_{\lambda} + t\lambda\},$$

where $t \in [0, \infty)$.

Recall that the star denotes the operation of taking the upper semicontinuous regularization.

Lemma 6.2. *Let ψ_{λ} be any test curve (not necessarily maximal). Then the Legendre transform $\widehat{\psi}_t$ is locally bounded for all t , and the map $t \mapsto \widehat{\psi}_t$ is a subgeodesic ray emanating from $\psi_{-\infty}$.*

Proof. By assumption, for some λ , ψ_{λ} is locally bounded, and trivially $\widehat{\psi}_t \geq \psi_{\lambda} + t\lambda$, thus $\widehat{\psi}_t$ is locally bounded. It is clear that for a fixed λ , the curve $\psi_{\lambda} + t\lambda$ is a subgeodesic. Clearly $\sup_{\lambda \in \mathbb{R}} \{\psi_{\lambda} + t\lambda\}$ is convex and Lipschitz in t , and the same is easily seen to hold for $\widehat{\psi}_t$. Thus $\widehat{\psi}_t$ is upper semicontinuous in the directions in X and also Lipschitz in t , which implies that it is upper semicontinuous on the product $X \times \mathbb{R}_{\geq 0}$. Therefore $\widehat{\psi}_t$ (thought of as a function on the product) coincides with the upper semicontinuous regularization of $\sup_{\lambda \in \mathbb{R}} \{\psi_{\lambda} + t\lambda\}$.

Now, taking the upper semicontinuous regularization of the supremum of subgeodesics yields a subgeodesic, as long as it is bounded from above. We observed above that $\psi_{\lambda} \leq \psi_{-\infty}$. Now for some constant C , $\psi_C \equiv -\infty$. It follows that $\widehat{\psi}_t \leq \psi_{-\infty} + tC$, so it is bounded from above and thus it is a subgeodesic.

Finally by definition $\widehat{\psi}_0 = \sup_\lambda^* \{\psi_\lambda\}$, which clearly is equal to $\psi_{-\infty}$ since $\psi_\lambda \leq \psi_{-\infty}$ (ψ_λ being decreasing in λ) and $\psi_{-\infty}$ itself being upper-semicontinuous. \square

One can also consider the inverse Legendre transform, going from subgeodesic rays to concave curves of positive metrics.

Definition 6.3. The Legendre transform of a subgeodesic ray ϕ_t , $t \in [0, \infty)$, denoted by $\widehat{\phi}_\lambda$, $\lambda \in \mathbb{R}$, is defined as

$$\widehat{\phi}_\lambda := \inf_{t \in [0, \infty)} \{\phi_t - t\lambda\}.$$

Remark 6.4. It follows from Kiselman's minimum principle (see [28]) that for any $\lambda \in \mathbb{R}$, $\widehat{\phi}_\lambda$ is a positive metric (we would like to thank Bo Berndtsson for this observation). Furthermore, it is clear that $\widehat{\phi}_\lambda$ is concave and decreasing in λ . From the involution property of the (real) Legendre transform it follows that the Legendre transform of $\widehat{\phi}_\lambda$ is ϕ_t , thus any subgeodesic ray is the Legendre transform of a concave curve of positive metrics.

The goal of this section is to prove that if ψ_λ is an *maximal* test curve then the Legendre transform $\widehat{\psi}_t$ of ψ_λ is a weak geodesic ray emanating from $\psi_{-\infty}$. By Lemma 6.2, we know $\widehat{\psi}_t$ is a subgeodesic ray emanating from $\psi_{-\infty}$. What remains then is to show that if ψ_λ is maximal then the Aubin–Mabuchi energy $\mathcal{E}(\widehat{\psi}_t, \widehat{\psi}_0)$ is linear in t , which we now do with an approximation argument.

For $N \in \mathbb{N}$ consider the approximation $\widehat{\psi}_t^N$ to $\widehat{\psi}_t$, given by

$$\widehat{\psi}_t^N := \sup_{k \in \mathbb{Z}} \{\psi_{k2^{-N}} + tk2^{-N}\}.$$

Since ψ_λ is concave it is continuous in λ at all points such that $\psi_\lambda(x) > -\infty$. From the continuity it follows that $\widehat{\psi}_t^N$ will increase pointwise to $\widehat{\psi}_t$ a.e. as N tends to infinity. Also let $\widehat{\psi}_t^{N,M}$ denote the curve

$$\widehat{\psi}_t^{N,M} := \sup_{k \in \mathbb{Z}, k \leq M} \{\psi_{k2^{-N}} + tk2^{-N}\},$$

and observe that $\widehat{\psi}_t^N$ and $\widehat{\psi}_t^{N,M}$ are all locally bounded.

Lemma 6.5. *Let $M < M'$ be two integers. Then*

$$\widehat{\psi}_t^{N,M'} = \psi_{M'2^{-N}} + tM'2^{-N}$$

implies that

$$\widehat{\psi}_t^{N,M} = \psi_{M2^{-N}} + tM2^{-N}.$$

Proof. Certainly $f(\lambda) := \psi_\lambda(x) + t\lambda$ is concave in λ . If

$$\widehat{\psi}_t^{N,M} > \psi_{M2^{-N}} + tM2^{-N}$$

at x , then f would be strictly decreasing at $\lambda = M2^{-N}$, so by concavity we would get that $f(M'2^{-N}) < f(M2^{-N}) < \widehat{\psi}_t^{N,M}(x)$, which would be a contradiction. \square

Lemma 6.6. *If ψ_λ is a maximal test curve then*

$$t2^{-N} \int_X MA(\psi_{(M+1)2^{-N}}) \leq \mathcal{E}(\widehat{\psi}_t^{N,M+1}, \widehat{\psi}_t^{N,M}) \leq t2^{-N} \int_X MA(\psi_{M2^{-N}}).$$

Proof. By Lemma 6.5 it follows that $\widehat{\psi}_t^{N,M} = \psi_{M2^{-N}} + tM2^{-N}$ on the support of $\widehat{\psi}_t^{N,M+1} - \widehat{\psi}_t^{N,M}$ and thus Lemma 3.9 yields

$$(6.1) \quad \mathcal{E}(\widehat{\psi}_t^{N,M+1}, \widehat{\psi}_t^{N,M}) = \mathcal{E}(\max\{\psi_{M2^{-N}}, \psi_{(M+1)2^{-N}} + t2^{-N}\}, \psi_{M2^{-N}}).$$

Since we assumed that ψ_λ is maximal, $\psi_{(M+1)2^{-N}}$ is maximal with respect to $\psi_{M2^{-N}}$, and thus the lemma follows immediately from Lemma 4.12. \square

Let ψ_λ be a maximal test curve, and let $F(\lambda)$ denote the function

$$F(\lambda) := \int_X MA(\psi_\lambda).$$

Whenever $\lambda < \lambda'$, $\psi_{\lambda'} \leq \psi_\lambda$ and therefore it follows from Theorem 3.7 that $F(\lambda)$ is decreasing in λ , hence $F(\lambda)$ is Riemann integrable.

Proposition 6.7. *If ψ_λ is a maximal test curve then*

$$(6.2) \quad \mathcal{E}(\widehat{\psi}_t, \widehat{\psi}_0) = -t \int_{\lambda=-\infty}^{\infty} \lambda dF(\lambda).$$

Proof. Suppose first $m \in \mathbb{Z}$ is such that $\psi_m = \psi_{-\infty}$. For a given $N \in \mathbb{N}$ set $M = m2^N$. Then

$$\widehat{\psi}_t^{N,M} = \psi_{-\infty} + tm = \widehat{\psi}_0 + tm.$$

By repeatedly using the cocycle property of the Aubin–Mabuchi energy in combination with Lemma 6.6 we get that

$$(6.3) \quad t \sum_{k>M} 2^{-N} F((k+1)2^{-N}) \leq \mathcal{E}(\widehat{\psi}_t^N, \widehat{\psi}_t^{N,M}) \leq t \sum_{k>M} 2^{-N} F(k2^{-N}).$$

We noted above that $\widehat{\psi}_t^N$ increases pointwise to $\widehat{\psi}_t$ a.e. as N tends to infinity. By the continuity of the Aubin–Mabuchi energy under a.e. pointwise increasing sequences (3.6),

$$\mathcal{E}(\widehat{\psi}_t, \widehat{\psi}_0 + tm) = t \int_{\lambda=m}^{\infty} \lambda F(\lambda) d\lambda,$$

since both the left- and the right-hand side of (6.3) converges to this. Again using the cocycle property we get that

$$(6.4) \quad \begin{aligned} \mathcal{E}(\widehat{\psi}_t, \widehat{\psi}_0) &= \mathcal{E}(\widehat{\psi}_t, \widehat{\psi}_0 + tm) + \mathcal{E}(\widehat{\psi}_0 + tm, \widehat{\psi}_0) = \\ &= t \int_{\lambda=m}^{\infty} \lambda F(\lambda) d\lambda + tm \int_X MA(\psi_{-\infty}) = t \int_{\lambda=m}^{\infty} F(\lambda) d\lambda + tmF(m). \end{aligned}$$

Since by our assumption the measure dF is zero on $(-\infty, m)$, integration by parts yields

$$\begin{aligned} -t \int_{\lambda=-\infty}^{\infty} \lambda dF(\lambda) &= -t\lambda F(\lambda)|_m^{\infty} + t \int_{\lambda=m}^{\infty} F(\lambda) d\lambda \\ (6.5) \qquad \qquad \qquad &= tmF(m) + t \int_{\lambda=m}^{\infty} F(\lambda) d\lambda. \end{aligned}$$

The proposition follows from combining equation (6.4) and equation (6.5). \square

Theorem 6.8. *The Legendre transform $\widehat{\psi}_t$ of a maximal test curve ψ_λ is a weak geodesic ray emanating from $\psi_{-\infty}$.*

Proof. That $\widehat{\psi}_t$ is a subgeodesic emanating from $\psi_{-\infty}$ was proved in Lemma 6.2. According to Proposition 6.7 the energy $\mathcal{E}(\widehat{\psi}_t, \widehat{\psi}_0)$ is linear in t , and therefore by Lemma 3.11 we get that $\widehat{\psi}_t$ is a weak geodesic ray. \square

These weak geodesics are continuous in ϕ in the following sense:

Proposition 6.9. *Let ψ_λ be a test curve and $\phi, \phi' \in \mathcal{H}(L)$. Suppose ϕ_λ is the maximal curve of ϕ (with respect to ψ_λ) and similarly for ϕ'_λ . If $\|\phi - \phi'\|_\infty < C$ then*

$$\|\widehat{\phi}_t - \widehat{\phi}'_t\|_\infty < C \quad \text{for all } t.$$

Proof. We claim that $\|\phi_\lambda - \phi'_\lambda\|_\infty < C$ for all λ . But this is clear since $\phi \leq \phi'$ implies that $\phi_\lambda \leq \phi'_\lambda$ for all λ . It is also clear that $(\phi + C)_\lambda = \phi_\lambda + C$ when C is a constant. Thus $\widehat{\phi}_t \leq \widehat{\phi}'_t$ for all t and $(\phi + C)_\lambda = \phi_\lambda + C$, so consequently $\widehat{\phi + C}_t = \widehat{\phi}_t + C$ which proves the lemma. \square

Let $[\psi_\lambda]$ be an analytic test configuration, and let ϕ_λ be the associated maximal test curve. Then $[\phi_\lambda]$ defines a new analytic test configuration. This could possibly differ from $[\psi_\lambda]$, but the following proposition tells us that the associated geodesic rays are the same.

Proposition 6.10. *Let $\phi' \in \mathcal{H}(L)$. Then the Legendre transform of $\phi'_{[\phi_\lambda]}$ coincides with the Legendre transform of $\phi'_\lambda := \phi'_{[\psi_\lambda]}$.*

Proof. Since $\phi'_\lambda \sim \phi_\lambda$ we get that $\phi'_{[\phi_\lambda]} = \phi'_{[\phi'_\lambda]}$, thus without loss of generality we can assume that $\phi' = \phi$. Recall that the critical value λ_c was defined as

$$\lambda_c := \inf\{\lambda : \phi_\lambda \equiv -\infty\}.$$

If $\lambda < \lambda_c$ there exists a λ' such that $\lambda < \lambda' < \lambda_c$, and thus by the assumption $\phi_{\lambda'}$ has small unbounded locus. Let C be a constant less than λ such that $\phi_C = \phi$. By concavity it follows that

$$\phi_\lambda \geq r\phi + (1-r)\phi_{\lambda'},$$

where $0 < r < 1$, is chosen such that

$$\lambda = rC + (1 - r)\lambda'.$$

If we let

$$\rho := r\phi + (1 - r)\phi_{\lambda'},$$

by the multilinearity of the Monge–Ampère operator it follows that $MA(\rho)$ dominates the volume form $r^n MA(\phi)$. Furthermore ρ has small unbounded locus and is more singular than ϕ_{λ} . Thus, by Proposition 4.7 we have

$$P_{\phi_{\lambda} + C\phi} \leq \phi_{\lambda}$$

for any constant C and therefore

$$(6.6) \quad \phi_{[\phi_{\lambda}]} = \phi_{\lambda},$$

whenever $\lambda < \lambda_c$. If $\lambda > \lambda_c$ then clearly equation (6.6) holds as well since both sides are identically equal to minus infinity. It follows that for any $\epsilon > 0$,

$$\phi_{\lambda} \leq \phi_{[\phi_{\lambda}]} \leq \phi_{\lambda - \epsilon},$$

which implies that

$$\widehat{(\phi_{\lambda})}_t \leq \widehat{(\phi_{[\phi_{\lambda}]})}_t \leq \widehat{(\phi_{\lambda - \epsilon})}_t = \widehat{(\phi_{\lambda})}_t + \epsilon t.$$

Since $\epsilon > 0$ was arbitrary the proposition follows. \square

7. Filtrations of the ring of sections

First, we recall what is meant by a filtration of a graded algebra.

Definition 7.1. A *filtration* \mathcal{F} of a graded algebra $\bigoplus_k V_k$ is a vector space-valued map from $\mathbb{R} \times \mathbb{N}$,

$$\mathcal{F} : (t, k) \longmapsto \mathcal{F}_t V_k,$$

such that for any k , $\mathcal{F}_t V_k$ is a family of subspaces of V_k that is decreasing and left-continuous in t .

In [12] Boucksom–Chen consider certain filtrations which behaves well with respect to the multiplicative structure of the algebra. They give the following definition.

Definition 7.2. Let \mathcal{F} be a filtration of a graded algebra $\bigoplus_k V_k$. We shall say that

(i) \mathcal{F} is *multiplicative* if

$$(\mathcal{F}_t V_k)(\mathcal{F}_s V_m) \subseteq \mathcal{F}_{t+s} V_{k+m}$$

for all $k, m \in \mathbb{N}$ and $s, t \in \mathbb{R}$.

(ii) \mathcal{F} is *(linearly) bounded* if there exists a constant C such that $\mathcal{F}_{-kC} V_k = V_k$ and $\mathcal{F}_{kC} V_k = \{0\}$ for all k .

The goal in this section is to associate an “analytic test configuration” $\phi_\lambda^{\mathcal{F}}$ to any bounded multiplicative filtration of the section ring $R(L) = \bigoplus_k H^0(kL)$, which will be used in the next section to construct an associated geodesic.

Example 7.3. An important example for our purpose will be the filtrations constructed from an algebraic test configuration (Section 8). Another kind of example to have in mind come when (X, L) is toric with moment polytope Δ . Let $f: \Delta \rightarrow \mathbb{R}$ be a bounded positive concave function. Recalling that $\Delta \cap k^{-1}\mathbb{Z}^n$ parametrizes a toric basis $\{s_\alpha^{(k)}\}$ for $H^0(kL)$ define

$$\mathcal{F}_t H^0(kL) = \text{linspan}\{s_\alpha^{(k)} : f(\alpha) \geq k^{-1}t\} \subset H^0(kL).$$

It is easy to check that the concavity of f implies that \mathcal{F} is multiplicative and that f is bounded implies that \mathcal{F} is linearly bounded. When f is rational piecewise linear it turns out that this is precisely the filtration associated to the algebraic test configuration defined by f as considered by Donaldson [24] (see also [38]). In this way, one sees how the analytic test configurations and associated geodesics considered in this section and the next generalize the algebraic picture.

Now fix $\phi \in \mathcal{H}(L)$, and let dV be a smooth volume form on X with unit mass. This gives the L^2 -scalar product on $H^0(kL)$ by letting

$$(s, t)_{k\phi} := \int_X s(z) \overline{t(z)} e^{-k\phi(z)} dV(z).$$

For any $\lambda \in \mathbb{R}$ let $\{s_{i,\lambda}\}$ be an orthonormal basis for $\mathcal{F}_{k\lambda} H^0(kL)$ and define

$$\phi_{k,\lambda} := \frac{1}{k} \ln \left(\sum_i |s_{i,\lambda}|^2 \right),$$

which is a positive metric on L .

Lemma 7.4. *For any λ , the sequence of metrics $\phi_{k,\lambda}$ converges to a limit as k tends to infinity, and the upper semicontinuous regularization of the limit*

$$\phi_\lambda^{\mathcal{F}} := \lim_{k \rightarrow \infty}^* \phi_{k,\lambda}$$

is a positive metric.

Proof. Since

$$K_\lambda(z, w) := \sum_i s_{i,\lambda}(z) \overline{s_{i,\lambda}(w)}$$

is a reproducing kernel of $\mathcal{F}_{k\lambda} H^0(kL)$ with respect to $(\cdot, \cdot)_{k\phi}$, as for the full Bergman kernel we have the following useful characterization (see, for example, [5, (4.3)])

$$(7.1) \quad \sum |s_{i,\lambda}|^2 = \sup\{|s|^2 : s \in \mathcal{F}_{k\lambda} H^0(kL), \|s\|_{k\phi}^2 \leq 1\}.$$

Let $\|s\|_\infty^2 := \sup_{z \in X} \{|s(z)|^2 e^{-k\phi}\}$ and define

$$F_{k,\lambda}(z) := \sup\{|s(z)|^2 : s \in \mathcal{F}_{k\lambda} H^0(kL), \|s\|_\infty^2 \leq 1\}.$$

We trivially have the upper bound

$$F_{k,\lambda}(z) \leq e^{-k\phi(z)}.$$

It follows that

$$\text{usc} \left(\frac{1}{k} \ln F_{k,\lambda} \right) = \sup^* \left\{ \frac{1}{k} \ln |s|^2 : s \in \mathcal{F}_{k\lambda} H^0(kL), \|s\|_\infty^2 \leq 1 \right\}$$

is a positive metric. Let λ be fixed, pick a point $z \in X$, and let for all k , $s_k \in \mathcal{F}_{k\lambda} H^0(kL)$ be such that $\|s_k\|_\infty = 1$ and

$$F_{k,\lambda}(z) = |s_k(z)|^2.$$

Since the product $s_k s_m$ lies in $\mathcal{F}_{(k+m)\lambda} H^0((k+m)L)$ by the multiplicativity of \mathcal{F} , and $\|s_k s_m\|_\infty \leq \|s_k\|_\infty \|s_m\|_\infty$, we get that

$$(7.2) \quad F_{k+m,\lambda}(z) \geq F_{k,\lambda}(z) F_{m,\lambda}(z),$$

so the map $k \mapsto F_{k,\lambda}(z)$ is multiplicative. The existence of a limit

$$\lim_{k \rightarrow \infty} \frac{1}{k} \ln F_{k,\lambda}(z)$$

thus follows from Fekete's lemma (see e.g., [9, p37]). Since we assumed that dV had unit mass, for any section s

$$\|s\|_{k\phi}^2 \leq \|s\|_\infty^2,$$

and thus by equation (7.1)

$$\sum |s_{i,\lambda}(z)|^2 \geq F_{k,\lambda}(z).$$

On the other hand, by the Bernstein–Markov property of any volume form dV we have that for any $\epsilon > 0$ there exists a constant C_ϵ so that

$$\|s\|_\infty^2 \leq C_\epsilon e^{\epsilon k} \|s\|_{k\phi}^2,$$

and thus

$$(7.3) \quad \sum |s_{i,\lambda}(z)|^2 \leq C_\epsilon e^{\epsilon k} F_{k,\lambda}(z).$$

It follows that the difference $\phi_{k,\lambda}(z) - \frac{1}{k} \ln F_{k,\lambda}(z)$ tends to zero as k tends to infinity, thus the convergence of $\phi_{k,\lambda}$ follows.

By the multiplicativity, for any $k \in \mathbb{N}$

$$\frac{1}{k} \ln F_{k,\lambda} \leq \lim_{l \rightarrow \infty} \frac{1}{l} \ln F_{l,\lambda} = \lim_{l \rightarrow \infty} \phi_{l,\lambda},$$

and thus

$$(7.4) \quad \text{usc} \left(\frac{1}{k} \ln F_{k,\lambda} \right) \leq \lim_{l \rightarrow \infty}^* \phi_{l,\lambda} =: \phi_\lambda^{\mathcal{F}}.$$

On the other hand, clearly

$$\lim_{l \rightarrow \infty} \phi_{l,\lambda} \leq \sup_k \left\{ \text{usc} \left(\frac{1}{k} \ln F_{k,\lambda} \right) \right\},$$

and it follows that

$$\phi_\lambda^{\mathcal{F}} = \sup_k^* \left\{ \text{usc} \left(\frac{1}{k} \ln F_{k,\lambda} \right) \right\}$$

so $\phi_\lambda^{\mathcal{F}}$ is indeed a positive metric. \square

Remark 7.5. Since all volume forms dV on X are equivalent, the limit ϕ_λ does not depend on the choice of volume form dV .

Lemma 7.6. *We have that*

$$\phi_{k,\lambda} \leq \phi_\lambda^{\mathcal{F}} + \epsilon(k),$$

where $\epsilon(k)$ is a constant independent of λ that tends to zero as k tends to infinity.

Proof. By combining the inequalities (7.3) and (7.4) from the proof of the previous lemma we see that for any $\epsilon > 0$ there exists a constant C_ϵ independent of λ such that

$$\phi_{k,\lambda} \leq \phi_\lambda^{\mathcal{F}} + \epsilon + (1/k) \ln C_\epsilon.$$

This yields the lemma. \square

Proposition 7.7. *The map $\lambda \mapsto \phi_\lambda^{\mathcal{F}}$ is a test curve.*

Proof. Let λ be such that $F_{k\lambda}H^0(kL) = H^0(kL)$ for all k . Then $\phi_{k,\lambda}$ is the usual Bergman metric, and by the result on Bergman kernel asymptotics due to Bouche–Catlin–Tian–Zelditch (see Section 3) we get that $\phi_{k,\lambda}$ converges to ϕ . Trivially we see that if $F_{k\lambda}H^0(kL) = \{0\}$ for all k then $\phi_\lambda^{\mathcal{F}} \equiv -\infty$. By the boundedness of the filtration we thus have $\phi_\lambda^{\mathcal{F}} = \phi$ for $\lambda < -C$ and $\phi_\lambda^{\mathcal{F}} \equiv -\infty$ for $\lambda > C$.

By the multiplicativity of the filtration $\phi_\lambda \equiv -\infty$ if and only if for all k ,

$$\mathcal{F}_{k\lambda}H^0(kL) = \{0\}.$$

Pick a λ such that $\phi_\lambda^{\mathcal{F}} \neq -\infty$, then for some k , $\mathcal{F}_{k\lambda}H^0(kL)$ is non-trivial. From Lemma 7.6 it follows that $\phi_\lambda^{\mathcal{F}}$ has small unbounded locus since $\phi_{k,\lambda}$ has small unbounded locus.

It remains to prove concavity. Let $\lambda_1, \lambda_2 \in \mathbb{R}$ and let t be a rational point in the unit interval. Let m be a natural number such that mt is an integer. Given a point $z \in X$, let $s_1 \in \mathcal{F}_{k\lambda_1}H^0(kL)$ and $s_2 \in \mathcal{F}_{k\lambda_2}H^0(kL)$ be two sections with $\|s_1\|_\infty = \|s_2\|_\infty = 1$ such that

$$F_{k,\lambda_1} = |s_1(z)|^2$$

and

$$F_{k,\lambda_2} = |s_2(z)|^2.$$

By the multiplicativity of the filtration,

$$s_1^{mt} s_2^{m(1-t)} \in F_{mk(t\lambda_1+(1-t)\lambda_2)} H^0(mkL),$$

and trivially $\|s_1^{mt} s_2^{m(1-t)}\|_\infty \leq 1$. It follows that

$$F_{mk,t\lambda_1+(1-t)\lambda_2}(z) \geq F_{k,\lambda_1}(z)^{mt} F_{k,\lambda_2}(z)^{m(1-t)}.$$

Taking the logarithm on both sides, dividing by mk , and taking the limit yields

$$(7.5) \quad \phi_{t\lambda_1+(1-t)\lambda_2}^{\mathcal{F}} \geq t\phi_{\lambda_1}^{\mathcal{F}} + (1-t)\phi_{\lambda_2}^{\mathcal{F}}$$

except possibly on the pluripolar set where the limits are not equal to their upper semicontinuous regularization.

Now if two plurisubharmonic functions ζ_1 and ζ_2 are equal almost everywhere then they are equal everywhere [29, 2.9.8]. Applying this to ζ_1 and $\max\{\zeta_1, \zeta_2\}$ we see that if $\zeta_1 \geq \zeta_2$ almost everywhere then this is true everywhere. Thus in fact (7.5) holds on the whole of X .

Recall that t was assumed to be rational. If $\lambda_1 \leq \lambda_2$, the left-hand side of (7.5) is decreasing in t since clearly $\phi_\lambda^{\mathcal{F}}$ is decreasing in λ . The right-hand side of (7.5) is continuous in t , so it follows that the equation (7.5) holds for all $t \in (0, 1)$, i.e., $\phi_\lambda^{\mathcal{F}}$ is concave in λ . \square

Lemma 7.8. *For any two $\phi, \psi \in \mathcal{H}(L)$ and any $\lambda \in \mathbb{R}$ we have $\phi_\lambda^{\mathcal{F}} \sim \psi_\lambda^{\mathcal{F}}$.*

Proof. If $\phi \leq \psi$ then for all k and λ we have that $\phi_{k,\lambda} \leq \psi_{k,\lambda}$, and so $\phi_\lambda^{\mathcal{F}} \leq \psi_\lambda^{\mathcal{F}}$. Also it is clear that $(\phi + C)_{k,\lambda} = \phi_{k,\lambda} + C$, which proves the lemma. \square

Definition 7.9. We call the map $\lambda \mapsto [\phi_\lambda^{\mathcal{F}}]$ the *analytic test configuration associated to the filtration \mathcal{F}* .

So by the previous lemma this analytic test configuration depends only on \mathcal{F} and not on the choice of $\phi \in \mathcal{H}(L)$. Our next goal is to show the curve $\phi_\lambda^{\mathcal{F}}$ is maximal for $\lambda < \lambda_c$, for which we will need a Skoda-type division theorem.

Theorem 7.10. *Let L be an ample line bundle. Assume that L has a smooth positive metric ϕ with the property that $dd^c\phi \geq dd^c\phi_{K_X}$ for some smooth metric ϕ_{K_X} on the canonical bundle K_X . Let $\{s_i\}$ be a finite collection of holomorphic sections of L and $m > n + 2$ where $n = \dim X$.*

Suppose s is a section of mL such that

$$\int_X \frac{|s|^2}{(\sum |s_i|^2)^m} dV < \infty.$$

Then there exists sections $h_\alpha \in H^0((n+1)L)$ such that

$$s = \sum_{\alpha} h_{\alpha} s^{\alpha},$$

where α is a multiindex $\alpha = (\alpha_i)$ with $\sum_i \alpha_i = m - n - 1$, and s^α are the monomials $s^\alpha := \prod_i s_i^{\alpha_i}$.

Proof. Let k be an integer such that $n + 2 \leq k \leq m$. Then given a section $t \in H^0(kL)$ with

$$\int_X \frac{|t|^2}{(\sum_i |s_i|^2)^k} dV < \infty$$

an application of the Skoda division theorem [47, Thm. 2.1] yields sections $\{t_i\}$ of $(k-1)L$ such that $t = \sum_i t_i s_i$ and

$$\int_X \frac{|t_i|^2}{(\sum_i |s_i|^2)^{k-1}} dV < \infty.$$

(To apply the cited theorem replace F, E, ψ, η with $kL - K_X, L, k\phi - \phi_{K_X}, \phi$ respectively and replace αq with $k - 1 > n + 1$.)

Now we first apply the above with $k = m$ to the section s , and then apply again with $k = m - 1$ to each of the sections t_i . Repeating this process with $k = m, m - 1, \dots, n + 2$ we see that s can be written as a linear sum of monomials in the s_i as required. \square

Proposition 7.11. *For λ less than the critical value λ_c ,*

$$\phi_{\lambda}^{\mathcal{F}} = \lim_{k \rightarrow \infty} \phi_{[\phi_k, \lambda]}.$$

Proof. Let $\phi_k := \phi_{k, -\infty}$, i.e., the Bergman metric $1/k \ln(\sum |s_i|^2)$, where $\{s_i\}$ is an orthonormal basis for the whole space $H^0(kL)$ with respect to $(\cdot, \cdot)_{k\phi}$. By the Bernstein–Markov property of any volume form dV (see e.g., [48]), or simply the maximum principle,

$$(7.6) \quad \phi_k \leq \phi + \epsilon_k,$$

where ϵ_k tends to zero as k tends to infinity. Since $\phi_{k, \lambda}$ is decreasing in λ , the inequality (7.6) still holds when ϕ_k is replaced by $\phi_{k, \lambda}$, i.e., $\phi_{k, \lambda} - \epsilon_k \leq \phi$. Therefore, $\phi_{k, \lambda} - \epsilon_k$ belongs to the class of metrics the supremum of which yields $P_{[\phi_k, \lambda]} \phi$, so

$$\phi_{k, \lambda} \leq P_{[\phi_k, \lambda]} \phi + \epsilon_k \leq \phi_{[\phi_k, \lambda]} + \epsilon_k,$$

so letting k tend to infinity

$$\phi_{\lambda}^{\mathcal{F}} \leq \phi_{[\phi_k, \lambda]}.$$

For the other inequality it is enough to show that for any constant C ,

$$(7.7) \quad P_{\phi_k, \lambda + C} \phi \leq \phi_{\lambda}^{\mathcal{F}}.$$

By the assumption that $\lambda < \lambda_c$ it follows that $\phi_\lambda^{\mathcal{F}} \not\equiv -\infty$. Let ψ be a positive metric dominated by both $\phi_{k,\lambda} + C$ and ϕ , where k is large enough so that kL fulfills the requirements of Theorem 7.10. We denote by $\mathcal{J}(k\psi)$ the multiplier ideal sheaf of germs of holomorphic functions locally integrable against $e^{-k\psi}$. Let $\{s_i\}$ be an orthonormal basis of $H^0(kL \otimes \mathcal{J}(k\psi))$, and denote by ψ_k the Bergman metric

$$\psi_k := \frac{1}{k} \ln \left(\sum |s_i|^2 \right).$$

By Theorem 3.2,

$$\psi \leq \psi_k + \delta_k$$

where δ_k tends to zero as k tends to infinity, and ψ_k converges pointwise to ψ . If s lies in $H^0(kL \otimes \mathcal{J}(k\psi))$, specifically we must have that

$$\int_X \frac{|s|^2}{\sum |s_{i,\lambda}|^2} dV < \infty,$$

since we assumed that ψ was dominated by $\phi_{k,\lambda} + C = 1/k \ln(\sum |s_{i,\lambda}|^2) + C$. Similarly if s lies in $H^0(kmL \otimes \mathcal{J}(km\psi))$ we have

$$\int_X \frac{|s|^2}{(\sum |s_{i,\lambda}|^2)^m} dV < \infty.$$

From Theorem 7.10 applied to the sections $\{s_{i,\lambda}\}$ it thus follows that

$$s = \sum h_\alpha s^\alpha,$$

where $h_\alpha \in H^0(k(n+1)L)$, and the s^α are monomials in the $\{s_{i,\lambda}\}$ of degree $m - n - 1$. Because of the multiplicativity of the filtration each s^α lies in

$$\mathcal{F}_{k(m-n-1)\lambda} H^0(k(m-n-1)L),$$

and by the boundedness of the filtration we also have that each h_α lies in

$$F_{-k(n+1)C} H^0(k(n+1)L)$$

for some fixed constant C . We thus get that $H^0(kmL \otimes \mathcal{J}(km\psi))$ is contained in

$$(7.8) \quad \begin{aligned} & (F_{-k(n+1)C} H^0(k(n+1)L)) (\mathcal{F}_{k(m-n-1)\lambda} H^0(k(m-n-1)L)) \\ & \subseteq \mathcal{F}_{k(m-n-1)\lambda - k(n+1)C} H^0(kmL). \end{aligned}$$

Since we assumed that $\psi \leq \phi$ we have that ψ_{km} is less than or equal to the Bergman metric using an orthonormal basis for $H^0(kmL \otimes \mathcal{J}(km\psi))$ with respect to ϕ . Because of (7.8) this Bergman metric is certainly less than or equal to $\phi_{km,\lambda'}$, where

$$\lambda' := \frac{1}{km} (k(m-n-1)\lambda - k(n+1)C).$$

Hence

$$\psi_{km} \leq \phi_{km,\lambda'}.$$

On the other hand, by Lemma 7.6 we get

$$\phi_{km,\lambda'} \leq \phi_{\lambda'}^{\mathcal{F}} + \epsilon(km),$$

where $\epsilon(km)$ is a constant independent of λ' that tends to zero as km tends to infinity. Since λ' tends to λ as m tends to infinity this implies $\psi \leq \lim_{\lambda' \rightarrow \lambda} \phi_{\lambda'}^{\mathcal{F}}$, and thus by Lemma 5.2 $\psi \leq \phi_{\lambda}^{\mathcal{F}}$. Taking the supremum over all such ψ completes the proof. \square

Corollary 7.12. *Suppose \mathcal{F} is a multiplicative linearly bounded filtration of $\oplus_k H^0(kL)$. Then the associated test curve $\phi_{\lambda}^{\mathcal{F}}$ is maximal for $\lambda < \lambda_c$ and its Legendre transform is a geodesic ray.*

Proof. Theorem 4.10 tells us that $\phi_{[\phi_{k,\lambda}]}$ is maximal with respect to $\phi = \phi_{-\infty}$. By Lemma 4.8 it follows that this is true for the limit $\phi_{\lambda}^{\mathcal{F}} = \lim_{k \rightarrow \infty} \phi_{[\phi_{k,\lambda}]}$ as well. Let ϕ_{λ} be the test curve defined by $\phi_{\lambda} := \phi_{\lambda}^{\mathcal{F}}$ for $\lambda < \lambda_c$ and $\phi_{\lambda} \equiv -\infty$ for $\lambda \geq \lambda_c$. Thus ϕ_{λ} is a maximal test curve, thus its Legendre transform is a geodesic ray. On the other hand, for every $\epsilon > 0$

$$\phi_{\lambda} \leq \phi_{\lambda}^{\mathcal{F}} \leq \phi_{\lambda-\epsilon},$$

and therefore

$$\widehat{\phi}_t \leq \widehat{(\phi^{\mathcal{F}})}_t \leq \widehat{\phi}_t + \epsilon t.$$

Since ϵ was arbitrary we get that the Legendre transform of $\phi_{\lambda}^{\mathcal{F}}$ coincides with that of ϕ_{λ} , and thus it is a geodesic ray. \square

Remark 7.13. Given an analytic test configuration $[\psi_{\lambda}]$ there is a naturally associated filtration \mathcal{F} of the section ring, defined as

$$\mathcal{F}_{k\lambda} H^0(kL) := H^0(kL \otimes \mathcal{J}(k\psi_{\lambda})).$$

This filtration is bounded, but in general not multiplicative.

8. Filtrations associated to algebraic test configurations

We recall briefly Donaldson's definition of a test configuration [24, 25]. In order to not confuse them with the our analytic test configurations, we will in this article refer to them as algebraic test configurations.

Definition 8.1. An *algebraic test configuration* \mathcal{T} for an ample line bundle L over X consists of:

- (i) a scheme \mathcal{X} with a \mathbb{C}^{\times} -action ρ ,
- (ii) a \mathbb{C}^{\times} -equivariant line bundle \mathcal{L} over \mathcal{X} ,
- (iii) and a flat \mathbb{C}^{\times} -equivariant projection $\pi : \mathcal{X} \rightarrow \mathbb{C}$ where \mathbb{C}^{\times} acts on \mathbb{C} by multiplication, such that \mathcal{L} is relatively ample, and such that if we denote by $X_1 := \pi^{-1}(1)$, then $\mathcal{L}|_{X_1} \rightarrow X_1$ is isomorphic to $rL \rightarrow X$ for some $r > 0$.

By rescaling we can without loss of generality assume that $r = 1$ in the definition. An algebraic test configuration is called a *product test configuration* if there is a \mathbb{C}^\times -action ρ' on $L \rightarrow X$ such that $\mathcal{L} = L \times \mathbb{C}$ with ρ acting on L by ρ' and on \mathbb{C} by multiplication. An algebraic test configuration is called *trivial* if it is a product test configuration with the action ρ' being the trivial \mathbb{C}^\times -action.

Since the zero-fiber $X_0 := \pi^{-1}(0)$ is invariant under the action ρ , we get an induced action on the space $H^0(kL_0)$, also denoted by ρ , where we have denoted the restriction of \mathcal{L} to X_0 by L_0 . Specifically, we let $\rho(\tau)$ act on a section $s \in H^0(kL_0)$ by

$$(8.1) \quad (\rho(\tau)(s))(x) := \rho(\tau)(s(\rho^{-1}(\tau)(x))).$$

By standard theory any vector space V with a \mathbb{C}^\times -action can be split into weight spaces V_{λ_i} on which $\rho(\tau)$ acts as multiplication by τ^{λ_i} , (see e.g., [24]). The numbers λ_i with non-trivial weight spaces are called the weights of the action. Thus we may write $H^0(kL_0)$ as

$$H^0(kL_0) = \bigoplus_{\lambda} V_{\lambda}$$

with respect to the induced action ρ .

In [35, Lemma 4] Phong–Sturm give the following linear bound on the absolute value of the weights.

Lemma 8.2. *Given a test configuration there is a constant C such that*

$$|\lambda_i| < Ck$$

whenever $\dim V_{\lambda_i} > 0$.

In [49] the second author showed how to get an associated filtration \mathcal{F} of the section ring $\bigoplus_k H^0(kL)$ given a test configuration \mathcal{T} of L which we now recall.

First note that the \mathbb{C}^\times -action ρ on \mathcal{L} via the equation (8.1) gives rise to an induced action on $H^0(\mathcal{X}, k\mathcal{L})$ as well as $H^0(\mathcal{X} \setminus X_0, k\mathcal{L})$, since $\mathcal{X} \setminus X_0$ is invariant. Let $s \in H^0(kL)$ be a holomorphic section. Then using the \mathbb{C}^\times -action ρ we get a canonical extension $\bar{s} \in H^0(\mathcal{X} \setminus X_0, k\mathcal{L})$ which is invariant under the action ρ , simply by letting

$$(8.2) \quad \bar{s}(\rho(\tau)x) := \rho(\tau)s(x)$$

for any $\tau \in \mathbb{C}^\times$ and $x \in X$.

We identify the coordinate z with the projection function $\pi(x)$, and we also consider it as a section of the trivial bundle over \mathcal{X} . Exactly as for $H^0(\mathcal{X}, k\mathcal{L})$, ρ gives rise to an induced action on sections of the trivial bundle, using the same formula (8.1). From this one sees

$$(8.3) \quad (\rho(\tau)z)(x) = \rho(\tau)(z(\rho^{-1}(\tau)x)) = \rho(\tau)(\tau^{-1}z(x)) = \tau^{-1}z(x),$$

where we used that ρ acts on the trivial bundle by multiplication on the z -coordinate. Thus

$$\rho(\tau)z = \tau^{-1}z,$$

which shows that the section z has weight -1 .

By this it follows that for any section $s \in H^0(kL)$ and any integer λ , we get a section $z^{-\lambda}\bar{s} \in H^0(\mathcal{X} \setminus X_0, k\mathcal{L})$, which has weight λ .

Lemma 8.3. *For any section $s \in H^0(kL)$ and any integer λ the section $z^{-\lambda}\bar{s}$ extends to a meromorphic section of $k\mathcal{L}$ over the whole of \mathcal{X} , which we also will denote by $z^{-\lambda}\bar{s}$.*

Proof. It is equivalent to saying that for any section s there exists an integer λ such that $z^{-\lambda}\bar{s}$ extends to a holomorphic section $S \in H^0(\mathcal{X}, k\mathcal{L})$. By flatness, which was assumed in the definition of a test configuration, the direct image bundle $\pi_*\mathcal{L}$ is in fact a vector bundle over \mathbb{C} . Thus it is trivial, since any vector bundle over \mathbb{C} is trivial. Therefore there exists a global section $S' \in H^0(\mathcal{X}, k\mathcal{L})$ such that $s = S'|_X$. On the other hand, as for $H^0(kL_0)$, $H^0(\mathcal{X}, k\mathcal{L})$ may be decomposed as a direct sum of invariant subspaces $W_{\lambda'}$ such that $\rho(\tau)$ restricted to $W_{\lambda'}$ acts as multiplication by $\tau^{\lambda'}$. Let us write

$$(8.4) \quad S' = \sum S'_{\lambda'},$$

where $S_{\lambda'} \in W_{\lambda'}$. Restricting the equation (8.4) to X gives a decomposition of s ,

$$s = \sum s_{\lambda'},$$

where $s_{\lambda'} := S'_{\lambda'}|_X$. From (8.2) and the fact that $S'_{\lambda'}$ lies in $W_{\lambda'}$ we get that for $x \in X$ and $\tau \in \mathbb{C}^\times$

$$\begin{aligned} \bar{s}_{\lambda'}(\rho(\tau)(x)) &= \rho(\tau)(s_{\lambda'}(x)) = \rho(\tau)(S'_{\lambda'}(x)) = (\rho(\tau)S'_{\lambda'})(\rho(\tau)(x)) = \\ &= \tau^{\lambda'} S'_{\lambda'}(\rho(\tau)(x)), \end{aligned}$$

and therefore $\bar{s}_{\lambda'} = \tau^{\lambda'} S'_{\lambda'}$. Since trivially

$$\bar{s} = \sum \bar{s}_{\lambda'}$$

it follows that $t^\lambda \bar{s}$ extends holomorphically as long as $\lambda \geq \max -\lambda'$. \square

Definition 8.4. Given a test configuration \mathcal{T} we define a vector space-valued map \mathcal{F} from $\mathbb{Z} \times \mathbb{N}$ by letting

$$(\lambda, k) \longmapsto \{s \in H^0(kL) : z^{-\lambda}\bar{s} \in H^0(\mathcal{X}, k\mathcal{L})\} =: \mathcal{F}_\lambda H^0(kL).$$

It is immediate that \mathcal{F}_λ is decreasing since $H^0(\mathcal{X}, k\mathcal{L})$ is a $\mathbb{C}[z]$ -module. We can extend \mathcal{F} to a filtration by letting

$$\mathcal{F}_\lambda H^0(kL) := \mathcal{F}_{\lceil \lambda \rceil} H^0(kL)$$

for non-integers λ , thus making \mathcal{F} left-continuous. Since

$$z^{-(\lambda+\lambda')}\overline{ss'} = (z^{-\lambda}\overline{s})(z^{-\lambda'}\overline{s'}) \in H^0(\mathcal{X}, k\mathcal{L})H^0(\mathcal{X}, m\mathcal{L}) \subseteq H^0(\mathcal{X}, (k+m)\mathcal{L})$$

whenever $s \in \mathcal{F}_\lambda H^0(kL)$ and $s' \in \mathcal{F}_{\lambda'} H^0(mL)$, we see that

$$(\mathcal{F}_\lambda H^0(kL))(\mathcal{F}_{\lambda'} H^0(mL)) \subseteq \mathcal{F}_{\lambda+\lambda'} H^0((k+m)L),$$

i.e., \mathcal{F} is multiplicative.

Recall that we had the decomposition of $H^0(kL_0)$ into weight spaces V_λ .

Lemma 8.5. *For each λ , we have that*

$$\dim \mathcal{F}_\lambda H^0(kL) = \sum_{\lambda' \geq \lambda} \dim V_{\lambda'}.$$

Proof. We have the following isomorphism:

$$(\pi_* k\mathcal{L})|_{\{0\}} \cong H^0(\mathcal{X}, k\mathcal{L})/zH^0(\mathcal{X}, k\mathcal{L}),$$

the right-to-left arrow being given by the restriction map (see e.g., [39, p12]).

Also, for $k \gg 0$, $(\pi_* k\mathcal{L})|_{\{0\}} = H^0(kL_0)$, therefore for large k

$$(8.5) \quad H^0(kL_0) \cong H^0(\mathcal{X}, k\mathcal{L})/zH^0(\mathcal{X}, k\mathcal{L}),$$

We also had a decomposition of $H^0(\mathcal{X}, k\mathcal{L})$ into the sum of its invariant weight spaces W_λ . By Lemma 8.3 it is clear that a section $S \in H^0(\mathcal{X}, k\mathcal{L})$ lies in W_λ if and only if it can be written as $z^{-\lambda}\overline{s}$ for some $s \in H^0(kL)$, in fact we have that $s = S|_X$. Thus

$$W_\lambda \cong \mathcal{F}_\lambda H^0(kL),$$

and by the isomorphism (8.5) then

$$V_\lambda \cong \mathcal{F}_\lambda H^0(kL)/\mathcal{F}_{\lambda+1} H^0(kL).$$

Therefore

$$(8.6) \quad \dim \mathcal{F}_\lambda H^0(kL) = \sum_{\lambda' \geq \lambda} \dim V_{\lambda'}.$$

□

Using Lemma 8.5 together with Lemma 8.2 shows that the filtration \mathcal{F} is bounded.

9. The geodesic rays of Phong and Sturm

In [35] Phong–Sturm show how to construct a weak geodesic ray, starting with a $\phi \in \mathcal{H}(L)$ and an algebraic test configuration \mathcal{T} (see also [44] for how this works in the toric setting). In the previous section we showed how to associate an analytic test configuration $[\phi_\lambda^{\mathcal{F}}]$ to an algebraic test configuration, and thus get a weak geodesic using the Legendre transform of its maximal envelope. Recall by Proposition 6.10 this geodesic is the same as

the Legendre transform of the original test curve $\phi_\lambda^{\mathcal{F}}$. The goal in this section is to prove that this ray coincides with the one constructed by Phong–Sturm.

To describe what we aim to show, recall that if V is a vector space with a scalar product, and \mathcal{F} is a filtration of V , there is a unique decomposition of V into a direct sum of mutually orthogonal subspaces V_{λ_i} such that

$$\mathcal{F}_\lambda V = \bigoplus_{\lambda_i \geq \lambda} V_{\lambda_i}.$$

Furthermore we allow for λ_i to be equal to λ_j even when $i \neq j$, so we can assume that all the subspaces V_{λ_i} are one dimensional. This additional decomposition is of course not unique, but it will not matter in what follows.

Let $\phi \in \mathcal{H}(L)$ and $H^0(kL) = \bigoplus V_{\lambda_i}$ be the decomposition of $H^0(kL)$ with respect to the scalar product $(\cdot, \cdot)_{k\phi}$ coming from the volume form $(dd^c\phi)^n$. Consider next the filtration coming from an algebraic test configuration (note that then the collection of λ_i will depend also on k but we omit that from our notation) and define the normalized weights to be

$$\bar{\lambda}_i := \frac{\lambda_i}{k},$$

which form a bounded family by Lemma 8.2.

Now if s_i is a vector of unit length in V_{λ_i} , then $\{s_i\}$ will be an orthonormal basis for $H^0(kL)$. Since the filtration \mathcal{F} encodes the \mathbb{C}^* -action on $H^0(k\mathcal{L})$ it is easy to see that the basis $\{s_i\}$ is the same one as in [35, Lem 7]. In terms of the notation in the previous sections

$$\phi_{k,\lambda} = \frac{1}{k} \ln \left(\sum_{\lambda_i \geq k\lambda} |s_i|^2 \right) \quad \text{and} \quad \phi_\lambda^{\mathcal{F}} = \lim_{k \rightarrow \infty}^* \phi_{k,\lambda}.$$

Definition 9.1. Let

$$\Phi_k(t) := \frac{1}{k} \ln \left(\sum_i e^{t\lambda_i} |s_i|^2 \right)$$

The *Phong–Sturm* ray is the limit

$$(9.1) \quad \Phi(t) := \lim_{k \rightarrow \infty}^* (\sup_{l \geq k} \Phi_l(t)).$$

Our goal is the following:

Theorem 9.2. *Let $\phi^{\mathcal{F}}$ be the analytic test configuration associated to the filtration \mathcal{F} from a test configuration. Then*

$$\Phi(t) = \widehat{(\phi^{\mathcal{F}})}_t.$$

In particular, the results from the previous section yield another proof of [35, Thm 1] which says that $\Phi(t)$ is a weak geodesic ray emanating from ϕ .

Lemma 9.3.

$$(9.2) \quad \Phi(t) = \lim_{k \rightarrow \infty}^* (\sup_{l \geq k} \Phi_l(t)) = \lim_{k \rightarrow \infty}^* (\sup_{l \geq k} \max_i \{\phi_{l, \bar{\lambda}_i} + t \bar{\lambda}_i\}).$$

Proof. Our proof will be based on the elementary fact that if $\{a_{l,i} : i \in I_l\}$ is a set of real numbers then

$$(9.3) \quad \max_{i \in I_l} a_{l,i} \leq \frac{1}{l} \ln \sum_{i \in I_l} e^{la_{l,i}} \leq \max_{i \in I_l} a_{l,i} + \frac{1}{l} \ln |I_l|.$$

Now pick $x \in X$ and $t > 0$. Let

$$a_{l,i} := \frac{1}{l} \ln |s_i(x)|^2 + t \bar{\lambda}_i$$

and I_l be the indexing set for the λ_i . Then $|I_l| = O(l^n)$ and

$$\Phi_l(t) = \frac{1}{l} \ln \left(\sum_i e^{la_{l,i}} \right).$$

Thus by (9.3)

$$(9.4) \quad \max_i \{a_{l,i}\} \leq \Phi_l(t) \leq \max_i \{a_{l,i}\} + \frac{|I_l|}{l}.$$

Now set

$$b_{l,i} := \phi_{l, \bar{\lambda}_i} + t \bar{\lambda}_i = \frac{1}{l} \ln \sum_{\lambda_j \geq \lambda_i} |s_j(x)|^2 + t \bar{\lambda}_i.$$

For fixed i , pick any j_0 such that

$$\max_{\lambda_j \geq \lambda_i} |s_j(x)|^2 = |s_{j_0}|^2 \quad \text{and} \quad \lambda_{j_0} \geq \lambda_i.$$

Then

$$b_{l,i} \leq \frac{1}{l} \ln (|I_l| |s_{j_0}|^2 + t \bar{\lambda}_i) \leq \frac{1}{l} \ln |s_{j_0}|^2 + t \bar{\lambda}_{j_0} + \frac{\ln |I_l|}{l} = a_{j_0, l} + \frac{\ln |I_l|}{l}.$$

Clearly $a_{l,i} \leq b_{l,i}$ for all i , so we in fact have

$$\max_i \{a_{l,i}\} \leq \max_i \{b_{l,i}\} \leq \max_i \{a_{l,i}\} + \frac{\ln |I_l|}{l},$$

which combined with (9.4) yields

$$\max_i \{b_{l,i}\} - \frac{\ln |I_l|}{l} \leq \Phi_l(t) \leq \max_i \{b_{l,i}\} + \frac{\ln |I_l|}{l}.$$

Now taking the supremum over all $l \geq k$ followed by the upper semicontinuous regularization and then the limit as k tends to infinity gives the result since $k^{-1} \ln |I_k|$ tends to zero. \square

Proof of Theorem 9.2. From Lemma 7.6 there is a constant $\epsilon(l)$ such that

$$\phi_{l,\bar{\lambda}_i} + t\bar{\lambda}_i \leq \phi_{\bar{\lambda}_i}^{\mathcal{F}} + t\bar{\lambda}_i + \epsilon(l),$$

where $\epsilon(l)$ is independent of λ_i and tends to zero as l tends to infinity. Thus

$$\max_i \{\phi_{l,\bar{\lambda}_i} + t\bar{\lambda}_i\} \leq \sup_{\lambda} \{\phi_{\lambda}^{\mathcal{F}} + t\lambda\} + \epsilon(l),$$

and so

$$\lim_{k \rightarrow \infty}^* \sup_{l \geq k} \max_i \{\phi_{l,\bar{\lambda}_i} + t\bar{\lambda}_i\} \leq (\widehat{\phi^{\mathcal{F}}})_t + \lim_{k \rightarrow \infty} \sup_{l \geq k} \epsilon(l),$$

so using Lemma 9.3 gives

$$\Phi(t) \leq (\widehat{\phi^{\mathcal{F}}})_t.$$

For the opposite inequality, let $\lambda \in \mathbb{R}$ be arbitrary. Trivially

$$\Phi_k(t) = \frac{1}{k} \ln \left(\sum_i e^{t\lambda_i} |s_i|^2 \right) \geq \frac{1}{k} \ln \left(\sum_{\lambda_i \geq k\lambda} e^{tk\lambda} |s_i|^2 \right) = \phi_{k,\lambda} + t\lambda.$$

Hence

$$\Phi(t) \geq \phi_{\lambda}^{\mathcal{F}} + t\lambda$$

for any λ , and thus

$$\Phi(t) \geq (\widehat{\phi^{\mathcal{F}}})_t. \quad \square$$

Remark 9.4. Phong–Sturm prove in [35] that the geodesic ray one gets from an algebraic test configuration \mathcal{F} is non-trivial if the norm of \mathcal{F} is non-zero. From the above we see that the weak geodesic ray is trivial if and only if the associated analytic test configuration is trivial, i.e., if there exists a number λ_c such that $\phi_{\lambda} = \phi$ when $\lambda < \lambda_c$ and $\phi_{\lambda} \equiv -\infty$ when $\lambda > \lambda_c$.

References

- [1] V. Apostolov, D. Calderbank, P. Gauduchon and C. Tønnesen-Friedman, *Hamiltonian 2-forms in Kähler geometry III, extremal metrics and stability*, Invent. Math **173**(3) (2008), 547–601.
- [2] C. Arezzo and G. Tian, *Infinite geodesic rays in the space of Kähler potentials*, Ann. Sc. Norm. Sup. Pisa (5) **2**(4) (2003), 617–630.

- [3] E. Bedford and B.A. Taylor, *A new capacity for plurisubharmonic functions*, Acta Math. **149**(1–2) (1982), 1–40.
- [4] E. Bedford and B.A. Taylor, *Fine topology, Silov boundary, and $(dd^c)^n$* , J. Funct. Anal. **72**(2) (1987), 225–251.
- [5] R. Berman, *Bergman kernels and equilibrium measures for line bundles over projective manifolds*, Amer. J. Math. **131**(5) (2009), 1485–1524.
- [6] R. Berman, *Bergman kernels and equilibrium measures for ample line bundles* (2007) [arXiv:0704.1640](https://arxiv.org/abs/0704.1640).
- [7] R. Berman and S. Boucksom, *Growth of balls of holomorphic sections and energy at equilibrium* Invent. Math. **181**(2) (2010), 337–394
- [8] R. Berman, S. Boucksom, V. Guedj and A. Zeriahi, *A variational approach to complex Monge–Ampère equations*, Publ. Math. Inst. Hautes Études Sci. **117** (2013), 179–245.
- [9] B. Bollobás and O. Riordan, *Percolation* Cambridge University Press, New York, 2006. x+323 pp. ISBN: 978-0-521-87232-4
- [10] T. Bouche, *Convergence de la métrique de Fubini–Study d’un fibré linéaire positif*, Ann. Inst. Fourier (1) **40** (1990), 117–130.
- [11] B. Berndtsson, *Probability measures associated to geodesics in the space of Kähler metrics*, (2009), [arXiv:math/0907.1806](https://arxiv.org/abs/math/0907.1806).
- [12] S. Boucksom and H. Chen, *Okounkov bodies of filtered linear series*, Compos. Math. **147**(4) (2011), 1205–1229
- [13] S. Boucksom, P. Eyssidieux, V. Guedj and A. Zeriahi, *Monge–Ampère equations in big cohomology classes*, Acta Math. **205**(2) (2010), 199–262.
- [14] D. Catlin, *The Bergman kernel and a theorem of Tian*, in ‘Analysis and geometry in several complex variables (Katata, 1997)’, Trends Math., 1–23. Birkhäuser Boston, Boston, MA, 1999.
- [15] X.X. Chen, *The space of Kähler metrics*, J. Differ. Geom. **56** (2000), 189–234.
- [16] X.X. Chen *Space of Kähler metrics III: on the lower bound of the Calabi energy and geodesic distance*, Invent. Math. **175**(3) (2009), 453–503.
- [17] X.X. Chen and S. Sun, *Space of Kähler metrics V–Kähler quantization* In Metric and Differential Geometry, the Jeff Cheeger anniversary volume, Progr. Math. **297** (2012), 19–42.
- [18] X.X. Chen and G. Tian, *Geometry of Kähler metrics and foliations by holomorphic discs*, Publ. Math. Inst. Hautes Études Sci. No. **107** (2008), 1–107.
- [19] J.-P. Demailly, *Complex analytic and differential geometry* OpenContent Book — available at www-fourier.ujf-grenoble.fr/~demailly/books.html
- [20] J.-P. Demailly, *Regularization of closed positive currents and intersection theory*, J. Algebr. Geom. **1**(3) (1992), 361–409.
- [21] J.-P. Demailly, *Appendix to I. Cheltsov and C. Shramov’s article “Log canonical thresholds of smooth Fano Threefolds”*, Russian Mathematical Surveys **63** (2008), 859–958.
- [22] J.-P. Demailly, T. Peternell and M. Schneider, *Pseudo-effective line bundles on compact Kähler manifolds* Int. J. Math. **12**(6) (2001), 689–741.
- [23] S.K. Donaldson, *Scalar curvature and projective embeddings. I*. J. Differ. Geom. **59**(3) (2001), 479–522.

- [24] S.K. Donaldson, *Scalar curvature and stability of toric varieties*, J. Differ. Geom. **62**(2) (2002), 289–349.
- [25] S.K. Donaldson, *Lower bounds on the Calabi functional*, J. Differ. Geom. **70**(3) (2005), 453–472.
- [26] S.K. Donaldson, *Symmetric spaces, Kähler geometry and Hamiltonian dynamics*. Northern California Symplectic Geometry Seminar, 13–33. Amer. Math. Soc. Transl. Ser. 2, 196, Amer. Math. Soc., Providence, RI, 1999.
- [27] V. Guedj and A. Zeriahi, *Intrinsic capacities on compact Kähler manifolds*, J. Geom. Anal. **15**(4) (2005), 607–639.
- [28] C. Kiselman, *The partial Legendre transformation for plurisubharmonic functions*, Invent. Math. **49**(2) (1978), 137–148.
- [29] M. Klimek, *Pluripotential Theory* London Mathematical Society Monographs. New Series, 6. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1991. ISBN: 0-19-853568-6
- [30] T. Mabuchi, *Some symplectic geometry on compact Kähler manifolds. I.*, Osaka J. Math. **24**(2) (1987), 227.
- [31] T. Mabuchi, *K-energy maps integrating Futaki invariants*, Tohoku Math. J. (2) **38**(4) (1986), 575–593.
- [32] A. Rashkovskii and R. Sigurdsson. *Green functions with singularities along complex spaces*, Int. J. Math. **16**(4) (2005), 333–355
- [33] R. Rockafellar, *Conjugates and Legendre transforms of convex functions*, Canad. J. Math. **19** (1967), 200–205.
- [34] D.H. Phong and J. Sturm, *The Monge–Ampère operator and geodesics in the space of Kähler potentials*, Invent. Math. **166**(1) (2006), 125–149.
- [35] D.H. Phong and J. Sturm, *Test configurations for K-stability and geodesic rays*, J. Symplec. Geom. **5**(2) (2007), 221–247.
- [36] D H Phong and J Sturm *Lectures on stability and constant scalar curvature*, in ‘Current Developments in Mathematics, 2007, 101–176, Int. Press, Somerville, MA, 2009
- [37] D.H. Phong and J. Sturm, *Regularity of geodesic rays and Monge–Ampère equations* Proc. Amer. Math. Soc. **138**(10) (2010), 3637–3650
- [38] G. Székelyhidi, *Filtrations and test-configurations*, (2011), [arXiv:1111.4986](https://arxiv.org/abs/1111.4986)
- [39] J. Ross and R.P. Thomas, *A study of the Hilbert–Mumford criterion for the stability of projective varieties*, J. Algebr. Geom. **16** (2007), 201–255.
- [40] J. Ross and D. Witt Nyström, *Envelopes of plurisubharmonic metrics with prescribed singularities* (2012), [arXiv:1210.2220](https://arxiv.org/abs/1210.2220)
- [41] Y.A. Rubinstein and S. Zelditch, *The Cauchy problem for the homogeneous Monge–Ampère equation, I. Toeplitz quantization*, J. Differ. Geom. **90**(2) (2012), 303–327
- [42] Y.A. Rubinstein and S. Zelditch, *The Cauchy problem for the homogeneous Monge–Ampère equation, II. Legendre transform*, Adv. Math. **228**(6) (2011), 2989–3025.
- [43] S. Semmes, *Complex Monge–Ampère and symplectic manifolds*, Amer. J. Math. **114**(3) (1992), 495–550.
- [44] J. Song and S. Zelditch, *Test configurations, large deviations and geodesic rays on toric varieties*, Adv. Math. **229**(4) (2012), 2338–2378.

- [45] G. Tian, *On a set of polarized Kähler metrics on algebraic manifolds*, J. Differ. Geom. **32** (1) (1990), 99–130.
- [46] G. Tian, *Kähler–Einstein metrics with positive scalar curvature*, Invent. Math. **130**(1) (1997), 1–37.
- [47] D. Varolin, *Division theorems and twisted complexes*, Math. Z. **259**(1) (2008), 1–20.
- [48] D. Witt Nyström, *Transforming metrics on a line bundle to the Okounkov body*, (2009), [arXiv:math/0903.5167](https://arxiv.org/abs/math/0903.5167).
- [49] D. Witt Nyström, *Test configurations and Okounkov bodies*, Comp. Math. **148**(6) (2012), 1736–1756.
- [50] S. Zelditch, *Szegő kernels and a theorem of Tian*, Int. Math. Res. Notices **6** (1998), 317–331.

DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS
UNIVERSITY OF CAMBRIDGE
UK
E-mail address: `j.ross@dpms.cam.ac.uk`

DEPARTMENT OF MATHEMATICAL SCIENCES
UNIVERSITY OF GÖTHENBURG
SWEDEN
E-mail address: `witnyst@chalmers.se`

Received 04/04/2011, accepted 04/08/2013

We would like to thank Robert Berman, Bo Berndtsson, Sebastian Boucksom, Yanir Rubinstein and Richard Thomas for helpful discussions and the referees for their constructive suggestions. We also thank Dano Kim for pointing out a mistake in a previous version of this paper. The first author also acknowledges support from a Marie Curie International Reintegration Grant within the 7th European Community Framework Program (PIRG-GA-2008-230920).

