

HIGH-ORDER GREEN OPERATORS ON THE DISK AND THE POLYDISC*

YANG LIU[†], ZHIHUA CHEN[‡], AND YIFEI PAN[§]

Abstract. In this paper, we give the explicit expressions of high-order Green operators on the disk and the polydisc, and hence the kernel functions of high-order Green operators are also presented. As applications, we present the explicit integral expressions of all the solutions for linear high-order partial differential equations in the disk.

Key words. Green operator, disk, polydisc, linear partial differential equation.

AMS subject classifications. 32W50.

1. Preliminaries. Singular integral is an important tool in harmonic analysis, complex analysis and Clifford analysis, et. al. There are many types of singular integrals investigated by numerous mathematicians. In [1], Calderón-Zygmund type singular integrals have been studied in theory and applications. For the singular integral operator, in [2], a boundedness criterion for the Cauchy singular integral operator in weighted grand Lebesgue spaces have been given. Moreover, semi-Fredholm properties of certain singular integral operators, asymptotic invertibility of Toeplitz operators, weighted uniform convergence of the quadrature method for Cauchy singular integral equations, and Toeplitz and singular integral operators on general Carleson Jordan curves have been discussed in [3]. The book [4] has been devoted to the Fredholm theory of singular integral operators with shifts on $L^p(\Gamma)$, ($1 < p < \infty$), where Γ is a Lyapunov curve in the complex plane which is homeomorphic to a circle or a segment.

In one complex dimension, the inverse operator (or Green operator) of the Cauchy-Riemann operator $\bar{\partial}$ plays an important role in finding a solution of $\bar{\partial}$ equation, see [5, 6, 7]. Especially, high-order Green operators are used in [7] to prove the general existence theorem for nonlinear partial differential systems of any order in one complex variable. In this paper, we will give the explicit expression of high-order Green operators on the disk and the polydisc, and the kernel functions of high-order Green operators will be presented as well.

Let D be the closed disk $\{z \in \mathbb{C} \mid |z| \leq R\}$ and C be its boundary $\{z \in \mathbb{C} \mid |z| = R\}$. Unless otherwise state, in this paper, functions we consider will be complex valued and integrable with D . $C^\alpha(D)$ denotes the set of all functions f on D , where

$$H_\alpha[f] = \sup\left\{\frac{|f(z) - f(z')|}{|z - z'|^\alpha} \mid z, z' \in D\right\}$$

is finite. For $f \in C^\alpha(D)$ we define

$$\|f\| = \|f\| + (2R)^\alpha H_\alpha[f],$$

*Received March 12, 2012; accepted for publication February 22, 2013. This work was supported by the NSF of Zhejiang Province of China (No. LY14A010008), NNSF of China (No. 11171255, 11101373), and Doctoral Program Foundation of the Ministry of Education of China (No. 20090072110053).

[†]Department of Mathematics, Zhejiang Normal University, Jinhua 321004, China (liuyang4740@gmail.com).

[‡]Department of Mathematics, Tongji University, Shanghai 200092, China (zzzhc@tongji.edu.cn).

[§]Department of Mathematical Sciences, Indiana University-Purdue University Fort Wayne, Fort Wayne, Indiana 46805, USA (pan@ipfw.edu).

where $|f|$ denotes $\sup_{z \in D} |f(z)|$. $C^k(D)$ is the set of all functions f on D whose k th order partial derivatives exist and are continuous, $k \geq 0$ is an integer. $C^{k+\alpha}(D)$ is the set of all functions f on D whose k th order partial derivatives exist and belong to $C^\alpha(D)$. For $f \in C^{k+\alpha}(D)$, we have the definition in terms of $\|\cdot\|$:

$$\|f\|^{(k)} = \max_{i+j=k} \{ \|\partial^i \bar{\partial}^j f\| \}.$$

It should be pointed out that the function $\|\cdot\|^{(k)}$ on $C^{k+\alpha}(D)$ is a semi-norm rather than a norm since $\|f\|^{(k)} = 0$ if and only if f is polynomial of degree of $k - 1$. The following operators are defined on D as in [5]:

$$\begin{aligned} (1) \quad T f(z) &= \frac{-1}{2\pi i} \int_D \frac{f(\zeta) d\bar{\zeta} \wedge d\zeta}{\zeta - z}, \quad \bar{T} f(z) = \frac{-1}{2\pi i} \int_D \frac{f(\zeta) d\bar{\zeta} \wedge d\zeta}{\bar{\zeta} - \bar{z}}, \\ {}^2 T f(z) &= \frac{-1}{2\pi i} \int_D \frac{f(\zeta) - f(z)}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta, \quad {}^2 \bar{T} f(z) = \frac{-1}{2\pi i} \int_D \frac{f(\zeta) - f(z)}{(\bar{\zeta} - \bar{z})^2} d\bar{\zeta} \wedge d\zeta, \\ S f(z) &= \frac{1}{2\pi i} \int_C \frac{f(\zeta) d\zeta}{\zeta - z}, \quad \bar{S} f(z) = \frac{-1}{2\pi i} \int_C \frac{f(\zeta) d\bar{\zeta}}{\bar{\zeta} - \bar{z}}, \\ S_b f(z) &= \frac{1}{2\pi i} \int_C \frac{f(\zeta) d\bar{\zeta}}{\zeta - z}, \quad \bar{S}_b f(z) = \frac{-1}{2\pi i} \int_C \frac{f(\zeta) d\zeta}{\bar{\zeta} - \bar{z}}, \end{aligned}$$

where $\bar{T}(f) = \overline{T(f)}$, $\bar{S}(f) = \overline{S(f)}$, and $\bar{S}_b(f) = \overline{S_b(f)}$. More generally, if Δ is a closed bounded domain, then $T_\Delta f$ and $S_\Delta f$ are defined for continuous f on Δ by

$$T_\Delta f(z) = \frac{-1}{2\pi i} \int_\Delta \frac{f(\zeta) d\bar{\zeta} \wedge d\zeta}{\zeta - z}, \quad S_{\partial\Delta} f(z) = \frac{1}{2\pi i} \int_{\partial\Delta} \frac{f(\zeta) d\zeta}{\zeta - z}.$$

The fundamental properties between operators T, S are given by [5] as follows.

LEMMA 1.1. [5] *If $f \in C^1(D)$, then*

$$\begin{aligned} T\bar{\partial}f &= f - S f \text{ on } \text{Int}(D), \\ \bar{T}\partial f &= f - \bar{S} f \text{ on } \text{Int}(D). \end{aligned}$$

The smoothness properties of integral operator T has been shown in [5].

LEMMA 1.2. [5] *If $f \in C^\alpha(D)$, then $Tf \in C^{1+\alpha}(D)$. Moreover,*

$$\bar{\partial} T f = f, \quad \partial T f = {}^2 T f,$$

and

$$H_\alpha[{}^2 T f] \leq C_0 H_\alpha[f]$$

where $C_0 = \frac{12}{\alpha(1-\alpha)}$. *If $f \in C^{k+\alpha}(D)$ ($k \geq 0$), then $Tf \in C^{k+1+\alpha}(D)$.*

Meanwhile, if replace $\partial, \bar{\partial}, T, {}^2 T$ with $\bar{\partial}, \partial, \bar{T}, {}^2 \bar{T}$, respectively, then one can get the similar result as above lemma for \bar{T} .

LEMMA 1.3. [7] *It holds for $l \geq 0$ that*

$$\int_\Delta \frac{(\bar{\zeta} - \bar{z}_0)^l}{\zeta - w} d\bar{\zeta} \wedge d\zeta = \frac{-2\pi i}{l+1} (\bar{w} - \bar{z}_0)^{l+1},$$

where $\Delta = \{\zeta \in \mathbb{C} \mid |\zeta - \bar{z}_0| \leq r\}$.

Proof. It is easy to get Lemma 1.1 if we replace D , C with Δ , $\partial\Delta$, respectively. Apply Lemma 1.1 to the function $(\bar{\zeta} - \bar{z}_0)^{l+1}$, we can obtain that

$$\begin{aligned} \int_{\Delta} \frac{(\bar{\zeta} - \bar{z}_0)^l}{\zeta - w} d\bar{\zeta} \wedge d\zeta &= \frac{-2\pi i}{l+1} T_{\Delta}(\bar{\partial}(\bar{\zeta} - \bar{z}_0)^{l+1})(w) \\ &= \frac{-2\pi i}{l+1} [(\bar{\zeta} - \bar{z}_0)^{l+1} - S_{\Delta}((\bar{\zeta} - \bar{z}_0)^{l+1})](w) \\ &= \frac{-2\pi i}{l+1} [(\bar{w} - \bar{z}_0)^{l+1} - \frac{1}{2\pi i} \int_{|\zeta - z_0|=r} \frac{(\bar{\zeta} - \bar{z}_0)^{l+1}}{\zeta - w} d\zeta] \\ &= \frac{-2\pi i}{l+1} [(\bar{w} - \bar{z}_0)^{l+1} - \frac{r^{2(l+1)}}{2\pi i} \int_{|\zeta - z_0|=r} \frac{1}{(\zeta - z_0)^{l+1}(\zeta - w)} d\zeta] \\ &= \frac{-2\pi i}{l+1} (\bar{w} - \bar{z}_0)^{l+1}, \end{aligned}$$

where the last equality comes from the residue theorem. The lemma is proved. \square

2. High-order Green operator on D . Denote $T^2 = TT$, $\bar{T}^2 = \bar{T}\bar{T}$, similar notations for $T^\mu, \bar{T}^\nu, T^\mu \bar{T}^\nu$ for any $\mu, \nu > 0$. In this section, we will get the explicit expression of $T^\mu \bar{T}^\nu f$.

THEOREM 2.1. *Given $f \in C^\alpha(D)$, $T^k f, \bar{T}^k f \in C^{k+\alpha}(D)$ with integer $k > 0$, and*

$$(2) \quad T^k f(z) = \frac{(-1)^k}{(k-1)! \cdot 2\pi i} \int_D \frac{(\bar{\zeta} - \bar{z})^{k-1} f(\zeta)}{\zeta - z} d\bar{\zeta} \wedge d\zeta,$$

$$(3) \quad \bar{T}^k f(z) = \frac{(-1)^k}{(k-1)! \cdot 2\pi i} \int_D \frac{(\zeta - z)^{k-1} f(\zeta)}{\bar{\zeta} - \bar{z}} d\bar{\zeta} \wedge d\zeta.$$

Proof. When $k = 1$, (2) is obvious.

Now we assume (2) is valid when for $k - 1$, i.e.,

$$T^{k-1} f(z) = \frac{(-1)^{k-1}}{(k-2)! \cdot 2\pi i} \int_D \frac{(\bar{\zeta} - \bar{z})^{k-2} f(\zeta)}{\zeta - z} d\bar{\zeta} \wedge d\zeta.$$

Hence, we can obtain with Lemma 1.3 that

$$\begin{aligned}
 T^k f(z) &= TT^{k-1} f(z) \\
 &= \frac{-1}{2\pi i} \int_D \frac{T^{k-1} f(\zeta)}{\zeta - z} d\bar{\zeta} \wedge d\zeta \\
 &= \frac{-1}{2\pi i} \int_D \frac{(-1)^{k-1}}{(k-2)! \cdot 2\pi i} \int_D \frac{(\bar{\eta} - \bar{\zeta})^{k-2} f(\eta)}{\eta - \zeta} d\bar{\eta} \wedge d\eta}{\zeta - z} d\bar{\zeta} \wedge d\zeta \\
 &= \frac{(-1)^k}{(k-2)! \cdot (2\pi i)^2} \int_D \int_D \frac{(\bar{\eta} - \bar{\zeta})^{k-2} d\bar{\zeta} \wedge d\zeta}{(\eta - \zeta)(\zeta - z)} f(\eta) d\bar{\eta} \wedge d\eta \\
 &= \frac{(-1)^k}{(k-2)! \cdot (2\pi i)^2} \int_D \frac{f(\eta) d\bar{\eta} \wedge d\eta}{\eta - z} \int_D (\bar{\eta} - \bar{\zeta})^{k-2} \left(\frac{1}{\eta - \zeta} + \frac{1}{\zeta - z} \right) d\bar{\zeta} \wedge d\zeta \\
 &= \frac{(-1)^k}{(k-2)! \cdot (2\pi i)^2} \int_D \frac{f(\eta) d\bar{\eta} \wedge d\eta}{\eta - z} \left[\int_D \frac{(\bar{\eta} - \bar{\zeta})^{k-2}}{\eta - \zeta} d\bar{\zeta} \wedge d\zeta \right. \\
 &\quad \left. + \int_D \frac{(\bar{\eta} - \bar{\zeta})^{k-2}}{\zeta - z} d\bar{\zeta} \wedge d\zeta \right] \\
 &= \frac{(-1)^k}{(k-2)! \cdot (2\pi i)^2} \int_D \frac{f(\eta) d\bar{\eta} \wedge d\eta}{\eta - z} \left[\int_D \frac{\sum_{l=0}^{k-2} \binom{k-2}{l} \bar{\eta}^l (-\bar{\zeta})^{k-2-l}}{\eta - \zeta} d\bar{\zeta} \wedge d\zeta \right. \\
 &\quad \left. + \int_D \frac{\sum_{l=0}^{k-2} \binom{k-2}{l} \bar{\eta}^l (-\bar{\zeta})^{k-2-l}}{\zeta - z} d\bar{\zeta} \wedge d\zeta \right] \\
 &= \frac{(-1)^k}{(k-2)! \cdot (2\pi i)^2} \int_D \frac{\sum_{l=0}^{k-2} \binom{k-2}{l} \bar{\eta}^l f(\eta) d\bar{\eta} \wedge d\eta}{\eta - z} \left[\int_D \frac{(-\bar{\zeta})^{k-2-l}}{\eta - \zeta} d\bar{\zeta} \wedge d\zeta \right. \\
 &\quad \left. + \int_D \frac{(-\bar{\zeta})^{k-2-l}}{\zeta - z} d\bar{\zeta} \wedge d\zeta \right] \\
 &= \frac{(-1)^k}{(k-2)! \cdot (2\pi i)^2} \int_D \frac{f(\eta)}{\eta - z} \sum_{l=0}^{k-2} \binom{k-2}{l} \bar{\eta}^l \left[\frac{2\pi i}{k-1-l} \bar{\eta}^{k-1-l} (-1)^{k-2-l} \right. \\
 &\quad \left. + \frac{-2\pi i}{k-1-l} \bar{z}^{k-1-l} (-1)^{k-2-l} \right] d\bar{\eta} \wedge d\eta \\
 &= \frac{(-1)^k}{(k-2)! \cdot (2\pi i)^2} \int_D \frac{-2\pi i f(\eta)}{(\eta - z)(k-1)} \sum_{l=0}^{k-1} \binom{k-1}{l} \bar{\eta}^l [(-\bar{\eta})^{k-1-l} - (-\bar{z})^{k-1-l}] d\bar{\eta} \wedge d\eta \\
 &= \frac{(-1)^k}{(k-2)! \cdot (2\pi i)^2} \int_D \frac{-2\pi i f(\eta)}{(\eta - z)(k-1)} [(\bar{\eta} - \bar{\eta})^{k-1} - (\bar{\eta} - \bar{z})^{k-1}] d\bar{\eta} \wedge d\eta \\
 &= \frac{(-1)^k}{(k-2)! \cdot (2\pi i)^2} \int_D \frac{2\pi i f(\eta)}{(\eta - z)(k-1)} (\bar{\eta} - \bar{z})^{k-1} d\bar{\eta} \wedge d\eta \\
 &= \frac{(-1)^k}{(k-1)! \cdot 2\pi i} \int_D \frac{(\bar{\eta} - \bar{z})^{k-1} f(\eta)}{\eta - z} d\bar{\eta} \wedge d\eta.
 \end{aligned}$$

Thus, (2) is proved. For (3), the proof is similar, we omit here. \square

REMARK 2.2. From Theorem 2.1, we have $\overline{T^k} \bar{f}(z) = \overline{T^k f(z)}$.

LEMMA 2.3. For any $a, b \in D$ and $a \neq b$, integer $k > 0$,

$$\int_D \frac{(\zeta - b)^{k-1} d\bar{\zeta} \wedge d\zeta}{(\zeta - a)(\bar{\zeta} - \bar{b})} = 2\pi i \left(C_1(a, b, k) + (a - b)^{k-1} \ln \frac{R^2 - a\bar{b}}{|a - b|^2} \right),$$

where $C_1(a, b, k) = \sum_{l=1}^{k-1} \binom{-b^l}{l} \left(\sum_{j=0}^{k-1-l} \binom{k-1}{j} a^{k-1-j} (-b)^j \right)$ for $k > 1$ and $C_1(a, b, 1) = 0$.

Proof. For any small $\varepsilon > 0$, denote $D_a = \{|\zeta - a| < \varepsilon\}$, $D_b = \{|\zeta - b| < \varepsilon\}$, $C_a = \{|\zeta - a| = \varepsilon\}$, $C_b = \{|\zeta - b| = \varepsilon\}$. Then for $k > 1$,

$$\begin{aligned}
 & \int_{D \setminus \{D_a \cup D_b\}} \frac{(\zeta - b)^{k-1} d\bar{\zeta} \wedge d\zeta}{(\zeta - a)(\bar{\zeta} - \bar{b})} \\
 &= \int_D d \left(\frac{(\zeta - b)^{k-1} \ln |\zeta - b|^2 d\zeta}{\zeta - a} \right) \\
 &\quad - \int_{D_a} d \left(\frac{(\zeta - b)^{k-1} \ln |\zeta - b|^2 d\zeta}{\zeta - a} \right) \\
 &\quad - \int_{D_b} d \left(\frac{(\zeta - b)^{k-1} \ln |\zeta - b|^2 d\zeta}{\zeta - a} \right) \\
 (4) \quad &= \int_C \frac{(\zeta - b)^{k-1} \ln |\zeta - b|^2 d\zeta}{\zeta - a} \\
 &\quad - \int_{C_a} \frac{(\zeta - b)^{k-1} \ln |\zeta - b|^2 d\zeta}{\zeta - a} \\
 &\quad - \int_{C_b} \frac{(\zeta - b)^{k-1} \ln |\zeta - b|^2 d\zeta}{\zeta - a},
 \end{aligned}$$

where the last equation comes from Stokes formula.

$$\begin{aligned}
 (5) \quad & \int_{D_a} \frac{(\zeta - b)^{k-1} d\bar{\zeta} \wedge d\zeta}{(\zeta - a)(\bar{\zeta} - \bar{b})} \\
 &= 2i \int_0^{2\pi} \int_0^\varepsilon \frac{(re^{i\theta} + a - b)^{k-1} r dr d\theta}{re^{i\theta}(re^{-i\theta} + \bar{a} - \bar{b})} \\
 &= 2i \int_0^{2\pi} \int_0^\varepsilon \frac{(re^{i\theta} + a - b)^{k-1} r dr d\theta}{r + (\bar{a} - \bar{b})e^{i\theta}} \\
 &= 2i \int_0^{2\pi} \int_0^\varepsilon \frac{e^{i(k-1)\theta} (r + (\bar{a} - \bar{b})e^{i\theta} - (\bar{a} - \bar{b})e^{i\theta} + a - b)^{k-1}}{r + (\bar{a} - \bar{b})e^{i\theta}} dr d\theta \\
 &= 2i \int_0^{2\pi} \int_0^\varepsilon e^{i(k-1)\theta} \sum_{l=0}^{k-1} \binom{k-1}{l} (r + (\bar{a} - \bar{b})e^{i\theta})^{l-1} (- (\bar{a} - \bar{b})e^{i\theta} + a - b)^{k-1-l} dr d\theta \\
 &= 2i \int_0^{2\pi} \int_0^\varepsilon e^{i(k-1)\theta} (r + (\bar{a} - \bar{b})e^{i\theta})^{-1} (- (\bar{a} - \bar{b})e^{i\theta} + a - b)^{k-1} dr d\theta \\
 &\quad + 2i \int_0^{2\pi} \int_0^\varepsilon e^{i(k-1)\theta} \sum_{l=1}^{k-1} \binom{k-1}{l} (r + (\bar{a} - \bar{b})e^{i\theta})^{l-1} (- (\bar{a} - \bar{b})e^{i\theta} + a - b)^{k-1-l} dr d\theta \\
 &= 2i \int_0^{2\pi} e^{i(k-1)\theta} (- (\bar{a} - \bar{b})e^{i\theta} + a - b)^{k-1} \ln (r + (\bar{a} - \bar{b})e^{i\theta})^{-1} \Big|_0^\varepsilon d\theta \\
 &\quad + 2i \int_0^{2\pi} \int_0^\varepsilon \frac{e^{i(k-1)\theta}}{l} \sum_{l=1}^{k-1} \binom{k-1}{l} (- (\bar{a} - \bar{b})e^{i\theta} + a - b)^{k-1-l} (r + (\bar{a} - \bar{b})e^{i\theta})^l \Big|_0^\varepsilon d\theta,
 \end{aligned}$$

which converges to 0 when $\varepsilon \rightarrow 0$.

$$\begin{aligned}
 (6) \quad & \int_{D_b} \frac{(\zeta - b)^{k-1} d\bar{\zeta} \wedge d\zeta}{(\zeta - a)(\bar{\zeta} - \bar{b})} \\
 &= 2i \int_0^{2\pi} \int_0^\varepsilon \frac{(re^{i\theta})^{k-1} r dr d\theta}{re^{-i\theta}(re^{i\theta} + b - a)} \\
 &= 2i \int_0^{2\pi} \int_0^\varepsilon \frac{(re^{i\theta} + (b - a) - (b - a))^{k-1} dr d\theta}{r + (b - a)e^{-i\theta}} \\
 &= 2i \int_0^{2\pi} \int_0^\varepsilon \frac{e^{i(k-1)\theta} (r + (b - a)e^{-i\theta} - (b - a)e^{-i\theta})^{k-1}}{r + (b - a)e^{-i\theta}} dr d\theta \\
 &= 2i \int_0^{2\pi} \int_0^\varepsilon e^{i(k-1)\theta} \sum_{l=0}^{k-1} \binom{k-1}{l} (r + (b - a)e^{-i\theta})^{l-1} (-(b - a)e^{-i\theta})^{k-1-l} dr d\theta \\
 &= 2i \int_0^{2\pi} \int_0^\varepsilon e^{i(k-1)\theta} (r + (b - a)e^{-i\theta})^{-1} (-(b - a)e^{-i\theta})^{k-1} dr d\theta \\
 &\quad + 2i \int_0^{2\pi} \int_0^\varepsilon e^{i(k-1)\theta} \sum_{l=1}^{k-1} \binom{k-1}{l} (r + (b - a)e^{-i\theta})^{l-1} (-(b - a)e^{-i\theta})^{k-1-l} dr d\theta \\
 &= 2i \int_0^{2\pi} e^{i(k-1)\theta} (-(b - a)e^{-i\theta})^{k-1} \ln(r + (b - a)e^{-i\theta})^{-1} \Big|_0^\varepsilon d\theta \\
 &\quad + 2i \int_0^{2\pi} \int_0^\varepsilon \frac{e^{i(k-1)\theta}}{l} \sum_{l=1}^{k-1} \binom{k-1}{l} (-(b - a)e^{-i\theta})^{k-1-l} (r + (b - a)e^{-i\theta})^l \Big|_0^\varepsilon d\theta,
 \end{aligned}$$

which also converges to 0 when $\varepsilon \rightarrow 0$.

$$\begin{aligned}
 (7) \quad & \int_{C_a} \frac{(\zeta - b)^{k-1} \ln |\zeta - b|^2 d\zeta}{\zeta - a} \\
 &= \int_0^{2\pi} \frac{(\varepsilon e^{i\theta} + a - b)^{k-1} \ln |\varepsilon e^{i\theta} + a - b|^2 \varepsilon e^{i\theta} i d\theta}{\varepsilon e^{i\theta}} \\
 &= \int_0^{2\pi} (\varepsilon e^{i\theta} + a - b)^{k-1} \ln |\varepsilon e^{i\theta} + a - b|^2 i d\theta,
 \end{aligned}$$

which converges to $2\pi i(a - b)^{k-1} \ln |a - b|^2$ when $\varepsilon \rightarrow 0$.

$$\begin{aligned}
 (8) \quad & \int_{C_b} \frac{(\zeta - b)^{k-1} \ln |\zeta - b|^2 d\zeta}{\zeta - a} \\
 &= \int_0^{2\pi} \frac{(\varepsilon e^{i\theta})^{k-1} \ln |\varepsilon|^2 \varepsilon e^{i\theta} i d\theta}{\varepsilon e^{i\theta} + b - a} \\
 &= \int_0^{2\pi} \frac{(\varepsilon e^{i\theta})^k \ln |\varepsilon|^2 i d\theta}{\varepsilon e^{i\theta} + b - a},
 \end{aligned}$$

which converges to 0 when $\varepsilon \rightarrow 0$.

Thus, it comes from (4)-(8) that

$$\begin{aligned}
 & \int_D \frac{(\zeta - b)^{k-1} d\bar{\zeta} \wedge d\zeta}{(\zeta - a)(\bar{\zeta} - \bar{b})} \\
 &= \int_C \frac{(\zeta - b)^{k-1} \ln|\zeta - b|^2 d\zeta}{\zeta - a} - 2\pi i(a - b)^{k-1} \ln|a - b|^2 \\
 (9) \quad &= \int_C \frac{(\zeta - b)^{k-1} \left(\ln|\zeta|^2 + \ln\left(1 - \frac{b}{\zeta}\right) + \ln\left(1 - \frac{\bar{b}}{\bar{\zeta}}\right) \right)}{\zeta - a} d\zeta - 2\pi i(a - b)^{k-1} \ln|a - b|^2 \\
 &= \int_C \frac{(\zeta - b)^{k-1} \ln R^2}{\zeta - a} d\zeta + \int_C \frac{(\zeta - b)^{k-1} \ln\left(1 - \frac{b}{\zeta}\right)}{\zeta - a} d\zeta \\
 &\quad + \int_C \frac{(\zeta - b)^{k-1} \ln\left(1 - \frac{\bar{b}}{\bar{\zeta}}\right)}{\zeta - a} d\zeta - 2\pi i(a - b)^{k-1} \ln|a - b|^2 \\
 &= 2\pi i(a - b)^{k-1} \ln R^2 + I_1 + I_2 - 2\pi i(a - b)^{k-1} \ln|a - b|^2.
 \end{aligned}$$

I_1 is given as follows,

$$\begin{aligned}
 I_1 &= \int_C \frac{(\zeta - b)^{k-1} \ln\left(1 - \frac{b}{\zeta}\right)}{\zeta - a} d\zeta \\
 &= \int_C \frac{(\zeta - b)^{k-1}}{\zeta - a} \sum_{l=1}^{\infty} \frac{-1}{l} \left(\frac{b}{\zeta}\right)^l d\zeta \\
 (10) \quad &= \sum_{l=1}^{\infty} \frac{-b^l}{l} \int_C \frac{(\zeta - b)^{k-1}}{\zeta - a} \left(\frac{1}{\zeta}\right)^l d\zeta \\
 &= 2\pi i \sum_{l=1}^{k-1} \left(\frac{-b^l}{l}\right) \left(\sum_{j=0}^{k-1-l} \binom{k-1}{j} a^{k-1-j} (-b)^j \right).
 \end{aligned}$$

In fact, we have applied Residue theorem to get the last equation. Consider $\phi(\zeta) = \frac{(\zeta - b)^{k-1}}{\zeta - a} \left(\frac{1}{\zeta}\right)^l$. The integral $\int_C \frac{(\zeta - b)^{k-1}}{\zeta - a} \left(\frac{1}{\zeta}\right)^l d\zeta$ equals $-2\pi i$ times the residue of ϕ at ∞ which is the opposite number of the coefficient of $\frac{1}{\zeta}$ in the Rolland expansion of $\phi(\zeta)$.

$$\begin{aligned}
 \frac{(\zeta - b)^{k-1}}{\zeta - a} \left(\frac{1}{\zeta}\right)^l &= \frac{(\zeta - b)^{k-1}}{\zeta\left(1 - \frac{a}{\zeta}\right)} \left(\frac{1}{\zeta}\right)^l \\
 &= \frac{1}{\zeta^{l+1}} \left(\sum_{p=0}^{\infty} \left(\frac{a}{\zeta}\right)^p \right) \left(\sum_{q=0}^{k-1} \binom{k-1}{q} \zeta^q (-b)^{k-1-q} \right),
 \end{aligned}$$

which implies that there is no $\frac{1}{\zeta}$ in the expansion unless $l \leq k - 1$. Thus for any $1 \leq l \leq k - 1$, we get the coefficient of $\frac{1}{\zeta}$ in the Rolland expansion of $\phi(\zeta)$ as

$$\sum_{j=0}^{k-1-l} \binom{k-1}{j} a^{k-1-j} (-b)^j.$$

Hence,

$$\int_C \frac{(\zeta - b)^{k-1}}{\zeta - a} \left(\frac{1}{\zeta}\right)^l d\zeta = 2\pi i \sum_{j=0}^{k-1-l} \binom{k-1}{j} a^{k-1-j} (-b)^j,$$

and (10) can be obtained.

On the other hand, by Cauchy integral, we can obtain I_2 as follows,

$$\begin{aligned}
 (11) \quad I_2 &= \int_C \frac{(\zeta - b)^{k-1} \ln(1 - \frac{\bar{b}}{\zeta})}{\bar{\zeta} - a} d\zeta \\
 &= \int_C \frac{(\zeta - b)^{k-1}}{\zeta - a} \sum_{l=1}^{\infty} \frac{-1}{l} \left(\frac{\bar{b}}{\bar{\zeta}}\right)^l d\zeta \\
 &= \int_C \frac{(\zeta - b)^{k-1}}{\zeta - a} \sum_{l=1}^{\infty} \frac{-1}{l} \left(\frac{\bar{b}}{R^2}\right)^l \zeta^l d\zeta \\
 &= \sum_{l=1}^{\infty} \frac{-1}{l} \left(\frac{\bar{b}}{R^2}\right)^l \int_C \frac{(\zeta - b)^{k-1}}{\zeta - a} \zeta^l d\zeta \\
 &= 2\pi i \sum_{l=1}^{\infty} \frac{-1}{l} \left(\frac{\bar{b}}{R^2}\right)^l (a - b)^{k-1} a^l \\
 &= 2\pi i (a - b)^{k-1} \ln\left(1 - \frac{a\bar{b}}{R^2}\right).
 \end{aligned}$$

Therefore, from (9), (10), and (11), we prove that

$$\begin{aligned}
 &\int_C \frac{(\zeta - b)^{k-1} \ln |\zeta - b|^2}{\zeta - a} d\zeta \\
 &= 2\pi i \sum_{l=1}^{k-1} \left(\frac{-b^l}{l}\right) \left(\sum_{j=0}^{k-1-l} \binom{k-1}{j} a^{k-1-j} (-b)^j\right) + 2\pi i (a - b)^{k-1} \ln(R^2 - a\bar{b}).
 \end{aligned}$$

For $k = 1$, one can easily yield from the above proof that $I_1 = 0$. Combining (9), we get the lemma. \square

LEMMA 2.4. For any $a, b \in D$ and $a \neq b, l > 0, \nu > 0$,

$$\int_C \frac{(\bar{\zeta} - \bar{b})^l (\zeta - b)^{\nu-1}}{\zeta - a} d\zeta = 2\pi i C_2(a, b, l, \nu),$$

where $C_2(a, b, l, \nu) = \sum_{0 \leq p \leq l, 0 \leq q \leq \nu-1, p \leq q} \binom{l}{p} \binom{\nu-1}{q} R^{2p} (-\bar{b})^{l-p} (-b)^{\nu-1-q} a^{q-p}$.

Proof. The computation is based on Residue theorem.

$$\begin{aligned}
 & \int_C \frac{(\bar{\zeta} - \bar{b})^l (\zeta - b)^{\nu-1}}{\zeta - a} d\zeta \\
 &= \int_C \frac{1}{\zeta - a} \left(\sum_{p=0}^l \binom{l}{p} \bar{\zeta}^p (-\bar{b})^{l-p} \right) \left(\sum_{q=0}^{\nu-1} \binom{\nu-1}{q} \zeta^q (-b)^{\nu-1-q} \right) d\zeta \\
 &= \int_C \frac{1}{\zeta - a} \left(\sum_{p=0}^l \binom{l}{p} \frac{R^{2p}}{\zeta^p} (-\bar{b})^{l-p} \right) \left(\sum_{q=0}^{\nu-1} \binom{\nu-1}{q} \zeta^q (-b)^{\nu-1-q} \right) d\zeta \\
 (12) \quad &= \int_C \frac{1}{\zeta - a} \left(\sum_{p=0}^l \sum_{q=0}^{\nu-1} \binom{l}{p} \binom{\nu-1}{q} R^{2p} (-\bar{b})^{l-p} (-b)^{\nu-1-q} \zeta^{q-p} \right) d\zeta \\
 &= \sum_{p=0}^l \sum_{q=0}^{\nu-1} \binom{l}{p} \binom{\nu-1}{q} R^{2p} (-\bar{b})^{l-p} (-b)^{\nu-1-q} \int_C \frac{1}{\zeta - a} \zeta^{q-p} d\zeta \\
 &= 2\pi i \sum_{0 \leq p \leq l, 0 \leq q \leq \nu-1, p \leq q} \binom{l}{p} \binom{\nu-1}{q} R^{2p} (-\bar{b})^{l-p} (-b)^{\nu-1-q} a^{q-p},
 \end{aligned}$$

since if $q - p < 0$, $\int_C \frac{1}{\zeta - a} \zeta^{q-p} d\zeta = 0$ and if $q - p \geq 0$, from the Residue theorem, $\int_C \frac{1}{\zeta - a} \zeta^{q-p} d\zeta = 2\pi i a^{q-p}$. \square

LEMMA 2.5. For any $a, b \in D$ and $a \neq b, \mu, \nu > 0$,

$$\int_D \frac{(\bar{\zeta} - \bar{a})^{\mu-1} (\zeta - b)^{\nu-1} d\bar{\zeta} \wedge d\zeta}{(\zeta - a)(\bar{\zeta} - \bar{b})} = 2\pi i C_3(a, b, \mu, \nu),$$

where $C_3(a, b, \mu, \nu) = (\bar{b} - \bar{a})^{\mu-1} \left(C_1(a, b, \nu) + (a - b)^{\nu-1} \ln \frac{R^2 - a\bar{b}}{|a-b|^2} \right) + \sum_{l=1}^{\mu-1} \binom{\mu-1}{l} (\bar{b} - \bar{a})^{\mu-1-l} \left(\frac{-1}{l} (\bar{a} - \bar{b})^l (a - b)^{\nu-1} - \frac{1}{l} C_2(a, b, l, \nu) \right)$ for $\mu > 1$ and $C_3(a, b, 1, \nu) = C_1(a, b, \nu) + (a - b)^{\nu-1} \ln \frac{R^2 - a\bar{b}}{|a-b|^2}$.

Proof. For $\mu > 1$, we have

$$\begin{aligned}
 & \int_D \frac{(\bar{\zeta} - \bar{a})^{\mu-1}(\zeta - b)^{\nu-1}d\bar{\zeta} \wedge d\zeta}{(\zeta - a)(\bar{\zeta} - \bar{b})} \\
 &= \int_D \frac{(\bar{\zeta} - \bar{b} + \bar{b} - \bar{a})^{\mu-1}(\zeta - b)^{\nu-1}d\bar{\zeta} \wedge d\zeta}{(\zeta - a)(\bar{\zeta} - \bar{b})} \\
 &= \int_D \frac{\sum_{l=0}^{\mu-1} \binom{\mu-1}{l} (\bar{\zeta} - \bar{b})^l (\bar{b} - \bar{a})^{\mu-1-l} (\zeta - b)^{\nu-1} d\bar{\zeta} \wedge d\zeta}{(\zeta - a)(\bar{\zeta} - \bar{b})} \\
 &= (\bar{b} - \bar{a})^{\mu-1} \int_D \frac{(\zeta - b)^{\nu-1} d\bar{\zeta} \wedge d\zeta}{(\zeta - a)(\bar{\zeta} - \bar{b})} \\
 &\quad + \sum_{l=1}^{\mu-1} \binom{\mu-1}{l} (\bar{b} - \bar{a})^{\mu-1-l} \int_D \frac{(\bar{\zeta} - \bar{b})^l (\zeta - b)^{\nu-1} d\bar{\zeta} \wedge d\zeta}{(\zeta - a)(\bar{\zeta} - \bar{b})} \\
 &= (\bar{b} - \bar{a})^{\mu-1} \int_D \frac{(\zeta - b)^{\nu-1} d\bar{\zeta} \wedge d\zeta}{(\zeta - a)(\bar{\zeta} - \bar{b})} \\
 &\quad + \sum_{l=1}^{\mu-1} \binom{\mu-1}{l} (\bar{b} - \bar{a})^{\mu-1-l} \int_D \frac{(\bar{\zeta} - \bar{b})^{l-1} (\zeta - b)^{\nu-1} d\bar{\zeta} \wedge d\zeta}{\zeta - a} \\
 &= I_3 + I_4.
 \end{aligned}$$

From Lemma 2.3,

$$I_3 = 2\pi i (\bar{b} - \bar{a})^{\mu-1} \left(C_1(a, b, \nu) + (a - b)^{\nu-1} \ln \frac{R^2 - a\bar{b}}{|a - b|^2} \right).$$

For $f(\zeta) = \frac{1}{l} (\bar{\zeta} - \bar{b})^l (\zeta - b)^{\nu-1}$, $\bar{\partial}f(\zeta) = (\bar{\zeta} - \bar{b})^{l-1} (\zeta - b)^{\nu-1}$. From Lemma 1.2,

$$\begin{aligned}
 I_4 &= \sum_{l=1}^{\mu-1} \binom{\mu-1}{l} (\bar{b} - \bar{a})^{\mu-1-l} \int_D \frac{(\bar{\zeta} - \bar{b})^{l-1} (\zeta - b)^{\nu-1} d\bar{\zeta} \wedge d\zeta}{\zeta - a} \\
 &= \sum_{l=1}^{\mu-1} \binom{\mu-1}{l} (\bar{b} - \bar{a})^{\mu-1-l} \int_D \frac{\bar{\partial}f(\zeta)}{\zeta - a} d\bar{\zeta} \wedge d\zeta \\
 &= -2\pi i \sum_{l=1}^{\mu-1} \binom{\mu-1}{l} (\bar{b} - \bar{a})^{\mu-1-l} T(\bar{\partial}f(\zeta))(a) \\
 &= -2\pi i \sum_{l=1}^{\mu-1} \binom{\mu-1}{l} (\bar{b} - \bar{a})^{\mu-1-l} (f(a) - S(f(\zeta)(a))) \\
 &= 2\pi i \sum_{l=1}^{\mu-1} \binom{\mu-1}{l} (\bar{b} - \bar{a})^{\mu-1-l} \left(\frac{-1}{l} (\bar{a} - \bar{b})^l (a - b)^{\nu-1} - \frac{1}{l \cdot 2\pi i} \int_C \frac{(\bar{\zeta} - \bar{b})^l (\zeta - b)^{\nu-1} d\zeta}{\zeta - a} \right) \\
 &= 2\pi i \sum_{l=1}^{\mu-1} \binom{\mu-1}{l} (\bar{b} - \bar{a})^{\mu-1-l} \left(\frac{-1}{l} (\bar{a} - \bar{b})^l (a - b)^{\nu-1} - \frac{1}{l} C_2(a, b, l, \nu) \right),
 \end{aligned}$$

where the last equation comes from Lemma 2.4. Furthermore, it is easy to see $I_4 = 0$ for $\mu = 1$. Let $C_3(a, b, \mu, \nu) = (\bar{b} - \bar{a})^{\mu-1} \left(C_1(a, b, \nu) + (a - b)^{\nu-1} \ln \frac{R^2 - a\bar{b}}{|a - b|^2} \right) + \sum_{l=1}^{\mu-1} \binom{\mu-1}{l} (\bar{b} - \bar{a})^{\mu-1-l} \left(\frac{-1}{l} (\bar{a} - \bar{b})^l (a - b)^{\nu-1} - \frac{1}{l} C_2(a, b, l, \nu) \right)$. Then we prove the lemma. \square

THEOREM 2.6. Given $f \in C^\alpha(D)$, $T^\mu \bar{T}^\nu f \in C^{\mu+\nu+\alpha}(D)$ with $\mu, \nu > 0$, and

$$T^\mu \bar{T}^\nu f(z) = \frac{(-1)^\mu}{(\mu - 1)!(\nu - 1)! \cdot 2\pi i} \int_D C_3(z, \eta, \mu, \nu) f(\eta) d\bar{\eta} \wedge d\eta,$$

where the kernel function C_3 is given by Lemma 2.5.

Proof. Given $f \in C^\alpha(D)$, it is obvious that $T^\mu \bar{T}^\nu f \in C^{\mu+\nu+\alpha}(D)$ by Lemma 1.2. From Theorem 2.1, we have

$$\begin{aligned} T^\mu \bar{T}^\nu f(z) &= \frac{(-1)^\mu}{(\mu - 1)! \cdot 2\pi i} \int_D \frac{(\bar{\zeta} - \bar{z})^{\mu-1} \bar{T}^\nu f(\zeta)}{\zeta - z} d\bar{\zeta} \wedge d\zeta \\ &= \frac{(-1)^\mu}{(\mu - 1)! \cdot 2\pi i} \int_D \frac{(\bar{\zeta} - \bar{z})^{\mu-1} \frac{(-1)^\nu}{(\nu-1)! \cdot 2\pi i} \int_D \frac{(n-\zeta)^{\nu-1} f(\eta)}{\eta - \zeta} d\bar{\eta} \wedge d\eta}{\zeta - z} d\bar{\zeta} \wedge d\zeta \\ &= \frac{(-1)^{\mu+\nu}}{(\mu - 1)!(\nu - 1)! \cdot (2\pi i)^2} \int_D \int_D \frac{(\bar{\zeta} - \bar{z})^{\mu-1} (\eta - \zeta)^{\nu-1} d\bar{\zeta} \wedge d\zeta}{(\zeta - z)(\bar{\eta} - \bar{\zeta})} f(\eta) d\bar{\eta} \wedge d\eta \\ &= \frac{(-1)^{\mu+\nu} (-1)^\nu}{(\mu - 1)!(\nu - 1)! \cdot (2\pi i)^2} \int_D \int_D \frac{(\bar{\zeta} - \bar{z})^{\mu-1} (\zeta - \eta)^{\nu-1} d\bar{\zeta} \wedge d\zeta}{(\zeta - z)(\bar{\zeta} - \bar{\eta})} f(\eta) d\bar{\eta} \wedge d\eta \\ &= \frac{(-1)^\mu}{(\mu - 1)!(\nu - 1)! \cdot 2\pi i} \int_D C_3(z, \eta, \mu, \nu) f(\eta) d\bar{\eta} \wedge d\eta. \end{aligned}$$

□

REMARK 2.7. From above theorem, we have $\bar{T}^\mu T^\nu \bar{f}(z) = \overline{T^\mu \bar{T}^\nu f(z)}$.

REMARK 2.8. Theorem 2.6 presents the expression of high-order Green operator and its kernel function on the disk. Here we give some special cases to shown the kernel in detail.

1. $C_3(z, \eta, 1, 1) = \ln \frac{R^2 - z\bar{\eta}}{|z - \eta|^2}$;
2. $C_3(z, \eta, 1, 2) = -z\eta + (z - \eta) \ln \frac{R^2 - z\bar{\eta}}{|z - \eta|^2}$;
3. $C_3(z, \eta, 2, 1) = (\bar{\eta} - \bar{z}) \ln \frac{R^2 - z\bar{\eta}}{|z - \eta|^2} + 2\bar{\eta} - \bar{z}$;
4. $C_3(z, \eta, 2, 2) = (\bar{\eta} - \bar{z}) (-z\eta + (z - \eta) \ln \frac{R^2 - z\bar{\eta}}{|z - \eta|^2}) - |z - \eta|^2 - |\eta|^2 + \bar{\eta}z - R^2$.

[7] has presented the relationship between the corresponding norms for f and $T^\mu \bar{T}^\nu f$ in the above theorem. We omit the proof here.

PROPOSITION 2.9. [7] If $f \in C^\alpha(D)$ and $\mu + \nu = m$, then

$$\|T^\mu \bar{T}^\nu f\|^{(m)} \leq 2^{\frac{(m-1)m}{2}} (C_4 m + C_0 + (m - 1)C_5)^m \|f\|,$$

where $C_0 = \frac{12}{\alpha(1-\alpha)}$, $C_4 = \frac{2^{\alpha+1}}{\alpha}$ and $C_5 = \frac{4}{\alpha(1-\alpha)}$.

3. High-order Green operator on D^n . Let D^n be the n -fold cartesian product of D , which is a closed ploydisc in \mathbb{C}^n with radius R . Suppose that f is a complex-valued function defined on D^n . We define $\Delta_i f$ as a function on that subset D_i of the $(n + 1)$ -fold product $D \times \dots \times D$ whose points $(z_1, \dots, z_{i-1}, (z_i, z'_i), z_{i+1}, \dots, z_n)$ satisfy $z_i \neq z'_i$. Then $\Delta_i f(z_1, \dots, z_{i-1}, (z_i, z'_i), z_{i+1}, \dots, z_n) := f(z_1, \dots, z_{i-1}, z_i, z_{i+1}, \dots, z_n) - f(z_1, \dots, z_{i-1}, z'_i, z_{i+1}, \dots, z_n)$. For any distinct integers $i_1, \dots, i_k \in \{1, \dots, n\}$, we define

$\Delta_{i_1 \dots i_k} f := \Delta_{i_k} \Delta_{i_1 \dots i_{k-1}} f$. For simplicity, denote $Z_{i_1 \dots i_k} = (z_1, \dots, (z_{i_1}, z'_{i_1}), \dots, (z_{i_k}, z'_{i_k}), \dots, z_n)$ satisfy $z_{i_j} \neq z'_{i_j}$, $j = 1, \dots, k$. A k th-order α -Hölder difference quotient is any expression $\delta_{i_1 \dots i_k} f$ defined by

$$\delta_{i_1 \dots i_k} f(Z_{i_1 \dots i_k}) = \frac{\Delta_{i_1 \dots i_k} f(Z_{i_1 \dots i_k})}{|z_{i_1} - z'_{i_1}|^\alpha \dots |z_{i_k} - z'_{i_k}|^\alpha}.$$

Set

$$H_\alpha^{(k)}[f] = \max\{|\delta_{i_1 \dots i_k} f| \mid i_1, \dots, i_k \text{ distinct}\}.$$

We define $\mathcal{C}^\alpha(D^n)$ as the set of those functions f defined on D^n for which that $H_\alpha^{(k)}[f]$ are finite for all $k = 0, \dots, n$, where $H_\alpha^{(0)}[f] = |f|$.

Define $\|\cdot\|$ on the space $\mathcal{C}^\alpha(D^n)$ by

$$\|f\| := \sum_{k=0}^n \frac{2R^{k\alpha}}{k!} H_\alpha^{(k)}[f].$$

It is proved in 7.1b of [5] that $\|\cdot\|$ defined on $\mathcal{C}^\alpha(D^n)$ is a norm.

Let $z = (z_1, \dots, z_n) \in D^n$, the following operators are defined on D^n :

$$T_j f(z) = \frac{-1}{2\pi i} \int_D \frac{f(z_1, \dots, z_{j-1}, \zeta, z_{j+1}, \dots, z_n) d\bar{\zeta} \wedge d\zeta}{\zeta - z_j},$$

$$\bar{T}_j f(z) = \frac{-1}{2\pi i} \int_D \frac{f(z_1, \dots, z_{j-1}, \zeta, z_{j+1}, \dots, z_n) d\bar{\zeta} \wedge d\zeta}{\bar{\zeta} - \bar{z}_j},$$

similar definitions are given for S_j, \bar{S}_j . From Lemma 1.1, it is easy to see that for any $f \in C^1(D^n)$,

$$T_j \bar{\partial}_j f = f - S_j f.$$

Furthermore, it is given in [5] that

LEMMA 3.1. *If $f \in \mathcal{C}^\alpha(D^n)$, then $T^i f \in \mathcal{C}^\alpha(D^n)$, $S^i f \in \mathcal{C}^\alpha(D^n)$ for all $i = 1, \dots, n$. And there exist constants C_6, C_7 such that*

$$\|T^i f\| \leq C_6 R \|f\|, \quad \|S^i f\| \leq C_7 \|f\|.$$

REMARK 3.2. *It should be noticed that the smoothness properties of the various integral operator T^i, \bar{T}^i , $i = 1, \dots, n$ defined for functions on D and D^n are different. T^i, \bar{T}^i are no longer smoothing order as in dimension one, see Lemma 1.2.*

Let vector index $\mu = (\mu_1, \dots, \mu_n)$, $\nu = (\nu_1, \dots, \nu_n)$, $\mathbf{1} = (1, \dots, 1) \in \mathbb{N}^n$. $|\mu| = \sum_{j=1}^n \mu_j$, $\mu! = \prod_{j=1}^n \mu_j!$. For $z \in D^n$, $f \in \mathcal{C}^\alpha(D^n)$, denote the operator $T^\mu \bar{T}^\nu$ as follows,

$$T^\mu \bar{T}^\nu f(z) = T_1^{\mu_1} \dots T_n^{\mu_n} \bar{T}_1^{\nu_1} \dots \bar{T}_n^{\nu_n} f(z).$$

From Theorem 2.6, it is easy to get the explicit expression and the kernel function of $T^\mu \bar{T}^\nu f$ on D^n .

COROLLARY 3.3. For $z \in D^n$, given $f \in \mathcal{C}^\alpha(D^n)$, then $T^\mu \bar{T}^\nu f \in \mathcal{C}^\alpha(D^n)$ and

$$T^\mu \bar{T}^\nu f(z) = C_8(\mu, \nu) \int_D \cdots \int_D \prod_{j=1}^n C_3(z_j, \eta_j, \mu_j, \nu_j) f(\eta) d\bar{\eta}_1 \wedge d\eta_1 \cdots d\bar{\eta}_n \wedge d\eta_n,$$

where $C_8(\mu, \nu) = \frac{(-1)^{|\mu|}}{(\mu-1)!(\nu-1)!(2\pi i)^n}$, and C_3 is given by Lemma 2.5.

Proof. By the definition of $T^\mu \bar{T}^\nu f(z)$,

$$\begin{aligned} T^\mu \bar{T}^\nu f(z) &= T_1^{\mu_1} \cdots T_n^{\mu_n} \bar{T}_1^{\nu_1} \cdots \bar{T}_n^{\nu_n} f(z) \\ &= T_1^{\mu_1} \bar{T}_1^{\nu_1} \cdots T_n^{\mu_n} \bar{T}_n^{\nu_n} f(z) \\ &= \frac{(-1)^{\mu_1}}{(\mu_1-1)!(\nu_1-1)! \cdot 2\pi i} \int_D C_3(z_1, \eta_1, \mu_1, \nu_1) \\ &\quad \times T_2^{\mu_2} \bar{T}_2^{\nu_2} \cdots T_n^{\mu_n} \bar{T}_n^{\nu_n} f(\eta_1, z_2, \dots, z_n) d\bar{\eta}_1 \wedge d\eta_1 \\ &= \frac{(-1)^{\mu_1+\mu_2}}{(\mu_1-1)!(\mu_2-1)!(\nu_1-1)!(\nu_2-1)! \cdot (2\pi i)^2} \\ &\quad \times \int_D \int_D C_3(z_1, \eta_1, \mu_1, \nu_1) C_3(z_2, \eta_2, \mu_2, \nu_2) \\ &\quad \times T_3^{\mu_3} \bar{T}_3^{\nu_3} \cdots T_n^{\mu_n} \bar{T}_n^{\nu_n} f(\eta_1, \eta_2, z_3, \dots, z_n) d\bar{\eta}_1 \wedge d\eta_1 \cdot d\bar{\eta}_2 \wedge d\eta_2 \\ &= \cdots \\ &= \frac{(-1)^{|\mu|}}{(\mu-1)!(\nu-1)! \cdot (2\pi i)^n} \\ &\quad \times \int_D \cdots \int_D \prod_{j=1}^n C_3(z_j, \eta_j, \mu_j, \nu_j) f(\eta) d\bar{\eta}_1 \wedge d\eta_1 \cdots d\bar{\eta}_n \wedge d\eta_n. \end{aligned}$$

□

By Lemma 3.1, for any $f \in \mathcal{C}^\alpha(D^n)$, one can easily get the following estimate for the norm of $f \in \mathcal{C}^\alpha(D^n)$.

PROPOSITION 3.4. If $f \in \mathcal{C}^\alpha(D^n)$ and $|\mu| + |\nu| = m$, then

$$\|T^\mu \bar{T}^\nu f\| \leq (C_6)^m \|f\|.$$

4. Applications. As applications of the integral expressions for high-order Green operators, we can present all the solutions for some high-order Laplace equations, moreover, express all the solutions for linear high-order partial differential equation with integrals.

Given $D = \{z \mid |z| \leq R\} = \{(x, y) \mid x^2 + y^2 \leq R^2\}$ as previous section. Let $A(x, y) : D \rightarrow \mathbb{R}$ be function of class $C^\alpha(D)$, $u(x, y) : D \rightarrow \mathbb{R}$ be unknown function. Since

$$\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} = 4\partial\bar{\partial}$$

and from Lemma 1.2

$$\bar{\partial}TA = A; \partial\bar{T}A = A,$$

then we have

$$\Delta^2 T^2 \bar{T}^2 A = 16A; \quad \Delta^2 \bar{T}^2 T^2 A = 16A.$$

$\Delta^2 u = 0$ is called biharmonic equation whose solutions can be described by $u = |z|^2 h_1 + h_2$, where h_1, h_2 are two harmonic functions satisfy Laplace's equation [8]. Then the solutions of 2-Laplace equation u :

$$\Delta^2 u(x, y) = A(x, y)$$

given by

$$\begin{aligned} u = \mathbf{Re} & \left(\frac{1}{32\pi i} \int_D [(\bar{\eta} - \bar{z})(-z\eta + (z - \eta) \ln \frac{R^2 - z\bar{\eta}}{|z - \eta|^2}) \right. \\ & \left. - |z - \eta|^2 - |\eta|^2 + \bar{\eta}z - R^2] A(\eta, \bar{\eta}) d\bar{\eta} \wedge d\eta \right) \\ & + |z|^2 h_1 + h_2. \end{aligned}$$

The integral expressions can also be used to give all the solutions for linear partial differential equations on D of any order. Let $H(D)$ be the set of all holomorphic functions on D and denote $T^0 f = f$. We have the following results.

LEMMA 4.1. *Given $\mu, \nu > 0$, the solutions of*

$$(13) \quad \partial^\mu \bar{\partial}^\nu u(z, \bar{z}) = 0$$

can be given by

$$u = \sum_{j=0}^{\nu-1} T^j(g_j) + T^\nu \left(\sum_{i=0}^{\mu-1} \bar{T}^i(\bar{f}_i) \right),$$

where $f_i, g_j \in H(D)$, $i = 0, \dots, \mu - 1$; $j = 0, \dots, \nu - 1$.

Proof. It is well known that all the solutions of $\bar{\partial} u = 0$ can be given by any $u \in H(D)$ and the solutions of $\partial u = 0$ can be given by u with any $\bar{u} \in H(D)$. Consider the equation (13), we have

$$\partial^{\mu-1} \bar{\partial}^\nu u = \bar{f}_0, \quad \forall f_0 \in H(D),$$

which means that

$$\partial^{\mu-2} \bar{\partial}^\nu u = \bar{f}_1 + \bar{T}(\bar{f}_0), \quad \forall f_0, f_1 \in H(D).$$

Similarly, we have

$$\partial^{\mu-3} \bar{\partial}^\nu u = f_2 + \bar{T}(\bar{f}_1) + \bar{T}^2(\bar{f}_0), \quad \forall f_0, f_1, f_2 \in H(D).$$

By iteration, one has

$$\bar{\partial}^\nu u = \sum_{i=0}^{\mu-1} \bar{T}^i(\bar{f}_{\mu-1-i}), \quad \forall f_i \in H(D), i = 0, \dots, \mu - 1.$$

Furthermore,

$$\bar{\partial}^{\nu-1}u = g_0 + T \sum_{i=0}^{\mu-1} \bar{T}^i(\bar{f}_{\mu-1-i}), \quad \forall g_0, f_i \in H(D), i = 0, \dots, \mu - 1,$$

and then

$$\bar{\partial}^{\nu-2}u = g_1 + T(g_0) + T^2 \sum_{i=0}^{\mu-1} \bar{T}^i(\bar{f}_{\mu-1-i}), \quad \forall g_0, g_1, f_i \in H(D), i = 0, \dots, \mu - 1.$$

By iteration, we can conclude

$$u = \sum_{j=0}^{\nu-1} T^j(g_{\nu-1-j}) + T^\nu \sum_{i=0}^{\mu-1} \bar{T}^i(\bar{f}_{\mu-1-i}), \quad \forall g_j, f_i \in H(D), i = 0, \dots, \mu - 1, j = 0, \dots, \nu - 1.$$

For simplicity, we replace the index and prove the lemma. Using Theorems 2.1 and 2.6, we can give the solutions of equation (13) by integral. \square

From Lemma 4.1, we can express all the solutions for linear high-order partial differential equation

$$\partial^\mu \bar{\partial}^\nu u(z, \bar{z}) = A(z, \bar{z})$$

as

$$(14) \quad u = \sum_{j=0}^{\nu-1} T^j(g_j) + T^\nu \sum_{i=0}^{\mu-1} \bar{T}^i(\bar{f}_i) + T^\nu \bar{T}^\mu(A),$$

where $g_j, f_i \in H(D), i = 0, \dots, \mu - 1, j = 0, \dots, \nu - 1$. Using Theorems 2.1 and 2.6, the integral expressions of solutions can be given.

For simplicity, we denote

$$(15) \quad G(z, \zeta, l) = \frac{(-1)^l (\bar{\zeta} - \bar{z})^{l-1}}{2\pi i (l-1)! (\zeta - z)};$$

$$G(z, \zeta, \mu, \nu) = \frac{(-1)^\mu}{2\pi i (\mu-1)! (\nu-1)!} C_3(z, \zeta, \mu, \nu).$$

From (14), we have

$$(16) \quad u = g_0 + \sum_{j=1}^{\nu-1} T^j(g_j) + T^\nu \bar{f}_0 + T^\nu \left(\sum_{i=1}^{\mu-1} \bar{T}^i(\bar{f}_i) \right) + T^\nu \bar{T}^\mu(A).$$

Combining (15) and (16), we have the following theorem.

THEOREM 4.2. *Given $\mu, \nu > 0$, the solutions of*

$$\partial^\mu \bar{\partial}^\nu u(z, \bar{z}) = A(z, \bar{z})$$

with $A(z, \bar{z}) \in C^\alpha(D)$ can be given by

$$u(z, \bar{z}) = g_0(z) + \int_D \left(\sum_{j=1}^{\nu-1} G(z, \zeta, j) g_j(\zeta) + G(z, \zeta, \nu) \bar{f}_0(\zeta) \right. \\ \left. + \sum_{i=1}^{\mu-1} G(z, \zeta, \nu, i) \bar{f}_i(\zeta) + G(z, \zeta, \nu, \mu) A(\zeta, \bar{\zeta}) \right) d\bar{\zeta} \wedge d\zeta,$$

where $f_i, g_j \in H(D), i = 0, \dots, \mu - 1; j = 0, \dots, \nu - 1$.

5. Conclusion. We have established the explicit expressions for high-order Green operators on the disk in \mathbb{C} and the polydisc in \mathbb{C}^n . As applications, we have presented all the solutions for biharmonic equations and high-order partial differential equations in the disk. The same method works identically in \mathbb{R}^n through Clifford analysis and the results will be presented in the forthcoming paper in a near future.

REFERENCES

- [1] M. CHRIST, *Lectures on singular integral operators*, CBMS Regional Conference Series in Mathematics, Washington, DC; Providence, RI, 1990.
- [2] V. KOKILASHVILI, *Boundedness criterion for the Cauchy singular integral operator in weighted grand Lebesgue spaces and application to the Riemann problem*, Proc. A. Razmadze Math. Inst., 151 (2009), pp. 129–133.
- [3] A. BÖTTCHER AND I. GOHBERG, *Singular integral operators and related topics*, Birkhäuser Verlag, Basel, 1996.
- [4] V. G. KRAVCHENKO AND G. S. LITVINCHUK, *Introduction to the theory of singular integral operators with shift*, (English summary) Kluwer Academic Publishers Group, Dordrecht, 1994.
- [5] A. NIJENHUIS AND W. WOOLF, *Some integration problems in almost-complex and complex manifolds*, Ann. of Math., 77:3 (1963), pp. 426–489.
- [6] A. NEULANDER AND L. NIRENBERG, *Complex analytic coordinates in almost complex manifolds*, Ann. of Math., 65:6 (1957), pp. 391–404.
- [7] Y. PAN, *On solvability of nonlinear partial differential systems of any order in the complex plane*, arXiv:1109.3325v2.
- [8] R. BHARGAVA, *Solution of a biharmonic equation*, Nature, 201(4918), 530, 1964.