

## AN APPLICATION OF THE RESULTANT TO OSCILLATORY INTEGRALS WITH POLYNOMIAL PHASE

DANIEL M. OBERLIN

Fix  $n = 2, 3, \dots$  and let  $p(t) = \sum_{j=1}^n a_j t^j$  be a polynomial with real coefficients. Let  $I \subseteq \mathbb{R}$  be an interval. This paper is concerned with estimates of the form

$$(1) \quad \left| \int_I e^{ip(t)} dt \right| \leq M(a_1, \dots, a_n),$$

where  $M(a_1, \dots, a_n)$  is independent of  $I$ . Let  $M_0(a_1, \dots, a_n)$  be the supremum over all  $I$  of the left hand side of (1). If  $q(t) = p(st)$  for  $s > 0$ , then

$$\int_a^b e^{iq(t)} dt = s^{-1} \int_{as}^{bs} e^{ip(t)} dt.$$

It follows that the homogeneity condition

$$(2) \quad M(sa_1, s^2 a_2, \dots, s^n a_n) = s^{-1} M(a_1, \dots, a_n), \quad s > 0,$$

is satisfied if  $M(a_1, \dots, a_n) = M_0(a_1, \dots, a_n)$ . Our interest here is in estimates (1) for which this natural equality (2) holds.

If  $1 \leq j_1 < j_2 \leq n$  and the derivatives  $p^{(j_1)}$  and  $p^{(j_2)}$  have no common zero in  $\mathbb{R}$ , then van der Corput's lemma ([S], p. 332) shows that

$$(3) \quad \left| \int_I e^{i\lambda p(t)} dt \right|$$

$O(|\lambda|^{-\frac{1}{j_2}})$  as the real parameter  $\lambda$  tends to infinity. We would like our estimates (1) to recover such information, with bounds on the relevant "big oh" constants as well. The paper [O2] contains such estimates in case either  $n \leq 4$  or  $j_2 = n - 1$ . (The case  $j_2 = n$  is covered by van der Corput's lemma.) Here we obtain, for each  $n$  and  $1 \leq j_1 < j_2 \leq n$ , an estimate (1) with right hand side  $M(a_1, \dots, a_n) = M_{j_1 j_2 n}(a_1, \dots, a_n)$  satisfying (2). If  $p^{(j_1)}$  and  $p^{(j_2)}$  have no common complex zero then  $M_{j_1 j_2 n}(\lambda a_1, \dots, \lambda a_n)$  is  $O(|\lambda|^{-(2n-j_1-j_2-1)/(n^2-j_1 j_2-n)})$  as  $|\lambda| \rightarrow \infty$

---

Received May 1, 1997.

Partially supported by a grant from the National Science Foundation.

when  $a_n \neq 0$ . If  $j_2 < n - 1$  this is weaker than the desired estimate of  $O(|\lambda|^{-\frac{1}{j_2}})$  for (3), even with the stronger hypothesis of no common complex, as opposed to real, zero. Still, our estimates here recover Theorem 1 and Corollary 3 of [O2] as special cases and, more importantly to our purposes, provide an extension valid for polynomials of degree  $n$  of that Corollary 3. (Corollary 3 of [O2] furnishes a Fourier transform estimate required for the proof of a convolution theorem dealing with measures on the curve  $(t, t^2, t^3, t^4)$  in  $\mathbb{R}^4$  - see [O1]. Its extension is the first step in our attempt to extend the work of [O1] to the  $n$ -dimensional setting.)

We begin by recalling the definition of the resultant of two polynomials. If  $q(t) = \sum_{\ell=0}^j b_\ell t^\ell$  and  $r(t) = \sum_{\ell=0}^k c_\ell t^\ell$  are polynomials, their resultant  $R = R(q, r)$  is the polynomial in the variables  $b_\ell$  and  $c_\ell$  defined by the determinant of the  $(j + k)$  by  $(j + k)$  matrix:

$$\begin{matrix} k \\ j \end{matrix} \left\{ \begin{matrix} \begin{matrix} b_j & b_{j-1} & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & b_0 \\ & b_j & b_{j-1} & \cdot & \cdot & \cdot & \cdot & \cdot & b_1 & b_0 \\ & & b_j & b_{j-1} & \cdot & \cdot & \cdot & \cdot & \cdot & b_1 & b_0 \\ & & & & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ c_k & c_{k-1} & \cdot & \cdot & \cdot & c_0 & & & & & \\ & c_k & c_{k-1} & \cdot & \cdot & c_1 & c_0 & & & & \\ & & c_k & c_{k-1} & \cdot & \cdot & c_1 & c_0 & & & \\ & & & & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \end{matrix} \end{matrix} \right\}.$$

Thus if  $n = 2, 3, \dots$ , if  $1 \leq j_1 < j_2 \leq n$ , and if  $p(t) = \sum_{j=0}^n a_j t^j$ , then  $R(p^{(j_1)}, p^{(j_2)})$  is a homogeneous polynomial of degree  $2n - j_1 - j_2$  in  $a_1, \dots, a_n$ . It is clear that if

$$P = P_{j_1 j_2 n}(a_1, \dots, a_n) = \frac{R(p^{(j_1)}, p^{(j_2)})}{a_n},$$

then  $P$  is a homogeneous polynomial of degree  $2n - j_1 - j_2 - 1$  in  $a_1, \dots, a_n$ . These polynomials  $P$  play a central role in our estimate.

**Theorem.** *With  $n, j_1$ , and  $j_2$  as above and  $P = P_{j_1 j_2 n}$ , there is a constant  $C$ , depending only on  $n, j_1$ , and  $j_2$ , such that if  $p(t) = \sum_{j=1}^n a_j t^j$  (with  $a_j \in \mathbb{R}$ ) then, for every interval  $I \subseteq \mathbb{R}$ ,*

$$(4) \quad \left| \int_I e^{ip(t)} dt \right| \leq \frac{C}{|P(a_1, \dots, a_n)|^{\frac{1}{n^2 - j_1 j_2 - n}}}.$$

As we will see, the polynomial  $P$  has the anisotropic homogeneity property

$$(5) \quad P(sa_1, s^2 a_2, \dots, s^n a_n) = s^{n^2 - j_1 j_2 - n} P(a_1, \dots, a_n)$$

so that (2) holds. Also,  $P$  is nonzero if and only if  $p^{(j_1)}$  and  $p^{(j_2)}$  have no common complex root. Then the (isotropic) homogeneity of  $P$  shows that (3) decays at least as fast as  $|\lambda|^{-(2n-j_1-j_2-1)/(n^2-j_1j_2-n)}$  as  $|\lambda| \rightarrow \infty$  if  $a_n \neq 0$ . The choices  $(n, j_1, j_2) = (n, j, n - 1)$  and  $(n, j_1, j_2) = (4, 1, 2)$  recover, respectively, Theorem 1 and Corollary 3 of [O2].

*Proof.* Our proof hinges on the following fact about the resultant  $R(q, r)$ : with  $q$  and  $r$  as above, if  $q_1, \dots, q_j$  are the (complex) zeros of  $q$  and  $r_1, \dots, r_k$  are the zeros of  $r$ , then

$$(6) \quad R(q, r) = b_j^k c_k^j \prod_{i=1}^j \prod_{\ell=1}^k (q_i - r_\ell).$$

(See, e.g., p. 137 in [L].) Thus, for example, it is clear that  $P$  is nonzero if and only if  $p^{(j_1)}$  and  $p^{(j_2)}$  have no common zero (assuming  $a_n \neq 0$ ). We will now observe two more consequences of (6): the first is that (5) holds and the second is

$$(7) \quad \text{if } t_0 \in \mathbb{R} \text{ and } p(t + t_0) = \sum_{j=0}^n b_j t^j, \text{ then } P(b_1, \dots, b_n) = P(a_1, \dots, a_n).$$

To prove (5), fix  $p(t) = \sum_{j=1}^n a_j t^j$  and  $s > 0$ . Let  $u(t) = p(st)$ . Then

$$(8) \quad P(sa_1, s^2 a_2, \dots, s^n a_n) = \frac{R(u^{(j_1)}, u^{(j_2)})}{s^n a_n}.$$

Since the zeros of  $u^{(j)}$  are the multiples by  $s^{-1}$  of the zeros of  $p^{(j)}$ , (6) shows that

$$R(u^{(j_1)}, u^{(j_2)}) = s^{n(n-j_1)+n(n-j_2)-(n-j_1)(n-j_2)} R(p^{(j_1)}, p^{(j_2)}).$$

With (8) this gives (5). The proof of (7) is somewhat similar: if we let  $w(t) = p(t + t_0)$ , then the zeros of  $w^{(j)}$  are the translates by  $-t_0$  of the zeros of  $p^{(j)}$ . Since  $w^{(j)}$  and  $p^{(j)}$  have the same leading coefficient, (7) follows from (6).

Our next goal is the inequality

$$(9) \quad |P(a_1, \dots, a_n)|^{\frac{1}{n^2-j_1j_2-n}} \leq C \inf_{t \in \mathbb{R}} \sum_{j=1}^n |p^{(j)}(t)|^{\frac{1}{j}}.$$

Define  $C$  to be

$$\sup \left\{ |P(\alpha_1, \dots, \alpha_n)|^{\frac{1}{n^2-j_1j_2-n}} : \sum_{j=1}^n |j! \alpha_j|^{\frac{1}{j}} = 1 \right\}.$$

Since for  $s > 0$

$$|P(sa_1, s^2a_2, \dots, s^na_n)|^{\frac{1}{n^2-j_1j_2-n}} = s |P(a_1, \dots, a_n)|^{\frac{1}{n^2-j_1j_2-n}}$$

by (5) and since

$$\sum_{j=1}^n |s^j j! a_j|^{\frac{1}{j}} = s \sum_{j=1}^n |j! a_j|^{\frac{1}{j}},$$

it follows from the definition of  $C$  that

$$|P(a_1, \dots, a_n)|^{\frac{1}{n^2-j_1j_2-n}} \leq C \sum_{j=1}^n |p^{(j)}(0)|^{\frac{1}{j}}$$

for any  $p(t) = \sum_{j=0}^n a_j t^j$ . Replacing  $p(t)$  by  $p(t + t_0)$  and applying (7) gives (9).

Now let  $N = N(n)$  be a positive integer so large that, for each  $p(t) = \sum_{j=1}^n a_j t^j$ , the cardinality of

$$\left\{ t \in \mathbb{R} : \left( p^{(k_1)}(t) \right)^{k_2} = \left( p^{(k_2)}(t) \right)^{k_1} \text{ for some } k_1, k_2 \text{ with } 1 \leq k_1 < k_2 \leq n \right\}$$

is bounded by  $N - 1$ . Then, given  $p$ ,  $\mathbb{R}$  can be written as a disjoint union of at most  $N$  intervals  $I_\ell$  such that for each  $I_\ell$  there is  $j' = j'(\ell) \in \{1, \dots, n\}$  with

$$\left| p^{(j')}(t) \right|^{\frac{1}{j'}} = \sup_{1 \leq j \leq n} \left| p^{(j)}(t) \right|^{\frac{1}{j}}, \quad t \in I_\ell.$$

Thus (9) and van der Corput's lemma give, for each  $\ell$ ,

$$\left| \int_{I \cap I_\ell} e^{ip(t)} dt \right| \leq \frac{C}{|P(a_1, \dots, a_n)|^{\frac{1}{n^2-j_1j_2-n}}}.$$

Since

$$\left| \int_I e^{ip(t)} dt \right| \leq \sum_{\ell=1}^{N(n)} \left| \int_{I \cap I_\ell} e^{ip(t)} dt \right|,$$

(4) follows.

## References

- [L] S. Lang, *Algebra*, Addison-Wesley, Reading, Mass., 1965.
- [O1] D. Oberlin, *A convolution estimate for a measure on a curve in  $\mathbb{R}^4$* , Proc. Amer. Math. Soc. **125** (1997), 1355–1361.
- [O2] ———, *Estimates for oscillatory integrals with polynomial phase*, Trans. Amer. Math. Soc. (to appear).
- [S] E. M. Stein, *Harmonic analysis: Real-variable methods, orthogonality, and oscillatory integrals*, Princeton Mathematical Series, 43. Monographs in Harmonic Analysis, III., Princeton University Press, Princeton, NJ, 1993.

DEPT. OF MATHEMATICS, FLORIDA STATE UNIVERSITY, TALLAHASSEE, FL 32306-3027  
*E-mail address:* oberlin@zeno.math.fsu.edu