

**SIMPLY NONDEGENERATE MULTILINEAR OSCILLATORY
INTEGRAL OPERATORS WITH SMOOTH PHASE**

MICHAEL GREENBLATT

1) Introduction As in [CLTaTh], in this paper we consider operators of the form

$$T_\lambda(f_1, \dots, f_n) = \int_{R^m} e^{i\lambda S(x)} \prod_{j=1}^n f_j(\pi_j(x)) \eta(x) dx \quad (1.1)$$

Here $S(x)$ is a smooth real-valued phase function, λ is a parameter, and $\eta(x)$ is a C^1 cutoff function supported in the unit ball centered at the origin. π_j denotes projection from R^m to some subspace V_j of dimension strictly less than m , and each f_j is an L^∞ function. A natural question to ask is for which $\{V_j\}_{j=1}^n$ and which smooth phase functions $S(x)$ do we get an estimate of the form

$$|T_\lambda(f_1, \dots, f_n)| \leq C(1 + |\lambda|)^{-\epsilon} \prod_{j=1}^n \|f_j\|_\infty \quad (1.2)$$

Here $\epsilon > 0$ is to be independent of $\eta(x)$. In [CLTaTh] this question was considered in considerable depth, and attention was focused on the case where $S(x)$ is a polynomial. Here we will extend several of their results to the general smooth case. When the π_j are all projections onto 1 dimensional spaces and one considers L^2 norms instead of L^∞ norms, much additional work has been done, such as [PS] [PSSt] [Se].

Note that if $S(x) = \sum_j p_j \circ \pi_j(x)$, then there can be no positive ϵ satisfying (1.2). For one can let $f_j = e^{-i\lambda p_j}$ and then (1.1) is equal to $\int \eta(x) dx$, a quantity independent of λ . In [CLTaTh], such a polynomial $S(x)$ is referred to as *degenerate relative to* $\{V_j\}_{j=1}^n$, and a polynomial $S(x)$ that has no realization of the form $S(x) = \sum_j p_j \circ \pi_j(x)$, p_j polynomials, is referred to as *nondegenerate relative to* $\{V_j\}_{j=1}^n$.

One way one can ensure that a polynomial is nondegenerate relative to $\{V_j\}_{j=1}^n$ is for there to exist a differential operator L of the form $L = \prod_{j=1}^n (w_j \circ \nabla)$, with each w_j a unit vector in V_j^\perp , such that $L(S)$ is not the zero function. For if one applies L to a polynomial of the form $S(x) = \sum_j p_j \circ \pi_j(x)$, each factor knocks out one of the terms and one is left with zero. [CLTaTh] refers to a polynomial S for which there exists such an L as a *simply nondegenerate* polynomial. Hence simply nondegenerate

Received by the editors May 23, 2007.

2000 *Mathematics Subject Classification.* 32C17, 32F29, 32Q20.

polynomials are all nondegenerate. The converse does not hold in general, as explained in [CLTaTh]. However for several important classes of $\{V_j\}_{j=1}^n$, simple nondegeneracy is equivalent to nondegeneracy. In [CLTaTh], the existence of an ϵ satisfying (1.2) is shown for simply nondegenerate polynomials $S(x)$ (among others), and therefore for all nondegenerate polynomials when simply nondegenerate and nondegenerate are equivalent. Their estimates are uniform in the sense that for a given family $\{V_j\}_{j=1}^n$, ϵ depends only on the degree of $S(x)$ and $\max_{|x|\leq 1} |L(S(x))|$, while C depends on these quantities as well as the C^1 norm of $\eta(x)$.

The purpose of this paper is to extend the simple nondegeneracy results of [CLTaTh] to the case of general C^∞ phase. Since the ϵ in the proofs in [CLTaTh] depend on the degree of $S(x)$, the results there do not carry over immediately. Some earlier results in this direction are contained in [CaW]. We also refer to Chapter 8 of [S] for relevant introductory material concerning Van der Corput-type lemmas.

Definition: Suppose $S(x)$ is a smooth function defined in a neighborhood of a point a . $S(x)$ is called degenerate at a if there are smooth functions $s_j : V_j \rightarrow \mathbb{R}$ for which $S(x) - \sum_{j=1}^n s_j(\pi_j(x))$ has a zero of infinite order at a . We say $S(x)$ is nondegenerate at a if there do not exist such functions.

Definition: Suppose $S(x)$ is a smooth function defined in a neighborhood of a point a . $S(x)$ is called simply nondegenerate at a if there exists a differential operator L of the form $L = \prod_{j=1}^n (w_j \circ \nabla)$, with each w_j a unit vector in V_j^\perp , such that $L(S(x))$ does not vanish to infinite order at a .

As in the polynomial case, in the smooth case it is immediate that if $S(x)$ is simply nondegenerate at a then it is nondegenerate at a .

Our main theorem gives uniform decay for simply nondegenerate $S(x)$ in the smooth case:

Theorem 1.1: Suppose $S(x)$ is simply nondegenerate at 0. Let L be as above and α be a multiindex such that $\partial^\alpha(L \circ S)(0) = c \neq 0$. Then there is an open U containing the origin, U depending on $c, m, n, |\alpha|$, and the $C^{|\alpha|+n+1}$ norm of S , such that if $\eta(x)$ is supported in U , for $\epsilon = \frac{2^{-n+1}}{|\alpha|+n}$ and any $\delta > 0$ we have

$$|T_\lambda(f_1, \dots, f_n)| \leq C(1 + |\lambda|)^{-\epsilon+\delta} \prod_{j=1}^n \|f_j\|_\infty \quad (1.3)$$

Here C depends on $c, m, n, |\alpha|, \|\eta\|_{C^1}$, and the C^l norm of $S(x)$ for some l depending on δ . For fixed values of the other parameters, C is $O(\|\eta\|_{C^1})$. In the case where S is a polynomial, we can take $\delta = 0$.

We will see in Lemma 1.4 below that if $\{V_j\}_{j=1}^n$ is such that simple nondegeneracy is equivalent to nondegeneracy for polynomial $S(x)$, then simple nondegeneracy is also equivalent to nondegeneracy for smooth $S(x)$ at any point. As a result, by Proposition 3.1 and Lemma 3.5 of [CLTaTh], Theorem 1.1 immediately implies the following smooth analogues of Theorem 2.2 and Theorem 2.4 of [CLTaTh]:

Theorem 1.2: Suppose each V_j has codimension 1. Then (1.2) holds for some $\epsilon > 0$ for all η with sufficiently small support if and only if $S(x)$ is nondegenerate at the origin.

Theorem 1.3: Suppose $\{V_j : 1 \leq j \leq n\}$ are each of dimension $k < n$. Suppose the orthocomplements V_j^\perp together span a space of dimension at most $(m - k)n$, and $(m - k)n \leq m$. Then (1.2) holds for some $\epsilon > 0$ for all η with sufficiently small support if and only if $S(x)$ is nondegenerate at the origin.

In view of Theorem 1.1, Theorems 1.2 and 1.3 are uniform in the sense that ϵ depends only on n and the order of vanishing of $L \circ S$ at the origin.

Lemma 1.4: Suppose $\{V_j\}_{j=1}^n$ are such that simple nondegeneracy is equivalent to nondegeneracy for any polynomial $S(x)$. Then for smooth $S(x)$, simple nondegeneracy is equivalent to nondegeneracy at any point.

Proof: Since simple nondegeneracy readily implies nondegeneracy, we assume that $S(x)$ is not simply nondegenerate at a point a , and we will show that $S(x)$ is degenerate at a . Clearly it suffices to assume that $a = 0$. Let $\sum_\alpha S_\alpha x^\alpha$ denote the (possibly divergent) Taylor expansion of $S(x)$ about the origin. For each positive integer i , we define

$$R_i(x) = \sum_{|\alpha|=i} S_\alpha x^\alpha \quad (1.4)$$

Suppose $w_j \in V_j^\perp$ for $1 \leq j \leq n$. Analogous to (1.4) we can write

$$\prod_{j=1}^n (w_j \circ \nabla) S = \sum_i \prod_{j=1}^n (w_j \circ \nabla) R_i(x) \quad (1.5)$$

Note that $\prod_{j=1}^n (w_j \circ \nabla) R_i(x)$ is the sum of the terms of (1.5) of degree $i - n$. Since $S(x)$ is not simply nondegenerate at the origin, for any such w_j the right hand side of (1.5) must have no nonvanishing terms. In other words, for all i and all choices of w_j we have

$$\prod_{j=1}^n (w_j \circ \nabla) R_i(x) = 0 \quad (1.6)$$

Since we are assuming simple nondegeneracy is equivalent to nondegeneracy for polynomials, this means we can write

$$R_i(x) = \sum_{j=1}^n p_{ij} \circ \pi_j(x) \quad (1.7)$$

Here p_{ij} are polynomials. Since the terms of $R_i(x)$ are all of degree i , the terms of the righthand sum of all other degrees must add to zero. Hence we may replace each p_{ij} by the sum of its monomials of degree i if necessary, and assume each p_{ij} is homogeneous of degree i .

Next, for a given j we let r_j be a rotation such that the image of $r_j \circ \pi_j$ is $\{(y, z) \in R^{k_j} \times R^{m-k_j} : z = 0\}$. Since $p_{ij} \circ \pi_j = p_{ij} \circ r_j^{-1} \circ (r_j \circ \pi_j)$ and $p_{ij} \circ \pi_j$ is homogeneous of degree i , $p_{ij} \circ r_j^{-1}$ is a homogeneous polynomial in the y variables of degree i . Thus the monomials of $p_{ij} \circ r_j^{-1}(y)$ for different i 's do not overlap. As a result, it makes sense to write $\sum_i p_{ij} \circ r_j^{-1}(y)$ as a (possibly nonconvergent) power series in the y variables.

By a famous theorem of Borel (see p.16 of [H]), we can let $q_j(y)$ be a C^∞ function whose Taylor series about the origin is given by $\sum_i p_{ij} \circ r_j^{-1}(y)$. So for a fixed i_0 , $q_j(y) - \sum_{i < i_0} p_{ij} \circ r_j^{-1}(y)$ has a zero of order i_0 at the origin. Similarly, $q_j \circ r_j \circ \pi_j(x) - \sum_{i < i_0} p_{ij} \circ \pi_j(x)$ has a zero of order i_0 at the origin. Adding this over all j , we get that $\sum_j q_j \circ r_j \circ \pi_j(x) - \sum_{i < i_0} R_i(x)$ has a zero of order i_0 at the origin. By (1.4), this means that $\sum_j q_j \circ r_j \circ \pi_j(x) - S(x)$ has a zero of infinite order at the origin. Since each $q_j \circ r_j$ is C^∞ , this means that $S(x)$ is degenerate to infinite order at the origin, and we are done.

2) Proof of Theorem 1.1

Case 1: $n = 1$, $S(x)$ a polynomial

Let g denote the degree of $S(x)$ and let k_1 denote the dimension of V_1 . Rotate coordinates such that $V_1 = \{x = (y, z) \in R^{k_1} \times R^{m-k_1} : z = 0\}$, and such that the vector w_1 in the definition of simply nondegenerate is in the z_1 direction. Our arguments will resemble the proof of the Van der Corput lemma in the z_1 direction. Namely, for some $a > 0$ to be determined we divide the domain of (1.1) into two parts. The first part is where $|\partial_{z_1} S(x)| < |\lambda|^{-a}$, and the second where $|\partial_{z_1} S(x)| \geq |\lambda|^{-a}$. On the first part, we take absolute values and integrate, obtaining a term of absolute value at most

$$m\{x \in \text{supp}(\eta) : |\partial_{z_1} S(x)| < |\lambda|^{-a}\} \|f_1\|_\infty \|\eta\|_\infty$$

Here m denotes Lebesgue measure. This in turn is bounded by

$$m\{x \in \text{supp}(\eta) : |\partial_{z_1} S(x)| < |\lambda|^{-a}\} \|f_1\|_\infty \|\eta\|_{C^1} \quad (2.1)$$

On the second part we do an integration by parts in z_1 on the oscillatory factor. We get

$$\frac{1}{i\lambda} \int_{|S_{z_1}(y,z)| \geq |\lambda|^{-a}} e^{i\lambda S(y,z)} \left[\frac{S_{z_1 z_1}(y,z)}{S_{z_1}(y,z)^2} \eta(y,z) - \eta_{z_1}(y,z) \right] f_1(y) dy dz \quad (2.2)$$

The expression (2.2) is of absolute value at most

$$\frac{1}{|\lambda|} \|f_1\|_\infty \|\eta\|_\infty \left(\int_{|(y,z)| < 1, |S_{z_1}(y,z)| \geq |\lambda|^{-a}} \left| \frac{S_{z_1 z_1}(y,z)}{S_{z_1}(y,z)^2} \right| dy dz \right) + \frac{1}{|\lambda|} \|f_1\|_\infty \|\eta\|_{C^1} \quad (2.3)$$

For fixed y and (z_2, \dots, z_n) , the domain of integration of (2.3) can be written as the union of at most $g - 1$ intervals on which S_{z_1} is monotonic as a function of z_1 . Consequently, on each of these intervals, one may perform the integration of $\left| \frac{S_{z_1 z_1}(y,z)}{S_{z_1}(y,z)^2} \right|$ as in the proof of the Van der Corput lemma, obtaining the difference of $\frac{1}{S_{z_1}(y,z)}$ at the endpoints, so that (2.3) is at most

$$\begin{aligned} & (2g - 2) |\lambda|^{a-1} \|f_1\|_\infty \|\eta\|_\infty + \frac{1}{|\lambda|} \|f_1\|_\infty \|\eta\|_{C^1} \\ & \leq (2g - 1) |\lambda|^{a-1} \|f_1\|_\infty \|\eta\|_{C^1} \end{aligned} \quad (2.4)$$

Combining (2.1) and (2.4), in the current situation the oscillatory integral (1.1) is bounded in absolute value by:

$$[m\{x \in \text{supp}(\eta) : |\partial_{z_1} S(x)| < |\lambda|^{-a}\} + (2g - 1)|\lambda|^{a-1}] \|f_1\|_\infty \|\eta\|_{C^1} \quad (2.5)$$

By the simple nondegeneracy of S , there is some multiindex α for which $\partial^\alpha \partial_{z_1} S(0) = c \neq 0$. As a result, by [C1] for example, if the support of η is sufficiently small, one has

$$m\{x \in \text{supp}(\eta) : |\partial_{z_1} S(x)| < |\lambda|^{-a}\} < C|\lambda|^{-\frac{a}{|\alpha|}} \quad (2.6)$$

Here C depends on c , $|\alpha|$, m , and the $C^{|\alpha|+2}$ norm of S . As a result, we can select $a = \frac{|\alpha|}{|\alpha|+1}$, and (2.5) is at most

$$(C + 2g - 1)|\lambda|^{-\frac{1}{1+|\alpha|}} \|f_1\|_\infty \|\eta\|_{C^1} \quad (2.7)$$

This gives us the estimate required by Theorem 1.1 and we are done when $n = 1$ and $S(x)$ is a polynomial.

Case 2: $n > 1$, $S(x)$ a polynomial.

To prove Theorem 1.1 in this case we will need the following procedure. Given a smooth function $Q(x)$ defined in a neighborhood of the origin, we successively define functions $Q^1(y, z)$, $Q^2(y, z, \zeta^2)$, ..., $Q^n(y, z, \zeta^2, \dots, \zeta^n)$. Let $Q^1(y, z)$ be $Q(x)$ in rotated coordinates such that V_1 is $\{(y, z) \in R^{k_1} \times R^{m-k_1} : z = 0\}$ and such that w_1 is the z_1 direction. Once Q^{j-1} is defined, one defines Q^j as follows. Consider coordinates such that $V_j = \{(y, z) \in R^{k_j} \times R^{m-k_j} : z = 0\}$ with w_j being the z_1 direction. Letting $Q_\zeta(y, z) = Q(y, z + \zeta) - Q(y, z)$ in these coordinates, we define $Q^j(y, z, \zeta^2, \dots, \zeta^j)$ to be $(Q_{\zeta^j})^{j-1}(y, z, \zeta^2, \dots, \zeta^{j-1})$. I claim that inductively we have

$$\frac{\partial^{j-1} Q^j}{\partial_{\zeta_1^j} \dots \partial_{\zeta_1^2}}(y, z, 0) = \prod_{i=2}^j (w_i \circ \nabla) Q(y, z) \quad (2.8)$$

For if we know the $j - 1$ case (the case $j = 1$ is a tautology) we have the following:

$$\begin{aligned} \frac{\partial^{j-2} Q^j}{\partial_{\zeta_1^{j-1}} \dots \partial_{\zeta_1^2}}(y, z, 0, \dots, 0, \zeta^j) &= \frac{\partial^{j-2} (Q_{\zeta^j})^{j-1}}{\partial_{\zeta_1^{j-1}} \dots \partial_{\zeta_1^2}}(y, z, 0) = \prod_{i=2}^{j-1} (w_i \circ \nabla) Q_{\zeta^j}(y, z) \\ &= \prod_{i=2}^{j-1} ((w_i \circ \nabla) Q)(y, z + \zeta^j) - \prod_{i=2}^{j-1} ((w_i \circ \nabla) Q)(y, z) \end{aligned}$$

Then taking directional derivatives in the ζ_1^j direction gives (2.8). We will be most interested in (2.8) when $Q = (w_1 \circ \nabla)S$. In fact, Theorem 1.1 in the polynomial case will follow readily from the following lemma:

Lemma 2.1: Suppose T_λ is of the form (1.1) with $S(x)$ a polynomial. Let δ be such that $|x| < \delta$ for all $x \in \text{supp}(\eta)$. Let $s(x)$ denote $(w_1 \circ \nabla)S(x)$, and let $m\{s^n(y, z, \zeta^2, \dots, \zeta^n) <$

$b\}$ denote $m\{(y, z, \zeta^2, \dots, \zeta^n) \in [-2\delta, 2\delta]^{m+\sum_{j>1}(m-k_j)} : s^n(y, z, \zeta^2, \dots, \zeta^n) < b\}$. For any $a > 0$ we have $|T_\lambda(f_1, \dots, f_n)| <$

$$[m\{(s^n(y, z, \zeta^2, \dots, \zeta^n) < |\lambda|^{-a}\} + (2g-1)|\lambda|^{a-1}]^{2^{1-n}} \|\eta\|_{C^1} \prod_{j=1}^n \|f_j\|_\infty \quad (2.9)$$

Proof: The case $n = 1$ follows from (2.5), so we can assume the lemma for $n - 1$ and prove it for n . We work in the general setup used in the inductive step in Theorem 2.3 of [CLTaTh], originating in [CaCW]. Like above and in [CLTaTh], we rotate to coordinates such that $V_n = \{(y, z) \in R^{k_n} \times R^{m-k_n} : z = 0\}$. As above, we assume the direction w_n in the definition of simply nondegenerate is the z_1 direction. We now have

$$T_\lambda(f_1, \dots, f_n) = \int f_n(y) \left(\int e^{i\lambda S(y,z)} \prod_{j=1}^{n-1} f_j(\pi_j(y, z)) \eta(y, z) dz \right) dy \quad (2.10)$$

This equals $\langle U_\lambda(f_1, \dots, f_{n-1}), \bar{f}_n \rangle$ for an appropriate operator U_λ . Hence we have

$$|T_\lambda(f_1, \dots, f_n)| \leq \|f_n\|_2 \|U_\lambda(f_1, \dots, f_{n-1})\|_2 \leq \|f_n\|_\infty \|U_\lambda(f_1, \dots, f_{n-1})\|_2 \quad (2.11)$$

As in [CLTaTh], we write $\|U_\lambda(f_1, \dots, f_{n-1})\|_2^2$ as

$$\int_{R^{m-k_n}} \left[\int_{R^m} e^{i\lambda(S(y,z) - S(y,z+\zeta))} \prod_{j<n} f_j(\pi_j(y, z)) \bar{f}_j(\pi_j(y, z+\zeta)) \eta(y, z) \bar{\eta}(y, z+\zeta) dz dy \right] d\zeta \quad (2.12)$$

For fixed ζ , the integral inside the brackets is of the form (1.1), with phase $S(y, z) - S(y, z + \zeta)$ and acting on the $n - 1$ functions $f_j(\pi_j(y, z)) \bar{f}_j(\pi_j(y, z + \zeta))$ for $j < n$. Thus one can apply the inductive hypothesis to the bracketed integral and say it is bounded in absolute value by

$$[m\{(s_\zeta)^{n-1}(y, z, \zeta^2, \dots, \zeta^{n-1}) < |\lambda|^{-a}\} + (2g-1)|\lambda|^{a-1}]^{2^{2-n}} \|\eta\|_{C^1}^2 \prod_{j<n} \|f_j\|_\infty^2 \quad (2.13)$$

Here $s_\zeta(y, z) = s(y, z + \zeta) - s(y, z) = (w_1 \circ \nabla)(S(y, z + \zeta) - S(y, z))$. We integrate (2.13) in ζ , using Jensen's inequality to push the exponent 2^{2-n} to the outside. Given that $(s_\zeta)^{n-1}(y, z, \zeta^2, \dots, \zeta^{j-1}) = s^n(y, z, \zeta^2, \dots, \zeta^{j-1}, \zeta)$, we get that (2.12) is bounded by

$$[m\{s^n(y, z, \zeta^2, \dots, \zeta^n) < |\lambda|^{-a}\} + (2g-1)|\lambda|^{a-1}]^{2^{2-n}} \|\eta\|_{C^1}^2 \prod_{j<n} \|f_j\|_\infty^2 \quad (2.14)$$

Putting this upper bound for $\|U_\lambda(f_1, \dots, f_{n-1})\|_2^2$ into (2.11) gives the lemma and we are done.

We now may prove Theorem 1.1 in the polynomial case. Applying (2.8) to $s(x)$ for $j = n$, the simple nondenergency hypothesis implies that for some α we have

$$\partial^\alpha \left(\frac{\partial^{n-1} s^n}{\partial \zeta_1^n \dots \partial \zeta_1^n} \right) (0, 0, 0) = c \neq 0$$

Thus s^n has a zero of order $|\alpha| + n - 1$ at the origin. Using [C] again for example, assuming the support of η is sufficiently small, depending on c , m , n , $|\alpha|$ and the $C^{|\alpha|+n+1}$ norm of S , we have that

$$m\{s^n(y, z, \zeta^1, \dots, \zeta^n) < |\lambda|^{-a}\} < C|\lambda|^{-\frac{a}{|\alpha|+n-1}} \quad (2.15)$$

Here C is a function of m , $|\alpha|$, n , and c . We choose $a = \frac{|\alpha|+n-1}{|\alpha|+n}$ in (2.15). Then (2.9) gives

$$|T_\lambda(f_1, \dots, f_n)| < C'|\lambda|^{-\frac{2^{1-n}}{|\alpha|+n}} \|\eta\|_{C^1} \prod_{j=1}^n \|f_j\|_\infty \quad (2.16)$$

Here C' is a function m , $|\alpha|$, n , c , and g . This completes the proof of Theorem 1.1 in the polynomial case.

Case 3: General smooth $S(x)$

Because the ϵ obtained in the polynomial case of Theorem 1.1 did not depend on the degree of $S(x)$, we will be able to extend to general smooth $S(x)$ by dividing the domain into cubes of radius $|\lambda|^{-e}$ for small e , and then approximating $S(x)$ by polynomials of sufficiently high degree on each of these cubes. To be precise, let l be some positive integer, and write the cutoff function $\eta(x)$ as $\sum_{i=1}^I \eta_i(x)$, where each $\eta_i(x)$ is supported on a set of diameter $< |\lambda|^{-\frac{1}{l}}$ and where $I < C_0|\lambda|^{\frac{m}{l}}$. We may also assume $\|\eta_i\|_{C^1} < C_0|\lambda|^{\frac{1}{l}} \|\eta\|_{C^1}$. Here C_0 is a uniform constant. We correspondingly write $T_\lambda = \sum_{i=1}^I T_\lambda^i$, where

$$T_\lambda^i(f_1, \dots, f_n) = \int_{R^m} e^{i\lambda S(x)} \prod_{j=1}^n f_j(\pi_j(x)) \eta_i(x) dx \quad (2.17)$$

We break each term (2.17) into two parts. For some fixed i , let x_0 be a point in the support of η_i . Let $\bar{S}(x)$ be the sum of the first $2l$ terms of the Taylor expansion of $S(x)$ taken about $x = x_0$. We write $T_\lambda^i = T_\lambda^{i,1} + T_\lambda^{i,2}$, where

$$T_\lambda^{i,1}(f_1, \dots, f_n) = \int_{R^m} e^{i\lambda \bar{S}(x)} \prod_{j=1}^n f_j(\pi_j(x)) \eta_i(x) dx \quad (2.18a)$$

$$T_\lambda^{i,2}(f_1, \dots, f_n) = \int_{R^m} [e^{i\lambda S(x)} - e^{i\lambda \bar{S}(x)}] \prod_{j=1}^n f_j(\pi_j(x)) \eta_i(x) dx \quad (2.18b)$$

We can apply the polynomial case treated above to (2.18a), and get that

$$|T_\lambda^{i,1}(f_1, \dots, f_n)| \leq C|\lambda|^{\frac{1}{l}} |\lambda|^{-\epsilon} \|\eta\|_{C^1} \prod_{j=1}^n \|f_j\|_\infty \quad (2.19)$$

Here $\epsilon = \frac{2^{1-n}}{|\alpha|+n}$, and C is a constant depending on c , l , m , n , and $|\alpha|$. For (2.18b), we use Taylor's theorem on the difference of exponentials, obtaining

$$|e^{i\lambda S(x)} - e^{i\lambda \bar{S}(x)}| \leq C'(|\lambda|^{-\frac{1}{l}})^{2l} = C'|\lambda|^{-2}$$

Here C' depends on the C^{2l+1} norm of S . Hence we have

$$|T_\lambda^{i,2}(f_1, \dots, f_n)| \leq C' |\lambda|^{-2+\frac{1}{i}} \|\eta\|_{C^1} \prod_{j=1}^n \|f_j\|_\infty \quad (2.20)$$

Adding up (2.19) and (2.20) over all i , we obtain

$$|T_\lambda(f_1, \dots, f_n)| \leq C'' (|\lambda|^{\frac{m+1}{i}} |\lambda|^{-\epsilon} + |\lambda|^{-2+\frac{m+1}{i}}) \|\eta\|_{C^1} \prod_{j=1}^n \|f_j\|_\infty \quad (2.21)$$

Here C'' depends on $c, l, m, n, |\alpha|$, and the C^{2l+1} norm of S . Given $\delta > 0$, by picking l large enough, one can make (2.21) bounded by $2C'' |\lambda|^{-\epsilon+\delta} \|\eta\|_{C^1} \prod_{j=1}^n \|f_j\|_\infty$. This completes the proof of Theorem 1.1.

References

- [CaCW] A. Carbery, M. Christ, and J. Wright, *Multidimensional van der Corput and sublevel set estimates*, J. Amer. Math. Soc. **12** (1999), 981–1015.
- [CaW] A. Carbery and J. Wright, *What is Van der Corput's lemma in higher dimensions?*, Proceedings of the 6th International Conference on Harmonic Analysis and PDEs, Publ. Mat. (2002), 13-26.
- [C] M. Christ, *Hilbert transforms along curves. I. Nilpotent groups*, Annals of Mathematics (2) **122** (1985), 575-596.
- [CLTaTh] M. Christ, X. Li, T. Tao, and C. Thiele, *On multilinear oscillatory integrals, nonsingular and singular*, Duke Math. J. **130** (2005), 321–351.
- [H] L. Hörmander, *The analysis of linear partial differential operators. I. Distribution theory and Fourier analysis*, 2nd ed. Springer-Verlag, Berlin, (1990), xii+440 pp.
- [PS] D. H. Phong and E. M. Stein, *The Newton polyhedron and oscillatory integral operators*, Acta Mathematica **179** (1997), 107-152.
- [PSSt] D. H. Phong, E. M. Stein, and J. Sturm, *Multilinear level set operators, oscillatory integral operators, and Newton diagrams*, Math. Annalen **319** (2001), 573-596.
- [Se] A. Seeger, *Radon transforms and finite type conditions*, Journal of the American Mathematical Society **11** (1998), 869-897.
- [S] E. M. Stein, *Harmonic analysis; real-variable methods, orthogonality, and oscillatory integrals*, Princeton Mathematics Series Vol. 43, Princeton University Press, Princeton, NJ, 1993.

DEPARTMENT OF MATHEMATICS, 244 MATHEMATICS BUILDING, UNIVERSITY AT BUFFALO, BUFFALO, NY 14260

E-mail address: mg62@buffalo.edu