

Photoacoustic and thermoacoustic tomography with an uncertain wave speed

LAURI OKSANEN AND GUNTHER UHLMANN

We consider the mathematical model of photoacoustic and thermoacoustic tomography in media with a variable sound speed. When the sound speed is known, the explicit reconstruction formula of [14] can be used. We study how a modeling error in the sound speed affects the reconstruction formula and quantify the effect in terms of a stability estimate.

1. Introduction

Coupled-physics imaging methods, also called hybrid methods, attempt to combine the high resolution of one imaging method with the high contrast capabilities of another through a physical principle. One important medical imaging application is breast cancer detection. Ultrasound provides a high, sub-millimeter resolution, but suffers from low contrast. On the other hand, many tumors absorb much more energy of electromagnetic waves, in some specific energy bands, than healthy cells.

Photoacoustic tomography (PAT) [19] consists of sending relatively harmless optical radiation into tissues that causes heating, with increases of the temperature in the millikelvin range, which results in the generation of propagating ultrasound waves. This is the photo-acoustic effect. The inverse problem then consists of reconstructing the optical properties of the tissue. In Thermoacoustic tomography (TAT), see e.g., [8], low-frequency microwaves, with wavelengths of the order of $1m$, are sent into the medium. The rationale for using the latter frequencies is that they are less absorbed than optical frequencies. PAT, TAT and other couple-physics imaging techniques offer potential breakthroughs in the clinical application of multi-wave methods to early detection of cancer, functional imaging and molecular imaging among others [19, 20].

1991 *Mathematics Subject Classification*. Primary: 35R30.

Key words and phrases. Inverse problems, photoacoustic tomography, thermoacoustic tomography, wave equation, stability.

There are two steps in PAT and TAT. The first one is to solve the inverse source problem for the wave equation. The source measures the absorbed radiation and varies from tissue to tissue. Once this is solved the second step is to determine the optical or electromagnetic parameters from the internal information obtained in the first step. Here we are concerned with the first step that we now describe.

Let u solve the initial value problem

$$(1) \quad \begin{cases} (\partial_t^2 - c^2 \Delta)u &= 0 & \text{in } (0, T) \times \mathbb{R}^n, \\ u|_{t=0} &= f, \\ \partial_t u|_{t=0} &= 0, \end{cases}$$

where $T > 0$ is fixed and c is a smooth and strictly positive function on \mathbb{R}^n . Let us assume that the source f is supported in M , where $M \subset \mathbb{R}^n$ is some bounded domain with smooth boundary. The measurements are modeled by the operator

$$(2) \quad \Lambda_c f := u|_{[0, T] \times \partial M}.$$

The inverse problem is to reconstruct the unknown f and c from Λ_c .

Most of the available results assume that the sound speed c is known. Concerning the constant speed case we refer to the survey paper [9] and references therein. In practice, there are many cases when the constant sound speed model is inaccurate. For instance in breast imaging, the different components of the breast, such as the glandular tissues, stromal tissues, cancerous tissues and other fatty issues, have different acoustic properties. The variations between their acoustic speeds can be as great as 10% [7]. The case of a variable sound speed was thoroughly investigated in [14] and a reconstruction method for the source was proposed. A numerical algorithm based on this method was developed in [13] and compared with the standard time-reversal method used for instance in [5, 6]. The case of a discontinuous sound speed was considered in [15] which arises, for instance, in brain imaging.

The paper [17] deals with the case where the source f is known and recovers the sound speed c with one measurement under the geometric assumption that the domain is foliated by strictly convex hypersurfaces with respect to the Riemannian metric $g = \frac{1}{c^2} dx^2$ where dx^2 denotes the Euclidean metric. In practice, the wave speed is known only up to some uncertainty and we would like to reconstruct both c and f given $\Lambda_c f$. This problem is linear in f and non-linear in c . It is shown in [16] that the linearized problem of recovering both parameters is unstable.

In this paper, we assume that our best guess for the wave speed c_0 is close to the true wave speed c . We will study how the modeling error $c - c_0$ affects the reconstruction of f by the modified time-reversal method introduced in [14].

2. Statement of the results

Let us begin by describing the reconstruction method introduced in [14]. To define the operator giving the reconstruction, we consider the time reversed wave equation on M ,

$$(3) \quad \begin{aligned} \partial_t^2 v - c^2 \Delta v &= 0, & \text{in } (0, T) \times M, \\ v|_{x \in \partial M} &= h, & \text{in } (0, T) \times \partial M, \\ v|_{t=T} &= \Delta^{-1} h|_{t=T}, \quad \partial_t v|_{t=T} = 0, & \text{in } M, \end{aligned}$$

where $\Delta^{-1} h|_{t=T}$ is the solution of the Dirichlet problem on M ,

$$\Delta \phi = 0 \text{ in } M \quad \text{and} \quad \phi = h|_{t=T} \text{ on } \partial M.$$

Moreover, we define the operators

$$A_c h = v|_{t=0} \quad \text{and} \quad K_c = 1 - A_c \Lambda_c,$$

where v is the solution of (3). The method of [14] gives a reconstruction by the formula

$$f = R_c \Lambda_c f, \quad \text{where} \quad R_c := \sum_{m=0}^{\infty} K_c^m A_c.$$

Let $c_0 \in C^\infty(\mathbb{R}^n)$ be strictly positive and let us consider the reconstruction operator R_{c_0} corresponding to c_0 . If c_0 is close to c , we expect the reconstruction

$$(4) \quad \tilde{f} = R_{c_0} \Lambda_c f$$

to be close to f . Indeed, in this paper we show that the difference $\tilde{f} - f$ is bounded by suitable norms of the measurement $\Lambda_c f$ and the modeling error $c - c_0$.

As mentioned above, the closely related problem to find c given the pair $(\Lambda_c f, f)$ was considered in [17], and the problem was shown to be stable

under the geometric assumption that M is foliated by strictly convex hypersurfaces with respect to the Riemannian metric $c^{-2}dx^2$. We will impose a similar convexity condition below. Furthermore, we will assume that ∂M is strictly convex with respect to the Riemannian metric $c^{-2}dx^2$. If this is the case, then the operator Λ_c has the following regularity property:

$$\Lambda_c : H_0^1(\mathcal{K}) \rightarrow H^1((0, T) \times \partial M),$$

where $\mathcal{K} \subset M^{\text{int}}$ is compact; see e.g., [14]. Without the convexity assumption, a loss of smoothness up to degree $1/4$ is possible [18]. We describe this phenomenon in more detail in the appendix below.

The operator K_c is compact if the state of the wave equation (1) is smooth at $t = T$ irrespective of f . This is guaranteed if all unit speed geodesics starting from M at $t = 0$ exit M before $t = T$ and do not re-enter before $t = T$. To avoid technicalities related to re-entering singularities we assume below that M is convex also in the sense that no geodesic exiting M re-enters M .

Theorem 1. *Let $c_0 \in C^\infty(\mathbb{R}^n)$ be strictly positive and suppose that the Riemannian manifold $(M, c_0^{-2}dx^2)$ has a strictly convex function with no critical points and that M is strictly convex with respect to the Riemannian metric $c_0^{-2}dx^2$. Let $\mathcal{K} \subset M^{\text{int}}$ be compact, let $C_0 > 0$ and consider the functions $f \in H^3(\mathbb{R}^n)$ and $c \in C^\infty(\mathbb{R}^n)$ satisfying*

$$\|c\|_{C^2(\mathcal{K})} \leq C_0, \quad \text{supp}(f) \subset \mathcal{K}, \quad c = c_0 \text{ in } \mathbb{R}^n \setminus \mathcal{K}.$$

There are $\epsilon_0, T, C > 0$ such that if c satisfies also

$$\|c - c_0\|_{C^1(\mathcal{K})} \leq \epsilon_0$$

then

$$(5) \quad \begin{aligned} & \| (R_{c_0} - R_c) \Lambda_c f \|_{H^1(M)} \\ & \leq C \|c - c_0\|_{L^\infty(M)} \| \Lambda_c f \|_{H^1((0, T) \times \partial M)}^{1/2} \| f \|_{H^3(\mathcal{K})}^{1/2}. \end{aligned}$$

We emphasize that $(R_{c_0} - R_c) \Lambda_c f = \tilde{f} - f$, whence (5) gives a bound for the reconstruction error under an a priori assumption on the size of the norm of f in $H^3(\mathcal{K})$.

3. The proof

The existence of a strictly convex function on $(M, c_0^{-2}dx^2)$ implies that the wave equation

$$\partial_t^2 v - c_0^2 \Delta v = 0, \quad \text{in } (0, T) \times M,$$

is exactly controllable for large $T > 0$; see e.g., [21]. This again implies that $(M, c_0^{-2}dx^2)$ is non-trapping; see e.g., [1]. Thus $\sum_{m=0}^\infty K_{c_0}^m$ is a bounded operator on $H_0^1(M)$ by [14]. Let us begin by showing that the reconstruction (4) is well defined. This is guaranteed by the next lemma.

Lemma 1. *Let $f \in H_0^1(M) \cap H^2(M)$. Then $A_{c_0} \Lambda_c f \in H_0^1(M)$.*

Proof. Notice that the solution u of (1) is in $C([0, T]; H^2(M))$. Let us consider the solution v of

$$\begin{aligned} \partial_t^2 v - c_0^2 \Delta v &= 0, & \text{in } (0, T) \times M, \\ v|_{x \in \partial M} &= \Lambda_c f, & \text{in } (0, T) \times \partial M, \\ v|_{t=T} &= \phi, \quad \partial_t v|_{t=T} = 0, & \text{in } M, \end{aligned}$$

where $\phi = \Delta^{-1} \Lambda_c f|_{t=T}$. The difference $w = u - v$ satisfies

$$\begin{aligned} \partial_t^2 w - c_0^2 \Delta w &= (c^2 - c_0^2) \Delta u, & \text{in } (0, T) \times M, \\ w|_{x \in M} &= 0, & \text{in } (0, T) \times \partial M, \\ w|_{t=T} &= u|_{t=T} - \phi, \quad \partial_t w|_{t=T} = \partial_t u|_{t=T}, & \text{in } M. \end{aligned}$$

Notice that $w|_{t=T} = 0$ on ∂M and $\Delta u \in C([0, T]; L^2(M))$. Thus we have $w \in C([0, T]; H_0^1(M))$, and $A_{c_0} \Lambda_c f = v|_{t=0} = f - w|_{t=0} \in H_0^1(M)$. \square

Let us recall that $A_{c_0} : H^1((0, T) \times \partial M) \rightarrow H^1(M)$ is bounded [11], and that $f = R_{c_0} \Lambda_{c_0} f$ [14]. Thus there is $C > 0$ depending only on c_0 and M such that

$$\begin{aligned} \left\| \tilde{f} - f \right\|_{H^1(M)} &= \|R_{c_0}(\Lambda_c f - \Lambda_{c_0} f)\|_{H^1(M)} \\ &\leq C \|\Lambda_c f - \Lambda_{c_0} f\|_{H^1((0, T) \times \partial M)}. \end{aligned}$$

Notice that $\Lambda_c f - \Lambda_{c_0} f = w|_{(0,T) \times \partial M}$, where w is the solution of the wave equation,

$$\begin{aligned} \partial_t^2 w - c_0^2 \Delta w &= (c^2 - c_0^2) \Delta u, & \text{in } (0, \infty) \times \mathbb{R}^n, \\ w|_{t=0} = 0, \quad \partial_t w|_{t=0} &= 0, & \text{in } \mathbb{R}^n, \end{aligned}$$

and u is the solution of (1).

By Theorem A.5 in the appendix below, there is a constant $C > 0$ depending only on c_0, T and M such that

$$\begin{aligned} \|w|_{(0,T) \times \partial M}\|_{H^1((0,T) \times \partial M)} &\leq C \|(c^2 - c_0^2) \Delta u\|_{L^2((0,T) \times \mathcal{K})} \\ &\leq C \|c^2 - c_0^2\|_{L^\infty(\mathcal{K})} \|\Delta u\|_{L^2((0,T) \times \mathcal{K})}. \end{aligned}$$

By applying the energy estimates, see e.g., [10, p. 153], on $\partial_t u$ we see that there is a constant $C > 0$ depending only on T and the bounds

$$(6) \quad \epsilon_1 \leq c(x), \quad \|c\|_{C^1(\mathbb{R}^n)} \leq C_1,$$

such that

$$\|\Delta u\|_{L^2((0,T) \times \mathcal{K})} \leq C \|f\|_{H^2(\mathcal{K})} \leq C \|f\|_{H^1(\mathcal{K})}^{1/2} \|f\|_{H^3(\mathcal{K})}^{1/2}.$$

Thus there is $C > 0$ depending only on c_0, T, M and the bounds (6) such that

$$\|\Lambda_c f - \Lambda_{c_0} f\|_{H^1((0,T) \times \partial M)} \leq C \|c^2 - c_0^2\|_{L^\infty(\mathcal{K})} \|f\|_{H^1(\mathcal{K})}^{1/2} \|f\|_{H^3(\mathcal{K})}^{1/2}.$$

By [14] there is a constant $C(c) > 0$ depending on c such that

$$(7) \quad \|f\|_{H^1(M)} \leq C(c) \|\Lambda_c f\|_{H^1((0,T) \times \partial M)}.$$

Theorem 1 follows after we show that $C(c)$ is uniformly bounded in a neighborhood of c_0 . We do not know how to prove this simply by perturbing the estimate (7). However, the uniform boundedness follows from a modification of the Carleman estimate in [12]. We will prove the modified estimate in the next section under the geometric assumptions formulated in Theorem 1; see Corollary 1 below.

The Carleman estimate in [12] is for a wave equation with homogeneous Dirichlet boundary conditions. Here we have no boundary conditions on ∂M and a study of boundary terms is required.

4. A Carleman estimate with explicit constants

As the proof of the Carleman estimate below is of geometric nature, we will consider the wave equation

$$\partial_t^2 u - \Delta_{g,\mu} u = 0, \quad \text{in } (0, T) \times M$$

on a smooth compact Riemannian manifold (M, g) with boundary. Here $\Delta_{g,\mu}$ is the weighted Laplace–Beltrami operator,

$$\Delta_{g,\mu} u = \mu^{-1} \operatorname{div}_g(\mu \nabla_g u),$$

where $\mu \in C^\infty(M)$ is strictly positive and div_g and ∇_g denote the divergence and the gradient with respect to the metric tensor g . In order to prove Theorem 1, we apply the results below to $g = c(x)^{-2} dx^2$ and $\mu(x) = c(x)^{n-2}$. Then $\Delta_{g,\mu} = c(x)^2 \Delta$, where Δ is the Euclidean Laplacian.

We denote by $|\cdot|_g$, $(\cdot, \cdot)_g$, dV_g , dS_g , ν_g and D_g^2 the norm, the inner product, the volume and the surface measures, the exterior unit normal vector and the Hessian with respect to g . Moreover, we write $\operatorname{div}_{g,\mu} X := \mu^{-1} \operatorname{div}_g(\mu X)$ for a vector field X , and will omit writing the subscript g when considering a fixed Riemannian metric tensor.

Let us recall the pointwise Carleman estimate from [12].

Theorem 2. *Let $\ell \in C^3(M)$ and $\rho > 0$ satisfy*

$$(8) \quad D^2 \ell(X, X) \geq \rho |X|^2, \quad X \in T_x M, \quad x \in M.$$

Let $\tau > 0$ and $u \in C^2(\mathbb{R} \times M)$. We define

$$\begin{aligned} v &:= e^{\tau \ell} u, \quad \vartheta := \tau((\Delta_\mu \ell - \rho)v + 2(\nabla v, \nabla \ell)), \\ Y &:= \tau((\partial_t v)^2 + |\nabla v|^2 - (\tau \rho - \tau^2 |\nabla \ell|^2) v^2) \nabla \ell. \end{aligned}$$

Then

$$\begin{aligned} &e^{2\tau \ell} (\partial_t^2 u - \Delta_\mu u)^2 / 2 - \partial_t (\vartheta \partial_t v) + \operatorname{div}_\mu (\vartheta \nabla v) + \operatorname{div}_\mu Y \\ &\geq e^{2\tau \ell} (\rho \tau - 1) ((\partial_t u)^2 + |\nabla u|^2) / 2 + e^{2\tau \ell} (2\rho |\nabla \ell|^2 \tau - C_1) \tau^2 u^2, \end{aligned}$$

where $C_1 = \rho^2 + \max_{x \in M} |\nabla(\Delta_\mu \ell(x))|^2$.

Lemma 2. *Let $T > 0$, $u \in C^2([0, T] \times M)$, $\tau > 1$, $\rho > 0$ and let $\ell \in C^3(M)$. We define v , ϑ and Y as in Theorem 2. Moreover, we write $dm := \mu dV$ and $dn := \mu dS$. Then*

$$\begin{aligned} & \int_0^T \int_M \operatorname{div}_\mu(\vartheta \nabla v) + \operatorname{div}_\mu Y \, dm \, dt \\ & \leq (6C_2 + C_3)e^{B_\ell \tau} \tau \int_0^T \int_{\partial M} (\partial_t u)^2 + |\nabla u|^2 \, dn \, dt \\ & \quad + (7C_2^3 \tau^2 + 5C_2 C_3 \tau)e^{B_\ell \tau} \tau \int_0^T \int_{\partial M} u^2 \, dn \, dt, \end{aligned}$$

where

$$C_2 = \max_{x \in M} (|\nabla \ell(x)| + 1), \quad C_3 = \max_{x \in M} (\rho + |\Delta_\mu \ell|)/2,$$

and $B_\ell = 2 \max_{x \in M} \ell(x)$. Moreover,

$$\begin{aligned} & \int_M |\vartheta \partial_t v| \, dm \\ & \leq C_F (2C_2^2 \tau + C_3) e^{B_\ell \tau} \tau \left(\int_M (\partial_t u)^2 + |\nabla u|^2 \, dm + \int_{\partial M} u^2 \, dn \right), \end{aligned}$$

where $C_F \geq 1$ is a constant satisfying the Friedrichs' inequality

$$(9) \quad \int_M \phi^2 \, dm \leq C_F \left(\int_M |\nabla \phi|^2 \, dm + \int_{\partial M} \phi^2 \, dn \right), \quad \phi \in C^\infty(M).$$

Proof. We have

$$\begin{aligned} \nabla v &= e^{\tau \ell} (\nabla u + \tau u \nabla \ell), \\ \vartheta &= \tau e^{\tau \ell} ((\Delta_\mu \ell - \rho + 2\tau |\nabla \ell|^2)u + 2(\nabla u, \nabla \ell)), \\ Y &= \tau e^{2\tau \ell} ((\partial_t u)^2 + |\nabla u|^2 + 2\tau u(\nabla u, \nabla \ell) + (2\tau^2 |\nabla \ell|^2 - \tau \rho)u^2) \nabla \ell. \end{aligned}$$

Thus

$$\begin{aligned} & \tau^{-1} e^{-2\tau \ell} (\vartheta(\nabla v, \nu) + (Y, \nu)) \\ &= [(\partial_t u)^2 + |\nabla u|^2 + 4\tau u(\nabla u, \nabla \ell) \\ & \quad + (4\tau^2 |\nabla \ell|^2 - 2\tau \rho + \tau \Delta_\mu \ell)u^2](\nabla \ell, \nu) \\ & \quad + ((\Delta_\mu \ell - \rho + 2\tau |\nabla \ell|^2)u + 2(\nabla u, \nabla \ell)) (\nabla u, \nu). \end{aligned}$$

We estimate the cross-terms as follows:

$$\begin{aligned} 4\tau|u(\nabla u, \nabla \ell)| &\leq 2\tau^2|\nabla \ell|^2u^2 + 2|\nabla u|^2, \\ |\Delta_\mu \ell - \rho||u(\nabla u, \nu)| &\leq |\Delta_\mu \ell - \rho|/2 u^2 + |\Delta_\mu \ell - \rho|/2 |\nabla u|^2, \\ 2\tau|\nabla \ell|^2|u(\nabla u, \nu)| &\leq \tau^2|\nabla \ell|^3u^2 + |\nabla \ell||\nabla u|^2, \end{aligned}$$

and get

$$\begin{aligned} \tau^{-1}e^{-2\tau\ell}|\vartheta(\nabla v, \nu) + (Y, \nu)| &\leq |\nabla \ell|(\partial_t u)^2 + (6|\nabla \ell| + (\rho + |\Delta_\mu \ell|)/2)|\nabla u|^2 \\ &\quad + (7\tau^2|\nabla \ell|^3 + (4\tau|\nabla \ell| + 1)(\rho + |\Delta_\mu \ell|)/2)u^2. \end{aligned}$$

The first claim follows from the divergence theorem.

For the second claimed inequality, notice that

$$\begin{aligned} \tau^{-1}e^{-2\tau\ell}|\vartheta\partial_t v| &= |((\Delta_\mu \ell - \rho)u + 2(\nabla u + \tau u\nabla \ell, \nabla \ell))\partial_t u| \\ &\leq |(\Delta_\mu \ell - \rho) + 2\tau|\nabla \ell|^2||u\partial_t u| + 2|\nabla \ell||\nabla u||\partial_t u| \\ &\leq |(\Delta_\mu \ell - \rho)/2 + \tau|\nabla \ell|^2|(u^2 + (\partial_t u)^2) + |\nabla \ell|(|\nabla u|^2 + (\partial_t u)^2). \end{aligned}$$

Hence

$$\begin{aligned} e^{-B_\ell\tau}\tau^{-1} \int_M |\vartheta\partial_t v| dm &\leq (\max|\Delta_\mu \ell - \rho|/2 + \tau \max|\nabla \ell|^2) \left(\int_M u^2 dm + \int_M (\partial_t u)^2 dm \right) \\ &\quad + \max|\nabla \ell| \int_M |\nabla u|^2 + (\partial_t u)^2 dm, \end{aligned}$$

and the second claimed inequality follows from (9) with $\phi = u$. □

Remark 1. Let $u \in C^2([0, \infty) \times M)$ be a solution of

$$(10) \quad \partial_t^2 u(t, x) - \Delta_\mu u(t, x) = 0, \quad (t, x) \in (0, \infty) \times M.$$

Then the energy,

$$E(t) := \int_M (\partial_t u(t))^2 + |\nabla u(t)|^2 dm,$$

satisfies

$$E(t) = E(0) + 2 \int_0^t \int_{\partial M} \partial_\nu u(s) \partial_t u(s) \, dn ds.$$

Theorem 3 (Observability inequality). *Suppose that there is a strictly convex function $\ell \in C^3(M)$ with no critical points. Let $\rho, r > 0$ satisfy*

$$(11) \quad D^2\ell(X, X) > \rho|X|^2, \quad |\nabla\ell(x)| > r,$$

for all $X \in T_xM$ and $x \in M$. We write $\beta_\ell = 2 \min_{x \in M} \ell(x)$ and suppose that

$$(12) \quad T > 2C_F(2C_2^2\tau + C_3)e^{(B_\ell - \beta_\ell)\tau}, \quad \text{where } \tau = \max\left(\frac{3}{\rho}, \frac{C_1}{2\rho r^2}, 1\right).$$

Let $u \in C^2([0, T] \times M)$ be a solution of (10). Then

$$E(0) \leq C \int_0^T \int_{\partial M} (\partial_t u)^2 + |\nabla u|^2 + u^2 dn dt,$$

where C depends only on $C_2, C_3, \beta_\ell, B_\ell, \tau$ and T .

Proof. We will integrate the inequality of Theorem 2. Notice that

$$\rho\tau - 1 \geq 2 \quad \text{and} \quad 2\rho|\nabla\ell|^2\tau - C_1 \geq 0,$$

whence

$$(13) \quad \begin{aligned} & e^{\beta_\ell\tau} \int_0^T \int_M (\partial_t u)^2 + |\nabla u|^2 dm dt \\ & \leq \int_0^T \int_M -\partial_t(\vartheta\partial_t v) + \operatorname{div}_\mu(\vartheta\nabla v) + \operatorname{div}_\mu Y dm dt. \end{aligned}$$

We estimate the left-hand side of (13) from below using Remark 1,

$$\begin{aligned} & e^{\beta_\ell\tau} \int_0^T \int_M (\partial_t u)^2 + |\nabla u|^2 dm dt \\ & \geq e^{\beta_\ell\tau} TE(0) - e^{\beta_\ell\tau} T \left(\|\partial_\nu u\|_{L^2((0,T) \times \partial M)}^2 + \|\partial_t u\|_{L^2((0,T) \times \partial M)}^2 \right). \end{aligned}$$

To estimate the right-hand side of (13) from above, we notice that

$$\begin{aligned} & C_0^{-1} \left| \int_0^T \int_M \partial_t(\vartheta\partial_t v) dm dt \right| \\ & \leq E(0) + E(T) + \int_{\partial M} u(0)^2 + u(T)^2 dn \\ & \leq 2E(0) + \|\partial_\nu u\|_{L^2((0,T) \times \partial M)}^2 + \|\partial_t u\|_{L^2((0,T) \times \partial M)}^2 \\ & \quad + \int_{\partial M} u(0)^2 + u(T)^2 dn, \end{aligned}$$

where $C_0 = C_F(2C_2^2\tau + C_3)e^{B_\ell\tau}\tau$. Hence

$$\begin{aligned} & (e^{\beta_\ell\tau}T - 2C_0)E(0) \\ & \leq C_0 \left(\|\partial_\nu u\|_{L^2((0,T)\times\partial M)}^2 + 2C_{Tr} \|u\|_{H^1((0,T)\times\partial M)}^2 \right) \\ & \quad + (6C_2 + C_3)e^{B_\ell\tau}\tau \left(\|\partial_t u\|_{L^2((0,T)\times\partial M)}^2 + \|\nabla u\|_{L^2((0,T)\times\partial M)}^2 \right) \\ & \quad + (7C_2^3\tau^2 + 5C_2C_3\tau)e^{B_\ell\tau}\tau \|u\|_{L^2((0,T)\times\partial M)}^2 \\ & \quad + e^{\beta_\ell\tau}T \left(\|\partial_\nu u\|_{L^2((0,T)\times\partial M)}^2 + \|\partial_t u\|_{L^2((0,T)\times\partial M)}^2 \right), \end{aligned}$$

where $C_{Tr} \geq 1$ is a constant satisfying the trace inequality

$$\|\phi(0)\|_{L^2(\partial M)}^2 + \|\phi(T)\|_{L^2(\partial M)}^2 \leq C_{Tr} \|\phi\|_{H^1((0,T)\times\partial M)}^2.$$

□

We recall the following trace regularity result by Lasiecka et al. [11].

Theorem 4. *Let $f \in C_0^\infty(M)$. There is $C > 0$ depending only on T , M and $c|_{\mathbb{R}^n \setminus M}$ such that*

$$\|\partial_\nu u\|_{L^2((0,T)\times\partial M)} \leq C \|\Lambda_c f\|_{H^1((0,T)\times\partial M)},$$

where u is the solution of (1).

Proof. Apply [11, Theorem 2.1] in the domain $B(0, R) \setminus M$, where $R > 0$ is large enough so that $u(t, x)$ vanish for $t \in [0, T]$ and $x \in \partial B(0, R)$. □

Corollary 1 (Stable observability). *Suppose that there is a strictly convex function $\ell \in C^3(M)$ with no critical points on the Riemannian manifold $(M, c_0^{-2}dx^2)$. Let U_0 be a bounded C^2 neighborhood of c_0 . Then there is a C^1 neighborhood U of c_0 and constants $C, T > 0$ satisfying the following: for all $c \in U_0 \cap U$ the solution of the wave equation (1) satisfies the observability inequality*

$$\|f\|_{H_0^1(M)} \leq C \|\Lambda_c f\|_{H^1((0,T)\times\partial M)}.$$

Proof. As in [12], we see that (11) holds for the Riemannian manifold $(M, c^{-2}dx^2)$ if $c \in U$ and U is small enough, and also that the constants $C_1, C_2, C_3, \beta_\ell, B_\ell, \tau$ and T for $(M, c^{-2}dx^2)$ stay bounded when c belongs to $U_0 \cap U$. □

Appendix A: on the regularity of traces

Let us define the projections

$$\begin{aligned} \pi_0^z &: T_z^*(\mathbb{R}^{1+n}) \rightarrow T_z^*({0} \times \mathcal{K}), \quad z \in {0} \times \mathcal{K}, \\ \pi_{\partial M}^z &: T_z^*(\mathbb{R}^{1+n}) \rightarrow T_z^*((0, T) \times \partial M), \quad z \in (0, T) \times \partial M, \end{aligned}$$

and write $\pi_0(z, \zeta) = (z, \pi_0^z(\zeta))$, $\pi_{\partial M}(z, \zeta) = (z, \pi_{\partial M}^z(\zeta))$. Furthermore, let us denote by $\gamma_{x,\xi}$ the geodesic satisfying $\gamma(0) = x$ and $\dot{\gamma}(0) = \xi$, where $\dot{\gamma}$ is the derivative of γ , and define

$$\tau(x, \xi) := \min\{t > 0; \gamma_{x,\xi}(t) \in \partial M\}.$$

The measurement operator

$$\Lambda_c : C_0^\infty(\mathcal{K}) \rightarrow C^\infty((0, T) \times \partial M)$$

is the sum of two zeroth order Fourier integral operators with canonical relations C_+ and C_- , and the union $C_+ \cup C_-$ consist of the points

$$(\pi_0(s, y, \sigma, \eta), \pi_{\partial M}(t, x, \tau, \xi))$$

such that $(t, x, \tau, \xi) \in T^*\mathbb{R}^{1+n}$ and $(s, y, \sigma, \eta) \in T^*\mathbb{R}^{1+n}$ lie on the same null bicharacteristic of $p(x, \tau, \xi) := \tau^2 - c(x)^2|\xi|^2$, $x \in \partial M$, $y \in \mathcal{K}$ and $s = 0$. Moreover, in a neighborhood of such a point $(y_0, \eta_0) \in T^*\mathcal{K}$ that

$$(A.1) \quad \dot{\gamma}_{y_0, \eta_0}(\tau(y_0, \eta_0)) \notin T^*(\partial M),$$

the canonical relation C_+ is given by the graph of the map

$$(y, \eta) \mapsto (\tau(y, \eta), \gamma_{y,\eta}(\tau(y, \eta)), |\eta|, \dot{\gamma}_{y,\eta}^\top(\tau(y, \eta))),$$

where $\dot{\gamma}^\top$ stands for the tangential projection on $T^*(\partial M)$. An analogous statement holds for C_- with η replaced by $-\eta$; see e.g., [14] for more details. In particular, when restricted near (y_0, η_0) satisfying (A.1) the measurement operator is continuous from H^1 to H^1 [4, Corollary 25.3.2].

The case of tangential intersection, i.e., the case when (A.1) does not hold, is more delicate and in general C_\pm can be a one-sided fold [18]. In this case, the general theory of Fourier integral operators implies continuity from H^1 to $H^{3/4}$ [2]. The following example illustrates how singularities in C_\pm arise.

Example A.1. *Suppose $M \subset \mathbb{R}^2$, $c = 1$ identically and that near $(1, 0)$ the boundary ∂M coincides with the unit circle*

$$S^1 := \{x \in \mathbb{R}^2; |x| = 1\}.$$

If \mathcal{K} intersects the vertical line through $(1, 0)$, then C_+ is not a local canonical graph.

Proof. Let us parametrize S^1 by $x(a) := (\cos a, \sin a)$, $a \in (-\pi, \pi)$. Then the cotangent space $T_{x(a)}^*S^1$ is spanned by $x^\perp(a) := (-\sin a, \cos a)$, and C_+ consist locally of the points $(y, \eta, t, a, \tau, \alpha)$ where

$$y = x(a) - t \frac{\eta}{|\eta|}, \quad \tau = |\eta|, \quad \alpha = \eta \cdot x^\perp(a).$$

Hence C_+ is locally parametrized by η, t and a . Let us consider the projection

$$\pi_L(t, a, \eta) = (y, \eta), \quad \pi_L : C_+ \rightarrow T^*(\mathcal{K}).$$

Its differential is of the form

$$d\pi_L = \begin{pmatrix} -\frac{\eta}{|\eta|} & x^\perp(a) & * \\ 0 & 0 & I \end{pmatrix},$$

where I is the identity matrix and the explicit form of $*$ is not needed for our purposes. Thus $d\pi_L(\delta t, \delta a, \delta \eta) = 0$ if and only if $\delta \eta = 0$ and

$$-\frac{\eta}{|\eta|} \delta t + x^\perp(a) \delta a = 0.$$

The latter equation has a non-trivial solution if and only if η is a scalar multiple of $x^\perp(a)$, i.e., the geodesic $\gamma(t) = y + t \frac{\eta}{|\eta|}$ intersects S^1 tangentially. In this case, $d\pi_L$ has a one-dimensional kernel and C_+ is not a local canonical graph (in fact, C_+ is a two-sided fold [18]). □

If ∂M is strictly convex with respect to the Riemannian metric $c^{-2}dx^2$, then (A.1) holds for all $(y_0, \eta_0) \in T^*\mathcal{K}$. In this case, we have the following regularity result.

Theorem A.5. *Suppose that ∂M is strictly convex with respect to the Riemannian metric $c^{-2}dx^2$ and let $\mathcal{K} \subset M^{int}$ be compact. Let $T > 0$ and let w*

be the solution of the wave equation

$$\begin{aligned} \partial_t^2 w - c^2 \Delta w &= F, & \text{in } (0, T) \times \mathbb{R}^n, \\ w|_{t=0} = 0, \quad \partial_t w|_{t=0} &= 0, & \text{in } \mathbb{R}^n, \end{aligned}$$

where $F \in C_0^\infty((0, T) \times \mathcal{K})$. Then there is $C > 0$ independent of F such that

$$\|w|_{(0, T) \times \partial M}\|_{H^1((0, T) \times \partial M)} \leq C \|F\|_{L^2((0, T) \times \mathcal{K})}.$$

Proof. The operator $F \mapsto w$ coincides with the forward parametrix of [4, Theorem 26.1.14] modulo an operator with a smooth kernel. The composition of the parametrix with the restriction on $(0, T) \times \partial M$ is a Fourier integral operator of order $-5/4$ with the canonical relation C consisting of such points

$$(t, x, |\xi|, \xi, s, \gamma_{x, \widehat{\xi}}(s - t), |\xi|, \dot{\gamma}_{x, \widehat{\xi}}^\top(s - t)),$$

that $\gamma_{x, \widehat{\xi}}(s - t) \in \partial M$, $(x, \xi) \in T^*\mathcal{K} \setminus 0$ and $t, s \in (0, T)$. Here $\widehat{\xi} = \xi/|\xi|$.

As above, the strict convexity of ∂M implies that s is locally a function of $(t, x, \widehat{\xi})$. In particular, the canonical relation is parametrized by (t, x, ξ) and the projection $C \rightarrow T^*((0, T) \times \mathcal{K})$ has the rank $2(n + 1) - 1$. We may apply [3, Theorem 4.3.2] to get the claimed continuity. \square

References

- [1] C. Bardos, G. Lebeau and J. Rauch, *Sharp sufficient conditions for the observation, control, and stabilization of waves from the boundary*, SIAM J. Control Optim. **30**(5) (1992), 1024–1065.
- [2] A. Greenleaf and A. Seeger, *Fourier integral operators with fold singularities*, J. Reine Angew. Math. **455** (1994), 35–56.
- [3] L. Hörmander, *Fourier integral operators. I*, Acta Math. **127**(1–2) (1971), 79–183.
- [4] ———, *The analysis of linear partial differential operators*. In, ‘IV Grundlehren der Mathematischen Wissenschaften’, **275**, Springer-Verlag, Berlin, 1985, ISBN 3-540-13829-3.
- [5] Y. Hristova, *Time reversal in thermoacoustic tomography — an error estimate*, Inverse Probl. **25**(5) (2009), 055008, 14.

- [6] Y. Hristova, P. Kuchment and L. Nguyen, *Reconstruction and time reversal in thermoacoustic tomography in acoustically homogeneous and inhomogeneous media*, Inverse Probl. **24**(5) (2008), 055006, 25.
- [7] X. Jin and L.V. Wang, *Thermoacoustic tomography with correction for acoustic speed variations*, Phys. Med. Biol. **51**(24) (2006), 6437.
- [8] R.A. Kruger, D.R. Reinecke and G.A. Kruger, *Thermoacoustic computed tomography—technical considerations*, Med. Phys. **26**(9) (1999), 1832–1837.
- [9] P. Kuchment and L. Kunyansky, *Mathematics of thermoacoustic tomography*, Eur. J. Appl. Math. **19**(2) (2008), 191–224.
- [10] O. A. Ladyzhenskaya, *The boundary value problems of mathematical physics*, in ‘Applied Mathematical Sciences’, **49**, Springer–Verlag, New York, 1985, ISBN 0-387-90989-3.
- [11] I. Lasiecka, J.-L. Lions and R. Triggiani, *Nonhomogeneous boundary value problems for second order hyperbolic operators*, J. Math. Pures Appl. (9) **65**(2) (1986), 149–192.
- [12] S. Liu and L. Oksanen, *A Lipschitz stable reconstruction formula for the inverse problem for the wave equation*, Trans. Amer. Math. Soc. (in press). Preprint arXiv:1210.1094 (2012).
- [13] J. Qian, P. Stefanov, G. Uhlmann and H. Zhao, *An efficient Neumann series-based algorithm for thermoacoustic and photoacoustic tomography with variable sound speed*, SIAM J. Imaging Sci. **4**(3) (2011), 850–883.
- [14] P. Stefanov and G. Uhlmann, *Thermoacoustic tomography with variable sound speed*, Inverse Probl. **25**(7) (2009), 075011, 16.
- [15] ———, *Thermoacoustic tomography arising in brain imaging*, Inverse Probl. **27**(4) (2011), 045004, 26.
- [16] ———, *Instability of the linearized problem in multiwave tomography of recovery both the source and the speed*, Inverse Probl. Imaging **7**(4) (2013), 1367–1377.
- [17] ———, *Recovery of a source term or a speed with one measurement and applications*, Trans. Amer. Math. Soc. **365**(11) (2013), 5737–5758.
- [18] D. Tataru, *On the regularity of boundary traces for the wave equation*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **26**(1) (1998), 185–206.

- [19] L. Wang, *Photoacoustic imaging and spectroscopy*, in ‘Optical Science and Engineering’, CRC Press, 2009, ISBN 9781420059922.
- [20] M. Xu and L.V. Wang, *Photoacoustic imaging in biomedicine*, Rev. Sci. Instrum. **77**(4) (2006), 041101.
- [21] P.-F. Yao, *On the observability inequalities for exact controllability of wave equations with variable coefficients*, SIAM J. Control Optim. **37**(5) (1999), 1568–1599 (electronic).

UNIVERSITY COLLEGE LONDON
DEPARTMENT OF MATHEMATICS
GOWER STREET, LONDON WC1E 6BT
UK
E-mail address: `l.oksanen@ucl.ac.uk`

UNIVERSITY OF WASHINGTON
DEPARTMENT OF MATHEMATICS
BOX 354350
SEATTLE, WA 98195-4350
E-mail address: `gunther@math.washington.edu`

RECEIVED AUGUST 12, 2013