

A view on elliptic integrals from primitive forms (Period integrals of type A_2 , B_2 and G_2)

KYOJI SAITO

Abstract: Elliptic integrals, since Euler’s finding of addition theorem 1751, has been studied extensively from various view points. The present paper gives a view point from primitive integrals of types A_2, B_2 and G_2 . We solve Jacobi inversion problem for the period maps by introducing generalized Eisenstein series of types A_2, B_2 and G_2 , which generate the ring of invariants functions on the period domain for the congruence subgroups $\Gamma_1(N)$ ($N = 1, 2$ and 3). Type A_2 case is classical. Type B_2 and type G_2 cases seems to be new. The goal of the paper is a partial answer to the discriminant conjecture: to show an existence of the cusp form of weight 1 with character of topological origin, which is a power root of the discriminant form (Aspects Math., E36, p. 265–320. 2004).

1	Introduction	908
2	Families of elliptic curves of type A_2, B_2 and G_2	910
3	Real elliptic curves and vanishing cycles	913
4	Fundamental group and monodromy representation	921
5	Periods for primitive forms	928
6	Jacobian variety	934
7	Laurent series solutions at infinities	936
8	Partial fractional expansions	941
9	Eisenstein series of type A_2, B_2 and G_2 (primitive automorphic forms)	945
10	Ring of modular forms and discriminant	951
11	Discriminant conjecture	957

Received August 12, 2019.

12 Concluding remarks	961
Acknowledgements	963
References	963

1. Introduction

We study period integrals for three families of elliptic curves: Weierstrass family, Legendre-Jacobi family and Hesse family. According to the lattice structure of vanishing cycles studied in §3, we call the families of type A_2, B_2 and G_2 , respectively. In order to treat these three types simultaneously, we use notation $I_2(p)$ for the dihedral groups [4] and identify $A_2 = I_2(3)$, $B_2 = I_2(4)$ and $G_2 = I_2(6)$. The theory, we will study, for the type A_2 is nothing but the classical well-known theory of elliptic integrals for the Weierstrass family of elliptic curves. The purpose of the present paper is to show that there exist some parallel worlds for types B_2 and G_2 , even in a somewhat deeper manner.

We refer the reader to §12 Concluding Remarks for the motivation and the background of the present study. Therefore, in the present introduction, we restrict ourselves to explanations of the contents down to the earth.

As mentioned above, we study the three families of types A_2, B_2 and G_2 , which are introduced in §2 as the families of affine elliptic curves $E_{\underline{g}}$, punctured by $N := [p/2]$ -points at infinity, parametrized by the two dimensional space $\underline{g} \in S_{I_2(p)}$ containing the discriminant loci $D_{I_2(p)}$. Here, the number N of punctures is equal to 1, 2 and 3 according to the types A_2, B_2 and G_2 , respectively.

Then in §3, by the help of some real structure of the families, the lattice $L_{\underline{g}}$ of vanishing cycles in the fiber $E_{\underline{g}}$ is described in terms of classical root lattice of type A_2, B_2 and G_2 . The fundamental group of the complement $S_{I_2(p)} \setminus D_{I_2(p)}$ is described in §4 in terms of the Artin group of type A_2, B_2 and G_2 . Then, the image of the monodromy action of the Artin groups on the lattice of vanishing cycles becomes the congruence modular group $\Gamma_1(N)$ of level $N = 1, 2$ and 3. The description of the modular group also gives a construction of its character $\vartheta_{I_2(p)}$ (4.10), which takes values in $2k(I_2(p)) (= 6, 4, 3$ for $p = 3, 4, 6$ (4.8))th roots of unity such that $\vartheta_{I_2(p)}^{k(I_2(p))}$ is the sign character: $\Gamma_1(N) \rightarrow \{\pm 1\}$. We shall come back to $\vartheta_{I_2(p)}$ at the final Theorem 11.1 of the present paper.

The primitive form ζ , as a relative de Rham cohomology class of degree 1 on the total space of the family, is introduced in §5. Then the period map associates to each point $\underline{g} \in S_{I_2(p)} \setminus D_{I_2(p)}$ a homomorphism in $\text{Hom}_{\mathbb{Z}}(L_{\underline{g}}, \mathbb{C})$, $\gamma \mapsto \int_{\gamma} \zeta$. Actually, the period map is a multivalued map from $S_{I_2(p)} \setminus D_{I_2(p)}$ to a

half subspace $\tilde{\mathbb{H}}$ in \mathbb{C}^2 , called the period domain. We ask to give a description of the univalent inversion map from the period domain $\tilde{\mathbb{H}}$ to $S_{I_2(p)}$. In the present paper, we shall call this question *Jacobi inversion problem*, whose answer will be given in §9.

In §§6–9, we study the analytic aspects of the theory. Namely, we consider the indefinite integral of the primitive form over open paths on the elliptic curve E_g starting from any fixed point at infinity as a map from the end-points of paths to the integral values in the complex plane \mathbb{C} . Actually, regarding this integral value z as the time variable, the inverse map of the integral (a pair of meromorphic doubly periodic functions in z) becomes a solution of the Hamilton equation (6.3) of the motion equipped with the energy constraint (6.4). Conversely, any formal meromorphic Laurent solution of the Hamilton equation is convergent. This gives a bijection between the set of formal solutions of the Hamilton equation with the set of points at infinity (§7 Lemma (7.1)).

This bijection between the sets of geometric solutions and formal solutions is a key step to solve the Jacobi inversion problem. Namely, in the next section §8, by a help of the Hamilton equations (6.3) and (6.4), we can expand those global meromorphic solutions on the full z -plane into partial fractions by the help of Weierstrass \mathfrak{p} -functions or zeta-functions, which depends only on the period variable $(\omega_1, \omega_0) \in \tilde{\mathbb{H}}$. Then in §9, again by expanding these global meromorphic functions into Laurent series at the origin $z = 0$, as coefficients of the expansion, we obtain an infinite sequence of functions in $(\omega_1, \omega_0) \in \tilde{\mathbb{H}}$, which we call the *Eisenstein series of type $I_2(p)$* . Comparing this expansion with that already given in §7, we see that the ring of Eisenstein series, as functions defined on $\tilde{\mathbb{H}}$, by pulling back by the period map, is identified with ring of the Cartesian coordinates (g_s, g_t) of the space $S_{I_2(p)}$ (Theorem 9.2). Thus, those Eisenstein series enable us to construct the inversion map $\tilde{\mathbb{H}} \rightarrow S_{I_2(p)}$, which gives the solution to the Jacobi inversion problem.

Next in the section §10, we identify the ring of Eisenstein series of types A_2 , B_2 and G_2 with the ring of modular forms of the congruence group $\Gamma_1(N)$, where the ring of modular forms were already determined explicitly by Aoki-Ibukiyama [2]. In order to determine the homomorphism exactly, we have to determine the exact values of the Eisenstein series at cusps of the modular group. Actually, Eisenstein series of type A_2 are classical, whose Fourier expansions are well known. Eisenstein series of types B_2 and G_2 are no-longer classical. However, when their weights are larger or equal than 3, then their expressions are still obtained by a “shift” of the constant term of the classical Eisenstein series (see (9.1)). Then, it is still possible to evaluate the values of the shifted classical Eisenstein series at cusps by using either the classical Riemann zeta function or Dirichlet’s L-function (see §10, Table 3).

However, Eisenstein series of types B_2 and G_2 of weights less than or equal to 2 have expression by special values of Weierstrass \mathfrak{p} -functions or by difference of special values of Weierstrass zeta-functions, which seem to be less known. We determine their values at cusps in a separate note joint with Aoki [2] (cf. [9, 16]). These determinations lead us to the identification of the ring Eisenstein series of types A_2, B_2 and G_2 with the ring of modular forms of $\Gamma_1(N)$ ($N = 1, 2$ and 3) (Theorem 10.1). It is marvelous that this identification induces further a one to one correspondence of the set of irreducible components of the discriminant of our family of elliptic curves with the set of modular form (up to constant factors) which vanishes exactly once at one $\Gamma_1(N)$ -orbit of cusps (§10 Lemma 10.3).

This modular form, generating the ideal vanishing at an equivalence class of cusps, on one hand as an equation for an irreducible component of the discriminant, is nowhere vanishing on $\tilde{\mathbb{H}}$. On the other hand as a modular form described by theta-series, it has integral Fourier coefficients. Such form can be expressed as suitable quotient of products of shifted Dedekind eta-functions (see Table 7). Doing this for all cusps, we determine the eta-product expressions of the discriminant (11.1) and the reduced discriminant (11.2) of our family of elliptic curves.

These expressions lead us to the final goal Theorem 11.1:

- (1) *There exists a cusp form $\lambda_{I_2(p)}(\tau)$ of weight 1 of the congruence group $\Gamma_1(N)$ with respect to the character $\vartheta_{I_2(p)}$ in §4 such the reduced discriminant of the family of elliptic curves for the type $I_2(p)$ is identified with the $2k(I_2(p))$ th power of this cusp form ($p = 3, 4$ and 6).*
- (2) *The non-reduced discriminants of the families of elliptic curves of all types A_2, B_2 and G_2 , up to a rational constant times the power π^{12} , are identified with the modular discriminant $q \prod_{n=1}^{\infty} (1 - q^n)^{24}$ of weight 12.*

These two statements give positive answers in our toy model cases to the general discriminant conjecture 6 in [23, §6], and we close the present paper.

2. Families of elliptic curves of type A_2, B_2 and G_2

We start with the three families, defined by the equations (2.1), of affine elliptic curves in the (x, y) -plane parametrized by two weighted homogeneous coordinates $\underline{g} = (g_s, g_l)$ (where s and l stands for small or large weights of the coordinates so that the equations become weighted homogeneous polynomials.

See Table 1).

$$\begin{aligned}
 (2.1) \quad A_2 & : F_{A_2}(x, y, \underline{g}) := y^2 - (4x^3 - g_s x - g_l) \\
 B_2 & : F_{B_2}(x, y, \underline{g}) := y^2 - (x^4 - g_s x^2 + g_l + g_s^2/8) \\
 G_2 & : F_{G_2}(x, y, \underline{g}) := x(y^2 - x^2) + g_s(3x^2 + y^2) - g_l - 2g_s^3.
 \end{aligned}$$

Historically, they are called Weierstrass, Legendre-Jacobi and Hesse family of elliptic curves (To be exact, compared with historical expressions, some coordinate change is done from a view point of primitive forms. See Footnote 1 and 2).

As given in the left side of the equations, we call the families by the names of root systems of rank 2, i.e. by A_2, B_2 and G_2 .¹ We shall justify this renaming of the families in §3 from a view point of vanishing cycles. In order to treat these three cases simultaneously, let us use the notation $I_2(p)$ for dihedral groups [4]. Namely, let us recall the following identifications.

$$(2.2) \quad A_2 = I_2(3), \quad B_2 = I_2(4), \quad G_2 = I_2(6)$$

These three cases are exactly the cases when the dihedral group of type $I_2(p)$ is crystallographic, corresponding to classical root systems defined over \mathbb{Z} which shall play crucial roles in the present paper.

Let us give Table of weights of the variables in the equation $F_{I_2(p)}$. We normalize them so that the total weight of the equation $F_{I_2(p)}$ is equal to 1.

Table 1: Weights of functions and coordinates

Type	$\text{wt}(F_{I_2(p)})$	$\text{wt}(x)$	$\text{wt}(y)$	$s=\text{wt}(g_s)$	$l=\text{wt}(g_l)$	$\text{wt}(\Delta_{I_2(p)})$	$\text{wt}(z)$
A_2	1	1/3	1/2	2/3	1	2	-1/6
B_2	1	1/4	1/2	1/2	1	3	-1/4
G_2	1	1/3	1/3	1/3	1	4	-1/3

¹To be exact, when we call a family of type A_2, B_2 or G_2 , we shall mean the family given in the equation (2.1) together with an action of an automorphism given in (3.11) of the family. They are subfamilies of the bigger families of affine elliptic curves of type A_2, A_3 and D_4 which admits an automorphism σ of order 1, 2 and 3, respectively, so that the present family is the subfamilies over the parameters which are fixed by σ (see Footnote 8). The study of the periods for the types A_3 and D_4 (unpublished) are beyond the scope of present paper and shall appear elsewhere. See also Footnote 10 and 18.

Here, $\Delta_{I_2(p)}$ and z are the discriminant introduced below (2.5) and the Hamilton time coordinate z of the elliptic curve $\tilde{E}_{I_2(p)}$ introduced in §6 (6.1), respectively. Note that the weight $\text{wt}(z) := \text{wt}(x) + \text{wt}(y) - \text{wt}(F_{I_2(p)})$ of the variable z is negative caused from a classification of vanishing cycles, where the negativity plays an essential role in the present paper.

The equations (2.1) define geometric families of affine elliptic curves. Namely, let us consider the morphisms:

$$(2.3) \quad \pi_{I_2(p)} : X_{I_2(p)} \longrightarrow S_{I_2(p)}, \quad p = 3, 4 \text{ or } 6$$

where $S_{I_2(p)}$ is the two dimensional complex parameter space of the coordinates $\underline{g} = (g_s, g_l)$,² $X_{I_2(p)}$ is the affine subvariety in $\mathbb{C}^2 \times S_{I_2(p)}$ defined by the equation $F_{I_2(p)} = 0$, and $\pi_{I_2(p)}$ is the morphism induced on $X_{I_2(p)}$ from the projection: $\mathbb{C}^2 \times S_{I_2(p)} \rightarrow S_{I_2(p)}$, respectively.

The relative critical point set C_F of the map $\pi_{I_2(p)}$ defined by $\frac{\partial F_{I_2(p)}}{\partial x} = \frac{\partial F_{I_2(p)}}{\partial y} = 0$ lies in $X_{I_2(p)}$ proper finite over the base parameter space $S_{I_2(p)}$. The image set $\pi_{I_2(p)}(C_F)$ in $S_{I_2(p)}$ is a one codimensional subvariety

$$(2.4) \quad D_{I_2(p)} \subset S_{I_2(p)},$$

called the *discriminant loci*, which is parametrizing singular elliptic curves in the family. The non-reduced defining equation $\Delta_{I_2(p)}$ of the discriminant loci together its multiplicity (up to a constant factor) is given by (cf. [19])

$$(2.5) \quad \begin{aligned} \Delta_{A_2} &= -27g_l^2 + g_s^3 = (\sqrt{27}g_l + g_s^{3/2})(-\sqrt{27}g_l + g_s^{3/2}), \\ \Delta_{B_2} &= (8g_l + g_s^2)(-8g_l + g_s^2)^2, \\ \Delta_{G_2} &= (g_l + 2g_s^3)(-g_l + 2g_s^3)^3. \end{aligned}$$

The fiber $E_{I_2(p),\underline{g}} := \pi_{I_2(p)}^{-1}(\underline{g})$ over a point \underline{g} in $S_{I_2(p)}$ is an affine open curve in the (x, y) -plane which is compactified to an elliptic curve by adding a single, two or three points at infinity according as the types A_2 , B_2 or G_2 .⁴

²The coordinates $\underline{g} = (g_s, g_l)$ are, up to constant factor, flat coordinates of the family [20, 23], whose weights are equal to exponents of $I_2(p)$ plus $1/p$ [4].

⁴A compactification of $E_{I_2(p),\underline{g}}$ is obtained by the curve defined in \mathbb{P}^2 by the homogenization of the equation $F_{I_2(p)}$. In case of type A_2 , it is tangent of order 3 to the infinite line at a single infinite point. In case of B_2 , it is twice tangent to the infinite line at the same infinite point (so that we need to normalize the compactified curve at the point to separate branches). In case of type G_2 , it is intersecting transversally with the infinite line at three distinct infinite points.

Using the index p in (2.2), we see that the number of the points at infinity for the family of the type $I_2(p)$ is given by $[p/2]$ ($= 1, 2$ and 3).

Let us denote the points at infinity by

$$(2.6) \quad \infty_1, \dots, \infty_{[p/2]}$$

(where exact labeling of them is fixed in §3), the compactified curve by

$$(2.7) \quad \overline{E}_{I_2(p), \underline{g}} = E_{I_2(p), \underline{g}} \cup \bigcup_{i=1}^{[p/2]} \{\infty_i\} \quad \text{and} \quad \overline{E}_{I_2(p), \underline{g}} \cap X_{I_2(p)} = E_{I_2(p), \underline{g}}$$

and the fiberwise compactified family by

$$(2.8) \quad \overline{\pi}_{I_2(p)} : \overline{X}_{I_2(p)} = X_{I_2(p)} \cup \bigcup_{i=1}^{[p/2]} \{\infty_i \times S_{I_2(p)}\} \longrightarrow S_{I_2(p)}.$$

Actually, $\overline{X}_{I_2(p)}$ is smooth at the points ∞_i and $\overline{\pi}_{I_2(p)}$ is transversal to the divisors $\infty_i \times S_{I_2(p)}$ (see Footnote 4). The weighted homogeneity of the equation $F_{I_2(p)}$ implies that there is the $t \in \mathbb{C}^\times$ -action $(x, y, g_s, g_l) \mapsto (t^{\text{wt}(x)}x, t^{\text{wt}(y)}y, t^{\text{wt}(g_s)}g_s, t^{\text{wt}(g_l)}g_l)$ leaving the space $X_{I_2(p)}$ invariant so that the family (2.3) is equivariant with the action. We note that that action on $X_{I_2(p)}$ extends to $\overline{X}_{I_2(p)}$ continuously so that the divisor $\infty_i \times S_{I_2(p)}$ is invariant and the morphism (2.8) is still equivariant.

The restriction of the family (2.3) over the complement $S_{I_2(p)} \setminus D_{I_2(p)}$ of the discriminant loci (i.e. the space of regular values of the family (2.3)) gives a locally topologically trivial family of punctured elliptic curves, which is relatively compactified to a topologically locally trivial family of compact smooth elliptic curves.

Note Compact elliptic curve has well-known abelian group structure (after choosing the origin). Then, we have the following elementary, but important fact, which we shall reformulate in (5.10) §5.

Fact 1. *The difference of two infinite points $[\infty_i]$ and $[\infty_j]$ ($1 \leq i, j \leq [p/2]$) is a torsion element of order $[p/2]$.*

3. Real elliptic curves and vanishing cycles

We study some real geometry of the family (2.3). It provides a description of the middle (i.e. one dimensional) homology groups of the affine elliptic curves $E_{I_2(p), \underline{g}}$ in terms of vanishing cycles of root systems of types A_2, B_2 and G_2 .

Let us denote by $X_{I_2(p)}^{\mathbb{R}}$ and $S_{I_2(p)}^{\mathbb{R}}$ the subsets of $X_{I_2(p)}$ and $S_{I_2(p)}$ consisting of the points where the coordinates (x, y) and \underline{g} take real values, and call them the real total space and the real parameter space of the family (2.3), respectively. In the real parameter space, we are interested in a particular connected component of the complement $S_{I_2(p)}^{\mathbb{R}} \setminus D_{I_2(p)}$ of the real discriminant loci $D_{I_2(p)} \cap S_{I_2(p)}^{\mathbb{R}}$, called the *totally real component* $\Gamma_{I_2(p)}$ ⁵, defined by

$$(3.1) \quad \Gamma_{I_2(p)} \quad : \quad -|g_s|^{p/2} < cgt < |g_s|^{p/2} \quad \text{and} \quad g_s > 0,$$

and its boundary as the union of two edges

$$(3.2) \quad \partial_{\pm}\Gamma_{I_2(p)} \quad : \quad cgt = \pm|g_s|^{p/2} \quad \text{and} \quad g_s > 0$$

and the origin $\{0\}$ (here $c = \sqrt{27}, 8$ or $1/2$ according as $p = 3, 4$ or 6 (2.5)). See Figure 1 and following explanation (see §4 for explanation of a, b and J).

Explanation The union of edges pathing through the origin 0 is a part of the real discriminant $D_{I_2(p)} \cap S_{I_2(p)}^{\mathbb{R}}$. The union of the lightly and darkly shaded areas, bounded by the edges, is the totally real component $\Gamma_{I_2(p)}$. The boundary $\partial\Gamma_{I_2(p)}$ is the union of the upper boundary $\partial_+\Gamma_{I_2(p)}$, lower boundary $\partial_-\Gamma_{I_2(p)}$ and the origin 0 . We choose points \underline{g}_0 and \underline{g}_{\pm} generically in $\Gamma_{I_2(p)}$ and in




Figure 1: Real discriminant $D_{I_2(p)}^{\mathbb{R}}$, totally real component $\Gamma_{I_2(p)}$ and $J_{I_2(p)}$.

⁵Actually, the discriminant loci $D_{I_2(p)}$ of the family (2.3) is identified with the discriminant loci in the quotient space of a vector space by the irreducible finite reflection group action of type $I_2(p)$. Then, there is the unique connected component Γ of the complement of the real discriminant loci such that the inverse image of a point in Γ is totally real in the original vector space of the representation [22]. Then, Γ is homeomorphic to a chamber of type $I_2(p)$ and $\partial_{\pm}\Gamma_{I_2(p)}$ are homeomorphic to its walls. These facts are used for a calculation of fundamental group of the complement of the discriminant loci in the next 4.

$\partial_{\pm}\Gamma_{I_2(p)}$, respectively. See §4 for the role of the other curves, the dark shaded area, called J , and its boundary component a and b .

We observe the following **Facts 1–5**. of vanishing cycles in the family of affine elliptic curves $E_{I_2(p)}$. Proof is achieved by explicit direct calculations (see Figure 2), and we omit the details of the proof.

Choose a generic point $\underline{g}_0 \in \Gamma_{I_2(p)}$ as the base point (recall Figure 1), and consider the complex affine elliptic curve $E_{I_2(p),\underline{g}_0} = \pi_{I_2(p)}^{-1}(\underline{g})$.

Fact 1. The real affine elliptic curve $E_{I_2(p),\underline{g}_0}^{\mathbb{R}} := E_{I_2(p),\underline{g}_0} \cap X_{I_2(p)}^{\mathbb{R}}$ (exhibited in the first row of Figure 2) consists of a single compact component (=an oval), which we shall call

$$(3.3) \quad \gamma_0$$

and $[p/2]$ -number of non-compact connected components (arcs), which we shall call $\delta_1, \dots, \delta_{[p/2]}$. The non-compact components are bounded by the points at infinity in the compactification $\overline{E}_{I_2(p),\underline{g}_0}$. We fix orientations of the arcs δ_i 's and the numbering of the points at infinity such that the cyclic union:

$$(3.4) \quad \{\infty_1\} \cup \delta_1 \cup \{\infty_2\} \cup \delta_2 \cup \dots \cup \delta_{[p/2]} \cup \{\infty_1\}$$

form an oriented closed cycle (i.e. $\partial\delta_1 = \infty_2 - \infty_1, \dots, \partial\delta_{[p/2]} = \infty_1 - \infty_{[p/2]}$). Then, this cycle is homologous to γ_0 after choosing an orientation of γ_0 accordingly. However, this condition determine only a cyclic ordering of arcs. So, there still remains also an ambiguity of reversing the orientation. We resolve this ambiguity in the following observation.

Fact 2. There exists a unique choice of an orientation and a cyclic numbering of arcs δ_i 's which satisfies the initial direction condition

$$(3.5) \quad (\delta_1, \infty_1) \subset (\mathbb{R}_{>0}, +\infty) \times (\mathbb{R}_{<0}, -\infty)$$

Here, (δ_1, ∞_1) in LHS means the germ of the arc δ_1 at ∞_1 , and the $(\mathbb{R}_{>0}, +\infty) \times (\mathbb{R}_{<0}, -\infty)$ in RHS means a neighborhood of $(+\infty, -\infty)$ in the real space $\mathbb{R}_{>0} \times \mathbb{R}_{<0} \subset \mathbb{C} \times \mathbb{C}$.⁶

⁶A priori, there exists neither a guarantee that there exists such pair (δ_i, ∞_i) satisfying the condition (3.5), nor a guarantee that the condition (3.5) choose the

	A_2	$B_2 = A_3^\sigma$	$G_2 = D_4^\sigma$
$E_{I_2(p), \underline{g}_0}^{\mathbb{R}}$ ($\underline{g}_0 \in \Gamma_{I_2(p)}$)			
$E_{I_2(p), \underline{g}_+}^{\mathbb{R}}$ ($\underline{g}_+ \in \partial_+ \Gamma_{I_2(p)}$)			
$E_{I_2(p), \underline{g}_-}^{\mathbb{R}}$ ($\underline{g}_- \in \partial_- \Gamma_{I_2(p)}$)			
$E_{I_2(p), 0}^{\mathbb{R}}$			

Figure 2: Real elliptic curves of type A_2 , B_2 and G_2 . The curves in the second and the third column of the table, called of type B_2 and G_2 , are also regarded as the real affine elliptic curve of type A_3 and D_4 (see Footnote 1) together with the action of an automorphism $\sigma_{I_2(p)}$ (3.11).

pair (δ_i, ∞_i) uniquely. Therefore, Fact 2. claims that the condition actually chooses the unique one (δ_1, ∞_1) . This is easily confirmed from Figure 2. Furthermore, we remark that the choice (3.5) in the real blow up space of $\mathbb{P}^2(\mathbb{R})$, in case of type B_2 , separates two infinity points ∞_1 and ∞_2 , which the $\mathbb{P}^2(\mathbb{C})$ -compactification did not separate (recall Footnote 3). Indeed, we have $(\delta_2, \infty_2) \subset (\mathbb{R}_{<0}, -\infty) \times (\mathbb{R}_{>0}, +\infty)$.

Fact 3. Let us consider to move the point \underline{g}_0 inside $\Gamma_{I_2(p)}$ to a point \underline{g}_+ on the upper boundary edge $\partial_+\Gamma_{I_2(p)}$ (e.g. move along the path a in Figure 1). Then, accordingly, the cycle γ_0 and each arc δ_i in the fiber $E_{I_2(p),\underline{g}_0}$ are getting close to each other, and, finally at $\underline{g}_+ \in \partial_+\Gamma_{I_2(p)}$ on the boundary, they intersect to a node p_i of the curve $E_{I_2(p),\underline{g}_+}$ ($i = 1, \dots, [p/2]$) (see the second row of Figure 2).

Fact 4. Let us consider to move the point \underline{g}_0 inside $\Gamma_{I_2(p)}$ to a point \underline{g}_- on the lower boundary edge $\partial_-\Gamma_{I_2(p)}$ (e.g. move along the path b in Figure 1). Then, accordingly, the cycle γ_0 in the fiber $E_{I_2(p),\underline{g}_0}$ pinches to a Morse singularity p_0 in the fiber $E_{I_2(p),\underline{g}_-}$ (see the third row of Figure 2). This implies that the cycle γ_0 is the vanishing cycle generated by the Morse singularity p_0 .

Fact 5. The vanishing cycle generated by the nodal point p_i in $E_{I_2(p),\underline{g}_0}$ shall be denoted by

$$(3.6) \quad \gamma_i$$

$i = 1, \dots, [p/2]$. It intersects with γ_0 and δ_i (recall Fact 1.) transversally. Since γ_i lies in the complex curve $E_{I_2(p),\underline{g}}$, we do not exhibit it in Figure 2, but its conceptual expression shall be given in the first row of Figure 3. We choose the orientation of γ_i by the following sign condition on the intersection:

$$(3.7) \quad \begin{aligned} \langle \gamma_0, \gamma_i \rangle &= \langle \delta_i, \gamma_i \rangle = 1 && (i = 1, \dots, [p/2]) \\ \langle \gamma_i, \gamma_j \rangle &= \langle \gamma_j, \gamma_i \rangle = 0 && (i, j = 1, \dots, [p/2]). \end{aligned}$$

Here we denote by $\langle \gamma, \gamma' \rangle$ the intersection number of paths γ and γ' whose sign convention is fixed as follows: The orientation of $E_{I_2(p),\underline{g}_0}$ as a real surface is fixed by its complex structure. If a path γ' crosses another path γ *counter-clockwisely*,⁷ then the local intersection number is $\langle \gamma, \gamma' \rangle = +1$.

Remark 3.1. The most degenerated real curve $E_{I_2(p),0}^{\mathbb{R}}$ is exhibited in the last row of Figure 2.

As a consequence of Facts 1–5, we obtain the following description of the homology group of the affine elliptic curve $E_{I_2(p),\underline{g}_0}$.

Fact 6. *The classes of γ_0 and γ_i ($i = 1, \dots, [p/2]$) in the first homology group of $E_{I_2(p),\underline{g}_0}$ form free basis (see the first row of Figure 3). We denote the classes*

⁷A path in the complex-plane crosses other path *counter-clockwisely*, if and only if their tangent vectors a and b at the crossing point satisfies $\text{Im}(a/b) > 0$.

by the same notation γ_i since we shall use notation $[\gamma_i]$ for another meaning below.

$$(3.8) \quad L := H_1(E_{I_2(p), \underline{g}_0}, \mathbb{Z}) = \mathbb{Z}\gamma_0 \oplus \bigoplus_{i=1}^{[p/2]} \mathbb{Z}\gamma_i.$$

The open embedding $E_{I_2(p), \underline{g}_0} \subset \overline{E}_{I_2(p), \underline{g}_0}$ induces a surjective homomorphism

$$(3.9) \quad L \longrightarrow H_1(\overline{E}_{I_2(p), \underline{g}_0}, \mathbb{Z})$$

whose kernel is the radical

$$\text{rad}(L) = \{\gamma \in L \mid \langle \gamma, \delta \rangle = 0, \forall \delta \in L\}$$

of the lattice L which is additively generated by homologous relations

$$\gamma_1 \sim \cdots \sim \gamma_{[p/2]}.$$

in the compactification $\overline{E}_{I_2(p), \underline{g}_0}$. Hence, $L/\text{rad}(L)$ is a rank 2 free abelian group generated by the equivalence classes $[\gamma_0]$ and $[\gamma_1] = \cdots = [\gamma_{[p/2]}]$.

$$(3.10) \quad L/\text{rad}(L) = \mathbb{Z}[\gamma_1] \oplus \mathbb{Z}[\gamma_0].$$

Here, we denote by $[\gamma]$ the equivalence class of $\gamma \in L$ in the quotient module. The map (3.9) preserves the intersection form, since it is the quotient morphism by the radical.

We next consider a SL_2 -linear automorphism of the (x, y) -plane by⁸

$$(3.11) \quad \begin{aligned} \sigma_{A_2}(x, y) &:= (x, y), \\ \sigma_{B_2}(x, y) &:= (-x, -y), \\ \sigma_{G_2}(x, y) &:= \left(\frac{y-x}{2}, \frac{-3x-y}{2}\right). \end{aligned}$$

It fixes the equation $F_{I_2(p)}$ and, hence, induces an automorphism of order $[p/2]$ of the fibers of the family (2.3) where the base space $S_{I_2(p)}$ is point-wisely

⁸There is an SL_2 -linear automorphism σ of order 1, 2 and 3 on bigger families of type A_2, A_3 and D_4 (see Footnote 1). Then, the action (3.11) is induced from σ as fiber-wise action on the subfamilies of type A_2, B_2 and G_2 , respectively. The σ -action has not only homological implications as discussed in this section, it has another important implication on a certain cohomology class called the primitive form (see Footnote 17).

fixed. Since $\sigma_{I_2(p)}$ leaves the real subspace $X_{I_2(p)}^{\mathbb{R}}$ invariant, it acts on each complex and real curve $E_{I_2(p),\underline{g}}$. One checks directly that $\sigma_{I_2(p)}$ -action induces the cyclic permutation of the cycles γ_i , oriented arcs δ_i and the points ∞_i at infinity for $i \in \mathbb{Z}/[p/2]\mathbb{Z}$, respectively. However, the cycle γ_0 is invariant by the $\sigma_{I_2(p)}$ -action.

Let us consider the sub-lattice of $H_1(E_{I_2(p),\underline{g}_0}, \mathbb{Z})$ consisting of $\sigma_{I_2(p)}$ -fixed elements.

$$(3.12) \quad L_{I_2(p)} := L^{\sigma_{I_2(p)}} = H_1(E_{I_2(p),\underline{g}_0}, \mathbb{Z})^{\sigma_{I_2(p)}}$$

It is immediate to see that $L_{I_2(p)}$ is a rank 2 sub-lattice generated by

$$(3.13) \quad \alpha := \sum_{i=1}^{[p/2]} \gamma_i \quad \text{and} \quad \beta := \gamma_0,$$

whose intersection number is counted by (3.7) as

$$(3.14) \quad \langle \alpha, \beta \rangle = -[p/2], \quad \langle \alpha, \alpha \rangle = 0 \quad \text{and} \quad \langle \beta, \beta \rangle = 0.$$

The composition of the embedding $L_{I_2(p)} \subset L$ with the radical quotient map (3.9) is again an isometric embedding of lattices:⁹

$$(3.15) \quad L_{I_2(p)} \subset L/\text{rad}(L), \quad \alpha \mapsto [p/2][\gamma_1], \quad \beta \mapsto [\gamma_0]$$

of finite index $[p/2]$.

We give below a root lattice theoretic interpretation of what we have calculated above (see also Footnotes 8, 9 and 11), which answers to the question on the naming: A_2, B_2 and G_2 of the family (2.1), posed in §2.

In the following Figure 3, we exhibit:

(1) The first row exhibits a conceptual description of the cycles γ_0 and γ_i ($i = 1, \dots, [p/2]$) in the surface $E_{I_2(p),\underline{g}_0}$. Here, $E_{I_2(p),\underline{g}_0}$ is a complexification of the real curve $E_{I_2(p),\underline{g}_0}^{\mathbb{R}}$ in the first row of Figure 2. Complexification of real curves in the second and the third row of Figure 2 can be obtained from this surface $E_{I_2(p),\underline{g}_0}$ by pinching either the cycles γ_i ($i = 1, \dots, [p/2]$) or the cycle γ_0 to points, respectively.

(2) The second row exhibits the intersection diagram of the basis γ_0 and γ_i ($i = 1, \dots, [p/2]$) of the homology group $L := H_1(E_{I_2(p),\underline{g}_0}, \mathbb{Z})$ (actually, they are known as diagrams of types A_2, A_3 and D_4 , respectively (note Footnote 1 and 10)).

⁹Since we shall no-longer use the lattice L in the present paper, so far as there is a no-confusion, we shall regard $L_{I_2(p)}$ as a sub-lattice of $L/\text{rad}(L)$ (e.g. (4.4)).




Figure 3: Vanishing cycles and associated diagrams.

(3) The third row exhibits the folding of the intersection diagram in previous (2) by the action of the automorphism $\sigma_{I_2(p)}$ which is the intersection diagram for the invariant basis α, β of the invariant homology group (3.12). Actually, they are known as diagrams of types A_2, B_2 and G_2 , respectively.

Consequently, we observe that the $\sigma_{I_2(p)}$ -invariant 1-cycles of the families (2.1) are indexed by the lattices of types A_2, B_2 and G_2 , respectively. This is the reason why we call the families according to the type of the root systems.¹⁰ The singularity at the origin $0 \in \mathbb{C}^2$ of the curve $E_{I_2(p), 0}$ for $\underline{g} = 0$

¹⁰To be exact, what we wrote here needs more explanations in the following sense. The intersection form $\langle \cdot, \cdot \rangle$ on the free abelian group $L := \sum_{i=0}^{\lfloor p/2 \rfloor} \mathbb{Z}\gamma_i$ is skew-symmetric so that the pair $(L, \langle \cdot, \cdot \rangle)$ is not a root lattice. In order to justify what we wrote above, we consider a pair (L, J) of the abelian group L with the non-symmetric Seifert form J on it defined by the Seifert matrices $\begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & -1 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

or $\begin{bmatrix} 1 & -1 & -1 & -1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$ (which is the table of linking numbers between the ordered basis γ_i (3.8) according to $p = 3, 4$ or 6 , respectively). Then, the difference $J_{I_2(p)} - {}^t J_{I_2(p)}$ is a skew symmetric form on L which is identified with the intersection form $\langle \cdot, \cdot \rangle$

(exhibited in the 4th row of Figure 2) equipped with the action of $\sigma_{I_2(p)}$ is called the singularity of type $I_2(p)$ (i.e. of type A_2 , B_2 and G_2 according as $p = 3, 4$ and 6) and the lattice $L_{I_2(p)}$ is called the lattice of vanishing cycles for the singularity. Then, the family (2.3) is the universal unfolding (by the parameter space $S_{I_2(p)}$) of the singularity of type $I_2(p)$.

4. Fundamental group and monodromy representation

We describe the fundamental group of the compliment of discriminant loci $S_{I_2(p)} \setminus D_{I_2(p)}$, and its monodromy representation in the first homology groups of $E_{I_2(p), \underline{g}}$ and that of its compactification $\overline{E}_{I_2(p), \underline{g}}$.

First, we sketch a geometric description of the fundamental group which is valid not only for types A_2, B_2 and G_2 but also for the regular orbit space of any type finite reflection group [5, 8, 22]. Since the formulation of [22] is suitable to the present work, we briefly recall it by specializing the setting.

Consider a translation action $\tau_\varepsilon : S_{I_2(p)} \rightarrow S_{I_2(p)}$, $(g_s, g_l) \mapsto (g_s, g_l + \varepsilon)$ for $\varepsilon \in \mathbb{C}$.¹¹ If $\varepsilon \in \mathbb{R}$, then the action preserves the real space $S_{I_2(p)}^{\mathbb{R}}$. In

on $H_1(E_{I_2(p), \underline{g}}, \mathbb{Z})$. On the other hand, the sum $I := J + {}^t J$ is a symmetric bilinear form so that the pair (L, I) is isomorphic to the root lattice of type A_2, A_3 and D_4 with the simple root basis $\gamma_0, \dots, \gamma_{[p/2]}$. The geometric automorphism (3.11) induces an automorphism of the lattice (L, J) , denoted again by $\sigma_{I_2(p)}$. Then the invariant sub-lattice $L_{I_2(p)} := L^{\sigma_{I_2(p)}}$ is spanned by $\alpha := \sum_{i=1}^{[p/2]} \gamma_i$ and $\beta := \gamma_0$ so that the Seifert form $J_{I_2(p)}$ on the basis gives the matrix $\begin{bmatrix} [p/2] - [p/2] & \\ 0 & 1 \end{bmatrix}$. Then $(L_{I_2(p)}, I|_{L_{I_2(p)}})$ is isomorphic to the root lattice of type A_2, B_2 and G_2 with the simple root basis α and β according as $p = 3, 4$ or 6 .

There are two constructions which realize the above formal justification (they are called mirror symmetric to each other): (1) Consider the category of matrix factorization of the singularity $F_{I_2(p)}(x, y, 0)$ (see [14]). Since the simple singularities are self-mirror, the category is isomorphic to the category of vanishing cycles in the Milnor fiber $F_{I_2(p)}(x, y, \underline{g}_0) = 0$. Actually, one finds strongly exceptional collections generating the category such that their images γ_i in the K -group of the category (\simeq the middle homology group of the Milnor fiber) gives the simple basis where the Euler form $\sum_k (-1)^k Ext^k(\gamma_i, \gamma_j)$ is identified with the Seifert form. (2) Consider the equation $F_{I_2(p)}(x, y, \underline{g}_0) + z^2$ in three variables (x, y, z) . Then the suspensions, denoted by $\Sigma\gamma_i$, of the basis γ_i ($0 \leq i \leq [p/2]$) of the vanishing cycles in the complex affine curve $F_{I_2(p)}(x, y, \underline{g}_0) = 0$ form vanishing cycle basis of the second homology group of the complex affine surface $F_{I_2(p)}(x, y, \underline{g}_0) + z^2 = 0$. Then the intersection form $-\langle \Sigma\gamma_i, \Sigma\gamma_j \rangle$ of the homology classes coincides with the symmetric form I .

¹¹The action τ is naturally obtained by integrating the vector field ∂_{g_l} , called the primitive vector field, which is, up to constant factor well-defined (see [22]).

Figure 1, we draw, together with the original discriminant loci $D_{I_2(p)}$, two shifted discriminant loci $\tau_\varepsilon(D_{I_2(p)})$ and $\tau_{-\varepsilon}(D_{I_2(p)})$ for some $\varepsilon \in \mathbb{R}_{>0}$. Then, the darkly shaded area J in $\Gamma_{I_2(p)} \subset S_{I_2(p)}^{\mathbb{R}}$ exhibit the component (which is homeomorphic to a rhombus) in $\Gamma_{I_2(p)}$ cut by the shifted discriminant loci $\tau_\varepsilon(D_{I_2(p)})$ and $\tau_{-\varepsilon}(D_{I_2(p)})$, where its boundary edges a and b are segments on $\tau_\varepsilon(D_{I_2(p)})$ and $\tau_{-\varepsilon}(D_{I_2(p)})$, respectively.

Let \tilde{a} and \tilde{b} be the path in $S_{I_2(p)} \setminus D_{I_2(p)}$ defined as follows. We choose the base point $\underline{g}_0 \in \Gamma_{I_2(p)}$ at the intersecting point of a and b as in Figure 1. Let $a_{\mathbb{C}}$ and $b_{\mathbb{C}}$ be complexifications of the real segments a and b , embedded in the complex shifted discriminants $\tau_\varepsilon D_{I_2(p)}$ and $\tau_{-\varepsilon} D_{I_2(p)}$, respectively. They intersect with the discriminant $D_{I_2(p)}$ in $S_{I_2(p)}$ transversally: $a_{\mathbb{C}} \cap D_{I_2(p)} = \{\underline{g}_+\}$ and $b_{\mathbb{C}} \cap D_{I_2(p)} = \{\underline{g}_-\}$. The Figure 4 shows how to choose paths \tilde{a} and \tilde{b} in the complexification $a_{\mathbb{C}}$ and in $b_{\mathbb{C}}$ (regarded as closed paths in $S_{I_2(p)} \setminus D_{I_2(p)}$).




Figure 4: Homotopy classes \tilde{a} and \tilde{b} .

Fact 7. *The fundamental group $\pi_1(S_{I_2(p)} \setminus D_{I_2(p)}, \underline{g}_0) =: G_{I_2(p)}$ is generated by the homotopy classes of \tilde{a} and \tilde{b} and determined by the relation*

$$(4.1) \quad \tilde{a}\tilde{b}\dots = \tilde{b}\tilde{a}\dots$$

where the both hand sides are words of alternating sequences of \tilde{a}, \tilde{b} of length p , which start either with \tilde{a} or with \tilde{b} .

Sketch of proof. 1. A direct proof is given by Zariski-van Kampen method [28] w.r.t. the pencils $\{g_s = \text{const.}\}$, which intersects with discriminant loci by two points, giving two generators of the fundamental group easily identified with \tilde{a}, \tilde{b} [22, §4.3] and gives the relation (4.1) by turning $g_s = e^{i\theta}$ ($0 \leq \theta \leq 2\pi$).

Let us sketch another approach in [5, 8, 22], where we regard the discriminant loci $D_{I_2(p)}$ as the discriminant loci for the orbit space for the dihedral group action on a two dimensional vector space. Namely, let $V_{\mathbb{R}}$ be the real 2-space on which the dihedral group $W(I_2(p))$ acts as the reflection group. We suitably (up to constant) identify the invariant ring $S(V_{\mathbb{R}}^*)^{W(I_2(p))}$ with the polynomial ring generated by two variables g_s and g_l (recall (2.1)). So, we do $W(I_2(p)) \setminus V_{\mathbb{C}}$ with $S_{I_2(p)}$ and $(W(I_2(p)) \setminus V_{\mathbb{C}})_{reg} = W(I_2(p)) \setminus (V_{\mathbb{C}} \setminus \cup H_\alpha)$

(=regular orbit space) with $S_{I_2(p)} \setminus D_{I_2}$ (where $\cup H_\alpha$ is the union of complexified reflection hyperplanes). By this identification, the real region $\Gamma_{I_2(p)}$ (3.1) is homeomorphic to any chamber in $V_{\mathbb{R}} \setminus \cup H_\alpha$. Then, the inverse image in $V_{\mathbb{R}}$ of the closure of $*$ is a curved polygon which is dual to the chamber decomposition, namely, hexagon, octagon and dodecagon according as $I_2(p)$ is of type A_2 , B_2 and G_2 . The boundary of the polygon is given by $(\dot{a}\dot{b})^p$ (see [22] Fig.). Here, each \dot{a} (resp. \dot{b}) is a double cover of the closed interval \bar{a} (resp. \bar{b}) which is crossing the reflection hyperplane at its central point. We now consider the inverse image in $V_{\mathbb{C}} \setminus \cup H_\alpha$ of the union $\tilde{a} \cup \tilde{b}$. Actually, the inverse image is free homotopic to the boundary to the curved dual polygon shifted by $\sqrt{-1}\delta$ for some $\delta \in V_{\mathbb{R}} \setminus \cup H_\alpha$ (corresponding to breadth of \tilde{a} and \tilde{b} , Figure 5), where each component of the inverse image of \tilde{a} (resp. \tilde{b}) is homotopic to a shifted edge $\dot{a} + \sqrt{-1}\delta$ (resp. $\dot{b} + \sqrt{-1}\delta$). On the other hand, the shifted real vector space $V_{\mathbb{R}} + \sqrt{-1}\delta$ does not intersect with any reflection hyperplane so that the shifted dual polygon is contractible in $V_{\mathbb{C}} \setminus \cup H_\alpha$. This creates the homotopy relation (4.1). That there is no-more relations follows from a dimension argument, which we omit here. \square

The group $G_{I_2(p)}$ with the presentation in Fact 7 is called the *Artin group of type $I_2(p)$* [6]. The element of the expression (4.1) is the least common multiple of the generators \tilde{a} and \tilde{b} inside the positive monoid of the group and is called the *fundamental element* [12, 6], denoted by¹²

$$\Delta = \Delta_{I_2(p)}.$$

It is well-known (e.g. [6]) that the center of the group $G_{I_2(p)}$ is an infinite cyclic group generated by Δ (types B_2 and G_2) or by Δ^2 (type A_2).

We determine the monodromy actions of the fundamental group $G_{I_2(p)}$ on the lattices L , $L_{I_2(p)}$ and $L/\text{rad}(L)$. Recall Fact 2, 3, 4 and 5 that the “degeneration” of the curve $E_{I_2(p),g}$ moving the parameter g along the path a (resp. b) pinches the cycles γ_i ($i = 1, \dots, [p/2]$) (resp. cycle γ_0) in $E_{I_2(p),g_0}$ to points p_i (resp. p_0). Then the actions $\rho = \rho_{I_2(p)}$ of \tilde{a} (resp. \tilde{b}) on the lattice $L = H_1(E_{I_2(p),g_0}, \mathbb{Z})$ is determined by Picard-Lefschetz formula (i.e. the transvections) of the vanishing cycles $\gamma_1, \dots, \gamma_{[p/2]}$ (resp. γ_0), taking the intersection number (3.7) in account. That is,

$$\rho(\tilde{a})(\gamma_j) = \begin{cases} \gamma_0 - \sum_{i=1}^{[p/2]} \gamma_i & \text{if } j = 0 \\ \gamma_j & \text{if } j = 1, 2, 3 \end{cases} \quad \rho(\tilde{b})(\gamma_j) = \begin{cases} \gamma_0 & \text{if } j = 0 \\ \gamma_j + \gamma_0 & \text{if } j = 1, 2, 3 \end{cases}$$

¹²Here, we have an unfortunate coincidence of notation of the fundamental element with that of the discriminant (2.5). Since we use them in different places, there shall be no confusion.

Then, the other actions on $L_{I_2(p)}$ and $L/\text{rad}(L)$, denoted by the same ρ , are induced from this action either by restriction to the sub-lattice or by the quotient out the radical of the lattice. In particular, the embedding (3.15) is equivariant with the monodromy action. Explicit formulae are given as follows.

Fact 8. 1) *The action of $G_{I_2(p)}$ on $L_{I_2(p)} = H_1(E_{I_2(p), \underline{g}_0}, \mathbb{Z})^{\sigma_{I_2(p)}}$ is given as follows.*

$$(4.2) \quad \rho(\tilde{a})(\alpha, \beta) = (\alpha, \beta) \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad \rho(\tilde{b})(\alpha, \beta) = (\alpha, \beta) \begin{bmatrix} 1 & 0 \\ [p/2] & 1 \end{bmatrix}.$$

2) *The action of $G_{I_2(p)}$ on $L/\text{rad}(L) = H_1(\overline{E}_{I_2(p), \underline{g}_0}, \mathbb{Z})$ is given as follows.*

$$(4.3) \quad \rho(\tilde{a})([\gamma_0], [\gamma_1]) = ([\gamma_0], [\gamma_1]) \begin{bmatrix} 1 & 0 \\ -[p/2] & 1 \end{bmatrix} \quad \text{and} \quad \rho(\tilde{b})([\gamma_0], [\gamma_1]) = ([\gamma_0], [\gamma_1]) \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}.$$

Here, the representations (4.2) and (4.3) are conjugate by the basis change:

$$(4.4) \quad (\alpha, \beta) = ([\gamma_0], [\gamma_1]) \begin{bmatrix} 0 & 1 \\ [p/2] & 0 \end{bmatrix}$$

(3.15), and, hence, they are equivalent.¹³ We determine the image and the kernel of the representations as follows.

Fact 9. *Let us identify $L/\text{rad}(L)$ with \mathbb{Z}^2 by the use of the basis $[\gamma_0]$ and $[\gamma_1]$, and regard the representation (4.3) as a homomorphism $\rho : G_{I_2(p)} \rightarrow \text{SL}_2(\mathbb{Z})$. Then, we have*

¹³In the sequel, we shall treat these two equivalent representations in parallel (e.g. (4.13)), which looks a bit redundant and cumbersome. This subtlety was caused since we, later on, want to study automorphic forms for congruence subgroups (see (4.6) and §6). More precisely, from a view point for the period map of a primitive form, it is natural to consider the lattice $L_{I_2(p)}$ of vanishing cycles. On the other hand, from a view point of classical elliptic integrals for a compact elliptic curve and to connect with classical elliptic modular function theory, it is natural to consider the lattice $L/\text{rad}(L)$. Therefore, in the sequel, when we talk about the matrix expression of the representation ρ , we shall mean those matrices with respect to the basis $[\gamma_0]$ and $[\gamma_1]$ but not the basis α and β . These cautious treatments are necessary for the cases of types B_2 and G_2 but not for A_2 , since the embedding (3.15) is already isomorphic.

1. The fundamental element Δ is represented by ρ as follows.

$$(4.5) \quad \rho_{A_2}(\Delta) = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad \rho_{B_2}(\Delta) = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \quad \text{and} \quad \rho_{G_2}(\Delta) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

In particular, the images $Im(\rho_{A_2})$ and $Im(\rho_{B_2})$ contain $-id$, but $Im(\rho_{G_2})$ does not.

2. The image of the representation ρ in $SL_2(\mathbb{Z})$ is equal to the subgroup

$$(4.6) \quad \Gamma_1([p/2]) := \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL_2(\mathbb{Z}) \mid \begin{bmatrix} a & b \\ c & d \end{bmatrix} \equiv \begin{bmatrix} 1 & b \\ 0 & 1 \end{bmatrix} \pmod{[p/2]} \right\},$$

called a congruence subgroup of $SL_2(\mathbb{Z})$ of level $[p/2]$ (see e.g. [15]):

3. The kernel of the representation ρ is an infinite cyclic group generated by

$$(4.7) \quad (\tilde{a}\tilde{b})^{k(I_2(p))}$$

where $k(I_2(p)) \in \mathbb{Z}_{\geq 2}$ is a number attached to any Coxeter system (see [23, §6, 6.1 ii]) such that

$$(4.8) \quad k(A_2) = 6, \quad k(B_2) = 4 \quad \text{and} \quad k(G_2) = 3.^{14}$$

4. Thus, we have the short exact sequence

$$(4.9) \quad 1 \rightarrow \mathbb{Z} \rightarrow G_{I_2(p)} \rightarrow \Gamma_1([p/2]) \rightarrow 1.$$

That is, the Artin group $G_{I_2(p)}$ is a central extension of an elliptic congruence modular group $\Gamma_1([p/2])$.

Proof. 1. This is a direct calculation for the cases $[p/2] = 1, 2$ and 3 . The fact that $Im(\rho_{A_2})$ does not contain $-id$ is a consequence of ii).

2. Let us show that ρ is surjective to $\Gamma_1([p/2])$. Set $A := \rho(\tilde{a})$ and $B := \rho(\tilde{b})$ in (4.3). Clearly $A, B \in \Gamma_1([p/2])$. We show that $\Gamma_1([p/2])$ is generated by A and B . Consider an element $C \in \Gamma_1([p/2])$ whose $(2, 1)$ (resp. $(2, 2)$) entry is c (resp. d). By definition, we set $c = \bar{c}[p/2]$ for some $\bar{c} \in \mathbb{Z}$. We also know by definition that $d \neq 0$. Then the $(2, 1)$ entry (resp. $(2, 2)$ entry) of CA^k ($k \in \mathbb{Z}$) is equal to $c + kd[p/2] = (\bar{c} + kd)[p/2]$ (resp. is unchanged d).

¹⁴The numbers $k(W)$ for any finite Weyl group W is defined in [23, 6.1] depending only on Coxeter diagram of W .

Then, by the Euclidean division algorithm, we can choose $k \in \mathbb{Z}$ such that $|\bar{c} - kd| \leq |d|/2$. Next, let us consider $C \in \Gamma_1([p/2])$ such that its (2,1) entry $c = \bar{c}[p/2]$ satisfies the condition $|\bar{c}| \leq |d|/2$. If $c = 0$ then the diagonal of C is $\pm(1, 1)$ where $(-1, -1)$ cannot occur for the case $[p/2] = 3$ by the definition of $\Gamma_1(3)$. By multiplying $-id$ if necessary for the cases $[p/2] = 1$ or 2 (recall the result in 1.), we see that C is already of the form B^l for some $l \in \mathbb{Z}$. Suppose $c \neq 0$. We consider CB^l for $l \in \mathbb{Z}$. Then, its (2,1)-entry is unchanged c , but the (2,2) entry is given by $d' := d - lc$. Then after a suitable choice of $l \in \mathbb{Z}$, we have $|d - lc| \leq |c|/2$. Combining the above two procedures, the (2,2) entry d' of the matrix $CA^k B^l$ satisfies $|d'| \leq |c|/2 = |\bar{c}[p/2]|/2 \leq |d|[p/2]/4$. Since $[p/2]/4 < 1$ in our case, this means $|d'| < |d|$. That is, if C is not generated by A and B , there are integers k and l such that the (2,2) entry of $CA^k B^l$ has the absolute value strictly smaller than that of C . This gives an induction proof of $\Gamma_1([p/2]) = \langle A, B \rangle$.

3. As a result of 1., we observe $\rho_{A_2}(\Delta^4)$, $\rho_{G_2}(\Delta^2)$ and $\rho_{G_2}(\Delta)$ are identity matrices. That is, $\Delta_{A_2}^4$, $\Delta_{B_2}^2$ and Δ_{G_2} belong to the kernel, where we have relations $\Delta_{A_2}^4 = (\tilde{a}\tilde{b})^6$, $\Delta_{B_2}^2 = (\tilde{a}\tilde{b})^4$ and $\Delta_{G_2} = (\tilde{a}\tilde{b})^3$. Thus the element (4.7) is contained in the $\ker \rho$. Conversely, let us show that $\ker(\rho_{\Gamma_2(p)}) \subset \text{center of } G_{\Gamma_2(p)}$. Then, we explicitly calculate $\Delta_{A_2}^4$, $\Delta_{B_2}^2$ and Δ_{G_2} generate the kernel. To show this, we use the fundamental domain of $\Gamma_1([p/2])$ (see [23, §6 Assertion 5, 6.]).

4. This is only the rewriting of the results 2. and 3. □

Remark 1. The following characterization of the congruence subgroup is well-known.

$$\begin{aligned} \Gamma_1([p/2]) &= \{m \in \text{SL}_2(\mathbb{Z}) \mid m \text{ preserves the subsets } L/\text{rad}(L) + \frac{1}{[p/2]}[\gamma_0]\} \\ &= \{m \in \text{SL}_2(\mathbb{Z}) \mid m \text{ preserves the subsets } L_{\Gamma_2(p)} \text{ and } L_{\Gamma_2(p)} + [\gamma_1]\} \\ &= \{m \in \text{SL}(L_{\Gamma_2(p)}) \mid m \text{ extends to } \text{SL}_2(\mathbb{Z}) \text{ and preserves } L_{\Gamma_2(p)} + [\gamma_1]\}. \end{aligned}$$

A sketch of proof. Reflections by the roots α and β , satisfy the conditions. Conversely, the conditions on $m = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \text{SL}_2(\mathbb{Z})$ that it preserves L (resp. $L_{\Gamma_2(p)} + [\gamma_1]$) implies $c \equiv 0 \pmod{[p/2]}$ (resp. $d \equiv 1 \pmod{[p/2]}$). □

As a consequence of the description of the congruence subgroup $\Gamma_1([p/2])$ in Fact 9, we introduce a character (see [23, §6 (6.1.6)]) on it, which shall be used to formulate the discriminant conjecture in §11:

$$(4.10) \quad \vartheta_{\Gamma_2(p)} : \Gamma_1([p/2]) \rightarrow \mathbb{C}^\times, \quad \tilde{a}, \tilde{b} \mapsto \exp\left(\frac{\pi\sqrt{-1}}{k(\Gamma_2(p))}\right).$$

Proof. Obviously, the relation (4.1) is satisfied by the images of $\vartheta_{I_2(p)}$. On the other hand, the $\vartheta_{I_2(p)}$ -image of (4.7) is equal to $\exp(2\pi\sqrt{-1})=1$. \square

Note that the $k(I_2(p))$ th power of the character $\vartheta_{I_2(p)}$

$$(4.11) \quad \theta_{I_2(p)} := \vartheta_{I_2(p)}^{k(I_2(p))} : \Gamma_1([p/2]) \rightarrow \{\pm 1\}, \quad \tilde{a}, \tilde{b} \mapsto -1$$

defines also a character for the anti-invariants, and $\vartheta_{I_2(p)}^{2k(I_2(p))}$ is trivial. Actually except for the type G_2 , $\theta_{I_2(p)}$ factor through the sign morphism $W_{I_2(p)} \rightarrow \{\pm 1\}$ of the Weyl group associated with $I_2(p)$.

For a use to describe the period map in §5, we prepare some notations. The monodromy ρ covering space $\tilde{S}_{I_2(p)}$ of $S_{I_2(p)} \setminus D_{I_2(p)}$ is defined by

$$(4.12) \quad \tilde{S}_{I_2(p)} := \ker(\rho_{I_2(p)}) \setminus (S_{I_2(p)} \setminus D_{I_2(p)})^\sim,$$

where $(S_{I_2(p)} \setminus D_{I_2(p)})^\sim$ is the universal covering of $S_{I_2(p)} \setminus D_{I_2(p)}$, on which the fundamental group $G_{I_2(p)}$ acts from left properly and discontinuously after choosing a copy $\tilde{\Gamma}_{I_2(p)}$ in $(S_{I_2(p)} \setminus D_{I_2(p)})^\sim$ of the base point loci $\Gamma_{I_2(p)} \subset S_{I_2(p)} \setminus D_{I_2(p)}$ (3.1) as the base point loci in $(S_{I_2(p)} \setminus D_{I_2(p)})^\sim$. Then we take the quotient of the universal covering by the kernel of the monodromy representation. The image of $\tilde{\Gamma}_{I_2(p)}$ in $\tilde{S}_{I_2(p)}$ is again denoted by $\tilde{\Gamma}_{I_2(p)}$ and called the base point loci of $\tilde{S}_{I_2(p)}$ (which is of homotopy type S^1).

We consider the liftings to $\tilde{S}_{I_2(p)}$ of the local systems of the homology groups $H_1(E_{I_2(p),g}, \mathbb{Z})^{\sigma_{I_2(p)}}$ and $H_1(\bar{E}_{I_2(p),g}, \mathbb{Z})$ over $g \in S_{I_2(p)} \setminus D_{I_2(p)}$ (equivalent to the data in **Fact 9**), and call them $\tilde{L}_{I_2(p)}$ and $\tilde{L}/\text{rad}(\tilde{L})$, respectively. By definition, they are trivial local systems on the space $\tilde{S}_{I_2(p)}$

$$(4.13) \quad \tilde{L}_{I_2(p)} \simeq L_{I_2(p)} \times \tilde{S}_{I_2(p)} \quad \text{and} \quad \tilde{L}/\text{rad}(\tilde{L}) \simeq L/\text{rad}(L) \times \tilde{S}_{I_2(p)}$$

of fibers $L_{I_2(p)}$ (3.12) and $L/\text{rad}(L)$ (3.10). Due to (3.15), we have natural inclusion relation $\tilde{L}_{I_2(p)} \subset \tilde{L}/\text{rad}(\tilde{L})$. For any element of $L_{I_2(p)} \subset L/\text{rad}(L)$, say γ , we shall denote by the same γ the global section of $\tilde{L}_{I_2(p)} \subset \tilde{L}/\text{rad}(\tilde{L})$ associated with it, so far as there may be no-confusion. So, the free basis α and β (3.12) (resp. $[\gamma_1]$ and $[\gamma_0]$ (**Fact 6**)) are lifted to global basis α and β (resp. $[\gamma_1]$ and $[\gamma_0]$) of $\tilde{L}_{I_2(p)}$ (resp. $\tilde{L}/\text{rad}(\tilde{L})$).

Let us show in the next **Fact 10** that not only those closed cycles $[\gamma_i]$ are lifted to global sections over $\tilde{S}_{I_2(p)}$ but also some “special arcs” δ_i on the curve

$\overline{E}_{I_2(p),\underline{g}}$, which originally defined only for the base point loci $\underline{g} \in \Gamma_{I_2(p)}$, have global sections over $\tilde{S}_{I_2(p)}$. Namely, let us first consider the pullback family

$$\tilde{\pi}_{I_2(p)} : \tilde{X}_{I_2(p)} \longrightarrow \tilde{S}_{I_2(p)}$$

of the family $\tilde{\pi}_{I_2(p)}$ (2.8) to $\tilde{S}_{I_2(p)}$. It carries some additional structures:

- (1) global sections $\{\infty_i \times \tilde{S}_{I_2(p)}\}$ ($1 \leq i \leq [p/2]$),
- (2) fiberwise automorphism $\tilde{\sigma}_{I_2(p)}$ such that global sections are cyclically permuted: $\sigma_{I_2(p)}(\infty_i) = \infty_{i+1}$ ($i \in \mathbb{Z}/[p/2]\mathbb{Z}$).

Recall further that if the parameter \underline{g} belongs to the base point locus $\Gamma_{I_2(p)}$, then the fiber curve $\overline{E}_{I_2(p),\underline{g}}$ contains oriented arcs δ_i ($i = 1, \dots, [p/2]$) such that

$$(4.14) \quad \begin{cases} a) & \partial(\delta_i) = \infty_{i+1} - \infty_i & \text{for } i \in \mathbb{Z}/[p/2]\mathbb{Z}, \\ b) & \sigma_{I_2(p)}(\delta_i) = \delta_{i+1} & \text{for } i \in \mathbb{Z}/[p/2]\mathbb{Z}, \\ c) & \sum_{i \in \mathbb{Z}/[p/2]\mathbb{Z}} \delta_i \sim \gamma_0 & \text{homologous in } \overline{E}_{I_2(p),\underline{g}}. \end{cases}$$

We extend these δ_i 's to global sections over $\tilde{S}_{I_2(p)}$ as follows.

Fact 10. 1. There exist global sections for $1 \leq i \leq [p/2]$

$$(4.15) \quad \delta_i : \underline{g} \in \tilde{S}_{I_2(p)} \mapsto \delta_i(\underline{g}) \in C_1(\overline{E}_{I_2(p),\underline{g}}, \mathbb{Z})$$

which coincide with δ_i in **Fact 1.** when $\underline{g} \in \Gamma_{I_2(p)}$, and satisfy the conditions (4.14). Here, $C_1(\overline{E}_{I_2(p),\underline{g}}, \mathbb{Z})$ is the module of singular 1-chains on $\overline{E}_{I_2(p),\underline{g}}$.

2. The sections are unique up to homologous zero. That is, if there exists other section δ'_i , then $\delta_i(\underline{g}) - \delta'_i(\underline{g}) \sim 0$ (homologous in $\overline{E}_{I_2(p),\underline{g}}$) for all $\underline{g} \in \tilde{S}_{I_2(p)}$.

The proof is an elementary topology and is left to the reader.

5. Periods for primitive forms

We study integrals over paths and cycles in the curves $E_{I_2(p),\underline{g}}$ of the family (2.3). Since we are interested in period integrals over the cycles in $H_1(E_{I_2(p),\underline{g}}, \mathbb{Z})$ (3.8), the integrant form should be holomorphic 1-form on $X_{I_2(p)}$ relative to the base space $S_{I_2(p)}$.¹⁵ Still, this condition is too weak to

¹⁵Forms on $X_{I_2(p)}$ relative to the base space $S_{I_2(p)}$ means the equivalence classes of forms on $X_{I_2(p)}$ modulo the $\mathcal{O}_{X_{I_2(p)}}$ -submodule generated by $dF_{I_2(p)}, dg_s$ and dg_l .

fix an integrant (e.g. the Betti number of the curves are larger than 1 so that just one choice of an integrant seems insufficient).

In present paper, we shall focus on the integrals only of the following form:

$$(5.1) \quad \zeta_{I_2(p)} := \text{Res} \left[\frac{dx dy}{F_{I_2(p)}(x, y, \underline{g})} \right]$$

(see Footnote 16 for explicit descriptions of the residue (5.1)¹⁶), which has a characterization, up to a constant factor, as the unique primitive form for the family (2.1).¹⁷

In the present section, we introduce the period map (5.3) by integrating $\zeta_{I_2(p)}$ over closed cycles in $E_{I_2(p), \underline{g}}$. The description of its inversion map (5.6)

¹⁶For each fixed $\underline{g} \in S_{I_2(p)}$, $dx dy / F_{I_2(p)}$ may be considered as a top degree meromorphic 2-form on (x, y) -plane of simple pole along the curve $E_{I_2(p), \underline{g}}$. Then, at the smooth point of $E_{I_2(p), \underline{g}}$, the symbol (5.1) defines the residue, which is a holomorphic one form on $\overline{E}_{I_2(p), \underline{g}}$. Actually, using the (x, y) -coordinates, the form is explicitly given by the relative differential forms $\frac{dx}{\partial F_{I_2(p)} / \partial y} \sim -\frac{dy}{\partial F_{I_2(p)} / \partial x}$ (which are equivalent modulo $dF_{I_2(p)}$). Using these expressions, we confirm that $\zeta_{I_2(p)}$, as a 1-form on $\overline{E}_{I_2(p), \underline{g}}$, is well-defined up to the singularity loci $\{(x, y) \in E_{I_2(p), \underline{g}} \mid \partial F_{I_2(p)} / \partial y = \partial F_{I_2(p)} / \partial x = 0\} =: \text{Sing}(E_{I_2(p), \underline{g}})$ when $\underline{g} \in D_{I_2(p)}$.

¹⁷As we shall see immediately in Fact 11, the form $\zeta_{I_2(p)}$ is, so called, the elliptic integral of the first kind for the compact elliptic curves $\overline{E}_{I_2(p), \underline{g}}$ for $\underline{g} \in S_{I_2(p)}$ (see, for instance, [27]). On the other hand, from a view point of integrals of vanishing cycles of the universal family (2.3) of type $I_2(p)$, the form $\zeta_{I_2(p)}$ is, up to an ambiguity of a constant factor, called the primitive form of the family (to be exact, this is a restriction of the primitive form ζ defined on the bigger family of type A_2 , A_3 or D_4 . Since the action of σ on the big family (recall Footnote 9) preserves the form ζ , it induces the form $\zeta_{I_2(p)}$ (5.1)) (see [19, 23] for primitive forms).

A primitive form in general has distinguished characterizations, which we shall implicitly (but not explicitly) use in the present paper (actually, we used already the action of the primitive vector field in §4, which is an important ingredient of the primitive form theory), and which we recall briefly as follows. Let us consider the twisted relative de-Rham cohomology group $\mathbb{R}\pi_{I_2(p), *}(\Omega_{X_{I_2(p)}/S_{I_2(p)}})$ of the family (2.3). It is a filtered \mathcal{O}_s -module equipped with the Gauss-Manin connection and higher residue pairings. Then, $\zeta_{I_2(p)}$ is an element in the 0th filter satisfying 1) primitivity, 2) homogeneity, 3) orthogonality, 4) holonomicity (see, for instance, [19]). In particular, the primitivity means that the covariant differentiations of $\zeta_{I_2(p)}$ generate all cohomology classes so that just a single choice of $\zeta_{I_2(p)}$ is sufficient.

by the use of Eisenstein series over the lattices $L_{I_2(p)} \subset L/\text{rad}(L)$ is the main subject of the present paper. In order to understand them, the study of integrals over the closed cycles are not sufficient, but we need to develop a study of integrals of $\zeta_{I_2(p)}$ over non closed paths in $\overline{E}_{I_2(p),\underline{g}}$. Precisely, we study integrals over the arcs δ_i in the present section and over indefinite paths in the next section.

We start with the integral of $\zeta_{I_2(p)}$ over a horizontal family of closed cycles $\gamma \in H_1(E_{I_2(p),\underline{g}}, \mathbb{Z})$ (where \underline{g} runs over a simply connected open subset of $S_{I_2(p)} \setminus D_{I_2(p)}$)

$$(5.2) \quad \omega_\gamma := \oint_\gamma \zeta_{I_2(p)}.$$

As is well-known that ω_γ is a holomorphic function in the parameter \underline{g} (which is proven by a use of Leray’s residue formula) on the domain where the family γ is defined. On the other hand, ω_γ depends only on its equivalence class $[\gamma] \in L/\text{rad}(L)$ due to the following fact.

Fact 11. *The form $\zeta_{I_2(p)}$ can be extended holomorphically on the compactified elliptic curves $\overline{E}_{I_2(p),\underline{g}}$ for $\underline{g} \in S_{I_2(p)}$. The extended form is nowhere vanishing on the smooth part $\overline{E}_{I_2(p),\underline{g}} \setminus \text{Sing}(\overline{E}_{I_2(p),\underline{g}})$ of the curve.*

Proof. If $(x, y) \in E_{I_2(p),\underline{g}}$ is a non-singular point, then either of $\partial F_{I_2(p)}(x, y, \underline{g})/\partial x$ or $\partial F_{I_2(p)}(x, y, \underline{g})/\partial y$ is non-zero. Then, one of the two explicit expressions of $\zeta_{I_2(p)}$ in Footnote 16 gives a non-vanishing and holomorphic expression of $\zeta_{I_2(p)}$ at the point (x, y) . At the infinity points, we have already observed in Footnote 3 that $\overline{E}_{I_2(p),\underline{g}}$ is smooth. Then using that expressions, we check again that $\zeta_{I_2(p)}$ is holomorphic and non-vanishing. The details are left to the reader. □

This means that $\zeta_{I_2(p)}$ may also be regarded as a relative de Rham cohomology class of the family of compactified elliptic curves $\overline{E}_{I_2(p),\underline{g}}$. The cycles $\gamma_1, \dots, \gamma_{[p/2]}$ are homologous to each other in $\overline{E}_{I_2(p),\underline{g}}$. This implies $\oint_{\gamma_1} \zeta_{I_2(p)} = \dots = \oint_{\gamma_{[p/2]}} \zeta_{I_2(p)}$, and, ω_γ depends only on the class $[\gamma] \in \tilde{L}/\text{rad}(\tilde{L})$. Therefore, we regard ω_γ as a holomorphic function defined on $\tilde{S}_{I_2(p)}$. In particular, in view of (3.13), we have expressions

$$\omega_\alpha := \oint_\alpha \zeta_{I_2(p)} = [p/2] \omega_{\gamma_1} \quad \text{and} \quad \omega_\beta := \oint_\beta \zeta_{I_2(p)} = \omega_{\gamma_0}.$$

Thus, in the present paper, we shall integrate either over a cycle in $\tilde{L}_{I_2(p)}$ and a cycle γ in $\tilde{L}/\text{rad}(\tilde{L})$.¹⁸ (in such situation, we shall say “integrate a cycle $\gamma \in \tilde{L}_{I_2(p)} \subset \tilde{L}/\text{rad}(\tilde{L})$ ”), or over “special arcs” δ_i (4.15).

For each point $\tilde{g} \in \tilde{S}_{I_2(p)}$ (4.12), we consider the linear map

$$(5.3) \quad \gamma \in \tilde{L}_{I_2(p)} \subset \tilde{L}/\text{rad}(\tilde{L}) \quad \mapsto \quad \omega_\gamma := \oint_\gamma \zeta_{I_2(p)} \in \mathcal{O}_{\tilde{S}_{I_2(p)}}$$

where we set $\mathcal{O}_{\tilde{S}_{I_2(p)}}$ = holomorphic functions on $\tilde{S}_{I_2(p)}$.

Using the trivialization (4.13) of $\tilde{L}_{I_2(p)} \subset \tilde{L}/\sim$, we obtain a holomorphic map

$$(5.4) \quad \begin{aligned} P_{I_2(p)} : \tilde{S}_{I_2(p)} &\rightarrow \text{Hom}_{\mathbb{Z}}(L_{I_2(p)}, \mathbb{C}) = \text{Hom}_{\mathbb{Z}}(L/\text{rad}(L), \mathbb{C}) \quad (\simeq \mathbb{C}^2) \\ \underline{g} &\mapsto \quad \gamma \mapsto \omega_\gamma := \oint_{\gamma(\underline{g})} \zeta_{I_2(p)} \end{aligned}$$

where the identification in RHS is canonically given by the change

$$(5.5) \quad \omega_\alpha = [p/2] \omega_{\gamma_1} \quad \text{and} \quad \omega_\beta = \omega_{\gamma_0}$$

of the basis in both vector spaces. We shall call $P_{I_2(p)}$ the *period map* associated with the primitive form $\zeta_{I_2(p)}$ (5.1).

Fact 12. *The period map $P_{I_2(p)}$ is locally bi-holomorphic.*

Proof. We show the Jacobian determinant of the map is no-where vanishing. We use an essential property: the *primitivity* of $\zeta_{I_2(p)}$ [19]. Since the proof uses relative de-Rham cohomology theory for the family (2.3) which is beyond the scope of present paper, a complete proof is left to the literature but we give here a brief sketch of its idea.

There exists, so called, the Gauss-Manin covariant differentiation operator ∇ on the module (over $\mathcal{O}_{S_{I_2(p)}}$) of relative de Rham cohomology classes

¹⁸It is rather restrictive view point to study integrals only over $\sigma_{I_2(p)}$ -invariant cycles or over equivalent class of cycles. This is caused by the fact that our family (2.1) is already the subfamily of the full families of type A_3 or D_4 , and is fixed by a cyclic action σ (recall Footnote 1 and 7) where the full (A_3 or D_4) lattice L does not play role. The studies of period integrals of the primitive form over the full lattice L in the big families of type A_3 and D_4 (unpublished) are by themselves interesting subject and should appear elsewhere.

of (2.3). Then, one basic property, called the primitivity, of a primitive form is that its covariant differentiations $\nabla_{\partial_{g_s}} \zeta_{I_2(p)}$ and $\nabla_{\partial_{g_l}} \zeta_{I_2(p)}$ generate the relative de-Rham cohomology module (here ∂_{g_s} and ∂_{g_l} stand for the partial derivatives w.r.t. the coordinate system g). Then, standard duality between the de-Rham cohomology group and the $(\sigma_{I_2(p)}$ -invariant) singular homology group of the curve $E_{I_2(p), \underline{g}}$ implies $\det \begin{bmatrix} \oint_{\gamma_0} \nabla_{\partial_{g_s}} \zeta_{I_2(p)} & \oint_{\gamma_0} \nabla_{\partial_{g_l}} \zeta_{I_2(p)} \\ \oint_{\gamma_1} \nabla_{\partial_{g_s}} \zeta_{I_2(p)} & \oint_{\gamma_1} \nabla_{\partial_{g_l}} \zeta_{I_2(p)} \end{bmatrix} \neq 0$. Since the integral \oint commutes with the derivation action ∂_g and the covariant differentiation ∇_{∂_g} , we see that the Jacobian determinant $\det \begin{bmatrix} \partial_{g_s} \oint_{\gamma_0} \zeta_{I_2(p)} & \partial_{g_l} \oint_{\gamma_0} \zeta_{I_2(p)} \\ \partial_{g_s} \oint_{\gamma_1} \zeta_{I_2(p)} & \partial_{g_s} \oint_{\gamma_1} \zeta_{I_2(p)} \end{bmatrix} \neq 0$ does not vanish. \square

Let us now formulate the first main theorem of the present paper.

Theorem 5.1. *The period map (5.4) induces a biholomorphic map*

$$(5.6) \quad P_{I_2(p)} : \tilde{S}_{I_2(p)} \simeq \tilde{\mathbb{H}},$$

where the RHS of (5.6), so called the period domain, is given as

$$(5.7) \quad \begin{aligned} \tilde{\mathbb{H}} &:= \{ \omega \in \text{Hom}_{\mathbb{Z}}(L_{I_2(p)}, \mathbb{C}) \mid \text{Im}(\omega_\alpha / \omega_\beta) > 0 \} \\ &= \{ \omega \in \text{Hom}_{\mathbb{Z}}(L / \text{rad}(L), \mathbb{C}) \mid \text{Im}(\omega_{\gamma_1} / \omega_{\gamma_0}) > 0 \}, \end{aligned}$$

(here we used again (5.5) for the identification of the first and the second lines). The map (5.4) is equivariant with the action of the congruence modular group $\Gamma_1([p/2]) \subset \text{SL}_{\mathbb{Z}}(L_{I_2(p)}) \cap \text{SL}_2(\mathbb{Z})$.

Proof. This result for the case of type $A_2 = I_2(3)$, is the well-known classic (e.g. see [27]). We want to show the parallel world for the other types B_2 and G_2 exists. The complete proof of this theorem can be given after solving the Jacobi-inversion problem for those period maps in Theorem 9.2 in §9. In the present section, we show only the following classical well-known fact:

Fact 13. *The image of the period map (5.4) is contained in RHS of (5.6). In particular, by the period integral (5.2), the lattice $\tilde{L} / \text{rad}(\tilde{L})_{\underline{g}}$ for $\underline{g} \in \tilde{S}_{I_2(p)}$ is embedded into a discrete lattice, called the period lattice:*

$$(5.8) \quad \Omega_{\tilde{L} / \text{rad}(\tilde{L}), \underline{g}} := \mathbb{Z}\omega_{\gamma_0} + \mathbb{Z}\omega_{\gamma_1}$$

in the complex plane \mathbb{C} .

Actually, this is well-known as a consequence of (1) the Riemann's inequality: $\frac{1}{2\sqrt{-1}} \int_{\overline{E_{I_2(p),g}}} \overline{\zeta_{I_2(p)}} \wedge \zeta_{I_2(p)} > 0$, due to the finite positivity of the real volume form $\frac{1}{2\sqrt{-1}} \overline{\zeta_{I_2(p)}} \wedge \zeta_{I_2(p)}$ and (2) the Stokes relation: $\int_{\overline{E_{I_2(p),g}}} \overline{\zeta_{I_2(p)}} \wedge \zeta_{I_2(p)} = \int_{\gamma_1} \zeta_{I_2(p)} - \int_{\gamma_0} \zeta_{I_2(p)} - \int_{\overline{\gamma_0}} \zeta_{I_2(p)} + \int_{\overline{\gamma_1}} \zeta_{I_2(p)}$, due to the fact that the cycles γ_0 and γ_1 give a canonical dissection of the real surface $\overline{E_{I_2(p),g}}$ where $\zeta_{I_2(p)}$ has no poles (i.e. it is only a topological but not analytical property. cf. the first row of Figure 3). \square

Before ending this section, let us consider also the integrals over the special arcs δ_i (4.15) constructed in Fact 1 in §2 and Fact 10 in §4. Namely, for every $\underline{g} \in \tilde{S}_{I_2(p)}$ and $1 \leq i \leq [p/2]$, set

$$(5.9) \quad \omega_{\delta_i} := \int_{\delta_i} \zeta_{I_2(p)}.$$

Recall that the δ_i 's are cyclically permuted by $\sigma_{I_2(p)}$ and their sum is homol-ogous to γ_0 (see (4.15) and Fact 10.1). On the other hand, one sees immedi-ately from the expression (5.1) that the form $\zeta_{I_2(p)}$ is invariant by the action of $\sigma_{I_2(p)}$. These together implies $\omega_{\delta_1} = \dots = \omega_{\delta_{[p/2]}}$ and $\omega_{\delta_1} + \dots + \omega_{\delta_{[p/2]}} = \omega_{\gamma_0}$. Thus those arc integrals (5.9) are expressed in terms of a classical period of a vanishing cycle as follows.

$$(5.10) \quad \omega_{\delta_1} = \dots = \omega_{\delta_{[p/2]}} = \frac{1}{[p/2]} \omega_{\beta} \quad (= \frac{1}{[p/2]} \omega_{\gamma_0})$$

Finally in this section, let us notice some elementary but useful facts.

Fact 14. *The periods ω_{γ_i} and ω_{δ_i} ($i = 1, \dots, [p/2]$) are weighted homogeneous functions on $\tilde{S}_{I_2(p)}$ of weights given in Table 1. That is, the \mathbb{C}^\times action on $S_{I_2(p)}$ is naturally lifted to that on $\tilde{S}_{I_2(p)}$, which, for an abuse of notation, we shall denote $\underline{g} \mapsto t^{\text{wt}(\underline{g})} \underline{g}$, so that we have the following equivariance w.r.t. the action of $t \in \mathbb{C}^\times$.*

$$\omega_{\gamma_i}(t^{\text{wt}(\underline{g})} \underline{g}) = t^{\text{wt}(\omega_{\gamma_i})} \omega_{\gamma_i}(\underline{g}) \quad \text{and} \quad \omega_{\delta_i}(t^{\text{wt}(\underline{g})} \underline{g}) = t^{\text{wt}(\omega_{\delta_i})} \omega_{\delta_i}(\underline{g})$$

where $\text{wt}(\omega_{\gamma_i}) = \text{wt}(\omega_{\delta_i}) = \text{wt}(z)$. Putting negatively graded structure on RHS's of (5.4) and (5.6) by multiplication $t^{\text{wt}(z)}$ for $t \in \mathbb{C}^\times$, the maps (5.4) and (5.6) are equivariant with respect to the \mathbb{C}^\times -action.

Proof. Recall that the \mathbb{C}^\times action on the space $X_{I_2(p)}$ extends to its partial compactification $\overline{X}_{I_2(p)}$ in such manner that the restriction of the action on the divisors $\infty_i \times S_{I_2(p)}$ are equivariant with that on $S_{I_2(p)}$. Then, the action

induces an action on the set of paths in $\overline{E}_{I_2(p),g}$ starting from $\infty_1 \times \underline{g}$, where the end point of the path is acted by the \mathbb{C}^\times . We replace the integral (6.1) over a path by the integral over the “acted” path. Due to the expression (5.1), we have $\text{wt}(z) = \text{wt}(x) + \text{wt}(y) - \text{wt}(F_{I_2(p)})$. □

6. Jacobian variety

We study integrals of the primitive form $\zeta_{I_2(p)}$ over paths in the smooth part of the curve $\overline{E}_{I_2(p),g}$ for each fixed $\underline{g} \in S_{I_2(p)} \setminus D_{I_2(p)}$ which may not necessarily be closed. That is, we study the Jacobian variety of the curve $\overline{E}_{I_2(p),g}$ and the Hamilton’s equation of the motion regarding the unfolding function $F_{I_2(p)}(x, y, \underline{g})$ as the Hamiltonian.¹⁹

Precisely, we focus on the integrals from the point at infinity ∞_1 :

$$(6.1) \quad z := \int_{\infty_1}^{(x,y)^\sim} \zeta_{I_2(p)} \in \mathbb{C}$$

where $(x, y)^\sim$ is a point in the universal covering (with respect to the base point ∞_1) of the curve $\overline{E}_{I_2(p),g}$ which lies over a point $(x, y) \in \overline{E}_{I_2(p),g}$, or, equivalently, a homotopy class of rectifiable paths in the curve $\overline{E}_{I_2(p),g}$ from ∞_1 to $(x, y) \in \overline{E}_{I_2(p),g}$ (as explained, the notation $(x, y)^\sim$ is ambiguous and we use it only in the present formula).

It is a classic that the integral (6.1) induces a biholomorphic map from the universal covering of $\overline{E}_{I_2(p),g}$ to the complex plane \mathbb{C} (*Proof.* The map is locally bi-regular (Fact 11) and is equivariant with the covering transformation of $\pi_1(\overline{E}_{I_2(p),g}, \infty_1)$ on $\overline{E}_{I_2(p),g}$ with the translation action by the full period lattice $\Omega_{\tilde{L}/\text{rad}(\tilde{L})_{\underline{g}}} = \mathbb{Z}\omega_{\gamma_0} \oplus \mathbb{Z}\omega_{\gamma_1}$ on \mathbb{C} (Fact 13)).

For a later use in the study of inversion problem, let us confirm the direction of the Hamiltonian $F_{I_2(p)}$ at the base point $\underline{g}_0 \in \Gamma_{I_2(p)}$ as follows.

Fact 15. *By the map (6.1) for $\underline{g}_0 \in \Gamma_{I_2(p)}$, the cycle (3.4) is mapped to the real interval $[0, \omega_\beta]$ in the z -complex plane.*

In particular, one has the correspondence: $\infty_i \leftrightarrow \frac{i}{|p/2|}\omega_\beta \pmod{\mathbb{Z}\omega_\beta}$.

Proof. Let’s first observe that the integral (6.1) over the arc δ_1 takes real increasing values. For the purpose, we only need to notify that δ_1 is a real path

¹⁹We can study in parallel the case when \underline{g} belongs to the discriminant $D_{I_2(p)}$ by replacing $\overline{E}_{I_2(p),g}$ by $\overline{E}_{I_2(p),g} \setminus \text{Sing}(\overline{E}_{I_2(p),g})$ and $L_{I_2(p)}$ by the lattices of rank 1 and 0. We omitt details which are left to the reader.




Figure 5: Period lattice $\Omega_{G_2, \underline{g}}$ and its fundamental domain. The big black spots indicate points in $\Omega_{G_2, \underline{g}}$, and other small spots indicate other places of the poles of the functions $x_{I_2(p)}(z, \underline{g})$ and $y_{I_2(p)}(z, \underline{g})$. Shaded area is a fundamental domain (=period parallelogram) for translation action of $\Omega_{G_2, \underline{g}}$.

as in Figure 2 and the integrand $\zeta_{I_2(p)} = \frac{dx}{\partial F_{I_2(p)}/\partial y}$ takes positive real values on the arc δ_1 , or equivalently, the function $\partial F_{I_2(p)}/\partial y$ takes negative real values when x is decreasing and takes positive real values when x is increasing on the path δ_1 (near at ∞_1). This can be confirmed directly depending on cases using the condition (3.5). The same argument works for the integral over the arcs $(\delta_2, \infty_2), \dots, (\delta_{[p/2]}, \infty_{[p/2]})$. Then, recall that the cycle (3.4) is homologous to $\gamma_0 = \beta$. \square

We consider the inverse map: $z = \int_{\infty_1}^{(x,y)} \zeta_{I_2(p)} \in \mathbb{C} \mapsto (x, y) \in \overline{E}_{I_2(p), \underline{g}}$. More precisely, we associate to $z \in \mathbb{C}$ the coordinate values of the corresponding point $(x, y) \in \overline{E}_{I_2(p), \underline{g}}$ by the relation (6.1) and denote it by

$$(6.2) \quad x_{I_2(p)}(z, \underline{g}) \quad \text{and} \quad y_{I_2(p)}(z, \underline{g}),$$

respectively.²⁰ To be careful, the values of the “functions” $x_{I_2(p)}$ and $y_{I_2(p)}$ may not be defined when (x, y) represents a point at infinity (2.7), however, the function is obviously holomorphic at other points and we see easily those undefined points are removable to a holomorphic or meromorphic function.²¹

²⁰Inversion expression of the coordinate (x, y) with respect to the integral value $w = \int_{\infty}^{(x,y)} \omega$ over the elliptic integral of the first kind $\omega = dx/\sqrt{4x^3 - g_3x + g_4}$ is well-known to be given by Weierstrass \mathbf{p} -function and its derivative as $x = \mathbf{p}(w), y = \mathbf{p}'(w)$. Since there exists a factor relation $\zeta_{A_2} = \omega/2$, we have the relation $w = 2z$ and the period lattice gets half size. Then, $x_{A_2}(z) = \frac{1}{4}\mathbf{p}(z), y_{A_2}(z) = \frac{1}{8}\mathbf{p}'(z)$.

²¹Here, one should be slightly cautious that $x_{I_2(p)}$ and $y_{I_2(p)}$ are (at present stage) as functions in $\underline{g} \in S_{I_2(p)} \setminus D_{I_2(p)}$ only pointwise. Their holomorphic dependence on \underline{g} can be shown again by a use of Leray’s residue formula (details are omitted).

Recalling the two local expressions in Footnote 16 of the de Rham class of $\zeta_{I_2(p)}$, we obtain the following Hamilton’s equation of motion:

$$(6.3) \quad \frac{\partial x_{I_2(p)}(z, \underline{g})}{\partial z} = \frac{\partial F_{I_2(p)}(x, y, \underline{g})}{\partial y} \quad \text{and} \quad \frac{\partial y_{I_2(p)}(z, \underline{g})}{\partial z} = -\frac{\partial F_{I_2(p)}(x, y, \underline{g})}{\partial x}.$$

However, this equation of the motion (which depends only on g_s but not on g_l) alone does not determine the solution (6.2) uniquely. In order to recover the functions (6.2) as a function in z , we need to put the following (obvious geometric) constraint on the energy level (depending on g_l) of the motion:

$$(6.4) \quad F_{I_2(p)}(x_{I_2(p)}, y_{I_2(p)}, \underline{g}) = 0$$

We remark also that the functions $x_{I_2(p)}$ and $y_{I_2(p)}$, which are no-longer polynomials but meromorphic in z , are still weighted homogeneous functions if we give the weights to x, y, z and \underline{g} as given in the Table 1. That is, we have the following equivariance w.r.t. the action of $t \in \mathbb{C}^\times$.

$$(6.5) \quad \begin{aligned} x_{I_2(p)}(t^{\text{wt}(z)}z, t^{\text{wt}(\underline{g})}\underline{g}) &= t^{\text{wt}(x)} x_{I_2(p)}(z, \underline{g}), \\ y_{I_2(p)}(t^{\text{wt}(z)}z, t^{\text{wt}(\underline{g})}\underline{g}) &= t^{\text{wt}(y)} y_{I_2(p)}(z, \underline{g}) \end{aligned}$$

(*Proof of (6.5).*) Recall that the \mathbb{C}^\times action on the space $X_{I_2(p)}$ extends to its partial compactification $\overline{X}_{I_2(p)}$ in such manner that the point ∞_1 at infinity of a curve stays at infinity ∞_1 of the curve whose parameter \underline{g} is acted by the \mathbb{C}^\times action. Then, the action induces an action on the set of paths starting from ∞_1 to a point in the curve, where the end point of the path is acted by the \mathbb{C}^\times . We replace the integral (6.1) over a path by the integral over the “acted” path. Due to the expression (5.1), we have $\text{wt}(z) = \text{wt}(x) + \text{wt}(y) - \text{wt}(F_{I_2(p)})$. \square .

One crucial fact here is that the only variable z has the negative weight (recall Table 1) and the functions $x_{I_2(p)}(z, \underline{g})$ and $y_{I_2(p)}(z, \underline{g})$ can be (and, actually, are) transcendental in z .

7. Laurent series solutions at infinities

We study formal Laurent series solutions of the equation of the motion (6.3) together with the constraint (6.4) of the energy level and the weight condi-

However, their extendability to $D_{I_2(p)}$ is not a priori obvious). Actually combining with the discussions in Footnote 15, it is possible to show that they actually are extendable to the functions on the whole $S_{I_2(p)}$. However, without using that logic, we show directly in Lemma 7.1 in §7.

tion (6.5). The solutions are exactly in one to one correspondence with the set of points at infinity of the curve $\tilde{E}_{I_2(p),\underline{g}}$, having the geometric origin.

Exactly, we shift the center of Laurent expansion: if $x_{I_2(p)}(z)$ or $y_{I_2(p)}(z)$ has a non trivial pole at $z = \omega(\underline{g})$ (where $\omega(\underline{g})$ is a function of $\underline{g} \in \tilde{S}_{I_2(p)}$ of weight = $\text{wt}(z)$), then we consider the Laurent expansion of the pair

$$(\mathbf{x}(\underline{\mathfrak{z}}), \mathbf{y}(\underline{\mathfrak{z}})) := (x_{I_2(p)}(\underline{\mathfrak{z}} + \omega(\underline{g})), y_{I_2(p)}(\underline{\mathfrak{z}} + \omega(\underline{g})))$$

with respect to the local formal coordinate $\underline{\mathfrak{z}}$ at 0. The pair $(\mathbf{x}(\underline{\mathfrak{z}}), \mathbf{y}(\underline{\mathfrak{z}}))$ satisfies the pair of equations

$$(6.3^*) \quad \frac{\partial \mathbf{x}(\underline{\mathfrak{z}}, \underline{g})}{\partial \underline{\mathfrak{z}}} = \frac{\partial F_{I_2(p)}(\mathbf{x}, \mathbf{y}, \underline{g})}{\partial \mathbf{y}} \quad \text{and} \quad \frac{\partial \mathbf{y}(\underline{\mathfrak{z}}, \underline{g})}{\partial \underline{\mathfrak{z}}} = -\frac{\partial F_{I_2(p)}(\mathbf{x}, \mathbf{y}, \underline{g})}{\partial \mathbf{x}}$$

since the equations (6.3) are invariant by the shift of the center. Next Lemma classify all formal Laurent solutions of (6.3*) with non-trivial poles in $\underline{\mathfrak{z}}$.

Lemma 7.1. *Consider the system of the equations (6.3) together with the constraint (6.4) and the weight condition (6.5). Then it has exactly $[p/2]$ -pairs of formal Laurent series solutions having non-trivial pole, which are in one to one correspondence with the set $\{\infty_1, \dots, \infty_{[p/2]}\}$ of the points at infinity (2.7) of the curve $\bar{E}_{I_2(p)}$.*

$$(7.1) \quad \begin{array}{l} \{ \text{Solutions with non-trivial pole} \} \\ (\mathbf{x}(\underline{\mathfrak{z}}), \mathbf{y}(\underline{\mathfrak{z}})) \end{array} \simeq \begin{array}{l} \{ \text{Points at infinity} \} \\ \lim_{\underline{\mathfrak{z}} \rightarrow 0} (\mathbf{x}(\underline{\mathfrak{z}}), \mathbf{y}(\underline{\mathfrak{z}})) \end{array}$$

The linear transformation $\sigma_{I_2(p)}$ (3.11) acts on the set of solutions cyclically equivariant with the bijection (7.1). The coefficients of the Laurent series are \mathbb{Q} -coefficients weighted homogeneous polynomials in \underline{g} so that the solution is a pair of weighted homogeneous functions of weight $(\text{wt}(x), \text{wt}(y))$.

Proof. The proof is divided into 4 steps.

Step 1. For each type $I_2(p)$, consider the pair of formal Laurent series:

$$(7.2) \quad \begin{array}{l} \mathbf{x}_{A_2}(\underline{\mathfrak{z}}, \underline{g}) = \sum_{n=-a}^{\infty} A_n \underline{\mathfrak{z}}^{2n} \quad \text{and} \quad \mathbf{y}_{A_2}(\underline{\mathfrak{z}}, \underline{g}) = \sum_{n=-b}^{\infty} B_n \underline{\mathfrak{z}}^{2n-1} \\ \mathbf{x}_{B_2}(\underline{\mathfrak{z}}, \underline{g}) = \sum_{n=-a}^{\infty} A_n \underline{\mathfrak{z}}^{2n+1} \quad \text{and} \quad \mathbf{y}_{B_2}(\underline{\mathfrak{z}}, \underline{g}) = \sum_{n=-b}^{\infty} B_n \underline{\mathfrak{z}}^{2n} \\ \mathbf{x}_{G_2}(\underline{\mathfrak{z}}, \underline{g}) = \sum_{n=-a}^{\infty} A_n \underline{\mathfrak{z}}^n \quad \text{and} \quad \mathbf{y}_{G_2}(\underline{\mathfrak{z}}, \underline{g}) = \sum_{n=-b}^{\infty} B_n \underline{\mathfrak{z}}^n \end{array}$$

of indeterminate coefficients A_n and B_n with non vanishing leading terms

$A_{-a}B_{-b} \neq 0$ ²² (according to the even property of x_{A_2} and y_{B_2} or odd property of y_{A_2} and x_{B_2} caused by the $\mathbb{Z}/2$ -symmetry $(x, y) \rightarrow (x, -y)$, the sum consist either only of even or odd powers in \mathfrak{z}). We assume that the pair has non-trivial pole, i.e. at least one of a or b is positive. Here the coefficients are weighted homogeneous functions in $\underline{g} \in \tilde{S}_{I_2(p)}$, a priori not necessarily polynomials, of weight

$$\text{wt}(A_n) = \text{wt}(B_n) = (1 + n)/d$$

where $d = 3, 2$ or 3 according as $I_2(p) = A_2, B_2$ or G_2 (use Table 1 for weights for x, y and z). Therefore, for each type $I_2(p)$, there exists a positive integer n_0 such that

$$\text{wt}(A_{n_0}) = \text{wt}(B_{n_0}) = 1 = \text{wt}(g_i)$$

(actually, $n_0 = d - 1 = 2, 1$ or 2 according as $I_2(p) = A_2, B_2$ or G_2).

Step 2. Using only the equations (6.3*) double inductively on A_n ($n \geq -a$) and B_n ($n \geq -b$), one can uniquely determine the coefficients until the degree $n < n_0$, where we observe two basic facts.

(1) The initial term (A_{-a}, B_{-b}) are constants independent of the parameter, and we have exactly $[p/2]$ -number of solutions. More exactly, according to each initial direction condition listed in the following table (7.3), there exists a unique solution satisfying it, where, in the last case, $\mathbf{x}_{I_2(p)}$ does not have non-trivial pole. The list of explicit solutions is given in (7.4), where one confirm that the coefficients are \mathbb{Q} -coefficient polynomials in g_s . The calculation is case by case and we omit details.

$$(7.3) \quad \begin{array}{ll} A_2 & \infty_1 : A_{-a} > 0 \quad \text{and} \quad B_{-b} < 0 \\ B_2 & \infty_1 : A_{-a} > 0 \quad \text{and} \quad B_{-b} < 0 \\ B_2 & \infty_2 : A_{-a} < 0 \quad \text{and} \quad B_{-b} > 0 \\ G_2 & \infty_1 : A_{-a} > 0 \quad \text{and} \quad B_{-b} < 0 \\ G_2 & \infty_2 : A_{-a} < 0 \quad \text{and} \quad B_{-b} < 0 \\ G_2 & \infty_3 : a \leq 0 \quad \text{and} \quad B_{-b} < 0 \end{array}$$

Already in this initial solutions level (under the assumption that they shall later extend to full solution), we can confirm the bijection (7.1) by the use of Figure 2, where the “roots” of paths δ_i ’s are the infinity points ∞_i . Therefore,

²²Here, we, unfortunately, use the notation A_n and B_n for the coefficients of the Laurent series, which have nothing to do with the classification names A_n and B_n for root systems. Since they are used only inside present proof, one should cautiously read them.

we indicated the point at infinity in the table (7.3) of initial conditions and in the table (7.4) of (partial) solutions.

(2) *The second nontrivial term $((A_1, B_1)$ for type A_2 and (A_0, B_0) for types B_2 and G_2) contains the variable g_s non-trivial linearly.*

Step 3. We next use the energy condition (6.4) to determine the coefficients A_{n_0}, B_{n_0} (actually, the equation (6.3) alone cannot determine the energy level). We confirm that g_l appears non-trivially in A_{n_0}, B_{n_0} . According to the 6 initial conditions listed in (7.3), the results are given below. In particular, we confirm that *the variable g_l appears as a non-trivial linear term in A_{n_0} and/or in B_{n_0} .*

$$\begin{aligned}
 \mathbf{x}_{A_2, \infty_1}(\mathfrak{z}) &= \frac{1}{4}\mathfrak{z}^{-2} + \frac{1}{5}g_s z^2 + \frac{4}{7}g_l\mathfrak{z}^4 + \sum_{n=3}^{\infty} A_n z^{2n} \\
 \mathbf{y}_{A_2, \infty_1}(\mathfrak{z}) &= -\frac{1}{4}\mathfrak{z}^{-3} + \frac{1}{5}g_s\mathfrak{z} + \frac{8}{7}g_l\mathfrak{z}^3 + \sum_{n=3}^{\infty} B_n \mathfrak{z}^{2n-1} \\
 \mathbf{x}_{B_2, \infty_1}(\mathfrak{z}) &= \frac{1}{2}\mathfrak{z}^{-1} + \frac{1}{3}g_s\mathfrak{z} + \left(\frac{1}{18}g_s^2 - \frac{4}{5}g_l\right)\mathfrak{z}^3 + \sum_{n=2}^{\infty} A_n \mathfrak{z}^{2n+1} \\
 \mathbf{y}_{B_2, \infty_1}(\mathfrak{z}) &= -\frac{1}{4}\mathfrak{z}^{-2} + \frac{1}{6}g_s + \left(\frac{1}{12}g_s^2 - \frac{6}{5}g_l\right)\mathfrak{z}^2 + \sum_{n=2}^{\infty} B_n \mathfrak{z}^{2n} \\
 \mathbf{x}_{B_2, \infty_2}(\mathfrak{z}) &= -\frac{1}{2}\mathfrak{z}^{-1} - \frac{1}{3}g_s\mathfrak{z} - \left(\frac{1}{18}g_s^2 - \frac{4}{5}g_l\right)\mathfrak{z}^3 + \sum_{n=2}^{\infty} A_n \mathfrak{z}^{2n+1} \\
 \mathbf{y}_{B_2, \infty_2}(\mathfrak{z}) &= \frac{1}{4}\mathfrak{z}^{-2} - \frac{1}{6}g_s - \left(\frac{1}{12}g_s^2 - \frac{6}{5}g_l\right)\mathfrak{z}^2 + \sum_{n=2}^{\infty} B_n \mathfrak{z}^{2n} \\
 \mathbf{x}_{G_2, \infty_1}(\mathfrak{z}) &= \frac{1}{2}\mathfrak{z}^{-1} + \frac{1}{2}g_s + \frac{3}{2}g_s^2\mathfrak{z} + (g_s^3 - \frac{1}{2}g_l)\mathfrak{z}^2 + \sum_{n=3}^{\infty} A_n \mathfrak{z}^n \\
 \mathbf{y}_{G_2, \infty_1}(\mathfrak{z}) &= -\frac{1}{2}\mathfrak{z}^{-1} + \frac{3}{2}g_s - \frac{3}{2}g_s^2\mathfrak{z} + (3g_s^3 - \frac{3}{2}g_l)\mathfrak{z}^2 + \sum_{n=3}^{\infty} B_n \mathfrak{z}^n \\
 \mathbf{x}_{G_2, \infty_2}(\mathfrak{z}) &= -\frac{1}{2}\mathfrak{z}^{-1} + \frac{1}{2}g_s - \frac{3}{2}g_s^2\mathfrak{z} + (g_s^3 - \frac{1}{2}g_l)\mathfrak{z}^2 + \sum_{n=3}^{\infty} A_n \mathfrak{z}^n \\
 \mathbf{y}_{G_2, \infty_2}(\mathfrak{z}) &= -\frac{1}{2}\mathfrak{z}^{-1} - \frac{3}{2}g_s - \frac{3}{2}g_s^2\mathfrak{z}^2 - (3g_s^3 - \frac{3}{2}g_l)\mathfrak{z}^2 + \sum_{n=3}^{\infty} B_n \mathfrak{z}^n \\
 \mathbf{x}_{G_2, \infty_3}(\mathfrak{z}) &= -g_s + (-2g_s^3 + g_l)\mathfrak{z}^2 + \sum_{n=2}^{\infty} A_{2n}\mathfrak{z}^{2n} \\
 \mathbf{y}_{G_2, \infty_3}(\mathfrak{z}) &= \mathfrak{z}^{-1} + 3g_s^2\mathfrak{z} + \sum_{n=2}^{\infty} B_{2n-1}\mathfrak{z}^{2n-1}
 \end{aligned}
 \tag{7.4}$$

Step 4. To determine the coefficients A_n and B_n for $n > n_0$, we use again the equation (6.3). By inserting (7.4) in (6.3*), compare coefficients of the Laurent expansions in BHS. Let C_n (resp. D_n) be the coefficient polynomial of the power of \mathfrak{z} in the RHS of (6.3) whose degree coincides with the term A_n (resp. B_n) in LHS. The C_n and D_n are rational coefficients weighted homogenous polynomials in g_s, g_l and A_m, B_m ($m \in \mathbb{Z}_{>n_0}$). Since the total weight of C_n (resp. D_n) is equal to $\text{wt}(A_n) = \text{wt}(B_n)$, the coefficient C_n (resp. D_n) cannot contain A_m, B_m for $m > n$ and A_n and B_n appear only linearly with constant coefficients. The linear coefficients are independent of n , since such terms appear in the expansion of RHS of (6.3*) only when the term A_n or B_n multiplied with the constant coefficient terms of $x_{I_2(p)}$ and $y_{I_2(p)}$, that is

lowest degree terms $A_{-a}z^{-a}$ and $B_{-b}z^{-b}$ (see (7.4)). But, such pattern does not depends on $n \in \mathbb{Z}_{>n_0}$. Calculating explicitly the linear coefficients, we obtain the equations:

$$\begin{aligned}
 & A_2 \text{ type : } 2nA_n = 2B_n + C'_n, & (2n - 1)B_n = 6A_n + D'_n \\
 (7.5) \quad & B_2 \text{ type : } (2n + 1)A_n = 2B_n + C'_n, & 2nB_n = 3A_n + D'_n \\
 & G_2 \text{ type : } nA_n = -A_n + B_n + C'_n, & nB_n = 3A_n + B_n + D'_n
 \end{aligned}$$

where C'_n and D'_n are the remaining part of C_n and D_n after subtracting the linear terms in A_n and B_n . We observe immediately that the determinants of coefficients of A_n and B_n in the two equation for three types are given by

$$\begin{aligned}
 A_2 : \quad \det \begin{bmatrix} 2n & -2 \\ -6 & 2n-1 \end{bmatrix} & \quad B_2 : \quad \det \begin{bmatrix} 2n+1 & -2 \\ -3 & 2n \end{bmatrix} & \quad G_2 : \quad \det \begin{bmatrix} n+1 & -1 \\ -3 & n-1 \end{bmatrix} \\
 = 2(2n+3)(n-2), & \quad = 2(2n-3)(n+2), & \quad = (n+2)(n-2)
 \end{aligned}$$

which takes positive values for $n > n_0$. Thus A_n and B_n are uniquely expressed as rational coefficients polynomials in A_m, B_m ($n_0 \leq m < n$) and g_s . This gives the inductive construction of the coefficients A_n and B_n for $n \in \mathbb{Z}_{>n_0}$.

This completes the proof of Lemma 7.1. □

As a result of Lemma 7.1, we can determine the principal parts of the Laurent expansions of the meromorphic functions $x_{I_2(p)}(z, \underline{g})$ and $y_{I_2(p)}(z, \underline{g})$ (6.2).

Fact 16. 1. *The formal Laurent series solutions in Lemma 7.1 are convergent.*
 2. *The following substitutions of \mathfrak{z} in the formal series solution (7.4):*

$$(7.6) \quad \mathbf{x}_{I_2(p), \infty_i} \left(z - \left(\frac{i-1}{[p/2]} \omega_{\gamma_0} + \omega_{\gamma} \right) \right), \quad \mathbf{y}_{I_2(p), \infty_i} \left(z - \left(\frac{i-1}{[p/2]} \omega_{\gamma_0} + \omega_{\gamma} \right) \right)$$

give the Laurent expansions of the meromorphic functions $x_{I_2(p)}(z, \underline{g})$ and $y_{I_2(p)}(z, \underline{g})$ at the place $\frac{i-1}{[p/2]} \omega_{\gamma_0} + \omega_{\gamma}$ for any $1 \leq i \leq [p/2]$ and $\gamma \in \tilde{L}/\text{rad}(\tilde{L})$.

Proof. 1. Can be shown as a consequence of the next 2.

2. We know already from geometry (recall a discussion after the definition (6.2) and the description of Fact 15) that the functions $x_{I_2(p)}(z, \underline{g})$ and $y_{I_2(p)}(z, \underline{g})$ may have poles only at the places $\frac{i-1}{[p/2]} \omega_0 + \omega_{\gamma}$ for $i = 1, \dots, [p/2]$ and $\gamma \in \tilde{L}/\text{rad}(\tilde{L})$. In view of the asymptotic behavior of paths δ_i 's at their starting points in the first row of Figure 1, we observe that all of them become poles except that the function x_{G_2} at the places $\frac{2}{3} \omega_0 + \omega_{\gamma}$ for any $\gamma \in \tilde{L}/\text{rad}(\tilde{L})$ does not have a pole.

Obviously, the Laurent expansions at those places should satisfy the equations (6.3) together with the constraint (6.4), which further satisfy the initial constraint (7.3) according to its location. Then, the uniqueness of the solution of the equations under the initial constraint implies that the formal solution should coincide with the expansion of $x_{I_2(p)}(z, \underline{g})$ or $y_{I_2(p)}(z, \underline{g})$. \square

We note that the proof of Lemma 7.1 for actually covers also the Laurent series expansions of the cases of $\underline{g} \in S_{I_2(p)} \setminus D_{I_2(p)}$ $\underline{g} \in D_{I_2(p)}$. The (7.3) express the Laurent series expansions of the coordinate (x, y) for the cases of $\underline{g} \in D_{I_2(p)}$ (recall Footnote 17 and 19) as trigonometric or rational functions in z .

Let us give explicitly the first few terms of the Laurent expansion of $x_{I_2(p)}(z, \underline{g})$ and $y_{I_2(p)}(z, \underline{g})$ at the origin $z = 0$ as follows.

Table 2: Laurent expansions of $x_{I_2(p)}$ and $y_{I_2(p)}$ at $z = 0$

A_2 case.

$$x_{A_2}(z) = \frac{1}{4}z^{-2} + \frac{1}{5}g_s z^2 + \frac{4}{7}g_l z^4 + \frac{4}{75}g_s^2 z^6 + \frac{48}{385}g_s g_l z^8 + \dots$$

$$y_{A_2}(z) = -\frac{1}{4}z^{-3} + \frac{1}{5}g_s z + \frac{8}{7}g_l z^3 + \frac{4}{25}g_s^2 z^5 + \frac{192}{385}g_s g_l z^7 + \dots$$

B_2 case.

$$x_{B_2}(z) = \frac{1}{2}z^{-1} + \frac{1}{3}g_s z + \left(\frac{1}{18}g_s^2 - \frac{4}{5}g_l\right)z^3 + \left(\frac{1}{27}g_s^3 - \frac{8}{35}g_s g_l\right)z^5 + \dots$$

$$y_{B_2}(z) = -\frac{1}{4}z^{-2} + \frac{1}{6}g_s + \left(\frac{1}{12}g_s^2 - \frac{6}{5}g_l\right)z^2 + \left(\frac{5}{54}g_s^3 - \frac{4}{7}g_s g_l\right)z^4 + \dots$$

G_2 case.

$$x_{G_2}(z) = \frac{1}{2}z^{-1} + \frac{1}{2}g_s + \frac{3}{2}g_s^2 z + (g_s^3 - \frac{1}{2}g_l)z^2 + (\frac{3}{2}g_s^4 - \frac{6}{5}g_l g_s)z^3 + \dots$$

$$y_{G_2}(z) = -\frac{1}{2}z^{-1} + \frac{3}{2}g_s - \frac{3}{2}g_s^2 z + (3g_s^3 - \frac{3}{2}g_l)z^2 - (\frac{3}{2}g_s^4 - \frac{6}{5}g_l g_s)z^3 + \dots$$

8. Partial fractional expansions

We come back to the global study of the meromorphic functions $x_{I_2(p)}(z, \underline{g})$ and $y_{I_2(p)}(z, \underline{g})$ in z (6.2). The goal of this section is to give the partial fractional expansion of them. The study belongs to classical elliptic function theory. In particular, A_2 -type case is well-known as Weierstrass \mathfrak{p} -function theory. We generalize it for the other two types B_2 and G_2 , since those descriptions in the present section, lead to “generalized Eisenstein series” expression of the modular forms for the congruence subgroups $\Gamma_1(2)$ and $\Gamma_1(3)$ in the next section, which seems to have been unknown (see also Remark 9.3 2 in §9).

We recall the classical Weierstrass’s \mathfrak{p} -function and ζ -function associated with any point $(\omega_0, \omega_1) \in \tilde{\mathbb{H}}$ as meromorphic functions on the z -plane with

double or simple poles (see, e.g. [13]).

$$\begin{aligned} \mathbf{p}(z) &= \frac{1}{z^2} + \sum_{\omega \neq 0 \in \Omega} \left(\frac{1}{(z-\omega)^2} - \frac{1}{\omega^2} \right) \\ \zeta(z) &= \frac{1}{z} + \sum_{\omega \neq 0 \in \Omega} \left(\frac{1}{z-\omega} + \frac{1}{\omega} + \frac{z}{\omega^2} \right), \end{aligned}$$

where we set $\Omega := \mathbb{Z}\omega_0 + \mathbb{Z}\omega_1$. Since they are compact uniform convergent on $\mathbb{C} \times \tilde{\mathbb{H}} \setminus \cup_{m,n \in \mathbb{Z}} \{z - (m\omega_1 + n\omega_0) = 0\}$, one may derivate them termwisely. In particular, one has the well-known relation: $\zeta'(z) = -\mathbf{p}(z)$.²³

Theorem 8.1. *The meromorphic functions $x_{\mathbb{I}_2(p)}(z, \underline{g})$ and $y_{\mathbb{I}_2(p)}(z, \underline{g})$ have the following partial fractional expansions.*

(8.1)

$$\begin{aligned} x_{A_2}(z, \underline{g}) &= \frac{1}{4}\mathbf{p}(z) \\ y_{A_2}(z, \underline{g}) &= \frac{1}{8}\mathbf{p}'(z) \\ x_{B_2}(z, \underline{g}) &= -\frac{1}{2}\zeta\left(\frac{1}{2}\omega_0\right) + \frac{1}{2}\zeta(z) - \frac{1}{2}\zeta\left(z - \frac{1}{2}\omega_{\gamma_0}\right) \\ y_{B_2}(z, \underline{g}) &= -\frac{1}{4}\mathbf{p}(z) + \frac{1}{4}\mathbf{p}\left(z - \frac{1}{2}\omega_{\gamma_0}\right) \\ x_{G_2}(z, \underline{g}) &= -\frac{1}{6}\zeta\left(\frac{1}{3}\omega_{\gamma_0}\right) - \frac{1}{6}\zeta\left(\frac{2}{3}\omega_{\gamma_0}\right) + \frac{1}{2}\zeta(z) - \frac{1}{2}\zeta\left(z - \frac{1}{3}\omega_{\gamma_0}\right) \\ y_{G_2}(z, \underline{g}) &= \frac{1}{2}\zeta\left(\frac{1}{3}\omega_{\gamma_0}\right) + \frac{1}{2}\zeta\left(\frac{2}{3}\omega_{\gamma_0}\right) - \frac{1}{2}\zeta(z) - \frac{1}{2}\zeta\left(z - \frac{1}{3}\omega_{\gamma_0}\right) + \zeta\left(z - \frac{2}{3}\omega_{\gamma_0}\right) \end{aligned}$$

where the \mathbf{p} -function and the ζ -function in RHS are those associated with the period $(\omega_{\gamma_0}, \omega_{\gamma_1}) \in \tilde{\mathbb{H}}$, and, hence, with the period lattice $\Omega = \Omega_{\tilde{L}/\text{rad}(\tilde{L}), \underline{g}}$ (5.8).

Proof. Owing to (7.4) and Fact 16, we know already the principal parts of poles of the functions $x_{\mathbb{I}_2(p)}(z, \underline{g})$ and $y_{\mathbb{I}_2(p)}(z, \underline{g})$ in z . Since the principal parts of poles of $\mathbf{p}(z)$, $\mathbf{p}'(z)$ and $\zeta(z)$ are $1/(z - \omega)^2$, $-2/(z - \omega)^3$ and $1/(z - \omega)$ for $\omega \in \Omega_{\mathbb{I}_2(p), \underline{g}}$, respectively, we see that the sum in the bracket of RHS of the following (8.2) give meromorphic functions on z , whose principal parts of poles coincide with those of $x_{\mathbb{I}_2(p)}(z, \underline{g})$ and $y_{\mathbb{I}_2(p)}(z, \underline{g})$ in z , respectively (here, we denote by $[f(z)]$ the set of all principal parts of a meromorphic function

²³The notation “ ’ ” or “ ” ” shall mean single or twice derivative with respect to the variable z .

$f(z)$ defined on z -plane).

$$\begin{aligned}
 [x_{A_2}(z, \underline{g})] &= [\frac{1}{4}\mathbf{p}(z)] \\
 [y_{A_2}(z, \underline{g})] &= [\frac{1}{8}\mathbf{p}'(z)] \\
 (8.2) \quad [x_{B_2}(z, \underline{g})] &= [\frac{1}{2}\zeta(z) - \frac{1}{2}\zeta(z - \frac{1}{2}\omega_{\gamma_0})] \\
 [y_{B_2}(z, \underline{g})] &= [-\frac{1}{4}\mathbf{p}(z) + \frac{1}{4}\mathbf{p}(z - \frac{1}{2}\omega_{\gamma_0})] \\
 [x_{G_2}(z, \underline{g})] &= [\frac{1}{2}\zeta(z) - \frac{1}{2}\zeta(z - \frac{1}{3}\omega_{\gamma_0})] \\
 [y_{G_2}(z, \underline{g})] &= [-\frac{1}{2}\zeta(z) - \frac{1}{2}\zeta(z - \frac{1}{3}\omega_{\gamma_0}) + \zeta(z - \frac{2}{3}\omega_{\gamma_0})]
 \end{aligned}$$

On the other hand, we remark that the functions in the bracket of the RHS of (8.2) are periodic functions w.r.t. the period lattice $\Omega_{I_2(p), \underline{g}}$, since (1) the functions $\mathbf{p}(z)$ and $\mathbf{p}'(z)$ are already periodic, and (2) the sum of coefficients of the linear combinations of the functions of the form $\zeta(z + *)$ in each formula is equal to zero and, then, it is well known that the linear combination is a periodic function (see, e.g. [13, Ch. 1§12], this follows from an elementary property of zeta function that $\zeta(z + m\omega_0 + n\omega_1) - \zeta(z) = m2\zeta(\omega_0/2) + n2\zeta(\omega_1/2)$ for $\omega \in \Omega$). Thus, due to Liouville's Theorem, the difference of meromorphic functions in the brackets of both hand sides of (8.2) are constants.

Actually, the data of the principal parts of poles are not sufficient to control the ambiguity of adding constant terms except for the case of type A_2 . Namely, the Laurent expansion for the type A_2 at $z = 0$ (see Table 2 at the end of §7) does not have constant terms but those for the other types B_2 and G_2 contain non-trivial constant terms, which are linear in g_s and which are still to be determined from the data of the lattice $\Omega_{I_2(p)}$. In order to overcome this issue, we use the Hamilton equation (6.3) of the motion. This is essentially the new feature for the types B_2 and G_2 to be cautious compared with the classical case of type A_2 .

Let us determine the constants depending on the type separately.

A_2 type case: Since the constant terms of the Laurent expansions at the origin of BHS are zero, the difference is zero, and we obtain already (8.1) for type A_2 .

B_2 type case: Set, for suitable constants (w.r.t. z) A and B ,

$$\begin{aligned}
 x_{B_2}(z, \underline{g}) &= A + \frac{1}{2}\zeta(z) - \frac{1}{2}\zeta(z - \frac{1}{2}\omega_{\gamma_0}) \\
 y_{B_2}(z, \underline{g}) &= B - \frac{1}{4}\mathbf{p}(z) + \frac{1}{4}\mathbf{p}(z - \frac{1}{2}\omega_{\gamma_0})
 \end{aligned}$$

In the first equality, since the constant terms of the Laurent expansions of $x_{B_2}(z, \underline{g})$ and $\zeta(z)$ are zero (recall (7.4) and the fact that $\zeta(z)$ is an odd function) the sum of the remaining terms $A - \zeta(0 - \frac{1}{2}\omega_{\gamma_0})$ is equal to zero. This determine $A = -\zeta(\frac{1}{2}\omega_{\gamma_0})$.

For the second row of the equality, recall the Hamilton's equation of the motion (6.3) $\frac{\partial x_{B_2}}{\partial z} = 2y_{B_2}$. The LHS is equal to $-\frac{1}{2}\mathfrak{p}(z) + \frac{1}{2}\mathfrak{p}(z - \frac{1}{2}\omega_{\gamma_0})$, and substituting y_{B_2} in the RHS, we see that $2B = 0$. These already gives (8.1) for type B_2 . Then, by comparing the constant terms of the Laurent expansion of the BHS of (8.1) in view of (7.4), we obtain $\frac{1}{6}g_s = \frac{1}{4}\mathfrak{p}(0 - \frac{1}{2}\omega_{\gamma_0})$. Hence,

$$(8.3) \quad \begin{aligned} A &= -\zeta(\frac{1}{2}\omega_{\gamma_0}) \\ B &= 0 \\ g_s &= \frac{3}{2} \mathfrak{p}(\frac{1}{2}\omega_{\gamma_0}). \end{aligned}$$

G₂ type case: Set, for suitable constants w.r.t. z , A and B ,

$$\begin{aligned} x_{G_2}(z, \underline{g}) &= A + \frac{1}{2}\zeta(z) - \frac{1}{2}\zeta(z - \frac{1}{3}\omega_{\gamma_0}) \\ y_{G_2}(z, \underline{g}) &= B - \frac{1}{2}\zeta(z) - \frac{1}{2}\zeta(z - \frac{1}{3}\omega_{\gamma_0}) + \zeta(z - \frac{2}{3}\omega_{\gamma_0}). \end{aligned}$$

Recall the Hamilton's equation of the motion (6.3)

$$\frac{\partial x_{G_2}}{\partial z} = 2x_{G_2}y_{G_2} + 2g_s y_{G_2}.$$

The LHS is equal to $-\frac{1}{2}\mathfrak{p}(z) + \frac{1}{2}\mathfrak{p}(z - \frac{1}{3}\omega_{\gamma_0})$ so the residue at any pole of LHS is equal to zero. Thus, we obtain two relations that the residues at $z = \frac{2}{3}\omega_{\gamma_0}$ and at $z = \frac{1}{3}\omega_{\gamma_0}$ of the meromorphic function

$$\begin{aligned} &2(A + \frac{1}{2}\zeta(z) - \frac{1}{2}\zeta(z - \frac{1}{3}\omega_{\gamma_0})) \times (B - \frac{1}{2}\zeta(z) - \frac{1}{2}\zeta(z - \frac{1}{3}\omega_{\gamma_0}) \\ &+ \zeta(z - \frac{2}{3}\omega_{\gamma_0})) + 2g_s(B - \frac{1}{2}\zeta(z) - \frac{1}{2}\zeta(z - \frac{1}{3}\omega_{\gamma_0}) + \zeta(z - \frac{2}{3}\omega_{\gamma_0})) \end{aligned}$$

are zero.

- (1) Residue at $z = \frac{2}{3}\omega_{\gamma_0}$: $2A + \zeta(\frac{2}{3}\omega_{\gamma_0}) - \zeta(\frac{2}{3}\omega_{\gamma_0} - \frac{1}{3}\omega_{\gamma_0}) + 2g_s = 0$
- (2) Residue at $z = \frac{1}{3}\omega_{\gamma_0}$: $-A - \frac{1}{2}\zeta(\frac{1}{3}\omega_{\gamma_0}) - B + \frac{1}{2}\zeta(\frac{1}{3}\omega_{\gamma_0}) - \zeta(\frac{1}{3}\omega_{\gamma_0} - \frac{2}{3}\omega_{\gamma_0}) - g_s = 0$

In addition to these, let us consider two more relations:

- (3) $\frac{1}{2}g_s = A - \frac{1}{2}\zeta(0 - \frac{1}{3}\omega_{\gamma_0})$
- (4) $\frac{3}{2}g_s = B - \frac{1}{2}\zeta(0 - \frac{1}{3}\omega_{\gamma_0}) + \zeta(0 - \frac{2}{3}\omega_{\gamma_0})$

obtained by comparing the constant terms of Laurent expansion of the equalities: $x_{G_2} = A + \frac{1}{2}\zeta(z) - \frac{1}{2}\zeta(z - \frac{1}{3}\omega_{\gamma_0})$ and $y_{G_2} = B - \frac{1}{2}\zeta(z) - \frac{1}{2}\zeta(z - \frac{1}{3}\omega_{\gamma_0}) + \zeta(z - \frac{2}{3}\omega_{\gamma_0})$.

Recalling the fact that ζ is an odd function, we see that (1), (2), (3) and (4) are overdetermined system for A, B and g_s , and we obtain the solution:

$$\begin{aligned}
 (8.4) \quad A &= -\frac{1}{6}\zeta(\frac{1}{3}\omega_{\gamma_0}) - \frac{1}{6}\zeta(\frac{2}{3}\omega_{\gamma_0}) \\
 B &= \frac{1}{2}\zeta(\frac{1}{3}\omega_{\gamma_0}) + \frac{1}{2}\zeta(\frac{2}{3}\omega_{\gamma_0}) \\
 g_s &= \frac{2}{3}\zeta(\frac{1}{3}\omega_{\gamma_0}) - \frac{1}{3}\zeta(\frac{2}{3}\omega_{\gamma_0})
 \end{aligned}$$

This completes the proof of Theorem 8.1. □

Remark 8.2. In order to get the equality (8.1), we have substituted the lattice Ω in the RHS by the period lattice $\Omega_{I_2(p),g}$. However, the expression in the RHS of (8.1) is defined in a self-contained manner for any point (ω_0, ω_1) in $\tilde{\mathbb{H}}$. Therefore, we shall hereafter regard RHS of (8.1) as meromorphic functions in z which are holomorphically parametrized by $\tilde{\mathbb{H}}$, where the holomorphicity follows from the compact uniform convergences of the series \mathfrak{p} and ζ also in the variable $(\omega_0, \omega_1) \in \tilde{\mathbb{H}}$, regardless whether it is in the image of the period map or not.

9. Eisenstein series of type A_2, B_2 and G_2 (primitive automorphic forms)

We come back to the solve the inversion problem posed at Theorem 5.1. For the purpose, we use some generalizations of Eisenstein series to obtain inversion maps (see Theorem 9.2). For type A_2 , this is classically well established theory. Our interest is to show that a generalization of the theory works for types B_2 and G_2 (which is the first main goal of the present paper).

Definition 9.1. For each type $I_2(p)$, the coefficients of the Laurent series expansion at $z = 0$ of the meromorphic functions in RHS of (8.1), as a weighted homogeneous holomorphic functions on $(\omega_0, \omega_1) \in \tilde{\mathbb{H}}$, shall be called Eisenstein series of type $I_2(p)$.

In the following, we determine explicitly all Eisenstein series of type $I_2(p)$. However, such explicit description is un-necessary to solve the inversion problem. So, some readers may skip the present paragraph till Theorem 9.2. However, the explicit description are unavoidably important in §10, when we study the Fourier expansions of the polynomials in $\mathbb{C}[g_s, g_t]$ as modular forms.

Set $\Omega := \mathbb{Z}\omega_0 + \mathbb{Z}\omega_1$ for $(\omega_0, \omega_1) \in \widetilde{\mathbb{H}}$. Depending on $m \in \mathbb{Z}_{\geq 3}$ and $a \in \mathbb{R}\omega_0 + \mathbb{R}\omega_1 = \mathbb{R} \otimes_{\mathbb{Z}} \Omega$, let us consider series:

$$(9.1) \quad G_m(a) := \begin{cases} \sum_{\omega \in \Omega \setminus \{0\}} \omega^{-m} & = \frac{1}{(m-1)!} \frac{d^{m-2}(\mathbf{p}-z^{-2})}{dz^{m-2}}(0) \\ & = -\frac{1}{(m-1)!} \frac{d^{m-1}(\zeta-z^{-1})}{dz^{m-1}}(0) \quad (\text{if } a \in \Omega) \\ \sum_{\omega \in \Omega} (\omega + a)^{-m} & = \frac{1}{(m-1)!} \frac{d^{m-2}\mathbf{p}}{dz^{m-2}}(-a) \\ & = -\frac{1}{(m-1)!} \frac{d^{m-1}\zeta}{dz^{m-1}}(-a) \quad (\text{if } a \notin \Omega). \end{cases}$$

The first series for $a \in \Omega$ are the classical well-known classical *Eisenstein series of weight m* (see, e.g. [H-C,E-Z]). However, the second series for $a \notin \Omega$ seem to be less explicitly known in literature. As we shall see, since both behaves in parallel to the classical Eisenstein series, we shall call the latter case *shifted classical Eisenstein series of weight m* .

It is absolute and locally uniformly convergent so that defines a holomorphic function on $\widetilde{\mathbb{H}}$ of weight $-m \cdot \text{wt}(z)$ ²⁴ parametrized by $a \in (\mathbb{R}\omega_0 + \mathbb{R}\omega_1)/\Omega$, such that $G_m(a) = (-1)^m G_m(-a)$. In particular, we have the relations:

$$G_m(a) = 0 \quad \text{for } a \in \frac{1}{2}\Omega \text{ and } m = \text{odd}.$$

Using (9.1), one get the following Laurent and Tayler expansions. The first two lines are standard (e.g. [13]), and the latter two for $a \in \mathbb{R} \otimes \Omega \setminus \Omega$ can be shown similarly.²⁵

$$(9.2) \quad \begin{aligned} \mathbf{p}(z) &= z^{-2} + \sum_{n=1}^{\infty} (2n+1)z^{2n}G_{2n+2}(0) \\ \zeta(z) &= z^{-1} - \sum_{n=1}^{\infty} z^{2n+1}G_{2n+2}(0) \\ \mathbf{p}(z-a) &= \mathbf{p}(a) + \sum_{m=1}^{\infty} (m+1)z^m G_{m+2}(a) \\ \zeta(z-a) &= -\zeta(a) - \mathbf{p}(a)z - \sum_{m=1}^{\infty} z^{m+1}G_{m+2}(a) \end{aligned}$$

Now, let us describe Eisenstein series for each type $I_2(p)$ separately. The calculation is straight forward from the formula (8.1), and we omit details of them.

²⁴We should be cautious about the use of the terminology “weight”. The weight $-m \cdot \text{wt}(z)$ of $G_m(a)$ as a function on $\widetilde{\mathbb{H}}$ comes from the \mathbb{C}^\times -action (recall Fact 14). It is proportional to the weight m as the Eisenstein series, but depends on the factor $\text{wt}(z)$ which depends on type $I_2(p)$ (recall Table 1) (cf. (10.3) and Table 5).

²⁵Actually, the last equality has meaning for the periodic variable a , even though the zeta function is not periodic, since ζ is still “semi-periodic” (see [13, §11]).

A_2 type: Set

$$(9.3) \quad \begin{aligned} x_{A_2}(z) &= \frac{1}{4}z^{-2} + \sum_{n=1}^{\infty} A_n z^{2n} \\ y_{A_2}(z) &= -\frac{1}{4}z^{-3} + \sum_{n=1}^{\infty} B_n z^{2n-1} \end{aligned}$$

Then, we have

$$(9.4) \quad \begin{aligned} A_n &= \frac{2n+1}{4}G_{2n+2}(0) \quad (n \geq 1) \\ B_n &= \frac{(2n+1)n}{4}G_{2n+2}(0) \quad (n \geq 1) \end{aligned}$$

B_2 type: Set

$$(9.5) \quad \begin{aligned} x_{B_2}(z) &= \frac{1}{2}z^{-1} + \sum_{n=0}^{\infty} A_n z^{2n+1} \\ y_{B_2}(z) &= -\frac{1}{4}z^{-2} + \sum_{n=0}^{\infty} B_n z^{2n} \end{aligned}$$

Then, we have

$$(9.6) \quad \begin{aligned} A_0 &= \frac{1}{2}\mathfrak{p}\left(\frac{1}{2}\omega_0\right) \\ A_n &= -\frac{1}{2}G_{2n+2}(0) + \frac{1}{2}G_{2n+2}\left(\frac{1}{2}\omega_0\right) \quad (n \geq 1) \\ B_0 &= \frac{1}{4}\mathfrak{p}\left(\frac{1}{2}\omega_0\right) \\ B_n &= -\frac{2n+1}{4}G_{2n+2}(0) + \frac{2n+1}{4}G_{2n+2}\left(\frac{1}{2}\omega_0\right) \quad (n \geq 1) \end{aligned}$$

G_2 type: Set

$$(9.7) \quad \begin{aligned} x_{G_2}(z) &= \frac{1}{2}z^{-1} + \sum_{n=0}^{\infty} A_n z^n \\ y_{G_2}(z) &= -\frac{1}{2}z^{-1} + \sum_{n=0}^{\infty} B_n z^n \end{aligned}$$

Then, we have

$$(9.8) \quad \begin{aligned} A_0 &= \frac{1}{3}\zeta\left(\frac{1}{3}\omega_{\gamma_0}\right) - \frac{1}{6}\zeta\left(\frac{2}{3}\omega_{\gamma_0}\right) = \frac{1}{2}\zeta\left(\frac{1}{3}\omega_{\gamma_0}\right) - \frac{1}{3}\zeta\left(\frac{1}{2}\omega_{\gamma_0}\right) \\ A_1 &= \frac{1}{2}\mathfrak{p}\left(\frac{1}{3}\omega_0\right) \\ A_n &= -\frac{1}{2}G_{n+1}(0) + \frac{1}{2}G_{n+1}\left(\frac{1}{3}\omega_0\right) \quad (n \geq 2) \\ B_0 &= \zeta\left(\frac{1}{3}\omega_{\gamma_0}\right) - \frac{1}{2}\zeta\left(\frac{2}{3}\omega_{\gamma_0}\right) = \frac{3}{2}\zeta\left(\frac{1}{3}\omega_{\gamma_0}\right) - \zeta\left(\frac{1}{2}\omega_{\gamma_0}\right) \\ B_1 &= \frac{1}{2}\mathfrak{p}\left(\frac{1}{3}\omega_0\right) - \mathfrak{p}\left(\frac{2}{3}\omega_0\right) = -\frac{1}{2}\mathfrak{p}\left(\frac{1}{3}\omega_0\right) \\ B_n &= \frac{1}{2}G_{n+1}(0) + \frac{1}{2}G_{n+1}\left(\frac{1}{3}\omega_0\right) - G_{n+1}\left(\frac{2}{3}\omega_0\right) \\ &= \frac{1}{2}G_{n+1}(0) + \left(\frac{1}{2} + (-1)^n\right)G_{n+1}\left(\frac{1}{3}\omega_0\right) \quad (n \geq 2) \end{aligned}$$

Remark 2. 1. The infinite sequence of Eisenstein series for each type $I_2(p)$ are not algebraically independent. More precisely, they are obeying recurrence relation (7.5) in Step 4. of the proof of Lemma 7.1 (which leads to the isomorphism (9.9), describing relations directly). The relations may be considered as the B_2 -type and G_2 -type generalizations of the classically well-known A_2 -type relations. However, in the present paper, we do not go into details of the relations.

2. We note that the first Eisenstein series A_0, B_0 in case of type B_2 and the first and the second Eisenstein series A_0, B_0 and A_1, B_1 in case of type G_2 do not have the description using the classical series (9.1). These exceptional behavior was caused by the fact that the classical Eisenstein series (9.1) do not converge absolutely in those low weights so that one need to make conditional convergent series by a help of \mathfrak{p} -function or ζ -function. This was made possible by the determination of the constant terms of fractional expansions in Theorem 8.1 using the energy condition (6.4) of the Hamilton’s equations of the motion.

In a forthcoming paper [2], we shall study systematically those “exceptional” Eisenstein series from a view point of modular forms.

Now we are to formulate the second main theorem of the present paper. The proof is essentially done already in previous sections so that we have only to coordinate them.

Theorem 9.2. *Consider the pull-back homomorphism $P_{I_2(p)}^* : \mathcal{O}_{\tilde{\mathbb{H}}} \rightarrow \mathcal{O}_{\tilde{S}_{I_2(p)}}$ from the ring of holomorphic functions on the period domain $\tilde{\mathbb{H}}$ to that on the monodromy covering space $\tilde{S}_{I_2(p)}$ of the base space $S_{I_2(p)}$ of the family (2.3) (recall (4.12) for the definition of $\tilde{S}_{I_2(p)}$ and (5.4) for the definition of $P_{I_2(p)}$). Then, it induces the ring isomorphism:*

$$(9.9) \quad \mathbb{Q}[\text{Eisenstein series of type } I_2(p)] \simeq \mathbb{Q}[g_s, g_l],$$

where LHS is the ring over \mathbb{Q} generated by all Eisenstein series of type $I_2(p)$ (recall Definition 9.1) and RHS is the coordinate ring of the space $S_{I_2(p)}$ (recall (2.3)) generated by the flat coordinates g_s and g_l over \mathbb{Q} .

In particular, the generators g_s and g_l are expressed by Eisenstein series $E_{I_2(p),s}$ and $E_{I_2(p),l}$ of type $I_2(p)$ as follows

A_2 type:

$$(9.10) \quad \begin{aligned} g_s &= E_{A_2,s} := \frac{15}{4} G_4(0) \\ g_l &= E_{A_2,l} := \frac{35}{16} G_6(0) \end{aligned}$$

B_2 type:

$$(9.11) \quad \begin{aligned} g_s &= E_{B_2,s} := \frac{3}{2}\mathfrak{p}\left(\frac{1}{2}\omega_0\right) \\ g_l &= E_{B_2,l} := \frac{5}{32}\mathfrak{p}^2\left(\frac{1}{2}\omega_0\right) + \frac{5}{8}G_4(0) - \frac{5}{8}G_4\left(\frac{1}{2}\omega_0\right) \end{aligned}$$

G_2 type:

$$(9.12) \quad \begin{aligned} g_s &= E_{G_2,s} := \zeta\left(\frac{1}{3}\omega_0\right) - \frac{2}{3}\zeta\left(\frac{1}{2}\omega_0\right) \\ g_s^2 &= E_{G_2,s}^2 := \frac{1}{3}\mathfrak{p}\left(\frac{1}{3}\omega_0\right) \\ g_l &= E_{G_2,l} := 2g_s^3 - G_3\left(\frac{1}{3}\omega_0\right) \end{aligned}$$

Proof. The explicit description of the function on $\widetilde{S}_{I_2(p)}$ corresponding to an Eisenstein series by the pull-back morphism $P_{I_2(p)}^*$ is obtained by the corresponding coefficient of the Laurent expansion at $z = 0$ of the meromorphic functions $x_{I_2(p)}(z, \underline{g})$ or $y_{I_2(p)}(z, \underline{g})$ (8.1). Then, it was already shown in Lemma 7.1 that they are rational coefficient polynomials in g_s and g_l (cf. Table 2 at the end of §7). This defines the homomorphism (9.9) from left to right.

The morphism is injective, since the period map $P_{I_2(p)}$ (5.4) is an open map between connected manifolds (Fact 12).

The morphism is surjective, since (1) the generator g_s is, up to a constant factor, given by the lowest weight Eisenstein series for each type (recall Step 2. of the proof of Lemma 7.1 and (7.4)), and (2) the generator g_l appear non-trivially and linearly in the coefficients A_{n_0} and B_{n_0} of the Laurent expansions of $x_{I_2(p)}(z, \underline{g})$ and $y_{I_2(p)}(z, \underline{g})$ (recall Step 3. of the proof of Lemma 7.1 and (7.4)). □

After the isomorphism (9.9), we shall sometimes identify the ring of Eisenstein series and the polynomial ring in g_s and g_l .

Proof of Theorem 5.1. We show by 4 steps that the period map $P_{I_2(p)}$ (5.6) is bi-holomorphic.

Step 1. Regardless, whether an element $(\omega_0, \omega_1) \in \widetilde{\mathbb{H}}$ belongs to the image of the period map or not, let us use the Eisenstein series of type $I_2(p)$ expressions (9.10), (9.11) and (9.12) to define a holomorphic map

$$(9.13) \quad E = (E_{I_2(p),s}, E_{I_2(p),l}) : \widetilde{\mathbb{H}} \longrightarrow S_{I_2(p)}.$$

The equality (8.1) in Theorem 8.1 implies that the following diagram is commutative

$$(9.14) \quad \begin{array}{ccc} \tilde{S}_{I_2(p)} & \xrightarrow{P_{I_2(p)}} & \tilde{\mathbb{H}} \\ & \searrow & \swarrow E \\ & S_{I_2(p)} & \end{array}$$

We remark that pull back of the polynomial ring on $S_{I_2(p)}$ by the morphism E (9.13) induces the same isomorphism (9.9), since (i) the period map $P_{I_2(p)}$ is a non-trivial open map, and (ii) any algebraic dependence relation among Eisenstein series on an open domain in $\tilde{\mathbb{H}}$ automatically extends on the whole $\tilde{\mathbb{H}}$ by analytic continuation, since $\tilde{\mathbb{H}}$ is *connected*.

Step 2. Let us show that the image of E is contained in the compliment of the discriminant: $E(\tilde{\mathbb{H}}) \subset S_{I_2(p)} \setminus D_{I_2(p)}$. For $(\omega_1, \omega_2) \in \tilde{\mathbb{H}}$, using RHS of (8.1), we define global meromorphic functions $x_{I_2(p)}$ and $y_{I_2(p)}$, which are periodic w.r.t. the lattice $\Omega = \mathbb{Z}\omega_0 + \mathbb{Z}\omega_1$. Let us see that the pair satisfies the relation (6.3) together with (6.4), where the parameter \underline{g} is given by (9.13). Actually, the both hand sides give doubly periodic function of the period $\Omega = \mathbb{Z}\omega_0 + \mathbb{Z}\omega_1$, where we can check they have the same principal parts of poles, and the constant term of Laurent expansions at 0 coincides.

This means that the time coordinate z is given by the integral (6.1) (up to a shift of a constant). That is, the image of the map $(x_{I_2(p)}, y_{I_2(p)})$ satisfies the equation (6.4). However, if \underline{g} belonged to the discriminant, then the associated curve defined by the equation (2.1) is a singular rational curve. The integral (6.1) (avoiding the singularity of the curve but admitting to go through points at infinity) cannot be doubly periodic (either one periodic for $\underline{g} \in D_{I_2(p)} \setminus \{0\}$, or no-periodic for $\underline{g} = 0$), where as the starting $(\omega_0, \omega_1) \in \tilde{\mathbb{H}}$ generates rank 2 lattices and $x_{I_2(p)}$ and $y_{I_2(p)}$ are doubly periodic. A contradiction!

Step 3. Let us show that the period map is surjective. Since $\tilde{\mathbb{H}}$ is connected, for any point $\omega \in \tilde{\mathbb{H}}$, consider any path, say p , in $\tilde{\mathbb{H}}$ connecting ω with the image $P_{I_2(p)}(\tilde{\Gamma}_{I_2(p)})$ in $\tilde{\mathbb{H}}$ of the base point loci of $\tilde{S}_{I_2(p)}$ (recall the definition (4.12)). Then, the projection image $E(p)$ is a path in $S_{I_2(p)} \setminus D_{I_2(p)}$ connecting the base point loci $\Gamma_{I_2(p)}$ with $E(\omega)$ (recall the commutative diagram (9.14)). Then, the monodromy lifting $\widetilde{E(p)}$ of the path $E(p)$ in the covering space $\tilde{S}_{I_2(p)}$ is a path connecting the base point loci $\tilde{\Gamma}_{I_2(p)}$ to a point $\widetilde{E(\omega)}$ which lies over $E(\omega)$. Then, the image $P_{I_2(p)}(\widetilde{E(p)})$ is a monodromy covering in $\tilde{\mathbb{H}}$ of the

path $E(p)$ connecting $P_{I_2(p)}(\tilde{\Gamma}_{I_2(p)})$ to a point $P_{I_2(p)}(\widetilde{E(\omega)})$. Since p is also a monodromy covering in \mathbb{H} of the same path $E(p)$ connecting the base point loci $P_{I_2(p)}(\tilde{\Gamma}_{I_2(p)})$ to the point p . So the two end points $P_{I_2(p)}(\widetilde{E(\omega)})$ and p of the paths should coincide each other. In particular, p is in the image of the period map. This shows also that the modular group $\Gamma_1([p/2])$, which is the monodromy representation ρ -image of the fundamental group of $S_{I_2(p)} \setminus D_{I_2(p)}$ (Fact 9, 2.), acts on any fiber of the map E transitively. That is, the modular group action quotient of \mathbb{H} is isomorphic to the discriminant compliment:

$$\Gamma_1([p/2]) \setminus \tilde{\mathbb{H}} \simeq S_{I_2(p)} \setminus D_{I_2(p)}$$

Step 4. Finally, let us show that the period map is injective. Since the period map is equivariant with the modular group action, it is sufficient that the modular group action on the period domain $\tilde{\mathbb{H}}$ is (generically) fixed point free. But this is trivially true, since the modular group is a subgroup of $GL_2(\mathbb{Z})$ so that its fixed points set is thin and $\tilde{\mathbb{H}}$ is an open subset of \mathbb{C}^2 . Since E is a covering map, if the action modular group is fixed point free in one fiber, it is fixed point free for all fibers.

This completes a proof of Theorem 5.1. □

Remark 9.3. 1.

$$(9.15) \quad \frac{\partial(g_s, g_t)}{\partial(\omega_0, \omega_1)} = c \Delta_{I_2(p)}^{red}$$

2. In [21, 23], we posed a general question to describe the inversion morphism to the period map defined by a primitive form. If a function on the parameter space of the family is described in terms of the coordinates of the period domain, we call the function (and its description on the period domain) a *primitive automorphic form*. In that sense, the generalized Eisenstein series of type B_2 and G_2 in this sections are the first examples of primitive automorphic forms beyond the classical case of type A_2 .

10. Ring of modular forms and discriminant

We identify the ring of Eisenstein series of type $I_2(p)$ with the ring of modular forms of the congruence group $\Gamma_1([p/2])$ (see [1] for $M_*(\Gamma_1([p/2]))$). Then we confirm that the set of irreducible components of the discriminant of the family (2.3) is in one to one correspondence with the set of cusps of the congruence group $\Gamma_1([p/2])$.

Theorem 10.1. *The ring of Eisenstein series of type $I_2(p)$ is identified with the ring of holomorphic modular forms of the congruence group $\Gamma_1([p/2])$, where the identification is given in following (10.3).²⁶*

$$(10.1) \quad \mathbb{C}[\text{Eisenstein series of type } I_2(p)] \simeq M_*(\Gamma_1([p/2])).$$

The correspondences of generators are given in (10.5), (10.6) and (10.7).

Proof. Proof of the theorem is divided into Steps 1–5.

Step 1. We explain the meaning of “identification”, and fix notation.

Recall the period domain $\tilde{\mathbb{H}} := \{(\omega_0, \omega_1) \in \mathbb{C}^2 \mid \text{Im}(\omega_1/\omega_0) > 0\}$ with its homogenous coordinates (ω_0, ω_1) . We introduce the inhomogeneous coordinate

$$(10.2) \quad \tau := \omega_1/\omega_0$$

Then the natural projection $\tilde{\mathbb{H}} \rightarrow \mathbb{H} := \{\tau \in \mathbb{C} \mid \text{Im}(\tau) > 0\}$, $(\omega_0, \omega_1) \mapsto \tau := \omega_1/\omega_0$ gives a principal \mathbb{C}^\times -bundle, say $(L^\times)^{-1}$, over \mathbb{H} , which we trivialize by the morphism $\tilde{\mathbb{H}} \simeq \mathbb{C}^\times \mathbb{H}$, $(\omega_0, \omega_1) \mapsto (\omega_0, \tau)$. The modular group $\Gamma_1([p/2])$ acts from the left on $\tilde{\mathbb{H}}$ and hence on L . For $k \in \mathbb{Z}_{>0}$, a holomorphic section of the L^k , say $s = s(\tau)$, such that $\gamma^*(s) := s \cdot \frac{a\tau+b}{c\tau+d}$ is equal to $(c\tau + d)^k s(\tau)$

for $\gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \Gamma_1([p/2])$, is called a modular form of weight k of $\Gamma_1([p/2])$ in a wide sense. Then, the correspondence $s(\tau) \mapsto \omega_0^{-k} \cdot s(\tau)$ defines the “identification”:

$$(10.3) \quad \begin{aligned} & \{\text{modular forms of weight } k \text{ of the group } \Gamma_1([p/2]) \text{ in a wide sense}\} \\ \leftrightarrow & \{\text{holomorphic functions on } \tilde{\mathbb{H}} \text{ of weight } -k \cdot \text{wt}(z) \text{ invariant by } \Gamma_1([p/2])\} \end{aligned}$$

Actually, we study more restricted class of modular forms which are holomorphic and taking finite (bounded) values at cusps, as we explain now.

Recall [15] that a point $x \in \mathbb{R} \cup \{\sqrt{-1}\infty\}$ ($= \partial\mathbb{H}$) is called a *cusps* of $\Gamma_1(N)$ if it is fixed by a hyperbolic element of $\Gamma_1(N)$. The isotropic subgroup of $\Gamma_1(N)$ fixing a cusp is an infinite unipotent group, and the set of

²⁶This is naturally an expected result. However, this should have been proven, since the Eisenstein series appeared in the context of the geometry of the period mapping, whereas the modular forms are defined independently by themselves. So, their coincidence is a non-trivial marvelous fact, which we need to work cautiously.

all cusps are invariant under the action of $\Gamma_1(N)$. For $\gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \text{SL}_2(\mathbb{Z})$, set $(s|_k\gamma)(\tau) := (\gamma^*s)(\tau)(c\tau + d)^{-k}$ (so that the modularity property of s is equivalent to $s|_k\gamma = s$ for all $\gamma \in \Gamma_1(N)$). Let $\gamma(x) = \sqrt{-1}\infty$ for a cusp x and $\gamma \in \text{SL}_2(\mathbb{Z})$. For a modular form s (in the wide sense), $s|_k\gamma$, as a periodic function in τ , develops into a Fourier series in τ . Then, s is called a *holomorphic* modular form if the Fourier series consists only of non-negative powers of $q = \exp(2\pi\sqrt{-1}\tau)$ at all cusps of $\Gamma_1(N)$. The constant term of the Fourier series is called the *value* of s at the cusp and denoted by $(s|_k\gamma)(\sqrt{-1}\infty)$.

Step 2. We recall the results by Aoki and Ibukiyama on the ring of modular forms. In [1], Aoki and Ibukiyama gave a simple unified description of the graded ring of holomorphic modular forms of $\Gamma_0(N)$ for $N = 1, 2, 3, 4$. From that description, we recover easily the ring $M_*(\Gamma_1(N))$ of holomorphic modular forms of $\Gamma_1(N)$. Namely, according as $N = 1, 2$ and 3 , the ring is generated by two (algebraically independent) modular forms e_4, e_6 of weight 4 and 6, α_2, β_4 of weight 2 and 4, and α_1, β_3 of weight 1 and 3, respectively. That is,

$$(10.4) \quad M_*(\Gamma_1(1)) = \mathbb{C}[e_4, e_6], \quad M_*(\Gamma_1(2)) = \mathbb{C}[\alpha_2, \beta_4], \quad M_*(\Gamma_1(3)) = \mathbb{C}[\alpha_1, \beta_3]$$

where explicit descriptions of the generators as theta-function, in particular, the first few Fourier coefficients at the cusps are given in [1].

Step 3. We define a morphism from the left hand side to the right hand side of (10.1). This is achieved by showing that the Eisenstein series are holomorphic at all cusps.

More explicitly, recall that, according as $N = 1, 2$ and 3 , the number of $\Gamma_1(N)$ -equivalence classes of cusps are 1, 2 and 2, whose representatives are given as follows.

$$A_2 \text{ type: } \sqrt{-1}\infty, \quad B_2 \text{ type: } \sqrt{-1}\infty \text{ and } 0, \quad G_2 \text{ type: } \sqrt{-1}\infty \text{ and } 0.$$

Let us consider the ring of Eisenstein series of type $I_2(p)$. Recall that the identification (9.9) was given by the composition with the period map $P_{I_2(p)}$, where the period map is $\Gamma_1([p/2])$ -equivariant. This means that any Eisenstein series is a $\Gamma_1([p/2])$ -invariant function on $\tilde{\mathbb{H}}$. So, by the identification (10.3), it gives arise a modular form. If we, further, show that the Eisenstein series are holomorphic at all cusps, we obtain a graded ring homomorphism from the left hand side to the right hand side of (10.1).

Lemma 10.2. *The modular forms associated with Eisenstein series of type $I_2(p)$ ($p = 1, 2, 3$) are holomorphic at their cusp(s).*

Proof. Let $E = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ and $S = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \in \text{SL}_2(\mathbb{Z})$, which transforms $\sqrt{-1}\infty$ and 0 to $\sqrt{-1}\infty$. We show that the series associated with $s|_k E$ and/or $s|_k S$ for an Eisenstein series s with the weight k converges absolute uniformly in a “neighborhood” of $\sqrt{-1}\infty$. This is classical for the case of Eisenstein series $s = G_m(0)$ for $m \geq 3$ (see [11]), and similar proof works for Eisenstein series of the form $s = G_m(a)$ for a suitable $a \in \Omega_{\mathbb{Q}}$. In case of Eisenstein series of the form $s = \mathfrak{p}(a)$ for a suitable $a \in \Omega_{\mathbb{Q}}$ in type B_2 and G_2 , we may either show directly the convergence in [2], or alternatively, we use the expression of the \mathfrak{p} -function as a proportion of Jacobi forms [10, Theorem 3.6] to show that the Fourier expansions at infinity consists only of positive powers. In case of $s = \zeta(\frac{1}{3}\omega_0) - \frac{2}{3}\zeta(\frac{1}{2}\omega_0)$, it will be shown in [2]. \square

Step 4. In the following table, we give the values at cusps of the additive summands in (9.10), (9.11) and (9.12). Calculations depend on cases: In case of the form $G_m(0)$, it is classical (e.g. [7, 11]). In case of the form $\mathfrak{p}(a)$ for some $a \in \Omega_{\mathbb{Q}}$ (see [2]). The cases of shifted series $G_4(\frac{1}{2}\omega_0)$ and $G_3(\frac{1}{3}\omega_0)$ at the cusp $\sqrt{-1}\infty$ are reduced to Riemann’s zeta function $\zeta_R(4)$ or Dirichlet’s L-function $L(3, \chi)$ of the character χ given by quadratic residues, respectively (see [3] for explicit values). Their values at the cusp 0 are directly shown to be zero. In case of $\zeta(\frac{1}{3}\omega_0) - \frac{2}{3}\zeta(\frac{1}{2}\omega_0)$, it will be shown in [2].

Table 3: Values of Eisenstein series of type $I_2(p)$ at cusps, I

A₂ type:

$$(\omega_0^4 \cdot G_4(0)|_4 E)(\sqrt{-1}\infty) = \frac{\pi^4}{45} \quad (\omega_0^6 \cdot G_6(0)|_6 E)(\sqrt{-1}\infty) = \frac{2\pi^6}{945}$$

B₂ type:

$$(\omega_0^2 \cdot \mathfrak{p}(\frac{1}{2}\omega_0)|_2 E)(\sqrt{-1}\infty) = \frac{2\pi^2}{3} \quad (\omega_0^2 \cdot \mathfrak{p}(\frac{1}{2}\omega_0)|_2 S)(\sqrt{-1}\infty) = -\frac{\pi^2}{3}$$

$$(\omega_0^4 \cdot \mathfrak{p}^2(\frac{1}{2}\omega_0)|_4 E)(\sqrt{-1}\infty) = \frac{4\pi^4}{9} \quad (\omega_0^4 \cdot \mathfrak{p}^2(\frac{1}{2}\omega_0)|_4 S)(\sqrt{-1}\infty) = \frac{\pi^4}{9}$$

$$(\omega_0^4 \cdot G_4(0)|_4 E)(\sqrt{-1}\infty) = \frac{\pi^4}{45} \quad (\omega_0^4 \cdot G_4(0)|_4 S)(\sqrt{-1}\infty) = \frac{\pi^4}{45}$$

$$(\omega_0^4 \cdot G_4(\frac{1}{2}\omega_0)|_4 E)(\sqrt{-1}\infty) = \frac{\pi^4}{3} \quad (\omega_0^4 \cdot G_4(\frac{1}{2}\omega_0)|_4 S)(\sqrt{-1}\infty) = 0$$

G₂ type:

$$(\omega_0 \cdot (\zeta(\frac{1}{3}\omega_0) - \frac{2}{3}\zeta(\frac{1}{2}\omega_0))|_1 E)(\sqrt{-1}\infty) = \frac{\pi}{\sqrt{3}} \quad (\omega_0 \cdot (\zeta(\frac{1}{3}\omega_0) - \frac{2}{3}\zeta(\frac{1}{2}\omega_0))|_1 S)(\sqrt{-1}\infty) = -\sqrt{-1}\frac{\pi}{3}$$

$$(\omega_0^2 \cdot \mathfrak{p}(\frac{1}{3}\omega_0)|_2 E)(\sqrt{-1}\infty) = \pi^2 \quad (\omega_0^2 \cdot \mathfrak{p}(\frac{1}{3}\omega_0)|_2 S)(\sqrt{-1}\infty) = -\frac{\pi^2}{3}$$

$$(\omega_0^3 \cdot G_3(\frac{1}{3}\omega_0)|_3 E)(\sqrt{-1}\infty) = \frac{2^2\pi^3}{3\sqrt{3}} \quad (\omega_0^3 \cdot G_3(\frac{1}{3}\omega_0)|_3 S)(\sqrt{-1}\infty) = 0$$

This completes a proof of Theorem 10.1. \square

Remark 3. In case of type A_2 , Fourier coefficients of Eisenstein series are well-known to be given by the divisor sum function $\sigma_p(n)$ with suitable constant factors given by special values of Riemann’s zeta function (e.g. [11, VII.1.3]). It is natural to ask for similar expressions of the Fourier coefficients of Eisenstein series for the types B_2 and G_2 .

Combining above calculations with the expressions (9.10), (9.11) and (9.12), we obtain the following table,

Table 4: Values of Eisenstein series of type $I_2(p)$ at cusps, II

A_2 type:

$$\begin{aligned} (\omega_0^4 \cdot g_s|_4E)(\sqrt{-1}\infty) &= \frac{1}{2^2 3} \pi^4, \\ (\omega_0^6 \cdot g_l|_6E)(\sqrt{-1}\infty) &= \frac{1}{2^3 3^3} \pi^6, \end{aligned}$$

B_2 type:

$$\begin{aligned} (\omega_0^2 \cdot g_s|_2E)(\sqrt{-1}\infty) &= \pi^2, & (\omega_0^2 \cdot g_s|_2S)(\sqrt{-1}\infty) &= -\frac{1}{2} \pi^2, \\ (\omega_0^4 \cdot g_l|_4E)(\sqrt{-1}\infty) &= -\frac{1}{2^3} \pi^4, & (\omega_0^4 \cdot g_l|_4S)(\sqrt{-1}\infty) &= \frac{1}{2^5} \pi^4, \end{aligned}$$

G_2 type:

$$\begin{aligned} (\omega_0 \cdot g_s|_1E)(\sqrt{-1}\infty) &= \frac{\pi}{\sqrt{3}}, & (\omega_0 \cdot g_s|_1S)(\sqrt{-1}\infty) &= -\frac{\sqrt{-1}}{3} \pi, \\ (\omega_0^3 \cdot g_l|_3E)(\sqrt{-1}\infty) &= \frac{2}{3\sqrt{3}} \pi^3, & (\omega_0^3 \cdot g_l|_3S)(\sqrt{-1}\infty) &= \frac{2\sqrt{-1}}{3^3} \pi^3. \end{aligned}$$

Step 5. This is the final step to obtain the isomorphism (10.1). We determine the linear relations between the generators of both hand sides by comparing their values at cusps.

First, we compare the weights (of the free generators of) the rings in both hand sides of (10.1) by the use of “weight factor” $\text{wt}(z)$ (recall (10.3)). Comparing Table 1 in §2 and the description of (10.4), we obtain the following “coincidences” of weights!

Table 5: Weights of the generators of the ring of modular forms

	$-\text{wt}(g_s)/\text{wt}(z)$,	$-\text{wt}(g_l)/\text{wt}(z)$	weights of $M_*(\Gamma_1([p/2]))$
A_2	$-(2/3)/(-1/6) = 4$,	$-1/(-1/6) = 6$	4, 6
B_2	$-(1/2)/(-1/4) = 2$,	$-1/(-1/4) = 4$	2, 4
G_2	$-(1/3)/(-1/3) = 1$,	$-1/(-1/3) = 3$	1, 3

This means that each generators g_s and g_l are mapped into the graded vector subspace of $M_*(\Gamma_1([p/2]))$ of the same degree as the corresponding generators α_i, β_j etc., respectively. For g_s , the dimension of the graded subspace containing it is equal to 1 so that we only need to fix the constant factor. In case of g_l , the dimension is either 1 for type A_2 or 2, spanned by g_l and a power of g_s , for types B_2 and G_2 .

Let us recall the values of the generators of the modular forms at cusps. The following values are taken from Aoki-Ibukiyama [1] Internat J. Math. 16-3(2005) 249–279, B_2 : p. 270 B_3 pp. 271–272.

Table 6: Values of Modular forms of $\Gamma_1([p/2])$ at cusps

$$\begin{aligned} (e_4|_2E)(\sqrt{-1}\infty) &= 1, \\ (e_6|_4E)(\sqrt{-1}\infty) &= 1, \\ (\alpha_2|_2E)(\sqrt{-1}\infty) &= 1, & (\alpha_2|_2S)(\sqrt{-1}\infty) &= -1/2, \\ (\beta_4|_4E)(\sqrt{-1}\infty) &= 0, & (\beta_4|_4S)(\sqrt{-1}\infty) &= 1/256, \\ (\alpha_1|_1E)(\sqrt{-1}\infty) &= 1, & (\alpha_1|_1S)(\sqrt{-1}\infty) &= -\sqrt{-1}/\sqrt{3}, \\ (\beta_3|_3E)(\sqrt{-1}\infty) &= 1, & (\beta_3|_3S)(\sqrt{-1}\infty) &= -\sqrt{-1}/3\sqrt{3}. \end{aligned}$$

Comparing Table 4 with Table 6, we obtain the following expressions of the isomorphism (10.1).

A₂ type:

$$(10.5) \quad \begin{aligned} g_s &= \frac{1}{2^2 3} \pi^4 e_4 \omega_0^{-4} \\ g_l &= \frac{1}{2^3 3^3} \pi^6 e_6 \omega_0^{-6} \end{aligned}$$

B₂ type:

$$(10.6) \quad \begin{aligned} g_s &= \pi^2 \alpha_2 \omega_0^{-2} \\ g_l &= 2^4 \pi^4 \beta_4 \omega_0^{-4} - \frac{1}{2^3} \pi^4 (\alpha_2 \omega_0^{-2})^2 \end{aligned}$$

G₂ type:

$$(10.7) \quad \begin{aligned} g_s &= \frac{1}{\sqrt{3}} \pi \alpha_1 \omega_0^{-1} \\ g_l &= -\frac{2}{3\sqrt{3}} \pi^3 \beta_3 \omega_0^{-3} \end{aligned}$$

Remark 4. As a consequence of the identification (10.1), we observe that the holomorphicity condition at cusps on modular forms are equivalent to the holomorphic extendability condition on functions on $S_{I_2(p)} \setminus D_{I_2(p)}$ to holomorphic functions on $S_{I_2(p)}$. Then the cusp form condition “should be” equivalent to the condition to vanish on the discriminant $D_{I_2(p)}$. This is the subject discussed in the following lemma.

Lemma 10.3. *The set of $\Gamma_1([p/2])$ -equivalence classes of cusp points corresponds naturally in one to one with the set of irreducible components of the discriminant loci $D_{I_2(p)}$ (2.4).*

We show that a generator of the ideal of the ring $M_*(\Gamma_1([p/2]))$ of modular forms vanishing at each equivalence class of cusp points, which can be easily

found from Table 4, is, by the pull back by the period map (5.6), up to a constant factor, identified with an irreducible component of the equation $\Delta_{I_2(p)} \in \mathbb{C}[g_s, g_t]$ (2.5) of the discriminant.

Here, we may recall Footnote 27 again.

A_2 type: Recall that there is a unique equivalence class of cusps which is represented by $\sqrt{-1}\infty$. The ideal vanishing at the class is generated by $e_4^3 - e_6^2$. We also recall the equation of the discriminant (2.5) of type A_2 . Then, the identification (10.5) induces the following identity:

$$(10.8) \quad -27g_t^2 + g_s^3 = \frac{\pi^{12}}{1728}(e_4^3 - e_6^2) \omega_0^{-12}$$

B_2 type: Recall that there are two equivalence classes of cusps, which are represented by $\sqrt{-1}\infty$ and by 0. The ideal vanishing at the class $\sqrt{-1}\infty$ is generated by β_4 , and the ideal vanishing at the class 0 is generated by $\alpha_2^2 - 64\beta_4$. We recall the irreducible factors of the equation of the discriminant (2.5) of type B_2 are $-8g_t + g_s^2$ and $8g_t + g_s^2$. Then, the identification (10.6) induces the following identities.

$$(10.9) \quad \begin{aligned} 8g_t + g_s^2 &= 128\pi^4\beta_4 \omega_0^{-4} \\ -8g_t + g_s^2 (= -G_2(\frac{1}{2}\omega_0)) &= 2\pi^4(\alpha_2^2 - 64\beta_4) \omega_0^{-4} \end{aligned}$$

G_2 type: Recall that there are two equivalence classes of cusps, which are represented by $\sqrt{-1}\infty$ and by 0. The ideal vanishing at the class $\sqrt{-1}\infty$ is generated by $\frac{1}{54}(\alpha_1^3 - \beta_3)$, and the ideal vanishing at the class 0 is generated by $\frac{1}{2}(\alpha_1^3 + \beta_3)$. We recall the irreducible factors of the equation of the discriminant (2.5) of type G_2 are $g_t + 2g_s^3$ and $g_t - 2g_s^3$. Then, the identification (10.7) induces the following identities.

$$(10.10) \quad \begin{aligned} g_t + 2g_s^3 &= \frac{2}{3\sqrt{3}}\pi^3(\alpha_1^3 - \beta_3) \omega_0^{-3} \\ g_t - 2g_s^3 (= -G_3(\frac{1}{3}\omega_0)) &= -\frac{2}{3\sqrt{3}}\pi^3(\alpha_1^3 + \beta_3) \omega_0^{-3} \end{aligned}$$

11. Discriminant conjecture

This is the last section of the study of the period map of types A_2 , B_2 and G_2 . We recover the classical *modular discriminant formula* for all types, and examine in **Theorem 11.1** that the discriminant conjecture in [23, §6] holds for these particular toy cases. Let us recall the conjecture in the original form.

Conjecture 6. *Let W be a crystallographic finite reflection group. Is the $k(W)$ th power root of δ_W , say λ_W (up to a constant factor), an automorphic*

form for the group $\Gamma(W)$ with the character ϑ_W ? Can one find an infinite product expression for λ_W compatible with Conjecture 4?

We first explain notation in the conjecture. In [23], W is a Weyl group for an irreducible finite root system of any type, but in the present paper, we restrict ourselves only to the types A_2, B_2 or G_2 . The number $k(W)$ and the character ϑ_W are defined in [23, §6], but, in the present paper, in (4.8) and (4.10), respectively. The δ_W is a generator of anti-invariants whose square is the discriminant $\Delta_{I_2(p)}$. We shall explain about δ_W and λ_W again in Theorem 11.1. The modular group $\Gamma(W)$ is, in [23, §6, 6.4 Example], unfortunately, wrongly stated to be equal to the congruence group $\Gamma_0([p/2])$. However, according to a result (4.6) in the present paper, the group $\Gamma(W)$ should be corrected to be $\Gamma_1([p/2])$.²⁷ The Conjecture 4 mentioned in the Conjecture 6 is something about the “liftability” of the ring of Eisenstein series to the ring of the “Galois” covering by the Weyl group of the type $I_2(p)$. However, we will not discuss on this in the present paper.

The answer to Conjecture is given by a use of Dedekind eta function $\eta(\tau) := q^{1/24} \prod_{n=1}^{\infty} (1 - q^n)$ ($q = \exp(2\pi\sqrt{-1}\tau)$). Recall that a function in $\tau \in \mathbb{H}$ is called an eta-quotient if it has a finite product/quotient expression $\prod_{i=1}^s \eta^{r_i}(m_i\tau)$ where $s, m_i \in \mathbb{Z}_{>0}, r_i \in \mathbb{Z}$. A holomorphic modular form that is non-vanishing on \mathbb{H} and has integer Fourier coefficients at infinity, is an integer multiple of an eta-quotient (see [18]).

Let us come back to the equalities (10.8), (10.9) and (10.10), and show that they admit eta-quotient expressions. LHSs, as defining equations of irreducible components of the discriminants, do not vanish on $S_{I_2(p)} \setminus D_{I_2(p)}$. So, after the identification (10.3), they do not vanish on $\tilde{\mathbb{H}}$. On the other hand, those generators $e_4, e_6, \alpha_2, \beta_4, \alpha_1, \beta_2$ in RHS, are described by theta-functions, and, therefore, have integral Fourier coefficients at infinity (see [1]). Then, we have the following expressions.

Table 7: Eta-quotients of irreducible components of the discriminant

$$\begin{aligned} \Gamma_1(1) &: \frac{e_4^3 - e_6^2}{1728} = \eta(\tau)^{24}, \\ \Gamma_1(2) &: \beta_4 = \frac{\eta(2\tau)^{16}}{\eta(\tau)^8}, \quad \alpha_2^2 - 64\beta_4 = \frac{\eta(\tau)^{16}}{\eta(2\tau)^8}, \\ \Gamma_1(3) &: \frac{1}{54}(\alpha_1^3 - \beta_3) = \frac{\eta(3\tau)^9}{\eta(\tau)^3}, \quad \frac{1}{2}(\alpha_1^3 + \beta_3) = \frac{\eta(\tau)^9}{\eta(3\tau)^3} \end{aligned}$$

Proof. Case of type A_2 is classical (e.g. [7, 11]).

²⁷In fact, $\Gamma_0(N) = \Gamma_1(N)$ for $N = 1, 2$, but $\Gamma_0(3)$ is a double extension of $\Gamma_1(3)$ by $\pm id$. Since in [23], one use the same generators as (4.3) in the present paper as for the generators of $\Gamma(I_2(p))$, practically the calculations in [23] are still meaningful, and we shall use them in the present paper.

Case of type B_2 : Because of level 2 condition, candidates of eta quotients are of the form $c\eta(\tau)^p\eta(2\tau)^q$ for unknowns $c \in \mathbb{C}$ and $p, q \in \mathbb{Z}$, satisfying $p + q = 2 \cdot \text{weight} = 8$. Recall Table 6, so that $\beta_4|4E(\sqrt{-1}\infty) = 0$ (simple zero in q), $\beta_4|4S(\sqrt{-1}\infty) = 1/256$ and $\alpha_2^2 - 64\beta_4|4E(\sqrt{-1}\infty) = 1$, $\alpha_2^2 - 64\beta_4|4S(\sqrt{-1}\infty) = 0$ (simple zero in q). Posing these constraints on the eta-quotients, we determine c, p and q , and obtain the expression.

Similar proof works for the type G_2 . □

Applying above expressions for (10.8), (10.9) and (10.10), we are now able to express the discriminant form $\Delta_{I_2(p)}$ (2.5) and its reduced form $\Delta_{I_2(p)}^{red}$ by some products of eta quotients. As is expected (since discriminant vanishes on both cusps), the results are no-longer *eta-quotients*, but are *eta-products*, i.e. they don't have denominators.

Table 8: Eta-product expressions of discriminants

$$\begin{aligned}
 \Delta_{A_2} &= -27g_l^2 + g_s^3 &= \pi^{12} \eta(\tau)^{24} \omega_0^{-12} \\
 \Delta_{B_2} &= (8g_l + g_s^2)(-8g_l + g_s^2)^2 &= 512 \pi^{12} \eta(\tau)^{24} \omega_0^{-12} \\
 \Delta_{G_2} &= (g_l + 2g_s^3)(-g_l + 2g_s^3)^3 &= -\frac{27}{4} \pi^{12} \eta(\tau)^{24} \omega_0^{-12}
 \end{aligned}
 \tag{11.1}$$

Table 9: Eta-product expressions of reduced discriminants

$$\begin{aligned}
 \Delta_{A_2}^{red} &= -27g_l^2 + g_s^3 &= \pi^{12} \eta(\tau)^{12} \eta(2\tau)^{12} \omega_0^{-12} \\
 \Delta_{B_2}^{red} &= -64g_l^2 + g_s^4 &= 256\pi^8 \eta(\tau)^8 \eta(2\tau)^8 \omega_0^{-8} \\
 \Delta_{G_2}^{red} &= -g_l^2 + 4g_s^6 &= -4\pi^6 \eta(\tau)^6 \eta(3\tau)^6 \omega_0^{-6}
 \end{aligned}
 \tag{11.2}$$

We formulate final Theorem of the present paper, where both formulae (11.2) and (11.1) give answers to the first half and the latter half of Conjecture, respectively.

Theorem 11.1. *1. For all three types A_2 , B_2 and G_2 , set*

$$\lambda_{I_2(p)}(\tau) := \eta(\tau)\eta([p/2]\tau).
 \tag{11.3}$$

Then, (i) $\lambda_{I_2(p)}(\tau)$ is a modular form of weight 1 of the group $\Gamma_1([p/2])$ with respect to the character $\vartheta_{I_2(p)}$ (4.10),

(ii) The power $\delta_{I_2(p)} := \lambda_{I_2(p)}(\tau)^{k(I_2(p))}$ is a generator of the module anti-invariant modular forms (recall (4.8) for $k(I_2(p))$),

(iii) The power $\lambda_{I_2(p)}(\tau)^{2k(I_2(p))}$, up to a non-zero constant factor, corresponds by (10.3) to the reduced discriminant form $\Delta_{I_2(p)}^{red}$.

2. For all three types A_2, B_2 and G_2 , the discriminant form $\Delta_{I_2(p)}$ (2.5), up to a non-zero constant factor, corresponds by (10.3) to

$$q \prod_{n=1}^{\infty} (1 - q^n)^{24}$$

called the modular discriminant (or discriminant function).²⁸

Proof. 1. (i) The calculations given in [23] to show the modularity of $\lambda_{I_2(p)}$ with the character ϑ are still valid. For a sake of completeness of the present paper, we recall it by adjusting notations.

For simplicity, we introduce a number $N \in \mathbb{Z}_{>0}$ called level, where $N = [p/2]$ in case of type A_2, B_2 and G_2 . Set $\zeta := \exp(\pi\sqrt{-1}/12)$ so that $\zeta^{N+1} = \exp(\pi\sqrt{-1}/k(I_2(p)))$ ($N = 1, 2, 3$ and $p = 3, 4, 6$). In view of (4.3), (4.6) and (4.10), it is sufficient to show the following.

Lemma 11.2. *The $\lambda_N := \eta(\tau)\eta(N\tau)$ for $N \in \mathbb{Z}_{>0}$ is a modular form of $\Gamma_1(N)$ with a character ϑ_N , where the character satisfies*

$$\vartheta_N : \tilde{a}_N := \begin{bmatrix} 1 & 0 \\ -N & 1 \end{bmatrix}, \tilde{b}_N := \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \mapsto \zeta^{N+1} \in \mathbb{C}^\times$$

Proof of Lemma 10.2. By definition, λ_N is automatically a modular form with a character of the group $\Gamma_1(N)$ (see, e.g. [18]). Recalling (10.2) and (10.3), we have only to show $(\tilde{a}_N^*)^{-1}(\lambda_N\omega_0^{-1}) = \zeta^{-N-1}(\lambda_N\omega_0^{-1})$ and $\tilde{b}_N^*(\lambda_N\omega_0^{-1}) = \zeta^{N+1}(\lambda_N\omega_0^{-1})$. In the following, we shall use a sign convention on the monodromy of the eta-function from [15, p. 121].

Recall (10.2), $\tilde{b}_N^*(\tau) = \tau + 1$ and $\tilde{b}_N^*(\omega_0) = \omega_0$. Then, using the transformation formula $\eta(\tau + 1) = \zeta\eta(\tau)$, we obtain:

$$\begin{aligned} \tilde{b}_N^*(\lambda_N\omega_0^{-1}) &:= \eta(\tau + 1)\eta(N(\tau + 1))\omega_0^{-1} \\ &= \zeta\eta(\tau) \zeta^N\eta(N\tau)\omega_0^{-1} = \zeta^{N+1}(\lambda_N\omega_0^{-1}) \end{aligned}$$

Recall (10.2) so that $\tilde{a}_N^*(\tau) = \frac{\tau}{1-N\tau}$ and $\tilde{a}_N^*(\omega_0) = \omega_0 - N\omega_1 = \omega_0(1 - N\tau)$. Then, using the transformation formula $\eta(-/\tau) = \sqrt{\frac{\tau}{\sqrt{-1}}}\eta(\tau)$, we obtain:

$$(\tilde{a}_N^*)^{-1}(\lambda_N\omega_0^{-1}) := \eta\left(\frac{\tau}{1 - N\tau}\right)\eta\left(\frac{N\tau}{N\tau + 1}\right)\frac{\omega_0^{-1}}{1 - N\tau}$$

²⁸All results are well-known for the type A_2 . However, for the other types B_2 and G_2 , at present, there is no geometric explanation why such power roots of the reduced discriminant $\Delta_{I_2(p)}^{red}$ form modular forms with character, and why all three geometric discriminants $\Delta_{I_2(p)}$ take the same modular discriminant form. What are $\lambda_{I_2(p)}$'s? Does mirror symmetric interpretation give a hint to understand them?

$$\begin{aligned}
 &= \sqrt{\frac{-(N\tau + 1)/\tau}{\sqrt{-1}}} \eta(-N - 1/\tau) \zeta \eta\left(-\frac{1}{N\tau + 1}\right) \frac{\omega_0^{-1}}{1 + N\tau} \\
 &= \sqrt{\frac{-(N\tau + 1)/\tau}{\sqrt{-1}}} \frac{\tau}{\sqrt{-1}} \frac{N\tau + 1}{\sqrt{-1}} \zeta^{-N} \eta(\tau) \zeta \eta(N\tau + 1) \frac{\omega_0^{-1}}{1 + N\tau} \\
 &= \zeta^{-N+2} \sqrt{\frac{1}{\sqrt{-1}}} \eta(\tau) \eta(N\tau) \omega_0^{-1} \\
 &= \zeta^{-N-1} \zeta(\tau) \eta(N\tau) \omega_0^{-1} = \zeta^{-N-1} (\lambda_N \omega_0^{-1})
 \end{aligned}$$

End of Proof of Lemma 10.2. □

(ii) Recalling the character θ (4.11), $\delta_{I_2(p)}$ is obviously an anti-invariant of the group $\Gamma_1(N)$.

(iii) This a paraphrase of (11.2).

2. This is only a paraphrase of (11.1). □

Remark 5. In [23, §6], we formulated 6 conjectures. The present paper gives positive answers to all conjectures for the types A_2, B_2 and G_2 . The conjectures seem to be still valid for all types of crystallographic reflection groups. That is, we ask for a construction of Eisenstein series for all types to answer to Jacobi inversion problem, where we may need special consideration for low weight cases as in the present paper.

12. Concluding remarks

The findings of the duplication formula for the Lemniscate arc length integral due to Fagnano (1718) and its generalization to the addition formula due to Euler (1751) were naturally understandable by inverting the variables, i.e. by parametrizing the Cartesian coordinates of the curve by the arc length. This led to the finding of new periodic functions, i.e. the elliptic functions, beyond the trigonometric or exponential functions. It is impressive to see the historical developments caused by these finding of the elliptic integrals, from the classical Abel-Jacobi theory through the modern mixed Hodge theory.

The author, however, was attracted by other aspects of elliptic integrals. Namely, covariant differentiations of the elliptic integral of the first kind by suitable deformation parameters give other kinds of elliptic integrals. That is, the elliptic integral of the first kind is a *potential for other periods*. He called this property the *primitivity* of the elliptic integral of the first kind.

The primitivity is combined with another remarkable property. Namely, the *rank* 2 of the lattice of cycles used for the elliptic integrals is equal to the

dimension 2 of the unfolding parameters (g_s, g_t) of the elliptic curve. That is, the map from the space of parameters to the space of periods of integrals of the lattice becomes an equidimensional morphism. Actually, it is locally bi-holomorphic for the elliptic integral of the first kind. The Schottky problem in classical Abelian integral theory to determine the image set of periods (cf. [27]) is resolved automatically in this setting! The author called this property the *equi-dimensionality* of the period map of elliptic integrals of the first kind.

Then, the inverse morphism from the space of integral values to the space of original family of curves were described by *theta-series* by Jacobi, *Eisenstein series* and later by *Weierstrass functions*. We call this procedure “solutions to *Jacobi’s inversion problem*”.

Following those classics, the author was inspired to look for (higher dimensional) analogs of the elliptic integral of the first kind, which carries the primitivity and the equi-dimensionality. That is the theory of integrals of *primitive forms* over Lefschetz vanishing cycles [19]. If we look back some historical works from the view point of primitive forms, the works by E. Picard [17] (1883) and by G. Shimura [25, 26] (1963, 1964) can be regarded already as some part of period integrals of primitive forms of type E_6 and $E_8, E_8^{(1,1)}$ and some others, respectively.

Nowadays, primitive forms become a driving force for constructing new integrable hierarchies, and play a role in mirror symmetry from complex geometric side. However, this is one aspect of the primitive form theory, i.e. algebraic analytic aspects. The transcendental aspects, i.e. the period integral theory over Lefschetz vanishing cycles is missing still. Primitive forms may be able to play their full original power only after they are integrated to period maps, and then the solutions to the Jacobi inversion problem should lead us to the study of new transcendental functions [24].

However, the integral theory over closed cycles is a quite hard subject, since they form a closed world which is rigid and inflexible. We first need to embed them in a big ocean of integrals over open cycles, where we have wide freedom of making new pictures and theories, as was done in the original works of arc-length integrals by Fagnano and Euler. Also, the classical abelian integral theory by Riemann was successful through integrals over open intervals, called the Jacobian variety theory, where the inversion maps are described by theta functions [27].

Thus, it was the pleasure to reinterpret the classical elliptic integrals of the families for Weierstrass, Legendre-Jacobi and Hesse in terms of integrals of primitive forms over vanishing cycles of types A_2, B_2 and G_2 as in the present paper (types D_4 and A_3 are ongoing). We tried to make clear the importance of integrals over open paths by showing that their inversion functions are the

solutions of *Hamilton's equation of motion* in §6. That is, open integrals are inverse to certain Dynamical systems. Actually, this fact was the key reason, why we could determine the inversion map in §7–9 to solve Jacobi's inversion problem by introducing *generalized Eisenstein series* for each type.

We do not know yet what are the higher dimensional analog of them: how to invert the period map to answer Jacobi's inversion problem, how to generate the inversion functions [21]. When the vanishing cycles form a finite root system, there are some conjectural descriptions [23]. The positive answer to the conjectures for types A_2 , B_2 and G_2 in the present paper by a use of the Dedekind *eta-function* is a toy model and did not give any new transcendental function. However, we may get hope that the conjectures still hold for all other types of root systems by extending to *higher dimensional eta-function*.

One may expect the mirror symmetry and the study of d-branes on open string theory in high energy physics may give some suggestions for the understanding of the period maps for primitive forms, since the study of the power roots of the discriminant, as done in the present paper, is explained not from the primitive form side by itself but from the structure of the root system of the vanishing cycles [23] which belongs to the mirror side, i.e. the symplectic geometry side.

Acknowledgements

The author expresses his gratitudes to Yoshihisa Obayashi, who draw Figures 1–5 of the present paper. Particular thanks go to Hiroki Aoki, without whose helps the author may have not accomplish the identification of the ring of Eisenstein series of type $I_2(p)$ with the ring of modular forms of the congruence group $\Gamma_1([p/2])$. He expresses also gratitudes to Yoshihisa Saito, Kenji Iohara, Takashi Takebe, Tomoyoshi Ibukiyama, Hiroyuki Yoshida, Masanobu Kaneko and Akio Fujii for helpful and inspiring discussions, and to Yota Shamoto for careful reading of manuscripts. This work was partially supported by JSPS KAKENHI Grant Number 18H01116.

References

- [1] HIROKI AOKI and TOMOYOSHI IBUKIYAMA. Simple graded rings of Siegel modular forms, differential operators and Borcherds products. *Internat. J. Math.*, **16**(3):249–279, 2005. [MR2130626](#)
- [2] HIROKI AOKI and KYOJI SAITO. Modular forms from Weierstrass \mathfrak{p} -function and zeta-function. Forthcoming.

- [3] TSUNEO ARAKAWA, MASANOBU KANEKO, and TOMOYOSHI IBUKIYAMA. *Bernoulli numbers and Zeta functions*. Springer Verlag, Tokyo, 2014. [MR3307736](#)
- [4] NICOLAS BOURBAKI. *Éléments de mathématique*. Masson, Paris, 1981. Groupes et algèbres de Lie. Chapitres 4, 5 et 6. [Lie groups and Lie algebras. Chapters 4, 5 and 6]. [MR0647314](#)
- [5] EGBERT BRIESKORN. Die Fundamentalgruppe des Raumes der regulären Orbits einer endlichen komplexen Spiegelungsgruppe. *Invent. Math.*, **12**:57–61, 1971. [MR0293615](#)
- [6] EGBERT BRIESKORN and KYOJI SAITO. Artin-Gruppen und Coxeter-Gruppen. *Invent. Math.*, **17**:245–271, 1972. [MR0323910](#)
- [7] JAN HENDRIK BRUINIER, GERARD VAN DER GEER, GÜNTER HARDER, and DON ZAGIER. *The 1-2-3 of modular forms*. Universitext. Springer-Verlag, Berlin, 2008. Lectures from the Summer School on Modular Forms and their Applications held in Nordfjordeid, June 2004, Edited by Kristian Ranestad. [MR2385372](#)
- [8] PIERRE DELIGNE. Les immeubles des groupes de tresses généralisés. *Invent. Math.*, **17**:273–302, 1972. [MR0422673](#)
- [9] FRED DIAMOND and JERRY SHURMAN. *A first course in modular forms*, volume 228 of Graduate Texts in Mathematics. Springer-Verlag, New York, 2005. [MR2112196](#)
- [10] MARTIN EICHLER and DON ZAGIER. *The theory of Jacobi forms*, volume 55 of Progress in Mathematics. Birkhäuser Boston, Inc., Boston, MA, 1985. [MR0781735](#)
- [11] EBERHARD FREITAG and ROLF BUSAM. *Complex analysis*. Universitext. Springer-Verlag, Berlin, second edition, 2009. [MR2513384](#)
- [12] F. A. GARSIDE. The braid group and other groups. *Quart. J. Math. Oxford Ser. (2)*, **20**:235–254, 1969. [MR0248801](#)
- [13] ADOLF HURWITZ and R. COURANT. *Vorlesungen über allgemeine Funktionentheorie und elliptische Funktionen*. Interscience Publishers, Inc., New York, 1944. [MR0011320](#)
- [14] HIROSHIGE KAJIURA, KYOJI SAITO, and ATSUSHI TAKAHASHI. Matrix factorization and representations of quivers. II. Type *ADE* case. *Adv. Math.*, **211**(1):327–362, 2007. [MR2313537](#)
- [15] NEAL KOBLITZ. *Introduction to elliptic curves and modular forms*, volume 97 of Graduate Texts in Mathematics. Springer-Verlag, New York, second edition, 1993. [MR1216136](#)
- [16] SERGE LANG. *Introduction to modular forms*. Grundlehren der mathematischen Wissenschaften, No. 222. Springer-Verlag, Berlin-New York, 1976. [MR0429740](#)

- [17] E. PICARD. Sur des fonctions de deux variables indépendantes analogues aux fonctions modulaires. *Acta Math.*, **2**:114–135, 1883. [MR1554595](#)
- [18] JEREMY ROUSE and JOHN J. WEBB. On spaces of modular forms spanned by eta-quotients. *Adv. Math.*, **272**:200–224, 2015. [MR3303232](#)
- [19] KYOJI SAITO. Period mapping associated to a primitive form. *Publ. Res. Inst. Math. Sci.*, **19**(3):1231–1264, 1983. [MR0723468](#)
- [20] KYOJI SAITO. On a linear structure of the quotient variety by a finite reflexion group. *Publ. Res. Inst. Math. Sci.*, **29**(4):535–579, 1993. [MR1245441](#)
- [21] KYOJI SAITO. Primitive automorphic forms. In *Mathematics Unlimited — 2001 and Beyond*, pages 1003–1018. Springer, 2001. [MR1852200](#)
- [22] KYOJI SAITO. Polyhedra dual to the Weyl chamber decomposition: a précis. *Publ. Res. Inst. Math. Sci.*, **40**(4):1337–1384, 2004. [MR2105710](#)
- [23] KYOJI SAITO. Uniformization of the orbifold of a finite reflection group. In *Frobenius manifolds*, vol. E36 of Aspects Math., pages 265–320. Friedr. Vieweg, Wiesbaden, 2004. [MR2115774](#)
- [24] KYOJI SAITO. Jugendtraum of a mathematician. *Asia Pac. Math. Newsl.*, **1**(3):1–6, 2011. [MR2857888](#)
- [25] GORO SHIMURA. On analytic families of polarized abelian varieties and automorphic functions. *Ann. of Math. (2)*, **78**:149–192, 1963. [MR0156001](#)
- [26] GORO SHIMURA. On purely transcendental fields automorphic functions of several variable. *Osaka Math. J.*, **1**(1):1–14, 1964. [MR0176113](#)
- [27] CARL LUDWIG SIEGEL. *Topics in complex function theory. Vol. I.* Wiley Classics Library. John Wiley & Sons, Inc., New York, 1988. Elliptic functions and uniformization theory, Translated from the German by A. Shenitzer and D. Solitar, With a preface by Wilhelm Magnus, Reprint of the 1969 edition, A Wiley-Interscience Publication. [MR1008930](#)
- [28] OSCAR ZARISKI. *Collected papers. Vol. III.* The MIT Press, Cambridge, Mass.-London, 1978. Topology of curves and surfaces, and special topics in the theory of algebraic varieties, Edited and with an introduction by M. Artin and B. Mazur, Mathematicians of Our Time. [MR0505104](#)

Kyoji Saito
Research Institute for Mathematical Sciences
Kyoto University
Sakyoku Kitashirakawa
Kyoto, 606-8502
Japan
Laboratory of AGHA
Moscow Institute of Physics and Technology
9 Institutskiy per.
Dolgoprudny, Moscow Region, 141700
Russian Federation
E-mail: kyoji.saito@ipmu.jp