

Surveys in Differential Geometry

- Vol. 1: Lectures given in 1990
edited by S.-T. Yau and H. Blaine Lawson
- Vol. 2: Lectures given in 1993
edited by C.C. Hsiung and S.-T. Yau
- Vol. 3: Lectures given in 1996
edited by C.C. Hsiung and S.-T. Yau
- Vol. 4: Integrable systems
edited by Chuu Lian Terng and Karen Uhlenbeck
- Vol. 5: Differential geometry inspired by string theory
edited by S.-T. Yau
- Vol. 6: Essays on Einstein manifolds
edited by Claude LeBrun and McKenzie Wang
- Vol. 7: Papers dedicated to Atiyah, Bott, Hirzebruch, and Singer
edited by S.-T. Yau
- Vol. 8: Papers in honor of Calabi, Lawson, Siu, and Uhlenbeck
edited by S.-T. Yau
- Vol. 9: Eigenvalues of Laplacians and other geometric operators
edited by A. Grigor'yan and S.-T. Yau
- Vol. 10: Essays in geometry in memory of S.-S. Chern
edited by S.-T. Yau
- Vol. 11: Metric and comparison geometry
edited by Jeffrey Cheeger and Karsten Grove
- Vol. 12: Geometric flows
edited by Huai-Dong Cao and S.-T. Yau
- Vol. 13: Geometry, analysis, and algebraic geometry
edited by Huai-Dong Cao and S.-T. Yau
- Vol. 14: Geometry of Riemann surfaces and their moduli spaces
edited by Lizhen Ji, Scott A. Wolpert, and S.-T. Yau

Volume IV

Surveys in
Differential Geometry

Integrable systems

edited by

Chuu-Lian Terng and Karen Uhlenbeck



International Press

www.intlpress.com

Surveys in Differential Geometry, Vol. 4
Integrable systems

Editors:
Chuu-Lian Terng
Karen Uhlenbeck

Copyright © 1998, 2010 by International Press
Somerville, Massachusetts, U.S.A.

All rights reserved. Individual readers of this publication, and non-profit libraries acting for them, are permitted to make fair use of the material, such as to copy a chapter for use in teaching or research. Permission is granted to quote brief passages from this publication in reviews, provided the customary acknowledgement of the source is given. Republication, systematic copying, or mass reproduction of any material in this publication is permitted only under license from International Press.

Excluded from these provisions is material in articles to which the author holds the copyright. In such cases, requests for permission to use or reprint should be addressed directly to the author.

ISBN 978-1-57146-215-2

*Paperback reissue 2010. Previously published in 1998 under ISBN 1-57146-066-7 (clothbound).
An error in the previous publication's volume title has been corrected.*

Printed in the United States of America.

Contents

| | |
|---|-----|
| Introduction | |
| Chuu-Lian Terng and Karen Uhlenbeck | 5 |
| Integrable systems in Riemannian geometry | |
| Nigel Hitchin | 21 |
| Seiberg-Witten Integrable Systems | |
| Ron Y. Donagi | 83 |
| Five Lectures on Soliton Equations | |
| Edward Frenkel | 131 |
| Geometry of the Space of Orbits of a Coxeter Group | |
| Boris Dubrovin | 181 |
| Differential Geometry of Moduli Spaces | |
| Boris Dubrovin | 213 |
| Symplectic Forms in the Theory of Solitons | |
| I.M. Krichever and D.H. Phong | 239 |
| Poisson Actions and Scattering Theory for Integrable Systems | |
| Chuu-Lian Terng and Karen Uhlenbeck | 315 |
| Loop Groups and Equations of KdV Type | |
| Graeme Segal and George Wilson | 403 |
| Scattering and Inverse Scattering for First Order Systems | |
| R. Beals and R. R. Coifman | 467 |

Introduction

Chuu-Lian Terng
Karen Uhlenbeck

I. The theory of integrable systems has been an active area of mathematics for the past thirty years. Different aspects of the subject have fundamental relations with mechanics and dynamics, applied mathematics, algebraic structures, theoretical physics, analysis and spectral theory and geometry. Most differential geometers have some knowledge and experience with finite dimensional integrable systems as they appear in symplectic geometry (mechanics) or ordinary differential equations, although the reformulation of part of this theory as algebraic geometry is not as commonly known. There are two quite separate methods of extension of these ideas to partial differential equations; one based on algebraic constructions and one based on spectral theory and analysis. These are less familiar still to differential geometers. This volume is a collection of papers intended to introduce and explain some of the more geometric aspects of integrable systems to the community.

This introduction contains a short historical discussion of the early geometers' ideas on "integrable geometric constructions" followed by a brief discussion of the material in the papers. An excellent historical treatment of finite dimensional integrable systems can be found in Moser's article [1980 Mo]. Many expositions on integrable equations (cf. [1991 AC], [1997 Pa]) discuss the history of the Korteweg de Vries (KdV) equation, beginning with the physical observation of Russell [1834 R] and the discovery of the equations by Boussinesq [1871 Bo], Korteweg and de Vries [1895 KdV]. In addition to describing water waves, the KdV equation also arises as a the model for the universal limit of lattice vibrations as the spacing goes to zero. Fermi-Pasta-Ulam's [1955 FPU] surprising numerical experiments on an anharmonic lattice, and the ingenious explanation by Zabuski-Kruskal [1965 ZK] of the results in terms of "solitons" of the KdV equation were quickly followed by a ground-breaking paper of Gardner, Greene, Kruskal and Miura [1967 GGKM] that introduced the method of solving KdV using the inverse scattering transform for the Hill's operator. This represents the approach of applied mathematics to discovering equations, solutions, and

their scattering properties. The local differential geometry of the 19th century, on the other hand, foreshadows to some extent modern algebraic constructions. Rather than giving a description of the development of the basic ideas, we describe some of the early geometry and include a bibliography which lists many of the important early papers and a selection from the years between 1975 and 1985 in chronological order. The few recent papers are those we specifically refer to in this introduction.

Many geometric equations are known to have integrable aspects, especially if one takes into account that most experts do not have a good definition of “integrable” as applied to partial differential equations, particularly elliptic examples. In addition to those we mention in our historical discussion, the equations for harmonic maps (sigma models) from surfaces into groups ([1978 ZMi], [1981 Ch], [1989 U]), harmonic tori in symmetric spaces [1993 BFD], constant mean curvature surfaces in space forms ([1989 PS], [1991 Bo]), isometric immersions of space forms in other space forms ([1919 Ca], [1980 TT]), and the theory of affine spheres ([1910 Tz], [1997 BS]) and affine minimal surfaces ([1980 CT]) are all examples of “elliptic” integrable systems. Likewise, the Yang-Mills equations on C^2 or $E^{2,2}$, their reductions to monopole equations in three variables, and self-dual equations on surfaces are new equations which originated in the physics literature of the late 1970’s [1987 Hi]. One atypical development illustrating the great variety in the geometric theory is the reduction of the global $SU(N)$ monopole equation on R^3 to an integrable ordinary differential equation by the physicist Nahm [1984, Do]. The ideas surrounding string theory result in a series of deep and not completely understood connections between representation theory of certain algebras and many of the more classical theories of integrable systems in mathematics. Most recently, supersymmetric quantum field theories produce in a natural way moduli spaces of vacua or ground states which have new geometry generated by the supersymmetry. Since the supersymmetry generalizes the classical symmetries which produce integrals for the Euler-Lagrange equations via Noether’s theorem, the connection with integrability is perhaps not surprising. However, this does not explain entirely the use of integrable systems in hyper-Kähler geometry ([1992 Hi]), Seiberg-Witten theory ([1996 DW]), special Kähler geometry ([1996 F], [1997 CRTP], [1997 Fr]) and quantum cohomology ([1991 W], [1994 KM], [1994 RT]).

The 19th century geometers were mainly interested in the local theory of surfaces in R^3 , which we might regard as the prehistory of these modern constructions. The sine-Gordon equation arose first through the theory of surfaces of constant Gauss curvature -1 , and the reduced 3-wave equation can be found in Darboux’s work on triply orthogonal systems of R^3 ([1866 Da]). In 1906, a student of Levi Civitas, da Rios, wrote a master’s thesis, in which he modeled the movement of a thin vortex filament in a viscous liquid using the equations of a curve propagating in R^3 along its binormal ([1906 dR]). It was much later that Hasimoto ([1971 Ha]) showed the equivalence of this system with the non-linear Schrödinger equation. Since the equations were rediscovered somewhat independently of their geometric history, the main contribution of the classical

geometers lies in the methods they developed for constructing explicit solutions of these equations. Ribaucour, Bianchi, Lie, and Bäcklund worked out the theory of what are now called Backlund transformations ([1883 Ba]). Lie developed a general theory of transformations known as “Lie transformations” which includes as examples scaling transformations and Lorentz transformations ([1979 L]). Darboux transformations for the Hills operator (the isospectral operator for KdV) were constructed by Darboux ([1882 Da]), but had been found even earlier by Moutard ([1875 M]). Most of the integrable partial differential equations are known to have Bäcklund, Lie and Darboux transformations.

II. The three-wave equation and the sine-Gordon equation arise out of the study of the classical differential geometry of R^3 . We take a short trip back to this subject. A triply orthogonal system on R^3 is a local coordinate system $X(x_1, x_2, x_3)$ on R^3 such that the Euclidean metric ds^2 on R^3 has the form

$$ds^2 = a_1(x)^2 dx_1^2 + a_2(x)^2 dx_2^2 + a_3(x)^2 dx_3^2$$

for some smooth functions a_1, a_2, a_3 , i.e., $X_{x_i} \cdot X_{x_j} = a_i^2 \delta_{ij}$ for $1 \leq i, j \leq 3$. Then the three one-parameter families of surfaces in R^3 , defined by setting the different x_i to be constant, are mutually orthogonal. Already in 1813, Dupin ([1813 Du]) published a proof that the x_i -curves are lines of curvature for these surfaces. The flatness of ds^2 implies that

$$\begin{cases} (\beta_{ij})_{x_j} + (\beta_{ji})_{x_i} + \beta_{ik}\beta_{kj} = 0, \\ (\beta_{ij})_{x_k} = \beta_{ik}\beta_{kj}, \quad i, j, k \text{ distinct,} \end{cases}$$

where $\beta_{ij} = (a_i)_{x_j}/a_j$ for $i \neq j$. These are the *Equations of Lamé* or *The Lamé System*. Given a solution (β_{ij}) of the Lamé system, finding a triply orthogonal system with these (β_{ij}) means finding the (a_1, a_2, a_3) that satisfy

$$(a_i)_{x_j} = \beta_{ij}a_j, \quad \text{for all } i \neq j.$$

Solutions of this latter system depend on three functions of one variable. Thus, given one triply orthogonal system, we can derive from it infinitely many others, parametrized by three functions of one variable. These transformations of a triply orthogonal system are referred to as Combescure transformations ([1867 Co]).

Triply orthogonal systems are easy to construct, and consequently many solutions to the equations of Lamé are known. For example, let $f(x_1, x_2)$ be a surface in R^3 parametrized by line of curvatures coordinates, and let N be the unit normal field. Then $X(x_1, x_2, x_3) = f(x_1, x_2) + x_3 N(x_1, x_2)$ is an orthogonal coordinate system for a domain in R^3 . Moreover, if the two fundamental forms of the surface f are given by

$$I = b_1^2 dx_1^2 + b_2^2 dx_2^2, \quad II = \lambda_1 b_1^2 dx_1^2 + \lambda_2 b_2^2 dx_2^2,$$

then the Euclidean metric of R^3 written in this coordinate is $ds^2 = \sum_{i=1}^3 a_i^2 dx_i^2$, where

$$a_i(x_1, x_2, x_3) = (1 - \lambda_i(x_1, x_2)x_3)b_i(x_1, x_2), \quad \text{for } i = 1, 2, \quad a_3 = 1.$$

In this case, the equations of Lamé for $\beta_{ij} = (a_i)_{x_j}/a_j$ are precisely the Gauss and Codazzi equations of the surface f .

Darboux considered the Lamé system restricted to the case when $\beta_{ij} = \beta_{ji}$. This condition is easily seen to be equivalent to the condition that $ds^2 = \frac{\partial\phi}{\partial x_1} dx_1^2 + \frac{\partial\phi}{\partial x_2} dx_2^2 + \frac{\partial\phi}{\partial x_3} dx_3^2$ for some function ϕ . Darboux called a metric with this property an *Egorof* metric. In this case, The Lamé system is equivalent to the reduced 3-wave equation studied more recently by [1973 ZM]. Local analytic solutions of this system were classified by Cartan (cf. [1945 Ca]) using the methods which he and Kähler developed, now usually referred to as the “Cartan-Kähler theory”. In fact, Cartan proved that (i) if c_1, c_2, c_3 are distinct, then the line $t(c_1, c_2, c_3)$ is non-characteristic, and (ii) that if L is a fixed, non-characteristic line, then given any three analytic functions f_1, f_2, f_3 defined on L , there exists a local analytic solution $\beta_{12}, \beta_{13}, \beta_{23}$ of the reduced 3-wave system such that $\beta_{12}, \beta_{13}, \beta_{23}$ are equal to f_1, f_2, f_3 on L . However, the ability to handle the global problem came only with the insights which come from scattering theory, and the solution of the problem with arbitrary smooth and rapidly decaying data is a recent development.

In the first construction we can find of a “Bäcklund transformation”, Ribaucour ([1870 Ri]) produced in 1870 an interesting triply orthogonal system. We include a description of this result to give a flavor of the geometry of that time. Note that classical geometers referred to a surface in R^3 of constant Gauss curvature -1 as *pseudospherical*, and we will keep this terminology. Start with a surface M in R^3 , and let C_p denote the unit circle in the physical tangent plane $T_p M$ in R^3 centered at p . Ribaucour proved that if there exist a one-parameter family of surfaces that meet these tangent circles orthogonally, then, first of all both M and all the surfaces in the one-parameter family are pseudospherical. In addition these pseudospherical surfaces are one of the families of surfaces in a triply orthogonal system. To describe this system, we let $f(u, v)$ denote the embedding of M in R^3 parametrized by line of curvature coordinates (u, v) , with e_1 and e_2 the unit vectors along the lines of curvature in the directions of f_u and f_v respectively. Fix a base point p_0 in M , and find for each angle τ the surface P_τ which is orthogonal to C_{p_0} at the point

$$f(u_0, v_0) + \cos \tau e_1(u_0, v_0) + \sin \tau e_2(u_0, v_0).$$

Define the function $\theta(u, v, \tau)$ to be the angle between $f(u, v) + e_1(u, v)$ and $P_\tau(u, v) \cap C_{f(u, v)}$. Then

$$X(u, v, \tau) = f(u, v) + \cos \theta(u, v, \tau)e_1(u, v) + \sin \theta(u, v, \tau)e_2(u, v)$$

is an orthogonal coordinate system in R^3 , and the surface defined by $\tau = \tau_0$ is P_{τ_0} .

The sine-Gordon equation first arose in the study of pseudospherical surfaces. If M is a surface of negative curvature, then there exist asymptotic coordinates (s, t) . The Codazzi equations of a pseudospherical surface implies that the (s, t) can be chosen to be arc length. There is a reduction of the Gauss equation in which the fact that the Gauss curvature is -1 translates into the sine-Gordon equation, $q_{st} = \sin q$, where q is the angle between the s and t -curves. Thus, by The Fundamental Theorem of surfaces, there is a local bijective correspondence between solutions of the sine-Gordon equation and pseudospherical surfaces, up to rigid motion. The surfaces typically have singularities and self-intersections. Modern computer graphics provide beautiful pictures of such surfaces which one believes the classical geometers would have appreciated very much.

Bäcklund transformations arose from a geometrical construction on pseudospherical surfaces, but to understand them we need to review yet more classical geometry. A *congruence of lines* is a two-parameter family of lines $L(u, v)$ in R^3 . We describe the line $L(u, v)$ as the line through $x(u, v)$ and in the direction of $\xi(u, v)$ parameterized by t ,

$$L(u, v) : \quad x(u, v) + t\xi(u, v).$$

A surface M given by $Y(u, v) = x(u, v) + t(u, v)\xi(u, v)$ for some smooth function t is called a *focal surface* of the congruence of lines if the line $L(u, v)$ is tangent to M at $Y(u, v)$ for all (u, v) . Hence $\xi(u, v)$ lies in the tangent plane of M at $Y(u, v)$, which is spanned by $x_u + t_u\xi + t\xi_u$ and $x_v + t_v\xi + t\xi_v$. This implies that t satisfies the following quadratic equation:

$$\det(\xi, x_u + t\xi_u, x_v + t\xi_v) = 0.$$

In general, this quadratic equation has two distinct solutions for t . Hence generically each congruence of lines has two focal surfaces, M and M^* . This results in a diffeomorphism $\ell : M \rightarrow M^*$ such that the line joining p and $p^* = \ell(p)$ is tangent to both M and M^* . We will call ℓ a *line congruence*. If you have difficulty understanding this construction, it is helpful to think through the construction of a congruence of lines (a one-parameter family) in R^2 and the single focal curve for this family.

A line congruence $\ell : M \rightarrow M^*$ is called *pseudospherical* with constant θ if the distance between p and $p^* = \ell(p)$ is $\sin \theta$ and the angle between the normal of M at p and the normal of M^* at p^* is θ for all $p \in M$. Bäcklund ([1883 Ba]) showed that if ℓ is a pseudospherical line congruence, then both M and M^* are pseudospherical, and ℓ maps asymptotic lines to asymptotic lines. However, the transformations come about from showing that this construction can always be realized. Given a pseudospherical surface, a constant θ , and a unit vector $v_0 \in TM_{p_0}$, not a principal direction, then there exists a unique surface M^* and a pseudospherical congruence $\ell : M \rightarrow M^*$ with constant θ such that $\overline{p_0 p_0^*} = \sin \theta v_0$. Analytically this is equivalent to the statement that if q is a solution of the sine-Gordon equation, then the following overdetermined system of ordinary differential equations is solvable for q^* :

$$\begin{cases} q_x^* &= q_x + 4s \sin\left(\frac{q^*+q}{2}\right), \\ q_t^* &= -q_t + \frac{1}{s} \sin\left(\frac{q^*-q}{2}\right), \end{cases} \quad (BT_{q,s})$$

where $s = \csc\theta - \cot\theta$. Moreover, a solution, q^* is again a solution of the sine-Gordon equation.

Bäcklund's theorem, but for $\theta = \frac{\pi}{2}$, was already known to Bianchi [1879 Bi]. In fact this case had been discovered even earlier (1870) in a different form by Ribaucour ([1870 Ri]). Note that if $X(u, v, \tau)$ is the Ribaucour triply orthogonal system defined above and τ is a fixed constant, then the map $\ell : M \rightarrow P_\tau$ defined by $\ell(f(u, v)) = X(u, v, \tau)$ is a pseudospherical congruence with $\theta = \pi/2$. However, Ribaucour did not realize that his theorem gave a method for constructing new pseudospherical surfaces from a given one.

The classic permutability Theorem for Bäcklund transformations is due to Bianchi (cf. [1909 Ei]). Given two pseudospherical congruences $\ell_i : M_0 \rightarrow M_i$ with angles θ_i respectively and $\sin\theta_1^2 \neq \sin\theta_2^2$, then there exist an algebraic construction of a unique surface M_3 , and two pseudospherical congruences $\tilde{\ell}_i : M_i \rightarrow M_3$, such that $\tilde{\ell}_2\ell_1 = \tilde{\ell}_1\ell_2$. The analytic reformulation of this permutability formula is the following. Suppose q is a solution of the sine-Gordon equation and q_1, q_2 are two solutions of the above system of ordinary differential equations $(BT)_{q,s}$ with constants $s = s_1, s_2$ respectively, where $s_i = \csc\theta_i - \cot\theta_i$. The Bianchi permutability theorem gives a third solution to the sine-Gordon equation

$$q_3 = q + 4 \tan^{-1} \left(\frac{s_1 + s_2}{s_1 - s_2} \tan \left(\frac{q_1 - q_2}{4} \right) \right).$$

What are now called Darboux transformations were discovered by Darboux during his investigation of Liouville metrics. A metric $ds^2 = A(x, y)(dx^2 + dy^2)$ is Liouville if there is a coordinate system (u, v) such that ds^2 is of the form

$$ds^2 = (f(u) - g(v))(du^2 + dv^2)$$

for some f and g of one variable. The classical geometers were interested in such metrics at least in part because Liouville had shown that all geodesics on such surfaces can be obtained by quadratures. The question of deciding whether a metric ds^2 is Liouville led to the study of the following special second order linear partial differential equation

$$w_{xy} = (f(x+y) - g(x-y))w.$$

Darboux was led to look for transformations of Hill's operators in the process of separating variables in this equation. The original analytic version of Darboux transformation ([1882 Da] and [1889 Da] Chap. 9) is the following: Let q be

a smooth function of one variable, λ_0 a constant, and suppose that f satisfies $f'' = (q + \lambda_0)f$. Set

$$q^\sharp = f(f^{-1})'' - \lambda_0.$$

If $y(x, \lambda)$ is the general solution of the Hills operator with potential q :

$$y'' = (q + \lambda)y,$$

then $z = y' - (f/f')y$ is the general solution of the Hills operator with potential q^\sharp :

$$z'' = (q^\sharp + \lambda)z.$$

Next, suppose that we factor

$$D^2 - q - \lambda_0 = (D + v)(D - v),$$

In other words, suppose that v satisfies $v_x + v^2 = q + \lambda_0$. (Here $D = d/dx$.) Choose f so that $f'/f = v$. Then

$$(D - v)(D + v) = D^2 - q^\sharp - \lambda_0.$$

This Darboux theorem gives an algebraic algorithm (without quadrature) to transform general solutions of $D^2 - q - \lambda$ to those of $D^2 - q^\sharp - \lambda$.

Examples where the classical geometry appears in the modern theory of integrable systems are common and easily found. For example, orthogonal systems and their Combescure transformations arise in the local Hamiltonian theory of hydrodynamics of weakly deformed soliton lattices, and are known as the Whitham equations. The latter is the work of Dubrovin and Novikov ([1983 DN]) and Tsarëv ([1991 Ts]). An Egorof metric that is homogeneous of degree m , i.e., $\phi_{x_i}(rx) = r^m \phi_{x_i}(x)$ for all i , gives rise to a Frobenius manifold in the sense of Dubrovin, and such metrics give solutions to the WDVV equation in the study of Gromov-Witten invariants ([1991 W], [1995 RT], [1996 Du]). One of the best known examples is perhaps the Darboux transformations. If $q(x, t)$ is a solution of KdV then the Hills operators with potential $q(\cdot, t)$ are isospectral. It follows that the Darboux transformations of the Hills operators induce transformations on the space of solutions of KdV. This is a critical observation due to Adler and Moser ([1978 AM]) and Deift ([1978 De]).

III. Many of the topics mentioned in the previous paragraphs will reappear in the papers in this volume, including the classical differential geometry. It is easy to understand the importance of the subject of integrable systems in modern mathematics once one notes that all of the papers except the last three mention either in passing or as fundamental motivation applications to constructions which are relevant in string theory and supersymmetric quantum field theory. Techniques from integrable systems, despite their limitations in classical geometry and applications, have provided one among the successful tools for rigorous mathematical interpretation of ideas in quantum field theory. It should be emphasized that it is mainly geometric ideas about classical geometric moduli spaces which can be treated. These spaces arise only as vacua for

the field theories, but the special properties of the various field theories induce geometry on the moduli spaces which is not the usual Riemannian or symplectic geometry familiar to differential geometers. Because of the nature of the questions, the smooth part of the moduli space is not necessarily compact, so local algebraic results reminiscent of the classical 19th century have relevance if a more complete global analysis is not feasible.

Nigel Hitchin's paper "Integrable systems in Riemannian Geometry" starts with a discussion of what integrability means, and of the contrast between this century and the previous one. He goes on to discuss three problems familiar to differential geometers: constant mean curvature surfaces (tori) in R^3 , Einstein metrics with certain symmetries in four dimensions, and hyper-Kähler manifolds. We meet loop groups in the section on tori, the flat Egorov metrics studied by Darboux in the section on Einstein manifolds, and a moduli space (for a supersymmetric quantum field theory) which has one of the special geometries, the hyper-Kähler one. The paper gives three different, beautiful illustrations of reduction of partial differential equations to finite dimensional flows and a nice explanation of "algebraically integrable".

The paper "Seiberg-Witten Integrable Systems" by Ron Donagi deals entirely with finite dimensional algebraically integrable systems which arise directly from a supersymmetric quantum field theory. The integrable system comes from the study of Higg's bundles on curves, and the paper is the only one to attempt a useful discussion of how an integrable system can relate to high energy physics. A host of references to the physics literature are given for the ambitious geometer, and a certain amount of algebraic geometry is needed to follow the details.

The paper "Five Lectures on Soliton Equations" by Edward Frenkel is motivated by the study of deformations of conformal field theories, and affine Lie algebras play a major role. The basic theory for the KdV equation, which is usually presented in terms of isospectral flows for a second order Hill's operator, is rederived for extensions of modified KdV , a system which is contained in the Non-linear Schrodinger hierarchy. This is particularly helpful in understanding the generality with which the algebraic constructions for KdV can be carried over to other classes of integrable systems and their Poisson brackets.

We are fortunate to include two papers by Boris Dubrovin which are most closely related to the classical geometry, but which in addition have important applications in two dimensional topological quantum field theory. In the paper "Differential Geometry of Moduli Spaces and Applications to Soliton Equations and Topological Conformal Field Theory", the relationship among Egorov metrics, n-wave equations, flat metrics on moduli spaces of curves with marked points and topological conformal field theories is discussed. The moduli space in the second paper "Differential Geometry of the Space of Orbits of a Coxeter Group" is a space of orbits, and carries the natural structure of a Frobenius manifold, a coordinate free description of the WDVV (Witten-Dijkgraaf, Verlinde, Verlinde) equations.

The emphasis in the paper "Symplectic Forms in the Theory of Solitons"

by Igor Krichever and Duong Phong lies on the construction of a universal symplectic form, which can be specialized to many specific cases. We meet, for the first time, global questions in the identification of scattering data for the initial data given in two dimensions for the *KP* equation. A description of the WKB or Whitham flows on the moduli spaces of exact solutions leads into a discussion of leafs of the various moduli spaces of curves and applications to the topological Landau-Ginsburg models treated by Dubrovin. The paper includes applications to the Seiberg-Witten moduli spaces discussed by Donagi. The careful discussion of the analytic versus algebraic constructions is particularly helpful to geometers in the habit of thinking in terms of actual solutions to partial differential equations.

The final recent paper in the collection, “Poisson Actions and Scattering Theory” is our own. While partially motivated by questions from modern physics, it represents an attempt to reconcile the local algebraic methods with an understanding of global inverse scattering theory on the line. Reality conditions consistent with a compact group permit this, but restrict the number of integrable systems to which the global theory is applicable. We also give an explanation of Bäcklund (or Darboux) transformations in terms of the action of a rational subgroup of a loop group. The implication is that not all constructions are best described in terms of Lie groups and infinitesimal actions, especially if one is interested in global questions.

We have reprinted two influential papers which were originally published in 1984 and 1985. The paper “Loop groups and Equations of KdV type” by Graeme Segal and George Wilson is an important introduction to both subjects. A description of the KP hierarchy as commuting flows on an infinite dimensional Grassman manifold was originally due to Sato ([1981 S], [1983 DJKM]), but the work of Segal and Wilson is written up with a geometric emphasis very much appreciated by some of us. We argued for the inclusion of the seminal paper “Scattering and Inverse Scattering for First Order Systems” by Richard Beals and Ronald Coifman. The algebraic geometry needed to treat the periodic theory is referred to by many authors in this volume, but it seems important for geometers to also have some understanding of analytic issues involved in constructing solutions to equations on non-compact domains.

Bibliography

- [1813 Du] Dupin, *Développements de Géométrie*, 1813.
- [1844 Ru] Russell, J.S., *Report on Waves*, 14th Mtg. of the British Assoc. for the Advance. of Science, John Murray, London (1844), 311–390 + 57 plates.
- [1859 La] Lamé, *Lecons sur les coordonnées curvilignes et leurs diverses applications*, 1859, Paris.

- [1866 Da] Darboux, G., *Sur les surfaces orthogonales*, Bull. Soci. Philomathique (1866), 16–.
- [1867 Co] Combescure, E., *Sur les déterminants fonctionnels et les coordonnées curvilignes*, Ann. de l'École Normale **4** (1867), 102–122.
- [1870 Ri] Ribaucour, *Note sur les développées des surfaces*, Comptes Rendus **70** (1870), 330–333.
- [1871 Bo] Boussinesq, M.J., *Théorie de l'intumescence appelée onde solitaire ou de translation se propageant dans un canal rectangulaire*, C. R. Acad. Sc. Paris **72** (1871), 755–759.
- [1875 M] Moutard, *Ricerche sulle superficie a curvatura e sulle elicoidi*, Ann. Scuola normale superiore di Pisa **2** (1879), 285–.
- [1879 Bi] Bianchi, *Ricerche sulle superficie a curvatura e sulle elicoidi*, Ann. Scuola normale superiore di Pisa **2** (1879), 285–.
- [1879 Li] Lie, S., *Ueber Flächen deren Krümmungsradien durch eine Relation Verknüpft sind*, Archiv Math. Naturvidenskab **4** (1879), 510–.
- [1882 Da] Darboux, G., *Sur une proposition relative aux équations linéaires*, Comptes Rendus **94** (1882), 1456–.
- [1883 Ba] Bäcklund, A.V., *Concerning surfaces with constant negative curvature*, New Era Printing Co., Lancaster, PA, original 1883, translation 1905.
- [1889 Da] G. Darboux, *Leçons sur la théorie générale des surfaces, v. II*, Gauthier-Villars, Paris, 1889.
- [1894 Da] G. Darboux, *Leçons sur la théorie générale des surfaces, v. III*, Gauthier-Villars, Paris, 1894.
- [1895 KdV] Koteweg, D.J., de Vries, G., *On the change of form of long waves advancing in a rectangular canal, and on a new type of long stationary waves*, Philos. Mag. Ser. 5 **39** (1895), 422–443.
- [1897 Da] G. Darboux, *Leçons systèmes orthogonaux et les coordonnées curvilignes*, Gauthier-Villars, Paris, 1897.
- [1906 dR] Da Rios, *Sul moto d'un liquido indefinito con un filetto vorticoso di forma qualunque (on the motion of an unbounded liquid with a vortex filament of any shape)*, Rend. Circ. Mat. Palermo **22** (1906), 117–135.
- [1909 Ei] Eisenhart, L.P., *A treatise on the differential geometry of curves and surfaces*, Ginn, 1909.

- [1910 Tz] Tzitzeica, G., *Sur une nouvelle classe des surfaces*, C.R. Acad. Sci. Paris **150** (1910), 955–956.
- [1919 Ca] Cartan, E., *Sur les variétés de courbure constante d'un espace euclidien ou non-euclidien*, Bull. Soc. Math. France **47** (1919), 132–208.
- [1945 Ca] Cartan, É., *Les systèmes différentiels extérieurs et leurs applications géométriques*, Hermann, Paris, 1945.
- [1955 FPU] Fermi, E., Pasta, J., Ulam, S., *Los Alamos Report LA1940* (1955).
- [1965 W] Whitham, G.B., *Non-linear dispersive waves*, Proc. Royal Soc. London **A. 139** (1965), 283–291.
- [1965 ZK] Zabusky, N.J., Kruskal, M.D., *Interaction of solitons in a collisionless plasma and the recurrence of initial states*, Physics Rev. Lett. **15** (1965), 240–243.
- [1967 GGKM] Gardner, C.S., Greene, J.M., Kruskal, M.D., Miura, R.M., *Method for solving the Korteweg-de Vries equation*, Physics Rev. Lett. **19** (1967), 1095–1097.
- [1968 La] Lax, P.D., *Integrals of nonlinear equations of evolution and solitary waves*, Comm. Pure. Appl. Math. **31** (1968) 467–490.
- [1971 Ha] Hasimoto, H. *Motion of a vortex filament and its relation to elastica*, J. Phys. Soc. Jap. **31** (1971) 293–.
- [1971 ZF] Zakharov, V.E., Faddeev, L.D., *Korteweg-de Vries equation, a completely integrable Hamiltonian system*, Func. Anal. Appl. **5** (1971), 280–287.
- [1972 ZS] Zakharov, V.E., Shabat, A.B., *Exact theory of two-dimensional self-focusing and one-dimensional interaction of waves in non-linear media*, Sov. Phys. JETP **34** (1972), 62–69.
- [1973 AKNS] Ablowitz, M.J., Kaup, D.J., Newell, A.C. and Segur, H., *Method for solving the sine-Gordon equation*, Phy. Rev. Lett. **30** (1973), 1262–1264.
- [1973 ZMa] Zakharov, V.E., Manakov, S.V., *On resonant interaction of wave packets in non-linear media*, JETP Letters **18** (1973) 243–247.
- [1974 Wh] Whitham, G., *Linear and non-linear waves*, Wiley, 1974.
- [1974 ZMa] Zakharov, V.E., Manakov, S.V., *The theory of resonant interaction of wave packets in non-linear media*, Sov. Phys. JETP **42** (1974), 842–850.

- [1974 FPU] Fermi, E., Pasta, J., Ulam, S., *Studies of Nonlinear Problems. I*, Lectures in Applied Math. **15** (1974), 143–155.
- [1975 Mo] Moser, J., *Three integrable Hamiltonian systems connected with isospectral deformations*, Adv. Math. **16** (1975), 197–220.
- [1976 GD_i] Gel'fand, I.M., Dikii, L.A., *Fractional Powers of Operators and Hamiltonian Systems*, Funkcional'nyi Analiz i ego Prilozhenia **10** (1976).
- [1977 Kr] Krichever, I.M., *Methods of algebraic geometry in the theory of non-linear equations*, Russian Math. Surveys **32** (1977), 185–213.
- [1978 AM] Adler, M., Moser, J., *On a class of polynomials connected with the Korteweg-deVries equation*, Comm. Math. Phys. **61** (1978), 1–30.
- [1978 De] Deift, P., *Applications of a commutation formula*, Duke. Math. J. **45** (1978), 267–310.
- [1978 ZM_i] Zakharov, V.E., Mikhailov, A.V., *Relativistically invariant 2-dimensional models of field theory which are integrable by means of the inverse scattering problem method*, Soviet Physics JETP **47** (1978), 1017–1027.
- [1979 Ad] Adler, M., *On a trace functional for formal pseudo-differential operators and the symplectic structure of the Korteweg-de Vries Type Equations*, Invent. Math. **50** (1979), 219–248.
- [1979 FRS] Frenkel, I.E., Reiman, A.G., Semenov-Tian-Shansky, M.A., *Graded Lie algebras and completely integrable dynamical systems*, Soviet Math. Dokl. **20** (1979), 811–814.
- [1979 Ko] Kostant, B., *The solution to a generalized Toda lattice and representation theory*, Adv. Math. **34** (1979), 195–338.
- [1979 ZS] Zakharov, V.E., Shabat, A.B., *Integration of non-linear equations of mathematical physics by the inverse scattering method, II*, Funct. Anal. Appl. **13** (1979), 166–174.
- [1980 AdM] Adler, M., van Moerbeke, P., *Completely integrable systems, Euclidean Lie algebras and curves*, Adv. Math. **38** (1980), 267–317.
- [1980 CT] Chern, S.S., Terng, C.L., *An analogue of Bäcklund's theorem in affine geometry*, Rocky Mountain J. Math. **10** (1980), 105–124.
- [1980 FFL] Flaschke, H., Forrest, M.G. and McLaughlin, D., *Multiphase averaging and the inverse spectral solution of the Korteweg-de Vries equation*, Comm. Pure Appl. Math **33** (1980), 739.

- [1980 Mo] Moser, J., *Various aspects of integrable Hamiltonian Systems*, Progress in Mathematics **8** (1980), 233–289, Birkhauser, Boston.
- [1980 RS] Reyman, A.G., Semenov-Tian-Shansky, *Current algebras and non-linear partial differential equations*, Sov. Math., Dokl. **21** (1980), 630–634.
- [1980 Sy] Symes, W.W., *Systems of Toda type, Inverse spectral problems, and representation theory*, Inventiones Math. **59** (1980), 13–51.
- [1980 TT] Tenenblat, K., Terng, C.L., *Bäcklund's theorem for n -dimensional submanifolds of R^{2n-1}* , Ann. Math. **111** (1980), 477–490.
- [1981 Ch] Cherednik, I.V., *Integration of the equation of a two dimensional chiral field and of its quantum analogue*, Soviet J. Nucl. Physics **33** (1981), 144–145.
- [1981a DJKM] Date, E., Jimbo, M., Kashiwara, M., Miwa, T., *Transformation groups for soliton equations I. The τ -function of the KP equation*, Proc. Japan Acad. Ser. A **57** (1981), 342–347.
- [1981 DS] Drinfel'd, V.G., and Sokolov, V.V., *Equations of Kortweg-de Vries type and simple Lie algebras*, Dokl. Akad. Nauk SSSR **258** (1981) 11–16, (Trans. as Soviet Math. Dokl. **23**, 457–462).
- [1981 KW] Kuperschmidt, B.A., Wilson, G., *Modifying Lax equations and the second Hamiltonian structure*, Invent. Math. **62** (1981), 403–436.
- [1981 S] Sato, M., *Soliton equations as dynamical systems on infinite dimensional Grassman manifolds*, RIMS Kokyuroku **439** (1982), 30–40.
- [1981 Wi] Wilson, G., *The modified Lax equations and two dimensional Toda lattice equations associated with simple Lie algebras*, Ergodic Theory and Dynamical Systems I **30** (1981), 361–380.
- [1983 Dr] Drinfel'd, V.G., *Hamiltonian structures on Lie groups, Lie bialgebras and the geometrical meaning of Yang-Baxter equations*, Sov. Math. Doklady **27** (1983), 68–71.
- [1983 DN] Dubrovin, B.A., Novikov, S.P., *Hamiltonian formalism of one dimensional systems of hydrodynamic type, and the Bogolyubov-Whitham averaging method*, Soviet Math. Dokl. **27** (1983), 781–785.
- [1984 AD] Arbarello, E. and DeConcini, C., *On a set of equations Characterizing Riemann matrices*, Ann. of Math. **120** (1984), 119–140.

- [1984 BC] Beals, R., Coifman, R.R., *Scattering and inverse scattering for first order systems*, Commun. Pure Appl. Math. **37** (1984), 39–90.
- [1984 Do] Donaldson, S., *Nahm's equations and the classification of monopoles*, Commun. Math. Physics **96** (1984), 387–407.
- [1984 DN] Dubrovin, B.A., Novikov, S.P., *On Poisson brackets of hydrodynamic type*, Soviet Math. Dokl. **30** (1984), 294–297.
- [1984 NMPZ] Novikov, S., Manakov, S., Pitaevskii, L.B. Zakharov, V.E., *Theory of Solitons*, Plenum. New York, 1984.
- [1985 BC] Beals, R., Coifman, R.R., *Inverse scattering and evolution equations*, Commun. Pure Appl. Math. **38** (1985), 29–42.
- [1985 SW] Segal, G., Wilson, G., *Loop groups and equations of KdV type*, Publ. Math. IHES **61** (1985), 5–65.
- [1987 ABT] Ablowitz, M.J., Beals, R., Tenenblat, K., *On the solution of the generalized wave and generalized sine-Gordon equations*, Stud. Appl. Math. **74** (1987), 177–203.
- [1987 Hi] Hitchin, N., *Stable bundles and integrable systems*, Duke Math. Journal **54** (1987), 91–1114.
- [1989 PS] Pinkall, U., Sterling, I., *On the classification of constant mean curvature tori*, Ann. of Math. **130** (1989), 407–451.
- [1989 U] Uhlenbeck, K., *Harmonic maps into Lie group (classical solutions of the Chiral model)*, J. Differential Geometry **30** (1989), 1–50.
- [1991 AC] Ablowitz, M.J., Clarkson, P.A., *Cambridge University*, 1991.
- [1991 Bo] Bobenko, A.I., *All constant mean curvature tori in R^3, S^3, H^3 in terms of theta functions*, Math. Ann. **290** (1991), 209–245.
- [1991 Ri] Ricca, R.L., *Rediscovery of Da Rios equations*, Nature **352** (1991), 561–562.
- [1991 Ts] Tsarëv, S.P., *The geometry of Hamiltonian systems of hydrodynamic type. The generalized Hodograph method*, Math. USSR Izvestiya **37** (1991), 397–419.
- [1991 W] Witten, E., *Two dimensional gravity and intersection theory on moduli space*, Survey in Differential Geometry **1** (1991), 243–310s.
- [1992 Hi] Hitchin, N., *Hyper-Kähler manifolds*, Astérisque **206** (1992), 137–166.

- [1993 BFPP] Burstall, F.E., Ferus, D., Pedit, F., Pinkall, U., *Harmonic tori in symmetric spaces and commuting Hamiltonian systems on loop algebras*, Annals of Math. **138** (1993), 173–212.
- [1994 KM] Kontsevich, M., Manin, Y., *Gromov-Witten classes, quantum cohomology, and enumerative geometry*, Comm. Math. Phys. **164** (1994), 625–562.
- [1994 RT] Ruan, Y.G., Tian, G., *A mathematical theory of quantum cohomology*, Math. res. Lett. **1** (1994), 269–278.
- [1995 RT] Ruan, Y.G., Tian, G., *A mathematical theory of quantum cohomology*, J. Differential Geom. **42** (1995), 259–367.
- [1996 DW] Donagi, R., Witten, E., *Supersymmetric Yang-Mills theory and integrable systems*, Nucl. Phys. **B.460** (1996), 299–334.
- [1996 D] Dubrovin, B.A., *Geometry of 2D topological field theories*, Integrable systems and quantum groups (1996), 299–334, Lecture notes in Math., v. 1620, Springer-Verlag.
- [1997 BS] Bobenko, A.I., Schief, W.K., *Affine spheres: discretization via duality relations* (to appear).
- [1997 F] Freed, D, *Special Kähler manifold* (to appear).
- [1997 Pa] Palais, R.S., *The symmetries of solitons*, Bulletin of the AMS **34** (1997), 339–403.